Title: Resting natural killer cells promote the progress of colon cancer liver metastasis by elevating tumor-derived sSCF

Author(s) and affiliation(s):
Chenchen Mao[1,2,#], Yanyu Chen[2,3,#], Dong Xing[2,#], Teming Zhang[2,#], Yangxuan Lin[4], Cong Long[2], Jiaye Yu[2], Yunhui Luo[2], Tao Ming[2], Wangkai Xie[2], Zheng Han[2], Dianfeng Mei[2], Dan Xiang[2,*], Mingdong Lu[1,*], Xian Shen[5,*], and Xiangyang Xue[1,2,*]

[1]Department of General Surgery, The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Wenzhou, China.
[2]Department of Microbiology and Immunology, Institute of Molecular Virology and Immunology, School of Basic Medical Sciences, Wenzhou Medical University, Wenzhou, China.
[3]Department of Pediatric Thoracic Surgery, The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Wenzhou, China.
[4]Department of Thoracic Surgery, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China.
[5]Department of General Surgery, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China.

[#]These authors contributed equally: Chenchen Mao, Yanyu Chen, Dong Xing and Teming Zhang.

Running title: Resting NK cells and colon cancer liver metastasis

*Correspondence should be addressed to,
Xiangyang Xue, Ph.D.
Tel: +15058788169
Chashan Higher Education Park, Ouhai District, Wenzhou City, Zhejiang Province
wzxxy@wmu.edu.cn

Xian Shen, Ph.D
Tel: +13968888872
Chashan Higher Education Park, Ouhai District, Wenzhou City, Zhejiang Province
shenxian@wmu.edu.cn

Mingdong Lu
Tel: +18757791206
No.1111 East Section of Wenzhou Avenue, Longwan District, Wenzhou City, Zhejiang Province
lumd@wmu.edu.cn

Dan Xiang, Ph.D.
Tel: +13587689767

45 Chashan Higher Education Park, Ouhai District, Wenzhou City, Zhejiang Province

46 xiangdan@wmu.edu.cn

Conflict of interest disclosure:

49 The authors declare no competing interests.

Abstract

Purpose: The abundance and biological contribution of Natural killer (NK) cells in cancer are controversial. Here, we aim to uncover clinical relevance and cellular roles of NK cells in colon cancer liver metastasis (CCLM)

Methods: We integrated single-cell RNA sequencing, spatial transcriptomics, and bulk RNA-sequencing datasets to investigate NK cells' biological properties and functions in the microenvironment of primary and liver metastatic tumors. Results were validated through an in vitro co-culture experiment based on bioinformatics analysis.

Results: We used single-cell RNA sequencing and spatial transcriptomics to map the immune cellular landscape of colon cancer and well-matched liver metastatic cancer. We discovered that GZMK+ resting NK cells increased significantly in tumor tissues and were enriched in the tumor regions of both diseases. After combining bulk RNA and clinical data, we observed that these NK cell subsets contributed to a worse prognosis. Meanwhile, KIR2DL4+ activated NK cells exhibited the opposite position and relevance. Pseudotime cell trajectory analysis revealed the evolution of activated to resting NK cells. In vitro experiments further confirmed that tumor-cell-co-cultured NK cells exhibited a decidual-like status, as evidenced by remarkable increasing CD9 expression. Functional experiments finally revealed that NK cells exhibited tumor-activating characteristics by promoting the dissociation of SCF (stem cell factor) on the tumor cells membrane depending on cell-to-cell interaction, as the supernatant of the co-culture system enhanced tumor progression.

Conclusion: Together, our findings revealed a population of protumorigenic NK cells that may be exploited for novel therapeutic strategies to improve therapeutic outcomes for patients with CCLM.

Keywords: Single cell RNA-sequencing, Spatial transcriptomics, Colon cancer liver metastasis, Natural killer cells

Introduction

Colon cancer is one of the most common malignancies, with more than 1 million new cases and 500 thousand deaths reported globally in 2020(1). Approximately 19–26% of patients with colon cancer present with synchronous metastatic diseases at the first diagnosis(2), among which 14.5–17.5% develop liver metastasis(2, 3). Although progress in the treatment of metastatic disease, including improved diagnosis and treatment strategies for liver metastases(4-6), increased cancer-directed surgery(7), and the development of targeted therapies(8), has greatly improved the survival of these patients in recent decades, the 2-year relative survival rate for patients diagnosed with distant-stage disease was 36%(2), significantly lower than those without metastasis. Thus, colon cancer-derived liver metastasis (CCLM) is a clinical challenge that requires urgent development of novel treatment methods.

The tumor microenvironment (TME) is a dynamic environment that governs tumor behavior and is required for tumor cell survival, growth, proliferation, and metastasis(9). Unlike previous understanding, it has recently been proposed that certain immune cells present in the TME (such as myeloid-derived suppressor cells(10), macrophages(11), neutrophils(12), and CD8$^+$ T cells(13)) favor tumor progression. Thus, the contradictory findings of the pro- and anti-tumor effects of tumor-infiltrating immune cells require a better understanding of the immune features and profile heterogeneity of colon cancer liver metastasis (CCLM) for the future development of immune-modulatory strategies to stratify and target immune cells.

Natural killer (NK) cells, an important component of tumor-infiltrating immune cells, are cytotoxic lymphocytes that play a key role in recognizing and eliminating malignant cells and are therefore considered the early responders against tumors(14). Among various immune cells, tumor-infiltrating NK cells are associated with prognosis in various cancers, such as colorectal cancer (CRC)(15), lung cancer(16), and gastric cancer(17). Recently, based on their anti-tumor potential, NK cells were exploited for treating malignancies and have proven to be highly promising for the treatment of certain hematologic malignancies(18, 19) but have limited efficacy for treating solid tumors(20-22). Additionally, intratumor NK cells differ phenotypically or functionally from peripheral NK cells in several malignancies(23, 24); a specific subset of tumor-infiltrating NK cells (CD11b$^-$ and CD27$^-$) as well as decidual-like NK (CD9$^+$ and cd49a$^+$) was associated with tumor progression in lung and hepatocellular cancers(25-27). Notably, the tumor-promoting transformation of NK cells may be educated by cancer cells(28, 29). However, their function remains elusive. In particular, the presence of specific NK cell subsets within colon cancer and their potential relationship to CCLM progression have not been fully characterized to date.

In this study, we combined previously published single-cell RNA sequencing, spatial transcriptomics (ST), and bulk RNA-sequencing data from public datasets, aiming to comprehensively chart the cellular landscape of TME in primary colon cancer and matched liver metastasis. Results of bioinformatics analysis were further validated through an in vitro co-culture experiment (Fig. 1A).

Materials and Methods

Data sources

In this study, we integrated four independent public datasets that contained single-cell RNA sequencing and ST data (10X Genomics) (derived from previously published research at the website: http://www.cancerdiversity.asia/scCCLM/)(30).   And two bulk transcriptomics data of colon cancer (downloaded from TCGA cohort COAD at the website: https://xena.ucsc.edu/ and NCBI's Gene Expression Omnibus with accession number GSE29623).

Single☐cell RNA Sequencing Analysis

The R (v4.0.5) package Seurat (v4.1.0)(31) was used to process the scRNA data. Since dataset quality control had been performed and Seurat objects had been created in previous studies, we did not further filter the scRNA-seq data or remove the batch effects. The SCTransform method was used to normalize the data. After selecting 2,000 highly variable genes using the Find Variable Features function in Seurat, principal component analysis was performed using these genes to reduce the data dimensions. Based on the ElbowPlot function in Seurat, we chose the top 30 principal components to run the FindNeighbors function. Next, the cells were clustered using the FindClusters function with a resolution of 0.1 for clustering immune cells. A uniform manifold approximation and projection algorithm was used for data visualization, as previously reported(32). Differentially expressed genes (DEGs) of each subset were identified using the FindAllMarkers function in Seurat. SingleR (v1.0.0) was used to name each cluster(33).   Additionally, the subgroup of NK cells was defined by the preferential expression marker of the resting and activated NK cells described in the CIBERSORT.

Pseudotime Analysis

To analyze the differentiations of NK cells, monocle2 (http://cole-trapnell-lab.github.io/monocle-release), which uses an algorithm to learn the changes in gene expression sequences that each cell must undergo as part of a dynamic biological process(34), was used for pseudotime trajectory analysis to identify the transitional relationships among different clusters. The cells were reduced dimensionally using the DDRTree method, sequenced in pseudotime, and finally visualized(35).

Bulk RNA☐seq Data Analysis

The bulk transcriptome RNA-seq data and corresponding clinical data were obtained from The Cancer Genome Atlas of colon adenocarcinoma through the UCSC Xena browser (GDC hub) (https://gdc.xenahubs.net)(36). In total, 459 colon cancer samples with survival information were downloaded, and 310 samples were finally enrolled (11 formalin-fixed samples were excluded, and an additional 134 and 4 samples were excluded due to deletion of metastasis information and sequencing matrix, respectively). Transcriptomic data from 65 colon cancer samples in GSE29623 were obtained as the validation cohorts. CIBERSORT(37), xCELL(38), EPIC(39), and MCP-counter(40), which use gene expression to infer the proportions

of tumor-infiltrating immune cells, were used to analyze the TME cell type. R packages survival (v3.2-10) and survminer (v0.4.9) were used for survival analysis. The Youden index was selected as the cutoff value to differentiate patients into distinct groups (high or low). The Kaplan–Meier survival curve was modeled using the survfit function. Subsequently, a Cox proportional hazards regression model was established to determine the independent risk factors. DEGs between the metastasis and non-metastasis groups were determined with the filtering condition of log2 |fold change|>1 and p-value <0.05, using the R package limma. GO and KEGG pathway functional enrichment analyses were performed using the clusterProfiler R package (v3.18.1) to assign various biological processes, molecular functions, cellular components, and pathways of identified marker genes in the cluster of interest(41), and p <0.05 was regarded as statistically enriched. To explore the different KEGG pathways and hallmark gene sets between the metastasis and non-metastasis groups, gene set enrichment analysis (GSEA) was performed using data from The Molecular Signatures Database (c2.cp.kegg.v7.3.symbols) and the fgsea R package (v1.12.0)(42). Pathways with an adjusted p-value below 0.05 were deemed to be significantly enriched.

ST Data Analysis

Seurat was also used for ST data processing and visualization. We used the SCT method to normalize the ST data; the functions SelectIntegrationFeatures, PrepSCTIntegration, FindIntegrationAnchors, and IntegrateData were used to integrate the ST data. An unsupervised clustering method was subsequently used to cluster similar ST spots. Cell population annotations were based on hematoxylin and eosin (H&E) staining sections and the highly variable genes in each cluster. Scores of cell-specific signatures (top 20 DEGs) from scRNA-seq were calculated using the AddModuleScore function. SpatialDimPlot and SpatialFeaturePlot were combined to visualize the cell expression level in the ST data(32). Since the tumor and non-tumor region was identified by immunohistochemistry as well as cell clusters in ST data, the differences of cell subgroup presence in tumor and non-tumor region were determined by visual inspection

Cell Lines and Co-cultures

Colon carcinoma cell lines HCT-116, DLD-1and Human Foreskin Fibroblast (HFF) were purchased from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China). HCT-116, DLD-1 were grown in a RPMI-1640 medium (GIBCO/BRL) supplemented with 10% FBS (GIBCO/BRL) while HFF were grown in the DMEM+10% FBS. The human NK cell line NK-92 was also purchased from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China) and maintained in RPMI-1640 supplemented with 12.5% horse serum (GIBCO), 12.5% FBS, 100 U/mL rhIL-2, 0.1 mmol/L b-mercaptoethanol, and 0.02 mmol/L folic acid. All the cells were cultured in a humidified chamber at 37 °C and 5% CO_2.

Colon cancer cells were plated at a density of 1×10^5 cells/well in 6-well plates for 24 h. Supernatants were collected and cells and debris therein were removed by

centrifugation. NK cells were co-cultured with cancer cells in a ratio of 1:1 in a fresh mixed medium for a further 24h (CN group). Additionally, NK cells were also cultured in a mixed medium of supernatant and NK medium in the ratio 1:1(SN group) as well as fresh RPMI-1640 10% FBS and NK medium in the same ratio (MN group) for 24h (Fig.S1A). When indicated, co-cultures were performed in Transwell devices (JET biofil), maintaining the same ratios and culture times. Finally, as shown in Fig.S1B, the co-cultured supernatant of CN, SN and MN group were collected, which determined as CNS, SNS and MNS, to incubate fresh HCT-116 to explore the NK cell-mediated inductive effect on colon cells.

Flow Cytometry Analysis

NK cells in each co-cultured group were collected and washed twice in $1 \times$ PBA. Then, they were incubated with APC-labeled CD56 (Biolegend), PE-labeled KIR2DL4 (Affinity Biosciences), FITC-labeled GZMK (Affinity Biosciences), FITC-labeled CD9 (Biolegend), PE-labeled PD-1 (Biolegend) and PerCP-labeled CD49a antibody for 30 min, rewashed, and resuspended in PBS. Fluorescence Minus One (FMO) control was set up to define the position of the positive gate as shown in Table. S1 and Table. S2. BD flow cytometry was used for detection.

Cell Counting Kit-8 (CCK-8) Cell Viability and Cell Colony Formation Assay

Colon cancer cells (5,000 cells/well) were cultured in 96-well plates with co-cultured supernatants from different co-culture groups alone or with both co-cultured supernatants and imatinib mesylate (2 μM) as previously reported(43) for 24 h. At pre-determined time points, 10 μL of CCK-8 reagent (Dojindo, Japan) was added and incubated for 2 h at 37 °C, and then the absorbance was measured using a microplate reader (Thermo Scientific) at 450 nm. All experiments were carried out in triplicate.

For the cell colony formation assay, colon cancer cells (500 cells/well) were seeded in 12□well plates with co-cultured supernatants alone or with both co-cultured supernatants and imatinib mesylate at 37 °C for 1 week. Then, cell colonies were fixed with 4% paraformaldehyde for 10 min and stained with 0.5% crystal violet for 5 min. Cell colonies containing >20 cells were counted. All experiments were carried out in triplicate.

Transwell Migration/Invasion Assays

The polycarbonate membrane in the transwell chambers was coated with Matrigel (Corning, NY, USA). Next, we transferred 1×10^5 cells from the serum-free medium with or without imatinib mesylate into the top chamber, added co-cultured supernatants alone or with both co-cultured supernatants and imatinib mesylate in the bottom chamber, and incubated at 37 °C for 24 h. Then, we removed the non-invading cells on the top side of the membrane by scrubbing, fixed the migrating or invading cells at the bottom side of the membrane with 4% paraformaldehyde, and stained with 0.5% crystal violet. The number of cells was counted under a microscope (Leica, London, UK) from four randomly chosen fields per well to determine the number of cells in each group.

Total RNA Extraction and Quantitative Real-Time PCR (qPCR)

Total RNA was extracted from cells using TRIZOL reagent (Invitrogen, USA). RNA was reverse transcribed into cDNA using a PrimeScript RT Reagent Kit (Takara, Japan). qPCR was performed using QuantStudio™ Test Development Software (Thermo Scientific, Waltham, MA, USA) with SYBR Green qPCR Master Mix (EZBioscience, Roseville, MN, USA). The sequence of KITLG and housekeeping gene GAPDH primers is listed in Table S3. KITLG mRNA data were normalized to that of GAPDH.

Luminex Liquid Suspension Chip Detection and ELISA

Luminex liquid suspension chip detection was performed by Wayen Biotechnologies (Shanghai, China). The Bio-Plex Pro Human Chemokine Panel 48-plex kit was used in accordance with the manufacturer's instructions. Briefly, supernatants from different co-cultured groups were incubated in 96-well plates embedded with microbeads for 1 h and then incubated with a detection antibody for 30 min. Subsequently, streptavidin-PE was added in each well for 10 min, and values were read using the BioPlex MAGPIX System (Bio-Rad). The Human SCF ELISA kit (Solarbio, China) was used according to the manufacturer's instructions.

Statistical Analysis and Visualization

All statistical analyses were performed using SPSS (v23.0; IBM SPSS, Chicago, IL, USA) and R (v4.0.5), and data visualization was performed on R packages Seurat (v4.1.0), ggplot2 (v3.3.5), ggsignif (v0.6.1), and pheatmap (v1.0.12).

Results

Integrated scRNA-seq and ST Precisely Quantified Immune Cell Diversity in CCLM

We used a previously published single-cell dataset containing 89 samples from paired samples of colon cancer, adjacent colon, liver metastasis, adjacent liver, lymph nodes along colons, and peripheral blood mononuclear cells from 20 patients to define the heterogeneous immune microenvironment landscapes of CCLM. Subsequently, 178,630 CD45+ cells were integrated.

To further define the main cell type, clustering analysis and SingleR were performed, and monocytes, neutrophil cells, native B cells, plasma cells, T cells, and NK cells were identified from the CCLM samples (Fig. 1B-C; Fig. S2A and B; Table.S4). Notably, T and NK cells were the major components of all immune cells and distinct across different tissue types, with NK cells significantly decreasing in tumor tissues (both primary colon cancer and liver metastasis cancer) compared to corresponding adjacent normal tissues, whereas T cells were the opposite (Fig. 1D and E; Fig. S2C).

To comprehensively analyze the spatial distribution profile of colon primary tumors and CCLM tumors, we collected ST data, which came from eight samples including paired primary colon cancer and liver metastasis. Unsupervised clustering

analysis was first performed to cluster similar ST spots, and the annotation of the
clusters was further determined according to cell marker genes (Fig. S3A-C,
Table.S5). Further, after combining the gene expression features of each sample (Fig.
S4) and H&E staining, six morphological regions, including tumor, fibroblast, smooth
muscle, B cells, hepatocytes, normal epithelium, and tumor and paratumor areas, were
identified. (Fig. 2A-C).

To integrate the scRNA-seq and ST data, we used the AddModuleScore function
in Seurat to quantify the main immune subpopulations. Consistent with scRNA-seq, T
cells were remarkably enriched in the tumor region of both primary colon cancer and
liver metastasis cancer, whereas NK cells were enriched in the non-tumor area (Fig.
2D).

Biological Relationship Between NK Cells and Metastasis of Colon Cancer
Revealed by Bulk RNA Transcriptomics
TME cell composition differences between metastasis and non-metastasis colon
cancer were evaluated using multiple tools that could robustly quantify the abundance
of cell populations based on transcriptomic datasets, including xCell, EPIC,
quanTIseq, and MCPcounter and bulk-seq datasets (TCGA COAD cohort). The
percentages of NK cells markedly decreased in metastasis colon cancer using
MCPcounter, EPIC, and xCell (Fig. 3A-D).

To further understand the potential triggers that induce colon cancer metastasis,
the DEGs were calculated between the metastasis and non-metastasis colon cancer
groups in the TCGA COAD cohort. A total of 1,378 DEGs were determined using the
limma package with cutoffs |log fold change|>1.5 and p <0.05, including 817
upregulated and 561 downregulated genes between metastasis and non-metastasis
colon cancer. Subsequently, Gene Ontology (GO) and Kyoto Encyclopedia of Genes
and Genomes (KEGG) enrichment analyses were performed to investigate the
correlative functions and pathways, As shown in Fig. 3E, cell killing, NK cell-
mediated immunity, NK cell-mediated cytotoxicity, NK cell activation, and MHC-
related GO terms and the KEGG pathways of NK cell-mediated cytotoxicity were
significantly enriched. Interestingly, nearly all the genes enriched in pathway of NK
cell-mediated cytotoxicity were markedly downregulated in the metastasis group
compared to the non-metastasis group (Fig. 3F). Similarly, Gene Set Enrichment
Analysis (GSEA) revealed that NK cell-mediated cytotoxicity was enriched in the
non-metastasis group (Fig. 3G and H).

The Landscape of NK Cells in CCLM Progression
Considering the differences in infiltrated NK cells between tumor-adjacent
tissues and tumor tissues as well as colon cancer with and without metastasis, we
further focused on NK cells to explore the detailed difference of NK subgroups in the
CCLM microenvironments. Unsupervised clustering analysis of NK cells was thus
performed and Eight NK cell subtypes were thus identified (Fig. 4A). Combining the
highly variable features of each NK cluster (Fig. 4B) and CIBERSORT-reported
canonical NK cell markers (Fig. 4C and D), NK cells were classified into three cell

types, including activated NK cells identified by the expression of KIR2DL4,
GPR183, GRP171, CD69, and IFNG, resting NK cells marked by GZMK, TTC38,
CD160, and PLEKNF1, and the other NK cells of which no characteristic gene was
identified. Although all three NK cell subtypes were presented in primary colon
tumors, adjacent normal colon tissues, liver metastasis tumors, and lymph nodes (Fig.
4E), the infiltration grade for each of these NK cell subsets was significantly different.
At the individual sample level, considerable variability was observed in the NK cell
subset composition. Where serial samples were available from individual patients, the
expression of resting NK cells increased, whereas that of activated NK cells decreased
during disease progression (Fig. 4F and G). Additionally, using the paired primary
colon tumor, adjacent normal colon tissues, and liver metastasis tumor samples, we
observed that the gradient of the resting NK cells increased, whereas the gradient of
the activated NK cells decreased (Fig. 4H).

To evaluate the clinical relationship between NK cell differences and CCLM,
CIBERSORT was performed using the TCGA COAD cohort. The percentages of
resting NK cells significantly increased (Fig. 5A), and the activated NK cells
decreased in the metastasis group (Fig. 5B), which is consistent with the scRNA-seq
results. Also, we observed that the resting NK cells were significantly increased,
whereas the activated NK cells decreased in higher T, N, and TNM stages (Fig. S5A).
In terms of survival, neutrophils and resting NK cells were associated with a worse
prognosis, whereas activated NK cells and M1 macrophages were associated with
better outcomes (Fig. S5B). Further, survival analysis showed that colon cancer
patients with lower and higher degrees of infiltration of activated NK cells and resting
NK cells, respectively, had a significantly worse prognosis in the TCGA COAD and
GSE29623 cohorts (Fig. 5C and D). Interestingly, resting NK cells were remarkably
enriched in the tumor region, whereas activated NK cells were enriched in the non-
tumor area of both primary colon cancer and liver metastasis (Fig. 5E).

371 Characterization and Developmental Course of Differential Subsets of NK Cells
372 in CCLM

To further study the heterogeneity of NK cells subgroup, the proportions of all
eight clusters of NK cells were calculated (Fig. S6). KIR2DL4+ GPR171+ activated
NK cells were lessened in proportion with the progression of the disease from normal
colon to colon cancer and liver metastasis cancer, whereas GZMK+ resting NK cells
increased (Fig. 6A and B), suggesting that these two NK subsets were associated with
metastasis. ST analysis also confirmed that there was more infiltration of the
KIR2DL4+ GPR171+ activated NK cell subset in the non-tumor area and GZMK+
resting NK cells in the tumor region, both primary colon cancer and liver metastasis
(Fig. 6C).

Pseudotime cell trajectory analysis of the two NK cell clusters was constructed to
investigate the evolutionary dynamics of metastasis-associated NK cells. NK cells
from the normal colon group were located at the top right corner of the trajectory
curve, suggesting the clear starting point of this evolving trajectory curve. After
confirmation of this starting point, developmental routes were determined to begin

with the KIR2DL4+ GPR171+ activated NK cells and then develop into GZMK+ resting NK cells (Fig. 6D).

To explore the transitional relationships of marker genes in NK cells in distinct clusters, we examined the expression level of these markers in different clusters through pseudotime analysis. Marker genes of activated NK cells (KIR2DL4 and GPR171) exhibited a gradient descent expression pattern, whereas the expression of resting NK cell marker genes (GZMK) exhibited a gradient rise pattern (Fig. 6E). BEAM analysis was performed to identify the cell cluster marker genes that change as NK cell subtypes move from the activated stage to the resting stage. The branched heatmap showed the expression dynamics of the top 200 significant genes in different cell fate branches. Corresponding GO enrichment analyses of these significant genes further demonstrated that a mass of leukocyte differentiation and immune system process-related GO terms were significantly enriched (Fig. 6F).

Interestingly, patients with colon cancer with higher infiltration of GZMK+ resting NK cells (Fig. 6G) and lower infiltration of KIR2DL4+ GPR171+ activated NK cells (Fig. 6H) exhibited shorter survival in the TCGA COAD cohort.

Tumor Cells-Educated NK Cells Shift Toward Tumor-Promoting Status Depends on Cell-to-Cell Interaction

To verify the NK cell-mediated inductive effect on colon cells, we set up mixed cell co-culture experiments using colon cell line HCT-116 and NK cell line NK-92 in the ratio 1:1 (CN group). As controls, NK cells were cultured alone, either in the tumor supernatant (SN group) or fresh medium (MN group). The co-cultured supernatant of each group were collected to incubate fresh HCT-116 after 24 h of co-culturing. Interestingly, CCK-8 and colony formation assays showed that colon cells in CNS group, which colon cells were cultured in the supernatant of co-cultured system that NK and cancer cell were in contact, underwent a significant increase in proliferation (Fig. 7A-C) compared to the CS group, where cancer cells were cultured directly in the supernatant of cancer cell after 24h culturing. Similarly, the migration and invasion of HCT-116 were also significantly increased in CNS group (Fig. 7D-F).

Furthermore, colony formation assays (Fig. 7G-H), CCK-8 (Fig. 7I), and transwell assays (Fig. 7J-L) showed colon cells in CNS group underwent a significant increase in proliferation, invasion and migration, compared to those cultured in the co-cultured supernatant (SNS group) and fresh medium (MNS group).

Additionally, another colon cancer cell line, DLD-1 was chosen to evaluate the effect of the supernatant in the different co-culture groups. DLD-1 cells in the CNS group did not undergo a prominent increase in proliferation compared to CS group (Fig. S7A-C). The same results were obtained for their migration and invasion (Fig. S7D-F). Similarly, no such cell functional changes were induced in the CNS group compared to SNS and MNS groups (Fig. S7G-L).

Given the characteristic spatial distribution of NK cell subpopulations, we hypothesized that the tumor region-enriched NK cells might be educated by tumor cells and thus functionally distinct. To verify the possible effect of tumor cells on the phenotypic switch of NK cells, NK cells were analyzed using FACS to determine the

expression of the NK cells markers above identified (GZMK and KIR2DL4) as well
as tumor infiltrated NK cells markers: CD9, CD49a and PD-1. FMO control was set
up to define the position of the positive gate as shown in Fig.S8. Upon exposure to
tumor cells, NK cells significantly decreased and slightly increased the expression of
activated marker KIR2DL4 (Fig. 7M) and the resting marker GZMK (Fig. 7N),
respectively. On the other way, for GZMK, which known as the secretory protein,
flow cytometric analyses were also performed with cell fixation and permeabilization.
However, nearly all the NK cells in different groups were GZMK positive and no
significant differences were found among each group (Fig.S9). Additionally, NK cells
in CN group underwent an increasing of tumor infiltrated NK cells markers, CD9 (Fig.
7O), CD49a (Fig. 7P) and PD-1(Fig. 7Q), especially CD9. However, no such effect
was observed after co-culturing with tumor supernatant and fresh medium.
On the other way, NK cells were also co-cultured with HFF, a normal human cell
line. The tissue infiltrated NK phenotype identified above (CD9, CD49a, PD-1) were
determined. When co-cultured with HFF in direct contact (CN group), NK cells were
also tending towards tissue infiltration state. However, the domestication effect was
significantly reduced compared to co-culturing with tumor cells (Fig.S10A and B).
Additionally, fresh HCT underwent a limited increase (no statistical significance was
found) in migration when cultured in CNS group, but not in the SNS group MNS
group (Fig.S10C-E).
451

Resting NK-induced Colon Cancer Malignant Phenotype Promotion Depends on
SCF Release

To study the potential mechanism by which the supernatant from the NK and
colon cells in direct contact with the co-culture system promoted colon cell migration,
Luminex liquid suspension chip detection was used to compare the differential
expression of 48 common chemotactic and inflammatory cytokines in the CNS, SNS,
and MNS groups. We observed that SCF was upregulated in all three repetitions in the
CNS group (Fig. 8A-B). Further, ELISA confirmed the upregulated level of SCF in
the CNS group (Fig. 8C).
We further assessed the inhibitory effects of imatinib mesylate on the enhanced
proliferation and invasion. CCK-8 (Fig. 8D-E) and colony formation assays (Fig. 8F-I)
showed that imatinib mesylate (2 μM) significantly inhibited supernatant-enhanced
proliferation of HCT-116. Similarly, the supernatant-enhance migration (Fig. 8J-K)
and invasion (Fig. 8L-M) were also inhibited by imatinib.
466

Discussion

CCLM is a multistep process and is mostly fatal(44). During disease progression,
functional interactions between tumor cells and the surrounding TME are critical for
influencing tumor growth, promoting angiogenesis, and finally resulting in
metastasis(45). Since the genomic divergence between primary and metastatic CRC
cells is relatively low(46, 47) and accumulating evidence has highlighted the key role
of tumor-infiltrating immune cells in dictating the fate of cancer cells(48, 49), we
presented a comprehensive cellular and spatial immune landscape of the primary and

liver metastatic tumors of CRC by using previously published scRNA-seq and ST data. Consistent with previous studies(30, 50), we observed that the immune microenvironment of primary and liver metastasis lesions of colon cancer had undergone extensive remodeling with a significantly decreased proportion of NK cells and a strong enrichment of T cells. However, the contrast between the decreased proportion of NK cells in tumor tissue and the enrichment of NK cells in tumor regions may indicate the specific effects of NK cells in CCLM.

Traditionally, NK cells within the TME were known as potential suppressors for tumor growth by directly killing cells and secreting proinflammatory cytokines(51, 52). However, the function of NK cells highly depends on their maturation status and localization(53). The peripheral blood $CD56^{dim}CD16^{high}$ NK cell population predominantly mediates the killing of target cells by secreting perforin and granzymes(54), whereas the $CD56^{bright}CD16^{low}$ NK cells that reside in secondary lymphoid tissues are immature and have reduced cytotoxic capability(55). Recently, tumor-associated NK cells, which are enriched in tumors with impaired anti-tumor functions, were discovered to be associated with an unfavorable prognosis and resistance to immunotherapy in multiple solid tumors (56), such as lung cancer, pancreatic cancer, and esophageal cancer. The potential impact of NK cells on CCML progression is still poorly investigated.

In this study, the proportion of NK cells was higher than in past cognition, similar to a previous study(57) in that NK cells occupied the largest immune cell compartment of nearly 50% in cholangiocarcinoma. In contrast, NK cells may not be well distinguished during automatic annotation using SingleR. Consistent with the study that revealed that treatment responders are associated with TME remodeling, including NK cell recruitment(58), the proportions of NK cells in PR patients were significantly higher among the LiverP and LiverT groups (Fig. S2D and E), which also increased the overall proportion of NK cells to some extent. Additionally, we observed that NK cells are decreased in primary colon cancer and liver metastatic using scRNA-seq, consistent with previous findings supporting that NK cells play a crucial role in anticancer immunity(59, 60). However, we further subdivided NK cells considering the conflicting result, in which NK cells were significantly enriched in the tumor area when using ST. Since this is the first study to comprehensively explore the heterogeneity of NK cells, no previous NK cell subpopulation markers were available; resting and activated NK cell markers in CIBERSORT were thus used to annotate NK cells. Consistent with a recent study that reported that mature and immature NK cells serve different functions(61), our results also provide evidence that resting NK cells in the colon cancer TME possess a potential tumor-promoting activity, whereas activated NK cells play an anti-tumor role as conventional cognition. Bulk RNA-sequencing and the corresponding clinical data further indicated that higher infiltration of resting NK cells and lower infiltration of activated NK cells were correlated with a worse prognosis in patients with CRC. However, since the resting and activated NK cells we identified were only the aggregation of multiple NK cell subpopulations, further studies are required to investigate the key NK cell subpopulations that play a crucial role in CCLM.

Further analysis of NK cell subtypes revealed a dramatic change in KIR2DL4+
GPR171+ activated NK cells and GZMK+ resting NK cells, whereas no other NK cell
subpopulations differed in colon cancer progression. KIR2DL4, also known as
CD158d, is an unusual member of the killer cell Ig-like receptor family expressed in
all NK cells(62). Studies revealed that KIR2DL4 activates the cytotoxicity of NK
cells(63), and NK-92 cells stimulated with KIR2DL4 had higher cytotoxicity against
breast cancer cells(64). Consistent with previous studies, we also observed that the
expression of the KIR2DL4+ GPR171+ activated NK cell subtype was decreased
during CCLM progression, possibly reflecting the anti-tumor activity of this
population. However, the GZMK+ resting NK cell subsets were first identified as
potential tumor promoters. Moreover, it has been proven that GZMK^{high} CD8^{+} T
effector memory cells, a cell subset that is particularly similar to GZMK+ NK cells,
were associated with poor clinical outcomes in patients with colorectal tumors. Our
results also showed that GZMK+ resting NK cells highly infiltrate the cancer region,
and this predicts a worse prognosis in patients with colon cancer. However, despite
the interesting and gratifying clinical relevance, the mechanism remains unknown.
Considering the evolutionary trajectory and the characteristic distribution of
these metastasis-associated NK cell subtypes, we predicted that activated NK cells
might differentiate into resting NK cells under the action of tumor cells. Consistent
with a recent study that indicated that exposure to cancer cells causes NK cells to lose
their cytotoxic ability(28), we identified the tendency of NK cells to differentiate to an
resting state with decreasing expression of activated marker KIR2DL4 as well as
increasing expression of decidual-like NK cells markers: CD9, CD49a and PD-1(27),
which are only triggered by cell-to-cell interactions as validated by FACS. It is worth
mentioning that decidual-like NK cells were the newly identified subset of NK cells
that promote tumors(27).
Additionally, NK-mediated tumor cell editing appears to partly depend on the
release of cytokines in in vitro co-culture experiments. SCF, the natural ligand of c-
Kit with membrane-bound (mSCF) and soluble forms (sSCF)(65), enhances the
growth and migration of cancer cells such as cervical cancer(66), Ewing's sarcoma(67)
and CRC(43). Although the dimeric mSCF is remarkably more active and induces a
more persistent activation of the receptor(65, 68), the tumor-promoting effect of the
supernatant from the co-culture system in the current study and the reversion effect of
imatinib for the co-culture supernatant induced pro-oncogenic effects suggests an
important role of sSCF on colon cancer cells. However, no up-regulation of KITLG
(the SCF encoding gene) was found in HCT-116 after co-culturing with NK cells (Fig.
S11A). In addition, when using DLD-1, a colon cancer cell with a nearly 15 times
lower expression of KITLG than HCT-116 (Fig. S11B), we also did not observe co-
cultured supernatant induced tumor-promoting effects. We thus hypothesized that in
our experimental system, NK cells may promote the cleavage and release of the
extracellular part of SCF on the tumor cell surface after direct contact with tumor cells,
and thus increase the concentration of sSCF, which induces the malignant phenotype
of colon cancer cells in a paracrine manner (Fig. 9). However, the detailed mechanism
still needs to be further studied.

In summary, we have unveiled the spatial and cellular profiles of TME from tumors and paratumor tissues of CCLM. Our analysis uncovered the different states, functions, and dynamic nature of NK cells in different CCLM settings, which can be used for further identification of regulatory mechanisms and for developing potential therapeutic targets.

Data Availability

All data generated or analyzed during this study are available from the corresponding author on reasonable request.

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	32070151	Xian Shen
Key R&D Program of Zhejiang Province	2020C03029	Xiangyang Xue
Key R&D Program of Zhejiang Province	2021C03120	Xian Shen
the Zhejiang Provincial Natural Science Foundation Project	Q24H160147	Wangkai Xie
the Postdoctoral Fellowship Program of CPSF	GZC20231956	Wangkai Xie
the Wenzhou Municipal Science and Technology Bureau Program	Y2020215	Dan Xiang
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication		

Acknowledgement

We thank Prof. Qiang Gao from Fudan University and Prof. Xiaoming Zhang from Shanghai Institute of Immunity and Infection, CAS for sharing the SC RNA sequencing and ST data. We thank Zhejiang Key Laboratory of New Techniques for Diagnosis and Treatment of Critical Diseases of Pancreas and Liver and Zhejiang International science and Technology Cooperation base for tumor transformation research for supporting this study. We thank Editage Group (https://www.editage.cn/) for polishing the draft of this manuscript.
]

Authors' Contributions:
CCM and **YYC**: Conceptualization, experimental design and execution, data curation, method development, formal analysis, statistics, writing original draft, and editing the manuscript. **DX** and **TMZ**: conceptualization, data curation and method development. **YXL, CL** and **JYY**: experimental design. **MHL** and **TM**: experimental design. **WKX, DFM** and **ZH**: Validation. **DX, MDL** and **XS**: Experimental design, supervision,

591   funding acquisition, and manuscript editing. **XYX XS**: Conceptualization, supervision,
592   writing manuscript, project administration.
593

References

1. Sung H, *et al.* (2021) Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 71(3):209-249.

2. Siegel RL, *et al.* (2020) Colorectal cancer statistics, 2020. *CA Cancer J Clin* 70(3):145-164.

3. Wang F, *et al.* (2023) Single-cell and spatial transcriptome analysis reveals the cellular heterogeneity of liver metastatic colorectal cancer. *Sci Adv* 9(24):eadf5464.

4. Chua TC, Saxena A, Chu F, Zhao J, & Morris DL (2011) Predictors of cure after hepatic resection of colorectal liver metastases: an analysis of actual 5- and 10-year survivors. *J Surg Oncol* 103(8):796-800.

5. Zampino MG, *et al.* (2016) Treatments for colorectal liver metastases: A new focus on a familiar concept. *Crit Rev Oncol Hematol* 108:154-163.

6. Ruers T, *et al.* (2017) Local Treatment of Unresectable Colorectal Liver Metastases: Results of a Randomized Phase II Trial. *J Natl Cancer Inst* 109(9).

7. Wancata LM, Banerjee M, Muenz DG, Haymart MR, & Wong SL (2016) Conditional survival in advanced colorectal cancer and surgery. *J Surg Res* 201(1):196-201.

8. Piawah S & Venook AP (2019) Targeted therapy for colorectal cancer metastases: A review of current methods of molecularly targeted therapy and the use of tumor biomarkers in the treatment of metastatic colorectal cancer. *Cancer* 125(23):4139-4147.

9. Yao J, Chen Y, & Lin Z (2023) Exosomes: Mediators in microenvironment of colorectal cancer. *Int J Cancer* 153(5):904-917.

10. Liu H, Wang Z, Zhou Y, & Yang Y (2023) MDSCs in breast cancer: an important enabler of tumor progression and an emerging therapeutic target. *Front Immunol* 14:1199273.

11. Ngambenjawong C, Gustafson HH, & Pun SH (2017) Progress in tumor-associated macrophage (TAM)-targeted therapeutics. *Adv Drug Deliv Rev* 114:206-221.

12. Li S, *et al.* (2019) Tumor-associated neutrophils induce EMT by IL-17a to promote migration and invasion in gastric cancer cells. *J Exp Clin Cancer Res* 38(1):6.

13. Tiberti S, *et al.* (2022) GZMK(high) CD8(+) T effector memory cells are associated with CD15(high) neutrophil abundance in non-metastatic colorectal tumors and predict poor clinical outcome. *Nat Commun* 13(1):6752.

14. Maskalenko NA, Zhigarev D, & Campbell KS (2022) Harnessing natural killer cells for cancer immunotherapy: dispatching the first responders. *Nat Rev Drug Discov* 21(8):559-577.

15. Zhong F, *et al.* (2022) Innate tumor killers in colorectal cancer. *Cancer Lett* 527:115-126.

16. Villegas FR, *et al.* (2002) Prognostic significance of tumor infiltrating natural killer cells subset CD57 in patients with squamous cell lung cancer. *Lung Cancer* 35(1):23-28.

17. Ishigami S, *et al.* (2000) Prognostic value of intratumoral natural killer cells in gastric carcinoma. *Cancer* 88(3):577-583.

18. Lamb MG, Rangarajan HG, Tullius BP, & Lee DA (2021) Natural killer cell therapy for hematologic malignancies: successes, challenges, and the future. *Stem Cell Res Ther* 12(1):211.

19. Liu E, *et al.* (2020) Use of CAR-Transduced Natural Killer Cells in CD19-Positive Lymphoid Tumors. *N Engl J Med* 382(6):545-553.

20. Zhang H, Yang L, Wang T, & Li Z (2024) NK cell-based tumor immunotherapy. *Bioact Mater* 31:63-86.

21. Hu W, Wang G, Huang D, Sui M, & Xu Y (2019) Cancer Immunotherapy Based on Natural

638   Killer Cells: Current Progress and New Opportunities. *Front Immunol* 10:1205.

22.   Oh S, Lee JH, Kwack K, & Choi SW (2019) Natural Killer Cell Therapy: A New Treatment
Paradigm for Solid Tumors. *Cancers (Basel)* 11(10).

23.   Bruno A, *et al.* (2013) The proangiogenic phenotype of natural killer cells in patients with
non-small cell lung cancer. *Neoplasia* 15(2):133-142.

24.   Carrega P, *et al.* (2008) Natural killer cells infiltrating human nonsmall-cell lung cancer are
enriched in CD56 bright CD16(-) cells and display an impaired capability to kill tumor cells.
Cancer 112(4):863-875.

25.   Zhang QF, *et al.* (2017) Liver-infiltrating CD11b(-)CD27(-) NK subsets account for NK-cell
dysfunction in patients with hepatocellular carcinoma and are associated with tumor
progression. *Cell Mol Immunol* 14(10):819-829.

26.   Jin J, *et al.* (2013) CD11b(-)CD27(-) NK cells are associated with the progression of lung
carcinoma. *PLoS One* 8(4):e61024.

27.   Albini A & Noonan DM (2021) Decidual-Like NK Cell Polarization: From Cancer Killing to
Cancer Nurturing. *Cancer Discov* 11(1):28-33.

28.   Chan IS, *et al.* (2020) Cancer cells educate natural killer cells to a metastasis-promoting cell
state. *J Cell Biol* 219(9).

29.   Huergo-Zapico L, *et al.* (2018) NK-cell Editing Mediates Epithelial-to-Mesenchymal
Transition via Phenotypic and Proteomic Changes in Melanoma Cell Lines. *Cancer Res*
78(14):3913-3925.

30.   Wu Y, *et al.* (2022) Spatiotemporal Immune Landscape of Colorectal Cancer Liver Metastasis
at Single-Cell Level. *Cancer Discov* 12(1):134-153.

31.   Hao Y, *et al.* (2021) Integrated analysis of multimodal single-cell data. *Cell* 184(13):3573-
3587 e3529.

32.   Peng Z, *et al.* (2023) Interactions between MFAP5 + fibroblasts and tumor-infiltrating
myeloid cells shape the malignant microenvironment of colorectal cancer. *J Transl Med*
21(1):405.

33.   Zheng H, Liu H, Ge Y, & Wang X (2021) Integrated single-cell and bulk RNA sequencing
analysis identifies a cancer associated fibroblast-related signature for predicting prognosis and
therapeutic responses in colorectal cancer. *Cancer Cell Int* 21(1):552.

34.   Hong B, *et al.* (2022) Single-cell transcriptional profiling reveals heterogeneity and
developmental trajectories of Ewing sarcoma. *J Cancer Res Clin Oncol* 148(12):3267-3280.

35.   Qiu X, *et al.* (2017) Single-cell mRNA quantification and differential analysis with Census.
Nat Methods 14(3):309-315.

36.   Goldman MJ, *et al.* (2020) Visualizing and interpreting cancer genomics data via the Xena
platform. *Nat Biotechnol* 38(6):675-678.

37.   Chen B, Khodadoust MS, Liu CL, Newman AM, & Alizadeh AA (2018) Profiling Tumor
Infiltrating Immune Cells with CIBERSORT. *Methods Mol Biol* 1711:243-259.

38.   Aran D, Hu Z, & Butte AJ (2017) xCell: digitally portraying the tissue cellular heterogeneity
landscape. *Genome Biol* 18(1):220.

39.   Racle J, de Jonge K, Baumgaertner P, Speiser DE, & Gfeller D (2017) Simultaneous
enumeration of cancer and immune cell types from bulk tumor gene expression data. *Elife* 6.

40.   Becht E, *et al.* (2016) Erratum to: Estimating the population abundance of tissue-infiltrating
immune and stromal cell populations using gene expression. *Genome Biol* 17(1):249.

41.   Yu G, Wang LG, Han Y, & He QY (2012) clusterProfiler: an R package for comparing
biological themes among gene clusters. *OMICS* 16(5):284-287.

42.   Liberzon A, *et al.* (2015) The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* 1(6):417-425.

43.   Yasuda A, *et al.* (2007) Stem cell factor/c-kit receptor signaling enhances the proliferation and
invasion of colorectal cancer cells through the PI3K/Akt pathway. *Dig Dis Sci* 52(9):2292-
2300.

44.   Manfredi S, *et al.* (2006) Epidemiology and management of liver metastases from colorectal
cancer. *Ann Surg* 244(2):254-259.

45.   Quail DF & Joyce JA (2013) Microenvironmental regulation of tumor progression and
metastasis. *Nat Med* 19(11):1423-1437.

46.   Wei Q, *et al.* (2017) Multiregion whole-exome sequencing of matched primary and metastatic
tumors revealed genomic heterogeneity and suggested polyclonal seeding in colorectal cancer
metastasis. *Ann Oncol* 28(9):2135-2141.

47.   Yaeger R, *et al.* (2018) Clinical Sequencing Defines the Genomic Landscape of Metastatic
Colorectal Cancer. *Cancer Cell* 33(1):125-136 e123.

48.   Kumar S, *et al.* (2018) DeltaNp63-driven recruitment of myeloid-derived suppressor cells
promotes metastasis in triple-negative breast cancer. *J Clin Invest* 128(11):5095-5109.

49.   Hanahan D & Coussens LM (2012) Accessories to the crime: functions of cells recruited to
the tumor microenvironment. *Cancer Cell* 21(3):309-322.

50.   Zhou SN, *et al.* (2021) Comparison of Immune Microenvironment Between Colon and Liver
Metastatic Tissue in Colon Cancer Patients with Liver Metastasis. *Dig Dis Sci* 66(2):474-482.

51.   Huntington ND, Vosshenrich CA, & Di Santo JP (2007) Developmental pathways that
generate natural-killer-cell diversity in mice and humans. *Nat Rev Immunol* 7(9):703-714.

52.   Freud AG & Caligiuri MA (2006) Human natural killer cell development. *Immunol Rev*
214:56-72.

53.   Cooper MA, Fehniger TA, & Caligiuri MA (2001) The biology of human natural killer-cell
subsets. *Trends Immunol* 22(11):633-640.

54.   Jacobs R, *et al.* (2001) CD56bright cells differ in their KIR repertoire and cytotoxic features
from CD56dim NK cells. *Eur J Immunol* 31(10):3121-3127.

55.   Moretta L (2010) Dissecting CD56dim human NK cells. *Blood* 116(19):3689-3691.

56.   Tang F, *et al.* (2023) A pan-cancer single-cell panorama of human natural killer cells. *Cell*.

57.   Xia T, *et al.* (2022) Immune cell atlas of cholangiocarcinomas reveals distinct tumor
microenvironments and associated prognoses. *J Hematol Oncol* 15(1):37.

58.   Kim R, *et al.* (2022) Early Tumor-Immune Microenvironmental Remodeling and Response to
First-Line Fluoropyrimidine and Platinum Chemotherapy in Advanced Gastric Cancer. *Cancer*
Discov 12(4):984-1001.

59.   Laskowski TJ, Biederstadt A, & Rezvani K (2022) Natural killer cells in antitumour adoptive
cell immunotherapy. *Nat Rev Cancer* 22(10):557-575.

60.   Myers JA & Miller JS (2021) Exploring the NK cell platform for cancer immunotherapy. *Nat*
Rev Clin Oncol 18(2):85-100.

61.   Thacker G, *et al.* (2023) Immature natural killer cells promote progression of triple-negative
breast cancer. *Sci Transl Med* 15(686):eabl4414.

62.   Andre P, *et al.* (2001) New nomenclature for MHC receptors. *Nat Immunol* 2(8):661.

63. Faure M & Long EO (2002) KIR2DL4 (CD158d), an NK cell-activating receptor with
inhibitory potential. *J Immunol* 168(12):6208-6214.

64. Kilic N, Dastouri M, Kandemir I, & Yilmaz E (2023) The effects of KIR2DL4 stimulated NK-
92 cells on the apoptotic pathways of HER2 + /HER-breast cancer cells. *Med Oncol* 40(5):139.

65. Hsu YR*, et al.* (1997) The majority of stem cell factor exists as monomer under physiological
conditions. Implications for dimerization mediating biological activity. *J Biol Chem*
272(10):6406-6415.

66. Aguilar C, Aguilar C, Lopez-Marure R, Jimenez-Sanchez A, & Rocha-Zavaleta L (2014) Co-
stimulation with stem cell factor and erythropoietin enhances migration of c-Kit expressing
cervical cancer cells through the sustained activation of ERK1/2. *Mol Med Rep* 9(5):1895-
1902.

67. Landuzzi L*, et al.* (2000) The metastatic ability of Ewing's sarcoma cells is modulated by stem
cell factor and by its receptor c-kit. *Am J Pathol* 157(6):2123-2131.

68. Langley KE*, et al.* (1993) Soluble stem cell factor in human serum. *Blood* 81(3):656-660.

740
741

Figure legends

Figure 1. Landscape of tumor immune microenvironment in CCLM revealed by
single-cell transcriptomics.
A. Schematic overview of the experimental design and analytical workflow. Colon T:
colon tumor tissue; Colon P: colon paratumor normal tissue; Liver T: liver metastasis
tumor tissue; Liver P: liver paratumor normal tissue; LN: lymph node metastasis
tissue. B. The UMAP plot of all main immune cell types. C. Dot plots showing
average expression of highly variable genes of each cell group. The dot size
represents percent of cells expressing the genes in each cluster and the color of dot
represents the expression intensity. D. The cell numbers of main immune cells across
tissues. E. Proportions of all main immune cells. P values were determined by the
paired nonparameter test.
754

Figure 2. Cellular identification in spatial transcriptomic samples.
A. UMAP visualization of cell clusters in spatial transcriptomic samples. B. Dot plots
showing average expression of markers in indicated cell clusters. C. Overview of the
spatial transcriptomic sections. H&E staining of spatial transcriptomic sections
(upper). Tumor tissue and paratumor tissue identification of each section (middle).
Spatial cluster distribution of each section (lower). PC: primary cancer; LM: liver
metastasis.
ST-P1: spatial transcriptomic dataset patient 1, ST-P1: spatial transcriptomic dataset
patient 1; ST-P2: spatial transcriptomic dataset patient 2; ST-P3: spatial transcriptomic
dataset patient 3; ST-P4: spatial transcriptomic dataset patient 4. D. The signature
scores of T cells (upper) and NK cells (lower) in colon cancer and liver metastasis in
the spatial transcriptomic sections.
767

Figure 3. Clinical and biological relationship between NK cells and metastasis of
colon cancer revealed by Bulk RNA transcriptomics.
A-D. The relationship of immune cell percentage determined by xCell, EPIC,
quanTIseq and MCPCounter between metastasis and non-metastasis tumor in TCGA
COAD cohort. E. Gene Ontology (GO) and Kyoto Encyclopedia of Genes and
Genomes (KEGG) enriched pathway bar chart of Differentially expressed genes
(DEGs) in metastasis versus non-metastasis colon cancer. F. Volcano plot showing
nearly all the genes enriched in pathway of NK cell-mediated cytotoxicity were
downregulated in the metastasis colon cancer. G. Gene set enrichment analysis
(GSEA) of KEGG gene set. H. Natural killer cell mediated cytotoxicity was enriched
in the non-metastasis colon cancer.
779

Figure 4: The landscape of NK cells in the disease progression of CCLM.
A. The UMAP plot of NK cells from CCLM. B. Unsupervised clustering identifies 8
subsets of NK cells. C. Expression of key markers that distinguish resting and
activated subsets of NK cells. D. Expression of key resting and activated NK cell
markers across all samples. E. The UMAP plot of distribution of resting and activated
NK cells from ColonP, ColonT, LiverT and LN. F. Cellular landscape of each sample

from the ColonP, ColonT, LiverT and LN. The proportion of NK cells subsets in total
immune cells (upper). The proportion of NK cells subsets in total NK cells (lower). G.
Number of cells identified from each group (ColonP, ColonT, LiverT and LN) by cell
type proportion. H. Proportions of resting (upper) and activated (lower) subsets of NK
cells. P values were determined by the paired nonparameter test.
791

Figure.5 Clinical relationship between NK cells subsets and metastasis of colon
cancer revealed by Bulk RNA and spatial transcriptomics.
A-B. The relationship of activated and resting NK cell percentage determined by
CIBERSORT and tumor metastasis in TCGA COAD cohort. M0: non-metastasis
colon cancer; M1: metastasis colon cancer. C-D. K-M survival plots show that high
resting NK cell and low activated NK cell predicted poor prognosis in TCGA COAD
and GSE29623 cohort. PFI: Progression Free Interval; OS: Overall Survival. E. The
signature scores of resting (upper) and activated NK cells (lower) in colon cancer and
liver metastasis in the spatial transcriptomic sections.
801

Figure 6. Characterization and developmental course of differential subsets of
NK cells in CCLM.
A. The UMAP plot of KIR2DL4+ GPR171+ Activated NK cells, GZMK+ resting NK
cells from CCLM. B. Proportions of KIR2DL4+ GPR171+ ActiVated NK cells,
GZMK+ resting NK cells in total immune cells. P values were determined by the
paired nonparameter test. C. The signature scores of GZMK+ resting NK cells (upper)
and KIR2DL4+ GPR171+ Activated NK cells (lower) in colon cancer and liver
metastasis in the spatial transcriptomic sections. D. Monocle analysis showing the
developmental trajectory of NK cells. Color as in Pseudotime, cell state, subsets of
NK cells and sample group. E. Pseudo□temporal change curve of marker genes in
each subsets of NK cells. F. The heatmap shows the expression patterns of the top 50
significant genes in branched expression analysis modeling, associated GO terms
(using DAVID v6.7) are given on the right of the corresponding gene clusters. G-H.
The Kaplan–Meier curve shows COAD patients survival with different GZMK+
resting NK cells and KIR2DL4+ GPR171+ Activated NK cells infiltration.
817

Figure.7 Colon cancer cells (HCT-116) educated NK cells shift toward tumor
promoting status depends on cell-to-cell interaction.
A-B. Clone formation assay showed the NK cell-mediated inductive effect on cell
proliferation of colon cancer cell (HCT-116). Colon cancer cells were cultured in the
supernatant from different co-culture system in transwell devices. CNS: colon cancer
cells were cultured in the supernatant from co-culture system that NK cells and cancer
cells were cultured in direct contact (CNS); CS: cancer cells cultured directly in
supernatant of cancer cells. C. Cell Counting Kit-8 (CCK-8) assay showed the NK
cell–mediated inductive effect on cell proliferation of colon cancer cell. D-F: The NK
cell–mediated inductive effect on migration and invasion of colon cancer cell. G-H:
Clone formation assay CCK-8 assay showed the NK cell–mediated inductive effect
on cell proliferation of colon cancer cell. SNS: colon cancer cells were cultured in the

supernatant from co-culture system that NK cells were cultured in supernatant of
cancer cells; MNS: colon cancer cells were cultured in the supernatant from co-
culture system that NK cells were cultured in fresh medium. M-Q. Phenotype switch
(KIR2DL4, GZMK, CD9, CD49a, PD-1) of NK cells was induced by cell-to-cell
interactions with cancer cells. CN: NK cells were co-cultured with colon cancer cells;
SN: NK cells were cultured in supernatant of cancer cells; MN: NK cells were
cultured in fresh medium.
837
Figure.8 The resting NK cell promote tumor malignant phenotype via elevating
tumor-derived sSCF.
A. luminex liquid suspension chip detection of 48 common chemotactic and
inflammatory cytokines in CNS, SNS and MNS group. B-C. Concentration of SCF
determined by luminex liquid suspension chip and Elisa in CNS, SNS and MNS
group. D-E. CCK-8 assay showed the proliferation of HCT-116 cells was inhibited by
imatinib mesylate, evaluated by a CCK-8 assay. Cells were incubated in the
supernatant from different co-culture system with DMSO or 2 µM imatinib mesylate.
F-I. Clone formation assay showed the proliferation of HCT-116 cells was inhibited
by imatinib mesylate. J-M. The NK cell–mediated inductive effect on migration and
invasion of HCT-116 was inhibited by imatinib mesylate.
849
850 **Figure.9. Schematic diagram.**
851 Colon cancer cells (HCT-116 cells) educate NK cells as resting status depends on cell-
852 to-cell interaction. Tumor-educated NK cells subsequent enhances tumor malignancy
853 in a paracrine manner by elevating tumor-derived sSCF.
854

Supplemental Figures legends:

Figure S1. Schematic overview of the in vitro experimental design.

A. Schematic overview of coculture experiments design. B. Schematic overview of

coculture experiments design in transwells.

Figure S2. Cluster characterization of the global landscape of CCLM revealed by

single-cell transcriptomics.

A. The UMAP plot of all 12 cell clusters. B. Dot plots showing average expression of

marker genes in indicated cell clusters. C. The pie chart of the proportion of main

immune cells across tissues. D. The proportion of NK cells in total immune cells of

each sample (including PR, PD/SD and untreated patients) from the Liver P, Liver T,

Colon P, Colon T. E. Proportions of NK cells in PR, PD/SD and untreated patients

among Liver P, Liver T, Colon P, Colon T groups.

Figure S3. Cellular identification in spatial transcriptomic samples.

A. The UMAP plot of all 11 cell clusters. B. Dot plots showing average expression of

marker genes in indicated cell clusters. C. Expression of key markers across all

samples.

Figure S4. Gene expression features of each sample. The feature plots showed the

expression distributions of ALB, CD24, COL1A1, MYH11, CXCL8, CD14,

CD79A AND CD3D in each spatial transcriptomic section.

Figure.S5 Clinical relationship between NK cells subsets and colon cancer

revealed by TCGA COAD cohort.

A. The relationship of activated and resting NK cell and clinical characteristics of

colon cancer in TCGA COAD cohort. B. The relationship of 22 immune cells

percentage determined by CIBERSORT and prognosis of colon cancer in TCGA

COAD cohort.

Figure.S6 The proportions of eight clusters of NK cells in Colon P, Colon T, Liver

T and LN.

Figure.S7 NK cell-mediated tumor promoting effect in colon cancer cells (DLD-1).

A. Clone formation assay showed the NK cell–mediated inductive effect on cell proliferation of DLD-1 cell between CNS and CS groups. B-C: CCK-8 assay showed the NK cell–mediated inductive effect on cell proliferation of DLD-1 cell between CNS and CS groups. D-F. The NK cell–mediated inductive effect on migration and invasion of DLD-1 cell between CNS and CS groups. G. CCK-8 assay showed the NK cell-mediated inductive effect on cell proliferation of DLD-1 cell among CNS, SNS, MNS groups. H-I. Clone formation assay showed the NK cell-mediated inductive effect on cell proliferation of DLD-1 cell among CNS, SNS, MNS groups. J-L The NK cell–mediated inductive effect on migration and invasion of DLD-1cell among CNS, SNS, MNS groups.

Figure.S8 The positive gate locations of CD56, GZMK, KIR2DL4, CD9, CD49a, PD-1 defined according to the FMO control.

Figure.S9 Phenotype switch (CD56+, GZMK+) of NK cells was analyzed by FACS after fixation and permeabilization in different co-cultured groups.

Figure.S10 Phenotype switch of NK cells in different co-cultured system and the corresponding NK cell-mediated effect on cell migration of fresh colon cancer cell (HCT-116).

A-B: NK cells underwent phenotype switch (high expression of CD9) when cocultured with HCT and HFF, the phenotype switch was more obvious when co-cultured with HCT. CN: NK cells cocultured with HCT/HFF; SN: NK cells cocultured with supernatant of HCT/HFF; MN: NK cells cocultured in fresh medium. C-E: Transwell assay showed the only tumor co-cultured NK mediated the inductive effect on cell migration of colon cancer cell (HCT-116). CNS: Colon cancer cells were cultured in the supernatant from co-culture system that NK and HCT/HFF were cultured in direct contact; SNS: Colon cancer cells were cultured in the supernatant from co-culture system that NK cocultured with supernatant of HCT/HFF; MNS: Colon cancer cells were cultured in the fresh medium.

Figure.S11 Relative KITLG expression in different groups.

A. RT-qPCR analysis of KITLG expression in HCT-116 with and without co-cultured with NK cells. B. RT-qPCR analysis of KITLG expression in HCT-116 and DLD-1 cells.