Genome-wide association study reveals multiple loci for nociception and opioid consumption behaviors associated with heroin vulnerability in outbred rats

Brittany N. Kuhn[1], Ph.D., Nazzareno Cannella[2], Ph.D., Apurva S. Chitre[3], M.S., Khai-Minh H. Nguyen[3], B.S., Katarina Cohen[3], B.S., Denghui Chen[3], B.S., Beverly Peng[3], B.S., Kendra S. Ziegler[3], B.S, Bonnie Lin[3], B.S., Benjamin B. Johnson[3], Ph.D., Thiago Missfeldt Sanches[3], Ph.D., Ayteria D. Crow[1], B.S., Veronica Lunerti[2], Ph.D., Arkobrato Gupta[4-6], M.S., Eric Dereschewitz[1], B.S., Laura Soverchia[2], Ph.D., Jordan L. Hopkins[1], M.S., Analyse T. Roberts[1], B.S., Massimo Ubaldi[2], Ph.D., Sarah Abdulmalek[7], Ph.D., Analia Kinen[2,7], B.S., Gary Hardiman[7,8], Ph.D., Dongjun Chung[4-6], Ph.D., Oksana Polesskaya[3], Ph.D., Leah C. Solberg Woods[9], Ph.D., Roberto Ciccocioppo[2], Ph.D., Peter W. Kalivas[1], Ph.D., and Abraham A. Palmer, Ph.D.[3,10]

[1]Department of Neuroscience, Medical University of South Carolina, Charleston, SC, USA

[2]School of Pharmacy, Center for Neuroscience, Pharmacology Unit, University of Camerino, Camerino, IT

[3]Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

[4]The Interdisciplinary Ph.D. Program in Biostatistics, The Ohio State University, OH, USA

[5]Department of Biomedical Informatics, The Ohio State University, Columbus, OH, USA

[6]Pelotonia Institute for Immuno-Oncology, The James Comprehensive Cancer Center, The Ohio State University, OH, USA

[7]School of Biological Sciences, Queen's University Belfast, Belfast, Northern Ireland, UK

[8]Departments of Medicine and Public Health Sciences, Medical University of South Carolina, Charleston, SC, USA

[9]Department of Internal Medicine, Wake Forest University, Winston-Salem, NC, USA

[10]Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA

Corresponding author:
Brittany N. Kuhn
Medical University of South Carolina
173 Ashley Avenue
BSB 403- MSC 510
Charleston, SC 29425
e-mail: kuhnb@musc.edu
Phone: 703-298-2596
Fax: +1-843-792-4423

Running Title: Genetic loci associated with heroin vulnerability

Abstract

The increased prevalence of opioid use disorder (OUD) makes it imperative to disentangle the biological mechanisms contributing to individual differences in OUD vulnerability. OUD shows strong heritability, however genetic variants contributing toward vulnerability remain poorly defined. We performed a genome-wide association study using over 850 male and female heterogeneous stock (HS) rats to identify genes underlying behaviors associated with OUD such as nociception, as well as heroin-taking, extinction and seeking behaviors. By using an animal model of OUD, we were able to identify genetic variants associated with distinct OUD behaviors while maintaining a uniform environment, an experimental design not easily achieved in humans. Furthermore, we used a novel non-linear network-based clustering approach to characterize rats based on OUD vulnerability to assess genetic variants associated with OUD susceptibility. Our findings confirm the heritability of several OUD-like behaviors, including OUD susceptibility. Additionally, several genetic variants associated with nociceptive threshold prior to heroin experience, heroin consumption, escalation of intake, and motivation to obtain heroin were identified. *Tom1*, a microglial component, was implicated for nociception. Several genes involved in dopaminergic signaling, neuroplasticity and substance use disorders, including *Brwd1*, *Pcp4, Phb1l2* and *Mmp15* were implicated for the heroin traits. Additionally, an OUD vulnerable phenotype was associated with genetic variants for consumption and break point, suggesting a specific genetic contribution for OUD-like traits contributing to vulnerability. Together, these findings identify novel genetic markers related to the susceptibility to OUD-relevant behaviors in HS rats.

Introduction

Opioid use disorder (OUD) is characterized by compulsive opioid use and relapse propensity, despite efforts to maintain abstinence. OUD diagnosis has increased substantially in the past decade (1), affecting approximately 16 million people per year worldwide (2). The rise in diagnosis is not demographically constrained (1), but is prevalent in individuals across socio-economic class, age and sex (3), increasing the urgency to better understand the biological mechanisms contributing to OUD vulnerability. In twin studies, OUD shows high heritability (~50%) (4, 5), indicating that specific genetic variants contribute to biological basis of OUD susceptibility.

Genome-wide association studies (GWAS) are a useful tool for identifying the genetic variants that influence heritable traits contributing to OUD. GWAS of OUD, though limited, have already yielded valuable insights into genetic contributions to diagnostic vulnerability (6-13). However, human GWAS of OUD are limited to observational data and are thus confounded by numerous poorly defined environmental factors. Understandably, due to a lack of empirical data from study participants, human GWAS have largely focused on OUD diagnosis, which is comprised of poorly quantified amounts of drug-taking, extinction and seeking behaviors. Using rodent models of OUD, we can quantify specific traits (e.g., consumption, escalation, reinstatement) associated with addiction-related behaviors to directly interrogate the relationship between these metrics and genetic variants. By doing so, we can discover novel genetic variants that may contribute to the different phases of OUD, which can then be compared to data from human studies. Crucially, such genes may also provide targets for the development of novel OUD treatments.

In the current study, we assessed several measures of heroin-taking, extinction and seeking behaviors in outbred male and female heterogeneous stock (HS) rats at two different testing locations. Additionally, anxiety- and stress-like behaviors, as well as nociceptive threshold, were assessed prior to and following heroin experience. HS rats are a diverse outbred population that was created by interbreeding 8 inbred

strains at NIH in 1984 to capture genetic and behavioral heterogeneity reflective of the human population (14-16). Compared to inbred and commercially available outbred rat lines, HS rats exhibit comparatively greater diversity (16), and have been used in several studies assessing addiction-related behaviors (17-23), including those focusing on genetic analyses (24-26). In addition, HS rats are supported by complementary expression quantitative trait loci datasets (27), furthering the translational utility of our findings. In this study, we found genetic variants specific to different OUD behavioral traits, many of which had known involvement in neurobiological processes associated with SUD. Additionally, we employed a novel non-linear network-based clustering approach to identify subpopulations of OUD vulnerable and resilient rats (23) and thereby identify genetic variants associated with OUD susceptibility.

Methods

Behavioral testing occurred at two geographically distinct locations: Medical University of South Carolina (MUSC, USA) and the University of Camerino (UCAM, Italy). Two testing locations rendered the advantage of accounting for environmental differences on behavior, akin to the human experience, while maintaining standardized experimental procedures and equipment used between the sites. All experimental procedures were approved by the MUSC Institutional Animal Care and Use Committee and the Italian Ministry of Health (UCAM). All testing procedures were compliant with the National Institute of Health Guide for the Care and Use of Laboratory Animals and the Assessment and Accreditation of Laboratory Animals Care, as well as the European Community Council Directive for Care and Use of Laboratory Animals.

Subjects

Male and female HS rats (NMcwiWFsm #13673907, RRID:RGD13673907) bred at Wake Forest University were shipped to MUSC and UCAM in batches of 40 (20 females and 20 males/site) at 5 weeks of age. Each site had an equal representation of littermates thereby creating genetically comparable testing cohorts. Whenever possible, each shipment contained only one male and one female per litter, to reduce genetic relationships, which can reduce power in GWAS. Upon arrival, rats were left undisturbed for 3 weeks in a climate-controlled vivarium room with a standard 12-hr light/dark cycle prior to testing. Rats were pair-housed and received ad libitum access to food and water throughout the duration of testing. Behavioral tests for stress, anxiety and nociception occurred during the light cycle, whereas all heroin-associated testing occurred during the dark phase. A total of 1,160 rats underwent testing from February of 2019 through December of 2022, with the following animals excluded from all analyses: 147 for behavioral control studies (e.g., did not undergo heroin self-administration procedures), 107 due to death (post-surgical illness, n=84; surgery, n=23); 29 for behavioral data collection technical issues, and 3 for spleen

identification issues. Final behavioral and genetic analyses were conducted on 874 rats (MUSC, n=479; UCAM, n=395).

Drugs

Heroin hydrochloride was supplied to both sites by the National Institute on Drug Abuse (Bethesda, MD) and dissolved in 0.9% sterile saline.

Behavioral testing

The experimental timeline is shown in Figure 1a. Experimental testing occurred as previously described (17, 23). Extensive behavioral analyses for these animals, including sex and site differences, have been conducted and reported in prior publications (17, 23), and we refer the reader to these manuscripts for further edification. The focus of this manuscript is on identifying the genetic variants that contribute to individual differences in these behaviors. A brief description of behavioral testing procedures is provided below with greater detail in the Supplemental Methods.

Behavioral testing

Stress and anxiety-related behaviors were assessed in the elevated-plus maze (EPM) and the open field test (OFT). Nociception under baseline (administration of 1 mg/kg saline, sc) and test (0.75 mg/kg heroin, sc) conditions were evaluated using the tail flick (TF) test. All tests were performed prior to and following heroin self-administration (see Supplemental Methods and also Kuhn et al., 2022 and 2024).

Heroin taking, extinction and seeking measures

After EPM, OFT and TF tests, rats were outfitted with an indwelling jugular catheter prior to undergoing heroin self-administration training procedures. During training, presses on the active lever resulted in presentation of a tone and cue light stimulus, as well an infusion of heroin (20 µg/kg/100 µl infusion over 3-sec). Training occurred for 3 weeks with four 12-hr sessions/week. Next, motivation to take heroin was

assessed using a progressive ratio test whereupon the number of lever presses needed to earn an infusion exponentially increased according to the following formula: $(5*e^{0.2n})-5$ (28). The maximum number of lever presses exerted to receive an infusion was considered the break point. Heroin self-administration training was then re-established for 3 days, followed by a within-session extinction-prime test lasting 6 hours. With two hours left in the session, animals received a heroin prime injection (0.25 mg/kg, sc). Animals next underwent 6 daily 2-h extinction training (i.e., no cue presentation or heroin delivery) sessions prior to a 2-h test for cue-induced reinstatement (i.e., active lever presses resulted in cue presentation but no heroin infusion). The EPM, OFT and TF tests were then repeated. See Supplemental Methods for further details.

Tissue collection and genotyping

Following all behavioral testing, animals were sacrificed and spleen samples were collected and shipped to University of California San Diego for genotyping. Genotyping-by-sequencing was employed as previously described to assess genotypes (29, 30). A total of 7,955,513 single nucleotide polymorphisms (SNPs) were identified with an error rate <1%. However, prior to GWAS analysis, the SNPs were filtered for missingness not above 10%, minor allele frequency of no less than 0.5% and the HWE deviation p-value no less than $1e^{-10}$, resulting in a final set of 5,308,726 SNPs for analysis.

Statistical analyses

Behavioral traits assessed in GWAS: Behaviors included in GWAS analysis included behavioral testing traits (i.e., EPM, OFT and TF) both prior to (time point 1) and following (time point 2) heroin experience, as well as the difference in individual testing measures between the time points (Supplemental Table 1). Traits representative of heroin taking, extinction and seeking measures were also analyzed (Supplemental Table 1). Heroin-taking behaviors included: escalation of intake in the first hour of the session across training (ug/kg; average consumption hour 1 days 1-3 subtracted from hour 1 days 10-12; Escalation of intake:

Hour 1), escalation of intake across the entire session (ug/kg; average consumption days 1-3 subtracted from days 10-12; Escalation of intake: Hour 1-12), total consumption (ug/kg) across training sessions 1-12, and break point achieved during the progressive ratio test. Extinction traits include active lever presses made during the following: the first (Extinction: Burst) and middle (Extinction: Prime) 2 hours of the within-session extinction-heroin prime reinstatement test, the last day of extinction training prior to the test for cued reinstatement (Extinction: Day 6) and the difference in lever presses between the first and last day of extinction training (Extinction de-escalation). Active lever presses made during the tests for heroin prime and cued reinstatement were assessed for seeking measures.

Behavioral clustering: In addition to specific traits, we wanted to determine if any genetic variants were associated with OUD susceptibility. To do so, we employed a non-linear network-based clustering approach (31) with rats characterized into OUD vulnerable, intermediate or resilient clusters. Behaviors included for analysis are indicated in Supplemental Table 1. Behavioral and neurobiological validation of this model are extensively detailed in prior publications (23). Cluster assignment was used as phenotypes (Vulnerable vs intermediate/resilient; Intermediate vs vulnerable/resilient; Resilient vs vulnerable/intermediate) for GWAS. Differences in cluster composition between the top SNP allelic variants were assessed using a Chi-square test.

Phenotypic correlations: To address both sex and site differences in raw behavioral data (23) as well as non-normal distributions, all data were quantile normalized separately for site (MUSC and UCAM) and sex (female and male) and then combined. For each site, covariates that explained more than 2% of variance in each trait within the data (Supplemental Table 2 and 3) were regressed out and data was again quantile normalized to maintain a normal distribution. Next, data from both MUSC and UCAM were merged and quantile normalized to maintain normality. Behavioral co-variance between traits was assessed using Spearman correlations with Bonferroni adjustment to correct for multiple comparisons using GraphPad Prism version 10. SNP heritability for behavioral traits was determined using GCTA-GREML analysis.

Quantitative trait loci (QTL) and SNP identification: To perform the GWAS we used GCTA software employing a genetic relatedness matrix and the Leave One Chromosome Out method to account for proximal contamination (32, 33). Because all data was quantile normalized, only one significance threshold, which was determined by permutation, was necessary (34). We report QTLs that surpassed the permutation-derived threshold of $-\log_{10}$ p>5.58 (p<0.05). Significant QTLs are only reported when there was at least one additional SNP located within 0.5 Mb with a significance within 2 $-\log_{10}$(p) units of the first SNP (35). Independence of QTLs on the same chromosome was established by making the top SNP in the most significant QTL a covariate and then repeating the GWAS analysis for the chromosome in question. This process was repeated until no more significant QTLs were identified (35). Linkage disequilibrium (LD; $r^2 \geq 0.6$) was used to define the boundaries of each QTL. SNPEff was used to identify coding SNPs within each QTL interval (36). To identify associations from other GWAS that have been performed in HS rats that map to the same loci, a phenome wide association study (PheWAS) was conducted. To explore functional mechanisms whereby SNPs may affect gene expression patterns, expression QTL (eQTL) and splice QTL (sQTL) within a 3 Mb window that were obtained from RatGTEx.org (27).

Principle component analysis: To assess whether variance in behavioral traits within significant QTLs could be explained by common factors, a principle component analysis was applied to the data. Number of components was determined using a minimum threshold of Eigenvalue of 1 and Varimax method with a minimum factor loading of 0.3 was used to rotate identified components. Analyses used SPSS version 27.

Results

Behavioral correlations

Behavioral traits showed weak to moderate correlations overall (Fig. 1b, Supplemental Table 4). However, behavior during the progressive ratio test, cued reinstatement and total heroin consumption were positively correlated ($r^2 \geq 0.48$), suggesting overlapping behavioral mechanisms guiding these behaviors.

Heritability estimates

SNP heritability (h^2) estimates the proportion of the phenotypic variance explained by the genetic variance and is dependent on the genetic diversity of the population and environmental sources of variability. Aspects of heroin taking (Consumption, $h^2=0.17$) and the inability to initially refrain from seeking (Extinction: Burst, $h^2=0.10$) were heritable (Fig. 2a, Supplemental Table 5). While some measures of heroin taking, extinction and seeking were not heritable, OUD susceptibility cluster assignment (23) was heritable (Vulnerable vs resilient/intermediate, $h^2=0.17$; Intermediate vs vulnerable/resilient, $h^2=0.09$; Resilient vs vulnerable/intermediate, $h^2=0.15$), suggesting a heritable genetic contribution to OUD susceptibility (Fig. 2a, Supplemental Table 5).

Several behavioral measures prior to and following heroin administration experience showed significant heritability (Fig. 2a, Supplemental Table 5). Nociceptive threshold under basal conditions both prior to (TF baseline 1, $h^2=0.11$) and following (TF baseline 2, $h^2=0.15$) heroin experience were both heritable. Heroin-induced nociception after prolonged heroin exposure was also heritable (TF test 2, $h^2=0.12$), reflecting the genetic contribution to the long-term effects of heroin exposure on nociceptive threshold. Aspects of anxiety- and stress-like behaviors were also heritable, including high anxiety prior to heroin experience (EPM closed 1, $h^2=0.08$), but lower levels following heroin self-administration (EPM open 2, $h^2=0.07$). Our results support previous findings that locomotor behavior is heritable (24, 25) (OFT distance 1, $h^2=0.14$;

OFT time 1, $h^2=0.10$) and show that this heritability persists even after exposure to heroin administration (OFT distance 2, $h^2=0.23$; OFT time 2, $h^2=0.16$).

GWAS

We identified genome-wide significant associations for several traits: escalation of intake across the entire session on Chromosome 10, total heroin consumption on Chromosome 11, break point achieved during the progressive ratio test on Chromosome 19, and baseline nociception prior to heroin experience (TF baseline 1) on Chromosomes 2 and 19 (Fig. 2b). Shared behavioral variance between these traits was assessed using PCA, with heroin taking traits loading onto one factor and baseline nociception onto its own factor, reflecting the independence of these measures in our data (Supplemental Fig. 1). While both nociception and break point were associated with loci on Chromosome 19, the associated loci are separated by approximately 4 Mb and are not in strong LD with one another, indicating that they are due to separate causal loci.

Escalation of intake: We identified a significant locus on Chromosome 10 for escalation of heroin intake (Fig. 3). This locus did not contain any coding variants that are in strong LD with the top SNP. Though eQTLs and sQTLs were identified, none are in strong LD with the top SNP, suggesting they are unlikely to be causally related to the escalation of heroin intake behavior. Identified eQTLs include metabotropic glutamate receptor 6 (*Grm6*) in whole brain ($r^2=0.63$), collagen type XXIII alpha 1 chain (*Col23a1*) in the orbitofrontal cortex (OFC; $r^2=0.63$) and zinc finger protein (*Zfp2*) in the prelimbic cortex (PrL; $r^2=0.64$). Two sQTLs were also identified, jade family PHD finger 2 (*Jade2*) in the nucleus accumbens core (NAcc) and basolateral amygdala (BLA; both $r^2=0.63$), and *Col23a1* in the PrL ($r^2=0.63$). The lack of any putatively causal variants in this interval suggests that the causal variant remains unknown, possibly because there is an eQTL or sQTL in a tissue that is not represented in the RatGTEx database, or an eQTL that was not

detected, for example, because it is significant at a different developmental time point or under different environmental conditions.

Total consumption: Total heroin consumed across training was associated with a locus on Chromosome 11 (Fig. 4a). Several significant causal coding variants were identified that are in strong LD with this locus, including bromodomain and WD repeat domain containing 1 (*Brwd1*: Leu2187Pro, r^2=0.99), Purkinje cell protein 4 (*Pcp4*: Lys12Arg, r^2=0.99) and two different amino acid coding variants for SH3 domain binding glutamate-rich protein (*Sh3bgr:* Glu169Lys, r^2=0.99; Glu177Lys, r^2=0.99). There were also eQTLs for *Sh3bgr* that was in high LD with this locus (r^2=0.99 for all) in whole brain, BLA, NAcc, PrL, lateral habenula (LHb) and OFC. Two additional genes with eQTLs in high LD with the locus were identified: *Pcp4* (PrL, r^2=0.88) and a long non-coding RNA (*ENSRNOG00000068065*; infralimbic cortex (IL), r^2=0.99). We also identified an sQTL for *ENSRNOG00000068065* in several brain regions (whole brain, BLA, LHb, OFC, NAcc, PrL, and PrL2; r^2=0.99-1.00). Lastly, there was an sQTL for the gene guided entry of tail-anchored proteins factor 1 (*Get1*; r^2=0.99) in the IL. Heroin consumption is likely mediated, in part, by these coding differences and or by heritable differences in expression of one or more of these genes. Because several plausible variants are in LD with one another, mechanistic or other complementary approaches will be needed to identify the causal gene. Additionally, PheWAS analysis showed that OUD vulnerability (Vulnerable vs Intermediate/resilient cluster) was associated with the top SNP at a suggestive level (-log(p)=4.5), with cluster composition differing between the allelic variants (x^2=18.12, p<0.0001; Fig. 4b,c).

Break point: A significant association on Chromosome 19 was observed for progressive ratio break point (Fig. 5a). Two potentially causal coding variants in prohibitin 1 like 2 (*Phb1l2*; Lys70Asn and Ile231Asn), which were in perfect LD with the top SNP (r^2=1.0), were identified. An eQTL with perfect LD with the locus for the gene matrix metallopeptidase 15 (*Mmp15*; r^2=1.0) was also identified in the whole brain. An additional eQTL (solute carrier family 38 member 7, *Slc38a7*) and sQTL (casein kinase 1 alpha 2, *Csnk2a2*) were identified in the BLA, however, the relationship between these genes and the locus are modest

($r^2=0.65$ for both) rendering them much less likely to be the causes of the association with break point behavior. Similar to the Chromosome 11 association for total heroin consumption, PheWAS analysis showed a suggestive association between the top SNP and the OUD vulnerable cluster (Vulnerable vs intermediate/resilient clusters; $-\log(p)=5.16$; Fig. 5b). OUD cluster composition differed between the alleles at the peak SNP ($x^2=18.91$, $p<0.0001$; Fig. 5c). While there is a small bias toward the AA variant conferring OUD resiliency, the minor variant, AC, strongly coincides with OUD vulnerability.

Baseline nociception prior to heroin experience: A locus on Chromosome 2 was identified for basal nociception prior to heroin taking procedures (Fig 6a). Several coding variants in modest LD with the causal SNP were identified in genes within this locus, including coiled-coil domain containing 152 (*Ccdc152*: Ala242Ser and Thr14Asn, both $r^2=0.77$), growth hormone receptor (*Ghr*: Ala546Val and Lys541Asn, both $r^2=0.77$), zinc finger protein 131 (*Zfp131*: Gly506Ser, $r^2=0.82$) and serine/threonine-protein kinase NIM1 (*Nim1k;* $r^2=0.64$). The gene selenoprotein P (*Selenop*) had an sQTL in both BLA and whole brain ($r^2=0.78$-0.79), and an eQTL for both whole brain and the NAcc (both $r^2=0.66$). These data provide some support of the involvement of these genes in the Chromosome 2 association with baseline nociception.

An additional significant association for this behavior was identified on Chromosome 19 (Fig. 6b). A potentially causal coding variant in modest LD with the causal SNP in the gene LARGE xylosyl- and glucuronyltransferase 1 *(Large1*: Ala92Val, $r^2=0.75$) was present. There were also several eQTLs for *Large1* (whole brain, $r^2=0.69$; BLA, $r^2=0.78$; IL, $r^2=0.68$; NAcc, $r^2=0.70$; OFC, $r^2=0.69$; PrL, $r^2=0.70$; and PrL2, $r^2=0.75$). In addition, there was an eQTL for ribosomal protein S18 (*Rps18l2*) within the LHb ($r^2=0.98$). An sQTL for the gene myb1 trafficking protein (*Tom1*) was in strong LD with this locus ($r^2=1.0$) within the PrL. Finally, there was a significant sQTL for metallothionein 3 (*Mt3*) in the BLA ($r^2=0.75$). These coding variants, eQTLs and sQTLs could point to heritable differences and underlying biological mechanisms for basal nociception prior to drug experience at the Chromosome 19 locus.

Discussion

The economic toll on society for OUD totals over one trillion dollars annually (37). Furthermore, over 1 million individuals seek treatment for OUD annually (38) and more than 80,000 people die from opioid overdoses every year in the US alone (39). Thus, there is an urgent need to understand the factors that influence vulnerability to OUD. Using a rat line emulating within species genetic and phenotypic variability (14) akin to humans, we assessed the genetic basis of several OUD-like behaviors and further extended analysis to account for OUD susceptibility. While heroin was used, we expect these findings will be relevant for other opioids of abuse whose primary site of action is mu opioid receptors. We found several traits related to OUD are heritable, including overall OUD susceptibility. Genetic variants associated with nociception prior to heroin experience as well as heroin consummatory behaviors were identified, several of which are either biomarkers for other SUDs or are involved in neuroplasticity. Using this model, we are able to identify genetic variants possibly associated with an OUD diagnosis, but also specific behavioral traits that comprise vulnerability or resiliency thereby further contributing to our knowledge of the genetic basis of OUD.

Heritability

OUD is heritable in humans and we found several OUD-like behaviors were also heritable in HS rats. Human twin studies have yielded heritability estimates between 23-54% for OUD (4, 5) and human GWAS of OUD have produced SNP heritability estimates, which are expected to be lower, ranging from 6-12.7% (9-13). However, given the complex and multi-symptomatic nature of OUD (40), assessing heritability of distinct behavioral traits associated with human OUD vulnerability is difficult. In a large sample of HS rats, we identified several heritable heroin addiction-like traits that model different aspects of human OUD. Both heroin consumption and initial extinction behavior were significantly heritable, with SNP heritability estimates for these traits similar to SNP heritability obtained from human studies of OUD. However,

behaviors following prolonged heroin use and forced abstinence, such as cued reinstatement, did not exhibit significant heritability, suggesting weaker genetics influences and/or limitations in our procedure's ability to detect the cascade of neuroplastic adaptations that occur following prolonged heroin self-administration (41, 42).

Novelty-induced locomotion, a heritable behavior (24, 25), has been associated with vulnerability in opioid addiction-related behaviors in both rodent models (17, 43-47) and humans (48). We confirmed the heritability of locomotor behavior both prior to and following heroin experience in HS rats, suggesting an enduring genetic contribution to this behavior that remains relatively unaffected by factors (e.g., epigenetic changes) associated with prolonged heroin experience. Lastly, in agreement with prior work in humans, we showed that nociceptive threshold to heat stimuli under baseline conditions is heritable (49). Following opioid use, both humans (50, 51) and rats (52) demonstrate opioid-induced hyperalgesia in response to a painful stimuli (17), a finding replicated in our study. However, the heritability of nociception in humans is unknown. Given the inherent behavioral and genetic heterogeneity present in HS rats, and the translational relevance of the findings reported here, HS rats offer an opportunity to disentangle the genetic factors contributing to opioid-induced nociception.

GWAS

Genetic variants associated with OUD-like behaviors in rats can contribute novel insights that improve our understanding of the neurobiological mechanisms underlying distinct behavioral traits associated with OUD in humans. This study implicated several loci and genes within those loci that had coding variants, eQTLs and/or sQTLs.

Escalation of heroin intake across the entire session: While no causal coding variants, eQTLs or sQTLs were in strong LD with the QTL on Chromosome 10 for escalation of heroin-taking behavior across the entire session, several genes in this interval are of potential interest. For example, *Adamts2*, an extracellular

matrix proteinase (53) localized within the mesolimbic and mesocortical regions within the brain (54), is a biomarker for heavy tobacco smoking and is downregulated following long-term morphine exposure and withdrawal (55). Similarly, the gene heterogeneous nuclear ribonucleoprotein H1 (*Hnrnph1*) is located in this interval. *Hnrnph1* was identified as the causal gene for a mouse QTL for methamphetamine response (56-59) and subsequent work has shown it also modulates the response to fentanyl (60) and alcohol (61). Further investigation is required to determine if there are overlooked coding variants or eQTLs for *Adamts2* or *Hnrnph1*.

Heroin consumption: The region on Chromosome 11 that was associated with total consumption of heroin across all training sessions contained a number of genes with coding and expression variants. Three genes harbored coding variants that were in strong LD with the peak SNP: *Brwd1*, *Pcp4* and *Sh3bgr*. In humans, *BRWD1* has been associated with potentially relevant traits including cigarette smoking (62-64), externalizing behaviors (65) and ADHD (66). *PCP4* mediates calcium signaling and through downstream mechanisms also affects dopamine and acetylcholine signaling [76]. *PCP4* levels are attenuated in post-mortem tissue of individuals diagnosed with alcohol use disorder [77] as well as in the NAc of rats following 2-weeks of nicotine withdrawal [78]. *PCP4* has also been linked to cigarette smoking (62). We identified an eQTL for *Pcp4* in the PrL, a critical component of drug-reward circuitry and plasticity (67, 68). The relationship between *PCP4* and OUD is unknown but our results suggest it may act to alter cortical to NAc transmission as a consequence of sustained heroin use. Finally, *Sh3bgr* had two coding variants and numerous eQTLs in whole brain, BLA, NAcc, OFC, LHb, and PrL, however the homologous human gene has not been associated with any psychiatric traits in prior human GWAS. In addition to these genes, a long non-coding RNA (ENSRNOG00000068065) and the gene *Get1* had sQTLs, but there is no supporting literature for their role in OUD-relevant behaviors.

Ets2, which contained the peak SNP for this locus, did not identify any coding variants or an e/sQTL. *Ets2* encodes a ubiquitous transcription factor that is found in many cells, and has be linked to molecular

processes (69) and cell types (70) that modulate synaptic remodeling, including those associated with OUD (71). Our results suggest *Ets2* and its allelic variants may impact OUD susceptibility, though further investigation is necessary.

While the functional association between heroin consumption and the other identified e/sQTLs are not known, *DSCAM* is correlated with the top SNP and encodes a neuronal cell adhesion molecule that regulates cellular connectivity (72) and glutamatergic plasticity associated with learning (73, 74). Glutamate-mediated neuroplasticity occurs following heroin experience, and is considered a driving force behind relapse vulnerability (42). *DSCAM* is necessary for tasks requiring memory formation (75, 76), including those associated with rewarded behaviors (76), and was identified in human GWAS as a biomarker for cigarette cessation success (77). Based on these data, *DSCAM* warrants further investigation for OUD.

Break point: The QTL on Chromosome 19 for break point achieved during the progressive ratio test contained two coding variants in strong LD with the causal variant for the poorly characterized gene *Phb1l2*. *Phb1l2* is localized to both neuronal and non-neural cells types in the central nervous system and helps to maintain mitochondria respiration by attenuating reactive oxygen species which contribute to cellular dysfunction (78). Chronic morphine administration disrupts mitochondrial respiration and restoring proper functioning attenuates opioid withdrawal (79). Additionally, *Phb1*, an analogue of *Phb1l2*, has been implicated in mediating glutamatergic and dopaminergic neural transmission and becomes dysregulated in several disorders, including alcohol use disorder (78). It is thus possible that differences in the coding sequence of *Phb1l2* influence break point through glutamate and dopamine dependent processes. Additionally, an eQTL for *Mmp15* in whole brain that was in strong LD with the causal SNP was identified, suggesting *Mmp15* expression could modulate break point. MMP15 activates MMP2 (80) which contributes to NAcc cell-specific regulation of extinction behavior following heroin self-administration (81). It is possible that attenuated *Mmp15*-induced *Mmp2* expression contributes to higher

break point levels. Furthermore, PheWAS analysis identified this locus as being associated with OUD vulnerability with the coding variant likely impacting OUD susceptibility.

Baseline nociceptive threshold prior to heroin experience: Genetic variants on two separate chromosomes were found for nociceptive threshold under baseline conditions, prior to heroin experience. On Chromosome 19, *Large1* was notable because it contained both putatively causal coding variants and an eQTL. Large1 has been associated with various behavioral traits in humans, including educational attainment (82), smoking (62) and neuroticism (83), however, no prior evidence that we are aware of link it to nociception. Heritable differences in splicing patterns (sQTLs) for both *Tom1* in the PrL and *Mt3* in the BLA are also candidates for the nociception, though neither has been previously directly linked to pain processing. *Mt3* is a brain zinc binding protein involved in several metabolic processes, and has been implicated in cellular dysfunction of diseases including Alzheimer and Huntington disease (84). *Tom1* is an integral component of microglial function (85), which is involved in neuropathic pain (86), but has yet to be implicated in acute painful stimuli.

On chromosome 2, two genes that operate in tandem with one another were also associated with nociception. There was modest LD between the top SNP for this QTL and splicing patterns in *Selenop* which encodes for a selenoprotein. This protein is secreted by the liver and acts to deliver selenium, a trace element necessary for proper brain function (87), via the bloodstream to regions including the brain. Selenium has been implicated in mediating dopamine release and transmission within the substantia nigra (88, 89) and the NAc (90). The presence of *Selenop* on NAc dopamine terminals is proposed to attenuate dopamine release into the synaptic cleft via direct mediation of the dopamine 2 autoreceptors, protecting against neurotoxic effects of drugs of abuse such as methamphetamine (90). *Ccdc152*, which also had coding variants that were in modest LD with this QTL, is encoded within the *Selenop* gene, and acts to attenuate *Selenop* levels via post-translational modification (91). Though the periaqueductal gray is the

predominant brain nuclei processing pain signaling (92), striatal dopamine signaling has recently been implicated in affecting pain processing and perception (93, 94) and warrants further investigation.

Conclusion and Future Directions

We used a rodent model to identify several heritable traits and genetic variants associated with OUD. Unlike GWAS performed in humans, we were able to identify genetic variants associated with distinct OUD behaviors that contribute toward diagnosis. Several of the identified genes are supported by prior relationships to SUDs and other relevant behavioral and psychiatric traits, suggesting that they warrant further investigation for OUD. Furthermore, using an established rodent model that characterizes rats using a non-linear compilation of behaviors across the SUD phases to confer OUD susceptibility, we identified genetic variants for heroin consumption and motivation that were associated specifically with OUD vulnerability. These findings strengthen the translational utility of the behavioral model, and can help to assess genetic variants and biomarkers associated with OUD vulnerability versus resiliency. However, complementing human OUD (5), phenotypic variance can only be partially explained by the genetic variants discovered here. Other genetic variants as well as epigenetic (95) and uncontrolled environmental factors also contribute to OUD behaviors is HS rats. Furthermore, both preclinical and clinical GWAS analysis focusing on genetic variants associated with OUD resiliency, a neurobiological active process (96, 97), that should be considered in future analyses.

Acknowledgements

The authors would like to thank the numerous technicians within the Kalivas and Ciccocioppo lab for their help performing the behavioral assays on the rats used in these analyses, as well as members of the Kalivas lab for their helpful feedback on the manuscript. This work was supported by the National Institute on Drug Abuse U01DA45300 (PWK), P50DA037844 (AAP), T32DA007288 (BNK) and K99DA057390 (BNK).

Conflict of Interest

The authors declare no conflict of interest.

References

1. Wide-ranging online data for epidemiologic research (WONDER) [Internet]. 2020. Available from: http://wonder.cdc.gov/.

2. Chang HY, Kharrazi H, Bodycombe D, Weiner JP, Alexander GC. Healthcare costs and utilization associated with high-risk prescription opioid use: a retrospective cohort study. BMC Med. 2018;16(1):69.

3. Jones CM, Logan J, Gladden RM, Bohm MK. Vital Signs: Demographic and Substance Use Trends Among Heroin Users - United States, 2002-2013. MMWR Morb Mortal Wkly Rep. 2015;64(26):719-25.

4. Berrettini W. A brief review of the genetics and pharmacogenetics of opioid use disorders. Dialogues Clin Neurosci. 2017;19(3):229-36.

5. Crist RC, Reiner BC, Berrettini WH. A review of opioid addiction genetics. Curr Opin Psychol. 2019;27:31-5.

6. Cheng Z, Zhou H, Sherva R, Farrer LA, Kranzler HR, Gelernter J. Genome-wide Association Study Identifies a Regulatory Variant of RGMA Associated With Opioid Dependence in European Americans. Biol Psychiatry. 2018;84(10):762-70.

7. Gelernter J, Kranzler HR, Sherva R, Koesterer R, Almasy L, Zhao H, et al. Genome-wide association study of opioid dependence: multiple associations mapped to calcium and potassium pathways. Biol Psychiatry. 2014;76(1):66-74.

8. Nelson EC, Agrawal A, Heath AC, Bogdan R, Sherva R, Zhang B, et al. Evidence of CNIH3 involvement in opioid dependence. Mol Psychiatry. 2016;21(5):608-14.

9. Sanchez-Roige S, Fontanillas P, Jennings MV, Bianchi SB, Huang Y, Hatoum AS, et al. Genome-wide association study of problematic opioid prescription use in 132,113 23andMe research participants of European ancestry. Mol Psychiatry. 2021;26(11):6209-17.

10. Zhou H, Rentsch CT, Cheng Z, Kember RL, Nunez YZ, Sherva RM, et al. Association of OPRM1 Functional Coding Variant With Opioid Use Disorder: A Genome-Wide Association Study. JAMA Psychiatry. 2020;77(10):1072-80.

11. Deak JD, Zhou H, Galimberti M, Levey DF, Wendt FR, Sanchez-Roige S, et al. Genome-wide association study in individuals of European and African ancestry and multi-trait analysis of opioid use disorder identifies 19 independent genome-wide significant risk loci. Mol Psychiatry. 2022;27(10):3970-9.

12. Polimanti R, Walters RK, Johnson EC, McClintick JN, Adkins AE, Adkins DE, et al. Leveraging genome-wide data to investigate differences between opioid use vs. opioid dependence in 41,176 individuals from the Psychiatric Genomics Consortium. Mol Psychiatry. 2020;25(8):1673-87.

13. Song W, Kossowsky J, Torous J, Chen CY, Huang H, Mukamal KJ, et al. Genome-wide association analysis of opioid use disorder: A novel approach using clinical data. Drug Alcohol Depend. 2020;217:108276.

14. Consortium S, Saar K, Beck A, Bihoreau MT, Birney E, Brocklebank D, et al. SNP and haplotype mapping for genetic analysis in the rat. Nat Genet. 2008;40(5):560-6.

15. Johannesson M, Lopez-Aumatell R, Stridh P, Diez M, Tuncel J, Blazquez G, et al. A resource for the simultaneous high-resolution mapping of multiple quantitative trait loci in rats: the NIH heterogeneous stock. Genome Res. 2009;19(1):150-8.

16. Hansen C, Spuhler K. Development of the National Institutes of Health genetically heterogeneous rat stock. Alcohol Clin Exp Res. 1984;8(5):477-9.

17. Kuhn BN, Cannella N, Crow AD, Roberts AT, Lunerti V, Allen C, et al. Novelty-induced locomotor behavior predicts heroin addiction vulnerability in male, but not female, rats. Psychopharmacology (Berl). 2022;239(11):3605-20.

18.   Kallupi M, Carrette LLG, Kononoff J, Solberg Woods LC, Palmer AA, Schweitzer P, et al. Nociceptin attenuates the escalation of oxycodone self-administration by normalizing CeA-GABA transmission in highly addicted rats. Proc Natl Acad Sci U S A. 2020;117(4):2140-8.

19.   Deal A, Cooper N, Kirse HA, Uneri A, Raab-Graham K, Weiner JL, et al. Early life stress induces hyperactivity but not increased anxiety-like behavior or ethanol drinking in outbred heterogeneous stock rats. Alcohol. 2021;91:41-51.

20.   Wang T, Han W, Chitre AS, Polesskaya O, Solberg Woods LC, Palmer AA, et al. Social and anxiety-like behaviors contribute to nicotine self-administration in adolescent outbred rats. Sci Rep. 2018;8(1):18069.

21.   King CP, Tripi JA, Hughson AR, Horvath AP, Lamparelli AC, Holl KL, et al. Sensitivity to food and cocaine cues are independent traits in a large sample of heterogeneous stock rats. Sci Rep. 2021;11(1):2223.

22.   Cannella N, Tambalo S, Lunerti V, Scuppa G, de Vivo L, Abdulmalek S, et al. Long-access heroin self-administration induces region specific reduction of grey matter volume and microglia reactivity in the rat. Brain Behav Immun. 2024.

23.   Kuhn BN, Cannella N, Crow AD, Lunerti V, Gupta A, Walterhouse SJ, et al. A multi-symptomatic model of heroin use disorder in rats reveals distinct behavioral profiles and neuronal correlates of heroin vulnerability versus resiliency. American Journal of Psychiatry (in press).

24.   Chitre AS, Hebda-Bauer EK, Blandino P, Bimschleger H, Nguyen KM, Maras P, et al. Genome-wide association study in a rat model of temperament identifies multiple loci for exploratory locomotion and anxiety-like traits. Front Genet. 2022;13:1003074.

25.   Gunturkun MH, Wang T, Chitre AS, Garcia Martinez A, Holl K, St Pierre C, et al. Genome-Wide Association Study on Three Behaviors Tested in an Open Field in Heterogeneous Stock Rats Identifies Multiple Loci Implicated in Psychiatric Disorders. Front Psychiatry. 2022;13:790566.

26.   Zhou JL, de Guglielmo G, Ho AJ, Kallupi M, Pokhrel N, Li HR, et al. Single-nucleus genomics in outbred rats with divergent cocaine addiction-like behaviors reveals changes in gene amygdala GABAergic inhibition. Nat Neurosci. 2023.

27.   Munro D, Wang T, Chitre AS, Polesskaya O, Ehsan N, Gao J, et al. The regulatory landscape of multiple brain regions in outbred heterogeneous stock rats. Nucleic Acids Res. 2022;50(19):10882-95.

28.   Richardson NR, Roberts DC. Progressive ratio schedules in drug self-administration studies in rats: a method to evaluate reinforcing efficacy. J Neurosci Methods. 1996;66(1):1-11.

29.   Parker CC, Gopalakrishnan S, Carbonetto P, Gonzales NM, Leung E, Park YJ, et al. Genome-wide association study of behavioral, physiological and gene expression traits in outbred CFW mice. Nat Genet. 2016;48(8):919-26.

30.   Gileta AF, Gao J, Chitre AS, Bimschleger HV, St Pierre CL, Gopalakrishnan S, et al. Adapting Genotyping-by-Sequencing and Variant Calling for Heterogeneous Stock Rats. G3 (Bethesda). 2020;10(7):2195-205.

31.   Allen C, Kuhn BN, Cannella N, Crow AD, Roberts AT, Lunerti V, et al. Network-Based Discovery of Opioid Use Vulnerability in Rats Using the Bayesian Stochastic Block Model. Front Psychiatry. 2021;12:745468.

32.   Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. Am J Hum Genet. 2011;88(1):76-82.

33.   Cheng R, Parker CC, Abney M, Palmer AA. Practical considerations regarding the use of genotype and pedigree data to model relatedness in the context of genome-wide association studies. G3 (Bethesda). 2013;3(10):1861-7.

34.   Cheng R, Palmer AA. A simulation study of permutation, bootstrap, and gene dropping for assessing statistical significance in the case of unequal relatedness. Genetics. 2013;193(3):1015-8.

35.   Chitre AS, Polesskaya O, Holl K, Gao J, Cheng R, Bimschleger H, et al. Genome-Wide Association Study in 3,173 Outbred Rats Identifies Multiple Loci for Body Weight, Adiposity, and Fasting Glucose. Obesity (Silver Spring). 2020;28(10):1964-73.

36.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, et al. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin). 2012;6(2):80-92.

37.   Florence C, Luo F, Rice K. The economic burden of opioid use disorder and fatal opioid overdose in the United States, 2017. Drug Alcohol Depend. 2021;218:108350.

38.   National Survey of Substance Abuse Treatment Services (N-SSATS): 2020 [Internet]. Substabce Abuse and Mental Health Services Administration. 2021. Available from: https://www.samhsa.gov/data/sites/default/files/reports/rpt35313/2020_NSSATS_FINAL.pdf.

39.   Drug Overdose death rates [Internet]. National Institute on Drug Abuse. 2023. Available from: https://nida.nih.gov/research-topics/trends-statistics/overdose-death-rates.

40.   Association AP. Diagnostic and statistical manual of mental disorders. 5 ed2013.

41.   Mazei-Robison MS, Nestler EJ. Opiate-induced molecular and cellular plasticity of ventral tegmental area and locus coeruleus catecholamine neurons. Cold Spring Harb Perspect Med. 2012;2(7):a012070.

42.   Kruyer A, Chioma VC, Kalivas PW. The Opioid-Addicted Tetrapartite Synapse. Biol Psychiatry. 2020;87(1):34-43.

43.   Ambrosio E, Goldberg SR, Elmer GI. Behavior genetic investigation of the relationship between spontaneous locomotor activity and the acquisition of morphine self-administration behavior. Behav Pharmacol. 1995;6(3):229-37.

44.   Lamarque S, Taghzouti K, Simon H. Chronic treatment with Delta(9)-tetrahydrocannabinol enhances the locomotor response to amphetamine and heroin. Implications for vulnerability to drug addiction. Neuropharmacology. 2001;41(1):118-29.

45.   Xigeng Z, Yonghui L, Xiaojing L, Lin X, Dongmei W, Jie L, et al. Social crowding sensitizes high-responding rats to psychomotor-stimulant effects of morphine. Pharmacol Biochem Behav. 2004;79(2):213-8.

46.   Swain Y, Muelken P, LeSage MG, Gewirtz JC, Harris AC. Locomotor activity does not predict individual differences in morphine self-administration in rats. Pharmacol Biochem Behav. 2018;166:48-56.

47.   Chang SE, Krueger LD, Flagel SB. Investigating individual differences in opioid-taking and opioid-seeking behavior in male rats. Psychopharmacology (Berl). 2022.

48.   Wingo T, Nesil T, Choi JS, Li MD. Novelty Seeking and Drug Addiction in Humans and Animals: From Behavior to Molecules. J Neuroimmune Pharmacol. 2016;11(3):456-70.

49.   Angst MS, Phillips NG, Drover DR, Tingle M, Ray A, Swan GE, et al. Pain sensitivity and opioid analgesia: a pharmacogenomic twin study. Pain. 2012;153(7):1397-409.

50.   Ren ZY, Shi J, Epstein DH, Wang J, Lu L. Abnormal pain response in pain-sensitive opiate addicts after prolonged abstinence predicts increased drug craving. Psychopharmacology (Berl). 2009;204(3):423-9.

51.   Carcoba LM, Contreras AE, Cepeda-Benito A, Meagher MW. Negative affect heightens opiate withdrawal-induced hyperalgesia in heroin dependent individuals. J Addict Dis. 2011;30(3):258-70.

52.   Marchette RCN, Gregory-Flores A, Tunstall BJ, Carlson ER, Jackson SN, Sulima A, et al. kappa-Opioid receptor antagonism reverses heroin withdrawal-induced hyperalgesia in male and female rats. Neurobiol Stress. 2021;14:100325.

53.   Shiomi T, Lemaitre V, D'Armiento J, Okada Y. Matrix metalloproteinases, a disintegrin and metalloproteinases, and a disintegrin and metalloproteinases with thrombospondin motifs in non-neoplastic diseases. Pathol Int. 2010;60(7):477-96.

54.   Ruso-Julve F, Pombero A, Pilar-Cuellar F, Garcia-Diaz N, Garcia-Lopez R, Juncal-Ruiz M, et al. Dopaminergic control of ADAMTS2 expression through cAMP/CREB and ERK: molecular effects of antipsychotics. Transl Psychiatry. 2019;9(1):306.

55.   Thiagarajan SK, Mok SY, Ogawa S, Parhar IS, Tang PY. Receptor-Mediated AKT/PI3K Signalling and Behavioural Alterations in Zebrafish Larvae Reveal Association between Schizophrenia and Opioid Use Disorder. Int J Mol Sci. 2022;23(9).

56.   Yazdani N, Parker CC, Shen Y, Reed ER, Guido MA, Kole LA, et al. Hnrnph1 Is A Quantitative Trait Gene for Methamphetamine Sensitivity. PLoS Genet. 2015;11(12):e1005713.

57.   Ruan QT, Yazdani N, Blum BC, Beierle JA, Lin W, Coelho MA, et al. A Mutation in Hnrnph1 That Decreases Methamphetamine-Induced Reinforcement, Reward, and Dopamine Release and Increases Synaptosomal hnRNP H and Mitochondrial Proteins. J Neurosci. 2020;40(1):107-30.

58.   Ruan QT, Yazdani N, Reed ER, Beierle JA, Peterson LP, Luttik KP, et al. 5' UTR variants in the quantitative trait gene Hnrnph1 support reduced 5' UTR usage and hnRNP H protein as a molecular mechanism underlying reduced methamphetamine sensitivity. FASEB J. 2020;34(7):9223-44.

59.   Borrelli KN, Langan CR, Dubinsky KR, Szumlinski KK, Carlezon WA, Jr., Chartoff EH, et al. Intracranial self-stimulation and concomitant behaviors following systemic methamphetamine administration in Hnrnph1 mutant mice. Psychopharmacology (Berl). 2021;238(7):2031-41.

60.   Bryant CD, Healy AF, Ruan QT, Coehlo MA, Lustig E, Yazdani N, et al. Sex-dependent effects of an Hnrnph1 mutation on fentanyl addiction-relevant behaviors but not antinociception in mice. Genes Brain Behav. 2021;20(3):e12711.

61.   Fultz EK, Coelho MA, Lieberman D, Jimenez-Chavez CL, Bryant CD, Szumlinski KK. Hnrnph1 is a novel regulator of alcohol reward. Drug Alcohol Depend. 2021;220:108518.

62.   Saunders GRB, Wang X, Chen F, Jang SK, Liu M, Wang C, et al. Genetic diversity fuels gene discovery for tobacco and alcohol use. Nature. 2022;612(7941):720-4.

63.   Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. Nat Genet. 2019;51(2):237-44.

64.   Xu K, Li B, McGinnis KA, Vickers-Smith R, Dao C, Sun N, et al. Genome-wide association study of smoking trajectory and meta-analysis of smoking status in 842,000 individuals. Nat Commun. 2020;11(1):5302.

65.   Karlsson Linner R, Mallard TT, Barr PB, Sanchez-Roige S, Madole JW, Driver MN, et al. Multivariate analysis of 1.5 million people identifies genetic associations with traits related to self-regulation and addiction. Nat Neurosci. 2021;24(10):1367-76.

66.   Rao S, Baranova A, Yao Y, Wang J, Zhang F. Genetic Relationships between Attention-Deficit/Hyperactivity Disorder, Autism Spectrum Disorder, and Intelligence. Neuropsychobiology. 2022;81(6):484-96.

67.   Hearing M. Prefrontal-accumbens opioid plasticity: Implications for relapse and dependence. Pharmacol Res. 2019;139:158-65.

68.   Kalivas PW, Volkow ND. The neural basis of addiction: a pathology of motivation and choice. Am J Psychiatry. 2005;162(8):1403-13.

69.   Trojanowska M. Ets factors and regulation of the extracellular matrix. Oncogene. 2000;19(55):6464-71.

70.   Cornell J, Salinas S, Huang HY, Zhou M. Microglia regulation of synaptic plasticity and learning and memory. Neural Regen Res. 2022;17(4):705-16.

71.   Green JM, Sundman MH, Chou YH. Opioid-induced microglia reactivity modulates opioid reward, analgesia, and behavior. Neurosci Biobehav Rev. 2022;135:104544.

72.   Ronn LC, Berezin V, Bock E. The neural cell adhesion molecule in synaptic plasticity and ageing. Int J Dev Neurosci. 2000;18(2-3):193-9.

73.   Li HL, Huang BS, Vishwasrao H, Sutedja N, Chen W, Jin I, et al. Dscam mediates remodeling of glutamate receptors in Aplysia during de novo and learning-related synapse formation. Neuron. 2009;61(4):527-40.

74.   Stachowicz K, Panczyszyn-Trzewik P, Sowa-Kucma M, Misztak P. Changes in working memory induced by lipopolysaccharide administration in mice are associated with metabotropic glutamate receptors 5 and contrast with changes induced by cyclooxygenase-2: Involvement of postsynaptic density protein 95 and down syndrome cell adhesion molecule. Neuropeptides. 2023;100:102347.

75.   Chen BE, Kondo M, Garnier A, Watson FL, Puettmann-Holgado R, Lamar DR, et al. The molecular diversity of Dscam is functionally required for neuronal wiring specificity in Drosophila. Cell. 2006;125(3):607-20.

76.   Keene AC, Krashes MJ, Leung B, Bernard JA, Waddell S. Drosophila dorsal paired medial neurons provide a general mechanism for memory consolidation. Curr Biol. 2006;16(15):1524-30.

77.   Uhl GR, Liu QR, Drgon T, Johnson C, Walther D, Rose JE, et al. Molecular genetics of successful smoking cessation: convergent genome-wide association study results. Arch Gen Psychiatry. 2008;65(6):683-93.

78.   Bernstein HG, Smalla KH, Keilhoff G, Dobrowolny H, Kreutz MR, Steiner J. The many "Neurofaces" of Prohibitins 1 and 2: Crucial for the healthy brain, dysregulated in numerous brain disorders. J Chem Neuroanat. 2023;132:102321.

79.   Kowalczyk P, Sulejczak D, Kleczkowska P, Bukowska-Osko I, Kucia M, Popiel M, et al. Mitochondrial Oxidative Stress-A Causative Factor and Therapeutic Target in Many Diseases. Int J Mol Sci. 2021;22(24).

80.   Wright JW, Harding JW. Contributions of matrix metalloproteinases to neural plasticity, habituation, associative learning and drug addiction. Neural Plast. 2009;2009:579382.

81.   Chioma VC, Kruyer A, Bobadilla AC, Angelis A, Ellison Z, Hodebourg R, et al. Heroin Seeking and Extinction From Seeking Activate Matrix Metalloproteinases at Synapses on Distinct Subpopulations of Accumbens Cells. Biol Psychiatry. 2021;89(10):947-58.

82.   Okbay A, Wu Y, Wang N, Jayashankar H, Bennett M, Nehzati SM, et al. Polygenic prediction of educational attainment within and between families from genome-wide association analyses in 3 million individuals. Nat Genet. 2022;54(4):437-49.

83.   Baselmans BML, Jansen R, Ip HF, van Dongen J, Abdellaoui A, van de Weijer MP, et al. Multivariate genome-wide analyses of the well-being spectrum. Nat Genet. 2019;51(3):445-51.

84.   Koh JY, Lee SJ. Metallothionein-3 as a multifunctional player in the control of cellular processes and diseases. Mol Brain. 2020;13(1):116.

85.   Wang H, Wang X, Shen Y, Wang Y, Yang T, Sun J, et al. SENP1 modulates chronic intermittent hypoxia-induced inflammation of microglia and neuronal injury by inhibiting TOM1 pathway. Int Immunopharmacol. 2023;119:110230.

86.   Inoue K, Tsuda M. Microglia in neuropathic pain: cellular and molecular mechanisms and therapeutic potential. Nat Rev Neurosci. 2018;19(3):138-52.

87.   Pillai R, Uyehara-Lock JH, Bellinger FP. Selenium and selenoprotein function in brain disorders. IUBMB Life. 2014;66(4):229-39.

88.   Bellinger FP, Raman AV, Rueli RH, Bellinger MT, Dewing AS, Seale LA, et al. Changes in selenoprotein P in substantia nigra and putamen in Parkinson's disease. J Parkinsons Dis. 2012;2(2):115-26.

89.   Romero-Ramos M, Venero JL, Cano J, Machado A. Low selenium diet induces tyrosine hydroxylase enzyme in nigrostriatal system of the rat. Brain Res Mol Brain Res. 2000;84(1-2):7-16.

90.   Torres DJ, Yorgason JT, Mitchell CC, Hagiwara A, Andres MA, Kurokawa S, et al. Selenoprotein P Modulates Methamphetamine Enhancement of Vesicular Dopamine Release in Mouse Nucleus Accumbens Via Dopamine D2 Receptors. Front Neurosci. 2021;15:631825.

91.   Mita Y, Uchida R, Yasuhara S, Kishi K, Hoshi T, Matsuo Y, et al. Identification of a novel endogenous long non-coding RNA that inhibits selenoprotein P translation. Nucleic Acids Res. 2021;49(12):6893-907.

92.   Averitt DL, Eidson LN, Doyle HH, Murphy AZ. Neuronal and glial factors contributing to sex differences in opioid modulation of pain. Neuropsychopharmacology. 2019;44(1):155-65.

93.   Taylor AMW, Becker S, Schweinhardt P, Cahill C. Mesolimbic dopamine signaling in acute and chronic pain: implications for motivation, analgesia, and addiction. Pain. 2016;157(6):1194-8.

94.   Harris HN, Peng YB. Evidence and explanation for the involvement of the nucleus accumbens in pain processing. Neural Regen Res. 2020;15(4):597-605.

95.   Werner CT, Altshuler RD, Shaham Y, Li X. Epigenetic Mechanisms in Drug Relapse. Biol Psychiatry. 2021;89(4):331-8.

96.   Ersche KD, Meng C, Ziauddeen H, Stochl J, Williams GB, Bullmore ET, et al. Brain networks underlying vulnerability and resilience to drug addiction. Proc Natl Acad Sci U S A. 2020;117(26):15253-61.

97.   Ersche KD, Williams GB, Robbins TW, Bullmore ET. Meta-analysis of structural brain abnormalities associated with stimulant drug dependence and neuroimaging of addiction vulnerability and resilience. Curr Opin Neurobiol. 2013;23(4):615-24.

Figure 1 Experimental timeline and correlational analyses. **a)** Rats underwent testing for stress- and

anxiety-related behaviors (OFT: open field test; EPM: elevated-plus maze) and nociceptive threshold (TF: tail flick) prior to jugular catheterization surgery, with tests repeated following heroine experience. Heroin self-administration occurred for 3 weeks (12-h/session, 4 days a week) followed by a progressive ratio test and an additional 3 days of self-administration training. Next, rats underwent a within-session extinction-heroin prime reinstatement test (6-h). Extinction training sessions (2-h/session) then occurred followed by a cue-induced reinstatement test (2-h). Spleen sample were collected for genetic analyses at the conclusion of training. **b)** Correlation matrix for behavioral trait covariance. Behavioral testing that occurred prior to heroin experience is designated with a 1, and testing following heroin training as 2. Traits in bold are those with at least one significant QTL identified in GWAS analysis. (n=874)

Figure 2 Heritability and genome-wide association results. **a)** SNP heritability (h^2) measure ± SEM for all behavioral traits assessed. Significant h^2 measures ($p<0.05$) are indicated in blue, and non-significant traits are in orange. **b)** Porcupine plot for selected behavioral traits. Top SNPs exceeding the genome-wide

significance threshold are highlighted and identified (red: -log(p)=5.36 and blue: -log(p)=5.58, which

correspond to permutation derived p values of 0.10 and 0.05, respectively. (n=874)

Figure 3 GWAS results for escalation of heroin intake across the entire session. Regional association plot for chromosome 10, with the top SNP in purple and chromosome location indicated on the plot. Other SNPs are colored based on their linkage disequilibrium with the top SNP. Known genes of SNPs are indicated below the plots. (n=874)

Figure 4 GWAS results for total heroin consumed across self-administration training. **a)** Regional association plot for chromosome 11, with the purple dot indicating the top SNP and location on chromosome indicated on the plot. The color of additional SNPS refers to extent of linkage disequilibrium with the top SNP. Known genes of SNPs are indicated below the plots. **b)** Effect plots for total heroin consumption during training (left: normalized data; right: raw data) for OUD phenotypes (Vulnerable: pink; Intermediate/resilient: yellow) according to peak SNP allelic variant. Half of a violin plot for each phenotype is shown, with the solid line indicating the median value of the data and the dash lines representing the upper and lower interquartile range. **c)** Top SNP allelic variant OUD phenotype cluster composition differ (x^2(1, 853)=18.12, p<0.0001), with the minor allele associated with greater OUD resiliency. (n=853; Vulnerable (pink), n= 361; Intermediate/resilient (yellow), n= 492)

Figure 5 GWAS results for break point achieved during the progressive ratio test. **a)** Regional association plot for chromosome 19, with the top SNP in purple and the color of other significant SNPs referring to level of linkage disequilibrium with the top SNPs. The location of the top SNP is indicated on the plot. Known genes of SNPs are indicated below the plots. **b)** Effect plots showing break point achieved during the progressive ratio test (left: normalized data, right: raw data) for OUD vulnerable (pink) and intermediate/resilient (yellow) rats in relation to the top SNP allelic variant. Half a violin plot is shown for each OUD phenotype, with the solid line indicating the median value of the data set, whereas the dashed lines represent the upper and lower interquartile range. **c)** OUD phenotypic cluster composition for both peak SNP allelic variants. Cluster composition between variants differed (x^2(1, 864)=18.91, p<0.0001), with the minor allele heavily biased toward an OUD vulnerable phenotype. (n=864; Vulnerable (pink), n= 345; Intermediate/resilient (yellow), n= 519)

Figure 6 GWAS results for analgesic threshold during the TF baseline session prior to heroin experience. Regional association plots for **a)** chromosome 2 and **b)** chromosome 19. The top SNP identified for each chromosome is in purple with chromosome location indicated on the plot. The color of other SNPs corresponds to extent of linkage disequilibrium with the top SNP. Known genes of SNPs are indicated below the plots. (n= 874)

Trait	Description
Prior to heroin experience	
OFT distance 1	Total distance travelled
OFT time 1	Time spent moving
EPM closed 1	Time spent in closed arms
EPM open 1	Time spent in open arms
TF baseline 1	Average latency of tail removal during baseline (i.e., saline) session
TF test 1	Average latency of tail removal during test (i.e., heroin) session
Heroin measures	
Escalation of intake: Hour 1	Average consumption (µg/kg) from hour 1 for sessions 1-3 subtracted from average hour 1 consumption for sessions 10-12
*Escalation of intake: Hour 1-12	Average consumption (µg/kg) from entire session for sessions 1-3 subtracted from average total consumption for sessions 10-12
*Consumption	Cumulative heroin consumption (µg/kg) for sessions 1-12
*Breakpoint	Break point (i.e., most active lever presses willing to expend to receive an infusion of heroin) during progressive ratio test
*Extinction: Burst	Active lever presses during the first 2 hr of the within-session extinction-prime reinstatement test
Extinction: Prime	Active lever presses between hour 2-4 in the within-session extinction-prime reinstatement test
*Extinction: Day 6	Active lever presses made during extinction session 6
Extinction de-escalation	Active lever presses during extinction session 1 subtracted from session 6
*Prime reinstatement	Active lever presses during prime reinstatement test
*Cued reinstatement	Active lever presses during cued reinstatement test
Following heroin experience	
OFT distance 2	Total distance travelled
OFT time 2	Time spent moving
OFT difference distance	Total distance travelled during OFT 1 subtracted from OFT 2
OFT time difference	Total time spent moving during OFT 1 subtracted from OFT 2
EPM closed 2	Time spent in closed arms
EPM open 2	Time spent in open arms
EPM difference closed	Time spent in closed arm during EPM 1 subtracted from EPM 2
EPM difference open	Time spent in open arm during EPM 1 subtracted from EPM 2
TF baseline 2	Average latency of tail removal during baseline (i.e., saline) session
TF test 2	Average latency of tail removal during test (i.e., heroin) session
TF difference baseline	Average latency of tail removal during TF baseline 1 subtracted from TF baseline 2
TF difference test	Average latency of tail removal during TF test 1 subtracted from TF test 2

Supplemental Table 1 Description of behavioral traits assessed in GWAS. Traits included those measured both prior to (time point 1) and following (time point 2) heroin experience, and the difference between the two time points, along with several behaviors associated with heroin taking, refraining and seeking.

Traits used in the non-linear clustering approach to categorize rats into OUD phenotypes are designated with an asterisk.

Trait	Covariates
Prior to heroin experience	
OFT distance 1	Weight at time of surgery; Batch 6
OFT time 1	Batch 6
EPM closed 1	Weight at time of surgery; Age at time of experiment; Sex; Batch 2, 3, 4, 6, 12, 13, 15 and 17; Two specific testing chambers
EPM open 1	Weight at time of surgery; Sex; Batch 2, 3, 4, 12 and 17 ; Coat color: black hood; Three specific testing chambers
TF baseline 1	Batch 3 and 4; Coat color: black and brown hood
TF test 1	Weight at time of surgery; Sex; Three specific testing chambers
Heroin measures	
Escalation of intake: Hour 1	None
Escalation of intake: Hour 1-12	None
Consumption	Weight at time of surgery; Age at time of first self-administration session; Sex; Batch 14 and 15; Two specific testing chambers
Break point	Age at time of experiment; Behavior for Batch 14 and 15; One specific testing chamber
Extinction burst	Weight at time of surgery; Batch 4 and 17; 1 specific testing chamber
Extinction: Prime	Weight at time of surgery; Batch 2, 4, 14, 17; Two specific testing chambers
Extinction: Day 6	Batch 3
Extinction de-escalation	Batch 2
Prime reinstatement	Weight at time of surgery; Weight at time of experiment; Sex; batch 3; One specific testing chamber
Cued reinstatement	Weight at time of surgery for; Age at time of experiment; Behavior for Batch 2, 3, 15 and 17; Two specific testing chambers
Following heroin experience	
OFT distance 2	Age at time of experiment for Batch 2
OFT time 2	Age at time of experiment; Batch 2
OFT difference distance	Batch 6
OFT time difference	Batch 6
EPM closed 2	Weight at time of surgery; Sex; Batch 2, 6, 12, 16. One specific testing chamber
EPM open 2	Weight at time of surgery; Sex; Batch 2, 6, and 16; Coat color: albino and black hood; Two specific testing chambers
EPM difference closed	Batch 2, 3, 4, 5, 6, 7, 9, 10, 11, 13 and 16
EPM difference open	Batch 7 and 16. Coat color: albino
TF baseline 2	Weight at time of surgery; Batch 2, 3, 4, 8, 14
TF test 2	Weight at time of surgery; Sex; Batch 3, 4, 7; Three specific testing chambers
TF difference baseline	Batch 2
TF difference test	Weight at time of surgery

Supplemental Table 2 Covariates for behavioral traits from rats tested at MUSC. Covariates that explained >2% of variance were regressed out prior to further analyses. Batches of 40 rats (20 males and 20 females) were shipped every few months from early 2019 through the end of 2022. Testing chamber refers to behavioral testing chamber that all heroin taking, extinction and seeking occurred in. Heterogeneous rats have 5 distinct coat colors: albino, black, brown, black hood and brown hood.

Trait	Covariates
Prior to heroin experience	
OFT distance 1	Weight at time of surgery; Sex; Four specific testing chambers
OFT time 1	Weight at time of surgery; Sex; Batch 8 and 14; Two specific testing chambers
EPM closed 1	Batch 3, 9, 11, 14 and 16
EPM open 1	Weight at time of surgery; Age at time of experiment; Sex; Batch 9 and 11; One specific testing chamber
TF baseline 1	Weight at time of surgery; Batch 5, 7, 8 and 14; Coat color: brown, albino and brown hood
TF test 1	Weight at time of surgery; Sex; Batch 3, 4, 5, 6, 7, 13
Heroin measures	
Escalation of intake: Hour 1	Batch 4, 5, 12 and 13
Escalation of intake: Hour 1-12	Batch 4, 5, 6, 12 and 14
Consumption	Weight at time of surgery; Age at time of first self-administration session; Sex; One specific testing chambers
Break point	Weight at time of surgery; Sex; Batch 4 and 11
Extinction burst	Weight at time of surgery; Sex
Extinction: Prime	Weight at time of surgery; Sex; Batch 6
Extinction: Day 6	Weight at time of surgery; Sex; Batch 6, 12 and 14; One specific testing chamber
Extinction de-escalation	Batch 6
Prime reinstatement	Weight at time of surgery; Sex; batch 8, 10 and 12; One specific testing chamber
Cued reinstatement	Weight at time of surgery for; Sex; Batch 12, 13, 14; One specific testing chamber
Following heroin experience	
OFT distance 2	Weight at time of surgery; Age at time of experiment; Sex; Two specific testing chambers
OFT time 2	Weight at time of surgery; Sex; Batch 8 and 16
OFT difference distance	Age at time of experiment; Sex; Batch 8; Two specific testing chambers
OFT time difference	Age at time of experiment; Sex; Two specific testing chambers
EPM closed 2	Weight at time of surgery; Batch 4 and 8
EPM open 2	Weight at time of surgery; Sex; Batch 6; One specific testing chamber
EPM difference closed	Batch 3 and 11
EPM difference open	Two specific testing chambers
TF baseline 2	Weight at time of surgery; Sex; Batch 7 and 16; Coat color: albino; One specific testing chamber
TF test 2	Weight at time of surgery; Age at time of experiment; Batch 4, 5, 8, 10, 11, 12, 13, 14 and 16
TF difference baseline	None
TF difference test	Batch 3, 5, 7, 8, 9, 10, 11, 12 and 14

Supplemental Table 3 Covariates for behavioral traits from rats tested at UCAM. All covariates that explained >2% of variance for a trait were regressed out prior to continuing analyses. Rats were shipped every few months from early 2019 through the end of 2022 in batches of 40 rats (20 males and 20 females). Testing chamber refers to behavioral testing chamber that all heroin taking, extinction and seeking occurred in. Heterogeneous rats have 5 distinct coat colors: albino, black, brown, black hood and brown hood.

Behavioral trait correlations

Trait 1	Trait2	Spearman r	p-value	n
Escalation of intake: Hour 1	Break point	0.39	1.98×10^{-33}	874
	Cued reinstatement	0.28	1.26×10^{-16}	866
	Escalation of intake: Hour 1-12	0.62	7.03×10^{-94}	874
	Extinction: Prime	0.17	4.93×10^{-7}	874
	Extinction: Burst	0.19	9.72×10^{-9}	874
	Extinction: Day 6	0.22	4.02×10^{-11}	874
	Consumption	0.39	1.02×10^{-32}	874
Escalation of intake: Hour 1-12	Break point	0.39	6.95×10^{-34}	874
	Cued reinstatement	0.27	2.87×10^{-16}	866
	Extinction: Prime	0.17	2.40×10^{-32}	874
	Extinction: Burst	0.21	2.36×10^{-10}	874
	Extinction: Day 6	0.21	8.70×10^{-10}	874
	Prime reinstatement	0.22	2.46×10^{-11}	874
	Consumption	0.29	3.67×10^{-18}	874
Consumption	Break point	0.57	7.83×10^{-78}	874
	Cued reinstatement	0.48	1.12×10^{-51}	866
	Extinction: Prime	0.31	1.33×10^{-20}	874
	Extinction: Burst	0.38	1.05×10^{-30}	874
	Extinction: Day 6	0.34	1.10×10^{-25}	874
	Prime reinstatement	0.39	1.20×10^{-33}	874
	OFT time 2	0.11	0.0009	868
	EPM closed 2	-0.12	0.0006	873
Break point	Cued reinstatement	0.51	1.96×10^{-58}	866
	Extinction: Prime	0.34	6.60×10^{-25}	874
	Extinction: Burst	0.45	3.64×10^{-45}	874
	Extinction: Day 6	0.41	1.54×10^{-36}	874
	OFT time 2	0.15	1.01×10^{-5}	868
	OFT distance 2	0.11	0.0001	868
Extinction: Burst	Cued reinstatement	0.47	5.72×10^{-49}	866
	Extinction: Prime	0.41	1.43×10^{-36}	874
	Extinction: Day 6	0.40	1.57×10^{-34}	874
	Extinction de-escalation	-0.11	0.001	874
	Prime reinstatement	0.39	6.52×10^{-31}	874
Extinction: Prime	Cued reinstatement	0.32	1.54×10^{-22}	866
	Extinction: Day 6	0.27	9.39×10^{-16}	874
	Extinction de-escalation	-0.11	0.001	874
	Prime reinstatement	0.36	1.71×10^{-27}	874
	EPM closed 1	-0.13	0.0002	873
	EPM closed 2	-0.12	0.0003	873
Prime reinstatement	Cued reinstatement	0.38	1.33×10^{-30}	866
	OFT time 2	0.12	0.00006	868

	EPM closed 2	-0.13	7.64×10^{-5}	873
Extinction: Day 6	Cued reinstatement	0.54	1.21×10^{-67}	866
	Extinction de-escalation	0.44	7.92×10^{-42}	874
	OFT time 2	0.13	0.0001	868
Cued reinstatement	OFT distance 1	0.13	0.0001	862
	OFT distance 2	0.12	0.0003	860
	OFT time 1	0.13	8.33×10^{-5}	862
	OFT time 2	0.15	9.46×10^{-6}	860
	EPM closed 2	-0.11	0.0008	865
OFT distance 1	OFT distance 2	0.45	5.13×10^{-44}	864
	OFT distance difference	-0.49	4.07×10^{-52}	864
	OFT time 2	0.39	2.64×10^{-33}	864
	OFT time difference	-0.51	4.28×10^{-58}	864
OFT distance 2	OFT distance difference	0.46	1.53×10^{-46}	864
	OFT time difference	0.40	1.29×10^{-33}	864
	OFT time 1	0.40	7.59×10^{-35}	864
	OFT time 2	0.89	5.40×10^{-291}	868
OFT distance difference	OFT time difference	0.92	0.00	864
	OFT time 1	-0.47	3.15×10^{-49}	864
	OFT time 2	0.45	2.54×10^{-44}	864
OFT time 1	OFT time difference	-0.54	8.73×10^{-68}	864
	OFT time 2	0.41	4.28×10^{-36}	864
OFT time 2	OFT time difference	0.45	2.52×10^{-43}	864
EPM open 1	EPM open 2	0.25	8.36×10^{-14}	872
	EPM open difference	-0.44	7.78×10^{-43}	872
	EPM closed 1	-0.58	3.65×10^{-78}	872
	EPM closed 2	-0.18	1.33×10^{-7}	872
	EPM closed difference	0.32	1.80×10^{-22}	872
EPM open 2	EPM open difference	0.61	1.82×10^{-90}	872
	EPM closed 1	-0.32	1.32×10^{-22}	873
	EPM closed 2	-0.67	9.29×10^{-166}	873
	EPM closed difference	-0.26	7.65×10^{-15}	873
EPM open difference	EPM closed 1	0.12	0.0005	872
	EPM closed 2	-0.41	5.88×10^{-36}	872
	EPM closed difference	-0.48	8.16×10^{-51}	872
EPM closed 1	EPM closed 2	0.37	2.77×10^{-30}	873
	EPM closed difference	-0.45	0.0005	873
EPM closed 2	EPM closed difference	0.47	2.38×10^{-48}	873
TF baseline 1	TF baseline 2	0.11	0.0012	874
	TF baseline difference	-0.60	7.67×10^{-88}	874
	TF test 2	0.11	0.0013	874
TF baseline 2	TF baseline difference	0.59	7.30×10^{-84}	874
	TF test 2	0.16	1.39×10^{-6}	874
TF test 1	TF test 2	0.17	9.78×10^{-7}	874
	TF test difference	-0.59	3.40×10^{-84}	874
TF test 2	TF test difference	0.57	2.29×10^{-77}	874

Supplemental Table 4 Data statistics for significant correlated behavioral traits. Heroin taking, refraining and seeking behaviors showed the tendency to co-vary with one another, whereas behaviors assessing stress- and anxiety-like behaviors as well as nociceptive threshold exhibited greater within-test covariance. Direction and magnitude of correlation (Spearman r) and total number of animals included in analysis (n) are included in the table. A significance threshold of $p<0.05$ with Bonferroni adjustment for correction for multiple comparisons was implemented, resulting in a $p<0.0018$ in order to attain significance.

Trait	h^{2}SNP	SE	p-value	n
OFT distance 1	0.139	0.048	*<0.001*	870
OFT time 1	0.103	0.046	*0.004*	870
EPM open 1	0.054	0.041	0.073	873
EPM closed 1	0.083	0.043	*0.012*	873
TF baseline 1	0.11	0.046	*0.003*	874
TF test 1	0.047	0.039	0.1	874
Escalation of intake: Hour 1	0.062	0.042	0.059	874
Escalation of intake: Hour 1-12	0.037	0.04	0.168	874
Consumption	0.172	0.052	*<0.001*	874
Break point	0.009	0.039	0.42	874
Extinction: Burst	0.098	0.045	*0.007*	874
Extinction: Prime	0.002	0.036	0.48	874
Extinction: Day 6	0	0.038	0.5	874
Extinction de-escalation	0.003	0.037	0.47	874
Prime reinstatement	0	0.036	0.5	874
Cued reinstatement	0.037	0.04	0.172	866
OFT distance 2	0.232	0.052	*<0.001*	868
OFT distance difference	0	0.033	0.5	864
OFT time 2	0.158	0.049	*<0.001*	868
OFT time difference	0.007	0.033	0.41	864
EPM open 2	0.072	0.042	*0.024*	873
EPM Open difference	0	0.035	0.5	872
EPM closed 2	0.016	0.035	0.31	873
EPM closed difference	0	0.034	0.5	873
TF baseline 2	0.146	0.049	*<0.001*	874
TF baseline difference	0.044	0.04	0.12	974
TF test 2	0.124	0.045	*<0.001*	874
TF test difference	0	0.035	0.5	874
Vulnerable vs resilient/intermediate	0.168	0.051	*<0.001*	862
Intermediate vs vulnerable/resilient	0.093	0.044	*0.007*	862
Resilient vs vulnerable/intermediate	0.15	0.053	*0.002*	862

Supplemental Table 5 SNP heritability table for behavioral traits. Significant relationships are

highlighted in bold. The heritability index (h^{2}SNP), standard error (SE), p-value and total number of

animals in analysis (n) for each trait is recorded.

Supplemental Figure 1 Principle component analysis for behavioral traits with significant QTLs. Two factors were identified and comprised 71.55% of total variance in the data. Heroin-taking behaviors accounted for 46.24% of variance (component 1, purple) with traits exhibiting strong loading (Break point: 0.86; Consumption: 0.81; and Escalation of intake: Hour 1-12: 0.67), and was mutually exclusive from baseline nociception prior to heroin experience (component 2, yellow) which accounted for 25.31% of variance with strong loading (0.99). Axes represent factor loading values for each trait within respected component. (n=874)

Supplemental Materials

Behavioral testing

Elevated-plus maze (EPM): Sessions were conducted in an elevated apparatus (San Diego Instruments; 4 arms at 110.5 cm long and 10.2 cm wide) that consisted of two "closed" arms (30.5 cm high walls) and two "open" arms (i.e., not enclosed). Animals were placed in the center of the maze to start, and movement was tracked automatically using ANY-maze behavioral tracking software (Stoelting, Wood Dale; version 6.17) over the course of the 5-min session. A minimum of 85% of the rat's body had to be within an arm for it to be counted. The percent time the animal spent in each arm was calculated.

Open field test (OFT): Animals were placed in the center of a Plexiglas chamber housed within a metal frame (Omnitech Electronics, Columbus, OH; 40.6 cm L x 40.6 cm W x 30.5 cm H) containing photocells that tracked vertical and horizontal movements. Sessions lasted for 60-min and data was recorded and analyzed for distance and time spent moving using Versamax (Omnitech Electronics, Columbus, OH; version 1.80-0142).

Tail flick test (TF): Testing occurred on a platform with the rat's tail over an infrared light beam equipped with a motion sensor that turned the beam off once the tail was removed, or after 10 seconds (Ugo Basile S.R.L., Gemonio, Italy). TF was comprised of two phases: baseline (1 mg/kg saline injection, s.c.) and test (0.75 mg/kg heroin, s.c.). Injections were administered 15 min prior to testing and the phases were separated by 1 hour. Each test consisted of 4 trials with the location of the tail tested being adjusted by 1 cm each trial to prevent tissue damage. Average latency to remove the tail from the beam was calculated.

Heroin taking, refraining and seeking measures

Prior to training, rats were outfitted with an indwelling jugular catheter and post-operatively administered an antibiotic (Cefazolin, 0.2 mg/kg, s.c.; or enrofloxacin, 1 mg/kg, i.v.) and analgesic (Ketorolac, 2 mg/kg, s.c.; or Meloxicam, 0.5 mg/rat, s.c.). Training occurred in standard behavioral testing chambers (Med

Associates, St. Albans, VT) housed within a sound attenuating box outfitted with a ventilation fan. Two levers with a light above each were on one chamber wall, and opposite was a house light and speaker. The house light turned on at the start of all sessions. A fixed ratio 1 schedule of reinforcement was used for heroin self-administration whereupon a press on the active lever resulted in an infusion of heroin (20 µg/kg/100 µl infusion over 3-sec) and a 5-sec presentation of a light/tone cue. The house light turned off for 20-sec at the time of infusion to signal a time out period during which additional active lever presses were recorded but without consequence. Four training sessions (12-h or 300 infusions earned) occurred per week (Monday-Friday) with one randomized day off per week, totaling 12 sessions. The progressive ratio test occurred next to assess heroin break point. Testing terminated after 12-h or 1-h of no earned infusion. Heroin self-administration training was then re-established for 3 days, followed by a within-session extinction-prime test lasting 6 hours. Testing occurred under extinction conditions (i.e., active lever presses no longer resulted in heroin infusion or light/tone cue presentation). With two hours left in the session, animals received a heroin prime injection (0.25 mg/kg, s.c.). Extinction training session and a test for cued reinstatement followed. Throughout testing, presses on the inactive lever were recorded but had no consequence.