CapP mediates the structural formation of biofilm-specific pili in the opportunistic

human pathogen *Bacillus cereus*

Ana Álvarez-Mena[1], Muhammed Bilal Abdul Shukkoor[2], Joaquín Caro-Astorga[1,3],

Melanie Berbon[2], María Luisa Antequera-Gómez[1], Axelle Grélard[2], Brice Kauffmann[4],

Estelle Morvan[4], Antonio de Vicente[1], Birgit Habenstein[2], Antoine Loquet[2,5], Diego

Romero[1,5,6]

[1]Departamento de Microbiología, Instituto de Hortofruticultura Subtropical y Mediterránea

La Mayora, Universidad de Málaga-Consejo Superior de Investigaciones Científicas,

Universidad de Málaga, Málaga, Spain

[2]Institut de Chimie et de Biologie des Membranes et des Nano-objets, Institut Européen

de Chimie et Biologie, CNRS, UMR5248, Université de Bordeaux, Pessac, France

[3]The Francis Crick Institute, London, NW1 1AT, United Kingdom

[4]Institut Européen de Chimie et Biologie, UAR3033 CNRS, Université de Bordeaux,

Inserm US01, Pessac F-33600, France

[5]Corresponding authors

[6]Lead contact: diego_romero@uma.es

X: @diegoromerohi

SUMMARY

Polymeric proteinaceous filaments are structural scaffolds that diversify the functionality

of the bacterial extracellular matrix. Here, we report a previously uncharacterized bacterial

factor called *bc1280* that is exclusive to *B. cereus* group and indispensable for the

establishment of a biofilm lifestyle. We propose that BC1280 is an essential chaperone

for the assembly of the filamentous platform that tightly controls the polymerization of

heteropili containing CalY and TasA as major subunits in a concentration-dependent

manner. Additionally, BC1280 modulates the expression of EPS via an uncharacterized

pathway that is activated by a protease and an ECF-type sigma factor. The pilus

biogenesis system described in this work highlights the complexity of extracellular matrix

assembly in *B. cereus* and introduces a singular three-part structuration mechanism

during biofilm formation and maturation.

KEYWORDS: fibrillary proteins, chaperone, pili, biofilm, extracellular matrix, *Bacillus*

cereus

INTRODUCTION

Bacillus cereus, a bacterium commonly found in soil, is capable of colonizing the gastrointestinal tracts of mammals or arthropods; *B. cereus* is contracted through the ingestion of contaminated food, including fresh vegetables and ready-to-eat meals.[1–4] Consequently, *B. cereus* strains are frequently identified as the etiological agent of gastrointestinal diseases outbreaks, and the diseases can progress through emetic or diarrheal symptoms.[5,6] As approximately 70% of microbial infections in humans are reportedly associated with biofilms, much effort has been dedicated to understanding the genetic basis and structural components that, upon activation, lead to the formation of these bacterial communities.[7,8] The transition from a planktonic to a sessile lifestyle is metabolically focused on the synthesis of proteins, polysaccharides and extracellular DNA (eDNA), which ultimately organize the extracellular matrix (ECM).[9,10] The ECM performs a wide range of functions, including adhesion to biotic and abiotic surfaces, regulation of nutrient and signaling molecule transport, defense against physical-chemical stress, and virulence, and provides support for biofilm architecture.[11,12] One of the most attractive components of the ECM is the fibril-forming proteins that serve as structural scaffolds, facilitating the assembly of the microbial community and contributing to surface adhesion, among other biological roles.[13,14] These proteins, termed functional amyloid-like proteins, share remarkable similarities with their pathogenic counterparts in humans, highlighted by their propensity to polymerize into highly ordered fibrillary aggregates, resistance to diverse environmental challenges, and their enrichment in β-sheet content.[15–18]

The development of *B. cereus* biofilms involves genetic pathways and structural

components of the ECM identical to those found in *B. subtilis*, a closely related species.

However, differences arise and are predicted to have ecological and pathological

implications.[19,20] Complementary reports have described the role of flagella, matrix

proteins, eDNA and exopolysaccharides in the formation of biofilms on *B. cereus.*[21–24] A

previous study identified a specific genomic region in *B. cereus* that is orthologous to the

corresponding region in *B. subtilis*, and it is expressed mainly in biofilm-associated

cells.[25,26] This region encodes an ortholog of *sipW;* two orthologs of *tasA*, called *tasA* and

calY; the main biofilm regulators *sinI* and *sinR*; and an uncharacterized locus called

bc1280. Interestingly, no orthologs for *tapA* are present in *B. cereus*. SipW is the signal

peptidase that processes TasA and TapA for efficient secretion, and both TasA and CalY

of *B. cereus* conserve the signal peptide with the canonical cleavage site recognized by

the cognate peptidase SipW.[27,28] The genes *sipW* and *tasA* constitute an operon, and

calY is expressed independently.[25] The deletion of *tasA* or *calY* resulted in differential

defects in biofilm formation, suggesting that the product of each gene might be involved

in cell-to-abiotic surface or cell-to-cell interactions, respectively.[25] TasA from *B. subtilis*

has been described as a globular protein in its monomeric state that polymerizes into

filamentous structures, which can be amyloid or non-amyloid, depending on the

experimental context.[29–33] TasA native fibers, observed in vivo in biofilm samples and also

in vitro upon extraction, exhibit an enrichment in β-sheet conformation while preserving

the β-rich globular structure of the monomer. Consequently, these TasA monomers

polymerize into filaments via a donor strand complementation mechanism, and as a

result, their structure diverges from the canonical cross-β organization associated with

amyloid proteins.[33] Nevertheless, recombinant TasA purified by heterologous expression

in *Escherichia coli* has been shown to form fibrils with typical amyloid properties, such as

the cross-beta signature.[29,32] TasA might therefore adapt its secondary and tertiary

structure in a context-dependent manner, forming either an amyloid conformation or a β-

rich globular fold. Extensive structural studies on TasA of *B. subtilis* exist, but only El-

Mammeri et al. have reported singularities in recombinantly produced TasA and CalY

from *B. cereus,* based on their secondary structure propensity, despite the cross-β

amyloid pattern displayed by these paralogous proteins.[32]

In this study, we provide the first evidence that the *bc1280* product is indispensable

for the growth of *B. cereus* biofilms, relying on two complementary functions. The first is

a structural role, i.e., the promotion of CalY polymerization, which is related to the correct

polymerization of TasA-CalY heterofibrils, essential for the formation and stability of the

ECM scaffold. The second is a regulatory function in which the expression of biofilm-

related EPSs is controlled via an undescribed sigma-antisigma pathway. Moreover, our

findings indicate that when BC1280 is expressed heterologously in *E. coli*, the N-domain

displays a predominantly β-sheet conformation within its rigid core, while the tandem

repetitions found in the C-domain potentially facilitate the interaction with CalY. From its

localization in the cell wall, BC1280 defines and triggers the polymerization of CalY, which

in turn incorporates TasA into nascent heterofilaments. Based on these findings, BC1280

was renamed CapP, which is a CalY-assisting polymerization protein. Our study provides

a new paradigm for the complex assembly of this proteinaceous appendage in *B. cereus*

biofilms, identifies CapP as a structural orchestrator for efficient ECM development, and

highlights CapP as a promising target for precisely targeted antibiofilm therapies.

RESULTS

***bc1280* is an essential locus in the developmental program of biofilm formation**

First, a phylogenetic analysis of BC1280 was performed using the amino acid sequence

of the type strain *B. cereus* ATCC14579 as a reference. The results revealed that BC1280

was present within the *B. cereus* group but not within the phylogenetically related species

B. subtilis (Figure S1A). The *bc1280* product was annotated as a hypothetical protein with

the following putative regions (Figure 1A): i) a signal peptide from position 1 to 39

predicted to be processed by a signal peptidase type I (SignalP5.0) (Figure S1B); ii) an

N-terminal domain from position 40 to 189 (the N-domain from now on) that contains a

highly conserved domain of unknown function (from I^{33} to E^{157}) annotated as DUF4047;

and iii) a C-terminal domain (the C-domain from now on) from position 190 to the C-

terminal end of the protein. This domain is characterized by a variable number of perfect

repetitions of QKKV/AEE motifs, depending on the *B. cereus* strains (Figure S1C).

A previous transcriptome analysis of the genetic features that distinguish

planktonic and biofilm cells in *B. cereus* showed that *bc1280* is specifically expressed in

biofilm-associated cells, and a complementary study revealed that the expression of

bc1280 comprised the regulon of the biofilm-specific master regulator SinR.[34] We

confirmed the biofilm-specific expression of *bc1280* via RT–qPCR analysis of samples

taken at 48 hours (Figure S1D) and determined that *bc1280* is transcribed as a

monocistronic unit independent of the previously reported transcriptional units *sipW-tasA*

or *calY* (Figure S1E).[25] The biofilm of the *B. cereus* wild-type strain ATCC14579 was

characterized by a white biomass ring that adhered to the plate wells; this ring was usually

visible at 24 hours and reached maturity at 72 hours (Figure 1B). A *bc1280* knockout

mutant strain (*Δbc1280*) was completely arrested during biofilm formation, a phenotype

that differed from the less adhesive or thinner biomass of *ΔtasA* or *ΔcalY,* respectively

(Figure 1B).[25] According to the distribution and conservation observed in our phylogenetic

analysis, deletion of *bc1280* in the emetic strain of *B. cereus* AH187, which belongs to a

different *B. cereus* subgroup, resulted in the complete elimination of pellicle formation at

the air–liquid interface compared to that in the wild-type strain (Figure S1F).

The wild-type phenotype for biofilm formation was completely restored through

complementation of *Δbc1280*, which was achieved through ectopic integration of the

bc1280 locus (*bc1280*::*Δbc1280*) under the control of its own putative promoter at a

neutral locus (Figure 1B). Moreover, these findings suggested that the promoter for the

expression of *bc1280* must be contained within the noncoding sequence between the

open reading frame of *tasA* and *bc1280*. BC1280 contains a distinctive C-domain with

perfect tandem repetition. As previously described, these characteristic regions often

have significant structural and functional importance for the stabilization of protein folding

overall.[35] Therefore, we investigated whether the C-domain is involved in biofilm

formation. The biofilm formation phenotype of *Δbc1280* was still rescued by a version of

BC1280 that lacked the C-domain from Q^{191}-E^{279} (Δ*C-domain*) (Figure 1B). Quantitative

measurements at 575 nm of the biomass stained with crystal violet reinforced the visual

differences in biofilm thickness (Figure 1C). No significant differences were observed

between the strains at 24 hours, when initial cell attachment precedes biofilm maturation;

however, differences became evident at 48 hours, coinciding with the initiation of ECM

assembly and the growth of biofilms. After cells are initially attached to surfaces, cell-to-

cell interactions occur that facilitate the subsequent growth of the biofilm, a cellular

phenomenon that can be monitored through auto-aggregation experiments. The wild-type

and *ΔC-domain* strains exhibited sharp aggregation kinetics; in contrast, *Δbc1280* did not

form sediments at the bottom of the tube at the end of the 24-hour experiment (Figure

1D). Taken together, these results indicate that the product encoded by *bc1280* is

involved in the developmental stages of bacterial cell-to-cell adhesion and the subsequent

assembly and maturation of the ECM, a function that may rely on the N-domain of the

protein.

The biofilm-defective phenotype of *bc1280* is independent of the expression level

of *tasA* or *calY*

The *Δbc1280* phenotype may result from changes in the expression levels of ECM

components or structural malfunctioning in the polymerization of TasA-CalY

heterofilaments; these two alternative explanations are not mutually exclusive. A whole-

transcriptome analysis of the *Δbc1280* strain revealed 35 genes that were overexpressed

and 4 genes that were repressed compared to those in the wild-type strain at 24 h.

However, none of these genes were predicted to be associated with biofilm formation

(Table S1). Notable transcriptional changes were observed at 48 hours, with 21 genes

overexpressed and 28 genes repressed in *Δbc1280*, including *tasA* and *calY* (log2FC

values of -3.13 and -3.14, respectively) (Table S2). This result was further supported by

RT–qPCR analysis, which revealed lower expression levels for the genes located within

the *eps1* and *eps2* genomic regions (Figure 2A). Our transcriptomic analysis additionally

revealed that the expression of *bc2793* and *bc2794* pairs (log2FC values of -8.11 and -

8.43, respectively) was significant repressed. These uncharacterized loci are annotated

as a putative Clp protease and an RNA polymerase ECF-type sigma factor, respectively.

ECF sigma factors are known to transduce changes in the cell envelope into a bacterial

genetic response[36,37]; thus, we hypothesized that under normal conditions, *bc2793* and

bc2794 positively regulate the gene expression of the main ECM components. The

overexpression of *bc2793* and *bc2794* in *Δbc1280* and induction with a 10 µM solution of

isopropyl β-D-thiogalactopyranoside (IPTG) significantly increased the biofilm biomass

(Figure S2A). Considering that the putative protease (*bc2793*) could regulate the activity

of *bc2794*, we overexpressed only the ECF factor, resulting in a six-fold increase in crystal

violet staining at 100 µM IPTG compared to the strain without IPTG induction (Figure

S2A). In contrast to these findings, the deletion of *bc2794* did not significantly affect

biofilm formation, whereas the deletion of *bc2793-bc2794* resulted in a slight reduction in

biofilm production compared to the wild-type strain, though this reduction was minimal

relative to the biofilm-impaired phenotype observed in *Δbc1280* (Figure S2A). RT–qPCR

analysis revealed no significant changes in the expression levels of *tasA* or *calY* in

Δbc1280-overexpressing *bc2794* or *bc2793-bc2794* compared to those in *Δbc1280*

(Figure 2B). However, the relative expression levels of *eps1* and *eps2* were significantly

upregulated in response to both genetic combinations. Thus, the ECF factor and its anti-

sigma factor were associated with the regulation of exopolysaccharide expression in *B.*

cereus, rather than the biofilm-related genes *tasA* and *calY*.

In accordance with our hypothesis on the structural role of BC1280 in the

polymerization of TasA or CalY, no reversion of phenotypes was observed in *Δbc1280*

cells overexpressing *tasA, calY*, or both loci (Figure S2B). TEM analysis of negatively

stained samples and immunolabeled with anti-TasA or anti-CalY antibodies revealed

(Figure 2C and D): i) the presence of filaments from both proteins on the surfaces of WT

cells, ii) aggregates reactive to anti-TasA antibodies in *Δbc1280* cells overexpressing

tasA, and iii) the absence of filaments in *Δbc1280* cells overexpressing *calY* but with an

extracellular signal associated with the anti-CalY antibody. We next examined whether

TasA or CalY, which were produced by heterologous expression in *E. coli* and added

exogenously to *Δbc1280*, might rescue the wild-type biofilm phenotype. The monomeric

conformation of TasA or CalY failed to rescue biofilm formation; however, ring biomass

formation was only partially restored when the proteins were incubated in vitro for one

week to promote filaments assembly and then added at a final concentration of 6 μM (Fig

2D). In line with these findings, TEM analysis revealed that upon addition of TasA and

CalY in their already assembled conformation, the cell surfaces were decorated with

specific immunoreactive aggregates of these proteins (Figure 2E). Taken together, these

data suggest that BC1280 is crucial for ECM maturation and raise the hypothesis that it

might be involved in the polymerization of CalY or TasA. However, to fully understand the

downregulation of *tasA* and *calY* over a longer period, additional research is required to

ascertain whether this is a direct result of the absence of *bc1280* or an indirect

consequence of an accumulation of TasA and CalY in the cytoplasm that are not being

secreted.

Our previous findings indicated that the N-domain might partially fulfill the function

of BC1280 in biofilm formation by specifically facilitating cell-to-cell interactions.

Therefore, this domain may influence the final localization of TasA or CalY within the

ECM. TEM analysis and immunolabeling of the *ΔC-domain* strain revealed that nanogold

particles associated with TasA extracellular filaments were exposed on the cell surfaces;

nevertheless, the morphologies of these filaments were different from those formed in the

wild-type cells (Figure 2C and F). Nanogold particles corresponding to CalY monomers

or oligomers, but not filaments, were observed in the extracellular matrix of the *ΔC-*

domain strain (Figure 2C and F). Based on the alterations observed in the polymerization

of TasA and especially CalY in the *ΔC-domain* strain, the C-domain of BC1280 might be

involved in the polymerization of these proteins or their initial association with the cell

surface.

BC1280 is a cell wall-associated protein that partially colocalizes with CalY in the

ECM

To precisely determine how BC1280 contributes to the assembly of the ECM, we further

studied the localization of the protein at the cellular level. Our preliminary data indicated

that BC1280 is likely expressed at very low levels and is undetectable by

immunocytochemistry assays. Therefore, we conducted these experiments using a

derivative *Δbc1280* strain that expressed a version of BC1280 fused with a histidine tag

at the carboxyl terminal domain. The biofilm formation of *Δbc1280* was rescued by

plasmid leakage without IPTG induction or upon induction with a 10 μM IPTG solution

(Figure S3B). Furthermore, fractionation and immunodetection experiments with an anti-

histidine antibody confirmed the presence of BC1280 in the extracellular matrix, cell wall,

and cellular fractions (membrane and cytosolic content) (Figure 3A). TEM analysis and

immunolabeling using anti-histidine antibodies revealed that foci of nanogold particles

related to BC1280 were distributed regularly along filaments that emerged from the cell

surfaces (Figure 3A). To better determine the localization of BC1280 in the cell wall, we

conducted confocal laser scanning microscopy (CLSM) and immunocytochemistry using

anti-histidine and secondary antibodies conjugated to Atto 488 and double staining with

wheat germ agglutinin (WGA) and Hoechst to visualize the cell wall and DNA,

respectively. According to the results of the fractionation experiments, a limited number

of foci corresponding to BC1280 proteins fused to the histidine tag per cell were observed

in association with the cell surface (Figure 3C and S3C). Considering the relevance of the

N-domain to the functionality of BC1280, we also studied its contribution to the cellular

localization of whole BC1280. Based on crystal violet staining of the biomass that had

adhered to the well, the amount of N-domain expressed without IPTG induction or after

induction with 10 µM IPTG solution was sufficient to rescue the biofilm phenotype of

Δbc1280 (Figure S3B)*.* Cellular fractionation and immunoblotting assays demonstrated a

specific reactive band in the extracellular matrix and cellular fractions and an undetectable

signal associated with the cell wall (Figure 3B). In addition, visualization via TEM

confirmed the presence of N-domain-related foci along the filaments emerging from the

cells, as previously observed for the entire protein, but no detectable fluorescent signals

associated with the cell wall were observed via CLSM (Figure 3B and C). The disparities

noted between the full-length and the truncated form of the protein suggest that the C-

domain contributes to the retention of BC1280 in the bacterial cell wall, likely before

BC1280 reaches the ECM.

To further define the connection between BC1280 and CalY or TasA at the cellular

level, we studied their colocalization via immunocytochemical analysis of cells expressing

pUTE657-*bc1280* fused to a histidine tag using specific primary antibodies. CLMS

analysis revealed that CalY and BC1280 colocalized along the fibrils that compose the

ECM (Pearson's coefficient of 0.58 and a p value of 0.0002) (Figures 3D and S3D). In

most cases, the signal corresponding to CalY was homogeneously distributed, and

BC1280 was detected as discrete foci dispersed along the filaments. However, no

colocalization signal was detected for TasA or BC1280, with a Pearson's coefficient of

0.19 and a p value of 0.0002 (Figure S3E and F). These findings were supported by

Mander's coefficient, which represents the percentage of overlapping pixels for each

fluorescent channel. For the BC1280-CalY pair, approximately 65% of the pixels

overlapped in both channels; for BC1280-TasA, the overlap decreased to 26% for Atto

488 and 21% for Alexa Fluor 647 (Figure S3F). The observation of BC1280 and CalY

colocalizing within the ECM hints at a potential structural relationship, including possible

direct interactions, between these proteins.

Heterologously produced BC1280 oligomerizes into macromolecular complexes

driven by the N-domain

BC1280 and the truncated version comprising the N-terminal domain (from A^{39} to G^{190})

were purified by heterologous expression in *E. coli* Lemo21(DE3) and subjected to further

structural analyses (Figure 4A). The two versions of the protein were purified under

soluble conditions by affinity chromatography followed by size exclusion chromatography,

and by tandem mass spectrometry analyses, the amino acid sequences of each protein

were confirmed (Figure S4A). The two fractions collected at different elution volumes in

the size exclusion chromatogram suggested that in buffer containing 20 mM Tris and 50

mM NaCl at pH 7, BC1280 and N-domain$_{39-190}$ could assemble into two main molecular

entities with distinct oligomeric states (Figure 4B and E). Dynamic light scattering (DLS)

analysis showed that both protein assemblies exhibited a degree of polydispersity, with

particles ranging in diameter from ~10 to 600 nm (Figure 4C-D and F-G for BC1280 and

N-domain$_{39-190}$, respectively; for further details, see Table S3). Remarkable differences

were observed in the second elution sample; specifically, the N-domain$_{39-190}$ exhibited a

greater proportion (~53%) of small oligomers than did BC1280 (~18%) (Figure 4D and

G). To complement the DLS results, each fraction was resolved in a native gel (Figure

4H). The sample corresponding to fraction 1 of each protein, which presumably contained

larger aggregates, failed to enter the gel, suggesting that the molecular complex was

larger than 1000 kDa. The second elution fraction was resolved as a band at 480 or 240

kDa for BC1280 or N-domain$_{39-190}$, respectively, which corresponds to oligomers of

approximately 15 subunits each. TEM analysis confirmed that fraction 1 contained large

aggregates of varying sizes for both proteins (Figure S4B). The samples corresponding

to the second elution fraction contained macromolecular complexes that were similar but

smaller in diameter (Figure S4B). Taken together, these results indicate that the variability

in the oligomerization of the full-length protein is preserved in the truncated version that

contains only the N-terminal region.

To further delineate the dynamics of oligomerization, we focused on the second

fraction, which was composed mainly of small oligomers and could promote the formation

of larger molecular entities. The fluorescence signal emitted upon binding to thioflavin T

(ThT)[38] showed a fast polymerization curve without a lag phase, aligning with the

presence of oligomers under native conditions (Figure 4I). Interestingly, the slope of the

exponential phase was more pronounced for the N-domain$_{39-190}$, and the fluorescence

intensity signal was twofold greater than that of the full-length protein after 150 hours of

incubation. These results correspond with the higher percentage of small oligomers

formed by the N-domain$_{39-190}$ (via DLS analysis), which suggested that the truncated

protein seeds rapidly. Furthermore, TEM analysis of negatively stained samples revealed

macroscopic particles of variable sizes and an amorphous aggregated morphology for

both proteins, resembling the aggregates observed for CalY but not the fibrillar structures

formed by TasA (Figure 4J).[32] These results strongly indicate that the N-domain is

involved in protein oligomerization and suggest that the absence of the C-domain in the

truncated protein might facilitate a faster aggregation rate.

The assembled N-domain$_{39-190}$ contains a rigid core with two β-strand-rich regions,

while the C-domain increases global rigidity

The amyloid-like nature of TasA and CalY[32], when heterologously produced in *E.coli*,

along with the association of BC1280 with these two proteins *in vivo* and its tendency to

oligomerize *in vitro*, led us to suspect an amyloid-like nature for BC1280. Since the N-

domain retains the functionality of BC1280 and the ability to form oligomers, this version

of the protein was considered for our initial structural studies. The X-ray diffraction pattern

of the centrifuged pellet of the N-domain$_{39-190}$ assemblies displayed typical features

reminiscent of the cross-β structure of amyloids, with strong signals at 4.7 Å and a slight

signal at ~10-11 Å, corresponding to the inter-strand and inter-sheet spacings,

respectively[39] (Figure 5A). In light of these findings, we predicted protein aggregation and

amyloidogenicity based on the amino acid sequence of BC1280 using the bioinformatics

tools AmylPred2,[40] FoldAmyloid,[41] MetAmyl,[42] PASTA 2.0[43] and TANGO.[44] At least two

algorithms predicted the following amino acid stretches as putative aggregative regions

and amyloid-forming segments: i) L^{51}STAIIFP^{58}, ii) I^{98}S^{99}, iii) L^{115}QNVYT^{120} and iv)

V^{144}SITVN^{149} (Figure S4C; letters in bold). Based on bioinformatics analysis using

FoldUnfold,[45] the N-domain appears intrinsically folded but the C-domain is predicted to

be unstructured, possibly due to the presence of repetitive amino acid sequences rich in

Gln, Lys, and Glu, which are disorder-promoter residues[35,46] (Figure S4C). As the

propensity to form β-sheet-rich self-assembled structures is often associated with regions

enriched in hydrophobic residues, we analyzed the hydrophobicity pattern of BC1280

using ProtScale.[47] The peptide T^{50}LSTAIIFPKT^{60}, previously identified as a putative

amyloidogenic segment, was found to be highly hydrophobic (Figure S4D). Finally,

AlphaFold[48,49] was used to obtain a tentative secondary structure. The model

predominantly exhibits an α-helix conformation with low confidence value, except for a

small central helical sandwich. This contrasts with the enrichment in β-sheet structural

elements derived from ThT polymerization kinetics and the cross-β structure observed by

X-ray diffraction (Figure S4E). We further performed fine structural characterization of the

molecular conformation of these protein assemblies using magic-angle spinning (MAS)

SSNMR, which has previously been employed to study non-crystalline self-assemblies,

such as amyloids.[50–52] Uniform $^{13}C/^{15}N$-labeled N-domain$_{39-190}$ assemblies were produced

to improve NMR sensitivity and perform multidimensional MAS SSNMR studies. To

investigate the presence of a central rigid core within the assemblies, we used dipolar-

based polarization transfer (cross-polarization) and two-dimensional ^{13}C-^{13}C

spectroscopy. A 50 ms mixing time proton-driven spin-diffusion (PDSD) experiment was

used to detect intra-residue ^{13}C-^{13}C correlations from the rigid core of the assembly. For

the spectral analysis, we first attempted to exploit the secondary structure model

predicted by AlphaFold,[48,49] and the Cα and Cβ chemical shifts were predicted using

SPARTA.[53] Subsequently, the predicted spectrum was superimposed with the

experimental 2D ^{13}C-^{13}C spectrum acquired for N-domain$_{39-190}$ (Figure S5A). No

correlation was found between the experimental and predicted NMR spectral patterns,

which reflects the discrepancies between the AlphaFold model and the structure of the

experimentally obtained multimers. These findings highlight that, while bioinformatics

predictions might reflect the relevant structures under certain conditions, differences

between predicted and experimentally observed structures can result from factors such

as the nature of the sample, the buffer composition (pH or salt), or oligomerization. The

NMR data provide evidence for the propensity of BC1280 to undergo structural changes

upon assembly under certain *in vitro* or environmental conditions. Thus, we manually

inspected the SSNMR cross-peaks and identified ~35-40 spin systems (i.e.,

corresponding to ~35-40 residues), as exemplified for an isoleucine in Figure 5B.

Following well-established statistical data for Cα and Cβ chemical shifts according to their

secondary structure,[54] we noticed that these residues are involved mainly in β-sheet

secondary structure elements. This finding was illustrated by the isoleucine spectral

pattern observed with Cα and Cβ chemical shift values of ~60 and 42 ppm, respectively,

which are indicative of a β-sheet conformation (Figure 5B). Approximately 35-40 out of

the 151 amino acids comprising the N-domain$_{39-190}$ were detected within the rigid core by

SSNMR, suggesting that the rigid core encompasses only a portion of the N-domain$_{39-190}$

sequence. However, the presence of residues undetectable in dipolar-based SSNMR

experiments is not surprising, as this observation has been made for most pathological

and functional amyloids and self-assemblies to date.[55,56]

In light of these findings, we observed that the spectral resolution, as measured by

the ^{13}C line width, is ~60-120 Hz (full width at half-height), which indicates a relatively

well-ordered of the subunits within the complex. These observations are consistent with

our previously reported ^{13}C line widths for *B. subtilis* TasA (~30-100 Hz), *B. subtilis* TapA

(~200 Hz) and *B. cereus* CalY (~200 Hz), which, respectively form well-ordered fibrils

(TasA) and polymorphic aggregates (TapA and CalY) in vitro.[32] To gain more site-specific

insights into the rigid core of the N-domain$_{39-190}$, the protein was incubated with proteinase

K, the bands corresponding to the rigid regions were resolved via SDS–PAGE, and the

amino acid sequences were determined through tandem mass spectrometry analysis

(Figure 5C and D). Interestingly, proteinase K digestion suggested that the following

resistant regions were present: the highly hydrophobic region V^{47}-K^{59}, which contains the

Ala-Ile-Ile segment, and the region A^{88}-K^{134}. Further examination of the SSNMR data

using an additional 200 ms PDSD experiment (Figure S5C) led us to identify several rigid

protein segments. Specifically, these segments correspond to the regions identified

through proteinase K digestion analysis, namely, K^{46}VEA^{49} and E^{93}IKQAISV^{100}. Notably,

the total number of residues identified as resistant to proteinase K was slightly greater

than the number of spin systems identified by SSNMR. This difference suggests that the

core visible in SSNMR contains fewer residues compared to the regions identified as

proteinase K resistant. Alternatively, this observation may result from insufficient spectral

resolution to detect all correlations. Based on these results, it can be inferred that the N-

domain$_{39-190}$ contains a structured core with a significant presence of rigid residues in a

β-sheet conformation.

As the full-length protein exhibits distinct polymerization kinetics from those of the

N-domain$_{39-190}$, we conducted a comprehensive biophysical characterization of full-length

BC1280 to identify putative differences from the truncated N-domain$_{39-190}$ in the context

of macromolecular assemblies. The full-length protein sample underwent a similar

treatment as the N-domain$_{39-190}$ sample, and upon ultracentrifugation, the following

notable differences were observed: i) BC1280 assembled into a transparent gel-like pellet

with high viscosity, while ii) the N-domain$_{39-190}$ pellet aggregated into a non-viscous white

precipitate. Consistent with previous findings for the N-domain$_{39-190}$, the diffraction pattern

of the BC1280 assemblies showed an intense signal at ~4.7 Å and a less pronounced

peak at ~10 Å (Figure S5D). The C-domain of BC1280 consists of fifteen tandem repeats

that were predicted to be disordered; thus, we wondered whether this domain participates

in the overall folding of the full-length protein assemblies. BC1280 was uniformly ^{13}C

labeled, and the resulting aggregates were analyzed via MAS SSNMR. To compare the

molecular conformation of N-domain$_{39-190}$ to that of full-length protein assemblies, we

superimposed 2D ^{13}C-^{13}C SSNMR spectral fingerprints and observed significant overlap,

suggesting that the rigid core of the N-domain is highly conserved in full-length protein

assemblies (Figure S5E). As only a few additional NMR signals were observed in the full-

length protein sample, we hypothesize that the C-domain is not a primary component of

the rigid structural fold of the BC1280 assemblies.

To complement the SSNMR analysis of the rigid BC1280 core, we used dynamics-

based spectral editing via J-coupling polarization transfer to detect dynamic protein

segments[57] and gain insights into any dissimilarities between the two protein versions.

The superimposition of 2D ^{1}H-^{13}C-insensitive nuclei was enhanced by polarization

transfer (INEPT) of the BC1280 and N-domain$_{39-190}$ assemblies, and the results revealed

a significant difference in the spectral fingerprint between the two samples (Figure 5E).

More INEPT signals were observed in the N-domain$_{39-190}$ than the full-length protein,

suggesting that the absence of tandem repetitions in the C-domain results in a more

flexible conformation overall. To confirm this hypothesis, we estimated the degree of

rigidity by comparing signal intensities from 1D ^{13}C-detected INEPT and CP experiments

(Figure 5F), respectively, by probing mobile and rigid protein segments through the

INEPT/CP intensity ratio. We observed a different behavior, with a decrease in mobile

signals in BC1280 (ratio ~0.1) compared to N-domain$_{39-190}$ (ratio ~0.25), suggesting fewer

dynamic regions in the assemblies of the full-length protein.

Considering the differences in the overall rigidity between BC1280 and N-

domain$_{39-190}$, we estimated the relative amount of secondary structural elements using

attenuated total reflection-Fourier transform infrared spectroscopy (ATR-FTIR) analysis.

To identify the minimum peaks within the amide I range according to well-established

assignments,[58] the second derivative was calculated from the raw data, and

deconvolution was performed to determine the percentage of contribution for each

secondary structure type (Figure 5G and S5F, G and H).Overall, due to the additional

structural elements, the ATR-FTIR spectra of BC1280 was more challenging to fit. The

N-domain$_{39-190}$ exhibited 33% antiparallel β-sheets, while the full-length protein had 25%

antiparallel β-sheets and a slightly greater proportion of parallel β-sheets. Both proteins

displayed an equal overall percentage of secondary structure in the random coil

conformation, while BC1280 contained slightly more α-helices and turns than the N-

domain$_{39-190}$.

BC1280 acts as a chaperone for CalY, with its functionality modulated by the C-

domain

The observed in vivo colocalization of BC1280-CalY in the extracellular matrix and the

potential involvement of BC1280 in CalY polymerization led us to ask whether BC1280

might act as an accessory protein that influences CalY assembly. We examined how

BC1280 affects CalY by studying the aggregation kinetics of CalY at different BC1280

molar ratios using ThT fluorescence emission. Surprisingly, we observed that even at

ratios (CalY:BC1280) of 70:30 or 50:50, the polymerization kinetics were significantly

affected compared to CalY incubated alone (Figure 6A). This negative impact, together

with the fact that BC1280 expression is likely tightly regulated, prompted us to investigate

the relative levels of BC1280 and CalY during biofilm formation using an isobaric tag for

relative and absolute quantification (iTRAQ) analysis (Figure 6B). At 24 hours, the relative

accumulation rate of CalY was approximately 3.3 times higher than that of BC1280;

nevertheless, BC1280 accumulated by 48 hours, reaching a ratio (CalY:BC1280) of 2:1.

This information was used to re-estimate the protein ratios for copolymerization assays

of CalY with BC1280. The presence of low concentrations of BC1280 (2 and 10 µM) did

not impact the polymerization kinetics of a 40 µM solution of CalY (Figure 6C). In contrast,

the presence of only the N-domain$_{39-190}$ significantly reduced the rate of polymerization

kinetics of CalY, resulting in a lower fluorescence intensity at the end of the experiment

compared to CalY alone (Figure 6C). In addition, the negative impact on CalY

polymerization was amplified in the presence of non-freshly purified full-length protein or

N-domain$_{39-190}$, which are enriched in preformed oligomers (Figure S6B and C). The

samples corresponding to CalY incubated with a freshly purified 10 µM solution of

BC1280 or N-domain$_{39-190}$ were visualized via TEM. Consistent with the ThT results,

incubation of 10 µM BC1280 with CalY resulted in macromolecular aggregates similar to

those formed by CalY (Figure 6D). However, the incubation of CalY with the N-domain$_{39-}$

$_{190}$ led to the accumulation of small oligomers (Figure 6D), aligning with the previous *in*

vivo results where the absence of the C-domain (*ΔC-domain*) arrested the formation of

CalY filaments (Figure 2F). The ability of BC1280 to regulate the CalY polymerization rate

points to a potential chaperone role for BC1280 in a concentration-dependent manner.

This hypothesis was further supported by circular dichroism analysis, which revealed

significant structural changes in CalY in the presence of BC1280 in a concentration-

dependent manner. After 16 hours of incubation under agitation, a 10 µM solution of

BC1280 induced an increase in the β-sheet conformation of CalY compared to CalY alone

or after incubation with 2 µM BC1280, which predominantly appeared as a random coil

or a disordered conformation (Figure 6E).

The experimental data shown in this work strongly suggest a direct interaction

between BC1280 and CalY, yet the contribution of TasA within the filaments that support

the ECM of *B. cereus* remains uncertain. In vitro co-assembly experiments have

demonstrated that CalY can function as an accessory protein in the assembly of TasA,

without altering its structural arrangement.[32] Moreover, previous findings from our

laboratory indicated that deletions in *tasA* or *calY*, while containing the *bc1280* gene, did

not completely arrest biofilm formation; instead, mutant strains displayed biofilms with

distinct phenotypes.[25] Therefore, we addressed the distribution of their respective

paralogue protein within the ECM in *ΔtasA* and *ΔcalY* mutants compared with the wild-

type strain. Immunochemistry and CLMS analysis revealed fibrillary structures reactive to

anti-TasA antibodies decorating the surfaces of wild-type cells, whereas discrete foci of

TasA fluorescence signals were observed in the ECM of *ΔcalY* (Figure 6F). Remarkably,

in *ΔtasA* mutant strain, CalY protein exhibited a higher degree of fibrillary organization compared to the immunofluorescence signal observed in *B. cereus* wild-type cells (Figure 6F). These results provide evidence that CalY can polymerize into filaments in *ΔtasA* in the presence of BC1280. In contrast, its paralogue TasA doesn't polymerize into filaments *in vivo* when CalY is absent, highlighting the high complexity of the mechanism underlying the ECM assembly in the human pathogen *B. cereus*.

DISCUSSION

Bacterial extracellular filaments are fascinating polymers from structural and biological perspectives. The diversity of structures, sizes, functions and mechanisms of polymerization reflects the specificity of their functions and need for tight regulation. In this work, we pinpointed BC1280 as a specific biofilm orchestrator factor that is exclusive to the *B. cereus* group. In addition, we propose an intriguing molecular mechanism by which this protein could contribute to the assembly of TasA-CalY filaments on the cell surface of this opportunistic human pathogen. The severe defective phenotype observed in *Δbc1280*, when contrasted with the phenotypes of *ΔtasA* and *ΔcalY*, substantiates the indispensable role of BC1280 in biofilm maturation.[25]

While BC1280 could be compared with the chaperone TapA from the phylogenetically related strain *B. subtilis*, notable differences between the two proteins highlight their distinct characteristics. First, BC1280 is exclusive to biofilm cells, whereas TapA is expressed in the planktonic population and from the early stages of biofilm formation. Upon *tapA* deletion, altered and fewer filaments are produced due to a drastic decrease in TasA levels, while the transcriptional expression of *tasA* remains

unaffected.[59] Remarkably, our findings reveal that *tasA* and *calY* levels are

transcriptionally downregulated in *Δbc1280*, resulting in the absence of filaments. It is

important to highlight that in a previous study by our group, we observed that the

heterologous expression of *B. cereus* alleles in the *B. subtilis ΔtapA* strain only reverts

biopellicle formation when *sipW-to-calY*, including *bc1280*, are overexpressed, indicating

only partial functional and mechanistic overlap of TapA and the proteins encoded in *sipW-*

to-calY.[25] Moreover, in the same study, it was demonstrated that in the *B. subtilis ΔtasA*

mutant, in which *tapA* is still expressed, when only *sipW-tasA* or *sipW-calY* from *B. cereus*

are overexpressed without the *bc1280* allele, the biopellicle is not formed, clearly

evidencing that the mechanism of assembly of TasA and CalY is specific of BC1280 and

cannot be replaced by TapA from *B. subtilis*.[25] The idea of a partial mechanistic overlap

between BC1280 and TapA is further corroborated by the structural differences observed

in the predicted model of BC1280 and the solved structure of TapA.[60] While considering

that structures exhibit context-dependent variations, they were partially predicted with

high confidence, supporting the notion that these structures should occur in a specific in

vivo configuration and perform functions such as membrane- or cell-wall interactions.

The results of this study demonstrate that BC1280 contains distinct structural and

functional regions: the N-terminal half, which is sufficient to fulfill the functionality of

BC1280; and the C-terminal half, which appears to be involved in attachment to the cell

surface and also controls the interaction of BC1280 with CalY. After establishing that

BC1280 is an ECM protein and its association with the cell surface, along with its

functional relationship with CalY, we propose its role as a chaperone for CalY based on

the following lines of evidence: i) the biofilm phenotype remains unchanged even after

the overexpression of *tasA* and/or *calY* in the *Δbc1280* mutant strain; ii) filaments of CalY

are not formed in the *ΔC-domain* strain; iii) colocalization of BC1280 and CalY within the

ECM occurs *in vivo;* and iv) BC1280 induces a structural rearrangement of CalY in vitro.

Collectively, these findings prompted us to rename BC1280 as CapP, a CalY-assisting

polymerization protein. In addition to its fundamental role in controlling pilus assembly,

CapP functions as a checkpoint of biofilm maturation by regulating the expression of EPS

via an uncharacterized pair of Clp-proteases and cognate ECF-type sigma factors.

However, this is a novel and specific biofilm regulatory pathway that deserves further

investigation.

We determined in vitro that the N-domain of CapP might trigger or support protein

aggregation, leading to the formation of a rigid core of ~ 40 amino acids enriched in β-

sheets. Through SSNMR, we determined at the atomic level that the rigid core contains

at least the following rigid regions: $K^{46}VEA^{49}$ and $E^{93}IKQAISV^{100}$. The role of each amino

acid stretch remains to be elucidated; nevertheless, we propose that the

$K^{46}VEATLSTAIIFP^{58}$ region significantly influences the aggregation propensity of CapP

in vitro, considering that i) the sequence is situated in the most hydrophobic region of the

protein, ii) the residues within this region are highly conserved, and iii) the region was

detected in the proteinase K-resistant fraction. Although the core is conserved in the full-

length protein, it exhibits slower polymerization kinetics and a greater polydispersity index

than that of the truncated N-domain. In addition, a higher global rigidity is generated by

the C-domain in CapP compared to the truncated N-domain. Together, these findings

strongly suggest that the entire protein has a more complex folding pattern, in which

tandem repetitions may play a role in regulating oligomerization and macromolecular

aggregate assembly. Because the in vitro-generated assemblies could represent only a

fraction of the structures adopted in vivo, the similar aggregation propensity of the

proteins into β-rich filaments supports the hypothesis that such interactions could promote

in vivo structuring.

These findings complement previously conducted structural studies on the

amyloid-like TasA and CalY proteins[32] and expand our knowledge of the coordinated

assembly of these proteins into filaments in the extracellular matrix of *B. cereus*.

Interestingly, CapP adds another level of structural complexity compared to the

orthologous pair TasA and CalY because it contains a C-domain region that is not

involved in the rigid core but could still mediates CalY polymerization. Moreover, CapP

(Figure S5B), similar to TasA and CalY,[32] is disordered in vitro prior to aggregation, a

notable difference from the TasA-TapA system of *B. subtilis*, which adopts a folded or

partially folded conformation before assembly into fibrils.[30,32,61] These distinctions suggest

that this mechanism in *B. cereus* depends on an external trigger, which could be chemical

or biophysical, such as the specific action of accessory proteins. This additional

requirement has been previously proposed for the *in vivo* polymerization of CalY, and

moreover, CalY has been described to enhance the polymerization of TasA without

perturbing the global filament architecture.[22,32] Based on these observations and our

findings, it is reasonable to propose that the three proteins cooperate in assembling

external pili and, consequently, organizing the ECM.

The diverse mechanisms that govern the assembly and attachment of non-flagellar

proteinaceous filaments to bacterial cell surfaces include various structures, such as

chaperone-usher pili, type V pili, type IV pili, curli, fap fibers, conjugative, type IV secretion

pili, e-pili in diderm bacteria, and sortase-mediated pili as well as type IV pili in monoderm

bacteria.[62,63] In the monoderm *B. cereus*, BcpA-BcpB fimbriae are covalently bound to

vegetative cells by sortase, while Ena appendages, exclusively found on spores, are

assembled through subunit crosslinking via disulfide bonding.[64–66] Nevertheless, our work

describes a new type of fibrillar assembly exclusive to the *B. cereus* group, orchestrated

by the chaperone function of CapP, with TasA and CalY serving as principal subunits.

The biphasic effect, dependent on the concentration of CapP in CalY polymerization, has

been reported for several extracellular chaperones.[67] Under conditions where its levels

are lower relative to the substrate, the chaperone acts to stabilize the conformation and

promote filament formation. Conversely, at high concentrations, the chaperone forms

soluble complexes with the substrate, completely inhibiting its assembly. Finally, it is

noteworthy that disordered regions in chaperones are critical for interacting with target

proteins.[68–70] Although we have demonstrated the involvement of the C-domain of CapP

in CalY polymerization, additional investigation is required to elucidate their precise

structural and molecular contacts.

Building upon the novel insights unveiled in this study and the current knowledge

on pili polymerization, we propose a revised model for the assembly of biofilm-specific pili

in *B. cereus* (see Graphical abstract), which involves the paralogous proteins TasA and

CalY, as well as the biofilm-specific protein CapP. Once the switch from a motile to sessile

lifestyle has occurred, further growth and maturation of the biofilm ECM rely upon several

factors, including the products of the operon *sipW-tasA* and the loci *capP* and *calY*, which

are expressed under the tight control of the master regulator SinR. TasA, CapP and CalY

are processed by the cognate signal peptidase SipW and secreted via the Sec pathway

in an unfolded conformation. At 24 hours, the relative levels of CapP compared to CalY

and TasA are lower, and associated with the cell surface, CapP interacts with unfolded

CalY, catalyzing the structural rearrangement necessary to initiate fibrillar polymer growth.

Subsequently, CalY facilitates the incorporation of TasA monomers into growing pili.

When the biofilm reaches a maturation stage at 48 hours, CapP levels rise, and it interacts

with CalY within the ECM to form stable chaperone-substrate complexes that arrest the

elongation of TasA-CalY heteropolymers. The ECM scaffold, composed of TasA-CalY

filaments, exopolysaccharides, and eDNA, will provide *B. cereus* with the chemical and

physical properties necessary for the correct assembly of the microbial community upon

adhesion to biotic or abiotic surfaces.

Our findings contribute to knowledge on the pivotal role of the biofilm-specific

protein CapP in orchestrating the formation and regulation of biofilms on *B. cereus* strains;

in addition, the results provide insights into a new type of pilus assembly mechanism in

monoderm bacteria. These remarkable advancements carry substantial implications for

the clinical and food industries, in which innovative and effective strategies are urgently

needed to address the critical challenge of *B. cereus* biofilm formation and its associated

risks.

ACKNOWLEDGMENTS

We thank Saray Morales Rojas for the technical support. We thank John Pearson from

the Nanoimaging Unit of the Plataforma Bionand, Instituto de Investigación Biomédica de

Málaga (IBIMA), for his technical support in confocal microscopy, data analysis and

sample preparation. We thank Juan Félix López from the Nanoimaging Unit of the

Plataforma Bionand, Instituto de Investigación Biomédica de Málaga (IBIMA), for his

technical assistance in transmission electron microscopy imaging and analysis. We also

thank Josefa Gómez Maldonado from the Ultrasequencing Unit of the SCBI-UMA for RNA

sequencing and Luis Díaz Martínez from the SCBI-UMA for the transcriptomic analysis.

The authors thank Carolina Lobo García and Casimiro Cárdenas García from the

Proteomic Unit of the SCAI-UMA for protein sequencing and MS analysis. We also thank

José Luis Zafra Paredes and Jose María Montenegro Martos from the Vibrational

Spectroscopy Unit and the Electronic Spectroscopy Unit, respectively, of the SCAI-UMA

for their technical support with the ATR-FITR spectroscopy and dynamic light scattering

experiments. We are grateful to Theresa M. Koehler (University of Texas Health Science

Center at Houston) for kindly providing the plasmid pUTE657 and to Stephen Leppla

(National Institute of Health) for kindly providing us with the rat anti-sigmaA.

This work was supported by grants from an ERC Starting Grant (BacBio 637971), the

Plan Nacional de I+D+i of the Ministerio de Ciencia e Innovación (PID2019-107724GB-

I00 and PID2022-141664NB-I00), and Junta de Andalucía (P20_00479). Ana Álvarez-

Mena is the recipient of an FPI contract (BES- 2017-081275). This work benefited from

the support of the French National Research Agency ANR (BH, grant No. ANR-23-CE11-

0005-01), the Biophysical and Structural Chemistry Platform at IECB, CNRS UAR 3033,

and INSERM US001.

AUTHOR CONTRIBUTIONS

D.R. conceived the study. D.R., A.L. and A.A.M. designed the experiments and method

analysis. A.A.M. performed the main experimental work and analyzed the data, with the

following exceptions: A.A.M. and M.B.A.S. analyzed the SS-NMR data; M.L.A.G.

performed some microbiological experiments; A.A.M., M.B. and B.K. performed

biophysical analysis; A.G., E.M., and A.L. performed NMR analysis. JCA contributed to

setting the initial hypothesis and performing iTRAQ experiments. A.A.M., M.B.A.S., B.H.,

A.L. and D.R. prepared the figures. A.A.M., B.H., A.A.L. and D.R. wrote the manuscript.

A.V. contributed critically to writing the final version of the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests

FIGURE TITLES AND LEGENDS

Figure 1. *bc1280* is an essential locus in the developmental program of biofilm formation. A) Schematic domain composition of BC1280. The signal peptide (M1-A39) is shown in red; the domain of unknown function, DUF4047 (I33-E157), which is located within the N-domain, is highlighted in blue; and the C-domain containing the repetitive region (Q190-E 279) is shown in yellow. The amino acid alignment of DUF4047 among different strains of the *B. cereus* group revealed that it is highly conserved, with the residues that are 100% conserved in all the strains marked in color. The color pattern is determined by the amino acid conservation. **B)** Biomasses formed by the *B. cereus* ATCC14579 wild-type strain and the mutants that adhered to abiotic surfaces were evaluated at different time points. Biofilms were stained with crystal violet to enable visualization and subsequent quantitative comparison. **C)** The amount of biofilm mass for each strain was estimated. Three independent experiments were conducted, each with three replicates. Average values are shown, and the error bars represent the standard deviation for each strain. Multiple comparisons were performed at each time point, with the wild-type serving as the control, and ordinary one-way ANOVA was used (** p value < 0.01 and **** p value < 0.0001). **D)** Auto-aggregation kinetics due to cell-to-cell interactions were studied for the wild-type, *Δbc1280* and *ΔC-domain* strains. The optical density at 600 nm at the air-liquid interface of the cell suspensions was measured each hour. The initial optical density was set to 3 at the start of the experiment, and the samples were incubated without agitation at room temperature. The error bars indicate the SDs calculated from three biological replicates.

Figure 2. The biofilm-defective phenotype of *bc1280* is independent of the expression level of *tasA* or *calY.* A) Relative expression levels of *tasA*, *calY*, *eps2* and genes belonging to the *eps1* operon (*bc5263*, *bc5268*, *bc5277* and *bc5279*) obtained by RT–qPCR in *Δbc1280* planktonic cells compared to wild-type cells at 48 hours. The average values of three biological replicates are shown with error bars indicating the standard deviation. **B)** Relative expression levels of *tasA*, *calY*, *eps1*, and *eps2* were measured in planktonic cells at 72 hours in the *Δbc1280* strain that overexpressed *bc2793-bc2794* or *bc2794* and compared to the wild-type strain. Average values from three biological replicates are presented, with error bars representing the standard deviation. **C)** Transmission electron micrographs of the wild-type and *Δbc1280* strains overexpressing *tasA* or *calY*, immunolabeled with specific anti-TasA or anti-CalY antibodies, respectively. Scale bar: 1 μm **D)** Extracellular complementation of *Δbc1280* after the addition of TasA or CalY at 6 μM in monomeric or fibrillar form. The biofilm mass was quantified by crystal violet staining, followed by absorbance measurement at 575 nm and comparison with the *Δbc1280* strain. Statistical analysis was performed using one-way ANOVA with Dunn's multiple comparisons test. Statistical significance is indicated as $p < 0.05$ (*) and $p < 0.01$ (**). **E)** Phenotype reversion of *Δbc1280* upon the addition of 6 μM polymerized TasA or CalY, visualized by immunolabeling with anti-TasA and anti-CalY antibodies, respectively. Scale bar: 1 μm **F)** Negative-stained micrographs of TasA and CalY immunodetection in the *ΔC-domain* strain. Scale bar: 1 μm

Figure 3. BC1280 is a cell wall-associated protein that partially colocalizes with CalY in the ECM. A) and **B)** Left: Immunodetection by Western blotting was performed for BC1280 (A) and the N-domain (B) fused to a histidine tag in different cellular fractions

of *Δbc1280*. This assay was conducted under conditions of overexpression using the

vector pUTE657 and with the addition of 10 µM IPTG. Biofilm samples were grown for 48

hours, after which the resulting biomass was recovered for fractionation into EM

(extracellular medium), CW (cell wall), and C (membrane and cytosol) fractions. The

immunoreactive bands were detected using an anti-His antibody (1:2500). The

immunoblot images were edited by cropping and splicing for better illustration. Right:

Negative stained images of BC1280 (A) and the N-domain (B) in the ECM of the *Δbc1280*

strain. The proteins were immunolabeled using an anti-His antibody (1:100) as the

primary antibody and a goat-antirabbit secondary antibody conjugated with 20 nm

nanogolds (1:100). Scale bar: 500 nm **C)** Immunolocalization of BC1280 and the N-

domain determined using CLSM. Biofilm samples were processed using

immunochemistry, with anti-His antibody (1:100) serving as the primary antibody and a

secondary antibody conjugated to Atto 488. The localization of BC1280 and the N-domain

is shown in green. The cell wall is labeled in red using WGA, and the DNA is stained in

blue using Hoechst. Scale bar: 5 µm **D)** Immunolocalization images of BC1280 and CalY

obtained through CLMS. An anti-His antibody (1:100) and a secondary antibody

conjugated with Atto 488 (1:400) were used for the immunolabeling of BC1280. A specific

primary anti-CalY antibody (1:100) and a secondary antibody labeled with Alexa Fluor

647 (1:400) were used for CalY detection. Additionally, DNA was stained with Hoechst

(1:1000) to visualize the bacteria, which are shown in blue. Scale bar: 3 µm

Figure 4. Heterologously produced BC1280 oligomerizes into macromolecular

complexes driven by the N-domain. A) Scheme illustrating the two constructs based

on the full-length sequence of BC1280. Each construct was fused to a poly-histidine tag

at the C-terminal domain, and the resulting two recombinant proteins were produced

through recombinant expression in *E. coli*. **B)** Size-exclusion chromatogram obtained

after BC1280 was purified under native conditions. **C)** and **D)** Size distribution histograms

obtained via DLS showing the diameters of the aggregates formed by BC1280 for peaks

1 and 2, respectively, as determined by SEC. **E)** Size-exclusion chromatogram obtained

after purifying N-domain$_{39\text{-}190}$ under native conditions. **F)** and **G)** Size distribution

histograms obtained via DLS, illustrating the diameters of the aggregates formed by N-

domain$_{39\text{-}190}$ for peaks 1 and 2, respectively, as determined by SEC. **H)** Native

polyacrylamide gel showing the oligomers formed by BC1280 and N-domain$_{39\text{-}190}$

following native purification through heterologous expression in *E. coli*. The numbers on

each line correspond to the first and second peaks, respectively, that were detected in

the size-exclusion chromatograms of each sample. The gel lanes were cropped and

spliced for illustrative purposes. All the lines superimposed on the image delineate the

lanes and boundaries of the spliced images. **I)** Aggregation kinetics of BC1280 (black

curve) and N-domain$_{39\text{-}190}$ (red curve) determined using ThT fluorescence. **J)**

Transmission electron micrographs of BC1280 and N-domain$_{39\text{-}190}$ at the end of the

polymerization experiment shown in I. Scale bar: 200 nm

Figure 5. The assembled N-domain$_{39\text{-}190}$ contains a rigid core with two β-strand-rich

regions, while the C-domain increases global rigidity. A) X-ray diffraction signals of

N-domain$_{39\text{-}190}$ at 4.7 and 10 Å, indicating interstrand and intersheet spacing, respectively.

B) 2D ^{13}C-^{13}C PDSD experiments at 50 ms for the N-domain$_{39\text{-}190}$. The chemical shifts

that correspond to isoleucine residues in the β-sheet conformation are highlighted. **C)** A

12% SDS–PAGE gel was obtained after the treatment of N-domain$_{39\text{-}190}$ aggregates with

proteinase K for 45 minutes. The three bands marked were analyzed by tandem mass spectrometry. **D)** Determination of the rigid core of N-domain$_{39-190}$ by tandem mass spectrometry. MS data were obtained from the bands selected after digestion of N-domain$_{39-190}$ with proteinase K. The amino acids outlined by a box are the rigid regions identified in the absence of proteinase-K, and letters in red correspond to the residues identified by SSNMR PDSD at 50 and 200 ms. **E)** Superimposition of the ^{1}H-^{13}C INEPT spectra of BC1280 (blue) and N-domain$_{39-190}$ (red), recorded at a 1H frequency of 600 MHz (276 K and 96 scans for BC1280; 300 K and 640 scans for N-domain$_{39-190}$) showing signals corresponding to the flexible part. **F)** Estimation of the global rigidity based on the INEPT:CP ratio of chemical shifts for each sample. **G)** Secondary structure content of the BC1280 and N-domain$_{39-190}$ aggregates probed via ATR-FTIR analysis.

Figure 6. BC1280 acts as a chaperone for CalY, with its functionality modulated by the C-domain. A) Polymerization kinetics of different CalY/BC1280 molar ratios measured by ThT fluorescence emission. Error bars represent the SEM. **B)** Relative levels of BC1280 and CalY in biofilm cells during biofilm formation using iTRAQ. **C)** Kinetics of ThT binding of CalY in the presence of BC1280 or the N-domain$_{39-190}$ at concentrations of 2 and 10 µM. Error bars represent the SEM. **D)** Transmission electron micrographs of negatively stained samples of CalY aggregates in the presence of BC1280 or N-domain$_{39-190}$ at a concentration of 10 µM after 30 hours of incubation at 37°C. Scale bar: 200 nm **E)** Circular dichroism (CD) spectra of CalY incubated for 16 hours under agitation alone or in the presence of BC1280 at 2 and 10 µM. **F)** Immunolocalization of TasA and CalY in wild-type and mutant strains using CLSM with

specific primary antibodies against TasA and CalY, and a secondary antibody conjugated

to Atto 488. Green fluorescence represents TasA and CalY localization, with the cell wall

stained red (WGA) and DNA stained blue (Hoescht). Scale bar: 5 μm

Graphical abstract. During the process of biofilm formation in *B. cereus*, TasA, CapP,

and CalY serve as the main components, in addition to other factors, ensuring that the

extracellular matrix correctly assembles. The three proteins are produced and processed

by the signal peptidase SipW and subsequently secreted through the Sec pathway in an

unfolded conformation. At early biofilm formation stages, CapP is initially present at low

levels, associates with the cell wall and is secreted into the ECM where it interacts with

unfolded subunits of CalY, facilitating CalY folding and initiating fibril growth. CalY serves

as a nucleator for incorporating TasA subunits into the pilus, forming TasA-CalY

heteropolymers. Once the ECM scaffold is established, CapP levels increase, forming

stable complexes with CalY that prevent its folding and arrest filament growth.

SUPPLEMENTAL INFORMATION TITLES AND LEGENDS

Figure S1. ***bc1280* is exclusively found in the *B. cereus* group and is essential for biofilm formation.** Related to Figure 1. **A)** Phylogenetic conservation of BC1280 among *B. cereus* group strains. Phylogenetic analysis was conducted on *B. cereus* group strains using the amino acid sequence of BC1280 (accession number in the KEGG database: BC1280) from *B. cereus* ATCC14579 as the reference. Evolutionary distances were calculated using the neighbor joining method. **B)** Signal peptide prediction by SignalP 5.0. A potential cleavage site was identified for the BC1280 protein between amino acid residues A^{39} and A^{40}. **C)** Amino acid alignment of the C-domain among different strains of the *B. cereus* group, with color indicating conservation level. **D)** The relative expression of *bc1280* in planktonic and biofilm cells of the wild-type strain at 48 hours is presented. The average values from three biological replicates are displayed, with error bars indicating the standard deviation. Statistical analysis was performed using a two-tailed t test, revealing a significant difference between planktonic and biofilm cells (** p value=0.0055). **E)** Top panel: Bands corresponding to the products of amplification using cDNA from RNA extracted from the wild-type strain grown for 48 hours as a template are shown. Bottom panel: To confirm the functionality of the primers, each fragment was amplified via PCR using genomic DNA from the wild-type strain *B. cereus* ATCC14579 as a template. **F)** Biofilm phenotypes formed at the air-liquid interphase by the emetic strain *B. cereus* AH187 wild-type and the *tasA*, *calY* and *bc1280* mutant strains. Biofilm formation was evaluated as the formation of a pellicle at the air-liquid interphase after 3 days of incubation at 28°C.

Figure S2. The genes *tasA*, *calY*, and an uncharacterized ECF sigma factor are

deregulated in the *Δbc1280* mutant at 48 hours. Related to Figure 2. **A)** Biofilm rings

formed after 72 hours by overexpressing *bc2793-bc2794* or *bc2794* in *Δbc1280* at various

IPTG concentrations, as well as in *Δbc2793-bc2794* or *Δbc2794* strains, with wild-type

and *Δbc1280* strains serving as controls. The biofilms were stained with crystal violet, and

the biomass was measured by absorbance at 575 nm. Statistical significance was

estimated using one-way ANOVA with Dunn's multiple comparisons test ($p < 0.05$ (*), p

< 0.01 (**), $p < 0.001$ (***), and $p < 0.0001$ (****). **B)** Biofilm phenotypes of the wild-type

and *Δbc1280* mutant strains were assessed after overexpressing *tasA*, *calY*, or both

genes using the plasmid pUTE657 with 10 µM IPTG for induction. As a negative control,

each strain containing the plasmid was grown without IPTG. Biofilm formation was

visualized by crystal violet staining. The biofilm mass was estimated by absorbance at

575 nm, and statistical significance was assessed by Dunn's multiple comparisons test.

No significant differences were detected between the control strain and upon induction

with IPTG. **C)** Evaluation of the specificity of the nanogold particles in immunolabeling

studies was conducted using the wild-type strain, with any primary antibody and the goat-

antirabbit secondary antibody conjugated with 20 nm nanogolds (1:100). Scale bar: 1 μm

D) Negative control showing TasA and CalY at 6 µM in monomer and fibrillar forms, grown

in TY medium without bacterial inoculation. The wells were stained with crystal violet, and

absorbance at 575 nm was measured to assess biofilm formation.

Figure S3. Biofilm formation is reverted upon overexpression of *bc1280* or *N-*

***domain$_{39-190}$* in *Δbc1280*, and BC1280 colocalizes with CalY but not with TasA.**

Related to Figure 3. **A)** and **B)** Overexpression of *bc1280* or the *N-domain* in the wild-

type and *Δbc1280* strain at 10 and 100 µM IPTG. The resulting biofilms were stained with

crystal violet and the absorbance at 575 nm was measured. **C)** Estimation of the

proportion of fluorescence from co-localized pixels in each channel was performed to

localize BC1280 in the cell wall using the Mander's split coefficient and from five different

fields of view. **D)** Pearson's coefficient (*** p value = 0.0002 using a two-tailed t-test with

Welch's correction) and Mander's split coefficient values obtained from BC1280-CalY

CLSM studies. **E)** For colocalization studies of BC1280 and TasA using CLSM, a specific

primary anti-TasA antibody (1:100) was utilized. Scale bar: 5 µm. **F)** Pearson's coefficient

(*** p value = 0.0002 using a two-tailed t-test with Welch's correction) and Mander's split

coefficient values derived from BC1280-CalY CLSM studies.

Figure S4. Preliminary studies indicate that BC1280 in vitro has a high tendency to

oligomerize, and only the N-domain appears to be folded. Related to Figure 4. **A)**

Mass spectrometry results obtained from the bands that corresponded to the elution

fractions of BC1280 and N-domain$_{39-190}$. **B)** Negative-staining micrographs of the first and

second peaks detected by SEC for BC1280 and N-domain$_{39-190}$. **C)** Prediction of the

folding state using FoldUnfold software. The amino acid residues are classified as folded

(black), folded (blue), or unfolded (red). The underlined amino acids are the regions

identified as putative amyloidogenic by at least two different algorithms. **D)** Hydrophobicity

prediction of residues in BC1280 using ProtScale software on the ExPASy server. **E)** The

PDB model was constructed with the AlphaFold package for BC1280. The color pattern

is displayed based on the confidence value.

Figure S5. **The structural rearrangement of the N-domain is conserved in the full-**

length protein. Related to Figure 5. **A)** Peaks predicted by SPARTA, based on the

structure predicted by AlphaFold, are represented by the cross. The red signal indicates

the experimental data obtained by SSNMR at a mixing time of 50 ms for the N-domain$_{39\text{-}}$

$_{190}$. **B)** Solution NMR ^{15}N-^{1}H spectrum of N-domain$_{39\text{-}190}$ protein demonstrates limited

chemical shift dispersion. **C)** Cross-peak identification for inter-residue interaction.

Overlay of the 2D ^{13}C-^{13}C PDSD correlation spectra obtained by SSNMR for N-domain$_{39\text{-}}$

$_{190}$ at 50 ms (black signals) and 200 ms (red signals) mixing times. The cross-peaks that

correspond to chemical shifts between residues are marked. **D)** X-ray diffraction signals

of BC1280 within 0–15 Å. **E)** Superimposition of the 2D ^{13}C-^{13}C PDSD correlation spectra

obtained by SSNMR for BC1280 (blue) and N-domain$_{39\text{-}190}$ (red) at a mixing time of 50

ms. **F)** Second derivative analysis of BC1280 (black) and N-domain$_{39\text{-}190}$ in the amide I

band using ATR-FTIR. **G)** and **H)** ATR-FTIR spectra of BC1280 and N-domain$_{39\text{-}190}$,

respectively, in the amide I band. The contribution of each peak was determined through

secondary derivative analysis of the raw data, followed by a deconvolution approach.

Figure S6. BC1280 levels have an impact on CalY polymerization. Related to Figure

6. **A)** Estimation of the relative levels of TasA, BC1280, and CalY in biofilm cells over 72

hours using iTRAQ. **B)** and **C)** Study of the effect of BC1280 and the N-domain39-190

without recent purification on CalY polymerization. ThT binding kinetics of CalY in the

presence of freshly and non-freshly purified BC1280 (B) or N-domain39-190 (C) samples.

Error bars indicate the standard error of the mean. **D)** Circular dichroism (CD) spectra

obtained for each sample without subtracting the contribution of BC1280 at 2 and 10 µM

when incubated with CalY for 16 hours under agitation.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Diego Romero (diego_romero@uma.es).

Materials availability

Plasmids generated in this study are available at the BacBio Lab. Please contact Diego Romero (diego_romero@uma.es) for access. This study did not generate new unique reagents.

Data and code availability

RNaseq data have been deposited at GEO database and are publicly available as of the date of publication. Accession numbers are listed in the Key Resources Table. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD010211. Microscopy data reported in this paper will be shared by the lead contact upon request.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDENT PARTICIPANT DETAILS

Bacterial strains and culture conditions

The bacterial strains used in this study are listed in the Key Resources Table. For experiments conducted using *B. cereus* wild-type or mutant strains, cultures were routinely grown at 28°C from frozen stocks on Luria–Bertani (LB) plates containing 1%

tryptone (Oxoid), 0.5% yeast extract (Oxoid), 0.5% NaCl and 1.5% agar before the

respective experiments were performed. Biofilm assays were performed in TY medium

supplemented with 1% tryptone (Oxoid), 0.5% yeast extract (Oxoid), 0.5% NaCl, 10 mM

$MgSO_4$ and 1 mM $MnSO_4$.

For cloning and plasmid replication, the *E. coli* DH5α strain was used. For

recombinant protein expression and purification, *E. coli* BL21(DE3) pLysS (Merck,

Kenilworth, NJ, USA) was used. All the strains were cultured in Luria–Bertani (LB) liquid

media. For the agar-solid plates, LB media was supplemented with 1.5% bacteriological

agar (Oxoid). The final antibiotic concentrations were 100 μg/ml ampicillin, 50 μg/ml

kanamycin, 100 μg/ml spectinomycin, and 5 μg/ml erythromycin.

For the purification of $^{13}C/^{15}N$-labeled proteins, the cultures were grown in M9

minimal medium (the final concentration was 48 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.6 mM

NaCl, 1 mM $MgSO_4$, 0.01 mM $ZnCl_2$, 0.001 mM $FeCl_3$, 0.1 mM $CaCl_2$ and 10 ml 100X

minimum essential medium vitamin solution) supplemented with 1 g/L $^{15}NH_4Cl$ as nitrogen

and 2 g/L $^{13}C_6$-D-glucose or 2-^{13}C glycerol as the carbon source.

METHOD DETAILS

Plasmid and strain construction

All the primers used to construct the strains used in this study are listed in Table S4.

B. cereus mutant strains were constructed via homologous recombination using

the upstream and downstream regions of the gene of interest, which were subsequently

cloned and inserted into the pMAD plasmid.[71] The constructs were designed using NEB

Builder HiFi DNA Assembly Master Mix (New England Biolabs, MA, USA) following the

manufacturer's instructions and utilizing specific primers. The pMAD vector was linearized

using the restriction enzyme SmaI (FastDigest, Thermo Fisher Scientific), and the

digested vector was incubated at 50°C for 1 hour along with the respective upstream and

downstream fragments. The combined total amount of fragments and plasmid was 0.2

pmol, and a proportion of 1:2 (vector: fragments) was maintained. The resulting reaction

mixture was subsequently transformed into *E. coli* DH5α, and positive colonies were

selected using ampicillin. Subsequently, the plasmids were purified, subjected to PCR

amplification and digestion for verification, and sequenced.

For the overexpression experiments in *B. cereus,* the gene of interest was cloned

and inserted into the plasmid pUTE657[72] under a promoter that was inducible by IPTG

using the primers listed in Table S4. Following gene amplification and the inclusion of a

specific RBS for *Bacillus*, the resulting PCR product was digested with SalI and SphI

(FastDigest, Thermo Fisher Scientific) and subsequently cloned and inserted into the

pUTE657 plasmid that was cut with the same restriction enzymes using T4 DNA ligase

(Thermo Fisher Scientific). Subsequently, the resulting reaction was transformed into *E.*

coli DH5α, and colonies were selected with 100 µg/ml ampicillin. After plasmid

purification, all the vectors were subjected to PCR amplification, digestion with restriction

enzymes, and sequencing.

The vectors were transformed into *B. cereus* by electroporation following

previously described methods with some modifications. A single colony from a pure

culture of *B. cereus* was inoculated in 5 ml of LB medium and incubated under shaking

conditions at 30°C for 16 hours. This culture was subsequently used to inoculate a flask

containing 100 ml of LB, which was incubated at 30°C with agitation until an OD_{600nm} of

0.3 was reached. The cell culture mixture was centrifuged (3000 rpm, 5 minutes, 4°C),

and the resulting pellet was resuspended in 12 ml of electroporation buffer (0.5 mM

K_2HPO_4, 0.5 mM KH_2PO_4, 0.5 mM $MgCl_2$, and 272 mM saccharose). The suspension was

centrifuged again and then resuspended in 250 µl of electroporation buffer. The

electrocompetent cells were mixed with 0.5–1 µg of plasmid and incubated on ice for 10

minutes. Next, the mixture was electroporated in a 0.2 cm cuvette using a voltage of 1.4

kV, a capacitance of 25 µF, and a resistance of 200 Ω. After electroporation, the

suspension was recovered and incubated with 1.5 ml of LB for 5 hours at 30°C with

agitation. The resulting culture was plated onto LB plates supplemented with the

corresponding antibiotic to select the colonies that had been successfully transformed

with the vector. For integration of the deletion, *B. cereus* transformants were incubated

without antibiotics at 40°C, facilitating the integration of the mutation into the genome

through homologous recombination. Finally, the culture was plated on LB agar plates,

and the colonies were picked onto LB agar plates supplemented with erythromycin.

Colonies that did not grow on LB supplemented with the antibiotic were selected, and the

mutation was confirmed through PCR amplification.

For the heterologous expression of BC1280, the N-domain$_{39-190}$ and the C-

domain$_{191-279}$ in *E. coli,* the open reading frames containing the *bc1280* gene, the region

from amino acids 39 to 190. The PCR products and the pET24b vector were digested

with NdeI and XhoI (FastDigest, Thermo Fisher Scientific) and subsequently cloned using

T4 DNA ligase (Thermo Fisher Scientific) following the instructions supplied by the

manufacturer. The reaction mixtures were subsequently transformed into *E. coli* DH5α,

after which the colonies were successfully transformed with 50 µg/ml kanamycin.

Phylogenetic tree

The amino acid sequence of BC1280 was obtained from the KEGG database using *B. cereus* ATCC14579 as a reference. BLASTP analysis was performed, and the top 50 hits were aligned using Clustal Omega.[73] The phylogenetic tree was then constructed using neighbor joining distances.

RT–qPCR

Quantitative reverse transcription PCR was used to estimate the transcription levels of i) *bc1280* in planktonic and biofilm cells of the wild-type strain; ii) *tasA*, *calY*, *eps1* and *eps2* in *Δb1280* compared to those in the wild-type; and iii) *tasA*, *calY*, *eps1* and *eps2* in *Δb1280* overexpressing *bc2794* or *bc2793-2794* compared to those in the wild-type. RT–qPCR was performed using an iCycler-iQ system (Bio-Rad) and Power SYBR Green Master Mix (Thermo Fisher Scientific). Primer3 software (https://primer3.ut.ee/) was used for primer design, maintaining the default parameters.[74] First, 1 µg of total RNA was reverse transcribed into cDNA using SuperScript III reverse transcriptase (Invitrogen) following instructions provided by the manufacturer. The reactions were performed in triplicate in 96-well plates with a total volume of 20 µl. The RT–qPCR cycle started with an initial denaturation at 95°C for 3 minutes, followed by 40 amplification cycles (95°C for 20 seconds, 56°C for 20 seconds, and 72°C for 30 seconds), and the final step was 95°C for 30 seconds. The target genes, including *bc1280*, which encodes the protein BC1280; *bc1279*, which encodes TasA (also known as spore coat-associated protein N); *bc1281*, which encodes CalY (annotated as cell envelope-bound metalloprotease, camelysin); *bc5263*, which encodes UDP-glucose 4-epimerase; *bc5268*, which encodes secreted

polysaccharide polymerase; *bc5277*, which encodes tyrosine-protein kinase; *bc5279*,

which encodes tyrosine-protein kinase; and *bc1583*, which encodes O-acetyl transferase,

were amplified using the primer pairs provided in Table S5. Primer efficiency and

amplification product specificity were assessed as previously described.[75] The estimation

of relative expression levels was calculated using the $\Delta\Delta$Ct threshold (Ct) method.[76] The

housekeeping gene *rpoA* was used as a reference for data normalization. The relative

expression value was calculated as the difference between the qPCR threshold cycle (Ct)

of the gene of interest and the Ct obtained for *rpoA* ($\Delta Ct = Ct_{target\ gene} - Ct_{rpoA}$). The results

obtained for the genes of interest were normalized to the values obtained for the wild-

type strain. The RT–qPCR analyses were conducted three times, each with three

independent biological replicates.

1020

Determination of the transcriptional unit by PCR

To determine whether *bc1280* constitutes an operon with *tasA* or *calY*, primers were

designed between the genes and inside each gene to amplify four specific fragments that

were 974 bp, 572 bp, 1668 bp, and 560 bp in size (primers listed in Table S6). cDNA was

used as the template, and PCR was performed using genomic DNA purified with the

commercial JetFlex™ Genomic DNA Purification Kit (Thermo Fisher Scientific, Bremen,

Germany) as a positive control. The fragments were amplified by PCR using Phusion™

High-Fidelity DNA Polymerase (Thermo Fisher Scientific) with the following program:

initial denaturation at 98°C for 30 seconds, followed by a 30-cycle amplification program

(98°C for 10 seconds, 56°C for 30 seconds, and 72°C for 1 minute), and a final extension

step at 72°C for 5 minutes. The samples were loaded onto an agarose gel and compared

to the fragments amplified from genomic DNA.

1033

Biofilm formation and extracellular complementation assays

To study the biofilm formation phenotype for each strain or condition, bacterial cultures

were grown in 24-well plates as previously described.[25] The strains were subsequently

grown on LB agar plates at 28°C for 24 hours. The bacterial mass was resuspended in 1

ml of TY medium, and the OD_{600} was adjusted to 1. Subsequently, 10 μl of the suspension

was inoculated into each well containing 1 ml of TY medium. The plates were then

incubated at 28°C without agitation for 72 hours.

For the extracellular complementation assay involving the addition of TasA or CalY

as monomers or in their fibrillated states, the proteins were previously purified *in vitro* from

inclusion bodies through heterologous expression in *E. coli* following a protocol

established by El-Mammeri et al., in 2019.[32] Subsequently, the proteins were dialyzed

against a 20 mM Tris and 50 mM NaCl solution at pH 7.4, after which the protein

concentrations were measured. The assembly of filaments was promoted at 37°C under

agitation at 200 rpm for one week. TasA or CalY protein, in both its monomeric and

fibrillated forms, was added at a final concentration of 6 μM to 24-well plates inoculated

with the *Δbc1280* strain. These plates were then incubated without agitation at 28°C for

72 hours, after which the resulting biomass was stained with crystal violet.

1051

Crystal violet adhesion assay

Bacterial adhesion to the well plate surface was quantitatively measured using crystal

violet staining, as previously described.[77] The biofilms were grown for 72 hours, after

which the medium was removed. Then, 1 ml of 1% crystal violet was added to each well

of the 24-well plate, and the cells were incubated for 15 minutes. The plates were washed

multiple times with water and left to dry for 45 minutes. Next, the biomass that was formed

in each well was resuspended in 1 ml of 50% acetic acid, and the absorbance at 575 nm

was measured for each sample using a plate reader (Tecan Infinite M1000 Pro; Tecan,

Männedorf, Switzerland). The statistical significance was evaluated using one-way

ANOVA with Dunnett´s multiple comparisons test. P values less than 0.05 were

considered to indicate statistical significance.

1063

Auto-aggregation assay

The effect of *bc1280* and *C-domain* deletion on the aggregation rate of *B. cereus* was

estimated using a kinetic method following a previously described protocol.[24] The strains

were inoculated into a flask containing 20 ml of TY medium and incubated overnight at

28°C with agitation. The OD_{600} was adjusted to 3 for a final volume of 10 ml, after which

the supernatant was centrifuged (5000 rpm, 10 minutes, 20°C) for dilution. The samples

were then incubated in vertical tubes under static conditions at room temperature for 24

hours, after which the OD_{600} was measured every hour at the air–liquid interface.

1072

RNA extraction

For RNA extraction, a previously described protocol was followed with several

modifications.[26] Bacterial strains were cultivated in 24-well plates to retrieve the desired

population—either planktonic or biofilm cells—at a specific incubation time. For the biofilm

cells, the supernatant was removed from each well, followed by three wash steps, and

the biomass was subsequently resuspended in 1 ml of PBS. To minimize variability

between replicates, each sample was prepared by extracting 500 µl from 8 distinct wells.

The bacterial suspensions were then centrifuged at 12000 × g for 5 minutes, and the

resulting pellets were frozen at -80°C for 30 minutes or until needed. The pellets were

resuspended in 900 µl of TRI-Reagent (Merck), and the cells were disrupted with 0.1 mm

beads using a TissueLyser (QIAGEN) for 3 minutes, followed by incubation at 55°C for 3

minutes. Then, 200 µl of chloroform was added to each tube, which was vortexed for 10

seconds and incubated at room temperature for 3 minutes. Finally, the samples were

centrifuged at 12000 × g for 10 minutes at 4°C. The resulting aqueous phase was

transferred to a new tube containing 500 µl of ice-cold isopropyl alcohol. The tubes were

inverted a few times, incubated for 10 minutes at room temperature, and then centrifuged

at 12000 × g for 10 minutes at 4°C. The supernatants were subsequently removed, and

the pellets were washed with 1 ml of ice-cold 75% ethanol, followed by centrifugation

(12000 × g, 5 minutes, 4°C). Briefly, the RNA pellets were dried and then resuspended in

50 µl of DEPC-treated water. The residual DNA was eliminated through treatment with

rDNAse, which is included in the Nucleo-Spin RNA Plant Kit (Macherey-Nagel), following

the instructions provided by the manufacturer. The quality and integrity of the total RNA

were assessed using an Agilent 2100 bioanalyzer (Agilent Technologies) and gel

electrophoresis.

1097

Whole-transcriptome analysis

A RNA-Seq analysis was performed on *Δbc1280* planktonic cells in comparison to the

wild-type after 24 and 48 hours of incubation at 28°C. RNA extraction was performed

following a previously described protocol, and each sample corresponding to a specific

condition was collected in duplicate. RNA sequencing was performed by the Omics Unit

of Supercomputing and Bioinnovation Center (SCBI, University of Malaga, Spain). rRNA

removal was achieved using the RiboZero rRNA Depletion Kit (Illumina, CA, USA), and

subsequently, 100-bp single-end read libraries were prepared using the TruSeq Stranded

Total RNA Kit (Illumina). Next, the libraries were sequenced using the NextSeq550

instrument (Illumina). To eliminate regions of low quality, ambiguity, and low complexity,

as well as potential contamination, the raw reads were preprocessed using SeqTrimNext

with the specific configuration parameters used for NGS technology.[78] Subsequently, the

clean reads were aligned and annotated using the *B. cereus* ATCC14579 genome

(NC_004722.1) as the reference. This alignment was conducted with Bowtie2, resulting

in BAM files that were subsequently sorted and indexed using the SAMtools v1.484

program.[79,80] To calculate the read count for each gene, the script Sam2counts

(https://github.com/vsbuffalo/sam2counts) was used to determine the number of uniquely

mapped reads. The analysis of differentially expressed genes (DEGs) between *Δbc1280*

and the wild-type strain was conducted using the R script DEgenes Hunter. This script

employs a combined p value, utilizing Fisher´s method to consider the nominal p values

obtained from edgeR and DEseq2.[81,82] The combined p value was adjusted using the

Benjamini–Hochberg (BH) test, which employs the false discovery rate approach. The

adjusted p value was subsequently used to rank all the DEGs obtained. A $p < 0.05$

and a log2FoldChange < -1 or > 1 were considered to indicate statistical significance.

Transmission electron microscopy and immunochemistry studies

To investigate the location of the proteins within the ECM using transmission electron

microscopy, we followed a protocol that was previously established and described, with

several modifications.[25,29,32] For TasA and CalY immunodetection, the strains were

cultured in multi-well plates at 28°C for 72 hours. Alternatively, for the localization of

BC1280 or the N-domain within *B. cereus* cells, the strains corresponding to *Δbc1280*

containing the plasmids pUTE657-*bc1280-6xHis* or pUTE657-*N-domain-6xHis* were

grown in multiwell plates with TY medium supplemented with a 10 µM IPTG solution. The

biofilm mass was subsequently harvested in 1 ml of PBS after 48 hours.

Then, 20 µl of each sample was applied to a copper grid and incubated for 2 hours.

The grids were subsequently washed with PBS and incubated for 5 minutes to remove

any excess bacterial cells. The cells that adhered to the grid were fixed using a 2%

paraformaldehyde solution diluted in PBS and incubated for 10 minutes. The grids were

washed again with PBS for 5 minutes and then blocked for 30 minutes using Pierce™

Protein-Free T20 (TBS) Blocking buffer (Thermo Scientific). Next, the samples were

incubated for 2 hours with the primary antibody at a concentration of 1:100, which was

diluted in blocking buffer. Specific primary antibodies against TasA and CalY were used

to detect TasA and CalY, respectively. For the immunolocation of BC1280-His and N-

domain-His, we employed the primary antibody anti-6-His produced in rabbits (Merck).

Subsequently, the grids were washed twice with TBS-T, each time with a 5-minute

incubation. The samples were incubated for 1 hour with a goat anti-rabbit 20-nm gold-

conjugated secondary antibody (Ted Pella, Redding, CA, USA) at a dilution of 1:100 in

blocking buffer. Next, the grids were subjected to two additional washes with TBS-T, each

lasting 5 minutes. The samples were fixed with 2.5% glutaraldehyde for 10 minutes. The

samples were washed with Milli-Q water for 5 minutes and subsequently negatively

stained with a 1% uranyl acetate solution. Finally, the grids were rinsed with a single drop

of water and dried under dark conditions.

As a negative control to check the specificity of the 20-nm gold secondary

antibody, we performed the same protocol, but instead of adding the primary antibody,

we added blocking buffer at that step.

The samples were examined using a Thermo Fisher Scientific Tecnai G^2 20 TWIN

transmission electron microscope at an accelerating voltage of 120 kV. The final images

were taken using an Olympus Veleta side-mounted CCD with a resolution of 2k × 2k Mpx.

1155

Cellular fractionation

For protein immunodetection by Western blotting, the samples were fractionated into

three components: the extracellular medium, cell wall-associated proteins, and cellular

contents (membrane and cytosol). Preliminary studies of BC1280 and the N-domain were

conducted through cellular fractionation using the *Δbc1280* strain harboring the plasmid

pUTE657-*bc1280*-6xHis or pUTE657-N-*domain*-6xHis, respectively. The strains were

subsequently grown in TY media supplemented with agitation at 28°C. When the cultures

reached an OD$_{600nm}$ of 0.4, 10 µM IPTG was added to each sample, and the cultures

were incubated for 4 hours.

All the samples were subjected to the following protocol. The samples were

centrifuged at 6000 × g for 5 minutes, after which the supernatant was passed through a

0.45 µm PES filter and retained as the extracellular medium fraction. The pellets were

resuspended in 10 ml of PBS containing 100 µg/ml lysozyme and incubated for 2 hours

at 37°C. Subsequently, the samples were centrifuged at 9000 × g for 20 min at 4°C, after

which the resulting supernatant and the pellet were retained as the cell wall and cellular

fractions, respectively. The cellular fraction was then resuspended in a small volume of

PBS. Finally, all the samples were precipitated with trichloroacetic acid (TCA) to a final

concentration of 10%. The mixture was then incubated on ice for 1 hour. Next, the

precipitated proteins were centrifuged at 13000 × g for 20 minutes at 4°C, and the

resulting pellets were washed twice with 1 ml of acetone. Finally, the samples were dried

at 37°C for 5 minutes, resuspended in 1x Laemmli buffer (Bio-Rad), and loaded onto a

12% SDS–PAGE gel for the immunodetection assay.

1178

Polyacrylamide gels and western blotting

The samples were diluted in 1x Laemmli buffer (Bio-Rad) and then heated at 100 °C for

5 minutes. The proteins were separated by 12% SDS–PAGE with the Spectra™

molecular weight marker (Thermo Fisher Scientific) and subsequently transferred to a

polyvinylidene fluoride (PVDF) membrane (Bio-Rad) using the Trans-Blot Turbo Transfer

System (Bio-Rad) at 25 V for 30 minutes. Next, the membrane was blocked for 1 hour

using 5% nonfat milk diluted in 50 mM Tris-HCl (150 mM NaCl, pH 7.5) containing 0.1%

Tween-20 (TBS-T). The membrane was then incubated with the primary antibody in a

solution of 3% nonfat milk in TBS-T. The membrane was washed three times with TBS-

T, with 10 minutes of incubation between each wash. Next, the membrane was incubated

with the secondary antibody against rabbit IgG conjugated to horseradish peroxidase

(Bio-Rad) for 2 hours at a concentration of 1:3000 and diluted in TBS-T. Following the

incubation, the membrane was washed twice with TBS-T and once with TBS. Finally, for

immunodetection, the membranes were exposed to Pierce ECL Western blotting

Substrate (Thermo Fisher Scientific).

For the immunodetection of BC1280-His or N-domain-His, we used an anti-6-His

(Merck) primary antibody produced in rabbits at a dilution of 1:2500.

To study the molecular weight of the oligomers formed by BC1280 and N-domain$_{39-}$

$_{190}$ under native conditions, the proteins were diluted in native sample buffer (Bio-Rad),

loaded onto polyacrylamide gels (Any kD™ Mini-PROTEAN® TGX™ Precast Protein

Gels, Bio-Rad), and analyzed using the NativeMark^{TM} (Invitrogen) marker as a reference.

The native gels were run in buffer containing 25 mM Tris and 192 mM glycine at a

constant voltage of 200 V. To visualize the proteins, the gels were stained with Coomassie

brilliant blue.

1203

Immunolocalization studies by fluorescence microscopy

For the immunofluorescence experiments, the samples were grown as previously

described for transmission electron microscopy studies. Then, 150 µl of each sample was

added to well slides coated with 0.1% poly-L-lysine (Sigma–Aldrich) and incubated for 2

hours to facilitate bacterial adhesion to the substrate. The sample was then removed, and

the bacteria that had attached to the slide surface were fixed with fixation buffer (3%

paraformaldehyde and 0.1% glutaraldehyde diluted in PBS) for 10 minutes. Briefly, the

wells were washed twice with PBS, and the samples were incubated for 1 hour with

blocking buffer (3% m/v bovine serum albumin (BSA) and 0.2% v/v Triton X-100 in PBS).

Then, the buffer was removed, and the wells were incubated for 3 hours with the primary

antibody at a concentration of 1:100 diluted in blocking buffer. The primary antibodies

used for visualization by TEM were also used for this experiment. The wells were then

rinsed three times with washing buffer (0.2% m/v BSA and 0.05% v/v Triton X-100 in

PBS), and each incubation lasted for 5 minutes. Next, the wells were incubated for 2

hours with the secondary antibody goat anti-rabbit IgG-Atto488 (manually labeled) at a

dilution of 1:400 in blocking buffer. The samples were washed once with washing buffer

and twice with PBS, for an incubation period of 5 minutes each. The immunostainings

were fixed for 5 minutes using fixation buffer, and the wells were subsequently rinsed with

PBS three times. To detect the cell wall, the slide was incubated with wheat germ

agglutinin labeled with Alexa Fluor 647 (WGA, Thermo Fisher Scientific) for 20 minutes

at a dilution of 1:100 in PBS. Finally, the bacterial DNA was stained for 20 minutes using

Hoechst at a dilution of 1:1000. As a negative control, immunostaining was performed

without incubation with the primary antibody.

The immunostaining was visualized using confocal laser scanning microscopy. For

the fluorescence corresponding to Atto-488 visualization, we employed an excitation

wavelength of 488 nm and detected the emission between 497 and 572 nm. The signal

corresponding to Alexa Fluor 647 was visualized with an excitation wavelength of 561

nm, and emission was detected between 576 and 686 nm. The Hoescht signal was

visualized independently using a specific dichroic filter.

For colocalization studies of BC1280-CalY/TasA, the same immunostaining

protocol was followed with a few modifications. First, BC1280-6xHis was labeled using

the anti-6-His antibody produced in rabbits (Merck), followed by goat anti-rabbit antibody

conjugated to Atto-488 as the primary and secondary antibody, respectively. After three

washes were performed with washing buffer, the samples were treated with quenching

solution (0.1 M glycine in PBS) for 10 minutes. Next, TasA or CalY were detected using

the following specific primary antibodies: anti-TasA and anti-CalY, which were produced

in rabbit, followed by anti-rabbit conjugated to Alexa-Fluor 647 produced in goat as the

secondary antibody. Following the final steps, the samples were fixed, stained with

Hoechst, and finally visualized via CLSM, as described previously.

The colocalization images were analyzed using the software ImageJ and the

colocalization threshold plugin. The region of interest (ROI) was defined for the region

corresponding to Alexa Fluor 647, and both Pearson´s and Mander´s coefficients were

calculated.[83] As a negative control, the band corresponding to Alexa Fluor 647

overlapped with itself, resulting in 100% colocalization. For each sample, a minimum of 5

different images were analyzed, and statistical significance was determined for Pearson´s

coefficient using an unpaired t test with Welch´s correlation, with a p value < 0.005.

1250

Protein expression and purification

BC1280 and N-domain$_{39-190}$ were purified under native conditions. For the heterologous

expression of BC1280, the plasmid pET24b-*BC1280* was transformed into *E. coli* BL21

(DE3) (Merck, Kenilworth, NJ, USA), and colonies were selected with 50 µg/ml

kanamycin. The plasmid pET24-*N-domain$_{39-190}$* was subsequently transformed into *E. coli*

Lemo21 (DE3)pLyss (New England Biolabs, USA) for N-domain$_{39-190}$ expression, after

which colonies were selected with 50 µg/ml kanamycin and 5 µg/ml chloramphenicol. The

following steps were performed for both proteins. A single fresh colony was picked and

cultured in 10 ml of LB supplemented with the corresponding antibiotics mentioned

previously. The preculture was incubated under shaking conditions for 6 hours at 37°C.

A total of 10% (v/v) of the precultures were inoculated into 1 L of LB supplemented with

the corresponding antibiotics. The culture was incubated at 37°C with agitation until the

optical density reached 0.6. For BC1280, the culture was induced with 400 μM IPTG and

incubated for 16 hours at 28°C. Otherwise, the culture corresponding to the

overexpression of N-domain$_{39-190}$ was induced with 400 μM IPTG and 100 μM rhamnose

and incubated for 3 hours at 28°C. Then, the cells were harvested (6000 × g, 30 minutes

at 4°C, JLA 8.1 rotor, Beckman Coulter, Brea, CA, USA), and the pellets were frozen at -

80°C until use.

The pellets were resuspended in 18 ml of buffer A (20 mM Na$_2$PO$_4$, 500 mM NaCl

and 20 mM imidazole, pH 8) with cOmplete™ EDTA-free Protease Inhibitor Cocktail

(Roche) and CelLytic^{TM} B-Cell Lysis reagent (Merck). The lysates were incubated for 30

min with agitation at room temperature. The samples were sonicated using a Branson

450 digital sonifier on ice with 3 pulses for 45 seconds at 40% amplitude and then

centrifuged at 15000 × g for 60 minutes at 4°C (F34-6-38 rotor; Eppendorf, Hamburg,

Germany). The supernatant was loaded onto a HisTrap HP 5 ml column (GE Healthcare)

and purified using an AKTA Start FPLC system (GE Healthcare). The column was

preequilibrated with buffer A, and after sample loading, it was washed with the same

buffer. Next, the proteins were eluted using a linear gradient of buffer B (20 mM Na$_2$PO$_4$,

500 mM NaCl and 500 mM imidazole, pH 8). Since the elution mixture was not completely

pure, the sample was loaded onto a HiLoad 16/600 Superdex 75 pg column (Cytiva), and

the pure protein was subsequently eluted in buffer containing 20 mM Tris and 50 mM

NaCl, pH 7.5. The elution fractions were loaded onto a 12% Tris-Tricine SDS–PAGE gel,

and the samples containing the purified protein were concentrated using Amicon Ultra15

centrifugal filter units (Millipore) with a 3 kDa cutoff. Subsequently, the purified fractions

were separated via 12% SDS–PAGE to assess their purity, which was determined

through Coomassie blue staining. Finally, the gel bands were analyzed using tandem

mass spectrometry.

CalY and TasA were purified via heterologous expression in *E. coli* BL21(DE3)

pLyss and from inclusion bodies under denaturing conditions following the protocol

established by El Mammeri et al. [32]

1291

Tandem mass spectrometry analysis of protein bands

The SDS–PAGE gel bands that corresponded to the protein purification and proteinase

K digestion experiments were subjected to analysis via tandem mass spectrometry using

a nanoion trap system [HPLC–electrospray ionization–tandem mass spectrometry (MS)].

The bands were cut and destained using a mixture of 50% acetonitrile (ACN) and 25 mM

ammonium bicarbonate. Then, the samples were dehydrated and dried using ACN.

Disulfide bridges were reduced with 10 mM dithiothreitol (DTT) diluted in 50 mM

ammonium bicarbonate, followed by incubation at 56°C for 30 minutes. Subsequently,

the excess DTT was removed, and the cysteine residues were carbamidomethylated

using 55 mM iodoacetamide diluted in 50 mM ammonium bicarbonate for 20 minutes at

room temperature in the dark. The gel bands were then dehydrated again, and the

proteins were digested using 10 ng/μl trypsin (Promega) at 30°C overnight. To extract the

peptides from the gel, the samples were incubated with a solution of 0.1% ACN/formic

acid (FA) for 30 minutes at room temperature. To eliminate ACN and ammonium

bicarbonate, the samples were dried using a SpeedVacTM. Next, the samples were

resuspended in a solution containing 0.1% FA, ultrasonicated for 3 minutes, and

centrifuged at 13,000 × g for 5 minutes. Subsequently, the samples were purified and

concentrated using C18 ZipTip® (Merck) following the instructions supplied by the

manufacturer. Finally, the samples were injected into an Easy nLC 1200 UHPLC system,

which was coupled to a Q Exactive HF-X Hybrid Quadrupole-Orbitrap mass spectrometer

(Thermo Fisher Scientific). The software versions used for acquisition and analysis were

Tune 2.9 and Xcalibur 4.1.31.9. The mobile phases used in the HPLC consisted of i)

buffer A, which contained 0.1% FA dissolved in water, and ii) buffer B, which contained

0.1% FA dissolved in 80% ACN. Peptides were loaded onto a precolumn (Acclaim

PepMap 100, 75 µm × 2 cm; C18, 3 µm; 100 A; Thermo Fisher Scientific) at a flow rate

of 20 µl/min, while elution was performed using a 50 cm analytical column (PepMap RSLC

C18, 2 µm; 100 A, 75 µm × 50 cm; Thermo Fisher Scientific). Elution was carried out

through a gradient concentration over 60 minutes, transitioning from 5% to 20% buffer B.

This was followed by a 5-minute gradient from 20% to 32% buffer B, resulting in a 10-

minute elution with 95% buffer B. Then, the column was equilibrated with 5% buffer B

using a constant flow rate of 300 µl/min. Before sample analysis, an external calibration

was performed using LTQ Velos ESI Positive Ion Calibration Solution (Pierce, IL, USA),

along with internal calibration using the polysiloxane ion signal at m/z 445.120024

obtained from ambient air. The MS1 scans were conducted within a m/z range of 375-

1.600 at a resolution of 120.000. Employing a data-dependent acquisition strategy, the

15 most intense precursor ions with a charge ranging from +2 to +5 within a window of

1.2 m/z were selected from fragmentation to generate the corresponding MS2 spectra.

The fragmentation ions were generated through high-energy collision-induced

dissociation with an initial mass set at 110 m/z and subsequently detected using a mass-

analyzed Orbitrap at a resolution of 30.000. The dynamic exclusion for the selected ions

was set to 30 seconds, and the maximum accumulation times for MS1 and MS2 were 50

ms and 70 ms, respectively. Finally, for protein sequence identification, the raw data were

analyzed using Proteome Discoverer 2.4 (Thermo Fisher Scientific) with the Sequest HT

search tool with mass tolerance parameters of 10 ppm and 0.02 Da for precursor and

fragment ions, respectively.

1337

Dynamic light scattering experiments

The samples corresponding to the elution peaks obtained by size exclusion

chromatography for BC1280 and N-domain$_{39-190}$ were filtered through a 0.46-µm syringe

filter. The size measurements were performed using a Malvern Zetasizer Nano ZS

(Malvern Panalytical, Malvern, United Kingdom) with a laser wavelength of 632.8 nm as

the excitation source and a 1 cm pathway polystyrene cuvette as the sample holder. A

total volume of 1 ml of the protein solution was transferred to the cuvette, and the dynamic

light scattering signal was measured with a count rate ranging from 100 to 230 kcps,

depending on the sample. Data collection and subsequent analysis were conducted using

Zetasizer software v.6.34 (Malvern Panalytical).

Transmission electron microscopy

To study the morphology of BC1280, N-domain$_{39-191}$ and C-domain$_{191-279}$ purified *in vitro*,

the samples were subjected to 2-fold serial dilutions and incubated on copper grids for 2

minutes. Then, the samples were stained with a 2% uranyl acetate solution for 1 minute

and dried under dark conditions.

The samples were examined using a Thermo Fisher Scientific Tecnai G^2 20 TWIN

transmission electron microscope at an accelerating voltage of 120 kV. The final images

were captured using an Olympus Veleta side-mounted CCD with a resolution of 2k × 2k

Mpx.

1357

Thioflavin T (ThT) assays

The polymerization kinetics were studied using thioflavin T staining. The experiment was

conducted in a 96-well plate, and 40 μM ThT was added to each well at a specific

concentration, after which the mixture was diluted in buffer containing 20 mM Tris and 50

mM NaCl at pH 7.4. The polymerization kinetics of CalY were assayed at 40 μM using

low concentrations of BC1280 and N-domain$_{39\text{-}190}$ (2 and 10 μM).

The plates were incubated at 37°C in a plate reader and shaken at 100 rpm before

each measurement (Tecan Infinite M1000 Pro; Tecan, Männedorf, Switzerland). The

fluorescence intensity was monitored every 30 minutes through the bottom of the plate

using an excitation wavelength of 440 nm and an emission wavelength of 480 nm. All

measurements were recorded in triplicate.

1369

Sequence analysis by different bioinformatics tools

The primary sequence of BC1280 was obtained from UniProt (https://www.uniprot.org/)

with the accession number Q81GC7. SignalP5.0 was used to predict the cleavage site

for the signal peptidase.[84] The hydrophobicity was analyzed using the ProtScale tool

provided by the ExPASy server, and the prediction of disordered regions was conducted

using FoldUnfold with default parameters.[45,47] Putative amyloidogenic regions were

predicted employing the following bioinformatics tools: AmylPred2, FoldAmyloid,

MetAmyl, PASTA2.0 and TANGO.[40–44]

1378

AlphaFold prediction

1380 The secondary structure model of BC1280 was predicted by AlphaFold using the

1381 sequence available in UniProt with the accession number Q81GC7 and default

1382 parameters.[48,49] The pdb model obtained was subsequently used for predicting chemical

1383 shifts via SPARTA.[53] Finally, the predicted correlations overlapped with the experimental

1384 data obtained by SSNMR.

1385

In vitro **assembly of oligomers**

1387 BC1280 and N-domain$_{39-190}$ were diluted to a final concentration of 50 µM in buffer

1388 containing 20 mM Tris and 50 mM NaCl at pH 7.5. The oligomers self-assembled under

1389 agitation for 2 weeks at 37°C and a speed of 200 rpm. The aggregates were then

1390 ultracentrifuged (40k rpm speed, 5 hours, 18°C, TLA 120.1 rotor; Beckman Coulter), and

1391 the resulting pellets were analyzed using X-ray diffraction, ATR-FTIR, and SSNMR.

1392

X-ray diffraction measurements

Filament diffraction patterns were obtained at 4°C using a Rigaku FR-X rotating anode X-

ray generator (Rigaku, Tokyo, Japan) equipped with an EIGER 1 M hybrid pixel detector

(Dectris, Baden, Switzerland) at the copper wavelength. The concentrated hydrated

samples were mounted in MicroLoops from Mitegen (Ithaca, NY, USA) on a goniometer

head under cold nitrogen flow. Each diffraction pattern represents a 360° rotation along

the φ axis, with an exposure time of 720 s. No correction, such as smooth filtering or

baseline correction, was applied to the data. WinPLOTR (https://cdifx.univ-

rennes1.fr/winplotr/winplotr.htm) was used for converting q values to reticular distance (d)

values before plotting.

1403

Proteinase K (PK) digestion

A 22.5 µg solution of aggregated N-domain$_{39\text{-}190}$ was treated for 45 minutes at 37°C with

3 µg/ml proteinase K in 20 mM Tris and 50 mM NaCl at pH 7.5. The proteinase activity

was stopped using 5 mM PMSF. The samples were mixed with an equal volume of

Laemmli buffer (Bio-Rad) and heated at 100°C for 5 minutes. Subsequently, the samples

were separated using 12% SDS–PAGE, and the bands were visualized by Coomassie

blue staining. The bands of interest were excised and subjected to analysis by tandem

mass spectrometry.

1412

Attenuated total reflection FTIR spectroscopy

The samples were treated under the same conditions as those used for the X-ray studies.

Infrared spectra for BC1280 and N-domain$_{39\text{-}190}$ were recorded with an FT-IR

spectrophotometer (model Vertex70; Bruker) in attenuated total reflectance (ATR) mode

utilizing a Golden Gate Single Reflection Diamond ATR System accessory (Specac). No

sample preparation was needed, and spectra were recorded with an air background and

64 scans for both the background and sample measurements. The spectral resolution

was set at 4 cm-1". The raw data within the spectral range of the amide I region were

used for estimating the total amount of secondary structure. To increase the resolution of

the minimal absorbance peaks, a second derivative was calculated. Next, a deconvolution

approach was employed using PeakFit software to determine the contribution percentage

for each type of secondary structure: parallel β-sheet (1634 cm^{-1}), antiparallel β-sheet

(1620 cm^{-1} and 1690 cm^{-1}), α-helix (1658 cm^{-1}), turns (1675 cm^{-1}) and random coil (1646

cm^{-1}), in accordance with the well-established assignments.[58]

1427

1428   **Proteomic analysis by the isobaric tag for relative and absolute quantification**

1429   **(iTRAQ) method**

1430   To estimate the relative levels of TasA, BC1280 and CalY during biofilm formation in the

1431   wild-type strain *B. cereus* ATCC14579, we conducted sample preprocessing and

1432   subsequent analysis following the protocols described by Caro-Astorga et al.,in 2020.[26]

1433

NMR spectroscopy

Solid-state NMR experiments were performed on a 600 MHz spectrometer (Bruker

Biospin, Germany) equipped with a triple resonance 4mm probe. Experiments were

performed at 11 kHz MAS frequency at a sample temperature of 283K. Chemical shifts

were referenced with DSS. For 2D ^{13}C-^{13}C PDSD spectra (mixing time of 50 and 200 ms),

an initial cross-polarization step of 0.5 ms was used and high-power SPINAL-64

decoupling was applied during acquisition times. 2D ^{13}C-^{13}C PDSD (50 ms) and ^{1}H-^{13}C

INEPT spectra were recorded for 1-3 days, and 6 days for the ^{13}C-^{13}C PDSD spectrum

(200 ms).

Solution NMR ^{1}H,^{15}N SOFAST-HMQC experiment was carried out at 298 K on a protein

sample containing 200 µM BC1280, 20mM Tris, 50mM NaCl, pH7.5, on a 700 MHz

spectrometer (Bruker Biospin, Germany) equipped with a 5mm TXI ^{1}H/^{13}C/^{15}N/^{2}H probe.

The NMR data were processed using the TOPSPIN 4.0.6 software and analyzed by

CCPNMR.[85]

Circular dichroism

CalY (40 µM) alone or in combination with BC1280 (at 2 or 10 µM) was incubated in 10

mM sodium phosphate buffer at pH 8 for 16 hours at 37ºC under agitation. Circular

dichroism spectra were recorded using a JASCO J-815 spectrometer, using a 0.1 cm

path-length cuvette in the range of 180-260 nm, with a 0.5 nm step and a 1 s collection

time per step. The scan rate was 50 nm/min. Final spectra were obtained as the average

of six scans after a blank correction. The spectrometer was continuously purged with dry

N2 gas.CD spectra were buffer subtracted, and the results were expressed as mean

residue ellipticity.

REFERENCES

1. Elhariry, H.M. (2011). Attachment strength and biofilm forming ability of Bacillus cereus on green-leafy vegetables: Cabbage and lettuce. Food Microbiology *28*, 1266–1274. https://doi.org/10.1016/j.fm.2011.05.004.

2. Hu, H.-J., Chen, Y.-L., Wang, Y.-F., Tang, Y.-Y., Chen, S.-L., and Yan, S.-Z. (2017). Endophytic *Bacillus cereus* Effectively Controls *Meloidogyne incognita* on Tomato Plants Through Rapid Rhizosphere Occupation and Repellent Action. Plant Disease *101*, 448–455. https://doi.org/10.1094/PDIS-06-16-0871-RE.

3. Cormontagne, D., Rigourd, V., Vidic, J., Rizzotto, F., Bille, E., and Ramarao, N. (2021). Bacillus cereus Induces Severe Infections in Preterm Neonates: Implication at the Hospital and Human Milk Bank Level. Toxins *13*, 123. https://doi.org/10.3390/toxins13020123.

4. Bottone, E.J. (2010). *Bacillus cereus* , a Volatile Human Pathogen. Clin Microbiol Rev *23*, 382–398. https://doi.org/10.1128/CMR.00073-09.

5. Thorsen, L., Azokpota, P., Munk Hansen, B., Rønsbo, M.H., Nielsen, K.F., Hounhouigan, D.J., and Jakobsen, M. (2011). Formation of cereulide and enterotoxins by Bacillus cereus in fermented African locust beans. Food Microbiology *28*, 1441–1447. https://doi.org/10.1016/j.fm.2011.07.003.

6. McKillip, J.L. (2000). Prevalence and expression of enterotoxins in Bacillus cereus and other Bacillus spp., a literature review. Antonie van Leeuwenhoek *77*, 393–399. https://doi.org/10.1023/A:1002706906154.

7. Dragoš, A., and Kovács, Á.T. (2017). The Peculiar Functions of the Bacterial Extracellular Matrix. Trends in Microbiology *25*, 257–266. https://doi.org/10.1016/j.tim.2016.12.010.

8. Jamal, M., Ahmad, W., Andleeb, S., Jalil, F., Imran, M., Nawaz, M.A., Hussain, T., Ali, M., Rafiq, M., and Kamil, M.A. (2018). Bacterial biofilm and associated infections. Journal of the Chinese Medical Association *81*, 7–11. https://doi.org/10.1016/j.jcma.2017.07.012.

9. Limoli, D.H., Jones, C.J., and Wozniak, D.J. (2015). Bacterial Extracellular Polysaccharides in Biofilm Formation and Function. Microbiol Spectr *3*, 3.3.29. https://doi.org/10.1128/microbiolspec.MB-0011-2014.

10. Lopez, D., Vlamakis, H., and Kolter, R. (2010). Biofilms. Cold Spring Harbor Perspectives in Biology *2*, a000398–a000398. https://doi.org/10.1101/cshperspect.a000398.

11. Lemon, K.P., Earl, A.M., Vlamakis, H.C., Aguilar, C., and Kolter, R. (2008). Biofilm Development with an Emphasis on Bacillus subtilis. In Bacterial Biofilms Current Topics in Microbiology and Immunology., T. Romeo, ed. (Springer Berlin Heidelberg), pp. 1–16. https://doi.org/10.1007/978-3-540-75418-3_1.

12. Bridier, A., Le Coq, D., Dubois-Brissonnet, F., Thomas, V., Aymerich, S., and Briandet, R. (2011). The Spatial Architecture of Bacillus subtilis Biofilms Deciphered Using a Surface-

Associated Model and In Situ Imaging. PLoS ONE *6*, e16177.
https://doi.org/10.1371/journal.pone.0016177.

13. Pham, C.L.L., Kwan, A.H., and Sunde, M. (2014). Functional amyloid: widespread in
Nature, diverse in purpose. Essays in Biochemistry *56*, 207–219.
https://doi.org/10.1042/bse0560207.

14. Álvarez-Mena, A., Cámara-Almirón, J., De Vicente, A., and Romero, D. (2020).
Multifunctional Amyloids in the Biology of Gram-Positive Bacteria. Microorganisms *8*,
2020. https://doi.org/10.3390/microorganisms8122020.

15. Blanco, L.P., Evans, M.L., Smith, D.R., Badtke, M.P., and Chapman, M.R. (2012).
Diversity, biogenesis and function of microbial amyloids. Trends in Microbiology *20*, 66–73.
https://doi.org/10.1016/j.tim.2011.11.005.

16. Fowler, D.M., Koulov, A.V., Balch, W.E., and Kelly, J.W. (2007). Functional amyloid –
from bacteria to humans. Trends in Biochemical Sciences *32*, 217–224.
https://doi.org/10.1016/j.tibs.2007.03.003.

17. Dragoš, A., Kovács, Á.T., and Claessen, D. (2017). The Role of Functional Amyloids in
Multicellular Growth and Development of Gram-Positive Bacteria. Biomolecules *7*, 60.
https://doi.org/10.3390/biom7030060.

18. Romero, D., and Kolter, R. (2014). Functional amyloids in bacteria. International
Microbiology, 65–73. https://doi.org/10.2436/20.1501.01.208.

19. Majed, R., Faille, C., Kallassy, M., and Gohar, M. (2016). Bacillus cereus Biofilms—Same,
Only Different. Front. Microbiol. *7*. https://doi.org/10.3389/fmicb.2016.01054.

20. Lin, Y., Briandet, R., and Kovács, Á.T. (2022). Bacillus cereus sensu lato biofilm formation
and its ecological importance. Biofilm *4*, 100070.
https://doi.org/10.1016/j.bioflm.2022.100070.

21. Houry, A., Briandet, R., Aymerich, S., and Gohar, M. (2010). Involvement of motility and
flagella in Bacillus cereus biofilm formation. Microbiology *156*, 1009–1018.
https://doi.org/10.1099/mic.0.034827-0.

22. Candela, T., Fagerlund, A., Buisson, C., Gilois, N., Kolstø, A., Økstad, O.A., Aymerich, S.,
Nielsen-Leroux, C., Lereclus, D., and Gohar, M. (2019). CalY is a major virulence factor
and a biofilm matrix protein. Molecular Microbiology *111*, 1416–1429.
https://doi.org/10.1111/mmi.14184.

23. Vilain, S., Pretorius, J.M., Theron, J., and Brözel, V.S. (2009). DNA as an Adhesin: *Bacillus*
cereus Requires Extracellular DNA To Form Biofilms. Appl Environ Microbiol *75*, 2861–
2868. https://doi.org/10.1128/AEM.01317-08.

24. Caro-Astorga, J., Álvarez-Mena, A., Hierrezuelo, J., Guadix, J.A., Heredia-Ponce, Z.,
Arboleda-Estudillo, Y., González-Munoz, E., De Vicente, A., and Romero, D. (2020). Two

genomic regions encoding exopolysaccharide production systems have complementary
functions in B. cereus multicellularity and host interaction. Sci Rep *10*, 1000.
https://doi.org/10.1038/s41598-020-57970-3.

25. Caro-Astorga, J., Pérez-García, A., De Vicente, A., and Romero, D. (2015). A genomic
region involved in the formation of adhesin fibers in Bacillus cereus biofilms. Front.
Microbiol. *5*. https://doi.org/10.3389/fmicb.2014.00745.

26. Caro-Astorga, J., Frenzel, E., Perkins, J.R., Álvarez-Mena, A., De Vicente, A., Ranea,
J.A.G., Kuipers, O.P., and Romero, D. (2020). Biofilm formation displays intrinsic offensive
and defensive features of Bacillus cereus. npj Biofilms Microbiomes *6*, 3.
https://doi.org/10.1038/s41522-019-0112-7.

27. Stöver, A.G., and Driks, A. (1999). Secretion, Localization, and Antibacterial Activity of
TasA, a *Bacillus subtilis* Spore-Associated Protein. J Bacteriol *181*, 1664–1672.
https://doi.org/10.1128/JB.181.5.1664-1672.1999.

28. Tjalsma, H., Bolhuis, A., Van Roosmalen, M.L., Wiegert, T., Schumann, W., Broekhuizen,
C.P., Quax, W.J., Venema, G., Bron, S., and Van Dijl, J.M. (1998). Functional analysis of
the secretory precursor processing machinery of *Bacillus subtilis* : identification of a
eubacterial homolog of archaeal and eukaryotic signal peptidases. Genes Dev. *12*, 2318–
2331. https://doi.org/10.1101/gad.12.15.2318.

29. Romero, D., Aguilar, C., Losick, R., and Kolter, R. (2010). Amyloid fibers provide structural
integrity to *Bacillus subtilis* biofilms. Proc. Natl. Acad. Sci. U.S.A. *107*, 2230–2234.
https://doi.org/10.1073/pnas.0910560107.

30. Diehl, A., Roske, Y., Ball, L., Chowdhury, A., Hiller, M., Molière, N., Kramer, R., Stöppler,
D., Worth, C.L., Schlegel, B., et al. (2018). Structural changes of TasA in biofilm formation
of *Bacillus subtilis*. Proc. Natl. Acad. Sci. U.S.A. *115*, 3237–3242.
https://doi.org/10.1073/pnas.1718102115.

31. Erskine, E., Morris, R.J., Schor, M., Earl, C., Gillespie, R.M.C., Bromley, K.M., Sukhodub,
T., Clark, L., Fyfe, P.K., Serpell, L.C., et al. (2018). Formation of functional, non-
amyloidogenic fibres by recombinant *Bacillus subtilis* TasA. Molecular Microbiology *110*,
897–913. https://doi.org/10.1111/mmi.13985.

32. Mammeri, N.E., Hierrezuelo, J., Tolchard, J., Cámara-Almirón, J., Caro-Astorga, J.,
Álvarez-Mena, A., Dutour, A., Berbon, M., Shenoy, J., Morvan, E., et al. (2019). Molecular
architecture of bacterial amyloids in *Bacillus* biofilms. FASEB j. *33*, 12146–12163.
https://doi.org/10.1096/fj.201900831R.

33. Böhning, J., Ghrayeb, M., Pedebos, C., Abbas, D.K., Khalid, S., Chai, L., and Bharat,
T.A.M. (2022). Donor-strand exchange drives assembly of the TasA scaffold in Bacillus
subtilis biofilms. Nat Commun *13*, 7082. https://doi.org/10.1038/s41467-022-34700-z.

34. Fagerlund, A., Dubois, T., Økstad, O.-A., Verplaetse, E., Gilois, N., Bennaceur, I., Perchat,
S., Gominet, M., Aymerich, S., Kolstø, A.-B., et al. (2014). SinR Controls Enterotoxin

Expression in Bacillus thuringiensis Biofilms. PLoS ONE *9*, e87532.
https://doi.org/10.1371/journal.pone.0087532.

35. Jorda, J., Xue, B., Uversky, V.N., and Kajava, A.V. (2010). Protein tandem repeats - the
more perfect, the less structured: Structural state of perfect protein repeats. FEBS Journal
277, 2673–2682. https://doi.org/10.1111/j.1742-464X.2010.07684.x.

36. Helmann, J.D. (2016). Bacillus subtilis extracytoplasmic function (ECF) sigma factors and
defense of the cell envelope. Current Opinion in Microbiology *30*, 122–132.
https://doi.org/10.1016/j.mib.2016.02.002.

37. Mascher, T. (2023). Past, Present, and Future of Extracytoplasmic Function σ Factors:
Distribution and Regulatory Diversity of the Third Pillar of Bacterial Signal Transduction.
Annu. Rev. Microbiol. *77*, 625–644. https://doi.org/10.1146/annurev-micro-032221-024032.

38. Gade Malmos, K., Blancas-Mejia, L.M., Weber, B., Buchner, J., Ramirez-Alvarado, M.,
Naiki, H., and Otzen, D. (2017). ThT 101: a primer on the use of thioflavin T to investigate
amyloid formation. Amyloid *24*, 1–16. https://doi.org/10.1080/13506129.2017.1304905.

39. Sunde, M., Serpell, L.C., Bartlam, M., Fraser, P.E., Pepys, M.B., and Blake, C.C.F. (1997).
Common core structure of amyloid fibrils by synchrotron X-ray diffraction 1 1Edited by F.
E. Cohen. Journal of Molecular Biology *273*, 729–739.
https://doi.org/10.1006/jmbi.1997.1348.

40. Tsolis, A.C., Papandreou, N.C., Iconomidou, V.A., and Hamodrakas, S.J. (2013). A
Consensus Method for the Prediction of 'Aggregation-Prone' Peptides in Globular Proteins.
PLoS ONE *8*, e54175. https://doi.org/10.1371/journal.pone.0054175.

41. Garbuzynskiy, S.O., Lobanov, M.Yu., and Galzitskaya, O.V. (2010). FoldAmyloid: a
method of prediction of amyloidogenic regions from protein sequence. Bioinformatics *26*,
326–332. https://doi.org/10.1093/bioinformatics/btp691.

42. Emily, M., Talvas, A., and Delamarche, C. (2013). MetAmyl: A METa-Predictor for
AMYLoid Proteins. PLoS ONE *8*, e79722. https://doi.org/10.1371/journal.pone.0079722.

43. Walsh, I., Seno, F., Tosatto, S.C.E., and Trovato, A. (2014). PASTA 2.0: an improved server
for protein aggregation prediction. Nucleic Acids Research *42*, W301–W307.
https://doi.org/10.1093/nar/gku399.

44. Fernandez-Escamilla, A.-M., Rousseau, F., Schymkowitz, J., and Serrano, L. (2004).
Prediction of sequence-dependent and mutational effects on the aggregation of peptides and
proteins. Nat Biotechnol *22*, 1302–1306. https://doi.org/10.1038/nbt1012.

45. Galzitskaya, O.V., Garbuzynskiy, S.O., and Lobanov, M.Yu. (2006). FoldUnfold: web server
for the prediction of disordered regions in protein chain. Bioinformatics *22*, 2948–2949.
https://doi.org/10.1093/bioinformatics/btl504.

46. Romero, P., Obradovic, Z., Li, X., Garner, E.C., Brown, C.J., and Dunker, A.K. (2001).
Sequence complexity of disordered protein. Proteins *42*, 38–48.
https://doi.org/10.1002/1097-0134(20010101)42:1<38::AID-PROT50>3.0.CO;2-3.

47. Wilkins, M.R., Gasteiger, E., Bairoch, A., Sanchez, J.-C., Williams, K.L., Appel, R.D., and
Hochstrasser, D.F. (1998). Protein Identification and Analysis Tools in the ExPASy Server.
In 2-D Proteome Analysis Protocols (Humana Press), pp. 531–552.
https://doi.org/10.1385/1-59259-584-7:531.

48. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596*, 583–589.
https://doi.org/10.1038/s41586-021-03819-2.

49. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D.,
Stroe, O., Wood, G., Laydon, A., et al. (2022). AlphaFold Protein Structure Database:
massively expanding the structural coverage of protein-sequence space with high-accuracy
models. Nucleic Acids Research *50*, D439–D444. https://doi.org/10.1093/nar/gkab1061.

50. Meier, B.H., Riek, R., and Böckmann, A. (2017). Emerging Structural Understanding of
Amyloid Fibrils by Solid-State NMR. Trends in Biochemical Sciences *42*, 777–787.
https://doi.org/10.1016/j.tibs.2017.08.001.

51. Loquet, A., El Mammeri, N., Stanek, J., Berbon, M., Bardiaux, B., Pintacuda, G., and
Habenstein, B. (2018). 3D structure determination of amyloid fibrils using solid-state NMR
spectroscopy. Methods *138–139*, 26–38. https://doi.org/10.1016/j.ymeth.2018.03.014.

52. Van Der Wel, P.C.A. (2017). Insights into protein misfolding and aggregation enabled by
solid-state NMR spectroscopy. Solid State Nuclear Magnetic Resonance *88*, 1–14.
https://doi.org/10.1016/j.ssnmr.2017.10.001.

53. Shen, Y., and Bax, A. (2010). SPARTA+: a modest improvement in empirical NMR
chemical shift prediction by means of an artificial neural network. J Biomol NMR *48*, 13–22.
https://doi.org/10.1007/s10858-010-9433-9.

54. Wang, Y., and Jardetzky, O. (2002). Probability-based protein secondary structure
identification using combined NMR chemical-shift data. Protein Science *11*, 852–861.
https://doi.org/10.1110/ps.3180102.

55. Ulamec, S.M., Brockwell, D.J., and Radford, S.E. (2020). Looking Beyond the Core: The
Role of Flanking Regions in the Aggregation of Amyloidogenic Peptides and Proteins. Front.
Neurosci. *14*, 611285. https://doi.org/10.3389/fnins.2020.611285.

56. Bhopatkar, A.A., and Kayed, R. (2023). Flanking regions, amyloid cores, and
polymorphism: the potential interplay underlying structural diversity. Journal of Biological
Chemistry *299*, 105122. https://doi.org/10.1016/j.jbc.2023.105122.

57. Matlahov, I., and Van Der Wel, P.C.A. (2018). Hidden motions and motion-induced
invisibility: Dynamics-based spectral editing in solid-state NMR. Methods *148*, 123–135.
https://doi.org/10.1016/j.ymeth.2018.04.015.

58. Goormaghtigh, E., Cabiaux, V., and Ruysschaert, J.-M. (1994). Determination of Soluble
and Membrane Protein Structure by Fourier Transform Infrared Spectroscopy. In
Physicochemical Methods in the Study of Biomembranes Subcellular Biochemistry., H. J.
Hilderson and G. B. Ralston, eds. (Springer US), pp. 405–450. https://doi.org/10.1007/978-
1-4615-1863-1_10.

59. Romero, D., Vlamakis, H., Losick, R., and Kolter, R. (2011). An accessory protein required
for anchoring and assembly of amyloid fibres in *B. subtilis* biofilms. Molecular
Microbiology *80*, 1155–1168. https://doi.org/10.1111/j.1365-2958.2011.07653.x.

60. Roske, Y., Lindemann, F., Diehl, A., Cremer, N., Higman, V.A., Schlegel, B., Leidert, M.,
Driller, K., Turgay, K., Schmieder, P., et al. (2023). TapA acts as specific chaperone in TasA
filament formation by strand complementation. Proc. Natl. Acad. Sci. U.S.A. *120*,
e2217070120. https://doi.org/10.1073/pnas.2217070120.

61. Abbasi, R., Mousa, R., Dekel, N., Amartely, H., Danieli, T., Lebendiker, M., Levi-Kalisman,
Y., Shalev, D.E., Metanis, N., and Chai, L. (2019). The Bacterial Extracellular Matrix
Protein TapA Is a Two-Domain Partially Disordered Protein. ChemBioChem *20*, 355–359.
https://doi.org/10.1002/cbic.201800634.

62. Lukaszczyk, M., Pradhan, B., and Remaut, H. (2019). The Biosynthesis and Structures of
Bacterial Pili. In Bacterial Cell Walls and Membranes Subcellular Biochemistry., A. Kuhn,
ed. (Springer International Publishing), pp. 369–413. https://doi.org/10.1007/978-3-030-
18768-2_12.

63. Shanmugasundarasamy, T., Karaiyagowder Govindarajan, D., and Kandaswamy, K. (2022).
A review on pilus assembly mechanisms in Gram-positive and Gram-negative bacteria. The
Cell Surface *8*, 100077. https://doi.org/10.1016/j.tcsw.2022.100077.

64. Pradhan, B., Liedtke, J., Sleutel, M., Lindbäck, T., Zegeye, E.D., O´Sullivan, K., Llarena, A.,
Brynildsrud, O., Aspholm, M., and Remaut, H. (2021). Endospore Appendages: a novel pilus
superfamily from the endospores of pathogenic Bacilli. The EMBO Journal *40*, e106887.
https://doi.org/10.15252/embj.2020106887.

65. Hendrickx, A.P.A., Poor, C.B., Jureller, J.E., Budzik, J.M., He, C., and Schneewind, O.
(2012). Isopeptide bonds of the major pilin protein BcpA influence pilus structure and
bundle formation on the surface of *Bacillus cereus*. Molecular Microbiology *85*, 152–163.
https://doi.org/10.1111/j.1365-2958.2012.08098.x.

66. Zegeye, E.D., Pradhan, B., Llarena, A.-K., and Aspholm, M. (2021). Enigmatic Pilus-Like
Endospore Appendages of Bacillus cereus Group Species. IJMS *22*, 12367.
https://doi.org/10.3390/ijms222212367.

67. Wilson, M.R., Yerbury, J.J., and Poon, S. (2008). Potential roles of abundant extracellular
chaperones in the control of amyloid formation and toxicity. Mol. BioSyst. *4*, 42–52.
https://doi.org/10.1039/B712728F.

68. Kovacs, D., and Tompa, P. (2012). Diverse functional manifestations of intrinsic structural
disorder in molecular chaperones. Biochemical Society Transactions *40*, 963–968.
https://doi.org/10.1042/BST20120108.

69. Foit, L., George, J.S., Zhang, B.W., Brooks, C.L., and Bardwell, J.C.A. (2013). Chaperone
activation by unfolding. Proc. Natl. Acad. Sci. U.S.A. *110*.
https://doi.org/10.1073/pnas.1222458110.

70. Díaz-Villanueva, J., Díaz-Molina, R., and García-González, V. (2015). Protein Folding and
Mechanisms of Proteostasis. IJMS *16*, 17193–17230.
https://doi.org/10.3390/ijms160817193.

71. Arnaud, M., Chastanet, A., and Débarbouillé, M. (2004). New Vector for Efficient Allelic
Replacement in Naturally Nontransformable, Low-GC-Content, Gram-Positive Bacteria.
Appl Environ Microbiol *70*, 6887–6891. https://doi.org/10.1128/AEM.70.11.6887-
6891.2004.

72. Pflughoeft, K.J., Sumby, P., and Koehler, T.M. (2011). *Bacillus anthracis sin* Locus and
Regulation of Secreted Proteases. J Bacteriol *193*, 631–639.
https://doi.org/10.1128/JB.01083-10.

73. Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O., Madhusoodanan,
N., Kolesnikov, A., and Lopez, R. (2022). Search and sequence analysis tools services from
EMBL-EBI in 2022. Nucleic Acids Research *50*, W276–W279.
https://doi.org/10.1093/nar/gkac240.

74. Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., and Rozen,
S.G. (2012). Primer3—new capabilities and interfaces. Nucleic Acids Research *40*, e115–
e115. https://doi.org/10.1093/nar/gks596.

75. Vargas, P., Felipe, A., Michán, C., and Gallegos, M.-T. (2011). Induction of *Pseudomonas
syringae* pv. *tomato* DC3000 MexAB-OprM Multidrug Efflux Pump by Flavonoids Is
Mediated by the Repressor PmeR. MPMI *24*, 1207–1219. https://doi.org/10.1094/MPMI-03-
11-0077.

76. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using
Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods *25*, 402–408.
https://doi.org/10.1006/meth.2001.1262.

77. O'Toole, G.A., Pratt, L.A., Watnick, P.I., Newman, D.K., Weaver, V.B., and Kolter, R.
(1999). [6] Genetic approaches to study of biofilms. In Methods in Enzymology (Elsevier),
pp. 91–109. https://doi.org/10.1016/S0076-6879(99)10008-9.

78. Falgueras, J., Lara, A.J., Fernández-Pozo, N., Cantón, F.R., Pérez-Trabado, G., and Claros,
M.G. (2010). SeqTrim: a high-throughput pipeline for pre-processing any type of sequence
read. BMC Bioinformatics *11*, 38. https://doi.org/10.1186/1471-2105-11-38.

79. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

80. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352.

81. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR : a Bioconductor package
for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–
140. https://doi.org/10.1093/bioinformatics/btp616.

82. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

83. Dunn, K.W., Kamocka, M.M., and McDonald, J.H. (2011). A practical guide to evaluating
colocalization in biological microscopy. American Journal of Physiology-Cell Physiology
300, C723–C742. https://doi.org/10.1152/ajpcell.00462.2010.

84. Almagro Armenteros, J.J., Tsirigos, K.D., Sønderby, C.K., Petersen, T.N., Winther, O.,
Brunak, S., Von Heijne, G., and Nielsen, H. (2019). SignalP 5.0 improves signal peptide
predictions using deep neural networks. Nat Biotechnol *37*, 420–423.
https://doi.org/10.1038/s41587-019-0036-z.

85. Skinner, S.P., Fogh, R.H., Boucher, W., Ragan, T.J., Mureddu, L.G., and Vuister, G.W.
(2016). CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis. J Biomol
NMR *66*, 111–124. https://doi.org/10.1007/s10858-016-0060-y.

1736

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Secondary structure element	Wavenumbers (cm⁻¹)	Percentage of structural elements (%)	
		BC1280	N-domain₃₉₋₁₉₀
β-sheet antiparallel	1620,1690	25%	33%
β-sheet parallel	1634	22%	19%
Random coil	1646	24%	24%
α-helix	1658	15%	12%
Turn	1675	14%	12%

Figure 6

A

Fluorescence intensity (A.U.) x 10³

CalY/BC1280 (%)
— 100/0
— 30/70
— 70/30
— 50/50
— 0/100

Time (hours)

B

Log2 (relative expression)

■ BC1280
■ CalY

Time (hours)

C

Fluorescence intensity (A.U.) x 10³

Time (hours)

— CalY 40 μM
— + BC1280 2 μM
— + BC1280 10 μM
— + N-domain$_{39-190}$ 2 μM
— + N-domain$_{39-190}$ 10 μM

D

| Negative control | + BC1280 10 μM | + N-domain$_{39-190}$ 10 μM |

E

mdeg

— CalY 40 μM
— +BC1280 2 μM
— +BC1280 10 μM

Absorbance

F

| WT | ΔcalY |

anti-TasA

| WT | ΔtasA |

anti-CalY

Initial stages of biofilm formation

Biofilm maturation

Cytoplasm

CapP unfolded

CalY unfolded

TasA unfolded

SEC

SipW

SEC

SipW

Periplasmic space

Extracellular medium

Low levels of CapP

cat.

cat.

High levels of CapP

CapP aggregates

X

Figure S1

Figure S2

Figure S3

Figure S4

A

```
                    10          20          30          40          50          60          70
         MGE I KMLKR PRNFKKML I LPCMCS I TFYLGSQ IMTHTEAAF IHETKVEATLSTA I I I FPKTVNTLKEQSEKHKQF I EREY
BC1280   - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - KVEATLSTA I I FPKT - - - - - - - - HKQF I EREY
N-domain₃₉₋₁₉₀  - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - AF IHETKVEATLSTA I I FPKTVNTLKEQSEKHKQF I EREY

                    80          90          100         110         120         130         140         150
         GTMKGKLKATS I EE I KQA I SVWQQGREK I VAEKEALQNVYTE I EAPYNQ I QEELKVNKDESMQQVS I YVNEGFRS I KEK
BC1280   GTMKGKLKATS I EE I KQA I SVWQQGREK - - - - - - ALQNVYTE I EAPYNQ I QEELKVNKDESMQQVS I YVNEGFRS I - - -
N-domain₃₉₋₁₉₀  GTMK - - LKATS I EE I KQA I SVWQQGREK I VAEKEALQNVYTE I EAPYNQ I QEELKVNKDESMQQVS I YVNEGFRS I K -

                    160         170         180         190         200         210         220         230
         RDY I EKE I SLKA I DEQ I QALQQQLNVA I EAEGQKKVEEQKKAEEQKKVEEQKKVEEQKKAEEQKKVEEQKKAEEQKKVE
BC1280   RDY I EKE I SLKA I DEQ I QALQQQLNVA I EAE - - - - VEEQ - - - - - - KKAEEQ - - - - - - - - - - - KKVEEQKKAEEQKKVE
N-domain₃₉₋₁₉₀  RDY I EKE I SLKA - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -

                    240         250         260         270
BC1280   EQKKAEEQKKAEEQKKVEEQKKVEEQKKAEEQKKVEEQKKAE
N-domain₃₉₋₁₉₀  EQKKAEEQK - - - - - - KKVEEQKKVEEQKKAEEQKKVEEQKK -
         - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
```

B

C

1   MGEIKMLKRPRNFKKMLILPCMCSITFYLGSQIMTHTEAAFIHETKVEATLSTAIIFPKTVNTLKEQSEKHKQFIEREYGTMKGKLKATSIEEIKQ   96

97 AISVWQQGREKIVAEKEALQNVYTEIEAPYNQIQEELKVNKDESMQQVSIYVNEGFRSIKEKRDYIEKEISLKAIDEQIQALQQQLNVAIEAEGQ   191

192 KKVEEQKKAEEQKKVEEQKKVEEQKKAEEQKKVEEQKKAEEQKKVEEQKKAEEQKKAEEQKKVEEQKKVEEQKKAEEQKKVEEQKKAE   279

XXX folded   XXX treated as folded   XXX unfolded
XXX putative amyloidogenic

D

E

Figure S5

Figure S6