Metabolic phenotyping of healthy and diseased human RPE cells

Saira Rizwan[1,2*], Beverly Toothman[1,2*], Bo Li[1,2,3*], Abbi J. Engel[4], Rayne R. Lim[4], Sheldon Niernberger[4],

Jinyu Lu[1,2], Cloe Ratliff[1,2], Yinxiao Xiang[1,2], Mark Eminhizer[1,2], Jennifer R. Chao[4&], and Jianhai Du[1,2&]

[1]Department of Ophthalmology and Visual Sciences, West Virginia University, Morgantown, WV 26506

[2]Department of Biochemistry and Molecular Medicine, West Virginia University, Morgantown, WV

26506

[3]Department of Ophthalmology, Affiliated Hospital of Yangzhou University, Yangzhou University,

Yangzhou, Jiangsu Province 225100, China.

[4]Department of Ophthalmology, University of Washington, Seattle, WA 98109, United States.

*Equal contributions

&Corresponding Authors: Jianhai Du, One Medical Center Dr, PO Box 9193, WVU Eye Institute,

Morgantown, WV 26505; Phone: (304)-293-9346; Fax; (304)-598-6928; Email: jianhai.du@hsc.wvu.edu;

or Jennifer R. Chao: 750 Republican Street, Box 358058, Seattle WA 98109; Phone: (206) 221-0594.

Email: jrchao@uw.edu.

Word count: 3473

Commercial relationships disclosures: None.

Funding: The funders had no role in the study design, data collection and analysis, decision on

publishing, or manuscript preparation. This work was supported by National Institutes of Health Grant

EY034364 (J. R. C. and J. D), EY03459 (J. R. C. and J. D), EY031324 (JD), EY032462(JD), the Retina

Research Foundation (JD), UW Ophthalmology Stem Cell Fund (JRC), RPB Unrestricted Grant (UW),

and funds for Core facilities P20 GM103434 and P20 GM144230 (WV INBRE grant).

Abstract (249 words)

Purpose: Metabolic defects in the retinal pigment epithelium (RPE) underlie many retinal degenerative diseases. This study aims to identify the nutrient requirements of healthy and diseased human RPE cells.

Methods: We profiled nutrient utilization of various human RPE cells, including differentiated and dedifferentiated fetal RPE (fRPE), induced pluripotent stem cell derived-RPE (iPSC RPE), Sorsby fundus dystrophy (SFD) patient-derived iPSC RPE, CRISPR-corrected isogenic SFD (cSFD) iPSC RPE, and ARPE-19 cell lines using Biolog Phenotype MicroArray Assays.

Results: Differentiated fRPE cells and healthy iPSC RPE cells can utilize 51 and 48 nutrients respectively, including sugars, intermediates from glycolysis and tricarboxylic acid (TCA) cycle, fatty acids, ketone bodies, amino acids, and dipeptides. However, when fRPE cells lose their epithelial phenotype through dedifferentiation, nutrient utilization becomes restricted to 17 nutrients, primarily sugar and glutamine-related amino acids. SFD RPE cells can utilize 37 nutrients; however, compared to cSFD RPE and healthy iPSC RPE, they are unable to utilize lactate, some TCA cycle intermediates, and short-chain fatty acids. Nonetheless, they show increased utilization of branch-chain amino acids (BCAAs) and BCAA-containing dipeptides. Dedifferentiated ARPE-19 cells grown in traditional culture media cannot utilize lactate and ketone bodies. In contrast, nicotinamide supplementation promotes differentiation towards an epithelial phenotype, restoring the ability to use these nutrients.

Conclusions: Epithelial phenotype confers metabolic flexibility to healthy RPE for utilizing various nutrients. SFD RPE cells have reduced metabolic flexibility, relying on the oxidation of BCAAs. Our findings highlight the potentially important roles of nutrient availability and utilization in RPE differentiation and diseases.

Introduction

Retinal pigment epithelium (RPE) is crucial in supporting photoreceptor function and survival through various functions, including the visual cycle, nutrient transport, light absorption, phagocytosis of outer segments, and formation of the blood-retina barrier[1]. To sustain these critical functions, the RPE relies on a robust mitochondrial metabolism by oxidizing fuels from the photoreceptors and choroidal supply[2]. Defects in RPE mitochondrial metabolism can cause epithelial-mesenchymal transition (EMT) and dedifferentiation, with loss of characteristic epithelial traits such as tight junctions, pigmentation and polarity, subsequently leading to photoreceptor death in retinal degenerative diseases, including Sorsby Fundus Dystrophy (SFD) and age-related macular degeneration (AMD)[2-4].

Cultured human RPE cells can be valuable models to investigate RPE function and retinal disease, as they exhibit typical RPE morphology, function and signature gene expression, which closely resemble native RPE cells[5-8]. Human fetal RPE (fRPE), patient-derived induced pluripotent stem cells (iPSC) RPE, and immortalized ARPE-19 cells are commonly used RPE cultures. Human fRPE cells have been rigorously characterized in their physiology, gene expression and metabolism, considered the gold standard in RPE culture[5, 9, 10]. iPSC RPE have been well-characterized and have the advantage of modeling inherited retinal diseases from patients. SFD, a rare early onset macular degeneration, results from mutations in the tissue inhibitor of metalloproteinase-3 (TIMP3) gene[11]. In our previous study, we showed that iPSC RPE derived from SFD patients carrying the S204C mutation in *TIMP3* have irregular extracellular matrices (ECM) and basal laminar deposits, while correction of the S204C variant in SFD iPSC RPE (cSFD) attenuates these findings[12]. ARPE-19 cells offer advantages over fRPE and iPSC RPE in terms of easier availability and low cost. ARPE-19 cells show fibroblast-like morphology under traditional culture media. However, supplementation with nicotinamide (NAM), a precursor for NAD, can rapidly induce differentiation of ARPE-19 cells into cells with many epithelial characteristics through revitalization of mitochondrial metabolism[13, 14].

In this study, we determined the metabolic phenotype of nutrient utilization in healthy and diseased human RPE. We conducted extensive metabolic screening of carbon and nitrogen sources across multiple human RPE cultures, including mature and dedifferentiated fRPE, healthy and SFD RPE, and ARPE-19 RPE cells under different culture media conditions. We found that healthy and mature RPE cells demonstrate robust metabolic flexibility, allowing them to utilize diverse nutrient sources. In contrast, dedifferentiated and SFD RPE cells have reduced metabolic flexibility, relying on specific nutrient sources. These cell-specific differences in nutrient utilization should provide insights into the underlying mechanisms of RPE differentiation and disease.

Methods

Reagent and key resources

All the reagents and key resources were detailed in Supplemental **Table S1** or methods.

Human RPE cell culture

Human fRPE was isolated from human fetal eye cups with no identifiers, obtained from two distinct donors to the Birth Defects Research Laboratory at the University of Washington (UW) and cultured as previously described[9, 15]. RPE cells at passages 4-6 were used for experiments, and they were seeded at a density of either 5×10^4 cells/well (regular density) or 5×10^3 cells/well (low density) in 96-well plates in MEM RPE media (See details in **Table S1**). iPSC RPE, one SFD patient-derived iPSC RPE cells and CRISPR-corrected isogenic cSFD RPE cells were generated in a stepwise procedure as reported (see details in Supplemental Methods)[12]. Human ARPE-19 cells obtained from the American Type Culture Collection (ATCC) were used at passages 5-8 for the experiments. The cells were seeded at 2×10^4 cells/well in 96-well plates and cultured in three different media to mimic dedifferentiated and differentiated states: 1) DMEM/F-12 media with 5% FBS, 2) MEM RPE media, and 3) MEM-NAM (MEM RPE media supplemented with 10 mM NAM as reported[13, 14]. All RPE cells were cultured for 4 weeks metabolic screening.

Metabolic screening with Biolog Phenotyping MicroArrays

Biolog Phenotyping MicroArrays (Biolog Inc.) are 96-well plate assays that measure cellular metabolic activity in utilizing various carbon and nitrogen substrates. Each well contains a specific nutrient at 100 µM concentration, except for the negative control. Cells are added to the 96-well in Biolog's proprietary nutrient-limiting media, followed by the addition of a proprietary redox dye. This dye is reduced by cellular NADH generated by substrate utilization, causing a color change measured by a plate reader. High absorbance indicates greater metabolic activity in utilizing specific substrate through NADH production. We used two Biolog Phenotype MicroArrays (PM-M1 for carbon sources and PM-M2 for nitrogen sources) to assess nutrient utilization in RPE cells (see details in **Table S1-3**). After pre-incubation with 60 uL of nutrient-limiting media (IFM 1, glutamine 200 µM, 1% dialyzed FBS, and 1% penicillin and streptomycin) for one hour, RPE cell-containing plates were washed with PBS and incubated with transferred pre-incubation media from PM-M1 or PM-M2 microarrays for 40 hours at 37°C in a 5% CO_2 incubator. The redox dye mix MA (10 µL) was then added, and the plates were sealed and read at an absorbance of 595 nm every 15 minutes for 4 hours.

^{13}C glutamine tracing in fRPE cells

Human fRPE cells seeded at regular and low densities were cultured for 4 weeks and changed into

DMEM with 5 mM glucose and 1 mM ^{13}C glutamine. After incubation for 48 hours, the RPE cells and

media were collected for metabolite extraction and analysis of ^{13}C glutamine-derived metabolites with gas

chromatography mass spectrometry (GC MS) as previously reported[9, 10].

Statistics

All data are expressed as the mean ± standard deviation. Fold change of absorbance over negative control

without nutrient source >1.5, or $p<0.05$ by Student's unpaired two-tailed t-tests using GraphPad Prism

9.0, was considered significant. Heat maps were generated in Microsoft Excel.

Results

Mature human fRPE cells demonstrate remarkable metabolic flexibility in nutrient utilization.

To study nutrient utilization in mature fRPE cells, fRPE were cultured for 4 weeks to maturity. RPE

demonstrated characteristic cobblestone morphology and pigmentation (Supplemental **Fig S1A-C**). The

mature RPE cultures were switched to nutrient-limiting media with or without a specific nutrient from the

PM-M1 plate to screen the utilization of 91 carbon sources (**Fig 1A, Table S2**). Remarkably, mature RPE

could utilize 23 carbon sources including sugars, glycolysis intermediates (such as sugar phosphates,

lactate, and pyruvate), TCA cycle intermediates, nucleosides, fatty acids and ketone bodies (**Fig 1B-C**). In

addition to glucose, mature RPE had the capacity to utilize other sugars such as fructose, mannose, and

galactose (**Fig 1B-C**). Nucleosides, containing ribose moiety, can serve as alternative energy sources.

Mature RPE could robustly utilize inosine, adenosine and uridine.

Similar to PM-M1 carbon source screening, we evaluated the utilization of 92 nitrogen sources, including

27 amino acids and 60 dipeptides in mature RPE using the PM-M2 plate as illustrated in **Fig 1D**. Mature

RPE could metabolize 7 amino acids and 17 dipeptides (**Fig 1E**). Except for methionine and tryptophan,

the remaining 5 amino acids (proline, ornithine, glutamine, glutamate, and aspartate) follow a similar

metabolic pathway in the generation of α-ketoglutarate (αKG) or oxaloacetate, which serve as fuels for

the TCA cycle (**Fig 1E, 1F**). Notably, glutamine, glutamate, aspartate, and proline are also present in the

17 dipeptides. These findings suggest that mature fRPE cells have remarkable metabolic flexibility,

enabling them to effectively utilize diverse nutrient sources to fuel their mitochondrial metabolism (**Fig

1C, 1E**).

Dedifferentiated human fRPE cells lose metabolic flexibility, primarily relying on sugars and glutamine

as their main nutrient sources.

Seeding fRPE at low density induces their dedifferentiation into a fibroblast-like phenotype[5]. To

understand nutrient utilization in dedifferentiated RPE cells, we seeded fRPE at 10% of regular density

and cultured them for 4 weeks. As expected, these cells transitioned from their characteristic cobblestone

structure to a fibroblast-like morphology, with upregulated expression of EMT markers including α-

smooth muscle actin (αSMA) and vimentin (**Fig 2A**, Supplemental **Fig S1D-F, Fig S2**). Total protein

content was similar in between mature and dedifferentiated RPE, indicating similar confluence between

culture conditions (**Fig S3**). Compared to mature fRPE, dedifferentiated fRPE utilized fewer nutrients,

with only 8 carbon and 6 nitrogen sources. Intriguingly, almost all carbon sources were sugar and sugar

phosphates, while all nitrogen sources contained glutamine (**Fig 2A-C**). These results suggest that

dedifferentiated RPE cells lose the ability to utilize multiple nutrient sources such as lactate, fatty acids,

ketone bodies and proline. Instead, they become largely restricted to consuming sugars and glutamine

(**Fig S4**). RPE seeded at low density for 1 week displayed a fibroblast-like morphology resembling that of

4-week cultured low-density fRPE cells but expressed less αSMA (**Fig S5A-B**, **Fig S1D-F, Fig S2**).

Nutrient utilization in 1-week cells was largely consistent with those of 4-week fRPE, except for the

capability to utilize galactose and adenosine at 1 week (**Fig S5B**), suggesting that time in culture has

minimal impact on the metabolic phenotype of nutrient utilization in RPE cells seeded at low density.

Consequently, we did not perform PM-M2 screening on 1-week fRPE.

To investigate how dedifferentiated RPE cells utilize glutamine, we incubated mature and dedifferentiated

RPE with ^{13}C glutamine for 48 hours and analyzed ^{13}C glutamine-derived intracellular and extracellular

metabolites (Fig 2D-E, Fig S6). Dedifferentiated RPE consumed more ^{13}C glutamine in the media, which

was utilized to produce mitochondrial intermediates and amino acids, especially succinate and proline

(Fig 2D-E, Fig S6). These results suggest that dedifferentiated RPE cells have an impaired electron

transport chain and reprogram their metabolism through reverse succinate dehydrogenase and proline

dehydrogenase towards succinate and proline production.

We next analyzed the expression of metabolic genes related to nutrient utilization in two RNA-Seq data

sets: 1) mature fRPE compared to dedifferentiated RPE induced by multiple passages[16] and 2) mature

primary adult RPE compared to dedifferentiated RPE induced by transforming growth factor-beta (TGFβ)

and tumor necrosis factor α (TNFα)[17]. Both dedifferentiated RPE models had significant downregulation

of genes involved in fatty acid oxidation, TCA cycle, transporters for lactate and ketone bodies,

nucleotide metabolism, and amino acid metabolism, especially proline catabolism and glutamine

synthetase (Fig 2F-G, **Table S4**). These data corroborate our findings that dedifferentiated RPE cannot

efficiently utilize fatty acids, lactate, ketone bodies, proline and other nucleosides for mitochondrial

metabolism. Importantly, genes involved in proline synthesis, glycolysis, collagen and heparin sulfate

were substantially upregulated (**Fig 2F**, **Table S4**), suggesting dedifferentiated RPE cells utilize sugar and

glutamine to activate proline synthesis and the hexosamine biosynthetic pathway to produce collagen and

glycan for ECM remodeling **(Fig 2F)**.

Human iPSC RPE cells have similar metabolic flexibility to fRPE cells but utilize more sugars and TCA

cycle intermediates.

To study nutrient utilization in human iPSC RPE cells, we cultured iPSC RPE to maturity at 4 weeks (**Fig**

S7) and then replaced media with carbon or nitrogen sources from PM-M1 or PM-M2 for metabolic

phenotyping (**Fig 3A**). iPSC RPE could utilize 31 carbon sources and 17 nitrogen sources (**Fig 3B-D**).

Similar to fRPE, iPSC RPE demonstrated robust flexibility in utilizing various sugars, fatty acids, lactate,

ketone bodies, nucleosides, amino acids, and dipeptides (**Fig 3B-D**). However, unlike fRPE, iPSC RPE

showed a utilization preference for more sugars and TCA cycle intermediates and fewer amino acids and

dipeptides (**Fig 3E-F**). These results suggest both healthy fRPE and iPSC RPE possess metabolic

flexibility in adapting to different substrates, yet also have cell type-specific preferences for certain

substrates.

SFD iPSC RPE cells utilize fewer intermediates from glycolytic and TCA cycle intermediates but more

branch-chain amino acids (BCAAs).

We reported SFD iPSC RPE cells harboring the *TIMP3* S204 mutation have increased extracellular

deposits and elevated intracellular 4-hydroxyproline[12]. These iPSC RPE and their isogenic controls

(cSFD) were grown to maturity at 4 weeks before undergoing metabolic phenotyping with PM-M1 and

PM-M2 (**Fig 4A**). Both SFD and cSFD had normal cobblestone structure and pigmentation (**Fig S8A-B**).

SFD RPE utilized 19 carbon sources and 19 nitrogen sources, while cSFD RPE utilized 28 carbon sources

and 7 nitrogen sources (**Fig 4B-E**). SFD RPE utilized different types of sugar, nucleosides, fatty acids,

ketone bodies, amino acids and dipeptides (**Fig 4B, 4D**). However, SFD RPE could not utilize lactate,

sugar phosphate, short-chain fatty acids (acetate and butyrate), succinate, αKG and gamma-aminobutyric

acid (GABA). Notably, the CRISPR-corrected RPE cells restored the utilization of these nutrients (**Fig**

4C-E). SFD RPE could use proline and dipeptides containing glutamine, alanine and arginine.

Intriguingly, many of the dipeptides utilized by SFD RPE cells contained BCAAs, leucine, isoleucine and

valine, along with the ability to catabolize free leucine (**Fig 4D**). These metabolic phenotypes are

different from fRPE and normal control iPSC RPE. While the cSFD RPE cells did not use the BCAAs,

they also utilized significantly fewer nitrogen sources (**Fig 4E, Fig S8C**). These results suggest that SFD

RPE cells have metabolic defects in recycling nutrients, such as lactate from the neural retina, and rely

more on BCAAs, which may contribute to the formation of sub-RPE deposits of lipids and ECM proteins

(**Fig 4F**).

Culture media composition controls the differentiation and nutrient utilization in ARPE-19 cells.

ARPE-19 cells, typically cultured in DMEM/F-12 media per manufacturer's instructions, undergo rapid

differentiation into a mature RPE-like phenotype grown MEM-α-based RPE culture media supplemented

with NAM, [13, 14]. To investigate the influence of media composition on nutrient utilization, we cultured

ARPE-19 cells under three media (DMEM/F12, MEM RPE media and MEM RPE media plus NAM) for

4 weeks and focused only on the screening of carbon sources using the PM-M1 plate (**Fig 5A**). DMEM/F-

12 cultured cells (fibroblast-like morphology, **Fig S9A**) utilized 16 carbon sources (**Fig 5B**) but, similar to

dedifferentiated fRPE, could not metabolize lactate, ketone bodies, or xylitol (**Fig 5B, Fig 6A**). MEM

RPE media partially differentiated ARPE-19 cells (cobblestone structure, **Fig S9B**) and increased their

carbon source utilization to 24 (covering 94% of those used by DMEM/F12 cells, Fig 5C). While they

gained the ability to utilize ketone bodies and some sugars, lactate and xylitol remained unutilized (**Fig

5C**). Supplementation with NAM in MEM RPE media induced mature RPE-like morphology (**Fig S9C**)

and allowed ARPE-19 cells to utilize lactate, ketone bodies, sugar phosphates, xylose, and tagatose (**Fig

5D-E**). This metabolic phenotype resembles that of mature fRPE and iPSC RPE, although ARPE-19 cells

reduced utilization of short-chain fatty acids and maltose (**Fig 6A**). These findings highlight the critical

role of nutrient availability and utilization in RPE differentiation.

Discussion

In this study, we identified distinctive metabolic phenotypes of nutrient utilization in healthy and diseased

human RPE cells (**Fig 6**). Mature fRPE and iPSC RPE are metabolically flexible, capable of adapting to

various nutrient sources. Conversely, dedifferentiated RPE cells and SFD patient-derived iPSC RPE cells

have reduced flexibility and are biased towards fewer, specific substrates. Our findings suggest the critical

roles of nutrient utilization in RPE differentiation and dedifferentiation.

The RPE must manage a substantial nutrient influx originating daily from phagocytosed lipid- and

protein-rich photoreceptor outer segments, "waste products" such as lactate produced by the neural retina,

and nutrient-laden blood through the choroidal circulation[1, 2]. In addition to the direct transport of

nutrients to the neural retina, RPE can recycle and generate nutrients via lysosomes and mitochondria to

support retinal metabolism, thereby establishing a metabolic ecosystem between the outer retina and

RPE[1, 2, 18]. The ability to utilize a variety of nutrients is crucial for maintaining this metabolic ecosystem,

and disruption of this balance due to mitochondrial and/or lysosomal dysfunction in RPE can lead to

retinal degeneration[3, 4, 19]. Healthy human RPE cells are known to be versatile in utilizing different

nutrients, including glucose, galactose, lactate, fatty acids, succinate, proline, glutamine and other amino

acids[9, 18, 20-23]. Our results confirm this versatility, demonstrating that both healthy fRPE and iPSC RPE

cells are capable of utilizing these nutrients.

Furthermore, we found that human RPE cells utilize a wide range of other nutrients including various

sugars, ketone bodies, nucleosides and dipeptides. Mannose and galactose, for example, are components

of glycoproteins found in abundance in outer segment proteins such as rhodopsin and peripherin 2[24-26].

The ability to utilize these sugars from the daily degradation of outer segments may allow RPE to

preserve more glucose to supply for photoreceptors. Additionally, like the liver, human RPE cells can

synthesize ketone bodies, which can be utilized by the neural retina[21, 27]. Interestingly, we found that

mature human RPE, but not dedifferentiated RPE cells, can readily utilize ketone bodies, suggesting that

the ability to utilize ketone bodies might be a characteristic of RPE metabolism. Consistent with this

metabolic capability, mature RPE expresses key genes involved in ketone body degradation[21], a feature

distinct from the liver, which cannot utilize ketone bodies. Our findings underscore the metabolic

flexibility of healthy mature RPE in nutrient utilization, which is an essential function for RPE to support

retinal metabolism and health.

RPE EMT and dedifferentiation have been implicated in the pathogenesis of proliferative

vitreoretinopathy and AMD[28]. We found dedifferentiated RPE cells were unable to utilize alternative

fuels like lactate, fatty acids, ketone bodies and proline, but, instead, prefer to utilize sugars and

glutamine. RNA-seq data from EMT models show reduced expression of genes involved in the TCA

cycle, lactate transport, ketone body utilization and proline catabolism and glutamine synthesis. In

contrast, gene expression associated with glycolysis, proline synthesis, collagen synthesis and

glycoproteins are upregulated (**Fig 2E, Table S4**). Consistently, dedifferentiated RPE cells have been

reported to switch from oxidative phosphorylation to glycolysis and lose the ability to utilize proline as a

fuel[5, 15]. Inhibition of mitochondrial metabolism in human RPE cells or mouse RPE is sufficient to cause

the activation of glycolysis, RPE dedifferentiation and retinal degeneration[4, 29]. Furthermore,

mitochondrial respiration is diminished in cultured primary RPE cells from AMD donors[30]. Inhibiting

mitochondrial respiration impedes proline consumption while increasing glucose consumption and lactate

production[31]. The inability of affected RPE to utilize exogenous lactate and other alternative nutrient

sources may be an important mechanism in AMD pathogenesis.

A key feature in dedifferentiated cells is their overproduction of ECM proteins, primarily collagens and

glycoproteins, through activation of cytokine signaling such as TGFβ and TNFα[32, 33]. In fibroblasts, TGFβ

stimulates the oxidation of glucose and glutamine to synthesize proline and other amino acids crucial for

ECM protein production[34]. Interestingly, proline synthesis acts as a vent for growth when cells are under

mitochondrial redox stress or hypoxia by recycling NADH into NAD$^+$, thus enhancing the oxidation of

glutamine and glucose[34, 35]. Proline synthesis may be particularly crucial in dedifferentiated RPE cells,

given their inhibited mitochondrial metabolism and heavy reliance on lactate production for NAD$^+$

regeneration. Similarly, to compensate for the deficiency in the electron transport chain or hypoxia,

fumarate can serve as an electron acceptor to produce succinate[36], which is accumulated in

dedifferentiated RPE. Strategies aimed at relieving mitochondrial redox stress and reverting proline

synthesis to proline catabolism or succinate production could hold promise for treating RPE

dedifferentiation.

TIMP3, a risk gene for AMD, is secreted by the RPE and plays a critical role in ECM remodeling by

inhibiting matrix metalloproteinases[37]. Specific mutations in *TIMP3* result in SFD, which is characterized

by a thickened layer of basal laminar drusen-like sub-RPE deposits containing lipids and ECM proteins[12,

38]. Unlike healthy or mutation-corrected RPE cells, SFD RPE cells utilize more free BCAAs or dipeptides

containing BCAAs (**Fig 4F, Fig 6**). Leucine and isoleucine are ketogenic amino acids, and their oxidation

promotes the synthesis and transport of fatty acids and cholesterol[39, 40]. BCAA oxidation is closely

associated with dysregulated glucose and lipid metabolism in diabetes and non-alcoholic fatty liver

diseases[39, 41, 42]. Interestingly, intermediates of the BCAA degradation pathway are significantly elevated

in the plasma of AMD patients[43], and *BCAT1*, a key gene in BCAA degradation, is upregulated in the

RPE/choroid from AMD donors[44]. Moreover, in addition to proline, ECM proteins are highly enriched in

BCAAs[45]. Our findings suggest that SFD RPE may upregulate the BCAA degradation pathway by

utilizing BCAAs from free amino acids or protein degradation, contributing to the formation of sub-RPE

deposits.

There are a few limitations in this study. We did not normalize nutrient utilization to specific cell

numbers, although total cellular protein content was similar between dedifferentiated and mature RPE by

4 weeks in culture. Nutrient utilization in additional dedifferentiation models need to be conducted in

future studies. Separately, high absorbance reflects greater reduction of NADH generated from a specific

nutrient. The Biolog assays cannot detect the reduction of FADH2 or other metabolic pathways such as

the synthesis of amino acids, nucleotides and fatty acids. In addition, the metabolic phenotyping assay

can only measure the capability and capacity of utilizing a specific nutrient but not preferences of nutrient

utilization because only one nutrient is available at a time. Therefore, the utilization might be different *in*

vivo, and additional functional studies with nutrient supplements or genetic manipulations of metabolic

pathways are needed to interpret these findings. Finally, testing additional iPSC RPE lines from a larger

cohort of normal and affected donors will help control for variability in genetic and environmental

backgrounds.

In conclusion, metabolic phenotyping offers a sensitive and powerful platform for measuring nutrient

utilization in healthy and diseased human RPE cells. The distinct patterns of nutrient utilization of various

RPE cells provide valuable insights into the mechanisms of RPE dedifferentiation and disease.

Acknowledgments

This work was supported by National Institutes of Health Grant EY034364 (J. R. C. and J. D), EY03459

(J. R. C. and J. D), EY031324 (JD), EY032462(JD), the Retina Research Foundation (JD), and funds for

Core facilities P20 GM103434 and P20 GM144230 (WV INBRE grant). Fetal RPE tissue samples were

made available through Dr. Ian Glass and the Birth Defects Research Laboratory at the University of

Washington (UWR24HD000836).

Conflicts of interest

The authors declare no conflicts of interest.

References

1.   Lakkaraju A, Umapathy A, Tan LX, et al. The cell biology of the retinal pigment epithelium. *Prog*
Retin Eye Res 2020;100846.
2.   Hurley JB. Retina Metabolism and Metabolism in the Pigmented Epithelium: A Busy Intersection.
Annu Rev Vis Sci 2021;7:665-692.
3.   Ferrington DA, Fisher CR, Kowluru RA. Mitochondrial Defects Drive Degenerative Retinal
Diseases. *Trends Mol Med* 2020;26:105-118.
4.   Zhao C, Yasumura D, Li X, et al. mTOR-mediated dedifferentiation of the retinal pigment
epithelium initiates photoreceptor degeneration in mice. *J Clin Invest* 2011;121:369-383.
5.   Adijanto J, Philp NJ. Cultured primary human fetal retinal pigment epithelium (hfRPE) as a model
for evaluating RPE metabolism. *Exp Eye Res* 2014;126:77-84.
6.   Bharti K, den Hollander AI, Lakkaraju A, et al. Cell culture models to study retinal pigment
epithelium-related pathogenesis in age-related macular degeneration. *Exp Eye Res* 2022;222:109170.
7.   Nazari H, Zhang L, Zhu D, et al. Stem cell based therapies for age-related macular degeneration:
The promises and the challenges. *Prog Retin Eye Res* 2015;48:1-39.

8.   Blenkinsop TA, Saini JS, Maminishkis A, et al. Human Adult Retinal Pigment Epithelial Stem Cell-
Derived RPE Monolayers Exhibit Key Physiological Characteristics of Native Tissue. *Invest Ophthalmol Vis*
Sci 2015;56:7085-7099.
9.   Chao JR, Knight K, Engel AL, et al. Human retinal pigment epithelial cells prefer proline as a
nutrient and transport metabolic intermediates to the retinal side. *J Biol Chem* 2017;292:12895-12905.
10.   Du J, Zhu S, Lim RR, Chao JR. Proline metabolism and transport in retinal health and disease.
Amino Acids 2021;53:1789-1806.
11.   Anand-Apte B, Chao JR, Singh R, Stohr H. Sorsby fundus dystrophy: Insights from the past and
looking to the future. *J Neurosci Res* 2019;97:88-97.
12.   Engel AL, Wang Y, Khuu TH, et al. Extracellular matrix dysfunction in Sorsby patient-derived
retinal pigment epithelium. *Exp Eye Res* 2022;215:108899.
13.   Hazim RA, Volland S, Yen A, Burgess BL, Williams DS. Rapid differentiation of the human RPE cell
line, ARPE-19, induced by nicotinamide. *Exp Eye Res* 2019;179:18-24.
14.   Hazim RA, Paniagua AE, Tang L, et al. Vitamin B3, nicotinamide, enhances mitochondrial
metabolism to promote differentiation of the retinal pigment epithelium. *J Biol Chem* 2022;298:102286.
15.   Yam M, Engel AL, Wang Y, et al. Proline mediates metabolic communication between retinal
pigment epithelial cells and the retina. *J Biol Chem* 2019;294:10278-10289.
16.   Radeke MJ, Radeke CM, Shih YH, et al. Restoration of mesenchymal retinal pigmented epithelial
cells by TGFbeta pathway inhibitors: implications for age-related macular degeneration. *Genome Med*
2015;7:58.
17.   Boles NC, Fernandes M, Swigut T, et al. Epigenomic and Transcriptomic Changes During Human
RPE EMT in a Stem Cell Model of Epiretinal Membrane Pathogenesis and Prevention by Nicotinamide.
Stem Cell Reports 2020;14:631-647.
18.   Kanow MA, Giarmarco MM, Jankowski CS, et al. Biochemical adaptations of the retina and
retinal pigment epithelium support a metabolic ecosystem in the vertebrate eye. *Elife* 2017;6.
19.   Boya P, Kaarniranta K, Handa JT, Sinha D. Lysosomes in retinal health and disease. *Trends*
Neurosci 2023;46:1067-1082.
20.   Liu B, Calton MA, Abell NS, et al. Genetic analyses of human fetal retinal pigment epithelium
gene expression suggest ocular disease mechanisms. *Commun Biol* 2019;2:186.
21.   Adijanto J, Du J, Moffat C, Seifert EL, Hurle JB, Philp NJ. The retinal pigment epithelium utilizes
fatty acids for ketogenesis. *J Biol Chem* 2014;289:20570-20582.
22.   Bisbach CM, Hass DT, Robbings BM, et al. Succinate Can Shuttle Reducing Power from the
Hypoxic Retina to the O(2)-Rich Pigment Epithelium. *Cell Rep* 2020;31:107606.
23.   Li B, Zhang T, Liu W, et al. Metabolic Features of Mouse and Human Retinas: Rods versus Cones,
Macula versus Periphery, Retina versus RPE. *iScience* 2020;23:101672.
24.   Skiba NP, Lewis TR, Spencer WJ, Castillo CM, Shevchenko A, Arshavsky VY. Absolute
Quantification of Photoreceptor Outer Segment Proteins. *J Proteome Res* 2023;22:2703-2713.
25.   Stuck MW, Conley SM, Naash MI. PRPH2/RDS and ROM-1: Historical context, current views and
future considerations. *Prog Retin Eye Res* 2016;52:47-63.
26.   Daniele LL, Han JYS, Samuels IS, et al. Glucose uptake by GLUT1 in photoreceptors is essential for
outer segment renewal and rod photoreceptor survival. *FASEB J* 2022;36:e22428.
27.   Reyes-Reveles J, Dhingra A, Alexander D, Bragin A, Philp NJ, Boesze-Battaglia K. Phagocytosis-
dependent ketogenesis in retinal pigment epithelium. *J Biol Chem* 2017;292:8038-8047.
28.   Zhou M, Geathers JS, Grillo SL, et al. Role of Epithelial-Mesenchymal Transition in Retinal
Pigment Epithelium Dysfunction. *Front Cell Dev Biol* 2020;8:501.
29.   Hu X, Calton MA, Tang S, Vollrath D. Depletion of Mitochondrial DNA in Differentiated Retinal
Pigment Epithelial Cells. *Sci Rep* 2019;9:15355.

30.   Fisher CR, Ferrington DA. Perspective on AMD Pathobiology: A Bioenergetic Crisis in the RPE.
Invest Ophthalmol Vis Sci 2018;59:AMD41-AMD47.
31.   Zhang R, Engel AL, Wang Y, et al. Inhibition of Mitochondrial Respiration Impairs Nutrient
Consumption and Metabolite Transport in Human Retinal Pigment Epithelium. *J Proteome Res*
2021;20:909-922.
32.   Scott LE, Weinberg SH, Lemmon CA. Mechanochemical Signaling of the Extracellular Matrix in
Epithelial-Mesenchymal Transition. *Front Cell Dev Biol* 2019;7:135.
33.   Shu DY, Butcher E, Saint-Geniez M. EMT and EndMT: Emerging Roles in Age-Related Macular
Degeneration. *Int J Mol Sci* 2020;21.
34.   Schworer S, Berisa M, Violante S, et al. Proline biosynthesis is a vent for TGFbeta-induced
mitochondrial redox stress. *EMBO J* 2020;39:e103334.
35.   Westbrook RL, Bridges E, Roberts J, et al. Proline synthesis through PYCR1 is required to support
cancer cell proliferation and survival in oxygen-limiting conditions. *Cell Rep* 2022;38:110320.
36.   Spinelli JB, Rosen PC, Sprenger HG, et al. Fumarate is a terminal electron acceptor in the
mammalian electron transport chain. *Science* 2021;374:1227-1237.
37.   Alexander JP, Bradley JM, Gabourel JD, Acott TS. Expression of matrix metalloproteinases and
inhibitor by human retinal pigment epithelium. *Invest Ophthalmol Vis Sci* 1990;31:2520-2528.
38.   Weber BH, Vogt G, Pruett RC, Stohr H, Felbor U. Mutations in the tissue inhibitor of
metalloproteinases-3 (TIMP3) in patients with Sorsby's fundus dystrophy. *Nat Genet* 1994;8:352-356.
39.   Neinast MD, Jang C, Hui S, et al. Quantitative Analysis of the Whole-Body Metabolic Fate of
Branched-Chain Amino Acids. *Cell Metab* 2019;29:417-429 e414.
40.   Jang C, Oh SF, Wada S, et al. A branched-chain amino acid metabolite drives vascular fatty acid
transport and causes insulin resistance. *Nat Med* 2016;22:421-426.
41.   White PJ, McGarrah RW, Grimsrud PA, et al. The BCKDH Kinase and Phosphatase Integrate BCAA
and Lipid Metabolism via Regulation of ATP-Citrate Lyase. *Cell Metab* 2018;27:1281-1293 e1287.
42.   White PJ, Lapworth AL, An J, et al. Branched-chain amino acid restriction in Zucker-fatty rats
improves muscle insulin sensitivity by enhancing efficiency of fatty acid oxidation and acyl-glycine
export. *Mol Metab* 2016;5:538-551.
43.   Haijes HA, Jans JJM, van der Ham M, van Hasselt PM, Verhoeven-Duif NM. Understanding acute
metabolic decompensation in propionic and methylmalonic acidemias: a deep metabolic phenotyping
approach. *Orphanet J Rare Dis* 2020;15:68.
44.   Newman AM, Gallo NB, Hancox LS, et al. Systems-level analysis of age-related macular
degeneration reveals global biomarkers and phenotype-specific functional networks. *Genome Med*
2012;4:16.
45.   Li P, Wu G. Roles of dietary glycine, proline, and hydroxyproline in collagen synthesis and animal
growth. *Amino Acids* 2018;50:29-38.
461

Figure legends

Figure 1. Metabolic phenotyping of mature human fRPE cells. (A) A schematic for metabolic

phenotyping with carbon sources. Human fRPE cells were grown for 4 weeks in a 96-well plate and then

switched into nutrient-limiting media containing different carbon sources in each well from PM-M1 plate.

The utilization of nutrients leads to NADH production, casuing a color change in a redox sensitive dye to

purple, which is quantified by a microplate reader at 595 nm. (B) The utilization of carbon sources by

mature human fRPE cells and (C) an illustration of the metabolic pathways. (D) A schematic for

metabolic phenotyping of nitrogen sources using PM-M2. (E) The utilization of nitrogen sources by

mature human fRPE cells and (F) an illustration of the metabolic pathways. N=3. Fold change over

negative control without nutrient source (Ctr)>1.5 or *P<0.05 over Ctr. Gluc, Glucose; Gal, galactose;

Mann, Mannose; F1P, fructose 1-phosphophate; R1P, ribose 1-phosphate, G6P, glucose 3-phosphate;

G3P, glycerol 3-phosphate; AcAc, Acetoacetic Acid; 3HB, β-Hydroxy Butyrate; MelA, Melibionic Acid.

Figure 2. **Metabolic phenotyping of dedifferentiated human fRPE.** (A) Human fRPE were seeded at

low density to induce dedifferentiation with a mesenchymal phenotype in a 96-well plate and then

switched into nutrient-limiting media containing different carbon (PM-M1) or nitrogen sources (PM-M2)

for metabolic phenotyping. (B) Nutrient utilization of carbon sources by dedifferentiated human fRPE.

N=4. Fold change>1.5 or P<0.05 over Ctr. (C) Nutrient utilization of nitrogen sources by dedifferentiated

human fRPE. N=3. Fold change>1.5 or P<0.05 over Ctr. (D-E) Dedifferentiated RPE consumes more

glutamine for their mitochondrial metabolism. ^{13}C-glutamine and its derived metabolites in media (D) and

RPE cells (E) were analyzed by GC MS and presented as fold change of ^{13}C labeled abundance over low

density RPE. N=4. *P<0.05 vs mature RPE. (F) Heat maps of altered metabolic genes in dedifferentiated

RPE cells induced by TNT (left panel) or multiple passage (right panel) (G) The dedifferentiated RPE

cells lose the ability to utilize multiple nutrient sources but switch to use glutamine, enhancing proline

and collagen synthesis. Magenta denotes an increase and blue denotes a decrease in gene expression or

metabolic pathways. *PFKFB4*, 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 4, *LDHA*, Lactate

dehydrogenase A; *OGDH*, Oxoglutarate dehydrogenase; *PRODH*; Proline dehydrogenase; *PYCR1*,

Pyrroline-5-carboxylate reductase 1; P4H4, Prolyl 4-hydroxylase subunit alpha or beta; *GMPPA*, GDP-

mannose pyrophosphorylase A. *PMM2*, Phosphomannomutase 2.

Figure 3. Metabolic phenotyping of healthy human iPSC RPE cells. (A) Human iPSC RPE cells were

differentiated in RPE media for 4 weeks and switched into nutrient-limit media from PM1 or PM2 for

metabolic phenotyping. (B-D) Nutrient utilization of carbon sources and nitrogen sources by human iPSC

RPE cells. N=3. Fold change>1.5 or P<0.05 over Ctr. (E) A comparison of number of nutrients that were

utilized between fRPE and iPSC RPE cells. (F) An illustration of nutrient utilization in iPSC RPE cells.

GlcNAc, N-Acetyl-D-Glucosamine; Tcba, Tricarballylic Acid; AlaN, L-Alaninamide.

Figure 4. Metabolic phenotyping of IPSC SFD and cSFD RPE cells. (A) Human iPSC SFD RPE

carrying TIMP3 S204C mutation and mutation corrected cSFD RPE cells were differentiated in RPE

media for 4 weeks and then switched to nutrient-limit media with carbon or nitrogen sources from PM-

M1 or PM-M2 for metabolic phenotyping. (B-E) Nutrient utilization of (B-C) carbon sources and (D-E)

nitrogen sources in iPSC SFD RPE cells. N=3. Fold change>1.5 or $P<0.05$ over Ctr. (F) An illustration of

altered nutrient utilization in SFD iPSC RPE cells. L-Mal, L-Malic Acid; mmSuc, mono methyl

succinate; 3HB, β-Hydroxy Butyrate; BCAAs, Branch-Chained Amino Acids.

Figure 5. Metabolic phenotyping of ARPE-19 cells cultured in different media.

(A) ARPE-19 cells were cultured in three different media: DMEM/F-12, MEM RPE media, and MEM

RPE media with 10 mM NAM for 4 weeks. All the cells were then changed into nutrient-limit media with

carbon sources from PM-M1 for metabolic phenotyping. (B-D) Nutrient utilization of carbon sources of

ARPE-19 cells cultured in (B) DMEM/F12 (C) MEM RPE media, (D) MEM RPE media with NAM.

Fold change>1.5 or $P<0.05$ over Ctr. (E) A comparison of number of nutrients that were utilized in

ARPE-19 cells grown in different culture media. (F-G) An illustration of altered nutrient utilization in

APRE-19 cells cultured between DMEM/F12 and MEM RPE media with NAM. Meso-Tar, Meso Tartatic

acid; mmSuc, mono methyl succinate; AcAc, Acetoacetic Acid; 3HB, β-Hydroxy Butyrate.

Figure 6. Comparison of metabolic phenotyping profiles of human fRPE cells, dedifferentiated

fRPE, iPSC RPE cells, SFD RPE, cSFD RPE and ARPE-19 cells grown under different culture

media. Heat map visualization of nutrients that were differentially utilized by different RPE cells from

(A) PM-M1 nutrients and (B) PM-M2 nutrient microarray plates. De-fRPE, dedifferentied fRPW with

low density seedings; ARPE, ARPE-19 cells; MelA, Melibionic Acid; GlcNAc, N-Acetyl-D-

Glucosamine; Tcba, Tricarballylic acid; mmSuc, mono methyl succinate; AlaN, Alaninamide.