Cell morphogenesis via shape-guided self-propelled treadmilling actin waves

Kio Yagami[1], Kentarou Baba[1], Shinji Misu[1], Hiroko Katsuno-Kambe[1], Takunori Minegishi[1], Kazunori Okano[2], Yuichi Sakumura[3], Yoichiroh Hosokawa[2] and Naoyuki Inagaki[1,*]

[1]Laboratory of Systems Neurobiology and Medicine, Division of Biological Science, Nara Institute of Science and Technology, Ikoma, Nara 630-0192, Japan.

[2]Bio-Process Engineering Laboratory, Division of Materials Science, Nara Institute of Science and Technology, Ikoma, Nara 630-0192, Japan.

[3]Laboratory of Data-driven Biology, Division of Biological Science, Nara Institute of Science and Technology, Ikoma, Nara 630-0192, Japan.

*e-mail: ninagaki@bs.naist.jp

Abstract

Actin dynamics mediate cell morphogenesis and motility. Actin filaments polymerise outward at cell protrusions such as the leading edge of migrating cells, thereby pushing the membrane to protrude. The current paradigm explains that actin dynamics are biochemically regulated by signalling pathways. However, it remains unclear whether these signal-driven mechanisms alone are sufficient to regulate local actin polymerisation at cell protrusions. Here, we show that cell shape can also regulate actin dynamics. We found that arrays of actin filaments emerge widely in glioma cells and translocate as actin waves in the direction of polymerisation through treadmilling, i.e. directional polymerisation and disassembly. The arrival of actin waves at the cell periphery pushes the plasma membrane to form protrusions (filopodia and lamellipodia). Furthermore, actin waves accumulate at protrusions even in the absence of local signalling cues, similar to self-propelled particles (active particles) colliding with a boundary, thereby leading to further expansion of protrusions for cell polarisation and migration. The shape-guided actin accumulation constitutes a positive feedback interaction with actin-mediated cell protrusion and mediates an effective lateral inhibition, thereby conferring robustness to the classical pathway of cellular morphogenesis. Because actin filaments undergo treadmilling in various protrusions, we propose that the cell shape-guided and self-propelled movement is a basic property of treadmilling actin filaments that mediates robust cell morphogenesis and motility.

Introduction

Spatial regulation of actin dynamics is crucial for controlling force, cell shape and motility[1-4]. Dynamic actin filaments are located in various cell protrusions, where they polymerise outward and push the membrane to protrude. In many cases, they disassemble proximally, which results in treadmilling of the filament arrays[1-5]. During this process, ATP-bound actin monomers assemble at the plus ends of actin filaments and ATP is hydrolysed within the filaments. The filaments disassemble from the minus end because they are destabilised by ADP-bound actin molecules. ADP in the subunit is replaced with ATP, which allows the subunit to be reused for polymerisation. Thus, actin filaments undergo directional polymerisation and disassembly using the energy derived from ATP hydrolysis[1-5]. The prevailing view is that actin filament nucleators and elongators, such as the Arp2/3 complex, formins, FMNL and Ena/VASP, control local actin assembly under Rho GTPase signalling[1,3,4]. On the other hand, actin disassembly-promoting proteins, such as ADF/cofilin, regulate actin turnover[2,5,6]. However, it remains unclear whether these signal-driven mechanisms alone are sufficient to regulate local actin dynamics at cell protrusions.

Accumulating evidence indicates that dynamic actin filaments undergo wave-like propagations in various cell types[7-10]. Many of these propagations have been explained by the reaction-diffusion mechanism[8-11]. On the other hand, we previously reported that actin treadmilling propels the translocation of actin waves and actin-binding proteins towards neurite tips[12]. In this study, we identified actin waves that translocate via treadmilling instead of the reaction-diffusion mechanism in non-neuronal cells. The arrival of actin waves at the cell periphery results in the formation of protrusions. Moreover, the actin waves spontaneously accumulated at cell protrusions even without local signalling in a manner similar to self-propelled particles (active particles), as described in the current literature on physics[13-15] and materials science[16]. This shape-guided and self-propelled movement mediates the accumulation of outward-polymerising actin filaments in protruding regions.

Results

Actin waves emerge widely in cells

To investigate the mechanism underlying the spatial regulation of actin dynamics, we

analysed a subline of glioma U251 cells that actively form lamellipodia (arrows,

Extended Data Fig. 1a) and filopodia (arrowheads). These cells spontaneously polarise

and migrate in the absence of growth factor stimulation (Extended Data Fig. 1b)[17]. Actin

dynamics were visualised by total internal reflection fluorescence (TIRF) microscopy

using the actin filament marker LifeAct (Fig. 1a and Supplementary Video 1). In

addition to prominent actin filaments at the leading edge (asterisk, Fig. 1a), arrays of

actin filaments that form filopodium-like structures (arrowheads, Fig. 1b) and

lamellipodium-like structures (arrows, Fig. 1c) emerged widely in the cytoplasm,

moved in random directions, and disappeared (Supplementary Video 1). Some

filopodium-like structures extended from the lamellipodium-like structures (yellow

arrowheads, Extended Data Fig. 1c, Supplementary Video 2), whereas

lamellipodium-like structures expanded from the filopodium-like structures (blue

arrow), indicating their interchangeability. Similar actin waves were also observed in

COS7 cells (Extended Data Fig. 1d, Supplementary Video 3), suggesting that they are

not cell type-specific. Herein, we analysed the filopodium-like and lamellipodium-like

structures that move within the cells as filopodium-type and lamellipodium-type actin

waves, respectively.

A combination of TIRF microscopy, which visualises the region near the

ventral plasma membrane, and epifluorescence microscopy, which visualises the entire

cell thickness, revealed a large fraction of waves translocating along the ventral plasma

membrane (arrowheads, Extended Data Fig. 1e, Supplementary Video 4), as well as

waves travelling apart from the adhesive substrate (arrowheads, Extended Data Fig. 1f,

Supplementary Video 4). In addition, 3D imaging with confocal deconvolution

microscopy revealed actin waves moving along the dorsal plasma membrane (Fig. 1d)

as well as waves translocating through the cytoplasm without interacting with the

plasma membrane (Fig. 1e, Supplementary Video 5).

Actin waves translocate by treadmilling

Next, we examined the actin dynamics within waves by labelling actin filaments with

EGFP-LifeAct, and monitoring actin molecules in waves by speckle imaging of

HaloTag-actin[18]. During the forward translocation of filopodium-type and

lamellipodium-type actin waves along the ventral membrane (green lines, Fig. 1f and

Extended Data Fig. 2a) individual actin molecules moved retrogradely within the

filaments (magenta lines, Supplementary Video 6). Similar retrograde movements of

actin molecules were observed in filopodium-type waves with no interaction with the

plasma membrane (Extended Data Fig. 2b), indicating that actin waves undergo

treadmilling and translocate in the direction of polymerisation.

We previously reported actin waves that translocate along neuronal nascent

axons through treadmilling[9,12]: their front and rear ends undergo polymerisation and

disassembly, respectively (black arrows, Fig. 1g, left)[9,12]. In addition, actin filaments in

waves are weakly anchored to the plasma membrane and adhesive substrate, through the

clutch-linker molecules shootin1a and cortactin as well as the cell adhesion molecule

L1[12,19] (Fig. 1g, left). This anchoring impedes the retrograde flow (slippage) of

treadmilling actin filaments (magenta arrow, Fig. 1g); thus, the actin

polymerisation/disassembly rate exceeds the retrograde flow rate, which results in the

forward translocation of the front and rear ends of waves (green arrow). During the

translocation, the local concentrations of actin subunits and actin-binding proteins

decrease at the front by polymerisation and increase at the rear by disassembly, resulting

in their directional diffusion along the local concentration gradients (black arrows, Fig.

1g). Thus, accompanied by the directional diffusion, treadmilling actin waves mediate

the translocation of actin and actin-binding proteins along axons[12,20].

Immunoblot analyses revealed that U251 cells express, shootin1b, cortactin and

L1 (Extended Data Fig. 2c); shootin1b is a splicing variant of shootin1a that also

mediates the actin-L1 linkage[21]. Shootin1b co-localised with actin filaments in

lamellipodia (Extended Data Fig. 2d) and waves (arrowheads, Extended Data Fig. 2e).

Inhibition of actin polymerisation in waves by cytochalasin B at a low concentration

(0.05 μM)[12] decreased the velocity of actin wave translocation (Extended Data Fig. 2f).

The Arp2/3 complex inhibitor CK666 (100 μM)[22] or a relatively non-specific formin

inhibitor SMIFH2 (5 μM)[23] also reduced the velocity of actin wave (Fig. 1h), indicating

that wave translocation is propelled by actin assembly. Furthermore, shootin1b

knockout (KO) resulted in an increased retrograde flow rate of actin filaments in waves

without affecting the polymerisation rate (Fig. 1i and Extended Data Fig. 2g), indicating

that shootin1b anchors treadmilling actin filaments to the plasma membrane and

adhesive substrate (Extended Data Fig. 2h). This in turn reduced the velocity of actin

wave translocation (Fig. 1i). Taken together, these results indicate that actin waves in

U251 cells translocate via their treadmilling and that shootin1b promotes their

translocation by impeding the retrograde flow (Fig. 1g, left)[12]. Actin waves may also

move via treadmilling, without anchoring to the membrane (Fig. 1e), when actin

filament retrograde flow is mechanically impeded in a viscous intracellular environment

(Fig. 1g, right; see supplementary discussion).

Actin waves as precursors of protrusions

Interestingly, we observed filopodium-type waves entering into pre-existing

lamellipodia and filopodia (arrowheads, Fig. 2a and b, Supplementary Video 7). The

wave entry resulted in local actin filament accumulation and lamellipodial expansion

(Fig. 2a, Supplementary Video 7). In addition, when lamellipodium-type and

filopodium-type waves arrived at the plasma membrane, which does not form a

protrusion, they pushed the membrane, resulting in the formation of lamellipodia as

reported[24,25] (yellow arrows, Fig. 2c, Supplementary Video 8) and filopodia (arrowheads,

Fig. 2d, Supplementary Video 9), respectively. Some filopodia and lamellipodia

continued to move laterally along the membrane (Fig. 2c, d), consistent with previous

reports that filopodia and lamellipodia move dynamically along the lateral side of

neuronal[20,26] and non-neuronal[27,28] cells. We also noted lamellipodium-type waves that

merged with the pre-existing lamellipodia through their lateral movement (yellow

arrows, Fig. 2c, Supplementary Video 8).

If the lateral movements of filopodia and lamellipodia are driven by the

mechanism of wave translocation, their velocity may be related to the angle θ of the

polymerising actin filaments with respect to the membrane (Fig. 2e). To examine this

possibility, we focused on the motion of filopodium-type waves because

filopodium-type and lamellipodium-type waves are interchangeable (Extended Data Fig.

1c) and it was difficult to monitor the direction of actin filaments in lamellipodium-type

waves (Fig. 1c). Time-lapse measurements of both the angle θ and the velocity of the

filopodial lateral movement (V_L) (Fig. 2f, Supplementary Video 10) demonstrated that

the velocity closely matched the cosine θ for the average velocity of the ventrally

translocating filopodium-type waves before arriving at the membrane (V_V) (2.0 ± 0.2

µm/min, Figs. 1i, 2f), indicating agreement between the data and the prediction based

on the mechanism. Together, these findings suggest that treadmilling actin waves and
filopodia/lamellipodia are equivalent structures and that actin waves serve as precursors
of filopodia and lamellipodia. Therefore, the filopodia and lamellipodia that move
laterally along the membrane are also termed actin waves in the present study.
167
Actin waves accumulate at protrusions
Intriguingly, we observed the accumulation of filopodium-type and lamellipodium-type
actin waves in protrusive regions owing to their lateral movement (Fig. 3a, b,
Supplementary Video 11), suggesting that actin dynamics are influenced by cell shape.
In addition, a previous study reported that different cell types confined to square
adhesive islands preferentially extend filopodia and lamellipodia from their corners[27].
To determine the potential influence of cell shape on actin dynamics, we prepared a
triangular pattern of adhesive laminin on the substrate (Extended Data Fig. 3a). U251
cells showed the triangular shape when cultured on the laminin-coated adhesive island
(Fig. 3c). While actin filaments were preferentially localised at the leading edge of
unconstrained cells (Extended Data Fig. 1a), they accumulated at the corners of
triangular cells (Fig. 3c). Live imaging of U251 cells labelled with LifeAct-mCherry
revealed the dynamic accumulation of actin filaments at the corners (Fig. 3d and
Supplementary Video 12). Actin waves travelled along the lateral edge, resulting in
local accumulation of actin filaments at the corners upon arrival (arrowheads). Notably,
in contrast to the current paradigm that the Arp2/3 complex promotes membrane
protrusion by activating actin polymerisation, a component of the Arp2/3 complex,
ARPC2, accumulated together with actin filaments at the protruding corners in the
absence of local signalling (Fig. 3c).
Since the arrival of actin waves increased actin filaments at the protrusive areas
(arrowheads, Fig. 3d), we hypothesised that actin waves mediate the accumulation of
actin filaments at cell protrusions. To examine this possibility, we constructed a
mathematical model to describe filopodium-type waves because actin filaments were
clearly observed in these waves (Fig. 1b). Our model consists of the following
properties of actin wave dynamics (see Methods).
193
1. Actin waves emerge at random locations within the cell and move in random

directions.

2. Actin waves frequently change their direction of translocation and disappear at the

end of their lifetime.

3. Actin waves re-emerge at the location where the preceding waves disappear.

4. Actin waves located at the cell periphery move laterally along the cell membrane.

The direction and velocity of the lateral movement conform to the $\cos\theta$ of the mean

velocity of ventrally translocating waves (V_V), where θ is the angle of the direction

of actin filaments relative to the cell membrane (Fig. 2f).

All the model parameters, i.e., lifetime, length, moving velocity and degree of the

change in moving direction, were obtained from our quantitative experimental data

(Extended Data Fig. 4, Extended Data Table 1). The actin waves described by the model

accumulated at the corners of the triangular cells (Fig. 3e and Supplementary Video 13),

as observed in U251 cells (Fig. 3c). Using this mathematical model, we further analysed

the movement of the actin wave in cells with different corner angles (i.e. 30°, 60°, 120°

or 180°; Extended Data Fig. 3b) and quantified the relative accumulation of actin

filaments at the cell periphery and corners (Extended Data Fig. 3c). The actin waves

accumulated at the cell periphery (Extended Data Fig. 3d). Furthermore, the

accumulation of actin waves increased as the corner angle decreased from 180° to 30°

(Fig. 3f). To validate this model, we cultured U251 cells using the same patterns. As

predicted, actin filaments accumulated at the periphery (Extended Data Fig. 3e) and

corners of the cells in a corner angle-dependent manner (Fig. 3g).

We further examined the effect of inhibiting the actin wave via knocking out

shootin1b. Interestingly, shootin1b KO not only reduced the translocation velocity (Fig.

1i) but also the lifetime (Extended Data Fig. 4a-c) of actin waves. Introduction of these

parameters (i.e. wave velocity and lifetime derived from shootin1b KO cells) into the

model disrupted actin accumulation at the cell periphery and corners (Fig. 3h and

Supplementary Video 13). Consistently, shootin1b KO inhibited actin accumulation at

the cell periphery and corners of U251 cells (Fig. 3i). Taken together, the findings

indicate that treadmilling actin waves mediate actin filament accumulation at the cell

periphery and protrusions.

Actin waves mediate cell morphogenesis

As shown above, the inhibition of actin wave movement by shootin1b KO disrupted the accumulation of actin filamentsat the periphery and corners of shape-constrained cells (Fig. 3h-i) without affecting the actin polymerisation (Fig. 1i). Finally, the role of the actin wave in cell morphogenesis was analysed using cells with a deformable plasma membrane. Shootin1b KO resulted in the fragmentation of lamellipodia (Fig. 4a), thereby decreasing the lamellipodial coverage rate at the cell periphery (Fig. 4b and Extended Data Fig. 5a) and reducing filopodial length (Extended Data Fig. 5b). Furthermore, shootin1b depletion disrupted the unilateral integration of lamellipodium, resulting in a decrease in the degree of cell polarity (Fig. 4c). To further analyse how actin waves drive cell polarisation, we monitored the recovery of cell morphology following latrunculin A-induced actin filament disassembly[29]. Treatment of wild-type (WT) U251 cells with 100 nM latrunculin A disrupted their polarised morphology (Fig. 4d and Supplementary Video 14). During recovery following latrunculin A washout, the cells formed multiple foci of small lamellipodia enriched with actin filaments (asterisks, Fig. 4d). The foci then grew (yellow arrows) and expanded laterally (blue arrows), eventually integrating into a large lamellipodia. In contrast, shootin1b deletion inhibited the expansion and integration of lamellipodium foci (asterisks, Fig. 4e), thereby delaying the onset of cell migration after recovery from the latrunculin A treatment (Fig. 4f).

These finding suggest that cell shape-guided movement of treadmilling actin waves boosts actin-dependent cell morphogenesis and motility. In glioma cells, actin waves arrived at the plasma membrane (green arrows, Extended Data Fig. 6) aligned randomly oriented actin filaments to outward polymerisation and pushed the membrane to protrude (white arrows). They accumulated spontaneously at cell protrusions (red arrows), resulting in their further enlargement. Through lateral expansion (yellow arrow), the protrusions were integrated into a large leading edge of the front. These processes would cooperate with other mechanisms, including myosin II-mediated contraction at the rear[17,30] to polarise cells and drive cell migration.

Discussion

Treadmilling is an inherent property of actin dynamics. We demonstrated that arrays of

treadmilling actin filaments emerge widely in glioma cells and move as actin waves in

the direction of polymerisation. Contrary to the classical pathway, in which actin

nucleators promote cell protrusion by activating actin polymerisation (black arrows, Fig.

5a), actin filaments and the Arp2/3 complex component ARPC2 accumulated in cell

protrusions in the absence of local signalling (Fig. 3a-c). Their corner angle-dependent

accumulation (Fig. 3e-g) is similar to the motion of self-propelled particles, which are

described in the fields of current physics[13,15] and soft matter[16]. Although we do not rule

out actin waves that travel via other mechanisms, this analogy supports our conclusion

that the treadmilling actin waves move as self-propelled machinery rather than via the

reaction-diffusion mechanism. Treadmilling actin waves are not stable particles, but

they continuously exchange actin subunits between the cytoplasm during translocation

(Fig. 1g)[12].

The shape-guided self-propelled movement of actin waves provides a new

framework for understanding the mechanism of cell morphogenesis (Fig. 5). First, the

shape-guided actin accumulation (green arrow, Fig. 5a) constitutes a positive feedback

interaction with actin-mediated cell protrusion (red box), thereby conferring robustness

to the classical pathway of cellular morphogenesis (black arrows) under the regulation

of signalling (blue arrow). Second, in the longstanding model of morphogenesis, i.e.

local-excitation-global-inhibition (LEGI) model[30,31], a local activator of protrusion

generates a global inhibitor (e.g. a molecule or membrane tension) that propagates

laterally to suppress lateral protrusion (blue, dotted lines, Fig. 5b). However, the

inhibitor would decay during the lateral propagation either by molecular diffusion or by

membrane stretching (blue line, Fig. 5b). Furthermore, it would also inhibit the

activated area, hampering the specific inhibition in the lateral regions (blue line). On the

other hand, the self-propelled model does not require a global inhibitor because the

translocation of the activator (actin wave) to the protrusion results in its lateral decrease,

thereby suppressing protrusive activities laterally (blue arrows, Fig. 5c). Thus, this

model offers a simple framework of effective lateral inhibition for cell morphogenesis

(blue line). Finally, the present cell shape-sensitive system can respond not only to

signalling but also to extracellular topographic cues that deform the plasma membrane

from outside (yellow arrow, Fig. 5a) (Fig. 3e-g). This would enable cells to navigate

flexibly in three-dimensional environments through mechanical barriers and tissue

pores[11,32-34].

292   Notably, recent simulation and microswimmer studies have reported that

self-propelled particles in vesicles induce various shape deformations and movements

similar to those observed in living cells[16,35-37]. Different actin-binding proteins and

physical interactions could control the mode of protrusion, either extension or

expansion, by altering the shape of the treadmilling filament arrays[3,4,38]. Because actin

filaments undergo treadmilling in various protrusions (e.g., lamellipodia and filopodia

of migrating cells[1,3,4], neurite tips and dendritic spines of neurons[39-42], microvilli and

cell-cell adhesion sites of epithelial cells[43-46], hair cell stereocilia[47], invadopodia[48,49], and

phagocytosis cups[50]), treadmilling actin waves may mediate various processes of cell

morphogenesis and motility.

References

1   Pollard, T. D. & Borisy, G. G. Cellular motility driven by assembly and disassembly of actin filaments. *Cell* **112**, 453-465 (2003).

2   Bernstein, B. W. & Bamburg, J. R. ADF/cofilin: a functional node in cell biology. *Trends Cell Biol.* **20**, 187-195 (2010).

3   Carlier, M. F. & Shekhar, S. Global treadmilling coordinates actin turnover and controls the size of actin networks. *Nat. Rev. Mol. Cell Biol.* **18**, 389-401, doi:10.1038/nrm.2016.172 (2017).

4   Rottner, K. & Schaks, M. Assembling actin filaments for protrusion. *Curr. Opin. Cell Biol.* **56**, 53-63 (2019).

5   Goode, B. L., Eskin, J. & Shekhar, S. Mechanisms of actin disassembly and turnover. *J. Cell Biol.* **222** (2023).

6   Vitriol, E. A., Wise, A. L., Berginski, M. E., Bamburg, J. R. & Zheng, J. Q. Instantaneous inactivation of cofilin reveals its function of F-actin disassembly in lamellipodia. *Mol. Biol. Cell* **24**, 2238-2247 (2013).

7   Carlsson, A. E. Actin dynamics: from nanoscale to microscale. *Annu. Rev. Biophys.* **39**, 91-110 (2010).

8   Allard, J. & Mogilner, A. Traveling waves in actin dynamics and cell motility. *Curr. Opin. Cell Biol.* **25**, 107-115 (2013).

9   Inagaki, N. & Katsuno, H. Actin waves: origin of cell polarization and migration? *Trends Cell Biol.* **27**, 515-526 (2017).

10   Beta, C., Edelstein-Keshet, L., Gov, N. & Yochelis, A. From actin waves to mechanism and back: How theory aids biological understanding. *eLife* **12** (2023).

11   Weiner, O. D., Marganski, W. A., Wu, L. F., Altschuler, S. J. & Kirschner, M. W. An actin-based wave generator organizes cell motility. *PLoS Biol.* **5**, e221 (2007).

12   Katsuno, H. *et al.* Actin migration driven by directional assembly and disassembly of membrane-anchored actin filaments. *Cell Rep.* **12**, 648-660 (2015).

13   Kaiser, A., Wensink, H. H. & Lowen, H. How to capture active particles. *Phys. Rev. Lett.* **108**, 268307 (2012).

14   Elgeti, J., Winkler, R. G. & Gompper, G. Physics of microswimmers--single particle motion and collective behavior: a review. *Rep. Prog. Phys.* **78**, 056601 (2015).

15   Bechinger, C. *et al.* Active particles in complex and crowded environments. *Rev. Mod. Phys.* **88**, 045006 (2016).

16   Vutukuri, H. R. *et al.* Active particles induce large shape deformations in giant lipid vesicles. *Nature* **586**, 52-56 (2020).

17   Vassilev, V., Platek, A., Hiver, S., Enomoto, H. & Takeichi, M. Catenins steer

cell migration via stabilization of front-rear polarity. *Dev. Cell* **43**, 463-479
(2017).

18 Minegishi, T., Fujikawa, R., Kastian, R. F., Sakumura, Y. & Inagaki, N. Analyses
of actin dynamics, clutch coupling and traction force for growth cone advance. *J.*
Vis. Exp., e63227 (2021).

19 Kubo, Y. *et al.* Shootin1-cortactin interaction mediates signal-force transduction
for axon outgrowth. *J. Cell Biol.* **210**, 663-676, doi:10.1083/jcb.201505011
(2015).

20 Flynn, K. C., Pak, C. W., Shaw, A. E., Bradke, F. & Bamburg, J. R. Growth
cone-like waves transport actin and promote axonogenesis and neurite branching.
Dev. Neurobiol. **69**, 761-779 (2009).

21 Minegishi, T. *et al.* Shootin1b mediates a mechanical clutch to produce force for
neuronal migration. *Cell Rep.* **25**, 624-639.e626 (2018).

22 Nolen, B. J. *et al.* Characterization of two classes of small molecule inhibitors of
Arp2/3 complex. *Nature* **460**, 1031-1034 (2009).

23 Nishimura, Y. *et al.* The formin inhibitor SMIFH2 inhibits members of the
myosin superfamily. *J. Cell Sci.* **134** (2021).

24 Bretschneider, T. *et al.* The three-dimensional dynamics of actin waves, a model
of cytoskeletal self-organization. *Biophys. J.* **96**, 2888-2900 (2009).

25 Zhang, K., Lyu, W., Yu, J. & Koleske, A. J. Abl2 is recruited to ventral actin
waves through cytoskeletal interactions to promote lamellipodium extension.
Mol. Biol. Cell **29**, 2863-2873 (2018).

26 Ruthel, G. & Banker, G. Actin-dependent anterograde movement of
growth-cone-like structures along growing hippocampal axons: a novel form of
axonal transport? *Cell Motil. Cytoskelet.* **40**, 160-173 (1998).

27 Parker, K. K. *et al.* Directional control of lamellipodia extension by constraining
cell shape and orienting cell tractional forces. *FASEB J.* **16**, 1195-1204 (2002).

28 Koestler, S. A., Auinger, S., Vinzenz, M., Rottner, K. & Small, J. V.
Differentially oriented populations of actin filaments generated in lamellipodia
collaborate in pushing and pausing at the cell front. *Nat. Cell Biol.* **10**, 306-313
(2008).

29 Gerisch, G. *et al.* Mobile actin clusters and traveling waves in cells recovering
from actin depolymerization. *Biophys. J.* **87**, 3493-3503 (2004).

30 Hadjitheodorou, A. *et al.* Directional reorientation of migrating neutrophils is
limited by suppression of receptor input signaling at the cell rear through myosin
II activity. *Nat. Commun.* **12**, 6619 (2021).

31 Xiong, Y., Huang, C. H., Iglesias, P. A. & Devreotes, P. N. Cells navigate with a
local-excitation, global-inhibition-biased excitable network. *Proc. Natl. Acad.*
Sci. U. S. A. **107**, 17079-17086 (2010).

32 Yamada, K. M. & Sixt, M. Mechanisms of 3D cell migration. *Nat. Rev. Mol.*
Cell Biol. **20**, 738-752 (2019).

33 Jasnin, M., Ecke, M., Baumeister, W. & Gerisch, G. Actin organization in cells
responding to a perforated surface, revealed by live imaging and cryo-electron
tomography. *Structure* **24**, 1031-1043 (2016).

34 Hadjitheodorou, A. *et al.* Leading edge competition promotes context-dependent
responses to receptor inputs to resolve directional dilemmas in neutrophil
migration. *Cell Syst.* **14**, 196-209 e196 (2023).

35 Paoluzzi, M., Di Leonardo, R., Marchetti, M. C. & Angelani, L. Shape and
displacement fluctuations in soft vesicles filled by active particles. *Sci. Rep.* **6**,
34146 (2016).

36 Li, Y. & Ten Wolde, P. R. Shape transformations of vesicles induced by swim
pressure. *Phys. Rev. Lett.* **123**, 148003 (2019).

37 Boudet, J. F. *et al.* From collections of independent, mindless robots to flexible,
mobile, and directional superstructures. *Sci. Robot.* **6** (2021).

38 Huber, L., Suzuki, R., Kruger, T., Frey, E. & Bausch, A. R. Emergence of
coexisting ordered states in active matter systems. *Science* **361**, 255-258 (2018).

39 Mallavarapu, A. & Mitchison, T. Regulated actin cytoskeleton assembly at
filopodium tips controls their extension and retraction. *J. Cell Biol.* **146**,
1097-1106 (1999).

40 Suter, D. M. & Forscher, P. Substrate-cytoskeletal coupling as a mechanism for
the regulation of growth cone motility and guidance. *J. Neurobiol.* **44**, 97-113
(2000).

41 Tatavarty, V., Kim, E. J., Rodionov, V. & Yu, J. Investigating sub-spine actin
dynamics in rat hippocampal neurons with super-resolution optical imaging.
PLoS ONE **4**, e7724 (2009).

42 Lin, W. H., Nebhan, C. A., Anderson, B. R. & Webb, D. J.
Vasodilator-stimulated phosphoprotein (VASP) induces actin assembly in
dendritic spines to promote their development and potentiate synaptic strength. *J.*
Biol. Chem. **285**, 36010-36020 (2010).

43 Loomis, P. A. *et al.* Espin cross-links cause the elongation of microvillus-type
parallel actin bundles in vivo. *J. Cell Biol.* **163**, 1045-1055 (2003).

44 Meenderink, L. M. *et al.* Actin dynamics drive microvillar motility and
clustering during brush border assembly. *Dev. Cell* **50**, 545-556 e544 (2019).

45 Kovacs, E. M. *et al.* N-WASP regulates the epithelial junctional actin
cytoskeleton through a non-canonical post-nucleation pathway. *Nat. Cell Biol.*
13, 934-943 (2011).

46 Efimova, N. & Svitkina, T. M. Branched actin networks push against each other
at adherens junctions to maintain cell-cell adhesion. *J. Cell Biol.* **217**, 1827-1845
(2018).

47 Rzadzinska, A. K., Schneider, M. E., Davies, C., Riordan, G. P. & Kachar, B. An
actin molecular treadmill and myosins maintain stereocilia functional
architecture and self-renewal. *J. Cell Biol.* **164**, 887-897 (2004).

424 48 Oser, M. *et al.* Cortactin regulates cofilin and N-WASp activities to control the
425   stages of invadopodium assembly and maturation. *J. Cell Biol.* **186**, 571-587
426   (2009).
427 49 Albiges-Rizo, C., Destaing, O., Fourcade, B., Planus, E. & Block, M. R. Actin
428   machinery and mechanosensitivity in invadopodia, podosomes and focal
429   adhesions. *J. Cell Sci.* **122**, 3037-3049 (2009).
430 50 Gerisch, G. Self-organizing actin waves that simulate phagocytic cup structures.
431   *PMC Biophys.* **3**, 7 (2010).
432 51 Okano, K. *et al.* In situ laser micropatterning of proteins for dynamically
433   arranging living cells. *Lab Chip* **13**, 4078-4086 (2013).
434 52 Baba, K. *et al.* Gradient-reading and mechano-effector machinery for
435   netrin-1-induced axon guidance. *eLife* **7**, e34593 (2018).
436 53 Toriyama, M. *et al.* Shootin1: A protein involved in the organization of an
437   asymmetric signal for neuronal polarization. *J. Cell Biol.* **175**, 147-157 (2006).
438 54 Higashiguchi, Y. *et al.* Identification of a shootin1 isoform expressed in
439   peripheral tissues. *Cell Tissue Res.* **366**, 75-87 (2016).
440 55 Cameron, L. A., Robbins, J. R., Footer, M. J. & Theriot, J. A. Biophysical
441   parameters influence actin-based movement, trajectory, and initiation in a
442   cell-free system. *Mol. Biol. Cell* **15**, 2312-2323 (2004).
443 56 Kalwarczyk, T. *et al.* Comparative analysis of viscosity of complex liquids and
444   cytoplasm of mammalian cells at the nanoscale. *Nano Lett.* **11**, 2157-2163
445   (2011).
446

Methods

Cell culture and transfection

A subline of glioma U251 cells[17] and COS7 cells were seeded on glass-bottom dishes (Matsunami, #D11130H) coated with 100 μg/ml poly-D-lysine (Sigma) and 5 μg/ml laminin (Lonza), and cultured in DMEM high glucose medium (Nacalai, #08458-16) supplemented with 10% fetal bovine serum (Japan Bio Serum) at 37°C and 5% CO_2. They were transfected with vectors using Polyethylenimine MAX (PEI MAX, Polysciences) following the manufacturer's protocol.

Generation of shootin1b KO U251 cells by CRISPR-Cas9

The guide oligonucleotide sequence (forward: 5′-CACCGGCAGCTCATTACCAGTCTGA-3′; reverse: 5′-AAACTCAGACTGGTAATGAGCTGCC-3′), corresponding to nucleotides 30-49 in the coding region of human shootin1b, were selected by an online CRISPR Cas9 Design tool (http://crispr.mit.edu/). The pSpCas9(BB)-2A-Puro (PX459) V2.0 plasmid (Addgene, #62988) was linearised using by BbsI-HF (New England Biolabs, #R3539S), then the guide oligonucleotides were annealed and inserted into the vector using Ligation high Ver.2 (TOYOBO, #LGK-201). U251 cells were transfected with the plasmid, cultured individually in 96-well plates, and screened using 4 μg/ml puromycin. Shootin1b depletion in KO cells was confirmed by immunoblot analyses (Extended Data Fig. 2g).

Micropatterning of laminin on the substrate for cell adhesion

Micropatterns of laminin for cell adhesion were produced on a glass surface following our previously established procedure[51], with modifications (Extended Data Fig. 3a). The glass-bottom dish (Iwaki, #3961-035) was cleaned using ultrasonic waves for 5 min with special grade ethanol, followed by rinsing with water and additional 5-min ultrasonic cleaning with pure water. After the cleaning, the dish was air-dried in a clean room at room temperature. 3-Phenylaminopropyltrimethoxysilane (2%, Shin-Etsu, #LS-4500) was dissolved in a 2% solution of acetic acid in water and then poured onto the glass-bottom dish, followed by incubation for 30 min. After being washed twice with water, the dish was dried at 70°C for 1 h. This process resulted in the covalent

bonding of the 3-phenylaminopropyl group with the glass surface, which made it

hydrophobic. Any residues resulting from the reaction were eliminated by ultrasonically

cleaning with special grade ethanol for 5 min.

The hydrophobic glass surface was coated with 100 µl of ethanol containing

0.2% of MPC (2-methacryloyloxyethyl phosphorylcholine) polymer (NOF,

#Lipiduro-CM5206). After the reaction for 10-15 sec, the coating solution was removed

and left to dry overnight at room temperature in the dark. This resulted in the formation

of an MPC polymer film on the glass. Then, the dish was incubated with 2 ml H_2O

overnight. In the film, the phosphatidylcholine residues of the MPC polymers are

oriented towards the aqueous solution (Extended Data Fig. 3a).

The cell non-adhesive MPC-polymer layer was removed by a programmed

raster scanning of a femtosecond laser (center wavelength 800 nm, pulse width 130 fs,

pulse energy 100 nJ, transmission frequency 5 kHz, Solstice, Spectra-Physics) focused

with an objective lens 20x 0.46 NA (Olympus, UMPlanFl) at a speed of 400 µm/s on

the glass surface. The raster scanning patterns are shown in Extended Data Fig. 3a, b.

Then, the cell-adhesive patterns of laminin were produced by coating the exposed glass

surface with poly-D-lysine (PDL) and laminin. The cell-adhesive patterns were

confirmed by coating with green fluorescent-HiLyte 488-Laminin (Cytoskeleton,

#LMN02-A) (Extended Data Fig. 3a).

Immunocytochemistry

Immunocytochemistry was performed as described previously[52]. Cultured cells were

fixed with 3.7% formaldehyde in Krebs buffer (118 mM NaCl, 4.7 mM KCl, 1.2 mM

KH_2PO_4, 1.2 mM $MgSO_4$, 4.2 mM $NaHCO_3$, 2 mM $CaCl_2$, 10 mM glucose, 400 mM

sucrose, 10 mM HEPES pH 7.0) for 20 min on ice and 10 min at room temperature,

followed by treatment for 15 min with 0.05% Triton X-100 in PBS on ice and 10% fetal

bovine serum in PBS for 1 h at room temperature. The cells were incubated with a

primary antibody diluted in PBS containing 10% fetal bovine serum overnight at 4°C.

They were washed with PBS, and then incubated with secondary antibody diluted in

PBS for 1 h at room temperature. After washing with PBS, the cells were stained with

Alexa Fluor 594 conjugated phalloidin (1:100, Thermo Fisher Scientific) and DAPI

(1:1000, Thermo Fisher Scientific) for 30 min at room temperature. The fluorescence

and phase-contrast images of U251 cells were acquired a fluorescence microscope (Carl
Zeiss, Axioplan2) equipped with an objective lens 40x 0.75 NA (Carl Zeiss,
plan-Neofluar), a charge-coupled device camera (Carl Zeiss, AxioCam MRm) and
imaging software (Carl Zeiss, ZEN), and a TIRF microscope (Olympus, IX81) equipped
with a CMOS camera (Hamamatsu, ORCA Flash4.0LT), an objective lens 100x 1.49
NA (Olympus, UAPON), and MetaMorph software.
517

Immunoblot
Cultured U251 cells were lysed with RIPA buffer (50 mM Tris-HCl (pH 8.0), 1 mM
EDTA, 150 mM NaCl, 1% Triton X-100, 0.1% SDS, 0.1% sodium deoxycholate, 1 mM
DTT, 1 mM PMSF, and 0.01 mM leupeptin), and incubated for 10 min at 4°C.
Subsequently, the cell lysate was centrifuged at 15000 rpm for 15 min at 4°C. The
supernatant was mixed with an equal volume of 2 x SDS sample buffer (131 mM
Tris-HCl, pH 6.8, 21% glycerol, 4% SDS, 0.05% bromophenol blue and 5%
β-mercaptoethanol). The mixture was incubated for 5 min at 95°C, followed by
SDS-polyacrylamide gel electrophoresis. Immunoblotting was performed as described
previously[21].
528

Antibodies
The following primary antibodies were used: anti-shootin1 (rabbit, 1:400)[53];
anti-shootin1b (rabbit, 1:2000)[54]; anti-NCAM-L1 (mouse, 1:1000, Santa Cruz
Biotechnology, #sc-514360); anti-cortactin (mouse, 1:1000, Millipore, #05-180);
anti-actin (mouse, 1:1000, Millipore, #MAB1501R); anti-ARPC2 (rabbit, 1:1000,
Millipore, #07-227). The following secondary antibodies were used: anti-rabbit (Alexa
594 conjugated donkey, 1:1000, Jackson ImmunoResearch Labs, #711-585-152);
anti-rabbit (Alexa 488 conjugated goat, 1:1000, Thermo Fisher Scientific, #A-11008);
anti-mouse (Alexa 488 conjugated goat, 1:1000, Thermo Fisher Scientific, #A-11029);
anti-rabbit (HRP conjugated donkey, 1:2000, GE Healthcare, #NA934); anti-mouse
(HRP conjugated goat, 1:5000, Thermo Fisher Scientific, #AP308P).
540

Chemicals
The following reagents were used: phalloidin (Alexa Fluor 594 conjugated, 1:100,

Thermo Fisher Scientific, #A12381); DAPI (1:1000, Thermo Fisher Scientific,

#D1306); Poly-D-lysine (100 μg/ml, Sigma, #P6407-5MG); Laminin (5 μg/ml, Lonza,

#120-05751); HaloTag TMR ligand (5mM, Promega, #G8251); LIPIDURE (0.2% NOF,

#CM5206); 3-phenylaminopropyltrimethoxysilane (2%, Shin-Etsu, #LS-4500)

DNA construction

To generate pEGFP-N1-LifeAct vector, the annealed LifeAct (forward:

5′-TCGAGATGGGTGTCGCAGATTTGATCAAGAAATTCGAAAGCATCTCAAAG

GAAGAAGGG-3′;   reverse:

5′-GATCCCTTCTTCCTTTGAGATGCTTTCGAATTTCTTGATCAAATCTGCGACA

CCCATC-3′) was fused to N-terminal of EGFP-N1 vectors (Clontech). To generate

pLifeAct-mCherry-N1 vector, the annealed LifeAct was fused to N-terminal of

mCherry-N1 vectors (Takara, #632523). Generation of pFN21A-HaloTag-actin was

described previously[21].

TIRF and epifluorescence microscopy

U251 cells were cultured on glass-bottom dishes coated with PDL and laminin. Actin

dynamics in close proximity to the ventral plasma membrane were visualised using the

actin filament marker EGFP-LifeAct or LifeAct-mCherry under TIRF or

epifluorescence microscopy. In addition, to differentially visualise actin waves

translocating in close proximity to the ventral plasma membrane and those detached

from the membrane, we combined TIRF microscopy, which visualises the region in

close proximity to the ventral plasma membrane, and epifluorescence microscopy,

which visualises the entire cell thickness (Extended Data Fig. 1e). U251 cells

expressing LifeAct-mCherry and EGFP-LifeAct were cultured on glass-bottom dishes

coated with PDL and laminin. Prior to observation, the cells were incubated in

DMEM/F-12 Ham medium (Sigma) supplemented with 10% FBS (Japan Bio Serum).

The TIRF laser was used to observe fluorescence images of LifeAct-mCherry, while the

epifluorescence laser was used to observe fluorescence images of EGFP-LifeAct. The

observations were made with a TIRF microscope (Olympus, IX81) equipped with a

CMOS camera (Hamamatsu, ORCA Flash4.0LT), an objective lens 100x 1.49 NA or

150x 1.45 NA (Olympus, UAPON), and MetaMorph software. Images were acquired

every 10 or 60 sec.

Fluorescent speckle imaging of actin wave by TERF microscopy

U251 cells expressing HaloTag-actin and EGFP-LifeAct were cultured on glass-bottom

dishes coated with PDL and laminin. Prior to observation, the cells were treated with

0.05 μM HaloTag TMR ligand (Promega) in the culture medium and incubated for 1 h

at 37°C. The ligand was washed with PBS, and the cells were then incubated in

DMEM/F-12 Ham medium (Sigma, #D8062) supplemented with 10% FBS (Japan Bio

Serum). The fluorescent speckles of HaloTag-actin and EGFP-LifeAct were observed

using a TIRF microscope (Olympus, IX81) equipped with a CMOS camera

(Hamamatsu, ORCA Flash4.0LT), an objective lens 150x 1.45 NA (Olympus, UAPON),

and MetaMorph software. Fluorescence images were acquired every 10 sec. Actin flow

velocity was calculated by tracing the speckles of HaloTag-actin in actin waves using

ImageJ (Fiji version). Actin polymerisation rate was calculated as the sum of actin wave

migration velocity, visualised by EGFP-LifeAct, and HaloTag-actin flow velocity, as

reported[18].

3D imaging by confocal deconvolution microscopy

U251 cells expressing HaloTag-actin and EGFP-LifeAct were cultured in 2D (on

glass-bottom dishes coated with PDL and laminin) or 3D (75% matrigel) environment.

Before observation, the cells were incubated with 0.05 μM HaloTag TMR ligand

(Promega) in the culture medium and incubated for 1 h at 37°C. The ligand was washed

with PBS, and the cells were then incubated in DMEM/F-12 Ham medium (Sigma)

supplemented with 10% FBS (Japan Bio Serum). The fluorescent speckles of

HaloTag-actin and EGFP-LifeAct were observed using a confocal microscope (Leica,

Stellaris 8) equipped with an objective lens 100x 1.40 NA (Leica, HC PL APO CS2) and

Las X software. Deconvolution of 3D image stacks were obtained by Las X software.

Statistical analyses

Statistical analyses were performed using Excel (Microsoft 365) or GraphPad Prism 7

(GraphPad Software). For samples with more than 7 data points, the D′Agostino–

Pearson normality test was used to determine whether the data followed a normal

distribution. In cases where the number of data points was between 3 and 7, the

Shapiro–Wilk test was used for the normality test. We also tested the equality of

variation with the F test for two independent groups that followed normal distributions.

Significance tests were performed as follows: (1) two-tailed unpaired Student′s t-test to

compare normally distributed data with equal variance from two independent groups;

(2) two-tailed unpaired Welch′s t-test to compare normally distributed data with unequal

variance from two independent groups; (3) two-tailed Mann–Whitney U-test to compare

nonnormally distributed data from two independent groups. The statistical information

and number of samples for each experiment are indicated in the Figure legends. For

detailed statistical results including the test statistics and exact p values, see the

statistical source data associated with each Figure. All data are shown as the mean ±

SEM. Statistical significance was defined as ***p < 0.01; **p < 0.02; *p < 0.05; ns, not

significant. All experiments were performed at least three times and reliably reproduced.

The investigators were blinded to the experimental groups for each analyses.

Mathematical modelling

To investigate whether actin waves mediate the accumulation of actin filaments at the

protrusions of cells confined to adhesive islands (Fig. 3c), we constructed a

mathematical model that describes the movement of intracellular actin waves. Our

model is based on quantitative data obtained from experimental observations: we

focused on filopodium-type waves because it is difficult to observe individual actin

filaments in lamellipodium-type waves (Fig. 1c), and filopodium-type and

lamellipodium-type waves are interchangeable (Extended Data Fig. 1c). For simplicity,

the present model describes the translocation of the actin filament arrays, not a detailed

mechanism of wave translocation[12]. The model depicts the processes of wave

emergence and disappearance, as well as changes in the direction of wave translocation.

The model parameters used in the model are summarised in Extended Data Table 1.

Emergence and disappearance of actin waves

Our experimental data indicate that actin waves emerge widely in the cytoplasm, move

in random directions, and then disappear (Supplementary Video 1). In our model, the

emergence locations and translocation directions of actin waves are randomly chosen at
the start of the simulation. The simulation is updated every 1 sec ($\Delta t = 1$ sec),
generating one actin wave in the cell with a probability of 10%. The lifetimes of actin
waves in WT and shootin1b KO cells, quantified by live cell imaging, were modeled
using gamma distributions (Extended Data Fig. 4a, b).

The length (L) and translocation velocity (V) of actin waves are also derived
from experimental data (Extended Data Fig. 4e and Fig. 1i); the model uses their
average values for simplicity (Extended Data Table 1). Actin waves located at the cell
periphery move laterally along the cell membrane (Fig. 2d). The velocity of the lateral
movement was modelled as $\cos\theta$ of the mean velocity of actin waves (Fig. 1i), where θ
is the angle of the direction of wave movement with respect to the cell membrane (Fig.
2f).

The local concentration of actin subunits increases immediately after the
disappearance (complete disassembly) of actin wave. Since actin polymerisation
depends on the local concentration of actin subunits, our model assumes that an actin
wave reemerges at the location where the preceding wave disappeared and translocates
in a random direction.

Change in the direction of wave translocation
Actin waves frequently change the direction of their translocation (Supplementary
Videos 1 and 4). The change in direction was quantified every 50 sec from live cell
imaging data (Extended Data Fig. 4d). As the angular distribution of these changes was
symmetric with a zero peak, we modelled it using a Gaussian distribution (Extended
Data Fig. 4d).

Analyses of actin filament accumulation and lamellipodial formation
Accumulation of actin filaments at the cell periphery was determined by dividing the
average fluorescence intensity of actin filaments within a 2 μm-wide region from the
cell periphery (green region, Extended Data Fig. 3c) by the average intensity of actin
filaments in the whole cell region (blue region). Accumulation of actin filaments at the
cell protrusions was determined by dividing the average intensity of actin filaments at
the corner regions (yellow region: 2 μm-wide and 5 μm-long regions in both sides of the

corner, Extended Data Fig. 3c) by the average intensity of actin filaments in the
peripheral region (green region). Lamellipodial coverage rate was calculated by
dividing the total length of lamellipodia (yellow line, Extended Data Fig. 5a) by the
perimeter of the cell (green line). The fluorescence intensity of actin filaments was
analysed using ImageJ (Fiji version). Lamellipodia was defined as the area of cell
periphery where the fluorescence intensity exceeded the mean intensity + SD of the
whole cell body. The polar histogram of the lamellipodial areas was generated using
MATLAB (Fig. 4d).

Data availability
680 Data will be made available upon reasonable request.

Code availability
683 The MATLAB codes used for mathematical simulations are available at GitHub
684 (https://github.com/KioYagami/actin-wave/tree/master). Any additional information will
685 be available from the corresponding author upon reasonable request.

Acknowledgments
We thank Drs. Masatoshi Takeichi (Riken) and Shintaro Suzuki (Kwansei Gakuin
Univ) for providing the subline of U251 cells, Drs. Hitoshi Sakano (Fukui Univ), Jim
Bamburg (CSU), Nir Gov (Weizmann Inst Sci), Carsten Beta (Potsdam Univ), Sean
Collins (UC Davis), Klemens Rottner (Tech Univ Braunschweig), Giovanni Volpe
(Univ Gothenburg), Shiro Suetsugu (NAIST), Naoki Watanabe (Kyoto Univ) and
Yusuke Maeda (Kyushu Univ) for valuable discussions, and Satoko Shimamura for kind
encouragements. This research was supported in part by AMED under Grant Number
JP17gm0810011 (N.I.), JSPS KAKENHI (JP19H03223, N.I.), JSPS Grants-in-Aid for
Early-Career Scientists (JP19K16127, K.B. and JP23K14181, T.M.), and the Osaka
Medical Research Foundation for Incurable Diseases (K.B., T.M.).

Author contributions
K.Y., K.B., S.M., H.K.-K. and K.O. performed the experiments and data analyses. K.Y.,
K.B., S.M., H.K.-K., T.M., K.O., Y.H. and N.I. designed the experiments. K.Y., Y.S. and
N.I. constructed the mathematical model. K.Y., Y.S., K.O. and N.I. wrote the

manuscript. N.I. supervised all the projects. All authors discussed the results and

commented on the manuscript.

Competing Interest Statement

The authors declare that they have no competing interests.

 Figure legends

Fig. 1. Actin waves emerge widely in cells and translocate by treadmilling. a, A

fluorescence image of a U251 cell expressing EGFP-LifeAct obtained by TIRF

microscopy. Asterisk indicates cellular leading edge. See Supplementary Video 1. **b-c**,

Fluorescence time-lapse images of filopodium-type (arrowheads, **b**) and

lamellipodium-type (arrows, **c**) actin waves in U251 cells expressing LifeAct-mCherry

obtained by TIRF microscopy. **d-e**, Fluorescence time-lapse images of actin waves on

the dorsal membrane (blue arrowheads and arrows, **d**) and filopodium-type actin waves

in cytoplasm (green arrowheads, **e**) in U251 cells expressing LifeAct-mCherry obtained

by 3D imaging with confocal deconvolution microscopy. See Supplementary Video 5. **f**,

A fluorescent speckle image of HaloTag-actin in a U251 cell obtained by TIRF

microscopy; actin filaments were also monitored by EGFP-LifeAct. See Supplementary

Video 6. Time-lapse montages of a filopodium-type actin wave in the rectangular region

at 10-sec intervals are shown to the right. Green and magenta lines indicate wave

forward translocation and actin filament retrograde flow, respectively. Actin

polymerisation rate can be calculated as the sum of the forward translocation rate and

actin retrograde flow rate (double-headed arrow). **g**, A diagram showing the

translocation mechanisms of actin waves travelling on the ventral plasma membrane

(left)[12]. Right panels show a possible mode of actin wave translocation observed in **e**

and Extended Data Fig. 1f. For explanations, see text and Supplementary Discussion. **h**,

Translocation velocities of filopodium-type actin waves (**f**, green line) before and after

the application of DMSO (control, N = 4 cells: before application, n = 34 waves; after

application, n = 32 waves), CK666 (N = 3 cells: before application, n = 49 waves; after

application, n = 33 waves) or SMIFH2 (N = 5 cells: before application, n = 52 waves;

after application, n = 37 waves). **i**, Velocities of actin filament retrograde flow (**f**,

magenta line), actin filament polymerisation (**f**, double-headed arrow), and

filopodium-type wave translocation (**f**, green line) in WT and shootin1b KO U251 cells

(WT, N = 11 cells, n = 71 waves; KO#1, N = 9 cells, n = 69 waves; KO#2, N = 10 cells,

n = 45 waves). Scale bars, 20 μm (**a**); 5 μm (**f** left): 1 μm (**b**, **c**, **d**, **e**, **f** right). Data

represent means ± SEM; ***p < 0.01; *p < 0.05; ns, not significant. Statistical analyses

were performed using the two-tailed Mann–Whitney *U*-test (**h**, **i**).

Fig. 2. Actin wave arrival at the periphery promotes membrane protrusion. a-b, Fluorescence time-lapse images of filopodium-type actin waves (arrowheads) which entered into pre-existing lamellipodium (arrows, **a**) and filopodium (asterisks, **b**) in U251 cells expressing EGFP-LifeAct obtained by TIRF microscopy. Entry of the waves into lamellipodia resulted in local actin filament accumulation and lamellipodial extension (arrows, **a**). See Supplementary Video 7. **c-d**, Fluorescence time-lapse images of lamellipodium-type actin wave (yellow arrows, **c**) and filopodium-type actin waves (arrowheads, **d**) which arrived at the cell periphery in U251 cells expressing LifeAct-mCherry obtained by TIRF microscopy (**c**) and expressing EGFP-LifeAct obtained by epifluorescence microscopy (**d**). They pushed the plasma membrane, resulting in the formation of lamellipodium (**c**) and filopodia (**d**), and continued to move laterally along the membrane. The former merged with the pre-existing lamellipodium (blue arrows, **d**). See Supplementary Videos 8 and 9. **e**, A diagram describing the relationship between the velocities of lateral movement (V_L) and ventral movement (V_V) of filopodium-type actin waves. If actin waves move laterally along the membrane driven by directional actin treadmilling, V_L is equal to $V_V \cos\theta$, where θ is the angle of the polymerising actin filaments with respect to the membrane. **f**, Fluorescence time-lapse images of a filopodium-type actin wave travelling along the lateral membrane (arrowheads) of a U251 cell expressing LifeAct-mCherry obtained by epifluorescence microscopy. See Supplementary Video 10. The right graph shows V_L and θ obtained by the time-lapse imaging; V_L closely matched $V_V\cos\theta$ (N = 10 cells, n = 21 waves). Scale bars, 5 μm (**c, d, f**); 1 μm (**a, b**).

Fig. 3. Treadmilling actin waves spontaneously accumulate at cell protrusions. a-b, Fluorescence time-lapse images of filopodium-type (arrowheads, **a**) and lamellipodium-type (arrows, **b**) actin waves which accumulated in protrusive regions (asterisks) in U251 cells expressing EGFP-LifeAct obtained by epifluorescence microscopy. See Supplementary Videos 11. **c**, A U251 cell cultured on a laminin-coated triangular adhesive island and stained with Alexa Fluor 594 conjugated phalloidin (for actin filament), anti-ARPC2 antibody, and DAPI (for nucleus). **d**, Fluorescence time-lapse images of a U251 cell expressing LifeAct-mCherry obtained by epifluorescence microscopy and cultured on a triangular adhesive island. Arrowheads

indicate an actin wave travelled along the lateral membrane, resulting in a local

accumulation of actin filaments at the corner upon arrival. See Supplementary Video 12.

e-f, Mathematical model data showing actin wave localisation in a triangular cell (**e**)

and cells with different corner angles, 30º, 60º, 120º or 180º (**f**), presented in arbitrary

unit (a.u.). See Supplementary Video 13. The right graph in **f** shows quantitative data for

the actin wave accumulation at the corners with different corner angles (30-180º, n = 20

simulations). **g**, U251 cells cultured on laminin-coated adhesive islands with the

different corner angles and stained with Alexa Fluor 594 conjugated phalloidin and

DAPI. The right graph shows quantitative data for the actin filament accumulation at the

corners (30º, N = 6 experiments, n = 22 cells; 60º, N = 8 experiments, n = 23 cells; 120º,

N = 7 experiments, n = 26 cells; 180º, N = 8 experiments, n = 18 cells). **h**, Mathematical

model data showing actin wave localisation in a triangular shootin1b KO cell. See

Supplementary Video 13. The right graphs show quantitative data for actin wave

accumulation at the edge and corners of WT and shoootin1b KO cells (WT, KO, n = 20

simulations). **i**, Shootin1b KO#1 U251 cells cultured on laminin-coated triangular

adhesive islands and stained with Alexa Fluor 594 conjugated phalloidin and DAPI. The

right graphs show quantitative data for actin filament accumulation at the edge and

corners of WT and shoootin1b KO cells (WT, N = 3 experiments, n = 13 cells; KO#1, N

= 6 experiments, n = 11 cells). Scale bars, 20 µm (**b**, **c**, **d**, **g**, **i**); 5 µm (**a**). Data represent

means ± SEM; ***p < 0.01; **p < 0.02; *p < 0.05; ns, not significant. Statistical

analyses were performed using the two-tailed unpaired Student′s *t*-test (**f**; 180º vs 120º,

120º vs 60º) (**g**; 180º vs 120º) (**h**, **i** right), two-tailed unpaired Welch′s *t*-test (**f**; 60º vs

30º) (**i** left), two-tailed Mann–Whitney *U*-test (**g**; 120º vs 60º, 60º vs 30º**).

Fig. 4. Actin waves drive cell polarisation and migration. a, WT, shootin1b KO#1

and KO#2 U251 cells stained with Alexa Fluor 594 conjugated phalloidin and DAPI.

b-c, Lamellipodial coverage rate (**b**) and histogram of lamellipodia localisation at cell

periphery (**c**) of WT, shootin1b KO#1 and KO#2 U251 cells in **a** (WT, KO#1, KO#2, N

= 3 experiments, n = 35 cells). **d-e**, Fluorescence time-lapse images of WT (**d**) and

shootin1b KO#1 (**e**) U251 cells expressing LifeAct-mCherry obtained by

epifluorescence microscopy: lower panels in (**d**) indicate enlarged time-lapse images of

the areas indicated by the dashed yellow boxes 1 and 2. The images were taken before and after the treatment with 100 nM latrunculin A (Lat A), and at the indicated times after the Lat A washout. The asterisks indicate foci of actin filament-enriched protrusions. Yellow and blue arrows indicate enlargement and lateral expansions of actin filament-enriched protrusions, respectively. See Supplementary Video 14. **f**, Time-lapse phase-contrast images of WT and shootin1b KO#1 U251 cells taken after the treatment with 100 nM Lat A and at the indicated times after the Lat A washout. Yellow boxes trace cell movement. The right graph indicates the time required to start cell migration after the Lat A washout (WT, N = 3 experiments, n = 28 cells; KO#1, N = 3 experiments, n = 40 cells; KO#2, N = 3 experiments, n = 35 cells). Scale bars, 20 μm (**a**, **d**, **e**, **f**); 5 μm (**c**, **d** enlarged views). Data represent means ± SEM; ***p < 0.01; **p < 0.02; *p < 0.05; ns, not significant. Statistical analyses were performed using the two-tailed unpaired Student′s *t*-test (**b**; KO#1 vs KO#2), two-tailed unpaired Welch′s t-test (**b**; WT vs KO#1, WT vs KO#2), two-tailed Mann–Whitney *U*-test (**f**).

Fig. 5. A model of cell morphogenesis driven by self-propelled actin waves. a, The shape-guided actin filament accumulation through actin waves (green arrow) constitutes a positive feedback interaction (red box) with actin-mediated cell protrusion, thereby conferring robustness to the classical pathway of cell morphogenesis (black arrows), under the regulation of signalling (blue arrow). This system can respond to extracellular topographic cues that deform the plasma membrane from outside (yellow arrow). **b**, In the LEGI model, a local activator of protrusion generates a global inhibitor (e.g. a molecule or membrane tension) to suppress protrusion at the lateral regions (blue dotted lines). The inhibitor would decay during the lateral propagation either by molecular diffusion or by membrane stretching; it would also inhibit the activated area, hampering the specific inhibition in the lateral regions (blue line). **c**, The self-propelled model does not require a global inhibitor because the translocation of the activator (actin wave) to the protrusion results in its lateral decrease, thereby suppressing protrusive activities laterally (blue arrows). Thus, this model offers a simple framework of effective lateral inhibition for cell morphogenesis (blue line).

 Supplementary Information

Supplementary Discussion

Translocation of treadmilling actin waves without anchoring to the plasma

membrane

As we previously reported in axonal shafts[12], the present study demonstrates that actin

waves translocate through their treadmilling and anchoring to the ventral plasma

membrane and adhesive substrate (Fig. 1g, left). However, we also observed actin

waves travelling apart from the adhesive substrate (arrowheads, Extended Data Fig. 1f,

Supplementary Video 4) as well as those moving along the dorsal plasma membrane

(Fig. 1d) and within the cytoplasm without interacting with the plasma membrane (Fig.

1e, Supplementary Video 5). In addition, inhibition of actin filament anchorage to the

plasma membrane by shootin1b KO did not completely inhibit wave translocation.

Given the highly viscous cytoplasmic environments containing cytoskeletal components

[55,56], we propose that actin waves can translocate by treadmilling without anchoring to

the membrane when retrograde actin filament flow is mechanically obstructed in

viscous intracellular environments (Fig. 1g, right). This mechanism could also explain

the actin waves moving on the ventral plasma membrane, with polymerising actin

filaments oriented to the membrane[33].

Extended Data Fig. 1. a, A U251 cell stained with Alexa Fluor 594 conjugated

phalloidin. An enlarged view of the rectangular region is shown to the right. Arrows and

arrowheads indicate lamellipodia and filopodia, respectively. **b**, Time-lapse

phase-contrast images of U251 cells. **c**, Fluorescence time-lapse images of

filopodium-type (arrowheads) and lamellipodium-type (arrows) actin waves in U251

cells expressing LifeAct-mCherry obtained by TIRF microscopy. See Supplementary

Video 2. **d**, A fluorescence image of a COS7 cell expressing LifeAct-mCherry obtained

by TIRF microscopy: right panels indicate enlarged time-lapse images in rectangular

regions 1 and 2. Arrows and arrowheads indicate lamellipodium-type and

filopodium-type waves, respectively. See Supplementary Video 3. **e-f**, Fluorescence

time-lapse images of a U251 cell expressing LifeAct-mCherry (obtained by TIRF

microscopy) and EGFP-LifeAct (obtained by epifluorescence microscopy): lower

panels and **f** indicate enlarged time-lapse images in rectangular regions 1, 2 and 3. See
Supplementary Video 4. Scale bars, 20 μm (**a**, **b**, **d**); 5 μm (**a**, **d** enlarged views, **c**, **e**); 1
μm (**e** enlarged views, **f**).

Extended Data Fig. 2. a, A Fluorescent speckle image of HaloTag-actin in a U251 cell
obtained by TIRF microscopy; actin filaments were also monitored by EGFP-LifeAct. **b**,
A Fluorescent speckle image of HaloTag-actin in a U251 cell obtained by 3D imaging
with confocal deconvolution microscopy; actin filaments were also monitored by
EGFP-LifeAct. **c**, Immunoblot analyses of U251 cells with anti-shootin1b,
anti-cortactin and anti-L1 antibodies. **d**, A fluorescence image of a U251 cell stained
with anti-shootin1b antibody, Alexa Fluor 594 conjugated phalloidin (for actin filament),
and DAPI (for nucleus). **e**, A fluorescence image of a U251 cell stained with
anti-shootin1b antibody and Alexa Fluor 594 conjugated phalloidin (for actin filament)
obtained by TIRF microscopy. An enlarged view of the area within the rectangle is
shown in the right panel. Arrowheads indicate shootin1b colocalisation with actin
filament. **f**, Translocation velocities of filopodium-type actin waves before and after the
application of 0.05 μM cytochalasin B (N = 4 cells, before application: n = 34 waves,
after application: n = 28 waves). **g**, Immunoblot analysis of shootin1b in WT U251 cells,
postnatal day 1 rate brain, and shootin1b KO U251 cells. The cell and tissue lysates
were immunoblotted with anti-shootin1 and anti-actin antibodies. The arrowhead and
asterisk indicate the bands corresponding to shootin1b and shootin1a, respectively.
Shootin1a is a splicing variant of shootin1b expressed in neurons. Immunoblots with
anti-actin antibody served as loading controls. **h**, A diagram showing the effects of
shootin1b KO on the velocities of actin filament retrograde flow (magenta arrows) and
actin wave translocation (green arrows). Scale bars, 20 μm (**b** left, **d**, **e** left); 5 μm (**b**
middle, **e** right); 1 μm (**a**, **b** right). Data represent means ± SEM; ***$p < 0.01$. Statistical
analyses were performed using the two-tailed Mann–Whitney U-test (**f**).

Extended Data Fig. 3. a, We created a triangular pattern of adhesive laminin substrate
with the edge length 35 μm and the area 530.4 μm^2. The cell-adhesive patterns
generated were confirmed by coating with green fluorescent-HiLyte 488-Laminin (see
Methods). **b**, The pattern of adhesive laminin substrate with different corner angles, 30º,

60º, 120º or 180º, and the same width 23.8 μm and the same area 530.4 μm^2 used in the

experiments in Fig. 3g, **c**, The accumulation of actin filaments at the cell periphery was

determined by dividing the average actin intensity in the periphery (2 μm width) by the

average actin intensity in the whole cell region. Similarly, the accumulation at corners

was determined by dividing the average of actin intensity in the corner regions (2 μm

width, 5 μm length) by the average actin intensity in the periphery. **d-e**, The

accumulation rates of actin filaments at cell periphery in simulations (**d**, n = 20) and

experiments (**e**, N = 3 experiments, n = 13 cells). Scale bars, 20 μm (**c**). Data represent

means ± SEM.

Extended Data Fig. 4. Quantitative experimental data used in the model

parameters. a-b, Distribution of the lifetime of filopodium-type waves in WT (**a**) and

shootin1b KO#1 U251 cells (**b**). White bars indicate the experimental data in Fig. 1i,

while red curves show gamma distributions with order α = 3.0 and β = 119 (**a**; WT),

and with α = 2.4 and β = 106 (**b**; shootin1b KO#1). The parameters were chosen to

give the best fit to the experimental data by maximum likelihood estimation. **c**, Lifetime

of filopodium-type waves in WT and shootin1b KO#1 U251 cells (WT, KO#1, N = 9

cells, n = 100 waves). **d**, Distribution of the degree of the change in travelling direction

of filopodium-type waves in 50 sec. White bars indicate the experimental data, while

red curves show Gaussian distributions with order mean; μ = -0.61 and SD; σ = 11.

The parameters were chosen to give the best fit to the experimental data by maximum

likelihood estimation (N = 8 cells, n = 86 waves). **e**, Length of filopodium-type waves

(N = 4 cells, n = 40 waves). Data represent means ± SEM; ***p < 0.01. Statistical

analysis was performed using the two-tailed Mann–Whitney U-test (**c**).

Extended Data Fig. 5. a, The lamellipodial coverage rate was determined by dividing

the total length of lamellipodia (yellow line) by the perimeter of the cell (green line). **b**,

Filopodium length of WT and shootin1b KO#1 U251 cells (WT, N = 6 cells, n = 51

filopodia; KO#1, N = 7 cells, n = 58 filopodia).

Extended Data Fig. 6. A model for cell polarisation and migration of U251 cells

mediated by actin waves. The arrival of treadmilling actin waves at the cell periphery

(green arrows) aligns randomly oriented actin filaments towards outward

polymerisation, which in turn pushes the membrane to protrude (white arrows).

Shootin1b facilitates this process by mechanically coupling treadmilling actin filaments

and the adhesive substrate[21]. Actin waves accumulate spontaneously at cell protrusions

(red arrows), leading to their further enlargement. Through lateral expansion (yellow

arrow), multiple protrusions are integrated into a large leading edge of the front. These

processes work together with other mechanisms, such as myosin II-mediated

contraction at the rear[17,30], to polarise cells and drive cell migration.

Extended Data Table 1. Model parameters estimated from experimental data.

Parameter	Value	Description	Figure
α	3.0 (WT) 2.4 (KO#1)	Parameters of the gamma distribution representing lifetime variation; estimated from quantitative data.	Extended Data Fig. 4a, b
β (sec)	119 (WT) 106 (KO#1)		
L (μm)	4.1	Actin wave length.	Extended Data Fig. 4e
V (μm/min)	2.0 (WT) 1.5 (KO#1)	Translocation velocity of actin wave.	Fig. 1i
μ, σ (degree)	$-0.61, 11$	Parameters of the Gaussian distribution representing actin wave directional change; quantified from experimental data.	Extended Data Fig. 4d

Legends for Supplementary Videos

Supplementary Video 1

A U251 cell expressing EGFP-LifeAct observed by TIRF microscopy. Filopodium-type and lamellipodium-type actin waves emerged widely in the cytoplasm, moved in random directions, and disappeared (see Fig. 1a). Time interval: 10 sec.

Supplementary Video 2

A Fluorescence time-lapse movie showing the interchangeability of filopodium-type (arrowheads) and lamellipodium-type (arrows) actin waves. A U251 cell expressing LifeAct-mCherry was observed by TIRF microscopy (see Extended Data Fig. 1c). Time interval: 60 sec.

Supplementary Video 3

Actin waves moving on the ventral plasma membrane of a COS7 cell expressing LifeAct-mCherry observed by TIRF microscopy (see Extended Data Fig. 1d). Time

interval: 60 sec.

Supplementary Video 4

A U251 cell expressing LifeAct-mCherry observed by TIRF microscopy and

EGFP-LifeAct observed by epifluorescence microscopy. Actin waves observed by

epifluorescence microscopy and TIRF microscopy are indicated by green and magenta

arrowheads, respectively (see Extended Data Fig. 1e, f). Time interval: 60 sec.

Supplementary Video 5

An actin wave translocating through the cytoplasm without interacting with the plasma

membrane (see Fig. 1e). A U251 cell expressing LifeAct-mCherry was observed by 3D

imaging with confocal deconvolution microscopy. Time interval: 13 sec. Z-stack

interval: 0.2 μm.

Supplementary Video 6

Fluorescent speckle dynamics of HaloTag-actin in a U251 cell observed by TIRF

microscopy; actin filaments were also monitored by EGFP-LifeAct. The actin wave

underwent treadmilling and translocated in the direction of polymerisation (see Fig. 1f).

A Time interval: 10 sec.

Supplementary Video 7

Filopodium-type actin waves entering into pre-existing lamellipodium and filopodium.

The wave entry resulted in local actin filament accumulation and lamellipodial

expansion (see Fig. 2a, b). U251 cells expressing EGFP-LifeAct were observed by

TIRF microscopy. Time interval: 10 sec.

Supplementary Video 8

Lamellipodium formation by a lamellipodium-type actin wave (see Fig. 2c). A U251

cell expressing LifeAct-mCherry was observed by TIRF microscopy. Time interval: 60

sec.

Supplementary Video 9

Filopodia formation by filopodium-type actin waves (see Fig. 2d). A U251 cell

expressing EGFP-LifeAct was observed by epifluorescence microscopy. Time interval:

10 sec.

Supplementary Video 10

Filopodium-type actin waves travelling along the lateral membranes (see Fig. 2f). U251

cells expressing LifeAct-mCherry were observed by epifluorescence microscopy. Time

interval: 10 sec.

Supplementary Video 11

Filopodium-type and lamellipodium-type actin waves accumulating at protrusive

regions (see Fig. 3a, b). U251 cells expressing EGFP-LifeAct were observed by

epifluorescence microscopy. Time interval: 10 sec.

1001

1002   **Supplementary Video 12**

1003   Actin wave accumulation at the corners of a triangular U251 cell (see Fig. 3d). A U251

1004   cell expressing LifeAct-mCherry was cultured on a triangular adhesive island and

1005   observed by epifluorescence microscopy. Time interval: 300 sec.

1006

1007   **Supplementary Video 13**

1008   Simulation of actin wave accumulation in WT and shootin1b KO triangular cells (see

1009   Fig. 3e and h). All the model parameters are provided by our quantitative experimental

1010   data (see Extended Data Fig. 4, Extended Data Table 1). Time interval: 10 sec.

1011

1012   **Supplementary Video 14**

1013   Fluorescence time-lapse movies of WT and shootin1b KO#1 U251 cells expressing

1014   LifeAct-mCherry and observed by epifluorescence microscopy. The images were taken

1015   before and after the treatment with 100 nM latrunculin A, and after the Lat A washout.

1016   Time interval: 60 sec.

a LifeAct

b 0″ 40″ 80″ 130″ 160″ 190″ 220″ 260″ 300″ 320″ 360″ 390″ 410″

c 0′ 2′ 6′ 14′

d 0″ 46″ 92″ 138″ 184″ 253″

Ventral

Dorsal

Cytoplasm

Adhesive substrate

e Top view

0″ 13″ 26″ 39″ 52″

Side view

f HaloTag-actin
EGFP-LifeAct

EGFP-LifeAct

HaloTag-actin

Merge

Actin wave forward translocation

Actin polymerisation

Actin filament retrograde flow

g Membrane anchoring (+)

F-actin retrograde flow

Actin wave translocation

HaloTag-actin

Membrane anchoring (−)

Adhesive substrate (ECM and neighboring cells)

Actin filament
Cortactin   Shootin1b   L1
Actin binding proteins
Directional diffusion
Viscous cytoplasm
Plasma membrane

h Before application   After application

Translocation velocity (μm/min)

DMSO   CK666   SMIFH2

n.s.   *   ***

i WT   KO#1   KO#2

Velocity (μm/min)

Retrograde flow   Polymerisation   Translocation

***   ***   n.s.

n.s.   n.s.   n.s.

***   *   n.s.

a

Signalling   **Topographic cue**

Actin polymerisation

F-actin accumulation

Actin wave

Cell protrusion

Cell morphogenesis and motility

Positive feedback loop

b   **LEGI model**

Activator

Inhibitor   Inhibitor

Inhibitor distribution

c   **Self-propelled model**

Actin wave accumulation
(F-actin & actin regulators)

Effective lateral inhibition

a

Coat glass with cell non-adhesive molecule MPC-polymer

Ablate non-adhesive MPC-polymer by laser

Femtosecond laser

Coat the ablated area by laminin

Seed cells on laminin

MPC-polymer

MPC   MPC

Hydrophobic side chain

60°

Area: 530.4 μm²

35 μm

Laminin

b

30°

60°

120°

180°

23.8 μm

c

F-actin
DAPI

Average intensity in whole cell region

Width: 2 μm

Average intensity in peripheral region

Length: 5 μm

Width: 2 μm

Average intensity in corner regions

F-actin intensity rate (periphery/cell) = $\dfrac{\text{Average intensity in peripheral region}}{\text{Average intensity in whole cell region}}$

F-actin intensity rate (corner/periphery) = $\dfrac{\text{Average intensity in corner regions}}{\text{Average intensity in peripheral region}}$

d   Model

F-actin intensity Rate (periphery/cell)

2.0
1.5
1.0
0.5
0.0

WT

e   Experiment

F-actin intensity Rate (periphery/cell)

2.0
1.5
1.0
0.5
0.0

WT

a

Lamellipodia

Perimeter

$$\text{Lamellipodial coverage rate} = \frac{\text{Length of lamellipodia}}{\text{Perimeter of cell}}$$

b