Title: Dysregulation of macrophage lipid metabolism underlies intracellular bacterial

neuroinvasion

Authors: Zhou Sha[1, #], Kun Yang[2, #], Shoupeng Fu[3, #], Xiaoyong Tong[4, #], Hui Yang[1],

Huiling Fang[1], Qianqian He[1], Ning Li[5, 6], Xinyu Shu[5, 6], Qi Liu[1], Yongliang Du[1],

Beibei Fu[1], Yan Xiong[1], Dong Guo[1], Jin Liu[4], Qian Li[7], Hao Zeng[8], Xiaokai Zhang[8],

Rui Yao[9], Xushuo Zhang[10], Wenjin Guo[3], Xuhu Mao[7], Mian Long[5, 6], Xiaoyuan Lin[7,

11, *], Quanming Zou[8, *], Haibo Wu[1, *]

1 School of Life Sciences, Chongqing University, Chongqing 401331, China

2 Chongqing Public Health Medical Center, Chongqing 400036, China

3 School of Veterinary Medicine, Jilin University, Changchun 130062, China

4 School of Pharmaceutical Sciences, Chongqing University, Chongqing 401331,

China

5 Center of Biomechanics and Bioengineering, Key Laboratory of Microgravity

(National Microgravity Laboratory), and Beijing Key Laboratory of Engineered

Construction and Mechanobiology, Institute of Mechanics, Chinese Academy of

Sciences, Beijing 100190, China

6 School of Engineering Sciences, University of Chinese Academy of Sciences,

Beijing 100049, China

7 Department of Clinical Microbiology and Immunology, College of Pharmacy and

Laboratory Medicine, Army Medical University (Third Military Medical University),

Chongqing 400038, China.

8 National Engineering Research Center of Immunological, Department of

Microbiology and Biochemical Pharmacy, College of Pharmacy and Laboratory

Medicine, Army Medical University (Third Military Medical University), Chongqing

400038, China.

9 Department of Pathology, Chongqing Hygeia Hospital, Chongqing 401331, China

10 Zhejiang Toyouvax Biopharming, Hangzhou 311100, China

11 Institut für Virologie, Freie Universität Berlin, Robert-von-Ostertag-Str. 7-13,

Berlin 14163, Germany.

# Equally Contributing Authors

* Corresponding authors

Xiaoyuan Lin, linxiaoyuan23@163.com

Quanming Zou, qmzou2007@163.com

Haibo Wu, hbwu023@cqu.edu.cn

Graphic abstract

Summary

Acute infection of the central nervous system is one of the deadliest diseases, but the mechanisms by which intracellular bacteria infiltrate the brain remain poorly understood. Phagocytic cells are usually recognized as the battlefield on which war is waged against intracellular bacteria; however, little is known about how the intracellular bacteria take advantage of infected phagocytes to access the brain. In this study, we find that a novel $CD36^+ Fabp4^+ Pparg^+$ macrophage subpopulation ($CD36^+$ macrophage) participates in penetration of the brain by intracellular bacteria without disruption of the brain-blood barrier. Biomechanical analysis reveals that abundant protrusions and adhesion molecules on $CD36^+$ macrophages confer significant resistance to the mechanical stress of blood flow, thereby providing more opportunities for these macrophages to adhere to the vascular endothelial surface. Through metabolomics analysis, we find that macrophage lipid metabolism is dysregulated during bacterial neuroinvasion, and that β-hydroxybutyrate promotes the formation and survival of $CD36^+$ macrophages. Importantly, ketogenesis exacerbates symptoms during bacterial neuroinvasion, which could be alleviated by supplementing with physiological level of glucose. Taken together, our findings uncover a pathway by which intracellular bacteria hijack macrophages to invade the brain, suggesting that lipid metabolism might play a role in the prevention or resolution of bacterial neuroinvasion.

Keywords: intracellular bacteria; neuroinvasion; CD36[+] macrophage; ketogenesis; β-hydroxybutyrate

Highlights

1. Intracellular bacteria hijack CD36[+] Mφ for brain invasion without disruption of BBB.

2. CD36[+] Mφ has enhanced adhesion to CVECs and increased resistance to shear flow.

3. Ketogenesis promotes bacterial neuroinvasion that can be halted by glucose treatment.

4. Serum β-hydroxybutyrate and CD36[+] Mφ accumulation fuel bacterial meningitis in human.

Introduction

Meningitis is an inflammation of the protective membranes covering the brain that caused by pathogen infection.[1,2] As a life-threatening infection with an extremely high mortality rate, acute bacterial meningitis usually occurs secondary to infected organs. The blood-brain barrier (BBB) is the foremost line of defense, which consists of brain endothelial cells working in concert with pericytes, smooth muscle cells, astrocytes, microglia and macrophages to protect the central nervous system from infections.[3] Currently, pathogens in the bloodstream are thought to penetrate the BBB via three

main pathways: transcellular entry, paracellular entry or so-called "Trojan horse"

spread.[4] It is known that some bacteria can invade the brain directly by accumulating

during transcytosis or disrupting the tight junction, accompanied by BBB leakage. For

example, *Streptococcus pneumoniae* (*S. pneumoniae*) interacts with host receptors on

endothelial cell monolayers to achieve brain invasion by transcytosis.[5-7] *Neisseria*

meningitidis hijacks specific host-pathogen interactions to redistribute the tight

junction or adhesion proteins, leading to BBB disruption.[8] However, the route of

"Trojan horse" spread by which bacteria circumvent without damaging the BBB has

rarely been reported. Some intracellular bacteria, including *Listeria monocytogenes*

(*L. monocytogenes*) and *Mycobacterium tuberculosis* (*M. tuberculosis*), can invade the

brain tissue with the assistance of host cells, [9,10] but the underlying mechanisms remain

unclear. Efforts to understand the factors that promote the neuroinvasion of

intracellular bacteria and host responses to these pathogens are urgently needed.

Phagocytes, such as neutrophils and macrophages, play dominant roles in host

response to intracellular bacteria.[11] The primary function of phagocytes is to uptake

and destroy microorganisms and cell debris through tissue circulation and migration.

[12,13] Via phagocytosis and antigen presentation, phagocytes act rapidly as the first line

of defense against bacterial infections. However, intracellular bacteria can employ

sophisticated strategies to replicate within phagocytes and evade host immune

surveillance.[14-16] Recently, Marc Lecuit et al. showed that the surface protein

internalin B of *L. monocytogenes* protects infected monocytes from immune killing

by CD8[+] T cells, thus prolonging the lifespan of infected cells and facilitating the

spread of infection throughout the central nervous system.[16] In this context,

phagocytes, especially monocytes and macrophages, might play important roles

during neuroinvasion, but exactly how intracellular bacteria take advantage of

infected macrophages is not well understood.

The metabolism of host immune cells is reprogrammed during infection. Metabolic

signaling helps awaken the immune cells to resist pathogenic invasion. For example,

glycolysis bolsters T cell immunity to prevent *L. monocytogenes* infection.[17] In

addition, many studies have shown that bacteria can actively or passively manipulate

host metabolites to evade clearance by the host immune system. [14,18,19] *Legionella*

pneumophila induces Warburg-like metabolism by secreting the type 4 secretion

system effector MitF and decreases oxidative phosphorylation (OXPHOS) in infected

macrophages to increase the pathogen's likelihood of survival.[18] A recent report

indicates that Nav1.8[+] nociceptors send signals to meningeal macrophages via the

calcitonin gene-related peptide, and this neuroimmune axis suppresses host defenses

and exacerbates bacterial meningitis.[19] These findings imply a potentially close

connection between host metabolism and bacterial brain invasion.

Ketogenesis, induced by fasting or dietary supplementation, is a metabolic pathway

that provides an alternative form of energy for the body and is closely related to host

immune response. [20,21] A recent study reports that in pneumonia caused by

Pseudomonas aeruginosa, ketone bodies are recycled to the lungs via OXPHOS to

maintain airway homeostasis. [22] Consistently, a ketogenic diet is reported to promotes

T cell responses in pulmonary viral infections.[23,24] However, researchers have also

found that pathogens can achieve intracellular survival and infection progression by

manipulating host ketogenesis.[25] Mice fed a ketogenic diet develop a

hyperinflammatory response to systemic lipopolysaccharide and an increased

mortality rate, phenomena that are independent of diet-induced microbiota and

hyperglycemia.[25] Therefore, the role of ketogenesis in the progression and migration

of infections is controversial.[26] β-hydroxybutyrate (BHB) produced by ketogenesis is

well known to promote the accumulation of intracellular lipid bodies, which serve as

both a source of nutrients and a secure niche for intracellular bacteria.[27] However, how

ketogenesis and BHB participate in regulating monocytic/macrophagic homeostasis

and their roles in bacterial neuroinvasion are poorly defined.

In this study, we discovered a novel subpopulation of $CD36^+$ macrophages that served

as immune " Trojan horses" that crosses but does not destroy the BBB, enabling

intracellular bacterial neuroinvasion. This macrophage subset has a foamy-like

phenotype, which improves the cells' ability to stay and migrate vertically on the

cerebrovascular endothelial surface under the mechanical stress of blood flow.

Furthermore, we found that serum BHB, which might be produced by a ketogenic

diet, exacerbated bacterial neuroinvasion by increasing the formation and promoting

the survival of $CD36^+$ macrophages in peripheral blood. More importantly, we

demonstrated that the perturbation of lipid metabolism during bacterial brain invasion

could be rectified by supplementing mice with physiological level of glucose, thereby

suppressing the generation of CD36$^+$ macrophages and greatly attenuating the clinical

symptoms of meningitis. Our study outlines a new common strategy employed by

intracellular bacteria in brain invasion, which could be a potential therapeutic target in

the treatment of bacterial meningitis.

Results

CD36$^+$ macrophages are potential carriers of bacteria during brain invasion

To study the role of immune cells in bacterial neuroinvasion, we constructed an acute

meningitis mice model of infection, and found that at 24 hours post infection, *L.*

monocytogenes massively invaded the brain and the mice began to die (Figure S1A

and S1B). Evans blue staining showed that no extravasation of dye in the whole brain,

indicating no disruption of BBB in this acute meningitis model (Figure S1C and

S1D). Then, peripheral blood mononuclear cells (PBMCs) were collected and single-

cell RNA-sequencing (scRNA-seq) was performed (Figure 1A). Immune cell lineages

were identified using canonical surface markers of dendritic cells (DCs: *Xcr1, Flt3,*

CD209a), NK cells (*Nkg7*), T cells (*CD3*), monocyte/macrophages (Mo/Mφ: *Fcgr1,*

Adgre1), B cells (*CD79*), neutrophils (*S100a9, S100a8*; Figure 1B and 1C).[28] Among

these lineages, Mo/Mφ are involved in phagocytosis and intracellular killing of

microorganisms; however, intracellular bacteria can also live in these cells to achieve

long-term latent infection.[29] In this case, Mo/Mφ (n=3535 cells) were extracted and

reclustered *in silico* to precisely delineate subpopulations and their distributions

before and after neuroinvasion. Based on differentially expressed cell surface markers

and transcripts, we annotated Mo/Mφ into 6 subpopulations (Figure 1D, S1E). Results

showed that the percentage of Cluster 2 cells was sharply increased in the

neurolisteriosis group (Figure S1F). Further, we found that Cluster 2 cells expressed

high levels of genes related to lipid accumulation, including *CD36*, peroxisome

proliferator–activated receptor gamma (*Pparg*), fatty-acid binding protein 4 (*Fabp4*),

CD9, apolipoprotein E epsilon (*ApoE*), and triggering receptor expressed on myeloid

cells 2 (*Trem2*; Figure. 1E).[30-34] These data implied that the functional role of Cluster 2

(CD36$^+$) macrophages was closely related to lipid metabolism.

To understand the characteristics of Cluster 2 macrophages, CD36$^+$ and CD36$^-$

macrophages were isolated from peripheral blood by fluorescence-activated cell

sorting. Compared with CD36$^-$ macrophages, the CD36$^+$ subpopulation exhibited

significantly higher granularity (SSC^{hi}; Figure S1G), which indicated the number of

proteins or organelles they contained and the protrusions on their surfaces. The sorted

macrophages were cultured *in vitro* overnight and then labeled with BODIPY FL, a

lipophilic probe for polar and neutral lipids. Results showed that the CD36$^+$

subpopulation accumulated significantly more lipid droplets than the CD36$^-$

macrophages (Figure S1H). We also performed Oil Red O staining to validate this

finding (Figure S1H). To confirm the lipid uptake ability of the CD36$^+$ macrophages,

we used fluorescently labeled 1,1′-dioctadecyl-3,3,3′,3′-tetramethylindocarbocyanine

perchlorate (Dil)–conjugated oxidized low-density lipoprotein (oxLDL) particles and

our finding showed that CD36$^+$ macrophages took up more than twice as much ox-

LDL as CD36$^-$ macrophages (Figure S1H). These data indicated that lipid metabolism

in Cluster 2 macrophages was dysregulated during brain invasion.

Next, we performed flow cytometry to detect percentage changes in CD36$^+$

macrophages over time during neurolisteriosis (Figure S1I). The results showed that

the percentage of CD36$^+$ macrophages in peripheral blood steadily increased over

time during neuroinvasion (Figure 1F). The CD36$^+$ subgroup could exceed 52% of all

macrophages (Figure 1G). In addition, the percentage of CD36$^+$ macrophages in

peripheral blood were positively associated with bacterial load in the brain (Figure

1H). These data implied that CD36$^+$ macrophages might be involved in bacterial

neuroinvasion. Furthermore, immunofluorescence co-localization analysis was used

to validate this hypothesis. The results showed that the number of CD36$^+$

macrophages attached to cerebrovascular endothelial cells (CVECs) increased as

neuroinvasion progressed in the acute meningitis model (Figure 1I). Taken together,

these results suggested that CD36$^+$ macrophage might play an important role in

bacterial brain invasion.

CD36$^+$ macrophages facilitate transport of intracellular bacteria across the

blood-brain barrier

Next, we sought to understand how necessary CD36$^+$ macrophages were to bacterial

neuroinvasion. Using mouse primary CVECs and neurons in a Transwell system, we

constructed an *in vitro* BBB model (Figure 2A). Ox-LDL has been reported to induce

a foamy-like phenotype by enhancing CD36 expression[35]; therefore, we converted

peritoneal macrophages isolated from mice into $CD36^+$ macrophages by treating them

with different concentrations of ox-LDL for 24 hours (Figure S2A). Subsequently,

macrophages were infected with *L. monocytogenes* (MOI=10) and then inoculated in

the upper Transwell chamber of the BBB model. We confirmed that intracellular

bacterial load increased with the transformation of peritoneal macrophages into

$CD36^+$ macrophages (Figure S2B). Of note, the assistance of ox-LDL–stimulated

macrophages significantly increased the number of *L. monocytogenes* organisms

crossing the BBB in a dose-dependent manner (Figure 2B). Using crystal violet

staining, we also demonstrated a dose-dependent increase in the number of

macrophages penetrating the endothelial monolayer upon stimulation with ox-LDL

(Figure 2C and 2D).

To understand whether the increase in $CD36^+$ macrophages favored the transfer of *L.*

monocytogenes into the brain *in vivo*, we performed adoptive macrophage transfer

experiments through tail vein injection based on a cyclophosphamide (CTX)-induced

immunosuppressive mouse model (Figure 2E). Consistent with previous studies, *L.*

monocytogene-induced clinical phenotype and mortality were relieved in the mice

transferred with purified peritoneal macrophages ($CD36^+$ Mφ group; Figure 2F, S2C

and S2D).[36,37] However, body weight was reduced, comprehensive clinical scoring was

increased, and survival time was shortened in the mice transferred with CD36$^+$

macrophages compared with the control (CD36$^+$ Mφ group; Figure 2F, S2C and S2D).

Furthermore, we showed that brain bacterial loads in mice transferred with CD36$^+$

macrophages were significantly higher than those in mice transferred with purified

peritoneal macrophages (Figure 2G). Interestingly, bacterial loads did not differ

among the lung, spleen, and liver (Figure S2E). We also tested proinflammatory factor

levels in plasma and brain homogenate. The results showed that levels of interleukin 6

(IL-6) and tumor necrosis factor alpha (TNF-α) were remarkably increased (Figure

2H and 2I), indicating a more severe phenotype of meningitis in the CD36$^+$

macrophage transferred group. However, brains of mice transferred with CD36$^+$

macrophages exhibited no Evans blue extravasation, suggesting that these

macrophages facilitated the transfer of *L. monocytogenes* into the brain without

disrupting the BBB (Figure S2F and S2G). Pathological histology analysis confirmed

the existence of more severe inflammation and neuronal damage in the cerebral

cortices in mice transferred with CD36$^+$ macrophages than in those of control (Figure

S2F and S2H). In addition, when mice were infected with a relatively low dose (5×

10^7 CFU/mouse) of *L. monocytogenes*, the difference in bacterial load in brain tissues

between CD36$^+$ macrophage transferred mice and the control group was even greater

than that under high dose condition (Figure S2I and S2J). These data suggested that

neuroinvasion mediated by CD36$^+$ macrophages were not restricted to an acute

meningitis model induced by high dose bacteria.

We then wondered whether CD36[+] macrophages might play the same role during

brain invasion by other intracellular bacteria. In the relevant experiment, a positive

correlation between CD36[+] macrophages and neuroinvasion was observed in

meningitis models of the intracellular bacterium *Salmonella typhimurium* (*S.*

typhimurium; Figure S3A and S3B). We observed the same result in neuroinvasion

model of Methicillin-resistant *Staphylococcus aureus* (MRSA; Figure S3C and S3D),

which, although not a classical intracellular pathogen, exhibits significant intracellular

habitation.[38] However, this finding was not validated in a neuroinvasion model of the

extracellular bacterium *S. pneumoniae* (Figure S3E and S3F). Next, we performed

adoptive macrophage transfer experiments to further verify the carrier role of CD36[+]

macrophages in intracellular bacterial invasion of the brain. We again observed

differences in brain tissue bacterial loads between the CD36[+] macrophage transferred

and control groups in the meningitis models of MRSA (Fig. S3G-S3K), but not in

models using *S. pneumoniae* (Figure S3L-S3O). Taken together, these data suggested

that CD36[+] macrophages greatly contributed to the process of intracellular bacterial

neuroinvasion.

Proprotein convertase subtilisin/kexin 9 (PCSK9) belongs to the proprotein

convertase family, whose members participate in cholesterol and fatty acid (FA)

metabolism. [39,40] Intracellular low-density lipoprotein cholesterol (LDL-C)

accumulates because LDL receptor located on the cell membrane cannot be degraded

normally in PCSK9[-/-] mice.[41,42] We found that PCSK9-deficient macrophages had

significantly higher lipid content compared with control (Figure S4A and S4B) and

that more *L. monocytogene* organisms parasitized PCSK9$^{-/-}$ macrophages than they

parasitized wild-type (WT) cells (Figure S4C). During neurolisteriosis, PCSK9$^{-/-}$ mice

had higher percentages of CD36$^+$ macrophages in peritoneal blood than their

littermates (Figure S4D). Therefore, we used PCSK9$^{-/-}$ mice as a model of

spontaneously forming CD36$^+$ macrophages, and verified the reliability of our

findings through adoptive macrophages transfer. After recovery from transfer for 12

hours, recipient mice were intraperitoneally injected with 2×10^8 CFU of *L.*

monocytogenes; 12 h later, samples were collected for further examinations (Figure

2J). We found that brain bacterial loads were significantly higher in PCSK9$^{+/+}$

recipients that had been transferred with PCSK9$^{-/-}$ macrophages than in those that had

received WT macrophages (Figure 2K). Similar results were observed in the PCSK9$^{-/-}$

recipients (Figure 2K). However, differences in bacterial load manifested in brain

tissues only, not in other organs such as the lung, spleen, and liver (Figure S4E).

Consistently, the mortality rate, and pro-inflammatory factors in the plasma and brain

homogenate were aggravated in both PCSK9$^{+/+}$ and PCSK9$^{-/-}$ recipients by transfer of

PCSK9$^{-/-}$ macrophages (Figure 2L, S4F and S4G). Since *L. monocytogene* is a

foodborne pathogen that can invade the body through the gastrointestinal tract, we

further confirmed the above-mentioned finding by oral administration of *L.*

monocytogenes in the same experimental model (Figure 2M and 2N, S4H-S4J). Taken

together, these data indicated that excessive lipid accumulation contributed to the

formation of CD36$^+$ macrophages, thereby facilitating neuroinvasion by intracellular

bacteria.

Shear flow analysis demonstrates increased adhesion and resistance of CD36$^+$

macrophages during neuroinvasion

We next asked how CD36$^+$ macrophages exploited their properties to accumulate on

the cerebral blood vessel wall and participate in the process of intracellular bacterial

neuroinvasion. Using a two-dimensional live-cell flow chamber assay with a

microfluidic chip [43], we deciphered the multistep process of how macrophages

adhered to and moved on the surface of vascular endothelium under physiological

flow (Figure 3A). Cellular adhesion ability was calculated by counting total

macrophages adhering to the endothelial monolayer under a given shear stress. We

found that as the percentage of CD36$^+$ macrophages increased, so did the number of

cells adhering to the endothelial monolayer (Figure 3B and 3C). Because adherent

cells could be detached by an increase in shear stress, we tested the dynamics of

CD36$^+$ macrophages' shear resistance. The results showed that such resistance was

significantly stronger than that of control macrophages: CD36$^+$ macrophages

outstandingly resisted high shear stress (Figure 3D). Further, we investigated the

effect of endothelial density on macrophagic adhesion. Results showed that

endothelial cell density, rather than cell type, greatly affected macrophagic adhesion,

and CD36$^+$ macrophages tended to adhere to the surface of dense endothelial cells

(Figure S5A-S5D).

Once attached to the endothelial monolayer, the macrophage crawls along it and finds

a place to cross it in the presence of blood flow. [44,45] We next tested the performance of

CD36$^+$ macrophages in post-arrest crawling on the endothelial monolayer. We

acquired sequential images of macrophages at a shear stress of 2 dyn/cm^2 and then

analyzed them by defining various parameters of crawling dynamics (Figure S5E).

The observations were presented by crawling trajectories of each individual cells

(Figure 3E), and we found that the crawling ability of adherent CD36$^+$ macrophages

was dampened, reflecting in shortened crawling displacement and slower crawling

speed (Figure 3F and 3G). However, we saw no obvious differences in crawling

duration (Figure S5F). Directionality is also a critical aspect of cell crawling

dynamics, it is quantified as the ratio of displacement along shear flow to the

perpendicular. Our results showed that the directionality of macrophages was slightly

but significantly enhanced by ox-LDL stimulation, implying that macrophagic

foaminess tended to drive cell crawling in the vertical direction (Figure 3H). We also

compared a key dynamic parameter of transmigration duration, which is defined as

the interval from the timepoint of cell arrival at the site to the endpoint of completion

of transmigration beneath the endothelial monolayer (Figure S5G). The result showed

that transmigration duration was significantly shortened in CD36$^+$ macrophage

(Figure 3I).

Considering that protrusion is critical for cell crawling and migration, we then

investigated whether intercellular lipid could affect the formation of protrusions. The

results showed that the numbers, as well as the lengths of protrusions were

significantly increased in CD36[+] macrophages (Figure 3J and 3K). We further

quantified macrophagic cell area and found that the mean area of CD36[+] macrophages

was significantly larger than the control (Figure 3L). Additionally, we examined

various integrins and adhesive ligands expressed on the cell surface, which have been

reported to be involved in adhesion and migration.[46] The results showed that the

expression levels of α-integrins, β-integrins and intercellular adhesion molecules were

significantly upregulated in the infected CD36[+] macrophages (Figure 3M).

Collectively, these data suggested that the foaminess of macrophages greatly

enhanced their ability to adhere under mechanical stress, predisposing them to stay

and to migrate vertically on the surface of CVECs; this might be attributable to the

formation of protrusions as well as the expression of integrins and adhesion ligands.

β-hydroxybutyrate promotes CD36[+] macrophage formation and exacerbates neuroinvasion

Next, we performed lipid metabolomics analysis of serum samples to reveal how

CD36[+] macrophages were formed during bacterial neuroinvasion. Principal

component analysis (PCA) showed that distinct metabolite genotype clusters were

generated in the neurolisteriosis group (Figure 4A). The volcano plot indicated that 70

FAs were upregulated and 71 FAs were downregulated ($p < 0.05$, $|\log_2(\text{fold change})| > 1$) in the neurolisteriosis group (Figure 4B, Table S1). The 70 upregulated FAs

were classified into four subclusters by the number of carbon atoms in the alkyl chain.

We found thatβ-hydroxybutyrate (BHB), which has been reported to favor the foamy-

like phenotype of macrophages [47], was the only short-chain FA among all upregulated

FAs (Figure 4C). Kyoto Encyclopedia of Genes and Genomes (KEGG)-based analysis

revealed that metabolites related to the synthesis and degradation of ketone bodies

were enriched (Figure S6A), and metabolomics data showed that BHB was

significantly increased during neuroinvasion (Figure S6B). Further, we found that

BHB levels in mice serum climbed steadily over time in the acute meningitis model of

L. monocytogenes infection (Figure 4D).

Then, we sought to explore the reason of enhanced ketogenesis during the brain

invasion. Liver tissues of acute meningitis mice model were collected for

transcriptome analysis by RNA-sequencing (RNA-seq). The results showed that

compared with control group, the genes associated with ketogenesis were sharply

upregulated in the liver of mice suffered from neurolisteriosis, while these genes were

moderately increased in the mild infection group (Figure S6C). KEGG-based analysis

and gene-set enrichment analysis (GSEA) among the neurolisteriosis and control

group revealed that the differentially expressed (DE) genes were enriched in

glycolysis or gluconeogenesis (Figure S6D and S6E). To validate this finding, we

measured the serum glucose levels of mice and found that glucose levels significantly

decreased and remained low during the *L. monocytogene*-induced neuroinvasion

(Figure S6F); meanwhile, we observed a significant decrease in cellular energy and an

increase in ADP/ATP ratios in mice with neurolisteriosis compared to the control

group (Figure S6G and S6H). These data implied that the decrease in glucose level

and increase in ketogenesis may be induced by excessive glycolysis during *L.*

monocytogene-induced neuroinvasion. Further, we showed that the extracellular

acidification rates (ECAR) were significantly increased in the livers of mice with

neurolisteriosis (Figure S6I). This data was confirmed by the measurement of lactate,

which is the end product of glycolysis (Figure S6J). Also, we found that mice

experienced weight loss and reduced food intake, which to a certain extent

contributed to the enhancement of ketogenesis after infection (Figure S6K and S6L).

Interestingly, there was no significant difference in these indicators between the mild

infection group and the control group during the experiments (Figure S6F-S6L).

Furthermore, a steady increase of serum BHB and decrease of glucose level was also

observed in the meningitis models established using *S. typhimurium* and MRSA

(Figure S6M and S6N). These data indicated that production of BHB was

significantly induced during bacterial neuroinvasion, and it was probably due to the

low serum glucose level induced by excessive glycolysis.

As previously reported, autocrine or paracrine BHB directly activates macrophages

function mainly via the hydroxycarboxylic acid receptor 2 (HCAR2).[27] Therefore, we

next used HCAR2 knockout mice to study whether BHB could promote bacterial

neuroinvasion. We confirmed that lipid accumulation and percentage of CD36[+]

macrophages all increased dramatically upon BHB stimulation, and there was no

significant difference when HCAR2 was knocked out (Figure 4E-4G). BHB inhibits

adenylyl cyclase activity through the HCAR2 receptor and reduces cellular cyclic

adenosine monophosphate (cAMP) levels, thereby protecting lipid droplets from

lipolysis by hormone-sensitive lipases (HSL).[27] As a result, we found that cAMP

pretreatment eliminated the decrease in HSL activity, the increase in lipid content and

intercellular bacterial burden stimulated by BHB; however, this phenomenon was not

found in HCAR2-deficient macrophages (Figure S6O-S6Q). By using the *in vitro*

BBB model (Figure 2A), the number of bacteria crossing the barrier increased

significantly in response to BHB stimulation in controls, but not in HCAR2$^{-/-}$

macrophages (Figure 4H). Next, peritoneal macrophages from HCAR2$^{+/+}$ and

HCAR2$^{-/-}$ mice were isolated to further assess the effects of BHB on the biophysical

dynamics of macrophages. The data demonstrated that BHB significantly altered the

static or shear-resistant adhesion of control macrophages, but not of HCAR2$^{-/-}$

macrophages (Figure 4I-4K). We also found that BHB greatly impaired the crawling

ability, including shortened crawling displacement, slowed down crawling speed and

unpredictable directionality of macrophages (Figure 4L-4O). In addition, the

transmigratory ability of macrophages was significantly increased when cells were

pretreated with BHB (Figure 4P). Consistently, the expression levels of the α-integrin

family were significantly upregulated by BHB treatment in controls, but not in

HCAR2-deficient macrophages (Figure 4Q). Taken together, these data indicated that

BHB facilitated the formation of CD36^{+} macrophages through the HCAR2 receptor,

thereby affecting the process of bacterial neuroinvasion.

BHB promotes the survival of macrophages via HCAR2-mediated calcium influx

Because the foamy stage appears to be a terminal point of macrophage differentiation [48], we wondered whether, in addition to converting peritoneal macrophages to the foamy phenotype, cell survival also contributed to the increase in absolute quantity of CD36$^+$ macrophages induced by BHB treatment. The results showed that the lactate dehydrogenase (LDH) activity in the culture medium was significantly reduced in response to BHB treatment, indicating a decrease in cell death (Figure 5A). Propidium iodide (PI) staining was used to evaluate the membrane integrity of infected macrophages, and we found that the proportion of PI$^+$ cells decreased from 59.7% to 24.5% under BHB stimulation, with no significant difference in cell viability between the BHB treatment and control groups when HCAR2 was knocked out (Figure 5B and 5C). Calcium influx is generally recognized as a signal that triggers cell membrane repair.[49] To investigate the impact of BHB-HCAR2 signaling on cytosolic calcium concentration, infected macrophages were loaded with Fluo-4 AM, a fluorescent Ca^{2+} indicator. The results showed that BHB induced remarkable accumulation of intracellular calcium, and HCAR2 knockdown eliminated calcium influx in macrophages (Figure 5D). Pretreatment of macrophages with BAPTA, a Ca^{2+} chelator, reversed the BHB-induced increase in cell survival (Figure 5E and 5F). Further, we tested whether calcium flux was associated with CD36$^+$ macrophage-mediated intracellular bacterial penetration of the BBB. As a result, inhibition of calcium flux had no effect on bacterial uptake by macrophages in the upper chamber (Figure S7A); however, we observed differences in bacterial loads in the macrophages

that had crossed the barrier (Figure 5G).

We next investigated how BHB maintained the survival of CD36[+] macrophages. As

previously reported, calcium influx initiates membrane repair mainly through

recruitment of the endosomal sorting complexes required for transport (ESCRT)

machinery.[50] Our data showed that BHB treatment significantly increased the number

of puncta that were positive for charged multivesicular body protein 4B (CHMP4B),

which indicated assembly of ESCRT in the infected macrophages; HCAR2

knockdown or BAPTA treatment could block this membrane repair process (Figure

5H and 5I, S7B and S7C).Cellular membrane repair also relies on lysosomes, which

can rapidly fuse with the plasma membrane when intracellular Ca^{2+} concentration

rises. Therefore, we examined lysosomal exocytosis in the infected macrophages

using exofacial lysosomal-associated membrane protein 1 (LAMP1) and the release of

lysosomal enzymes, such as acid phosphatase (ACP), β-galactosidase (β-gal) and β-

hexosaminidase (β-hex). The results confirmed that BHB promoted the lysosomal

exocytosis in infected macrophages through the HCAR2 receptor (Figure 5J-5L, S6D-

S6F). These data collectively indicated that BHB signaling activated Ca^{2+}-mediated

membrane repair, thereby maintaining the survival of CD36[+] macrophages via the

HCAR2 receptor.

Ketogenesis exacerbates intracellular bacterial neuroinvasion

BHB can be synthesized via the metabolism of FAs and ketogenesis.[51] To understand

the relationship between ketogenesis and bacterial meningitis, we gave mice different

doses of ketone ester drink for 4 weeks, and then infected them with *L.*

monocytogenes via intraperitoneal injection (Figure 6A). The data showed that ketone

ester elevated the levels of serum BHB and the proportion of CD36$^+$ macrophages in

peritoneal blood (Figure 6B and S8A), accompanied by remarkable increases in the

clinical scores, mortality rate and bacterial load in brain tissues (Figure 6C and 6D,

S8B). Dietary modulation can increase and maintain circulating metabolites in human

circulation, and a growing amount of evidence demonstrates that a ketogenic diet

elevates BHB levels.[52] The ketogenic diet is a low-carb, high-fat diet that causes

weight loss and provides numerous health benefits, including alleviating symptoms of

neurodegenerative diseases, cardiovascular diseases, and cancers.[21,53-55] We next asked

whether a ketogenic diet was associated with neurolisteriosis. HCAR2$^{-/-}$ mice and

their littermates were fed a normal or ketogenic diet for 4 weeks respectively, and

then infected with *L. monocytogenes* by intraperitoneally injection (Figure 6E). As a

result, the lifespan of HCAR2$^{+/+}$ mice fed a ketogenic diet was significantly shortened

under *L. monocytogenes* infection, and the survival disadvantage caused by dietary

modulation was abrogated by HCAR2 knockout (Figure 6F). Consistently, the clinical

scores (Figure. S8C), percentages of CD36$^+$ macrophages in peritoneal blood (Figure

6G), the intracellular bacterial burden of macrophages (Figure 6H), and bacterial

loads in brain tissue (Figure 6I) were all greatly increased in HCAR2$^{+/+}$ mice on a

ketogenic diet. Interestingly, mice in this group lost less weight than mice in the

control group, and experienced greater spread of bacteria to other organs (Figure 6I

and S8D). We then examined the levels of pro-inflammatory factors in the plasma and

brain homogenate. The results showed that a ketogenic diet significantly increased the

levels of pro-inflammatory cytokines and induced more severe neuronal damage in

$HCAR2^{+/+}$ mice, while it had no effect on $HCAR2^{-/-}$ mice (Figure 6J-6M). Notably,

Evans blue extravasation was not detected in the brain of mice fed a ketogenic diet,

ruling out the possibility of direct damage to the BBB (Figure 6L and S8E). These

data were confirmed by oral administration of *L. monocytogenes* in the same

experimental model (Figure S8F-S8I). We also used MRSA to further verify our

findings. The data indicated that, as expected, the ketogenic diet promoted brain

invasion by MRSA and exacerbated symptoms of meningitis (Figure S8J-S8L). Taken

together, our data showed that a ketogenic diet could exacerbate bacterial

neuroinvasion and hasten the demise of mice, primarily depending on the elevation in

serum BHB via the HCAR2 receptor.

Our data thus far suggested that BHB played an important role in bacterial

neuroinvasion in mice, and the enhancement in ketogenesis was probably induced by

the low serum glucose level. We then evaluated the effects of restoring blood glucose

level on the outcomes of infected mice. The results showed that under the acute

meningitis model of *L. monocytogenes* infection, supplementing mice with glucose

every 12 hours to restore blood glucose levels resulted in a significant decrease in

BHB levels and $CD36^+$ macrophages percentages (Figure S9A-S9D). Surprisingly, we

showed that four rounds of glucose supplementation significantly improved the

survival rate of mice, reduced their brain bacterial burden and pro-inflammatory

cytokines levels (Figure S9E-S9I). These data indicated that glucose homeostasis

restricted the brain invasion of *L. monocytogenes* infection and alleviated meningitis

symptoms in mice.

Serum BHB fuels the symptoms of human bacterial meningitis

A retrospective study that based on the Medical Information Mart for Intensive Care

IV (MIMIC-IV) was performed to explore the risk factors of bacterial meningitis in

human. As a result, ferritin (OR 5.09, 95% CI 1.55-16.71), fibrinogen (OR 3.57, 95%

CI 1.13-11.25), LDL (OR 3.31, 95% CI 1.01-10.83), glucose (OR 2.17, 95% CI 1.04-

4.50) and aspartate amino transferase (OR 2.16, 95% CI 1.01-4.63) were identified as

independent risk factors, whereas potassium (OR 0.42, 95% CI 0.20-0.87), alkaline

phosphatase (AP, OR 0.38, 95% CI 0.18-0.83), phosphate (OR 0.23, 95% CI 0.11-

0.50), high-density lipoprotein (HDL, OR 0.19, 95% CI 0.05-0.66) and transferrin

(OR 0.14, 95% CI 0.04-0.50) were identified as protective factors for patients with

bacterial meningitis (Figure S10A). Next, we asked whether the accumulation of

CD36$^+$ macrophages or BHB could also be identified and were associated with

bacterial meningitis in human clinical samples. Gender- and age-matched bacterial

meningitis subjects and healthy individuals were recruited and peripheral blood

samples were collected for analysis (Figure 7A). Since blood glucose is a risk factor

for meningitis and BHB level can be dramatically affected by glycolysis, subjects

with abnormal blood glucose or insulin concentrations were excluded from the two

groups (Figure S10B-S10E). As a result, serum BHB levels were significantly

elevated in meningitis patients compared with healthy controls, suggesting an

infection-induced ketogenesis (Figure 7B). Interestingly, serum cholesterol levels,

including total cholesterol (TC) and LDL-C were increased in meningitis patients

compared with healthy controls, while HDL-C was slightly but not significant

reduced (Figure S10F-10H). This finding was consistent with data from our

retrospective study based on MIMIC-IV (Figure S10A). To investigate whether BHB

was associated the severity of infection, we further divided patients into three groups,

high BHB (in the top 1/3 among samples tested, n=21), moderate (middle 1/3, n=20)

and low (bottom 1/3, n=21; Figure 7A). Clearly, patients with high BHB levels

exhibited higher clinical scores than patients with low BHB levels (Figure 7C).

Consistently, high-BHB patients had higher cell counts and more protein in the

cerebrospinal fluid (CSF; Figure 7D and 7E); higher infection indices such as white

blood cell counts (WBC; Figure 7F) and procalcitonin (PCT; Figure 7G); and higher

values for markers of inflammatory conditions such as erythrocyte sedimentation rate

(ESR; Figure 7H) and C-reactive protein (CRP; Figure 7I), suggesting increased

disease severity. These data implied that infection-induced ketogenesis might provide

important fuel for severe bacterial neuroinvasion.

To explore the dynamics of CD36$^+$ macrophages and BHB in the pathogenesis of

bacterial meningitis, we followed-up with 7 individuals during their hospitalization.

We found that CD36$^+$ macrophages and BHB levels in patients tended to increase

during disease progression, and decrease during the recovery stage (Figure 7J and

7K). To further confirm the relationship between CD36[+] macrophages and bacterial

meningitis pathology, we collected PBMCs from patients or healthy controls for

analysis. Compared with healthy controls, the expression levels of genes involved in

cholesterol uptake were significantly increased, while those of genes related to

cholesterol utilization and excretion were decreased in macrophages from meningitis

patients (Figure 7L). Taken together, these data indicated that BHB fueled human

bacterial meningitis by promoting the development of CD36[+] macrophages.

Discussion

Our findings suggested that CD36[+] macrophage might serve as a carrier for

intracellular bacteria that colonize in the brain, facilitating the penetration of these

bacteria across the BBB. Generally, the BBB prevents pathogens from crossing unless

there is significant trauma to the brain. However, specific bacteria can break through

the BBB via their unique mechanisms and cause severe brain infections[1,16]. In this

study, we discovered that intracellular bacteria may prefer to exploit a "Trojan horse"

pathway, which causes little scathe to the BBB but enables acute brain invasion.

CD36[+] macrophages could be hijacked by intracellular bacteria and serve as vehicles

for neuroinvasion. Compared with peritoneal macrophages, CD36[+] macrophages tend

to have more protrusions and greater cell area, which make them more conducive to

colonization on CVECs under the pressure of blood flow. Based on the above-

mentioned remarkable characteristics, CD36[+] macrophages have the potential to

become carriers of engineered intracellular bacteria for bacteriotherapy.

Pathogens can exploit most of the host's nutrients, including glycerol, FAs, and

several other carbon substances, to reprogram immune cells.[56] Our findings

demonstrated that intracellular bacteria can actively influence host free FA

metabolism (Figure 4A-4C) and reprogram macrophages, thereby exacerbating

neuroinvasion. The foamy-like phenotype of macrophages is formed through

dysregulation of lipid metabolism, and it impairs the immune function of

macrophages and contributes to pathogenesis. For example, foamy macrophage

necrosis was previously reported to expand tuberculous lesions in lung parenchyma,

causing progressive destruction of lung tissue and loss of lung function in patients

with active disease.[57] In this study, we discovered that BHB produced by a ketogenic

diet promoted the formation of a foamy-like phenotype in $CD36^+$ macrophages in

peripheral blood, thereby facilitating bacterial neuroinvasion. Interestingly, we found

that serum cholesterol levels, including TC and LDL-C were significantly higher in

meningitis patients than in healthy controls (Figure S10F and S10H). We hypothesize

that cholesterol or other lipid metabolites might also contribute to the formation of

$CD36^+$ macrophages. However, this hypothesis should be further verified

experimentally.

Our biomechanical data indicated that $CD36^+$ macrophages adhered more strongly to

the endothelial surface than normal macrophages, and that this adhesion-enhancing

effect was more pronounced when endothelial cells were dense (Figure 3B and 3C,

S5A and S5B). This suggested that CD36$^+$ macrophage-mediated bacterial invasion

affected the brain to a greater extent than other tissues. Accordingly, we showed that

the increase in CD36$^+$ macrophages in peripheral blood led to the transfer of bacteria

mainly to the brain, rather than to other organs (Figure 2G and S2E). However, we

found that a ketogenic diet and ketone ester supplementation significantly increased

bacterial load not only in the brain, but also in other tissues such as the liver and lung

(Figure 6D and 6I). This suggested that ketogenesis, in addition to promoting the

generation of CD36$^+$ macrophage, might play other unknown roles in the intracellular

bacterial infections. Therefore, the role of ketogenesis in the formation of CD36$^+$

macrophages and the contribution of ketogenesis to bacterial neuroinvasion should be

further determined.

The host's basal nutritional status may also greatly influence the outcome of infection.

A recent study reported that glucose supplementation in mice has a protective effect

against *Yersinia pseudotuberculosis* infection.[58] Our results indicated that the decrease

in blood glucose level and enhancement of ketogenesis in the liver may be induced by

excessive glycolysis during bacterial neuroinvasion (Figure S6C-S6I). Interestingly,

we found that mice experienced weight loss and decreased appetite during the

bacterial meningitis, and the two events occurred almost simultaneously (Figure S6K

and S6L). These data together imply that maintenance of normal blood sugar levels

may be critical for the development, progression, and prognosis of meningitis.

Therefore, we attempted to restore the blood sugar levels of acute meningitis mice by

intraperitoneal injection of glucose, and surprisingly we found that glucose

supplementation significantly alleviated the clinical symptoms of meningitis, reduced

bacterial burden in the brain and improved the survival rate of mice (Figure S9).

These findings indicate that in addition to antibacterial therapy, nutritional therapy

plays a vital role in the clinical treatment of bacterial meningitis.

Overall, this study demonstrated that a BHB-rich environment created by the

ketogenic diet rendered mice more susceptible to *L. monocytogene*-induced

neuroinvasion. Mechanistically, BHB protected CD36$^+$ macrophages from senescence

via HCAR2-mediated membrane repair. However, inhibition of HCAR2 signaling or

calcium influx reversed the improvement in CD36$^+$ macrophage survival but did not

completely reverse the exacerbation in bacterial neuroinvasion induced by BHB

(Figure 5E-5G). These data suggested that BHB-induced bacterial brain invasion was

not solely dependent on HCAR2 signaling but that other potential mechanisms might

also be involved in this process. However, this hypothesis requires further

experimental verification.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No.

92369115), the Fundamental Research Funds for the Central Universities (No.

2023CDJXY-009 and 2019CDYGZD009), Natural Science Foundation of Chongqing,

Author Contributions

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1. Identification of CD36$^+$ macrophage accumulation during

neurolisteriosis.

(A) Summary of experimental design (n=5 control mice and n=7 mice infected with

2×10^8 CFU *L. monocytogenes* for 24 hours).

(B) Uniform Manifold Approximation and Projection (UMAP) representation of

integrated transcriptome-based clustering. A total of 17351 peripheral blood

mononuclear cells (PBMCs) were clustered and major immune cell lineages were

identified.

(C) Dot plot showing the expression of selected transcripts in immune cell clusters.

All transcripts were significantly enriched in the pertinent clusters with adjusted p <

0.05.

(D) A t-distributed Stochastic Neighbor Embedding (tSNE) analysis of the re-

clustered cells corresponding to monocytes and macrophages (baseline, n=1804;

neurolisteriosis, n=1731).

(E) Expression of the indicated transcripts (encoding genes associated with lipid

accumulation) was projected onto a monocyte/macrophage tSNE plot.

(F-G) An acute mouse neurolisteriosis model was generated by intraperitoneal

injection of 2×10^8 CFU *L. monocytogenes*. Quantification of CD36$^+$ macrophages in

PBMCs (F) and total macrophages (G) was detected by flow cytometry (n=5 for each

group). Experiments were performed in duplicate. Data are presented as mean ± SEM.

*p < 0.05, **p < 0.01.

(H) Correlation between CD36$^+$ macrophages and bacterial burdens in brain during

the neurolisteriosis. R, correlation coefficient.

(I) Immunofluorescence staining of CD36$^+$ macrophages in cerebral vasculature of

mice with neurolisteriosis. CD36[+] macrophages were stained with CD11b (red) and

CD36 (green). The boundary of vascular was depicted in dashed lines. Scale bar, 100

μm.

Abbreviations: PBMC, peripheral blood mononuclear cell; i.p., intraperitoneally;

CFU, colony forming units.

Figure 2. Infected CD36[+] macrophages transfer bacteria into brain parenchyma

***in vitro* and *in vivo*.**

(A) Schematic diagram of an *in vitro* BBB model based on Transwell system.

(B) Quantification of bacterial burden in the macrophages penetrating into the lower

chamber of Transwell system under different concentrations of ox-LDL.

(C-D) (C) Representative image of macrophages that penetrated into the lower

chamber. Scale bar, 100 μm. (D) Quantification of the penetrating cells in each field

of view.

(E) Experimental diagram of adoptive macrophage transfer.

(F-I) After the adoptive macrophage transfer, recipient mice were intraperitoneally

infected with 2×10^8 CFU *L. monocytogenes*. (F) Survival of recipient mice (n=10 for

each group). (G-I) Bacterial burdens in brains (G), and cytokine levels in plasma (H)

and brain homogenate (I) of recipient mice on day 1.25 (n=5 for each group).

(J-N) (J) Schematic diagram of adoptive macrophage transfer and infection

experiments among PCSK9[+/+] (WT) and PCSK9[-/-] (KO) mice. Peritoneal

macrophages from PCSK9[+/+] and PCSK9[-/-] mice were injected into PCSK9[+/+] and

PCSK9$^{-/-}$ mice, respectively, followed by *L. monocytogenes* infection.

(K, M) Bacterial burdens in the brains of recipient mice after intraperitoneal injection

(K) or oral administration (M) of *L. monocytogenes* (n=5 for each group).

(L, N) Survival of recipient mice after intraperitoneal injection (L) or oral

administration (N) of *L. monocytogenes* (n=10 for each group).

Experiments were performed in duplicate (E-N) or triplicate (A-D). Data are

presented as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ox-LDL, oxidized

low density lipoprotein; FOV, field of view; NS, normal saline; PMφ, purified

peritoneal macrophages; CD36$^+$ Mφ, CD36$^+$ macrophages; i.p., intraperitoneally; ns,

nonsignificant.

Figure. 3 Two-dimensional live-cell flow chamber assay of the migration

characteristics of infected CD36$^+$ macrophages.

(A) Schematic diagram of live-cell flow chamber experiment. Primary peritoneal

macrophages were pretreated with ox-LDL for 24 hours and then infected with *L.*

monocytogenes (MOI=10) for 6 hours.

(B-C) (B) Optical images of ox-LDL pretreated macrophages adhering to the surface

of monolayer endothelial cells under shear flow. Scale bar, 200 μm. (C)

Quantification of adherent cells in each field of view.

(D) Resistance to shear flow after adhesion of ox-LDL pretreated macrophages to

endothelial cells. Upon 10-min adherence of equal macrophages to the monolayer

endothelial cells at 1 dyn/cm^2, the shear stress was stepwise increased to 2, 4, 8, 16,

and 32 dyn/cm^2 for 30 s sequentially to detach those adhered cells.

(E-H) Crawling dynamics of ox-LDL pretreated macrophages under shear flow. (E)

Typical trajectories of ox-LDL pretreated macrophages crawling on monolayer

endothelial cells. For each trajectory, the arrested sites of macrophages were overlaid

at the origin of the diagram (x/y=0/0) and the corresponding end points were indicated

with dots. Arrows indicate the flow direction. (F-H) Quantification of crawling

displacement (F), velocity (G) and directness (H).

(I) Transmigration duration of ox-LDL pretreated macrophages after adhesion to the

surface of monolayer endothelial cells.

(J-K) Quantification of the number (J) and mean length (K) of protrusions in ox-LDL

pretreated macrophages.

(L) Quantification of cell area in ox-LDL pretreated macrophages.

(M) qRT-PCR analysis of relative expression of genes associated with intercellular

adhesion molecules.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ox-LDL, oxidized low density lipoprotein.

Figure 4. β-hydroxybutyrate aggravates bacterial neuroinvasion by promoting

the formation of CD36$^+$ macrophages via HCAR2 receptor.

(A-C) Lipid metabolomic profiling analyses of serum from the mice with or without

neurolisteriosis. The mice of neurolisteriosis group were infected with 2×10^8 CFU *L.*

monocytogenes for 24 hours (n=6 for each group). (A) Principal coordinates analysis

(PCA) analysis of differences between groups. (B) Expression of fatty acids (FAs) in

the serum was shown in the volcano plot. Differentially expressed FAs were selected

with |log2 fold change|>1 and adjusted P < 0.05. The dark blue dots represent down-

regulated FAs and the orange dots represent up-regulated FAs. (C) Classification of

the up-regulated FAs (SCFA, short chain fatty acids; MCFA, medium chain fatty

acids; LCFA, long chain fatty acids; VLCFA, very long chain fatty acids) shown in

the pie chart.

(D) Biochemical analysis of serum BHB levels in mice during the neurolisteriosis.

(E-Q) Primary peritoneal macrophages were extracted from 8-week-old HCAR2$^{+/+}$

and HCAR2$^{-/-}$ mice in preparation for further functional analysis. HCAR2$^{+/+}$ and

HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) for 24 hours and then

infected with *L. monocytogenes* (MOI=10) for 6 hours.

(E-F) (E) Flow cytometry analysis of BODYPY FL 493/503 in BHB-treated

macrophages (0, 5, 10 mM). (F) The histogram represents normalized MFI.

(G) Flow cytometry analysis of CD36^{+} macrophages in BHB-treated macrophages (0,

5, 10 mM).

(H) Quantification of bacterial burden in the macrophages penetrating into the lower

chamber of Transwell system. The macrophages stimulated with ox-LDL (80 ng/ml)

were used as positive control.

(I-J) (I) Optical images of adherent macrophages on the surface of monolayer

endothelial cells. Scale bar, 200 μm. (J) Quantification of adherent cells in each field

of view.

(K) Resistance to shear flow after adhesion of macrophages to endothelial cells.

(L-P) Crawling and migrating dynamics of infected macrophages under shear flow.

(L) Typical trajectories of infected macrophages crawling on monolayer endothelial

cells. Quantification of crawling displacement (M), velocity (N), directness (O) and

transmigration duration (P).

(Q) mRNA expression of intercellular α-integrin in infected macrophages.

Experiments were performed in duplicate (D) or triplicate (E-Q). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β -

hydroxybutyrate; MFI, mean value of fluorescence intensity.

Figure 5. β-hydroxybutyrate promotes the survival of macrophage via HCAR2-

mediated calcium influx.

(A-L) Primary peritoneal macrophages were extracted from 8-week-old HCAR2$^{+/+}$

and HCAR2$^{-/-}$ mice, cells were then infected with *L. monocytogenes* (MOI=10) for 6

hours.

(A-C) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM)

before infection. (A) Quantification of the percentage of released lactate

dehydrogenase (LDH) in cell culture medium. (B) Immunofluorescence of Propylene

glycol (PI)-stained macrophages. DAPI (blue) labels the nucleus and PI (red) labels

the cell membrane. (C) Quantification of the percentage of PI^{+} cells after infection.

(D) Flow cytometry analysis of dynamics in intracellular calcium concentration.

Infected macrophages were preloaded with Fluo-4 AM. BHB (10 mM) was added to

the buffer after a span of 30 s.

(E-G) Macrophages were pretreated with BHB (10 mM) or BAPTA (10 mM) for 24

hours before infection. (E) Quantification of the percentage of released lactate

dehydrogenase (LDH) in cell culture medium. (F) Quantification of the percentage of

PI$^+$ cells after infection. (G) Quantification of bacterial burden in the macrophages

penetrating into the lower chamber of Transwell system.

(H-I) (H) Infected macrophages were treated with BHB (10 mM). Time-lapse images

of CHMP4B-EGFP in macrophages were collected at 0, 5, 10, 15, 20, 30 min post

BHB treatment with confocal microscopy. (I) Quantification of CHMP4B-EGFP

puncta.

(J-L) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) for

24 hours before infection. (J) Flow cytometry analysis of LAMP1 expression on the

cell surface. (K) The histogram representing normalized MFI. (L) Quantification of

the lysosomal enzyme (acid phosphatase [ACP], β-galactosidase [β-gal] and β-

hexosaminidase [β-hex]) activity released into the culture medium.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate.

Figure 6. Ketogenesis exacerbates neurolisteriosis through β-hydroxybutyrate-

HCAR2 signaling.

(A-D) (A) Schematic diagram of *in vivo* KE supplementation. C57BL/6 mice were

supplemented with KE (10, 20 mg/ml) in drinking water for 2 weeks and then

challenged with $2×10^8$ CFU *L. monocytogenes*. Serum BHB levels (B) and bacterial

burdens in the parenchymal organs (D) were detected (n=5 for each group). (C)

Survival of mice exposed *L. monocytogenes* (n=10 for each group).

(E-M) (E) Schematic diagram of *in vivo* ketogenic diet. HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice were treated with ketogenic or control diets, respectively, for 4 weeks, followed

by intraperitoneal infection with $2×10^8$ CFU *L. monocytogenes*. (F) Survival of mice

exposed to *L. monocytogenes* (n=10 for each group). (G) Flow cytometry analysis of

CD36$^+$ macrophages before and after neurolisteriosis. (H-M) (H-I) Bacterial burdens

in peripheral blood macrophages (H) and parenchymal organs (I) of mice on day 29.

(J-K) Cytokine levels in the plasma (J) and brain homogenate (K). (L) Representative

images of Evans blue leakage of mouse brains and Hematein & Eosin staining of

cerebral cortex tissue. (M) Pathology scores. Scale bar, 10 μm. (n=5 for each group).

Experiments were performed in duplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant; KE, Ketone ester; CD, control

diet; KD, ketogenic diet; i.p., intraperitoneally.

Figure 7. Levels of β-hydroxybutyrate are decreased in bacterial meningitis in

humans, which correlated with the severity of disease.

(A) Schematic diagram of HC (n=60) and BM (n=62) samples. The BM patients were

divided into three groups according to serum BHB levels.

(B) Serum BHB level in HC and BM patients.

(C-I) Clinical scores (C), cell counts (D) and protein levels (E) in cerebrospinal fluid,

white blood cells (WBC, F), procalcitonin (PCT, G), erythrocyte sedimentation rate

(ESR, H) and C-reactive protein (CRP, I) in BM patients with different BHB levels.

(J-K) Dynamics of CD36$^+$ macrophages (J) and serum BHB (K) in 7 BM donors after

admission. The time after admission was divided into symptomatic and convalescent

periods according to clinical and laboratory indexes of infection. The scatter plot on

the right panel represents the statistics of AUC.

(L) Relative expression of genes involved in lipid metabolism in PBMCs from HC

and BM patients was detected by qRT-PCR.

Data are presented as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns,

nonsignificant; HC, healthy control; BM, bacterial meningitis; AUC, area under the

curve; BHB, β-hydroxybutyrate .

Figure S1. Additional results of identification of CD36$^+$ macrophage

accumulation during the neurolisteriosis, related to Figure 1.

(A-B) C57BL/6 mice were intraperitoneally infected with 2×10^8 CFU *L.*

monocytogenes. (A) Bacterial burdens in brains. (B) Survival of the mice (n=10 for

each group). The yellow shading represents the range of confidence interval.

(C-D) Evans blue assay of the mouse challenged with *L. monocytogene*s (2×10^8

CFU/mouse) or LPS (2 mg/mouse). Representative images (C) and quantification of

Evans blue leakage in the mouse brains (D).

(E) Cell clusters identified in 10× scRNA-seq dataset. The top 10 enriched genes for

each cluster are highlighted. Normalized z-score is used to illustrate gene expression.

(F) Proportion of each cluster according to experimental conditions in Figure 1D.

(G) Gating strategy for analysis of CD36$^+$ macrophages in PBMCs.

(H) CD36$^+$ and CD36$^-$ macrophages were sorted from peripheral blood by flow cytometry and then stained with BODIPY FL 493/503 and oil Red O. Scale bar, 10 μm.

(I) Normalized intake of Dil-oxLDL by CD36$^+$ and CD36$^-$ macrophages.

Experiments were performed in duplicate (A-D) or triplicate (H-I). Data are presented as mean ± SEM. *p < 0.05, **p < 0.01.

Figure S2. Additional results of adoptive transfer of CD36$^+$ macrophages *in vitro* and *in vivo*, related to Figure 2.

(A) Conversion of mice peritoneal macrophages into CD36$^+$ macrophages by treating them with different concentrations of ox-LDL for 24 hours.

(B) Intracellular bacterial burden was quantified in macrophages challenged with *L. monocytogenes* (MOI=10) in the upper chamber of Transwell system under different concentrations of ox-LDL.

 (C-H) The adoptive macrophage transfer experiment was shown in Figure 2E-2I (n=5 for each group). Recipient mice were intraperitoneally infected with 2×10^8 CFU *L. monocytogenes*. (C-D) Body weight changes (C) and clinical scores (D) of recipient mice during neurolisteriosis. (E) Bacterial burdens in other tissues (lung, spleen, liver) of recipient mice on day 1.25. (F-H) (F) Representative images of Evans blue leakage of mouse brains and Hematein & Eosin staining of cerebral cortex tissue.

Quantification of extravasation (G) and (H) pathology scores. Scale bar, 20 μm.

(J-K) After adoptive macrophage transfer, recipient mice were intraperitoneally

infected with $5×10^7$ CFU *L. monocytogenes*. Bacterial burdens in brains (J) and other

tissues (K) of recipient mice on day 1.25 were detected.

Experiments were performed in duplicate (C-J) or triplicate (A-B). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; NS, normal

saline; PMφ, purified peritoneal macrophages; $CD36^+$ Mφ, $CD36^+$ macrophages; i.p.,

intraperitoneally.

Figure S3. Accumulation of $CD36^+$ macrophages exacerbate intracellular

bacteria neuroinvasion.

(A-F) Mice were intravenously infected with *S. Typhimurium* (A-B), MRSA (C-D) or

S. pneumoniae (E-F), respectively (n=5 for each group). (A, C, E) Flow cytometry

analysis of $CD36^+$ macrophages during neuroinvasion. (B, D, F) Correlation between

$CD36^+$ macrophages and bacterial burden in the brain during bacteria neuroinvasion.

R, correlation coefficient.

(G-K) After the adoptive macrophage transfer, recipient mice were intraperitoneally

infected with $2×10^8$ CFU MRSA. (G) Survival of recipient mice (n=10 for each

group). (H-I) Bacterial burdens in the brains (H) and other tissues (lung, spleen, liver).

(J-K) Cytokine levels in plasma (J) and brain homogenate (K) of recipient mice on

day 1.5 (n=5 for each group).

(L-O) Mice were intraperitoneally infected with $1×10^8$ CFU *S. pneumoniae* (n=5 for

each group). (L) Survival rate of mice (n=10). (M) Bacterial burdens in tissues. (N-O)

Cytokine levels in plasma (N) and brain homogenate (O) of recipient mice on day 2

(n=5 for each group).

Experiments were performed in duplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant.

Figure S4. Additional results of bidirectional transfer of macrophages from

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice, related to Figure 2.

(A-B) (A) Flow cytometry analysis of lipid droplets in peritoneal macrophages of

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice. (B) The histogram represents normalized MFI.

(C) Peritoneal macrophages of PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice were infected with *L.*

monocytogenes (MOI=10) for 2 hours and then incubated in a fresh solution

containing 5 μg/mL lysostaphin for 2 or 12 hours. Intracellular bacterial burden of

infected macrophages was quantified

(D) Flow cytometry analysis of CD36^{+} macrophages during neurolisteriosis in

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice, respectively (n=5 for each group).

(E-G) Recipient mice in Figure 2J were intraperitoneally infected with 2×10^{8} CFU *L.*

monocytogenes. (E) Bacterial burdens in other tissues (lung, spleen, liver). (F-G)

Cytokine levels in plasma (F) and brain homogenate (G) (n=5 for each group).

(H-J) Recipient mice in Figure 2J were orally infected with 2×10^{8} CFU *L.*

monocytogenes. (H) Bacterial burdens in other tissues (lung, spleen, liver). (I-J)

Cytokine levels in plasma (I) and brain homogenate (J) (n=5 for each group).

Experiments were performed in duplicate (D-J) or triplicate (A-C). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; i.p.,

intraperitoneally.

Figure S5. Supplemental data for the live-cell flow chamber assays, related to

Figure 3.

(A-D) Optical images of *L. monocytogenes* infected macrophages (with or without ox-

LDL stimulation) on the surface of endothelial monolayers with different densities

(A) or endothelial monolayers derived from different tissues (C). Scale bar, 200 μm.

(B, D) Quantification of adherent cells in each field of view.

(E) Schematic trajectory used for defining macrophage crawling parameters.

(F) Crawling duration of *L. monocytogene*-infected macrophages after adhesion to

endothelial monolayer surface.

(G) Relative fluorescence intensity of a macrophage on endothelial surface over time.

It is a random example to illustrate the definition of transmigration duration. A 70%

decrease in fluorescence intensity after a macrophage crosses a monolayer is

considered successful transmigration.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: CVEC, cerebrovascular endothelial cell; PVEC,

pulmonary vascular endothelial cell; LSEC, liver sinusoidal endothelial cell; TEM,

transendothelial migration; ns, nonsignificant.

Figure S6. Additional results from the influence of β-hydroxybutyrate on

macrophages during neuroinvasion, related to Figure 4.

(A) KEGG enrichment analysis of lipid metabolites.

(B) Mass spectrometry analysis of serum BHB levels.

(C-E) Transcriptomics analysis of liver tissues from mice (n=3 for each group). The

mice of mild infection group were infected with 2×10^6 CFU *L. monocytogenes* for 24

hours while those of neurolisteriosis group were 2×10^8 CFU. (C) Heatmap of

differentially expressed (DE) genes related to ketogenesis. (D) Top20 KEGG

enrichment analysis of function of DE genes between neurolisteriosis group and

baseline group. (E) Gene-set enrichment analysis of glycolysis/gluconeogenesis

between neurolisteriosis group and baseline group.

(F) Biochemical analysis of serum glucose levels in mice during the *L.*

monocytogenes infection.

(G-J) Levels of ATP generation, (G) ratio of ADP/ATP (H), seahorse analysis of the

ECAR (I) and relative abundances of lactate (J) in the liver tissues from mice with or

without *L. monocytogenes* infection for 24 hours.

(K-L) Body weight changes (K) and food uptake (L) were monitored every 6 hours

during neuroinvasion.

(M-N) Biochemical analysis of serum BHB and glucose levels in mice during

neuroinvasion by *S. typhimurium* (M) and MRSA (N).

(O-Q) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) and

cAMP (1 mM) for 24 hours and then infected with *L. monocytogenes* (MOI=10) for 6

hours. Quantification of the HSL activity (O), normalized MFI of BODIPY FL

493/503 (P) and the bacterial burden in the macrophages penetrating into the lower

chamber of Transwell system (Q).

Experiments were performed in duplicate (F-N) or triplicate (O-Q). Data are

presented as mean ± SEM. *$p < 0.05$, **$p < 0.01$. Abbreviations: BL, baseline; NL,

neurolisteriosis; ns, nonsignificant; MFI, mean fluorescence intensity; HSL, hormone-

sensitive lipase.

Figure S7. Additional results on the role of β-hydroxybutyrate in regulating

macrophage survival, related to Figure 5.

(A-F) Primary peritoneal macrophages were extracted and then infected with *L.*

monocytogenes (MOI=10) for 6 hours.

(A) Macrophages were pretreated with BHB (10 mM) or BAPTA (10 mM) for 24

hours before infection. Intracellular bacterial burden of macrophages in the upper

chamber of Transwell system was quantified.

(B-C) (B) Infected macrophages were treated with BHB (10 mM). Time-lapse images

of CHMP4B-EGFP in macrophages were collected at 0, 5, 10, 15, 20, 30 min post

BHB treatment with confocal microscopy. (C) Quantification of CHMP4B-EGFP

puncta.

(D-F) Peritoneal macrophages were pretreated with BHB (10 mM) or BAPTA (10

mM) for 24 hours before infection. (D) Flow cytometry analysis of LAMP1

expression on the cell surface. (E) The histogram representing normalized MFI. (F)

Quantification of the lysosomal enzyme activity released into the culture medium.

Experiments were performed in triplicate. Data are presented as mean ± SEM. $*p <$

0.05, $**p < 0.01$. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate; MFI,

mean fluorescence intensity.

1016

Figure S8. Additional results on the role of ketogenesis in bacterial

neuroinvasion, related to Figure 6.

(A-B) The KE supplemental experiment was shown in Figure 6A-6D. (A) Flow

cytometry analysis of $CD36^+$ macrophages in mice supplemented with KE. (B)

Clinical scores of mice supplemented with KE.

(C-D) The ketogenic diet experiment was shown in Figure 6E-6M. Body weight

changes (C) and clinical scores (D) of mice during *L. monocytogenes* infection.

(E) Quantification of Evans blue leakage of mouse brains in Figure 6L.

(F-I) $HCAR2^{+/+}$ and $HCAR2^{-/-}$ mice were treated with ketogenic and control diets,

respectively, for 4 weeks, followed by oral infection with $2×10^8$ CFU *L.*

monocytogenes. (F) Survival of mice exposed to *L. monocytogenes* (n=10 for each

group). (G) Bacterial burdens in organs (brain, lung, spleen, liver). Cytokine levels in

plasma (H) and brain homogenate (I) (n=5 for each group).

(J-L) C57BL/6 mice were treated with ketogenic and control diets, respectively, for 4

weeks, followed by intravenous infection with $2×10^8$ CFU MRSA. (J) Survival of

mice exposed to MRSA (n=15 for each group). (K) Bacterial burdens in organs (brain,

lung, spleen, liver). (L) Clinical scores of mice during neuroinvasion (n=5 for each

group).

Experiments were performed in duplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant; i.p., intraperitoneally.

1037

Figure S9. Increased glucose levels alleviate bacterial neuroinvasion.

(A) Schematic diagram of *in vivo* glucose supplementation. C57BL/6 mice were

challenged with 2×10^8 CFU *L. monocytogenes* and then injected intraperitoneally

with glucose (100 mg/kg) every 12 hours.

(B-D) Serum glucose (B), BHB levels (C) and CD36$^+$ macrophages percentage (D)

were detected every 12 hours (n=5 for each group).

(E) Survival of mice exposed to *L. monocytogenes* with or without glucose

supplementation (n=10 for each group).

(F) Bacterial burdens in parenchymal organs of mice after 48 hours of infection.

(G-H) Cytokine levels in the plasma (G) and brain homogenate (H).

(I) Clinical scores of mice during L. monocytogenes infection.

Experiments were performed in duplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant.

1051

Figure S10. Additional results of high levels of BHB and human bacterial

meningitis, related to Figure 7.

(A) Risk factors of bacterial meningitis from the patients in the MIMIC IV database.

(B-E) Glucose (B) and insulin (C) levels in serum from the recruited HC and BM

patients. Correlation between serum glucose level and BHB (D), as well as serum

insulin levels and BHB (E) in BM patients. R, correlation coefficient

(F-H) Serum triglyceride (TC, F), high density lipoprotein (HDL-C, G) and low

density lipoprotein (LDL-C, H) from HC and BM patients.

Data are presented as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns,

nonsignificant, HC, health control, BM, bacterial meningitis.

1062

1063

Materials and methods

Clinical samples

Bacterial meningitis (BM) disease subjects (n = 62, 32 males and 30 females, aged

41.62 ±15.68 years) and healthy donors (n = 60, 30 males and 30 females, aged

38.32±9.17 years) were recruited from Chongqing Public Health Medical Center.

Isolation of PBMCs and single-cell suspensions was performed using PBMC Isolation

Kit (P8680, Solarbio) by density gradient centrifugation. Plasma was collected from

each patient and donor and then used for quantification of BHB. This study was

carried out in strict accordance with the Guidelines for the Care and Use of Animals

of Chongqing University, and was compliant with the "Guidance of the Ministry of

Science and Technology (MOST) for the Review and Approval of Human Genetic

Resources". Healthy individuals and patients providing blood samples were given

informed consent. The ethics committee approved this consent procedure. The clinical

symptoms of the BM patients included burnout, headache, irritability, vomiting, fever,

neck stiffness, convulsion, and consciousness disorder. Each symptom is worth 1

point.

The retrospective study was using Medical Information Mart for Intensive Care IV

(MIMIC-IV) version 1.0 data (https://physionet.org/content/mimiciv/1.0/), which

includes the medical information of ICU patients in a Boston hospital in the United

States over a 10-year study period. The establishment of the database was approved

by the Massachusetts Institute of Technology (Cambridge, MA) and the Institutional

Review Boards of Beth Israel Deaconess Medical Center (Boston, MA). Data were

extracted from the MIMIC-IV database using Structured Query Language (SQL) with

Navicat Premium (version 12.0.28). A total of 128 records of admission (49 patients /

79 controls) in the MIMIC IV database were included in this work. The patients with

benign intracranial hypertension were regarded as the control group.

Animal

All animal experiments were performed following institutional guidelines at

Chongqing University. In all experiments, control and experimental mice were 6-8-

week-old littermates of male C57BL/6 mice. Littermates of the same sex were housed

together regardless of genotype. All mice were maintained under a 12-hour light-dark

cycle with ad libitum access to regular food and water.

Bacterial strains

L. monocytogenes (19115) and *S. pneumoniae* 49619 were grown in Brain Heart

1100 Infusion Broth. Methicillin-resistant *Staphylococcus aureus* USA300 and *S.*

1101 *Typhimurium* were grown in Luria-Bertani Broth.

1102

1103 **Data and code availability**

1104 Single-cell RNA-seq data (accession number GEO: GSE264020, secure token for

1105 reviewer: stchicgorhozbkb) and RNA-seq data (accession number GEO: GSE273820,

1106 secure token for reviewer: gbohyqeanrmzhyn) have been deposited at GEO and are

1107 publicly available as of the date of publication.

1108

Single cell RNA-sequencing analysis

Isolated PBMCs from the mice were transported in in sterile culture dish with 10 ml

Dulbecco's Phosphate-Buffered Saline (DPBS, Thermo Fisher Scientific) on ice to

remove the residual tissue storage solution, then minced on ice. We used dissociation

enzyme 0.25% Trypsin (Thermo Fisher Scientific) and 10 ug/mL DNase I (Sigma)

dissolved in PBS with 5% Fetal Bovine Serum (FBS, GIBCO) to digest the tissues.

PBMCs were dissociated at 37 ℃ under a shaking speed of 50 rpm for about 40 min.

We repeatedly collected the dissociated cells at interval of 20 min to increase cell

yield and viability. Cell suspensions were filtered using a 40-μm nylon cell strainer

and red blood cells were removed by red blood cell lysis solution (Thermo Fisher

Scientific). Dissociated cells were washed with DPBS containing 2% FBS.

10× beads were then subject to second-strand cDNA synthesis, adaptor ligation, and

universal amplification. Sequencing libraries were prepared using randomly

interrupted whole-transcriptome amplification products to enrich the 3' end of the

transcripts linked with the cell barcode and UMI. All the remaining procedures

including the library construction were performed according to the standard

manufacturer's protocol (Chromium Single Cell 3' v3). Sequencing libraries were

quantified using a High Sensitivity DNA Chip (Agilent) on a Bioanalyzer 2100 and

the Qubit High Sensitivity DNA Assay (Thermo Fisher Scientific). The libraries were

sequenced on NovaSeq6000 (Illumina) using 2×150 chemistry. The sequencing and

bioinformatic analysis were performed on platform of Majorbio Co., Ltd (Shanghai,

China).

Reads were processed using the Cell Ranger (v7.1.0) pipeline with default and

recommended parameters. FASTQs generated from Illumina sequencing output were

aligned to the mouse genome, version GRCm38, using the STAR algorithm. Gene-

barcode matrix containing the barcoded cells and gene expression counts were

generated. This output was then imported into the Seurat (v3.2.0) R toolkit for quality

control and downstream analysis of our single cell RNAseq data.

Cells were clustered using graph-based clustering of the PCA reduced data with the

Louvain Method after computing a shared nearest neighbor graph. For sub-clustering,

we applied the same procedure of scaled, dimensionality reduction, and clustering to

the specific set of data. For each cluster, we used the Wilcoxon Rank-Sum Test to find

significant deferentially expressed genes comparing to the remaining clusters.

To identify differential expression genes (DEGs) between two different samples or

clusters, was performed using the function FindMarkers in Seurat, using a likelihood

ratio test. Essentially, DEGs with $|log2FC| > 0.25$ and Q value $\leqslant 0.05$ were

significantly different expressed genes. In addition, functional-enrichment analysis

GO were performed to identify which DEGs were significantly enriched in GO terms

and metabolic pathways at Bonferroni-corrected P-value $\leqslant 0.05$ compared with the

whole-transcriptome background. GO functional enrichment analysis were carried out

by Goatools (https://github.com/tanghaibao/Goatools).

1150

Mouse model of acute neuroinvasion

Mice were challenged with pathogenic bacteria (*L. monocytogenes* and MRSA, 2×10^8

CFU; *S. Typhimurium* and *S. pneumonia*, 1×10^8 CFU) to construct a model of acute

bacterial meningitis. Mice challenged with LPS (2 mg/mouse) were used as positive

controls for BBB damage. The progression of acute bacterial meningitis was

determined by clinical scores as previously described with a few adaptations.[59]

Clinical scoring was performed based on the Table S2, including activity (0–2), time

to return to upright position (0–2), eyes (0–2) and neurological examination (0–2).

Mice were deeply anaesthetized by an intraperitoneal injection of tribromoethanol

solution (Avertin, 500 mg/kg) and transcranial perfused with saline solution for brain

tissue pathomorphology. Blood was sampled immediately before perfusion. Tissue

samples, including brain, lung, liver, and spleen, were dissected and weighed.

1163

Analysis of bacterial load in cell or tissue

Tissue samples of mice were weighted and then transferred into 2 ml tubes containing

4 mm stainless steel beads (Qiagen) and 1 ml of ice-cold sterile saline.

Homogenization was carried out in TissueLyser II (Qiagen) for 5 min at 70 Hz. Serial

dilutions of the homogenates were plated on the plates incubated at 37℃ and CFU

were counted.

The infected macrophages were incubated in a fresh solution containing 5 μg/mL

lysostaphin for the indicated periods and then lysed by 0.2% Triton to release

intracellular bacteria. The lysates were diluted to appropriate dilutions with sterile

PBS for CFU count on agar plates.

1174

BBB permeability assays

After infection, mice were retro-orbitally injected with 2% Evans blue. After 2 h of

circulation, mice were deeply anaesthetized, and the brains were delicately dissected

after transcardial PBS perfusion. Whole-brain images were acquired using a digital

camera. To quantify Evans blue leakage, mouse brains were longitudinally cut,

weighted and then homogenized in 0.5 ml of 50% trichloroacetic acid. The lysates

were centrifuged at 15,000 g for 30 min. The amount of extravasated Evans blue in

the supernatants was measured at a Multi-Mode Microplate Reader (SpetraMax i3x)

at 620/680 nm emission, calculated by a standard curve, and normalized to the weight

of each brain.

1185

Brain histopathology and Immunostaining

Brain samples were collected from infected mice 24 hours after injection. Mice were

deeply anaesthetized with tribromoethanol solution (Avertin, 500 mg/kg,

intravenously) and transcranial perfused with 30 ml saline solution followed by 10 ml

of 4% Paraformaldehyde (PFA) in saline. After dissection, brain samples were

maintained in 4% PFA at 4 ℃. Fixed brains were embedded in paraffin, sectioned and

half of the slides were stained using haematoxylin and eosin. Each field was assigned

a score from 0 to 3 based on neuronal morphology: grade 0 (not altered); grade 1 (no

vacuolation with small numbers of pyknotic cells); grade 2 (moderate vacuolation and

pyknosis); and grade 3 (extensive vacuolation, pyknosis and tissue loss or liquefactive

necrosis).

Fresh frozen brain tissue sections of 5-10 mm thickness were mounted on slides, air

dried for 10 minutes at room temperature and fixed in ice cold 10% neutral-buffered

formalin for 5-10 min. Slices were washed in PBS, then incubated for 2 hours in 3%

bovine serum albumin (BSA) and then 0.1% Triton X-100 in PBS. The slides were

labelled with the appropriate primary antibodies of CD11b and CD36 overnight at

4 ℃ in mild blocking conditions. Next day, the slices were washed in PBS, and then

incubated with secondary antibodies and DAPI. Tissues were washed in PBS and then

mounted on glass slides under coverslips in mounting medium. The slides were left in

darkness overnight before observation under a Zeiss LSM900 or LM710 microscope.

1206

Cell sorting and flow cytometry.

Isolated mouse PBMCs were cleaned twice with PBS. After calculating and

determining the appropriate concentration, appropriate amount of fluorescent-coupled

primary antibody (F4/80-APC, CD11b-FITC, as the marker of macrophage, CD36-

FITC) was added respectively. Incubate at 4 ℃ for 30 min away from light. The

living macrophages were collected according to the expression of CD36 and then

culture in vitro overnight. The sorting cells were detected the lipid content using

BODIPY FL and oil Red O. To evaluate the ability of lipid uptake, macrophages were

incubated with 40 μg/ml DiI-ox-LDL particles for 6h and then detect the fluorescence

intensity by flow cytometry.

To determine the proportion of CD36$^+$ macrophages population, cells were stained

with fluorescent-coupled primary antibody (F4/80-APC, CD11b-PE, CD36-FITC) at

4 ℃ for 30 min away from light. To detect the expression of LAMP1 on the cell

surface, infected macrophages were cleaned twice with PBS and then stained with

LAMP1-FITC at 4 ℃ for 30 min away from light. The cells were washed three times

and then re-suspended in 200μl of the staining buffer for final FACS analysis. Data

was analyzed with FlowJo software.

1224

Isolation and culture of primary cells

To generate peritoneal macrophages, 8-week-old mice were injected intraperitoneally

with 3% thioglycolate broth. After 72 hours, peritoneal cells were harvested, and

macrophages were enriched by quick adherence.

Primary endothelial cells were isolated from 8 to 12-week-old male C57BL/6 mice.

The cerebral cortex in the brains were separated and rolled on the filter paper to

remove the meninges and choroid plexuses. A mechanical reduction was performed

by mincing the brain pieces using surgical blades. The homogenate was centrifuged

(800 g, 8 min, 4℃) and the precipitate was digested in DMEM containing 1 mg/ml

type II collagenase and 58.5 U/ml type I DNase for 75 min on a benchtop orbital

shaker at 37 ℃. The supernatant was finally collected and centrifuged at 500 g for 8

min. The isolated cerebrovascular endothelial cells were purified by the anchoring

technique. The liver and lung from an anaesthetized mouse were perfused via the

portal vein sequentially with balanced salt solution containing 5 mM glucose, 0.01%

sodium heparin, and 5 mM EGTA and high glucose DMEM medium containing 4

mM $CaCl_2$, 2% FBS and 0.05% collagenase IV (Sigma-Aldrich, St. Louis, MO) at a

flow rate of 5 ml/min for 5 min, respectively. The following steps are similar in order

to separate the liver sinusoids endothelial cells and pulmonary vascular endothelial

cells.

Isolation of primary embryonic neurons and single-cell suspensions was performed

using Pierce Primary Neuron Isolation Kit (88280, Thermo Scientific) by density

gradient centrifugation.

1247

***In vitro* blood-brain barrier**

The *in vitro* BBB model was constructed with primary endothelial cells using a

Transwell cell culture system. Briefly, CVECs were seeded onto the upper chamber of

the Transwell pre-coated with gelatine, and cultured with the medium containing 10%

FBS. The integrity of the monolayer cell was evaluated by measuring the TEER

values using a Millicell-ERS voltohmmeter (Millipore). The cell monolayers with

TEER values higher than 300 $\Omega.cm^2$ were used as the BBB model for the

transmigration studies. The neurons cells were seeded onto the lower chamber of the

Transwell for bacterial load.

Macrophages were exposed to *L. monocytogenes* (MOI=10) for 6 hours. After the

BBB model has been constructed, the infected macrophages were digested for cell

counting and added into the upper chamber. The conventional incubation time was 12

hours and then the neurons cells in the bottom chamber were lysed for penetrating

bacterial quantification.

1262

Adoptive macrophages transfer in mouse bacteria brain neuroinvasion

Peritoneal macrophages from donor mice were collected via intraperitoneal injection

of thioglycolate. Following further cultivation, purified macrophages were obtained.

To collect CD36$^+$ macrophages, purified macrophages were stimulated with 80 μg/ml

ox-LDL over 24 hours. Recipient mice were pretreated with 3 days of CTX and then

intraperitoneally given 0.2 ml normal saline or cell suspension (1×10^6 per mouse) on

Day 0. After adoptive macrophages transfer, recipient mice were intraperitoneally

injected with 2×10^8 CFU of *L. monocytogenes* in 12 hours and euthanized after 18

hours.

Purified peritoneal macrophages were harvested from PCSK9$^{+/+}$ and PCSK9$^{-/-}$ donor

mice respectively, and then transferred to recipient mice as depicted in Figure 2J.

Recipient mice were infected with 2×10^8 CFU *L. monocytogenes* 12 hours after

macrophages transfer and euthanized after 24 hours.

1276

Live-cell flow chamber assay

1277

A flow cell system was used to test macrophages adhesion, shear resistance, crawling,

and transmigration, on endothelial monolayer, which was performed as previously

described.[43] Shear flow was applied by a syringe pump (PHD22/2000, Harvard

Apparatus) into a microfluidic chip with a 100 μm × 1 mm× 15 mm main flowing

zone. Different shear stresses were acquired by setting corresponding volumetric flow

rates. Isolated macrophages were stimulated under different concentration of ox-LDL

for 24 hours and then infected with *L. monocytogenes* (MOI=10) for 6 hours.

Macrophages were re-suspended and stained with Hoechst. The cells were then

perfused over pre-formed endothelial monolayer for 8 min at a shear stress of 0.1

dyne/cm^2. Then, the adhesion and crawling dynamics of macrophages on endothelial

monolayer was recorded at 2 dyn/cm^2 for 10 min at 20× objective by a CCD camera.

Images were taken from the video every second and analyzed the crawling parameters

and transmigrating duration with Image J software. Here a macrophage crossing the

monolayer with a 70% fluorescence intensity reduction was deemed transmigrated.

For testing shear-resistant macrophage adhesion, primed macrophages were re-

suspended and counted. The same number of cells ($2×10^4$) were completely perfused

over pre-formed endothelial monolayer within 10 min. The shear stress was increased

stepwise from 1 to 2, 4, 8, 16, 32 dyn/cm^2 for 30s each. Remaining adherent

macrophages on endothelial monolayer were recorded at 20× objective by a CCD

camera and counted using Image J software. The normalized adhesion fraction was

calculated by dividing the number of adherent macrophages at the endpoint of each

shear phase by the number at the endpoint of 1 dyn/cm^2. For defining the macrophage

transmigration dynamics, the time-lapse images were analyzed offline frame-by-frame

every second.

1302

Quantification of *L. monocytogenes* protrusion formation

Isolated peritoneal macrophages were stimulated under different concentration of ox-

LDL for 24 hours and then infected with CFSE-labeled *L. monocytogenes* (MOI=10)

for 6 hours. At 4 hours post-infection, cells were fixed, permeabilized, and labeled

with mouse anti-ezrin and phalloidin-Alexa 647. Samples were analyzed using

confocal microscopy. Protrusions were identified as bacteria with ezrin tails that

project from host cells into neighboring. In each experiment, ten cells were scored for

each condition. The lengths of ezrin comet tails were measured using ImageJ

software.

1312

Lipid metabolomics

Metabolite extraction and mass spectrometry detection was processed using Ultra-

High Performance Liquid Chromatography-mass spectrometry (UHPLC-MS) of

Biomarker Technologies (Beijing, China). The raw data collected using MassLynx

V4.2 is processed by Progenesis QI software for peak extraction, peak alignment, and

other data processing operations, based on the Progenesis QI software online

METLIN database and Biomark's self-built library for identification. The theoretical

fragment identification and mass deviation are within 100 ppm. After normalizing the

original peak area information with the total peak area, the follow-up analysis was

performed. A total of 4,625 named metabolites were retained for datasets. The R

language package ropls was used to perform OPLS-DA modeling, and 200 times

permutation tests was performed to verify the reliability of the model.

1325

RNA-seq (RNA Sequencing)

RNA was extracted from mouse liver tissues using a Trizol extraction and Purelink

RNA kit (ThermoFisher Scientific). Bioanalyzer chips were used to assess RNA

quality prior to RNA sequencing. A total amount of 1 μg RNA per sample was used as

input material for the RNA sample preparations. Sequencing libraries were generated

using Hieff NGS Ultima Dual-mode mRNA Library Prep Kit for lumina following

manufacturer's recommendations and index codes were added to attribute sequences

to each sample. The libraries were sequenced on an Illumina NovaSeq platform to

generate 150 bp paired-end reads, according to the manufacturer's instructions.

1335

Measurement of ECAR

ECAR was measured using the XF Cell Mito Stress Test kit (103020-100, Seahorse

Biosciences) and XF Glycolysis Stress Test kit (103020-100, Seahorse Biosciences),

respectively. In brief, liver tissue were seeded into XFe24 Cell Culture Microplates

and cultured in a 37 °C incubator containing 5% CO_2 overnight. Three injections (10

mM glucose, 1.5 μM oligomycin and 50 mM 2-DG) were performed at the indicated

time. ECAR values were normalized to the total amount of proteins. ECAR

parameters were calculated using Wave software (v.2.4.3, Seahorse Bioscience,

Agilent Technologies) according to manufacturer instructions.

1345

Fura-4 Ca^{2+} imaging

Cells were removed from incubator and washed with PBS and then incubated with 5

μM Fluo-4 AM (Beyotime) for 60 min at 37 ℃ with 0.04% Pluronic-127. After

incubation, cells were washed with PBS and incubated with PBS for 30 min at 37 ℃.

Fluorescence was detected with a CytoFLEX Flow Cytometer (Beckman Coulter)

equipped with 488 nm excitation wavelength. As indicated, BHB (10 mM) were

added into the cells pre-loaded with Fluo-4 30 s after the record. Data was analyzed

with Flow Jo to determine the fluorescence intensity.

1354

Diets for *in vivo* assays

When indicated, mice were administered a supplement of 20 mg/ml ketone ester (D-

β-hydroxybutyrate-(R)-1,3 butanediol monoester) in the drinking water for 4 weeks

before the infection. Ketogenic diets (# XTKD01, Xietong Pharma Co.) contained

10% protein and 90% fat while the control diets contained 23.2% protein, 2.8% fat

and 74% sugar. Mice were consequently placed on either the ketogenic or control diet

for an additional 4 weeks before infection.

For the glucose supplementation experiment, mice were injected intraperitoneally

with saline or glucose (100 mg/kg) every 12 hours after infection. For blood glucose

level measurement, a drop of blood was collected by snipping the very end of the tail,

and the blood in the strip was analysed using the glucometer to measure the blood

glucose levels.

1367

Real-time qRT-PCR

Total RNA was isolated using an RNeasy Mini Kit (QIAGEN). The mRNA was

reverse-transcribed and quantified using a HiScript II 1st Strand cDNA Synthesis Kit

(Vazyme). qRT-PCR was performed on an ABI StepOnePlus PCR system (Thermo

Fisher Scientific), and results were normalized to GAPDH mRNA levels. Data were

analyzed using the 2-DDct method.

1374

Statistical analysis

Statistical analysis was performed using GraphPad Prism Software. Two-group

comparisons were made using two-tailed unpaired Student's t-test. One-way analysis

of variance (ANOVA) with appropriate multiple comparisons tests was used to

compare three independent groups. For comparisons of multiple factors, two-way

ANOVA with appropriate multiple comparisons tests was used. Exact p values for

specific group comparisons in all the figures. A value of $p < 0.05$ was considered

significant. Specific details of statistical analysis are presented above or in associated

figure legends.

Principal component analysis (PCA) and Spearman correlation analysis were used to

judge the repeatability of the samples within group and the quantity control samples.

Differential expression was tested for normalized count data using the R package edge

R, with a generalized linear model. Genes with $|\log2 (\text{fold change})| \geqslant 1$ and $P \leqslant$

0.05 were identified as differentially expressed genes. The identified compounds are

searched for classification and pathway information in KEGG and HMDB databases.

The difference genes of KEGG pathway enrichment significance were calculated

using hypergeometric distribution test.

1392

1393

1394

Reference

1.   Doran, K.S., Fulde, M., Gratz, N., Kim, B.J., Nau, R., Prasadarao, N.,

Schubert-Unkmeir, A., Tuomanen, E.I., and Valentin-Weigand, P. (2016).

Host-pathogen interactions in bacterial meningitis. Acta Neuropathol *131*,

185-209. 10.1007/s00401-015-1531-z.

2.   Hasbun, R. (2022). Progress and Challenges in Bacterial Meningitis: A

Review. JAMA *328*, 2147-2154. 10.1001/jama.2022.20521.

3.   Langen, U.H., Ayloo, S., and Gu, C. (2019). Development and Cell Biology of

the Blood-Brain Barrier. Annu Rev Cell Dev Biol *35*, 591-613.

10.1146/annurev-cellbio-100617-062608.

4.   Forrester, J.V., McMenamin, P.G., and Dando, S.J. (2018). CNS infection and

immune privilege. Nat Rev Neurosci *19*, 655-671. 10.1038/s41583-018-0070-

8.

1408 5. Loh, L.N., Gao, G., and Tuomanen, E.I. (2017). Dissecting Bacterial Cell Wall

1409   Entry and Signaling in Eukaryotic Cells: an Actin-Dependent Pathway

1410   Parallels Platelet-Activating Factor Receptor-Mediated Endocytosis. mBio *8*.

1411   10.1128/mBio.02030-16.

1412 6. Tabusi, M., Thorsdottir, S., Lysandrou, M., Narciso, A.R., Minoia, M.,

1413   Srambickal, C.V., Widengren, J., Henriques-Normark, B., and Iovino, F.

1414   (2021). Neuronal death in pneumococcal meningitis is triggered by

1415   pneumolysin and RrgA interactions with beta-actin. PLoS Pathog *17*,

1416   e1009432. 10.1371/journal.ppat.1009432.

1417 7. Coureuil, M., Lecuyer, H., Bourdoulous, S., and Nassif, X. (2017). A journey

1418   into the brain: insight into how bacterial pathogens cross blood-brain barriers.

1419   Nat Rev Microbiol *15*, 149-159. 10.1038/nrmicro.2016.178.

1420 8. Coureuil, M., Lecuyer, H., Scott, M.G., Boularan, C., Enslen, H., Soyer, M.,

1421   Mikaty, G., Bourdoulous, S., Nassif, X., and Marullo, S. (2010).

1422   Meningococcus Hijacks a beta2-adrenoceptor/beta-Arrestin pathway to cross

1423   brain microvasculature endothelium. Cell *143*, 1149-1160.

1424   10.1016/j.cell.2010.11.035.

1425 9. Huynh, J., Donovan, J., Phu, N.H., Nghia, H.D.T., Thuong, N.T.T., and

1426   Thwaites, G.E. (2022). Tuberculous meningitis: progress and remaining

1427   questions. Lancet Neurol *21*, 450-464. 10.1016/S1474-4422(21)00435-X.

1428 10. Drevets, D.A., Dillon, M.J., Schawang, J.S., Van Rooijen, N., Ehrchen, J.,

1429   Sunderkotter, C., and Leenen, P.J. (2004). The Ly-6Chigh monocyte

subpopulation transports Listeria monocytogenes into the brain during

systemic infection of mice. J Immunol *172*, 4418-4424.

10.4049/jimmunol.172.7.4418.

11.   Levin, B.R., Baquero, F., Ankomah, P.P., and McCall, I.C. (2017). Phagocytes,

Antibiotics, and Self-Limiting Bacterial Infections. Trends Microbiol *25*, 878-

892. 10.1016/j.tim.2017.07.005.

12.   Boada-Romero, E., Martinez, J., Heckmann, B.L., and Green, D.R. (2020).

The clearance of dead cells by efferocytosis. Nat Rev Mol Cell Biol *21*, 398-

414. 10.1038/s41580-020-0232-1.

13.   Locati, M., Curtale, G., and Mantovani, A. (2020). Diversity, Mechanisms,

and Significance of Macrophage Plasticity. Annu Rev Pathol *15*, 123-147.

10.1146/annurev-pathmechdis-012418-012718.

14.   Zhai, L., Huang, C., Ning, Z., Zhang, Y., Zhuang, M., Yang, W., Wang, X.,

Wang, J., Zhang, L., Xiao, H., et al. (2023). Ruminococcus gnavus plays a

pathogenic role in diarrhea-predominant irritable bowel syndrome by

increasing serotonin biosynthesis. Cell Host Microbe *31*, 33-44 e35.

10.1016/j.chom.2022.11.006.

15.   Van der Merwe, S., Chokshi, S., Bernsmeier, C., and Albillos, A. (2021). The

multifactorial mechanisms of bacterial infection in decompensated cirrhosis. J

Hepatol *75 Suppl 1*, S82-S100. 10.1016/j.jhep.2020.11.029.

16.   Maudet, C., Kheloufi, M., Levallois, S., Gaillard, J., Huang, L., Gaultier, C.,

Tsai, Y.H., Disson, O., and Lecuit, M. (2022). Bacterial inhibition of Fas-

1452 mediated killing promotes neuroinvasion and persistence. Nature *603*, 900-

1453 906. 10.1038/s41586-022-04505-7.

1454 17. Xu, K., Yin, N., Peng, M., Stamatiades, E.G., Shyu, A., Li, P., Zhang, X., Do,

1455 M.H., Wang, Z., Capistrano, K.J., et al. (2021). Glycolysis fuels

1456 phosphoinositide 3-kinase signaling to bolster T cell immunity. Science *371*,

1457 405-410. 10.1126/science.abb2683.

1458 18. Escoll, P., Song, O.R., Viana, F., Steiner, B., Lagache, T., Olivo-Marin, J.C.,

1459 Impens, F., Brodin, P., Hilbi, H., and Buchrieser, C. (2017). Legionella

1460 pneumophila Modulates Mitochondrial Dynamics to Trigger Metabolic

1461 Repurposing of Infected Macrophages. Cell Host Microbe *22*, 302-316 e307.

1462 10.1016/j.chom.2017.07.020.

1463 19. Yang, D., Jacobson, A., Meerschaert, K.A., Sifakis, J.J., Wu, M., Chen, X.,

1464 Yang, T., Zhou, Y., Anekal, P.V., Rucker, R.A., et al. (2022). Nociceptor

1465 neurons direct goblet cells via a CGRP-RAMP1 axis to drive mucus

1466 production and gut barrier protection. Cell *185*, 4190-4205 e4125.

1467 10.1016/j.cell.2022.09.024.

1468 20. Loft, A., Schmidt, S.F., Caratti, G., Stifel, U., Havelund, J., Sekar, R., Kwon,

1469 Y., Sulaj, A., Chow, K.K., Alfaro, A.J., et al. (2022). A macrophage-hepatocyte

1470 glucocorticoid receptor axis coordinates fasting ketogenesis. Cell Metab *34*,

1471 473-486 e479. 10.1016/j.cmet.2022.01.004.

1472 21. Lu, J.F., Zhu, M.Q., Xia, B., Zhang, N.N., Liu, X.P., Liu, H., Zhang, R.X.,

1473 Xiao, J.Y., Yang, H., Zhang, Y.Q., et al. (2023). GDF15 is a major determinant

of ketogenic diet-induced weight loss. Cell Metab *35*, 2165-2182 e2167.

10.1016/j.cmet.2023.11.003.

22.   Tomlinson, K.L., Chen, Y.T., Junker, A., Urso, A., Wong Fok Lung, T., Ahn,

D., Hofstaedter, C.E., Baskota, S.U., Ernst, R.K., Prince, A., and Riquelme,

S.A. (2023). Ketogenesis promotes tolerance to Pseudomonas aeruginosa

pulmonary infection. Cell Metab *35*, 1767-1781 e1766.

10.1016/j.cmet.2023.09.001.

23.   Goldberg, E.L., Molony, R.D., Kudo, E., Sidorov, S., Kong, Y., Dixit, V.D.,

and Iwasaki, A. (2019). Ketogenic diet activates protective gammadelta T cell

responses against influenza virus infection. Sci Immunol *4*.

10.1126/sciimmunol.aav2026.

24.   Karagiannis, F., Peukert, K., Surace, L., Michla, M., Nikolka, F., Fox, M.,

Weiss, P., Feuerborn, C., Maier, P., Schulz, S., et al. (2022). Impaired

ketogenesis ties metabolism to T cell dysfunction in COVID-19. Nature *609*,

801-807. 10.1038/s41586-022-05128-8.

25.   Seufert, A.L., Hickman, J.W., Traxler, S.K., Peterson, R.M., Waugh, T.A.,

Lashley, S.J., Shulzhenko, N., Napier, R.J., and Napier, B.A. (2022). Enriched

dietary saturated fatty acids induce trained immunity via ceramide production

that enhances severity of endotoxemia and clearance of infection. Elife *11*.

10.7554/eLife.76744.

26.   Wilhelm, C., Surendar, J., and Karagiannis, F. (2021). Enemy or ally? Fasting

as an essential regulator of immune responses. Trends Immunol *42*, 389-400.

1496 10.1016/j.it.2021.03.007.

27. Singh, V., Jamwal, S., Jain, R., Verma, P., Gokhale, R., and Rao, K.V. (2012). Mycobacterium tuberculosis-driven targeted recalibration of macrophage lipid homeostasis promotes the foamy phenotype. Cell Host Microbe *12*, 669-681. 10.1016/j.chom.2012.09.012.

28. McFarland, A.P., Yalin, A., Wang, S.Y., Cortez, V.S., Landsberger, T., Sudan, R., Peng, V., Miller, H.L., Ricci, B., David, E., et al. (2021). Multi-tissue single-cell analysis deconstructs the complex programs of mouse natural killer and type 1 innate lymphoid cells in tissues and circulation. Immunity *54*, 1320-1337 e1324. 10.1016/j.immuni.2021.03.024.

29. Ahmad, F., Rani, A., Alam, A., Zarin, S., Pandey, S., Singh, H., Hasnain, S.E., and Ehtesham, N.Z. (2022). Macrophage: A Cell With Many Faces and Functions in Tuberculosis. Front Immunol *13*, 747799. 10.3389/fimmu.2022.747799.

30. Hill, D.A., Lim, H.W., Kim, Y.H., Ho, W.Y., Foong, Y.H., Nelson, V.L., Nguyen, H.C.B., Chegireddy, K., Kim, J., Habertheuer, A., et al. (2018). Distinct macrophage populations direct inflammatory versus physiological changes in adipose tissue. Proc Natl Acad Sci U S A *115*, E5096-E5105. 10.1073/pnas.1802611115.

31. Jaitin, D.A., Adlung, L., Thaiss, C.A., Weiner, A., Li, B., Descamps, H., Lundgren, P., Bleriot, C., Liu, Z., Deczkowska, A., et al. (2019). Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-

Dependent Manner. Cell *178*, 686-698 e614. 10.1016/j.cell.2019.05.054.

32.   Luis, G., Godfroid, A., Nishiumi, S., Cimino, J., Blacher, S., Maquoi, E.,

Wery, C., Collignon, A., Longuespee, R., Montero-Ruiz, L., et al. (2021).

Tumor resistance to ferroptosis driven by Stearoyl-CoA Desaturase-1 (SCD1)

in cancer cells and Fatty Acid Biding Protein-4 (FABP4) in tumor

microenvironment promote tumor recurrence. Redox Biol *43*, 102006.

10.1016/j.redox.2021.102006.

33.   Victor, M.B., Leary, N., Luna, X., Meharena, H.S., Scannail, A.N., Bozzelli,

P.L., Samaan, G., Murdock, M.H., von Maydell, D., Effenberger, A.H., et al.

(2022). Lipid accumulation induced by APOE4 impairs microglial

surveillance of neuronal-network activity. Cell Stem Cell *29*, 1197-1212

e1198. 10.1016/j.stem.2022.07.005.

34.   Chen, Y., Zhang, J., Cui, W., and Silverstein, R.L. (2022). CD36, a signaling

receptor and fatty acid transporter that regulates immune cell metabolism and

fate. J Exp Med *219*. 10.1084/jem.20211314.

35.   Botham, K.M., and Wheeler-Jones, C.P. (2013). Postprandial lipoproteins and

the molecular regulation of vascular homeostasis. Prog Lipid Res *52*, 446-464.

10.1016/j.plipres.2013.06.001.

36.   Jaggi, U., Matundan, H.H., Yu, J., Hirose, S., Mueller, M., Wormley, F.L., Jr.,

and Ghiasi, H. (2021). Essential role of M1 macrophages in blocking cytokine

storm and pathology associated with murine HSV-1 infection. PLoS Pathog

17, e1009999. 10.1371/journal.ppat.1009999.

37. Pasula, R., Martin, W.J., 2nd, Kesavalu, B.R., Abdalla, M.Y., and Britigan,

B.E. (2017). Passive transfer of interferon-gamma over-expressing

macrophages enhances resistance of SCID mice to Mycobacterium

tuberculosis infection. Cytokine *95*, 70-79. 10.1016/j.cyto.2017.02.009.

38. Peyrusson, F., Varet, H., Nguyen, T.K., Legendre, R., Sismeiro, O., Coppee,

J.Y., Wolz, C., Tenson, T., and Van Bambeke, F. (2020). Intracellular

Staphylococcus aureus persisters upon antibiotic exposure. Nat Commun *11*,

2200. 10.1038/s41467-020-15966-7.

39. Ding, Z., Pothineni, N.V.K., Goel, A., Luscher, T.F., and Mehta, J.L. (2020).

PCSK9 and inflammation: role of shear stress, pro-inflammatory cytokines,

and LOX-1. Cardiovasc Res *116*, 908-915. 10.1093/cvr/cvz313.

40. Da Dalt, L., Castiglioni, L., Baragetti, A., Audano, M., Svecla, M., Bonacina,

F., Pedretti, S., Uboldi, P., Benzoni, P., Giannetti, F., et al. (2021). PCSK9

deficiency rewires heart metabolism and drives heart failure with preserved

ejection fraction. Eur Heart J *42*, 3078-3090. 10.1093/eurheartj/ehab431.

41. Katsuki, S., P, K.J., Lupieri, A., Nakano, T., Passos, L.S.A., Rogers, M.A.,

Becker-Greene, D., Le, T.D., Decano, J.L., Ho Lee, L., et al. (2022).

Proprotein Convertase Subtilisin/Kexin 9 (PCSK9) Promotes Macrophage

Activation via LDL Receptor-Independent Mechanisms. Circ Res *131*, 873-

889. 10.1161/CIRCRESAHA.121.320056.

42. Qi, Z., Hu, L., Zhang, J., Yang, W., Liu, X., Jia, D., Yao, Z., Chang, L., Pan,

G., Zhong, H., et al. (2021). PCSK9 (Proprotein Convertase Subtilisin/Kexin

9) Enhances Platelet Activation, Thrombosis, and Myocardial Infarct

Expansion by Binding to Platelet CD36. Circulation *143*, 45-61.

10.1161/CIRCULATIONAHA.120.046290.

43. Yang, H., Li, N., Du, Y., Tong, C., Lu, S., Hu, J., Zhang, Y., and Long, M.

(2017). Neutrophil adhesion and crawling dynamics on liver sinusoidal

endothelial cells under shear flow. Exp Cell Res *351*, 91-99.

10.1016/j.yexcr.2017.01.002.

44. Neupane, A.S., Willson, M., Chojnacki, A.K., Vargas, E.S.C.F., Morehouse,

C., Carestia, A., Keller, A.E., Peiseler, M., DiGiandomenico, A., Kelly, M.M.,

et al. (2020). Patrolling Alveolar Macrophages Conceal Bacteria from the

Immune System to Maintain Homeostasis. Cell *183*, 110-125 e111.

10.1016/j.cell.2020.08.020.

45. Hernandez, G.E., and Iruela-Arispe, M.L. (2020). The many flavors of

monocyte/macrophage--endothelial cell interactions. Curr Opin Hematol *27*,

181-189. 10.1097/MOH.0000000000000573.

46. Pang, X., He, X., Qiu, Z., Zhang, H., Xie, R., Liu, Z., Gu, Y., Zhao, N., Xiang,

Q., and Cui, Y. (2023). Targeting integrin pathways: mechanisms and advances

in therapy. Signal Transduct Target Ther *8*, 1. 10.1038/s41392-022-01259-6.

47. Deng, L., Kersten, S., and Stienstra, R. (2023). Triacylglycerol uptake and

handling by macrophages: From fatty acids to lipoproteins. Prog Lipid Res *92*,

101250. 10.1016/j.plipres.2023.101250.

48. Patterson, M.T., Firulyova, M.M., Xu, Y., Hillman, H., Bishop, C., Zhu, A.,

Hickok, G.H., Schrank, P.R., Ronayne, C.E., Caillot, Z., et al. (2023). Trem2

promotes foamy macrophage lipid uptake and survival in atherosclerosis.

Nature Cardiovascular Research *2*, 1015-1031. 10.1038/s44161-023-00354-3.

49.   Andrews, N.W., and Corrotte, M. (2018). Plasma membrane repair. Curr Biol

28, R392-R397. 10.1016/j.cub.2017.12.034.

50.   Jimenez, A.J., Maiuri, P., Lafaurie-Janvore, J., Divoux, S., Piel, M., and Perez,

F. (2014). ESCRT machinery is required for plasma membrane repair. Science

343, 1247136. 10.1126/science.1247136.

51.   Newman, J.C., and Verdin, E. (2017). beta-Hydroxybutyrate: A Signaling

Metabolite. Annu Rev Nutr *37*, 51-76. 10.1146/annurev-nutr-071816-064916.

52.   Youm, Y.H., Nguyen, K.Y., Grant, R.W., Goldberg, E.L., Bodogai, M., Kim,

D., D'Agostino, D., Planavsky, N., Lupfer, C., Kanneganti, T.D., et al. (2015).

The ketone metabolite beta-hydroxybutyrate blocks NLRP3 inflammasome-

mediated inflammatory disease. Nat Med *21*, 263-269. 10.1038/nm.3804.

53.   Duking, T., Spieth, L., Berghoff, S.A., Piepkorn, L., Schmidke, A.M.,

Mitkovski, M., Kannaiyan, N., Hosang, L., Scholz, P., Shaib, A.H., et al.

(2022). Ketogenic diet uncovers differential metabolic plasticity of brain cells.

Sci Adv *8*, eabo7639. 10.1126/sciadv.abo7639.

54.   Matsuura, T.R., Puchalska, P., Crawford, P.A., and Kelly, D.P. (2023). Ketones

and the Heart: Metabolic Principles and Therapeutic Implications. Circ Res

132, 882-898. 10.1161/CIRCRESAHA.123.321872.

55.   Shah, U.A., and Iyengar, N.M. (2022). Plant-Based and Ketogenic Diets As

Diverging Paths to Address Cancer: A Review. JAMA Oncol *8*, 1201-1208.

10.1001/jamaoncol.2022.1769.

56. Steeb, B., Claudi, B., Burton, N.A., Tienz, P., Schmidt, A., Farhan, H., Maze,

A., and Bumann, D. (2013). Parallel exploitation of diverse host nutrients

enhances Salmonella virulence. PLoS Pathog *9*, e1003301.

10.1371/journal.ppat.1003301.

57. Peyron, P., Vaubourgeix, J., Poquet, Y., Levillain, F., Botanch, C., Bardou, F.,

Daffe, M., Emile, J.F., Marchou, B., Cardona, P.J., et al. (2008). Foamy

macrophages from tuberculous patients' granulomas constitute a nutrient-rich

reservoir for M. tuberculosis persistence. PLoS Pathog *4*, e1000204.

10.1371/journal.ppat.1000204.

58. Yang, Y., Fang, H., Xie, Z., Ren, F., Yan, L., Zhang, M., Xu, G., Song, Z.,

Chen, Z., Sun, W., et al. (2024). Yersinia infection induces glucose depletion

and AMPK-dependent inhibition of pyroptosis in mice. Nat Microbiol.

10.1038/s41564-024-01734-6.

59. Koopmans, M.M., Engelen-Lee, J., Brouwer, M.C., Jaspers, V., Man, W.K.,

Vall Seron, M., and van de Beek, D. (2018). Characterization of a Listeria

monocytogenes meningitis mouse model. J Neuroinflammation *15*, 257.

10.1186/s12974-018-1293-3.

1625

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

A Upper Chamber

B CHMP4B-EGFP

C

E

D

F

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9