Title: Dysregulation of macrophage lipid metabolism underlies intracellular bacterial

neuroinvasion

Authors: Zhou Sha[1, #], Kun Yang[2, #], Shoupeng Fu[3, #], Xiaoyong Tong[4, #], Hui Yang[1],

Huiling Fang[1], Qianqian He[1], Ning Li[5, 6], Xinyu Shu[5, 6], Qi Liu[1], Yongliang Du[1],

Beibei Fu[1], Yan Xiong[1], Dong Guo[1], Jin Liu[4], Qian Li[7], Hao Zeng[8], Xiaokai Zhang[8],

Rui Yao[9], Xushuo Zhang[10], Wenjin Guo[3], Xuhu Mao[7], Mian Long[5, 6], Xiaoyuan Lin[7,

11, *], Quanming Zou[8, *], Haibo Wu[1, *]

1 School of Life Sciences, Chongqing University, Chongqing 401331, China

2 Chongqing Public Health Medical Center, Chongqing 400036, China

3 School of Veterinary Medicine, Jilin University, Changchun 130062, China

4 School of Pharmaceutical Sciences, Chongqing University, Chongqing 401331,

China

5 Center of Biomechanics and Bioengineering, Key Laboratory of Microgravity

(National Microgravity Laboratory), and Beijing Key Laboratory of Engineered

Construction and Mechanobiology, Institute of Mechanics, Chinese Academy of

Sciences, Beijing 100190, China

6 School of Engineering Sciences, University of Chinese Academy of Sciences,

Beijing 100049, China

7 Department of Clinical Microbiology and Immunology, College of Pharmacy and

Laboratory Medicine, Army Medical University (Third Military Medical University),

Chongqing 400038, China

8 National Engineering Research Center of Immunological, Department of

Microbiology and Biochemical Pharmacy, College of Pharmacy and Laboratory

Medicine, Army Medical University (Third Military Medical University), Chongqing

400038, China

9 Department of Pathology, Chongqing Hygeia Hospital, Chongqing 401331, China

10 Zhejiang Toyouvax Biopharming, Hangzhou 311100, China

11 Institut für Virologie, Freie Universität Berlin, Robert-von-Ostertag-Str. 7-13,

Berlin 14163, Germany

# Equally Contributing Authors

* Corresponding authors

Xiaoyuan Lin, linxiaoyuan23@163.com

Quanming Zou, qmzou2007@163.com

Haibo Wu, hbwu023@cqu.edu.cn

Summary

Acute infection of the central nervous system is one of the most lethal diseases, yet the mechanisms by which intracellular bacteria infiltrate the brain remain poorly understood. Phagocytic cells are typically regarded as the primary site of conflict in the battle against intracellular bacteria; however, little is known about how the intracellular bacteria exploit infected phagocytes to gain access to the brain. In this study, we show that a novel $CD36^+Fabp4^+Pparg^+$ macrophage subpopulation ($CD36^+$ macrophage) participates in penetration of the brain by intracellular bacteria without disruption of the blood-brain barrier. Biomechanical analysis reveals that the abundance of protrusions and adhesion molecules on $CD36^+$ macrophages confers significant resistance to the mechanical stress of blood flow, thereby providing increased opportunities for these macrophages to adhere to the vascular endothelial surface. Metabolomics analysis identifies that macrophage lipid metabolism is dysregulated during bacterial neuroinvasion, and that β-hydroxybutyrate promotes the formation and survival of $CD36^+$ macrophages. Importantly, ketogenesis exacerbates symptoms during bacterial neuroinvasion, which could be halted by supplementing with physiological levels of glucose. Collectively, our findings elucidate a pathway by which intracellular bacteria hijack macrophages to invade the brain, suggesting that glycolipid metabolic homeostasis may play a role in the prevention or resolution of bacterial neuroinvasion.

Keywords: intracellular bacteria; infection; neuroinvasion; CD36[+] macrophage;

ketogenesis; β-hydroxybutyrate

Highlights

1. Intracellular bacteria hijack CD36[+] Mφ for brain invasion without disrupting the

BBB

2. CD36[+] Mφ has enhanced adhesion to BMECs and increased resistance to shear

flow

3. Ketogenesis promotes bacterial neuroinvasion that can be halted by glucose

treatment

4. Accumulation of peripheral BHB and CD36[+] Mφ fuels bacterial neuroinvasion in

humans

Introduction

Infections of the central nervous system (CNS) are among the most serious infectious

diseases caused by pathogens.[1] As a life-threatening infection with an extremely high

mortality rate, acute bacterial neuroinvasion are usually secondary to infected organs.

The blood-brain barrier (BBB) is the foremost line of defense, which consists of brain

endothelial cells working in concert with pericytes, smooth muscle cells, astrocytes,

microglia and macrophages to protect the CNS from infection.[2] Currently, pathogens

in the bloodstream are thought to penetrate the BBB via three main pathways:

transcellular entry, paracellular entry or "Trojan horse" spread.[1] It is known that some

bacteria can enter the brain directly by accumulation during transcytosis or by

disruption of tight junctions, accompanied by BBB leakage. For example,

Streptococcus pneumoniae (*S. pneumoniae*) interacts with host receptors on

endothelial cell monolayers to achieve brain invasion by transcytosis.[3-5] *Neisseria*

meningitidis hijacks specific host-pathogen interactions to redistribute tight junctions

or adhesion proteins, leading to BBB disruption.[6] *L. monocytogenes* employs the

invasion protein internalin F to facilitate the crossing of the blood-brain barrier [7,8], and

LLO and ActA have been identified to play a significant role during the invasion of

the olfactory epithelium for neurolisteriosis in infants via the intranasal route of

infection.[9] However, the "Trojan horse" route by which the bacteria circumvent the

BBB without damaging it has been rarely reported. Efforts to understand the host cells

that facilitate the neuroinvasion of intracellular bacteria are urgently needed.

Phagocytes, such as neutrophils and macrophages, play a dominant role in the host

response to intracellular bacteria.[10] The primary function of phagocytes is to engulf

and destroy microorganisms and cellular debris through tissue circulation and

migration. [11,12] By means of phagocytosis and antigen presentation, phagocytes act

rapidly as the first line of defense against bacterial infection. However, intracellular

bacteria can employ sophisticated strategies to replicate within phagocytes and evade

host immune surveillance.[13-15] Recently, *S. pneumoniae* was reported to hijack

receptor activity modifying protein 1 signaling in meningeal macrophages to facilitate

brain invasion.[16] Lecuit et al. showed that the surface protein internalin B of *L.*

monocytogenes protects infected monocytes from immune killing by CD8$^+$ T cells,

thereby prolonging the lifespan of infected cells and facilitating the spread of

infection throughout the CNS.[15] In this context, phagocytes, particularly monocytes

and macrophages, may play an important role in neuroinvasion, but how intracellular

bacteria take advantage of infected macrophages is not well understood.

The metabolism of host immune cells is reprogrammed during infection. Metabolic

signaling, such as glycolysis, helps to awaken the immune cells to resist pathogenic

invasion.[17] In addition, many studies have shown that bacteria manipulate host

metabolites to evade clearance by the host immune system.[13,18,19] *Legionella*

pneumophila induces Warburg-like metabolism by secreting the type 4 secretion

system effector MitF and decreases oxidative phosphorylation (OXPHOS) in infected

macrophages to increase the likelihood of pathogen survival.[18] A recent report

indicates that Nav1.8$^+$ nociceptors signal meningeal macrophages via the calcitonin

gene-related peptide, and this neuroimmune axis suppresses host defenses and

exacerbates bacterial meningitis.[16,19] Ketogenesis, induced by fasting or dietary

supplementation, is a metabolic pathway that provides the body with an alternative

form of energy and is closely related to the host immune response.[20,21] A recent study

reported that in pneumonia caused by *Pseudomonas aeruginosa*, ketone bodies are

recycled to the lungs via OXPHOS to maintain airway homeostasis.[22] However,

studies have demonstrated that mice fed a ketogenic diet develop a

hyperinflammatory response to systemic lipopolysaccharide and an increased

mortality rate, which are not attributable to diet-induced microbiota or

hyperglycemia.[23] The role of ketogenesis in the progression and migration of

infections remains a topic of contention.[24] Furthermore, the precise manner in which

ketogenesis contributes to the regulation of monocytic/macrophagic homeostasis and

the role in bacterial neuroinvasion is poorly defined.

In this study, we discovered a previously unidentified subpopulation of $CD36^+Fabp4^+$

$Pparg^+$ macrophages ($CD36^+$ macrophages) that serve as immune "Trojan horse" that

crosses the BBB without disrupting it, enabling intracellular bacterial neuroinvasion.

This subset of macrophages exhibits a foamy-like phenotype, which enhances the

ability of the cells to adhere and migrate vertically on the cerebrovascular endothelial

surface under the mechanical stress of blood flow. Furthermore, we found that serum

BHB, which could be produced by a ketogenic diet, exacerbated bacterial

neuroinvasion by increasing the generation and promoting the survival of $CD36^+$

macrophages in the peripheral blood. More importantly, we demonstrated that

supplementation of mice with physiological levels of glucose rescued the disturbance

in lipid metabolism, suppressed the generation of $CD36^+$ macrophages, and

significantly attenuated the clinical symptoms of encephalitis during bacterial brain

invasion. Our study outlines a new common strategy employed by intracellular

bacteria to invade the brain, which may be a potential therapeutic target for the

treatment of bacterial neuroinvasion.

Results

CD36$^+$ macrophages are potential carriers of bacteria during brain invasion

To investigate the role of immune cells in bacterial neuroinvasion, we constructed an

acute encephalitis mouse model caused by *L. monocytogenes*. This was achieved by

testing different infection routes and different doses, including intravenous injection

(1×10^5, 1×10^6, and 1×10^7 CFU) (Figure S1A) and intraperitoneal injection (2×10^6,

2×10^7, and 2×10^8 CFU) (Figure S1B). A comprehensive analysis of the survival rate,

the bacterial burden in brain tissues, and the pro-inflammatory cytokines in

cerebrospinal fluid (CSF) and brain homogenate revealed that both the intravenous

injection of 1×10^7 CFU and the intraperitoneal injection of 2×10^8 CFU *L.*

monocytogenes were successful in establishing an acute neuroinvasion mouse model

(Figure S1A and S1B). Additionally, histopathology confirmed the presence of

neuronal damage in the cerebral cortices, which is a hallmark of encephalitis (Figure

S1C). Behavioral changes, such as diminished activity, paresis, and loss of body

weight, were also observed (Figure S1D and S1E). To eliminate the possibility of

septic shock resulting from bacteremia, 1×10^7 CFU of heat-killed and Δhly *L.*

monocytogenes were intravenously administered to mice, respectively. The results

showed that neither heat-killed nor attenuated *L. monocytogenes* was capable of

causing acute neuroinvasion in the same manner as live bacteria, as evidenced by the

bacterial loads in the brain, the survival rates, and the clinical scores (Figure S1F-

S1H). To further substantiate the hypothesis that intracellular bacteria may utilize

phagocytes as a means of gaining access to the brain, gentamicin was administered to

mice four hours post-infection with *L. monocytogenes* (Figure S1I). The results

showed that when gentamicin was administered prior to infection, there was a

substantial remission in blood and brain bacterial loads, as well as an improvement in

survival rate of mice (Figure S1I-S1L). However, when gentamicin was administered

4 hours post infection, we found that only the bacteria in the circulation were

eradicated, and the brain bacterial loads and survival rate of mice were not

significantly different from those of the control group (Figure S1J-S1L). Importantly,

Evans blue staining revealed no appreciable extravasation of dye in these *L.*

monocytogenes-infected mice (Figure S1M). This finding effectively excludes the

possibility that the bacteria gain direct access to the brain by disrupting the BBB.

Collectively, these data suggested that intracellular bacteria may exploit host cells to

gain access to the brain in acute neuroinvasion.

To identify the specific carrier in the acute bacterial neuroinvasion model, peripheral

blood mononuclear cells (PBMCs) were collected and single-cell RNA sequencing

(scRNA-seq) was performed (Figure 1A). Immune cell lineages were identified using

canonical surface markers of dendritic cells (DCs: *Xcr1, Flt3, CD209a*), NK cells

(*Nkg7*), T cells (*CD3*), monocyte/macrophages (Mo/Mφ: *Fcgr1, Adgre1*), B cells

(*CD79*), neutrophils (*S100a9, S100a8*; Figure 1B and 1C).[25] Among these lineages,

Mo/Mφ is involved in phagocytosis and intracellular killing of microorganisms;

however, intracellular bacteria can also reside within these cells, thereby establishing

a long-term latent infection.[26] In this case, Mo/Mφ (n=3535 cells) were extracted and

reclustered *in silico* to precisely delineate subpopulations and their distributions

before and after neuroinvasion. Based on differentially expressed cell surface markers

and transcripts, we annotated Mo/Mφ into 6 subpopulations (Figure 1D, S2A). The

results showed that the percentage of Cluster 2 cells was sharply increased in the

neurolisteriosis group (Figure S2B). Further, we found that Cluster 2 cells expressed

high levels of genes related to lipid accumulation, including *CD36*, peroxisome

proliferator-activated receptor gamma (*Pparg*), fatty-acid binding protein 4 (*Fabp4*),

CD9, apolipoprotein E epsilon (*ApoE*), and triggering receptor expressed on myeloid

cells 2 (*Trem2*; Figure 1E).[27-31]

In order to gain insight into the characteristics of Cluster 2 macrophages, $CD36^+$ and

$CD36^-$ macrophages were isolated from peripheral blood by fluorescence-activated

cell sorting. In comparison to $CD36^-$ macrophages, the $CD36^+$ subpopulation

exhibited a markedly elevated level of granularity (SSC^{high}; Figure S2C), indicative of

an increased number of proteins or organelles within their intracellular compartment

and the presence of protrusions on their surfaces. The sorted macrophages were

cultured *in vitro* overnight and then labeled with BODIPY FL, a lipophilic probe for

polar and neutral lipids. The results showed that the $CD36^+$ subpopulation exhibited a

significantly greater accumulation of lipid droplets than the $CD36^-$ macrophages

(Figure S2D). Furthermore, Oil Red O staining was conducted to corroborate this

finding (Figure S2D). To confirm the lipid uptake ability of the $CD36^+$ macrophages,

we employed the use of fluorescently labeled 1,1′-dioctadecyl-3,3,3′,3′-

tetramethylindocarbocyanine perchlorate (DiI)-conjugated oxidized low-density

lipoprotein (ox-LDL), and the uptake of DiI-conjugated ox-LDL particles by CD36$^+$

and CD36$^-$ macrophages was quantified. Our findings revealed that CD36$^+$

macrophages internalized more than twice the amount of ox-LDL compared to CD36$^-$

macrophages (Figure S2E). In addition to CD36, other scavenger receptors, including

MARCO, SRA1, LDLR and MSR1, have also been implicated in the accumulation of

lipids.[32] Consequently, qRT-PCR assays were performed in order to examine the

expression of these receptors in CD36$^+$ macrophages. The results demonstrated that

only the expression of LDLR was significantly increased in this subpopulation

(Figure S2F), a finding that is consistent with the scRNA-seq data.

Subsequently, flow cytometry was performed to ascertain the percentage alterations in

CD36$^+$ macrophages over time in the context of neurolisteriosis (Figure S2C). The

results showed that the percentage and absolute number of CD36$^+$ macrophages in the

peripheral blood increased gradually over time during neuroinvasion (Figure 1F, S2G),

with the CD36$^+$ subgroup constituting more than 52% of all macrophages (Figure 1G).

Further, the CD36$^+$ subpopulation demonstrated the capacity to harbor a greater

quantity of bacteria (Figure S2H and S2I), and the percentage of CD36$^+$ macrophages

in peripheral blood was observed to be positively correlated with the bacterial load in

the brain (Figure 1H). These data implied that CD36$^+$ macrophages may be involved

in bacterial neuroinvasion. In addition, immunofluorescence co-localization analysis

was used to validate this hypothesis. The results showed that CD36$^+$ macrophages

attached to brain microvascular endothelial cells (BMECs) increased with the

progression of neuroinvasion in the neurolisteriosis mice (Figure S2J). Further,

transection of cerebral blood vessels was performed, and the vascular endothelial cells

were identified using an anti-CD31 antibody. The results showed that a substantial

number of CD36$^+$ macrophages and *L. monocytogenes* co-localized and aggregated on

the surface of the vascular endothelial cells (Figure 1I). In addition, CD36$^+$ and

CD36$^-$ macrophages were isolated from brain tissues of the neurolisteriosis mice. The

results obtained by flow cytometry, CFU counting, and immunofluorescence indicated

that the *L. monocytogenes* was located in the CD36$^+$ macrophages rather than CD36$^-$

macrophages (Figure 1J and 1K, S2K and S2L). As there was a clear inflammatory

response in the brain of mice during neuroinvasion, we subsequently examined the

inflammatory profile of the CD36$^+$ macrophages. The results showed that CD36$^+$

macrophage subpopulation isolated from mice exhibited an anti-inflammatory feature

(Figure S2M), suggesting that the inflammatory phenotype of the mice brain is not

dependent on the characteristics of CD36$^+$ macrophages themselves. Taken together,

these results suggested that CD36$^+$ macrophages may play a pivotal role in bacterial

brain invasion.

CD36$^+$ macrophages facilitate transport of intracellular bacteria across the

blood-brain barrier

Ox-LDL has been reported to induce a foamy-like phenotype by enhancing CD36

expression.[33] Consequently, to obtain CD36$^+$ macrophages in large quantities *in vitro*

in a rapid and efficient manner, we developed a method to induce the conversion of

peritoneal macrophages to CD36$^+$ macrophages by stimulation with different

concentrations of ox-LDL for 24 hours (Figure S3A). To investigate whether *in vitro*

generated CD36$^+$ macrophages are analogous to those isolated from peripheral blood,

the purified peritoneal macrophages from mice were treated with ox-LDL (80 μg/mL)

and then CD36$^+$ macrophages were sorted with an APC-conjugated CD36 antibody

(Figure S3B). Through Western blotting and RNA-seq analysis, we found that CD36$^+$

macrophages generated by ox-LDL treatment *in vitro* exhibited Fabp4 and Pparg

expression patterns (Figure S3C), and transcriptomic profiling that were highly

similar to those of CD36$^+$ macrophages isolated from peripheral blood (Figure S3D

and S3E). Furthermore, we showed that ox-LDL treatment influenced the

development of CD36$^+$ macrophages (Figure S3A), however this signal did not

impact the inflammatory profile once CD36$^+$ macrophages had been formed (Figure

S3F and S3G).

Next, to ascertain the role of CD36$^+$ macrophages in bacterial neuroinvasion, we

constructed an *in vitro* BBB model comprising mouse primary BMECs and neurons in

a Transwell system (Figure 2A). The integrity of the cell monolayer was evaluated by

measuring the transepithelial electrical resistance (TEER) values using a Millicell-

ERS voltohmmeter, and cell monolayers with TEER values above 300 Ω/cm^2 were

considered intact and functional.[34] We showed that the integrity of the *in vitro* BBB

model was not significantly affected by high dose and short period of bacterial

infection (Figure S3H). Subsequently, macrophages were infected with *L.*

monocytogenes (MOI=10) and then inoculated into the upper Transwell chamber of

the BBB model. The transformation of peritoneal macrophages into $CD36^+$

macrophages was confirmed to result in an increase in the intracellular bacterial load

(Figure S3I). Additionally, the assistance of ox-LDL-stimulated macrophages was

observed to significantly increase the number of *L. monocytogenes* organisms

crossing the BBB in a dose-dependent manner (Figure 2B). Furthermore, crystal

violet staining also demonstrated a dose-dependent increase in the number of

macrophages penetrating the endothelial monolayer upon stimulation with ox-LDL

(Figure 2C and 2D). To validate whether *L. monocytogenes* reside in the $CD36^+$

macrophages and cross the BBB, bacterial staining of these penetrating macrophages

was performed. The results showed that bacteria were present in the cells and

successfully crossed the BBB in the *in vitro* model (Figure S3J and S3K). In order to

conduct further investigation into the capacity of the $CD36^-$ but with high lipid

content subpopulation ($CD36^-$ $lipid^{high}$) to promote neuroinvasion, we tested whether

these $CD36^-$ $lipid^{high}$ macrophages could facilitate bacterial crossing of the barrier

using an *in vitro* BBB model. Interestingly, we did not observe significant differences

in the bacterial burden between $CD36^-$ $lipid^{high}$ and $CD36^-$ $lipid^{low}$ macrophages

(Figure S3L-S3N). This result suggested that, despite the crucial role of lipid

metabolism in the formation and function of $CD36^+$ macrophages, the CD36 receptor

may have a hitherto unknown function during the bacterial neuroinvasion.

To ascertain whether the rise in CD36$^+$ macrophages facilitated the transfer of *L.*

monocytogenes into the brain *in vivo*, we performed adoptive macrophage transfer

experiments via tail vein injection based on an irradiated mouse model (Figure 2E).

Bone marrow irradiation-based immunosuppression resulted in mice being more

susceptible to bacterial infection, which necessitated a reduction of the infectious dose

to 1×10^6 CFU of *L. monocytogenes.* The results showed that the survival time was

shortened, the body weight was reduced, and the comprehensive clinical score was

increased in the mice receiving CD36$^+$ macrophages compared with the control group

(Figure 2F, S3O and S3P). Furthermore, we showed that the brain bacterial loads in

mice receiving CD36$^+$ macrophages were significantly higher than those in mice

receiving CD36$^-$ macrophages (Figure 2G). Interestingly, no significant differences

were observed in bacterial loads among the lung, spleen, and liver (Figure S3Q).

Additionally, we examined the levels of pro-inflammatory factors in the CSF and

brain homogenate. The results showed a notable elevation in interleukin 6 (IL-6) and

tumor necrosis factor alpha (TNF-α) levels (Figure 2H and 2I), indicating a more

severe encephalitis phenotype in the CD36$^+$ macrophage transfer group. Importantly,

the brains of mice transferred with CD36$^+$ macrophages exhibited no discernible

Evans blue extravasation, suggesting that these macrophages facilitated the transfer of

L. monocytogenes into the brain without disrupting the BBB (Figure S3R and S3S).

The results of the pathological histology analysis confirmed the existence of more

severe inflammation and neuronal damage in the cerebral cortices of the mice

transferred with CD36$^+$ macrophages than in the control group (Figure S3R and S3T).

Furthermore, when mice were infected with a relatively low dose (1×10^5 CFU/mouse)

of *L. monocytogenes*, the difference in bacterial load in brain tissues between the

CD36$^+$ macrophage transferred mice and the control group was also observed at day 2

post infection (Figure S3U-S3W). These data suggest that neuroinvasion mediated by

CD36$^+$ macrophages is not limited to an acute infection model induced by a high dose

of bacteria.

We subsequently investigated whether CD36$^+$ macrophages may perform a

comparable function during the invasion of the brain by other intracellular bacteria. In

the pertinent experiments, a positive correlation was observed between CD36$^+$

macrophages and neuroinvasion in mice infection models of the intracellular

bacterium *Salmonella typhimurium* (*S. typhimurium*; Figure S4A and S4B). The same

results were observed in an infection model of Methicillin-resistant *Staphylococcus*

aureus (MRSA; Figure S4C and S4D), which, although not a classical intracellular

pathogen, exhibits significant intracellular habitation.[35] The results of the

immunofluorescence and bacterial burden analyses of the macrophages isolated from

PBMCs of infected mice showed that *S. typhimurium* and MRSA were predominantly

present in the CD36$^+$ macrophages (Figure S4E and S4F). However, this finding was

not validated in a neuroinvasion model of the extracellular bacterium *S. pneumoniae*

(Figure S4G and S4H). Next, adoptive macrophage transfer experiments were

conducted to further substantiate the hypothesis that CD36$^+$ macrophages serve as

carriers in the intracellular bacterial invasion of the brain. Once more, disparities in

brain tissue bacterial loads were discerned between the CD36$^+$ macrophage-

transferred and control groups in the neuroinvasion model of MRSA (Fig. S4I-S4K),

though not in the model utilizing *S. pneumoniae* (Figure S4L-S4N). Taken together,

these data suggested that CD36$^+$ macrophages contributed to the process of

intracellular bacterial neuroinvasion.

Proprotein convertase subtilisin/kexin 9 (PCSK9) belongs to the proprotein

convertase family, members of which participate in cholesterol and fatty acid (FA)

metabolism.[36,37] The intracellular accumulation of low-density lipoprotein cholesterol

(LDL-C) in PCSK9$^{-/-}$ is attributed to the inability of the LDL receptor, which is

located on the cell membrane, to undergo normal degradation.[38,39] The mRNA and

protein levels of PCSK9 in CD36$^+$ macrophage subsets promoted by *L.*

monocytogenes infection and in peritoneal macrophages exposed to ox-LDL were

examined. The results showed that PCSK9 levels were significantly reduced in both

circulating and oxLDL-induced CD36$^+$ macrophages (Figure S5A and S5B). Further,

we found that macrophages lacking PCSK9 exhibited markedly elevated lipid content

in comparison to the control group (Figure S5C), a finding that was corroborated by

the expression of FABP4 and PPARG (Figure S5D). Also, it was observed that a

greater number of *L. monocytogene* organisms were able to parasitize PCSK9$^{-/-}$

macrophages than those parasitizing wild-type (WT) cells (Figure S5E). In the context

of neurolisteriosis, PCSK9$^{-/-}$ mice exhibited a higher percentage of CD36$^+$

macrophages in peritoneal blood than their littermates (Figure S5F). In addition, the

results of survival rate, bacterial burden in brain tissues, and pro-inflammatory

cytokines in CSF revealed that PCSK9$^{-/-}$ mice developed a more pronounced bacterial

neuroinvasion in comparison to their littermates following tail vein injection of 1×10^7

CFU of *L. monocytogenes* (Figure S5G-S5I). Consequently, PCSK9$^{-/-}$ mice were

utilized as a model for the spontaneous formation of CD36$^+$ macrophages, and the

veracity of our findings was validated through adoptive macrophage transfer.

Subsequent to the irradiation and a 12-hour recovery period post transfer, recipient

mice were intravenously inoculated with 1×10^6 CFU of *L. monocytogenes*.

Gentamicin was used to eliminate extracellular bacteria. Two days later, samples were

collected for further examination (Figure 2J). We found that the brain bacterial loads

were significantly higher in PCSK9$^{+/+}$ recipients transferred with PCSK9$^{-/-}$

macrophages than those that receiving WT macrophages (Figure 2K). Similar results

were obtained in the PCSK9$^{-/-}$ recipients (Figure 2K). However, the differences in

bacterial were observed exclusively in the brain tissues, and not in other organs such

as the lung, spleen, and liver (Figure S5J). Consistently, the mortality rate, and pro-

inflammatory factors in the CSF and brain homogenate were exacerbated in both

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ recipients by the transfer of PCSK9$^{-/-}$ macrophages (Figure

2L, S5K and S5L). We further confirmed the aforementioned finding by intravenous

injection of 1×10^5 CFU of *L. monocytogenes* in the same experimental model (Figure

2M and 2N, S5M-S5O).

Furthermore, in order to test the impact of PCSK9 function on the pathology driven

by CD36$^+$ macrophages during *L. monocytogenes* infection, adoptive macrophage

transfer of CD36$^+$ and CD36$^-$ macrophages sorted from the PCSK9$^{-/-}$ mice were

performed (Figure S5P). The results showed that the brain bacterial loads were

significantly higher in recipients receiving PCSK9$^{-/-}$CD36$^+$ macrophages compared

with those receiving PCSK9$^{-/-}$CD36$^-$ macrophages (Figure S5Q and S5R). This

finding was further corroborated by elevated mortality rates and pro-inflammatory

factors in the CSF (Figure S5S and S5T). These results suggested that, while PCSK9

regulates CD36$^+$ macrophage formation by modulating intracellular lipids, it does not

function directly in CD36$^+$ macrophage-driven pathology during bacterial

neuroinvasion. Taken together, these data indicated that excessive lipid accumulation

contributed to the formation of CD36$^+$ macrophages, thereby facilitating

neuroinvasion by intracellular bacteria.

Shear flow analysis demonstrates increased adhesion and resistance of CD36$^+$

macrophages during neuroinvasion

We next asked how CD36$^+$ macrophages exploited their properties to accumulate on

the cerebral blood vessel wall and participate in the process of intracellular bacterial

neuroinvasion. Using a two-dimensional live-cell flow chamber assay with a

microfluidic chip[40], we deciphered the multistep process of how macrophages adhered

to and moved on the surface of vascular endothelium under physiological flow

(Figure 3A). Cellular adhesion ability was calculated by counting total macrophages

adhering to the endothelial monolayer under a given shear stress. We found that an

increase in the percentage of CD36$^+$ macrophages was concomitant with an increase

in the number of cells adhering to the endothelial monolayer (Figure 3B and 3C).

Immunofluorescence was performed to show that *L. monocytogenes* were residing in

the CD36$^+$ macrophages in the biomechanical analysis (Figure S6A). Because

adherent cells could be detached by an increase in shear stress, we tested the

dynamics of CD36$^+$ macrophages' shear resistance. The results showed that such

resistance was significantly stronger than that of control macrophages: CD36$^+$

macrophages generated either from ox-LDL treatment or bacterial infection, exhibited

remarkable resistance to high shear stress (Figure 3D). Further, we investigated the

effect of endothelial density on macrophagic adhesion. The results showed that

endothelial cell density, rather than cell type, greatly affected macrophagic adhesion,

and CD36$^+$ macrophages tended to adhere to the surface of dense endothelial cells

(Figure S6B-S6E).

Once attached to the endothelial monolayer, the macrophage crawls along it and finds

a place to cross it in the presence of blood flow.[41,42] We next tested the performance of

CD36$^+$ macrophages in post-arrest crawling on the endothelial monolayer. We

acquired sequential images of macrophages at a shear stress of 2 dyn/cm^2 and then

analyzed them by defining various parameters of crawling dynamics (Figure S6F).

The observations were presented by crawling trajectories of each individual cells

(Figure 3E), and we found that the crawling ability of adherent CD36$^+$ macrophages

was dampened, reflecting in shortened crawling displacement and slower crawling

speed (Figure 3F and 3G). However, we saw no obvious differences in crawling

duration (Figure S6G). Directionality is also a critical aspect of cell crawling

dynamics, which is quantified as the ratio of displacement perpendicular to shear flow

to that along shear flow. Our results showed that the directionality of macrophages

was slightly but significantly enhanced by ox-LDL stimulation, implying that $CD36^+$

macrophages exhibited a propensity to migrate in the vertical direction (Figure 3H).

We also compared a key dynamic parameter of transmigration duration, which is

defined as the interval from the timepoint of cell arrival at the site to the endpoint of

completion of transmigration beneath the endothelial monolayer (Figure S6H). The

result showed that transmigration duration was significantly shortened in $CD36^+$

macrophage (Figure 3I).

Considering that protrusion is critical for cell crawling and migration, we then

investigated whether intracellular lipid could affect the formation of protrusions. The

number and the length of protrusions, and cell area were measured using the ImageJ

software (Figure S6I). The results showed that the numbers, as well as the lengths of

protrusions were significantly increased in $CD36^+$ macrophages (Figure 3J and 3K).

We further quantified macrophagic cell area and found that the mean area of $CD36^+$

macrophages was significantly larger than the control (Figure 3L). Additionally, we

examined various integrins and adhesive ligands expressed on the cell surface, which

have been reported to be involved in adhesion and migration.[43] The results showed

that the expression levels of α-integrins, β-integrins and intercellular adhesion

molecules were significantly upregulated in the infected CD36$^+$ macrophages (Figure

3M and S6J). Additionally, these data were corroborated by the circulating CD36$^+$

macrophages sorted from peripheral blood (Figure S6K). Further, we tested the

adhesion and movement ability of CD36$^+$ macrophages when adhesion molecule

ICAM-1 was blocked. The data showed that the cellular adhesion ability, crawling

displacement and crawling speed were significantly impaired, whereas the

directionality and transmigration duration of macrophages were barely affected by

ICAM1 antibody treatment (Figure S6L-S6S). Collectively, these data suggested that

the foaminess of macrophages greatly enhanced their ability to adhere under

mechanical stress, predisposing them to stay and to migrate vertically on the surface

of BMECs; this might be attributable to the formation of protrusions as well as the

expression of integrins and adhesion ligands.

β-hydroxybutyrate promotes CD36$^+$ macrophage formation and exacerbates

neuroinvasion

Next, we performed lipid metabolomics analysis of serum samples to reveal how

CD36$^+$ macrophages were formed during bacterial neuroinvasion. Principal

component analysis (PCA) showed that distinct metabolite genotype clusters were

generated in the neurolisteriosis group (Figure 4A). The volcano plot indicated that 70

fatty acids (FAs) were upregulated and 71 FAs were downregulated ($p < 0.05$,

$|\log_2(\text{fold change})| > 1$) in the neurolisteriosis group (Figure 4B, Table S1). The 70

upregulated FAs were classified into four subclusters by the number of carbon atoms

in the alkyl chain. The synthesis and metabolism cycle of long-chain FAs is longer

than that of short-chain FAs. Based on the consideration of rapid progression of

bacterial neuroinvasion, we focused on short-chain FAs and found that β-

hydroxybutyrate (BHB), which has been reported to favor the foamy-like phenotype

of macrophages [44], was the only short-chain FA among all upregulated FAs (Figure

4C). Kyoto Encyclopedia of Genes and Genomes (KEGG)-based analysis revealed

that metabolites related to the synthesis and degradation of ketone bodies were

enriched (Figure S7A), and metabolomics data showed that BHB was significantly

increased during neuroinvasion (Figure S7B). Further, we found that BHB levels in

mice serum climbed steadily over time in the acute neuroinvasion model of *L.*

monocytogenes infection (Figure 4D). This finding is consistent with the increases in

the bacterial load in the brain and the proportion of CD36$^+$ macrophages in the

PBMCs (Figure S7C and S7D).

Subsequently, we sought to explore the reasons for the enhanced ketogenesis during

bacterial brain invasion. Liver tissues from acute neuroinvasion mice model were

collected for transcriptome analysis by RNA-seq. The results showed that compared

with controls, the genes associated with ketogenesis were sharply upregulated in the

livers of mice suffered from neurolisteriosis, whereas these genes were moderately

increased in the mildly infected group (Figure S7E). KEGG-based analysis and gene-

set enrichment analysis (GSEA) between the neurolisteriosis and control group

revealed that the differentially expressed (DE) genes were enriched in glycolysis or

gluconeogenesis (Figure S7F and S7G). To validate this finding, we measured serum

glucose levels in mice and found that glucose levels decreased significantly and

remained low during *L. monocytogene*-induced neuroinvasion (Figure S7H);

meanwhile, we observed a significant decrease in cellular energy and an increase in

ADP/ATP ratios in mice with neurolisteriosis compared to controls (Figure S7I and

S7J). These data implied that the decrease in glucose levels and increase in

ketogenesis may be induced by excessive glycolysis during *L. monocytogene*-induced

neuroinvasion. Further, we showed that the extracellular acidification rates (ECAR)

were significantly increased in the livers of mice with neurolisteriosis (Figure S7K).

This data was confirmed by the measurement of lactate, the end product of glycolysis

(Figure S7L). Also, we found that mice experienced weight loss and reduced food

intake, which to a certain extent contributed to ketogenesis after infection (Figure

S7M and S7N). Interestingly, there was no significant difference between the mildly

infected group and the control group in these indicators during the experiments

(Figure S7H-S7N). Furthermore, a steady increase in serum BHB and a decrease in

glucose levels were also observed in the neuroinvasion models established using *S.*

typhimurium and MRSA (Figure S7O and S7P). These data indicated that BHB

production was significantly induced during bacterial neuroinvasion, possibly due to

low serum glucose levels induced by excessive glycolysis.

As previously reported, autocrine or paracrine BHB directly activates macrophages

function mainly via the hydroxycarboxylic acid receptor 2 (HCAR2).[45] Therefore, we

next used primary peritoneal macrophages isolated from HCAR2 knockout mice to

study whether BHB could promote bacterial neuroinvasion. We confirmed that lipid

accumulation and percentage of CD36^{+} macrophages all increased dramatically upon

BHB stimulation, and there was no significant difference when HCAR2 was knocked

out (Figure 4E-4G, S7Q). BHB inhibits adenylyl cyclase activity through the HCAR2

receptor and reduces cellular cyclic adenosine monophosphate (cAMP) levels, thereby

protecting lipid droplets from lipolysis by hormone-sensitive lipases (HSL).[45] As a

result, we found that cAMP pretreatment eliminated the decrease in HSL activity, the

increase in lipid content and intracellular bacterial burden stimulated by BHB;

however, this phenomenon was not found in HCAR2-deficient macrophages (Figure

S7R-S7T). By using the *in vitro* BBB model (Figure 2A), the number of bacteria

crossing the barrier increased significantly in response to BHB stimulation in controls,

but not in HCAR2$^{-/-}$ macrophages (Figure 4H). Next, peritoneal macrophages from

HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice were isolated to further assess the effects of BHB on

the biophysical dynamics of macrophages. The data demonstrated that BHB

significantly altered the static or shear-resistant adhesion of control macrophages, but

not of HCAR2$^{-/-}$ macrophages (Figure 4I-4K). We also found that BHB greatly

impaired the crawling ability, including shortened crawling displacement, slowed

down crawling speed and unpredictable directionality of macrophages (Figure 4L-4O).

In addition, the transmigratory ability of macrophages was significantly increased

when cells were pretreated with BHB (Figure 4P). Consistently, the expression levels

of the α-integrin family were significantly upregulated by BHB treatment in controls,

but not in HCAR2-deficient macrophages (Figure 4Q). Taken together, these data

indicated that BHB facilitated the formation of CD36$^+$ macrophages through the

HCAR2 receptor, thereby affecting the process of bacterial neuroinvasion.

BHB promotes the survival of macrophages via HCAR2-mediated calcium influx

Because the foamy stage appears to be a terminal point of macrophage differentiation

[46], we wondered whether, in addition to converting peritoneal macrophages to the

foamy phenotype, cell survival also contributed to the increase in absolute quantity of

CD36$^+$ macrophages induced by BHB treatment. Immunofluorescence was performed

with the objective of showing the presence of *L. monocytogenes* within the

macrophages (Figure S8A). The results showed that the lactate dehydrogenase (LDH)

activity in the culture medium was significantly reduced in response to BHB

treatment, indicating a decrease in cell death that induced by infection (Figure 5A).

Propidium iodide (PI) staining was used to evaluate the membrane integrity of

infected macrophages, and we found that the proportion of PI$^+$ cells decreased from

59.7% to 24.5% under BHB stimulation, with no significant difference in cell

viability between the BHB treatment and control groups when HCAR2 was knocked

out (Figure 5B and 5C). Calcium influx is generally recognized as a signal that

triggers cell membrane repair.[47] To investigate the impact of BHB-HCAR2 signaling

on cytosolic calcium concentration, infected macrophages were loaded with Fluo-4

AM, a fluorescent Ca^{2+} indicator. The results showed that BHB induced remarkable

accumulation of intracellular calcium, and HCAR2 knockdown eliminated calcium

influx in macrophages (Figure 5D). Pretreatment of macrophages with BAPTA, a

Ca^{2+} chelator, reversed the BHB-induced increase in cell survival (Figure 5E and 5F).

Further, we tested whether calcium flux was associated with CD36$^+$ macrophage-

mediated intracellular bacterial penetration of the BBB. As a result, inhibition of

calcium flux had no effect on bacterial uptake by macrophages in the upper chamber

(Figure S8B); however, we observed differences in bacterial loads in the macrophages

that had crossed the barrier (Figure 5G).

We next investigated how BHB maintained the survival of CD36$^+$ macrophages. As

previously reported, calcium influx initiates membrane repair mainly through

recruitment of the endosomal sorting complexes required for transport (ESCRT)

machinery.[48] Our data showed that BHB treatment significantly increased the number

of puncta that were positive for charged multivesicular body protein 4B (CHMP4B),

which indicated assembly of ESCRT in the infected macrophages; HCAR2

knockdown or BAPTA treatment could block this membrane repair process (Figure

5H and 5I, S8C and S8D). Cellular membrane repair also relies on lysosomes, which

can rapidly fuse with the plasma membrane when intracellular Ca^{2+} concentration

rises. Therefore, we examined lysosomal exocytosis in the infected macrophages

using exofacial lysosomal-associated membrane protein 1 (LAMP1) and the release of

lysosomal enzymes, such as acid phosphatase (ACP), β-galactosidase (β-gal) and β-

hexosaminidase (β-hex). The results confirmed that BHB promoted the lysosomal

exocytosis in infected macrophages through the HCAR2 receptor (Figure 5J-5L, S8E-

S8G). As thioglycolate has been demonstrated to initiate an inflammatory response

and alter lysosomal enzymes during the induction of PBMCs into macrophages [49], we

further detected the lysosomal exocytosis of infected resident peritoneal macrophages

and found that BHB indeed promoted the lysosomal exocytosis, thereby maintaining

the survival of resident CD36$^+$ macrophages (Figure S8H-S8J). Lipophagy has been

reported to implicate in regulating lipid metabolism and cell survival, particularly

during intracellular bacterial infections [50], we asked whether lipophagy was involved

in modulating phagocyte survival during infection. The results showed that lipophagy

was significantly activated in CD36$^+$ macrophages following BHB treatment.

However, this activation was abrogated in the HACR2 knockout group (Figure S8K).

Importantly, we found that the proportion of CD36$^+$PI$^+$ macrophages was reduced

after BHB treatment, and that the effect of BHB was significantly inhibited by

pretreatment with 100 nM lipophagy inhibitor Baf-A1 for 6 hours (Figure S8L). The

results suggested that BHB promotes macrophage survival, which is likely to be

dependent on lipophagy. Collectively, these data indicated that BHB signaling

activated Ca^{2+}-mediated membrane repair, thereby maintaining the survival of CD36$^+$

macrophages via the HCAR2 receptor.

Ketogenesis exacerbates intracellular bacterial neuroinvasion

BHB can be synthesized via the metabolism of FAs and ketogenesis.[51] To elucidate

the correlation between ketogenesis and bacterial neuroinvasion, mice were

administered varying doses of a ketone ester drink for a period of two weeks, after

which they were infected with 2×10^8 CFU of *L. monocytogenes* via intraperitoneal

injection (Figure 6A). The data showed that the ketone ester elevated the levels of

serum BHB and the proportion of CD36$^+$ macrophages in peritoneal blood (Figure 6B

and 6C), accompanied by remarkable increases in the clinical scores, mortality rate

and bacterial load in brain tissues (Figure S9A, 6D and 6E). It has been established

that dietary modulation can increase and maintain circulating metabolites in the

human circulation, and a growing body of evidence demonstrates that a ketogenic diet

elevates BHB levels.[52] The ketogenic diet is a low-carbohydrate, high-fat diet that has

been demonstrated to result in weight loss and confer a multitude of health benefits,

including the alleviation of symptoms associated with neurodegenerative diseases,

cardiovascular diseases, and cancers.[21,53-55] We proceeded to investigate whether a

ketogenic diet was associated with neurolisteriosis. The HCAR2$^{-/-}$ mice and their

littermates were fed a normal or ketogenic diet for four weeks, respectively, and then

infected with *L. monocytogenes* via intravenous injection (Figure 6F). Consequently,

the lifespan of HCAR2$^{+/+}$ mice fed a ketogenic diet was markedly reduced in the

presence of *L. monocytogenes* infection, and the detrimental impact of dietary

modulation was abrogated by HCAR2 knockout (Figure 6G). Consistently, the

clinical scores (Figure S9B), percentages of CD36$^+$ macrophages in peritoneal blood

(Figure 6H), the intracellular bacterial burden of macrophages (Figure 6I), and

bacterial loads in brain tissues (Figure 6J) were all significantly increased in

HCAR2$^{+/+}$ mice on a ketogenic diet. Interestingly, mice in this group exhibited less

weight loss than mice in the control group, and experienced greater dissemination of

bacteria to other organs (Figure 6J and S9C). Subsequently, the levels of pro-

inflammatory factors in the CSF and brain homogenate were examined. The results

showed that a ketogenic diet significantly increased the levels of pro-inflammatory

cytokines and precipitated more pronounced neuronal injury in HCAR2$^{+/+}$ mice,

whereas it had no discernible impact on HCAR2$^{-/-}$ mice (Figure 6K-6N). Notably, no

extravasation of Evans blue was observed in the brains of mice fed a ketogenic diet,

thereby ruling out the possibility of direct damage to the BBB (Figure 6M and 6O).

These data were corroborated by an intraperitoneal injection of 2×10^8 CFU of *L.*

monocytogenes in the same experimental model (Figure S9D-S9G). Further, the

inflammatory profile of the CD36$^+$ macrophages generated from mice fed a ketogenic

diet was examined. It was observed that these CD36$^+$ macrophages also exhibited an

anti-inflammatory phenotype (Figure S9H), and the inflammatory signature was not

affected by BHB-HCAR2 signaling once CD36$^+$ macrophages was formed (Figure

S9I and S9J). These characteristics were identical to those of CD36$^+$ macrophages

isolated from mice following infection (Figure S2M, S3F and S3G).

Further, to exclude the possibility of BHB exerting its effects by preventing the

clearance of bacteria before they reach the brain, we isolated peritoneal macrophages

from HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice to perform adoptive macrophage transfer

experiments (Figure S9K). Gentamicin was used to eliminate circulating bacteria that

had not yet entered the phagocytes. The results showed that mice transferred with

macrophages obtained from HCAR2$^{+/+}$ mice fed a ketogenic diet had a shortened

survival time in comparison to mice fed a control diet. However, this difference in

survival time between different diets was eliminated in the HCAR2$^{-/-}$ group (Figure

S9L). The observation was further confirmed by examining bacterial loads in the

brain tissues, and inflammatory factor levels in the CSF of recipient mice (Figure

S9M and S9N). In addition to HCAR2, BHB has been shown to act through other

mechanisms, therefore a 3-hydroxy-3-methylglutaryl-CoA synthase 2 (HMGCS2, the

rate-limiting enzyme during ketogenesis) knockout mouse line was used to validate

our findings (Figure S9O). The results showed that intravenous infection with 1×10^7

CFU of *L. monocytogenes* resulted in a dramatic decrease in blood BHB levels and

the proportion of CD36$^+$ macrophages in the peritoneal blood of the Hmgcs2

knockout mice compared with the controls (Figure S9P and S9Q), accompanied by a

significant reduction in mortality rates, brain tissue bacterial loads, and CSF

inflammatory factor levels (Figure S9R-S9T). These data suggested that BHB may

exert its effects though promoting the formation of CD36$^+$ macrophages via HCAR2

receptor, thus exacerbating bacterial neuroinvasion. We also employed MRSA to

further verify our findings. The data indicated that, as expected, the ketogenic diet

facilitated brain invasion by MRSA and intensified the phenotypes of neuroinvasion

(Figure S9U-S9W). Taken together, our data demonstrated that a ketogenic diet could

exacerbate bacterial neuroinvasion and hasten the demise of mice, primarily

depending on the elevation in serum BHB via the HCAR2 receptor.

Our data thus far suggest that BHB plays an important role in bacterial neuroinvasion

in mice, and that the enhancement in ketogenesis is probably induced by low serum

glucose levels. Subsequently, the impact of restoring blood glucose levels on the

outcomes of infected mice was evaluated. The results showed that in an acute

neuroinvasion model of *L. monocytogenes* infection, the supplementation of mice

with glucose at 12-hour intervals successfully restored blood glucose levels and

inhibited hepatic glycolysis (Figure S10A-S10F), leading to significant decreases in

BHB levels and CD36$^+$ macrophage percentages (Figure S10G-S10H). Surprisingly,

we found that four rounds of glucose supplementation significantly increased the

survival of mice, and reduced their clinical scores, brain bacterial burden and pro-

inflammatory cytokines levels (Figure S10I-S10M). These data indicate that glucose

homeostasis may limit the brain invasion by *L. monocytogenes* infection and alleviate

neuroinvasion symptoms in mice.

Serum BHB fuels the symptoms of human bacterial neuroinvasion

A retrospective study based on the Medical Information Mart for Intensive Care IV

(MIMIC-IV) was performed to explore the risk factors for bacterial neuroinvasion in

humans. As a result, ferritin (OR 5.09, 95% CI 1.55-16.71), fibrinogen (OR 3.57, 95%

CI 1.13-11.25), LDL (OR 3.31, 95% CI 1.01-10.83), glucose (OR 2.17, 95% CI 1.04-

4.50) and aspartate amino transferase (OR 2.16, 95% CI 1.01-4.63) were identified as

independent risk factors, whereas potassium (OR 0.42, 95% CI 0.20-0.87), alkaline

phosphatase (AP, OR 0.38, 95% CI 0.18-0.83), phosphate (OR 0.23, 95% CI 0.11-

0.50), high-density lipoprotein (HDL, OR 0.19, 95% CI 0.05-0.66) and transferrin

(OR 0.14, 95% CI 0.04-0.50) were identified as protective factors in patients with

bacterial neuroinvasion (Figure S11A). Next, we asked whether the accumulation of

BHB or CD36[+] macrophages could also be identified and were associated with

bacterial infection of the CNS in human clinical samples. Gender- and age-matched

bacterial neuroinvasion subjects and healthy individuals were recruited and peripheral

blood samples were collected for analysis (Figure 7A). Since blood glucose is a risk

factor for bacterial neuroinvasion and BHB level can be dramatically affected by

glycolysis, subjects with abnormal blood glucose or insulin concentrations were

excluded from the two groups (Figure S11B-S11E). As a result, serum BHB levels

were significantly elevated in patients compared with healthy controls, suggesting an

infection-induced ketogenesis (Figure 7B). Interestingly, serum cholesterol levels,

including total cholesterol (TC) and LDL-C were increased in patients compared with

healthy controls, while HDL-C was slightly but not significant reduced (Figure S11F-

11H). This finding is consistent with data from our retrospective study based on

MIMIC-IV (Figure S11A). To investigate whether BHB was associated the severity of

infection, we further divided patients into three groups, high BHB (in the top 1/3

among samples tested, n=21), moderate (middle 1/3, n=20) and low (bottom 1/3,

n=21; Figure 7A). Clearly, patients with high BHB levels exhibited higher clinical

scores than patients with low BHB levels (Figure 7C). Consistently, high-BHB

patients had higher cell counts and more protein in the CSF (Figure 7D and 7E);

higher infection indices such as white blood cell counts (WBC; Figure 7F) and

procalcitonin (PCT; Figure 7G); and higher values for markers of inflammatory

conditions such as erythrocyte sedimentation rate (ESR; Figure 7H) and C-reactive

protein (CRP; Figure 7I), suggesting increased disease severity. These data implied

that infection-induced ketogenesis might provide important fuel for severe bacterial

neuroinvasion.

To explore the dynamics of CD36$^+$ macrophages and BHB in the pathogenesis of

bacterial neuroinvasion, we followed-up with 7 individuals during their

hospitalization. We found that CD36$^+$ macrophages and BHB levels in patients tended

to increase during disease progression, and decrease during the recovery stage (Figure

7J and 7K). The intracellular bacterial load of CD36$^+$ macrophages sorted from

patients was also calculated. The results showed that bacteria were predominantly

present in CD36$^+$ macrophages, but not in CD36$^-$ macrophages (Figure S11I). To

further confirm the relationship between macrophage lipid metabolism and bacterial

neuroinvasion pathology, we collected PBMCs from patients or healthy controls for

analysis. Compared with healthy controls, the expression levels of genes involved in

cholesterol uptake were significantly increased, while those of genes related to

cholesterol utilization and excretion were decreased in macrophages from patients

(Figure 7L). Taken together, these data indicated that BHB fueled human bacterial

neuroinvasion by promoting the development of CD36$^+$ macrophages.

Discussion

Our findings suggested that CD36$^+$ macrophages may serve as a carrier for

intracellular bacteria that colonize the brain, thereby facilitating the penetration of

these bacteria across the BBB. Generally, the BBB prevents pathogens from crossing

the brain tissue unless there is significant trauma to the brain. Nevertheless, specific

bacteria are capable of breaking through the BBB via their distinctive mechanisms,

leading to severe brain infections.[15,56]. In this study, we discovered that intracellular

bacteria may prefer to exploit a "Trojan horse" pathway, which causes minimal

damage to the BBB but enables acute brain invasion. It is possible that $CD36^+$

macrophages may be hijacked by intracellular bacteria and serve as vehicles for

neuroinvasion. In comparison to peritoneal macrophages, $CD36^+$ macrophages exhibit

a greater number of protrusions and a larger cell area, which renders them more

conducive to colonization on BMECs under the pressure of blood flow (Figure 3E-

3L). Based on the aforementioned remarkable characteristics, $CD36^+$ macrophages

have the potential to become carriers of engineered intracellular bacteria for

bacteriotherapy.

Pathogens have the capacity to exploit the majority of the host's nutrients, including

glycerol, FAs, and a number of other carbon-based substances, with the objective of

reprogramming immune cells.[57] Our findings have demonstrated that intracellular

bacteria are capable to actively influence host free FA metabolism (Figure 4A-4C) and

reprogram macrophages, thereby exacerbating neuroinvasion. The foamy-like

phenotype of macrophages is formed through dysregulation of lipid metabolism,

which in turn impairs the immune function of macrophages and contributes to

pathogenesis. For example, foamy macrophage necrosis has previously been observed

to expand tuberculous lesions in lung parenchyma, resulting in progressive

destruction of lung tissue and loss of lung function in patients with active disease.[58] In

this study, we discovered that BHB produced by a ketogenic diet promoted the

formation of a foamy-like phenotype of macrophages in peripheral blood, thereby

facilitating bacterial neuroinvasion. Interestingly, we found that serum cholesterol

levels, including TC and LDL-C were significantly higher in patients than in healthy

controls (Figure S11F and S11H). We hypothesize that cholesterol and other lipid

metabolites may also be involved in the formation of $CD36^+$ macrophages. However,

this hypothesis requires further experimental verification.

The biomechanical data indicated that $CD36^+$ macrophages exhibited a greater degree

of adhesion to the endothelial surface than normal macrophages, and this adhesion-

enhancing effect was more pronounced when endothelial cells were dense (Figure 3B

and 3C, S6B and S6C). Accordingly, we found that an elevation in $CD36^+$

macrophages in peripheral blood resulted in the spread of bacteria predominantly to

the brain, as opposed to other organs (Figure 2G and S3Q). This suggested that

$CD36^+$ macrophage-mediated bacterial invasion affected the brain to a greater extent

than other tissues. Interestingly, it was observed that a ketogenic diet and ketone ester

supplementation significantly increased bacterial load not only within the brain, but

also in other tissues such as the liver and lung (Figure 6E and 6J). This suggested that,

in addition to promoting the generation of $CD36^+$ macrophages, ketogenesis might

play a role in the intracellular bacterial infections that is currently unknown. Further

studies are therefore necessary to determine the role of ketogenesis in the formation of

CD36$^+$ macrophages and the contribution of ketogenesis to bacterial neuroinvasion.

The basal nutritional status of the host may also have an important influence on the

outcome of infection. In this study, we showed that neurolisteriosis mice experienced

a decrease in blood glucose levels and an increase of ketogenesis in the liver (Figure

S7A-S7H). We also found that during the course of bacterial neuroinvasion, mice

exhibited a reduction in body weight and diminished appetite (Figure S7M and S7N).

Interestingly, the fluctuations in glucose levels appeared to precede a decline in food

intake in neurolisteriosis mice (Figure S7H and S7N). In this context, the decrease in

glucose levels and increase in ketogenesis cannot be attributed exclusively to anorexia,

and further studies are therefore required to elucidate the underlying mechanisms of

the fluctuations in glycolysis and glucose levels during neuroinvasion.

Our findings suggest that the maintenance of optimal blood sugar levels may be a key

factor in the onset, progression, and prognosis of bacterial neuroinvasion. Several

recent reports provide support for this hypothesis. For example, glucose

supplementation has been shown to exert a protective effect against *Yersinia*

pseudotuberculosis infection in mice.[59] In addition, reduced BHB levels in patients

with sepsis have been demonstrated to diminish glucose production, so that the blood

glucose levels in non-survivor sepsis patients are likely to be considerably lower than

in the survivors. [60-62] Interestingly, another study by Wang et al. showed that glucose

supplementation was detrimental in bacterial sepsis. However, in their study, there

was no discernible difference in the kinetics of blood glucose following glucose

treatment [63], which further suggests the pivotal role of blood glucose levels in

bacterial infections. Consequently, we endeavored to replenish blood sugar levels in

mice afflicted with acute neuroinvasion via intraperitoneal administration of glucose,

and surprisingly found that glucose supplementation resulted in a marked alleviation

of the clinical symptoms, a reduction in the bacterial burden within the brain, and an

improvement in the survival rate of the mice (Figure S10). These findings highlight

the crucial role of nutritional therapy, in addition to antibacterial therapy, in the

clinical treatment of bacterial neuroinvasion. In addition, *L. monocytogenes* has been

shown to induce the accumulation of lipid droplets in macrophages, thereby

contributing to bacterial survival and evasion of innate immunity. [64] Given that

intracellular lipid is the key regulator of $CD36^+$ macrophage function, targeting this

lipid utilization pathway may be a promising therapeutic strategy for bacterial

neuroinvasion.

Overall, this study demonstrated that a BHB-rich environment, as created by the

ketogenic diet, rendered mice more susceptible to *L. monocytogene*-induced

neuroinvasion. From a mechanistic perspective, BHB protected $CD36^+$ macrophages

from the onset of senescence via HCAR2-mediated membrane repair. However, the

inhibition of HCAR2 signaling or calcium influx was observed to reverse the

improvement in CD36$^+$ macrophage survival, yet did not fully reverse the

exacerbation in bacterial neuroinvasion induced by BHB (Figure 5E-5G). These data

suggested that bacterial brain invasion induced by BHB is not solely dependent on

HCAR2 signaling, and that other potential mechanisms may also be involved in this

process. Nevertheless, this hypothesis requires further experimental confirmation.

Limitations of the study

The role of other cell clusters or FAs in bacterial neuroinvasion, beyond CD36$^+$

macrophages and BHB, remains to be elucidated. To circumvent the potential for

BHB to exert its effects through alternative target cell populations and organs, the

utilization of a HCAR2 conditional knockout mouse line is imperative. Moreover,

additional human neuroinvasion samples induced by intracellular bacteria other than

MTB should be further collected and analyzed.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No.

92369115, 82422048 and 92469110), the National Key Research and Development

Program of China (No. 2024YFC2310800 and 2021YFC2301405), the Fundamental

Research Funds for the Central Universities (No. 2024CDJXY-016), Natural Science

Foundation of Chongqing, China (No. CSTB2023NSCQMSX0402), and Chongqing

Talents: Exceptional Young Talents Project (No. cstc2021ycjh-bgzxm0099). The

funders had no role in study design, data collection and analysis, decision to publish,

or preparation of the manuscript.

Author Contributions

H.W., Q.Z., X.L. and Z.S. conceived and designed the study. Z.S., H.Y., H.F., Q.H.,

B.F., Y.X. and D.G. performed the animal experiments. Z.S., K.Y., Q.H., Q.L., Y.D.

and Xushuo Z. carried out healthy volunteer and patient recruitment and sample

collection. Z.S., N.L. and X.S. carried out the live-cell flow chamber assays. H.W.,

Z.S., K.Y., H.Y. and Xiaokai Z. collected and analyzed the data. X.T., S.F., J. L., R.Y.,

W.G., X.M. and M.L. contributed to experimental material and insightful suggestions.

H.W., X.L., Xushuo Z. and Z.S. wrote the manuscript. H.W. and Q.Z. coordinated the

project. All authors read and approved the final manuscript.

Declaration of interests

871　The authors declare no competing interests.

Figure Legends

Figure 1. Identification of CD36^{+} macrophage during neurolisteriosis.

(A) Summary of the experimental design (control mice, n=5; neurolisteriosis mice,

n=7).

(B) Uniform Manifold Approximation and Projection (UMAP) representation of

integrated transcriptome-based clustering. A total of 17,351 peripheral blood

mononuclear cells (PBMCs) were clustered and major immune cell lineages were

identified.

(C) Dot plot showing the expression of selected transcripts in immune cell clusters.

All transcripts were significantly enriched in the pertinent clusters with adjusted p <

0.05.

(D) t-distributed Stochastic Neighbor Embedding (tSNE) analysis of the re-clustered

cells corresponding to monocytes and macrophages (baseline, n=1,804;

neurolisteriosis, n=1,731).

(E) Expression of the indicated transcripts (encoding genes associated with lipid

accumulation) was projected onto a monocyte/macrophage tSNE plot.

(F-G) Mice was intraperitoneally injected with 2×10^7 CFU *L. monocytogenes* (n=5 for

each group). Flow cytometry was performed to assess the percentages of CD36$^+$

macrophages in PBMCs (F) and total macrophages (G).

(H) Correlation between CD36$^+$ macrophages and bacterial burdens in the brain

during neurolisteriosis. R, correlation coefficient.

(I) Immunofluorescence staining of infected CD36$^+$ macrophages in cerebral

vasculature of mice with neurolisteriosis. Infected CD36$^+$ macrophages were stained

with CD36 (red) and *L. monocytogenes* (green), while vascular endothelial cells were

stained with CD31 (white). Scale bar, 100 μm.

(J-K) FACS of CD36$^-$ and CD36$^+$ macrophages in the brain tissues (J). The

intracellular bacterial burdens were evaluated by flow cytometry (J) and plate dilution

method (K).

Experiments were performed in duplicate (F-K). Data are presented as mean ± SEM

[n=3 in (K), n=5 in (F, G)]. *p < 0.05, **p < 0.01. Abbreviations: PBMCs, peripheral

blood mononuclear cells; CD36$^+$ Mφs, CD36$^+$ macrophages; CFU, colony forming

unit.

Figure 2. Infected CD36$^+$ macrophages transfer bacteria into brain.

(A-D) (A) Schematic diagram of the *in vitro* BBB model based on Transwell system.

Macrophages were exposed to *L. monocytogenes* (MOI=10) for 6 hours and then

treated with 100 µg/ml gentamicin for 2 hours before being added into the upper

chamber. After 12 hours of penetration, neuron cells in the lower chamber were lysed

for the quantification of penetrating bacteria. (B) Quantification of penetrating

bacterial burden in the neuron cells in the lower chamber. (C) Representative images

of macrophages penetrating into the lower chamber. Scale bar, 100 µm. (D)

Quantification of the penetrating cells in each field of view.

(E-I) (E) Experimental diagram of adoptive macrophage transfer. Recipient mice were

anesthetized for irradiation (Co 60, 10 Gy). Twelve hours later, adoptive macrophages

transfer was performed. After recovery for another 12 hours, recipient mice were

intravenously injected with 1x10^6 CFU of *L. monocytogenes*. Gentamicin (30 mg/kg)

was administered intraperitoneally 4 hours after infection. Samples were collected for

examination at 24 hours post infection. (n=10 for each group) (F) Survival of

recipient mice. (G) Bacterial burdens in brains. (H-I) Pro-inflammatory factor levels

in CSF (H) and brain homogenates (I) of recipient mice.

(J-N) (J) Schematic diagram of adoptive macrophage transfer experiments in

PCSK9$^{+/+}$ (WT) and PCSK9$^{-/-}$ (KO) mice. Recipient mice were irradiated and

adoptive macrophages transfer was performed, and then mice were intravenously

infected with 1×10^6 CFU (K-L) or 1×10^5 CFU (M-N) of *L. monocytogenes*.

Gentamicin was administered intraperitoneally 4 hours after infection. (n=10 for each

group) (K, M) Bacterial burdens in the brains of recipient mice after intravenous

injection of 1×10^6 CFU *L. monocytogenes* 1 day post infection (K) and 1×10^5 CFU *L.*

monocytogenes 2 days post infection (M). (L, N) Survival of recipient mice after

intravenous injection of 1×10^6 CFU (L) and 1×10^5 CFU (N) *L. monocytogenes*.

Experiments were performed in duplicate (E-N) or triplicate (A-D). Data are

presented as mean ± SEM [n=3 in (B, D), n=5 in (G-I, K, M)]. *p < 0.05, **p < 0.01.

Abbreviations: ox-LDL, oxidized low density lipoprotein; FOV, field of view; NS,

normal saline; PMφ, peritoneal macrophages; CD36$^-$ Mφ, CD36$^-$ macrophages;

CD36$^+$ Mφ, CD36$^+$ macrophages; i.v., intravenously; i.p., intraperitoneally; ns,

nonsignificant.

Figure. 3 Characterization of infected CD36$^+$ macrophages.

(A) Schematic diagram of live-cell flow chamber experiment. Primary peritoneal

macrophages were pretreated with ox-LDL for 24 hours and then infected with *L.*

monocytogenes (MOI=10) for 6 hours. Then cells were perfused over pre-formed

endothelial monolayer for 8 minutes at a shear stress of 0.1 dyn/cm^2 and recorded at 2

dyn/cm^2 for 10 minutes unless otherwise indicated.

(B-C) (B) Optical images of ox-LDL pretreated macrophages adhering to the surface

of monolayer endothelial cells under shear flow. Scale bar, 200 μm. (C)

Quantification of adherent cells in each field of view.

(D) Macrophages resistance to shear flow after adhesion to endothelial cells.

Peritoneal macrophages were pretreated with ox-LDL for 24 hours and then

challenged with *L. monocytogenes* (MOI=10) for 6 hours. Upon 10-minute adherence

of equal macrophages to the monolayer endothelial cells at 1 dyn/cm^2, the shear stress

was stepwise increased to 2, 4, 8, 16, and 32 dyn/cm^2 for 30 seconds sequentially to

detach those adhered cells.

(E-H) Crawling dynamics of ox-LDL pretreated macrophages under shear flow. (E)

Typical trajectories of ox-LDL pretreated macrophages crawling on monolayer

endothelial cells. For each trajectory, the arrested sites of macrophages were overlaid

at the origin of the diagram (x/y=0/0) and the corresponding end points were indicated

with dots. Arrows indicate the flow direction. (F-H) Quantification of crawling

displacement (F), velocity (G) and directionality (H).

(I) Transmigration duration of ox-LDL pretreated macrophages after adhesion to the

surface of monolayer endothelial cells.

(J-K) Quantification of the number (J) and mean length (K) of protrusions in the

macrophages pretreated with ox-LDL.

(L) Quantification of cell area in the macrophages pretreated with ox-LDL.

(M) qRT-PCR analysis of relative expression of genes associated with intercellular

adhesion molecules.

Experiments were performed in triplicate. Data are presented as mean ± SEM [n=8 in

(C, I), n=10 in (F-H), n=18 in (L), n=20 in (K)]. *$p < 0.05$, **$p < 0.01$. Abbreviations:

ox-LDL, oxidized low density lipoprotein.

Figure 4. β-hydroxybutyrate aggravates bacterial neuroinvasion by promoting

the formation of CD36$^+$ macrophages via HCAR2 receptor.

(A-C) Lipid metabolomic profiling analyses of serum from mice with or without

neurolisteriosis (n=6 for each group). (A) Principal coordinates analysis (PCA)

analysis of differences between groups. (B) Expression of FAs in serum was shown in

the volcano plot. Differentially expressed FAs were selected with |log$_2$ fold change|>1

and adjusted $P < 0.05$. The dark blue dots represent down-regulated FAs and the

orange dots represent up-regulated FAs. (C) Classification of up-regulated FAs (SCFA,

short chain FAs; MCFA, medium chain FAs; LCFA, long chain FAs; VLCFA, very

long chain FAs) shown in the pie chart.

(D) Biochemical analysis of serum BHB levels in mice during neurolisteriosis.

(E-Q) Primary peritoneal macrophages were extracted from HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice in preparation for further functional analysis. HCAR2$^{+/+}$ and HCAR2$^{-/-}$

macrophages were pretreated with BHB (10 mM) for 24 hours and then infected with

L. monocytogenes (MOI=10) for 6 hours.

(E-F) Flow cytometry analysis (E) of BODIPY FL and normalized MFI (F) in BHB-

treated macrophages (0, 5, 10 mM).

(G) Percentage of CD36$^+$ macrophages in BHB-treated macrophages by flow

cytometry.

(H) Quantification of penetrating bacterial burden in the neuron cells in the lower

chamber of Transwell system. The macrophages stimulated with ox-LDL (80 ng/ml)

were used as positive control.

(I-J) Optical images (I) of adherent macrophages on the surface of monolayer

endothelial cells. Scale bar, 200 μm. Quantification of adherent cells in each field of

view (J).

(K) Macrophages resistance to shear flow after adhesion to endothelial cells.

(L-P) Crawling and migrating dynamics of infected macrophages under shear flow. (L)

Typical trajectories of infected macrophages crawling on monolayer endothelial cells.

Quantification of crawling displacement (M), velocity (N), directionality (O) and

transmigration duration (P).

(Q) MRNA expressions of intercellular α-integrin in infected macrophages.

Experiments were performed in duplicate (D) or triplicate (E-Q). Data are presented

as mean ± SEM [n=3 in (F-H, Q), n=10 in (J, M-P)]. *$p < 0.05$, **$p < 0.01$.

Abbreviations: ns, nonsignificant; FAs, fatty acids; BHB, β-hydroxybutyrate; MFI,

mean value of fluorescence intensity.

1007

Figure 5. β-hydroxybutyrate promotes the survival of macrophages via HCAR2-

mediated calcium influx.

(A-L) Primary peritoneal macrophages were extracted from HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice. Cells were then infected with *L. monocytogenes* (MOI=10) for 6 hours.

(A-C) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM)

before infection. (A) Quantification of the released LDH in cell culture medium. (B)

Immunofluorescence of propidium iodide (PI)-stained macrophages. Scale bar, 100

μm. (C) Quantification of the PI$^+$ cells after infection.

(D) Flow cytometry analysis of dynamics in intracellular calcium concentrations.

Infected macrophages were preloaded with Fluo-4 AM. BHB (10 mM) was added to

the buffer 30 seconds later.

(E-G) Macrophages were pretreated with BHB (10 mM) or BAPTA (10 μM) 24 hours

before infection. (E) Quantification of the released LDH in cell culture medium. (F)

Quantification of the PI$^+$ cells post infection. (G) Quantification of penetrating

bacterial burden in the neuron cells in the lower chamber of Transwell system.

(H-I) Infected macrophages were treated with BHB (10 mM). (H) Time-lapse images

of CHMP4B-EGFP in macrophages were collected every 5 minutes post BHB

treatment with confocal microscopy. (I) Quantification of CHMP4B-EGFP puncta.

(J-L) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) for

24 hours before infection. Flow cytometry analysis (J) and the normalized MFI (K) of

LAMP1 expression on the cell surface. (L) Quantification of the lysosomal enzyme

[acid phosphatase (ACP), β-galactosidase (β-gal) and β-hexosaminidase (β-hex)]

activities released into the culture medium.

Experiments were performed in triplicate. Data are presented as mean ± SEM (n=3).

*p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate;

LDH, lactate dehydrogenase.

1034

Figure 6. Ketogenesis exacerbates neurolisteriosis through β-hydroxybutyrate-

HCAR2 signaling.

(A-D) (A) Schematic diagram of KE supplementation in mice. C57BL/6 mice were

supplemented with KE (10, 20 mg/ml) in drinking water for 2 weeks and then

intraperitoneally challenged with 2×10^8 CFU *L. monocytogenes*. Gentamicin (30

mg/kg) was administered intraperitoneally 4 hours after infection. (n=15 for each

group) (B) Serum BHB levels. (C) Percentage of CD36$^+$ macrophages. (D) Survival

of mice exposed to *L. monocytogenes*. (E) Bacterial burdens in the organs.

(F-L) (F) Schematic diagram of ketogenic diet in mice. HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice

were treated with ketogenic or control diet for 4 weeks, followed by intravenous

infection of 1×10^7 CFU *L. monocytogenes*. Gentamicin (30 mg/kg) was administered

intraperitoneally 4 hours after infection. Samples were collected for examination 24

hours post infection. (n=10 for each group) (G) Survival of mice exposed to *L.*

monocytogenes. (H) Percentage of CD36$^+$ macrophages by flow cytometry. (I-J)

Bacterial burdens in PBMCs (I) and parenchymal organs (J). (K-L) Pro-inflammatory

factor levels in CSF (K) and brain homogenates (L). (M-O) Representative images (M)

of Evans blue leakage in mouse brains and HE staining in cerebral cortex tissues.

Scale bar, 10 μm. Quantification of pathological score (N) and Evans blue leakage (O).

Experiments were performed in duplicate. Data are presented as mean ± SEM (n=5).

*p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; KE, Ketone ester; GEN,

gentamicin; CD, control diet; KD, ketogenic diet; i.p., intraperitoneally; i.v.,

intravenously.

1057

Figure 7. β-hydroxybutyrate levels are increased during bacterial neuroinvasion

in humans.

(A) Schematic diagram of samples from healthy controls (n=60) and patients with

bacterial CNS infection (n=62). The patients were divided into three groups according

to serum BHB levels.

(B) Serum BHB levels in healthy controls and patients.

(C-I) Clinical scores (C), cell counts (D) and protein levels (E) in CSF. White blood

cells (WBC) (F), procalcitonin (PCT) (G), erythrocyte sedimentation rate (ESR) (H)

and C-reactive protein (CRP) (I) in the peripheral blood of patients with different

BHB levels.

(J-K) Dynamics of $CD36^+$ macrophages (J) and serum BHB (K) in 7 donors after

admission. The time after admission was divided into symptomatic and convalescent

periods according to clinical and laboratory indexes of infection. The scatter plot on

the right panel represents the statistics of AUC.

(L) Relative expression of genes involved in lipid metabolism in PBMCs from healthy

controls and patients was detected by qRT-PCR.

Data are presented as mean ± SEM. $*p < 0.05$, $**p < 0.01$. Abbreviations: ns,

nonsignificant; AUC, area under the curve; BHB, β-hydroxybutyrate.

1076

Figure S1. Acute bacterial neuroinvasion mouse model induced by _L._

monocytogenes.

(A) Mice were intravenously infected with 1×10^5, 1×10^6, or 1×10^7 CFU of *L.*

monocytogenes. Survival, bacterial burdens, and pro-inflammatory factor levels in the

CSF and brain homogenates were evaluated. (n=10 for each group)

(B) Mice were intraperitoneally infected with 2×10^6, 2×10^7, or 2×10^8 CFU of *L.*

monocytogenes. Survival, bacterial burdens, and pro-inflammatory factor levels in the

CSF and brain homogenates were evaluated. (n=10 for each group)

(C-E) Mice were intravenously infected with 1×10^5, 1×10^6, or 1×10^7 CFU of *L.*

monocytogenes. (C) Representative images of HE staining of the cerebral cortex

tissues and pathology scores. (D) Clinical scores. (E) Body weight changes.

(F-H) Mice were intravenously infected with 1×10^7 CFU of live, heat-killed or

attenuated (Δhly) *L. monocytogenes*. (n=10 for each group) (F) Bacterial burdens of

brains. (G) Survival rate. (H) Clinical scores.

(I-J) (I) Experimental diagram of mice treated with gentamicin. Mice were

intravenously infected with 1×10^7 CFU of *L. monocytogenes*. Gentamicin (30 mg/kg)

was administered to mice daily during the infection or 4 hours post-infection. (n=15

for each group) (J-K) Bacterial burdens in the peripheral blood (J) and brain tissues

(K). (L) Survival of the mice.

(M) Representative images and quantification of Evans blue leakage in mouse brains.

Evans blue assay was performed on mice challenged with *L. monocytogene*s (1×10^7

CFU intravenously) or LPS (2 mg/mouse).

Experiments were performed in duplicate. Data are presented as mean ± SEM (n=5).

*p < 0.05, **p < 0.01. Abbreviations: L. m, *L. monocytogene*; i.v., intravenous; i.p.,

intraperitoneal; hpi, hours post infection; GEN, gentamicin.

1102

Figure S2. Additional results of identification of CD36$^+$ macrophages during

neurolisteriosis, related to Figure 1.

(A) Cell clusters identified in 10× scRNA-seq dataset. The top 10 enriched genes for

each cluster are highlighted. Normalized z-score is used to illustrate gene expressions.

(B) Proportion of each cluster according to experimental conditions in Figure 1D.

(C) Gating strategy for analysis of CD36$^+$ macrophages in PBMCs.

(D-F) (D) FACS of the CD36$^+$ and CD36$^-$ macrophages from peripheral blood and

staining of BODIPY FL 493/503 and oil Red O. Scale bar, 10 μm. Quantification of

the normalized intake of Dil-oxLDL (E) and mRNA expressions of the scavenger

receptors (F) in sorted CD36$^+$ and CD36$^-$ macrophages.

(G) Quantification of the absolute numbers of CD36$^+$ macrophages in PBMCs

according to experimental conditions in Figure 1F.

(H-I) Quantification of the intracellular bacterial burdens of the CD36$^+$ and CD36$^-$

macrophages sorted from peripheral blood by flow cytometry (H) and plate dilution

method (I).

(J) Immunofluorescence staining of CD36$^+$ macrophages in cerebral vasculature of

mice with neurolisteriosis. CD36$^+$ macrophages were stained with CD11b (red) and

CD36 (green). The boundary of vascular was depicted by dashed lines. Scale bar, 100

μm.

(K-L) CD36$^-$ and CD36$^+$ macrophages were sorted from brain tissues of infected mice

according to experimental conditions in Figure 1J. (K) Percentages of CD36$^-$ and

CD36$^+$ macrophages in the brain. (L) Immunofluorescence staining of infected CD36$^-$

and CD36$^+$ macrophages. *L. monocytogenes* (green) and actin (red) were stained.

Scale bar, 10 μm.

(M) MRNA levels of the pro-inflammatory factors in control macrophages and sorted

CD36$^+$ macrophages.

Experiments were performed in duplicate (A-D, G-L) or triplicate (E-F, M). Data are

presented as mean ± SEM [n=3 in (E-F, M), n=5 in (G, I-K)]. *$p < 0.05$, **$p < 0.01$.

Abbreviations: PMφ, peritoneal macrophages; CD36$^-$ Mφ, CD36$^-$ macrophages;

CD36$^+$ Mφ, CD36$^+$ macrophages.

1133

Figure S3. Additional results of adoptive transfer of CD36$^+$ macrophages, related

to Figure 2.

(A) Mouse peritoneal macrophages were treated with different concentrations of ox-

LDL for 24 hours, and then the percentage of CD36$^+$ macrophages was assessed by

flow cytometry.

(B-E) (B) Schematic diagram of isolation of CD36$^+$ macrophages from two different

sources. (C) Immunoblot analysis of FABP4 and PPARG expressions. (D-E)

Transcriptomics analysis of CD36$^+$ macrophages from different sources. (n=3 for each

group) (D) Differentially expressed genes (DEGs) were shown in the volcano plot. (E)

PCA analysis of differences between groups.

(F-G) MRNA expression of pro-inflammatory (F) and anti-inflammatory (G) factors

in CD36$^+$ and control macrophages sorted from PBMCs.

(H-K) Experiments based on the *in vitro* BBB model shown in Figure 2A. (H) TEER

values of the BBB model before and after bacterial infection. (I) Quantification of the

intracellular bacterial burdens in macrophages in the upper chamber of Transwell

system. (J-K) Representative images (J) and quantification (K) of the bacteria inside

the macrophages that penetrating into the lower chamber. Scale bar, 10 μm.

(L-N) (L) Gating strategy for sorting CD36$^-$ lipid^{high} and CD36$^-$ lipid^{low} macrophages

in PBMCs. (M-N) Quantification of intracellular bacterial burdens in the upper

chamber (M) and penetrating bacterial burdens in the neuron cells in the lower

chamber of Transwell system (N).

(O-W) Additional results of adoptive macrophage transfer experiments shown in

Figure 2E. (O-T) Body weight changes (O). Clinical scores (P). Bacterial burdens in

other organs of recipient mice at 24 hours post infection (Q). Representative images

(R) of Evans blue leakage in mouse brains and HE staining in cerebral cortex tissues.

Quantification of extravasation (S) and pathology scores (T). Scale bar, 20 μm.

(U-W) After adoptive macrophage transfer, recipient mice were intravenously

infected with 1×10^5 CFU of *L. monocytogenes*. (n=10 for each group) Survival of

recipient mice (U). Bacterial burdens in brains (V) and other organs (W) of recipient

mice at 2 days post infection.

Experiments were performed in duplicate (O-W) or triplicate (A, F-N). Data are

presented as mean ± SEM [n=3 in (A, F-I, K, M-N), n=5 in (Q, S-T, V-W)]. *p < 0.05,

**p < 0.01. Abbreviations: ns, nonsignificant; NS, normal saline; PMφ, purified

peritoneal macrophages; CD36⁻ Mφ, CD36⁻ macrophages; CD36⁺ Mφ, CD36⁺

macrophages; ox-LDL, oxidized low densitylipoprotein; i.v., intravenous; i.p.,

intraperitoneal.

1170

Figure S4. Accumulation of CD36$^+$ macrophages exacerbates intracellular

bacterial neuroinvasion.

(A-H) Mice were intraperitoneally infected with 1×10^8 CFU of *S. Typhimurium* (A-B),

2×10^8 CFU of MRSA (C-D) or 1×10^8 CFU of *S. pneumoniae* (G-H), respectively.

(n=5 for each group) (A, C, G) Flow cytometry analysis of CD36$^+$ macrophages

during neuroinvasion. (B, D, H) Correlation between CD36$^+$ macrophages and

bacterial burdens in the brain during bacterial neuroinvasion. R, correlation

coefficient. (E-F) CD36$^-$ and CD36$^+$ macrophages were sorted from brain tissues of

mice infected with *S. Typhimurium* or MRSA. (E) Immunofluorescence staining of

CD36$^-$ and CD36$^+$ macrophages. Bacteria (green) and actin (red) in macrophages

were stained. Scale bar, 10 μm. (F) Quantification of the intracellular bacterial burden

of the CD36$^+$ and CD36$^-$ macrophages by plate dilution method.

(I-K) After adoptive macrophage transfer, recipient mice were intravenously infected

with 2×10^6 CFU of MRSA. Vancomycin (10 mg/kg) was administered

intraperitoneally 4 hours post infection. (n=10 for each group) (I) Survival of recipient

mice. (J) Bacterial burdens in tissues. (K) Pro-inflammatory factor levels in CSF of

recipient mice on day 1 post infection.

(L-N) After adoptive macrophage transfer, recipient mice were intravenously infected

with 1×10^6 CFU of *S. pneumoniae*. (n=10 for each group) (L) Survival rates of mice.

(M) Bacterial burdens in tissues. (N) Pro-inflammatory factor levels in CSF of

recipient mice on day 2 post infection.

Experiments were performed in duplicate. Data are presented as mean ± SEM (n=5).

*p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; CD36$^-$ macrophages; CD36$^+$

Mφ, CD36$^+$ macrophages.

1195

Figure S5. Additional results of bidirectional transfer of macrophages from

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice, related to Figure 2.

(A-B) Immunoblot (A) and qRT-PCR (B) analysis of expression of PCSK9 in CD36$^+$

macrophages.

(C) Flow cytometry analysis of lipid droplets in peritoneal macrophages of PCSK9$^{+/+}$

and PCSK9$^{-/-}$ mice.

(D) Immunoblot analysis of FABP4 and PPARG expressions.

(E) Intracellular bacterial burdens of infected macrophages. Peritoneal macrophages

of PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice were infected with *L. monocytogenes* (MOI=10) for 2

hours, and then cells were washed by PBS containing gentamicin. Intracellular

bacterial burdens were calculated at 2 or 12 hours post infection.

(F-I) PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice were intravenously infected with 1×10^7 CFU of *L.*

monocytogenes. (n=10 for each group) (F) Percentage of CD36$^+$ macrophages by flow

cytometry during neurolisteriosis. (G) Survival rates of mice. (H-I) Bacterial burdens

in the brain (H) and pro-inflammatory factor levels in CSF (I) of mice at 24 hours post

infection.

(J-L) Additional results of adoptive macrophage transfer experiments shown in Figure

2K-2L. (J) Bacterial burdens in other organs (lung, spleen, and liver). (K-L) Pro-

inflammatory factor levels in CSF (K) and brain homogenates (L) at day 1 post

infection.

(M-O) Additional results of adoptive macrophage transfer experiments shown in

Figure 2M-2N. (M) Bacterial burdens in other tissues (lung, spleen, and liver). (N-O)

Cytokine levels in CSF (N) and brain homogenates (O) on day 2 post infection.

(P-T) (P) Schematic diagram of adoptive macrophage transfer experiments. PCSK9$^{-/-}$

CD36^{+} and PCSK9$^{-/-}$CD36^{-} macrophages were sorted from the peritoneal

macrophages of PCSK9$^{-/-}$ mice, and then injected into irradiated recipient mice.

Gentamicin (30 mg/kg) was administered intraperitoneally 4 hours after infection.

(n=10 for each group) (Q-R) Bacterial burdens in the brain (Q) and other organs (lung,

spleen, and liver) (R). (S) Survival of recipient mice. (T) Pro-inflammatory factor

levels in the CSF of recipient mice at 24 hours post infection.

Experiments were performed in duplicate (F-T) or triplicate (A-E). Data are presented

as mean ± SEM [n=3 in (B-C, E), n=5 in (F, H-O, Q-R, T)]. *p < 0.05, **p < 0.01.

Abbreviations: ns, nonsignificant; i.p., intraperitoneally; dpi, days post infection;

CD36^{-} macrophages; CD36^{+} Mφ, CD36^{+} macrophages.

1230

Figure S6. Supplemental data for the live-cell flow chamber assays, related to

Figure 3.

(A) Immunofluorescence staining of infected macrophages. Primary peritoneal

macrophages were pretreated with ox-LDL for 24 hours and then infected with *L.*

monocytogenes (MOI=10) for 6 hours. Intracellular bacteria (green) in macrophages

were stained afterwards. Scale bar, 10 μm.

(B-E) Optical images of *L. monocytogenes*-infected macrophages on the surface of

endothelial monolayers with different densities (B) or endothelial monolayers derived

from different tissues (D). Scale bar, 200 μm. (C, E) Quantification of adherent cells

in each field of view.

(F) Schematic trajectory used for defining macrophage crawling parameters.

(G) Crawling duration of *L. monocytogenes*-infected macrophages after adhesion to

endothelial monolayer surface.

(H) Relative fluorescence intensity of a macrophage on endothelial surface over time.

It is a random example to illustrate the definition of transmigration duration. A 70%

decrease in fluorescence intensity after a macrophage crosses a monolayer is

considered successful transmigration.

(I) Representative images for quantifying the cell area and protrusions of

macrophages.

(J-K) Flow cytometry analysis of intercellular adhesion molecules in peritoneal

macrophages (J) and sorted CD36$^-$ and CD36$^+$ macrophages (K).

(L-S) Primary peritoneal macrophages were pretreated with ox-LDL for 24 hours and

then infected with *L. monocytogenes* (MOI=10) for 6 hours. ICAM1 antibody (50

μg/mL) blockade was performed for 30 minutes prior to ox-LDL stimulation. (L)

Optical images of adherent macrophages on the surface of monolayer endothelial cells.

Scale bar, 200 μm. (M) Quantification of adherent cells in each field of view. (N)

Macrophages resistance to shear flow after adhesion to endothelial cells. (O-S)

Crawling and migrating dynamics of infected macrophages under shear flow. (O)

Typical trajectories of infected macrophages crawling on monolayer endothelial cells.

Quantification of crawling displacement (P), velocity (Q), directionality (R) and

transmigration duration (S).

Experiments were performed in triplicate. Data are presented as mean ± SEM [n=8 in

(C, E, G), n=5 in (M, P-S)]. *$p < 0.05$, **$p < 0.01$. Abbreviations: BMEC, brain

microvascular endothelial cell; PVEC, pulmonary vascular endothelial cell; LSEC,

liver sinusoidal endothelial cell; TEM, transendothelial migration; ns, nonsignificant.

1266

Figure S7. Additional results of β-hydroxybutyrate promote bacterial

neuroinvasion, related to Figure 4.

(A) KEGG enrichment analysis of lipid metabolites. The horizontal axis represents

the number of -$\log_{10}$(p-Value), and the dashed line represents p-Value=0.05. Five of

the top six enrichment pathways are above the significance level, namely

Sphingolipid metabolism (Sphingosine, Sphinganine, Phytosphingosine. p-Value=

0.0005), Ubiquinone and other terpenoid-quinone biosynthesis (Vitamin K2, Vitamin

K1. p-Value= 0.002), Butanoate metabolism [3-hydroxybutyrate (BHB), Succinic

acid. p-Value=0.006], Synthesis and degradation of ketone bodies [3-hydroxybutyrate

(BHB), Acetyl coenzyme A. p-Value=0.04], Fatty acid biosynthesis (Dodecanoic acid.

p-Value=0.04).

(B) Mass spectrometry analysis of serum BHB levels.

(C-N) Mice in the mild infection group were intraperitoneally infected with 2×10^6

CFU, whereas the neurolisteriosis group was intraperitoneally infected with 2×10^8

CFU of *L. monocytogenes*. (n=5 for each group)

(C-D) Bacterial burdens in the organs (C) and percentages of CD36$^+$ macrophage (D)

were detected at 24 hours post infection.

(E-G) Transcriptomics analysis of liver tissues from mice (n=3 for each group). (E)

Heatmap of differentially expressed (DE) genes related to ketogenesis. (F) Top 20

KEGG enrichment analysis of function of DE genes in neurolisteriosis group and

baseline group. (G) Gene-set enrichment analysis for glycolysis/gluconeogenesis in

neurolisteriosis group and baseline group.

(H) Biochemical analysis of serum glucose levels in mice during *L. monocytogenes*

infection.

(I-L) Levels of ATP generation (I), ADP/ATP ratios (J), seahorse analysis of the

ECAR (K) and relative abundances of lactate (L) in liver tissues of mice were

detected at 24 hours (unless otherwise indicated) post infection.

(M-N) Changes in body weight (M) and food intake (N) of the mice post infection.

(O-P) Biochemical analysis of serum BHB and glucose levels in mice

intraperitoneally infected with 1×10^8 CFU of *S. typhimurium* (O) and 2×10^8 CFU of

MRSA (P). (n=5 for each group)

(Q) PBMCs were isolated form HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice, and then treated with

BHB (10 mM) for 24 hours. Percentages of CD36^{+} macrophages were analyzed by

flow cytometry.

(R-T) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) or

cAMP (1 mM) for 24 hours and then infected with *L. monocytogenes* (MOI=10) for 6

hours. HSL activity (R), normalized MFI of BODIPY (S) and the penetrating bacterial

burden in the lower chamber of Transwell system (T) were quantified.

Experiments were performed in duplicate (C-P) or triplicate (Q-T). Data are presented

as mean $\pm$ SEM [n=3 in (Q-T), n=5 in (B-D, H-P)]. *$p < 0.05$, **$p < 0.01$.

Abbreviations: BL, baseline; NL, neurolisteriosis; ns, nonsignificant; MFI, mean

fluorescence intensity; HSL, hormone-sensitive lipase.

1309

Figure S8. Additional results on the role of β-hydroxybutyrate in regulating

macrophage survival, related to Figure 5.

(A) Immunofluorescence staining of infected macrophages. The HCAR2$^{+/+}$ and

HCAR2$^{-/-}$ macrophages were isolated and infected with *L. monocytogenes* (MOI=10)

for 6 hours, and then cells were washed by PBS containing gentamicin. Intracellular

bacteria (green) in macrophages were stained afterwards. Scale bar, 10 μm.

(B-G) Primary peritoneal macrophages were extracted from HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice and then treated with BHB (10 mM) or BAPTA (10 μM) 24 hours before

infection (MOI=10, 6 hours). (B) Intracellular bacterial burdens of macrophages in

the upper chamber of Transwell system. (C) Time-lapse images of CHMP4B-EGFP in

macrophages were collected every 5 minutes post BHB treatment with confocal

microscopy. (D) Quantification of CHMP4B-EGFP puncta. Flow cytometry analysis

(E) and normalized MFI (F) of LAMP1 expression on the cell surface. (G)

Quantification of the activities of lysosomal enzymes released into the culture

medium.

(H-J) Resident peritoneal macrophages were isolated and then infected with *L.*

monocytogenes (MOI=10) for 6 hours. Percentage of PI$^+$ cells (H) and the normalized

MFI of LAMP1 expression on the cell surface (I) were assessed by flow cytometry. (J)

Quantification of the activities of lysosomal enzymes released into the culture

medium.

(K) Primary peritoneal macrophages were isolated from HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice and then infected with *L. monocytogenes* (MOI=10) for 6 hours. The

colocalization of BODIPY (green) and LC3 (red) in infected macrophages was

examined by immunofluorescence. Scale bar, 10 μm.

(L) Percentage of CD36$^+$PI$^+$ cells at 24 hours post infection was quantified by flow

cytometry. CD36$^+$ macrophages were treated with BHB (10 mM) 24 hours before

infection. Baf-A1 (100 nM) was added into the culture medium 6 hours before BHB

stimulation.

Experiments were performed in triplicate. Data are presented as mean ± SEM (n=3).

*p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate;

MFI, mean fluorescence intensity; rPMφs, resident peritoneal macrophages.

1341

Figure S9. Additional results of ketogenesis increases bacterial neuroinvasion,

related to Figure 6.

(A) Clinical scores of mice supplemented with KE as shown in Figure 6A.

(B-C) Clinical scores (B) and changes in body weight (C) of mice in the ketogenic

diet experiments as shown in Figure 6F.

(D-G) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice were treated with ketogenic or control diets,

respectively, followed by intraperitoneal infection of 2×10^8 CFU *L. monocytogenes*.

(n=10 for each group) (D) Survival of mice exposed to *L. monocytogenes*. (E)

Bacterial burdens in the organs. Pro-inflammatory factor levels in CSF (F) and brain

homogenates (G).

(H-J) MRNA expressions of inflammatory factors in control macrophages and CD36^{+}

macrophages sorted from peripheral blood of mice with ketogenic diets (H) or with

BHB (10 mM) pretreatment (I-J).

(K-N) (K) Schematic diagram of adoptive macrophage transfer experiments.

Peritoneal macrophages isolated from HCAR2$^{-/-}$ and HCAR2$^{+/+}$ mice with ketogenic

or control diets were injected into irradiated recipient mice. After recovery for 12

hours, recipient mice were intravenously injected with 1×10^6 CFU of *L.*

monocytogenes. Gentamicin (30 mg/kg) was administered intraperitoneally 4 hours

after infection. (n=10 for each group) (L) Survival of recipient mice. (M) Bacterial

burdens in organs. (N) Pro-inflammatory factor levels of recipient mice at 24 hours

post infection.

(O-T) (O) Anabolic mechanism of BHB. (P-T) HMGCS2$^{+/+}$ and HMGCS2$^{-/-}$ mice

were intravenously infected with 1×10^7 CFU of *L. monocytogenes*. Samples were

collected for examination 24 hours post infection. (n=10 for each group) (P) Serum

BHB levels. (Q) Percentages of CD36$^+$ macrophages. (R-T) Survival rates of mice (R),

bacterial burdens in the brain (S) and pro-inflammatory factor levels in the CSF (T) of

infected mice.

(U-W) Mice were treated with ketogenic or control diets, respectively, followed by

intraperitoneal infection of 2×10^8 CFU MRSA. (n=15 for each group) (U) Survival of

mice exposed to MRSA. (V) Bacterial burdens in the organs (brain, lung, spleen, and

liver). (W) Pro-inflammatory factor levels in the CSF.

Experiments were performed in duplicate (A-G, K-W) or triplicate (H-J). Data are

presented as mean ± SEM [n=3 in (H-J), n=5 in (A-C, E-G, M-N, P-Q, R-T, U-V)]. *p

< 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; i.v., intravenous; i.p.,

intraperitoneal; PMφ, peritoneal macrophages; ctrl, control CD36$^+$ Mφ, CD36$^+$

macrophages; GEN, gentamicin; CD, control diet; KD, ketogenic diet.

1378

Figure S10. Increased glucose levels alleviate bacterial neuroinvasion.

(A-M) (A) Schematic diagram of *in vivo* glucose supplementation. C57BL/6 mice

were intraperitoneally challenged with 2×10^8 CFU *L. monocytogenes*, and glucose

(100 mg/kg) was injected 1 hour post infection, with repeated injections every 12

hours. Samples were collected at 48 hours (unless otherwise indicated) post infection.

(n=15 for each group) (B) Serum glucose levels. (C-F) Levels of ATP generation (C),

ADP/ATP ratios (D), relative abundances of lactate (E) and seahorse analysis of the

ECAR (F) in liver tissues of mice at 48 hours post infection. (G) BHB levels. (H)

Percentages of CD36$^+$ macrophages in the PBMCs of infected mice. (I) Survival of

mice exposed to *L. monocytogenes* with or without glucose supplementation. (J)

Clinical scores. (K) Bacterial burdens in parenchymal organs. (L-M) Pro-

inflammatory factor levels in CSF (L) and brain homogenates (M) of mice at 48 hours

post infection.

Experiments were performed in duplicate. Data are presented as mean ± SEM (n=5).

*p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; GEN, gentamicin; GLU,

glucose.

1395

Figure S11. Additional results of high levels of β-hydroxybutyrate associate with

bacterial neuroinvasion in humans, related to Figure 7.

(A) Risk factors for patients with bacterial CNS infection in the MIMIC IV database.

(B-E) Glucose (B) and insulin (C) levels in the serum from recruited healthy controls

and bacterial neuroinvasion patients. Correlation between serum glucose levels and

BHB levels (D), and correlation between serum insulin levels and BHB levels (E) in

bacterial neuroinvasion patients were calculated. R, correlation coefficient.

(F-H) Serum triglyceride (TC) (F), high density lipoprotein (HDL-C) (G), and low

density lipoprotein (LDL-C) (H) from healthy controls and patients.

(I) The intracellular bacterial burdens of CD36$^+$ and CD36$^-$ macrophages sorted from

peripheral blood of patients by were examined by flow cytometry.

Data are presented as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns,

nonsignificant.

1409

1410

1411   **Materials and methods**

1412   **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Human/Mouse CD11b, FITC	Multi Sciences	Cat# F11011B01
Anti-Mouse F4/80, PE	Multi Sciences	Cat# F21480A02
Anti-Mouse CD36, APC	eBioscience	Cat# 17-0362-82
Anti-Mouse ITGA2, FITC	eBioscience	Cat# 11-5971-82
Anti-Mouse ITGB1, FITC	eBioscience	Cat# 11-0291-82
Anti-Mouse ICAM1, PE	eBioscience	Cat# 11-0541-82
Anti-Mouse CD107a, FITC	Proteintech	Cat# 65050-1-Ig
Anti-Mouse CD36	CST	Cat# 28109
Anti-Mouse CD31	CST	Cat# 15585
Anti-Mouse PPARG	CST	Cat# 2435
Anti-Mouse β-Actin	CST	Cat# 4970
Anti-Mouse IgG, HRP	CST	Cat# 7076
Anti-rabbit IgG, HRP	CST	Cat# 7074
Anti-Listeria monocytogenes	This paper	N/A
Anti-Mouse FABP4	Abcam	Cat# ab92501
Anti-Mouse PCSK9	Abcam	Cat# ab185194
Anti- Mouse CD11b	Abcam	Cat# ab8878
Anti-Mouse LC3B	Abcam	Cat# ab192890
Anti-ICAM1	Abcam	Cat# ab ab222736
Anti-Mouse α-TUBULIN	Santa Cruz	Cat# sc-398103
Bacterial strains		
Listeria monocytogenes	ATCC	Cat# 19115
MRSA USA300	ATCC	Cat# TCH1516
Salmonella Typhimurium	ATCC	Cat# SL1438
Streptococcus pneumoniae	ATCC	Cat# 49619
Chemicals and Reagents		
Thioglycollate medium	Sigma-Aldrich	Cat# 9032
Collagenase IV	Sigma-Aldrich	Cat# C4-BIOC
Oxidized Low Density	Invitrogen	Cat# L34357

Lipoprotein (ox-LDL)		
DiI-OxLDL	Invitrogen	Cat# L34358
DNase I	Invitrogen	Cat# 18047019
Hoechst 34580	Invitrogen	Cat# H21486
Propidium iodide	Invitrogen	Cat# P1304MP
Fluo-4 AM	Invitrogen	Cat# F14217
Lipofectamine 2000 Transfection Reagent	Invitrogen	Cat# 11668030
Dulbecco's modified Eagle's medium (DMEM)	GIBCO	Cat# C11965500BT
Fetal bovine serum (FBS)	GIBCO	Cat# 10091148
Penicillin Streptomycin (Pen Strep)	GIBCO	Cat# 15140122
Collagenase II	GIBCO	Cat# 17101015
Phalloidin-Alexa 647	Thermo Fisher	Cat# A22287
BODIPY 493/503	MedChemExpress	Cat# HY-W090090
Gentamycin	MedChemExpress	Cat# HY-A0276A
Vancomycin	MedChemExpress	Cat# HY-B0671
BAPTA-AM	MedChemExpress	Cat# HY-100545
Evans Blue	MedChemExpress	Cat# HY-B1102
BHI broth	MERCK	Cat# 53286
Puromycin	InvivoGen	Cat# ant-pr
Critical commercial assays		
Peripheral blood mononuclear Cell Isolation Kit	Solarbio	Cat# P8680
Pierce Primary Neuron Isolation Kit	Thermo Fisher	Cat# 88280
Mouse IL-6 Uncoated ELISA Kit	Thermo Fisher	Cat# 88-7064-77
Mouse TNF-α Uncoated ELISA Kit	Thermo Fisher	Cat# 88-7324-88
CyQUANT LDH Cytotoxicity Assay Kit	Thermo Fisher	Cat# C20301
β-galactosidase Assay Kit	Thermo Fisher	Cat# 75707
Glucose Colorimetric Detection Kit	Thermo Fisher	Cat# EIAGLUC
Acid Phosphatase Assay	Abcam	Cat# ab83367

β-hexosaminidase Assay Kit	Nanjing Jiancheng Bioengineering Institute	Cat# H485-1-2
ATP Assay Kit	Sigma-Aldrich	Cat# MAK473
ADP/ATP Ratio Assay Kit	Sigma-Aldrich	Cat# MAK135
Lactate Assay Kit	Sigma-Aldrich	Cat# MAK570
HSL ELISA Kit	Mlbio	Cat# ml026143
Hematoxylin and Eosin Staining Kit	Beyotime	Cat# C0105S
Oil Red O Staining Kit	Beyotime	Cat# C0157
Non-endotoxin plasmid small extraction medium dose kit	TIANGEN	Cat# DP118
HiScript II 1st Strand cDNA Synthesis Kit	Vazyme	Cat# R212
XF Glycolysis Stress Test kit	Seahorse Biosciences	Cat# 103020-100
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Charles River	Cat# 219
Recombinant DNA		
pLV3-CMV-Chmp4b(mouse)-3×FLAG-CopGFP-Puro	Miaoling Biology	Cat# P44204
Software and algorithms		
GraphPad Prism 9	GraphPad Software	https://www.graphpad.com
ImageJ	NIH	https://imagej.nih.gov/ij/
FlowJo V10	FlowJo LLC	https://www.flowjo.com
R studio	r-project.org	
Other		
Ketone ester	Guangao Biotechnology	Cat# GA8901
Ketogenic diet (KD)	Xietong Pharma	Cat# XTKD01

1413

Clinical samples

1414

Patients diagnosed with bacterial CNS infection (n = 62, 32 males and 30 females,

aged 41.62 ± 15.68 years) and healthy donors (n = 60, 30 males and 30 females, aged

38.32 ± 9.17 years) were recruited from Chongqing Public Health Medical Center.

Patients were all exposed to *Mycobacterium tuberculosis* without immunosuppression.

Patients were gender- and age-matched and were not concurrently infected with other

pathogens, such as influenza or HIV. Meanwhile, subjects were recruited without any

metabolic disorders, such as obesity or diabetes, in order to exclude their direct effects

on BHB and $CD36^+$ macrophages levels. Since blood glucose is a risk factor for

bacterial neuroinvasion, subjects with abnormal blood glucose or insulin

concentrations were also excluded. Clinical symptoms exhibited by the patients

included a state of exhaustion, headaches, irritability, vomiting, fever, neck stiffness,

convulsions, and consciousness disorder. Each symptom was assigned a value of one

point. The clinical data of the patients were collected within 12 hours of admission.

The entire process of seven hospitalized patients from admission to recovery and

discharge was followed up. Blood samples were collected and $CD36^+$ macrophages

and BHB levels were detected in all seven patients within 12 hours of admission, and

changes in the $CD36^+$ macrophages and BHB levels were measured continuously

throughout the hospitalization period. According to the medical records, the entire

hospitalization period was divided into a symptomatic period and a convalescent

period, based on the recovery time points at which indicators of CNS infection

(including white blood cell counts, protein levels and bacterial counts in the

cerebrospinal fluid), and clinical symptoms of the patients returned to normal. All data

were normalized to the parameters of first detection on admission.

Isolation of PBMCs and single-cell suspensions was performed by density gradient

centrifugation using PBMC Isolation Kit. Plasma from each patient and donor was

collected and then used for quantification of BHB. This study was reviewed and

approved by the Ethics Committee at the Chongqing Public Health Medical Center,

and was compliant with the "Guidance of the Ministry of Science and Technology

(MOST) for the Review and Approval of Human Genetic Resources". Healthy

individuals and patients providing blood samples were given informed consent. The

ethics committee approved this consent procedure.

The data in retrospective study were obtained from the Medical Information Mart for

Intensive Care IV (MIMIC-IV) version 1.0

(https://physionet.org/content/mimiciv/1.0/), which includes medical information on

patients in the intensive care unit of a hospital in Boston, USA, over a 10-year study

period. The database was established with the approval of the Massachusetts Institute

of Technology (Cambridge, MA) and the Institutional Review Boards of Beth Israel

Deaconess Medical Center (Boston, MA). Data were extracted from the MIMIC-IV

database using Structured Query Language (SQL) with Navicat Premium (version

12.0.28). A total of 128 admission records (49 patients /79 controls) from the MIMIC

IV database were included in this study. Patients with benign intracranial hypertension

were considered as controls.

1457

Animals

Animal experimental procedures were approved by the Laboratory Animal Welfare

and Ethics Committee of Chongqing University. All mice were housed in a specific

pathogen-free (SPF) facility according to standard humane animal husbandry

protocols. In all experiments, experimental and control mice were 6-8-week-old

littermates of C57BL/6 mice. HCAR2$^{-/-}$, PCSK9$^{-/-}$, HMGCS2$^{-/-}$ mice and littermate

controls obtained from heterozygote crosses were used for all experiments. Each in

vivo experiment was performed using 50% female and 50% male animals, and results

were not expected to be influenced by sex. All mice were maintained under a 12-hour

light-dark cycle with ad libitum access to regular food and water.

1468

Bacterial strains

All procedures related to pathogenic bacteria were conducted in accordance with

Biosafety Level 2 protocols and guidelines. The *L. monocytogenes* (ATCC 19115) and

S. pneumoniae (ATCC 49619) were grown in Brain Heart Infusion Broth. The

Methicillin-resistant *Staphylococcus aureus* USA300 (ATCC TCH1516) and *S.*

Typhimurium (ATCC SL1438) were grown in Luria-Bertani Broth. Bacteria were

grown at 37℃ with 5% CO_2 to mid-log phase, and growth was evaluated by

monitoring at an optical density of 600 nm. Frozen stocks of bacteria in 20% glycerol

were prepared and kept at -80℃ until use.

1478

Data and code availability

Single-cell RNA-seq data (accession number GEO: GSE264020, secure token for

reviewer: stchicgorhozbkb) and RNA-seq data (accession number GEO: GSE273820,

secure token for reviewer: gbohyqeanrmzhyn; accession number GEO: GSE283713,

secure token for reviewer: gbujwcoejdchtwz) have been deposited in GEO and are

publicly available as of the date of publication.

1485

Mouse model of acute neuroinvasion

Mice were injected intravenously of 1×10^7 CFU or intraperitoneally of 2×10^8 CFU *L.*

monocytogenes to establish an acute neuroinvasion mouse model. Gentamicin (30

mg/kg) was administered intraperitoneally 4 hours post infection to eliminate

circulating bacteria that did not enter the phagocytes. Mice challenged with LPS (2

mg/mouse) served as positive controls for BBB damage. For other intracellular

bacteria induced neuroinvasion models, mice were infected with MRSA (2×10^7 CFU

intravenously or 2×10^8 CFU intraperitoneally), *S. Typhimurium* (1×10^7 CFU

intravenously or 1×10^8 CFU intraperitoneally), or *S. pneumonia* (1×10^7 CFU

intravenously or 1×10^8 CFU intraperitoneally), respectively. In the adoptive

macrophages transfer experiments, bone marrow irradiation-based

immunosuppression resulted in mice being more susceptible to bacterial infection,

which necessitated a reduction of the infectious dose. After recovery from transfer for

12 hours, recipient mice were challenged with *L. monocytogenes* (1×10^6 CFU

intravenously), MRSA (2×10^6 CFU intravenously), or *S. pneumonia* (1×10^6 CFU

intravenously) unless otherwise indicated.

The progression of acute bacterial meningitis was determined by clinical scoring as

previously described with adjustments.[65] Clinical scoring was performed based on the

Table S2, including activity (0-2), time to return to upright position (0-2), eyes (0-2)

and neurological examination (0-2). Mice were deeply anaesthetized by

intraperitoneal injection of tribromoethanol solution (Avertin, 500 mg/kg), and

perfused transcranially with saline solution for brain tissue pathomorphology. Blood

was sampled immediately before perfusion. Tissue samples, including brain, lung,

liver, and spleen, were dissected and weighed.

1510

Adoptive macrophage transfer experiments

Primary peritoneal macrophages from donor mice were collected 72 hours after

intraperitoneal injection of thioglycolate. To collect CD36$^+$ macrophages, purified

macrophages were stimulated with 80 μg/ml ox-LDL for 24 hours and then sorted by

an APC-conjugated CD36 antibody. Recipient mice were weighed 25-28 g and

anesthetized for irradiation (Co 60, 10 Gy). 12 hours later, adoptive macrophages

transfer was performed. After recovery for another 12 hours, recipient mice were

challenged with bacteria (see mouse model of acute neuroinvasion section for

infectious routes and doses), and extracellular bacteria were eliminated by gentamicin

(30 mg/kg) or vancomycin (10 mg/kg) treatment 4 hours post infection. Samples were

collected at 24 hours post infection for subsequent examinations unless otherwise

indicated.

1523

Mouse diet Study and glucose supplementation

Mice were administered a supplement of 20 mg/ml ketone ester (D-β-

hydroxybutyrate-(R)-1,3 butanediol monoester) in the drinking water for 4 weeks

before the infection. The ketogenic diet contained 10% protein and 90% fat, whereas

the control diet contained 23.2% protein, 2.8% fat and 74% sugar. Mice were placed

on either the ketogenic or control diet for an additional 4 weeks before infection.

The food consumption was estimated by a subtraction method with the automated

weighing scales as previously reported.[66] A computer analyzed the changes in weight

of each cage and outputs data on food intake by blocks of hours. The value of

consumption per mouse was the amount of food reduced per cage divided by the

number of mice per cage.

In the glucose supplementation experiment, mice were injected intraperitoneally with

saline or glucose (100 mg/kg) every 12 hours after infection. To measure blood

glucose levels, a drop of blood was collected by snipping the very end of the tail, and

the blood in the strip was analyzed using a glucometer to measure blood glucose

levels.

1540

Analysis of bacterial load in cells or tissues

Tissue samples of mice were weighted and then transferred into 2 ml tubes containing

4 mm stainless steel beads and 1 ml of ice-cold sterile saline. Homogenization was

carried out in TissueLyser II (Qiagen) for 5 minutes at 70 Hz. Serial dilutions of the

homogenates were plated on the plates incubated at 37℃ and CFUs were counted.

The infected macrophages were incubated in fresh solution containing 5 μg/mL

lysostaphin for the indicated periods and then lysed by 0.2% Triton to release

intracellular bacteria. The lysates were diluted to appropriate dilutions with sterile

PBS for CFU count on agar plates.

1550

BBB permeability assays

After infection, mice were retro-orbitally injected with 2% Evans blue. After 2 hours

of circulation, mice were deeply anaesthetized, and the brains were delicately

dissected after transcardial PBS perfusion. Whole-brain images were acquired using a

digital camera. To quantify Evans blue leakage, mouse brains were sectioned

longitudinally, weighted and then homogenized in 0.5 ml of 50% trichloroacetic acid.

Lysates were centrifuged at 15,000 g for 30 minutes. The amount of extravasated

Evans blue in the supernatant was measured at 620/680 nm emission using a Multi-

Mode Microplate Reader (SpetraMax i3x), calculated by a standard curve, and

normalized to the weight of each brain.

1561

Brain histopathology and Immunostaining

Brain samples were collected from infected mice 24 hours after injection. Mice were

deeply anaesthetized with tribromoethanol solution (Avertin, 500 mg/kg,

intravenously), followed by transcranial perfusion with 30 ml of saline solution, and

then of 4% Paraformaldehyde (PFA) in 10 ml of saline. After dissection, brain

samples were maintained in 4% PFA at 4℃. Fixed brains were embedded in paraffin,

sectioned and half of the slides were stained using haematoxylin and eosin. Each field

was assigned a score from 0 to 3 based on neuronal morphology: grade 0 (no

alteration); grade 1 (no vacuolation with small numbers of pyknotic cells); grade 2

(moderate vacuolation and pyknosis); and grade 3 (extensive vacuolation, pyknosis

and tissue loss or liquefactive necrosis).

Fresh frozen brain tissue sections of 5-10 mm thickness were mounted on slides, air

dried for 10 minutes at room temperature, and fixed in ice cold 10% neutral-buffered

formalin for 5-10 minutes. Slices were washed in PBS, incubated in 3% bovine serum

albumin (BSA) for 2 hours, and then washed with 0.1% Triton X-100 in PBS. The

slides were labeled with the appropriate primary antibodies overnight at 4℃ in mild

blocking conditions. Then next day, the slices were washed in PBS, and then

incubated with secondary antibodies and DAPI. Tissues were washed in PBS and then

mounted on glass slides under coverslips in mounting medium. The slides were left in

darkness overnight before observation under a Zeiss LSM900 or LM710 microscope.

1582

Cell sorting and flow cytometry

Isolated mouse PBMCs were washed twice with PBS. The proportion of CD36$^+$

macrophages population was determined by adding appropriate amount of

fluorescent-coupled primary antibodies (F4/80-PE, CD11b-FITC, CD36-APC). The

cells were incubated at 4℃ for 30 minutes away from light. The living macrophages

were collected according to the expression of CD36 and then cultured overnight *in*

vitro. The lipid content in sorting cells was detected using BODIPY FL and oil Red O.

To evaluate the lipid uptake capacity, macrophages were incubated with 40 µg/ml DiI-

ox-LDL particles for 6 hours, and then the fluorescence intensity was measured by

flow cytometry. To detect the expression of LAMP1 on the cell surface, infected

macrophages were washed twice with PBS and then stained with LAMP1-FITC at 4℃

for 30 minutes away from light. The cells were washed three times and then re-

suspended in 200μl of the staining buffer for final FACS analysis. Data was analyzed

with FlowJo software.

1597

Isolation and culture of primary cells

To generate mouse peritoneal macrophages, 8-week-old mice were injected

intraperitoneally with 3% thioglycolate broth. After 72 hours, peritoneal cells were

harvested and macrophages were enriched by centrifuge. To obtain primary mouse

brain microvascular endothelial cells (BMECs), the cerebral cortex in the brains were

separated and rolled on the filter paper to remove the meninges and choroid plexuses.

A mechanical reduction was performed by mincing the brain pieces using surgical

blades. The homogenate was centrifuged (800 g, 8 minutes, 4℃) and the precipitate

was digested in DMEM containing 1 mg/ml collagenase II and 58.5 U/ml DNase I for

75 minutes on a benchtop orbital shaker at 37℃. The supernatant was collected and

centrifuged at 500 g for 8 minutes. The isolated cerebrovascular endothelial cells were

purified by the anchoring technique. The livers and lungs from anaesthetized mice

were perfused via the portal vein sequentially with balanced salt solution containing 5

mM glucose, 0.01% sodium heparin, and 5 mM EGTA and DMEM medium

containing 4 mM $CaCl_2$, 2% FBS and 0.05% collagenase IV at a flow rate of 5 ml/min

for 5 minutes, respectively. The following steps are similar in order to separate the

liver sinusoids endothelial cells and pulmonary vascular endothelial cells. Isolation of

primary embryonic neurons and single-cell suspensions was performed using Pierce

Primary Neuron Isolation Kit by density gradient centrifugation.

1617

***In vitro* blood-brain barrier model**

The *in vitro* BBB model was constructed with primary endothelial cells using a

Transwell cell culture system. Briefly, BMECs were seeded onto the upper chamber

of a Transwell pre-coated with gelatine, and cultured with the medium containing 10%

FBS. The integrity of the monolayer cell was evaluated by measuring the TEER

values using a Millicell-ERS voltohmmeter (Millipore). The cell monolayers with

TEER values higher than 300 $\Omega.cm^2$ were used as BBB models for the transmigration

studies. The neurons cells were seeded onto the lower chamber of the Transwell for

bacterial loading.

Macrophages were exposed to *L. monocytogenes* (MOI=10) for 6 hours and then

treated with 100 µg/ml gentamicin for 2 hours to kill the extracellular bacteria. After

construction of the BBB model, the infected macrophages were trypsinized for cell

counting and added into the upper chamber. The incubation time was 12 hours (unless

otherwise indicated) and then the neurons cells in the bottom chamber were lysed for

penetrating bacterial quantification.

1633

Live-cell flow chamber assays

The adhesion, shear resistance, crawling, and transmigration of macrophages on

endothelial monolayer were tested using a flow cell system as previously described.[40]

Shear flow was applied by a syringe pump (PHD22/2000, Harvard Apparatus) into a

microfluidic chip with a 100 μm $\times$ 1 mm $\times$ 15 mm main flowing zone. Different shear

stresses were acquired by setting corresponding volumetric flow rates. Isolated

macrophages were cultured with different concentrations of ox-LDL for 24 hours and

then infected with *L. monocytogenes* (MOI=10) for 6 hours. Macrophages were re-

suspended and stained with Hoechst. The cells were then perfused over pre-formed

endothelial monolayer at a shear stress of 0.1 dyn/cm^2 for 8 minutes. Then, the

adhesion and crawling dynamics of macrophages on endothelial monolayer were

recorded at shear stress of 2 dyn/cm^2 for 10 minutes, by using a CCD camera with

20$\times$ objective. Images were taken from the video every second and analyzed for the

crawling parameters and transmigrating duration with Image J software.

The post-arrest crawling of macrophages on the microfluidic chip can be factorized

into two metrics: the forward migration index (FMI) towards the x-axis (along with

flow, xFMI), and the y-axis (perpendicular to flow, yFMI). xFMI and yFMI were

accurately measured by a CCD camera, and these two indices constituted the crawling

trajectories of each individual cell shown in Figure 3E. Directionality was quantified

as the ratio of yFMI to xFMI. Directionality indicates the angle at which the direction

of cell movement deviates from the shear flow. The larger the value of directionality,

the less the direction of cell movement is affected by the shear flow, indicating that

the cell has a stronger ability to fight against the shear flow. Transmigration duration,

which is defined as the interval from the timepoint of cell arrival at the site to the

endpoint of completion of transmigration beneath the endothelial monolayer (Figure

S6H), was used to indicate the transmigration capacity of the cell. Here a macrophage

crossing the monolayer with a 70% fluorescence intensity reduction was deemed

transmigrated.

For testing shear-resistant macrophage adhesion, primed macrophages were re-

suspended and counted. The same number of cells (2×10^4) was completely perfused

over pre-formed endothelial monolayer within 10 minutes. The shear stress was

increased stepwise from 1 to 2, 4, 8, 16, 32 dyn/cm^2 for 30 seconds each. Remaining

adherent macrophages on endothelial monolayer were recorded at 20× objective by a

CCD camera and counted using Image J software. The normalized adhesion fraction

was calculated by dividing the number of adherent macrophages at the endpoint of

each shear phase by the number at the endpoint of 1 dyn/cm^2. For defining the

macrophage transmigration dynamics, the time-lapse images were analyzed offline

frame-by-frame every second.

1672

Quantification of cell area and protrusions

The cell area and protrusions of macrophages were measured using a quantitative cell

morphology analysis system adapted from Young et al.[67] In brief, peritoneal

macrophages were stimulated with different concentrations of ox-LDL for 24 hours,

and then cells were fixed, permeabilized, and labeled with phalloidin-Alexa 647.

Samples were analyzed using a Zeiss LSM900 confocal microscope. 8-bit images

were acquired by using ImageJ software and converted to grayscale to best visualize

all positive staining. Brightness and contrast were adjusted and a despeckle step was

performed as needed until cell edges were clearly visible. For fractal analysis, images

were skeletonized using the plugin AnalyzeSkeleton (2D/3D). Skeleton fragments

generated during image acquisition were removed to facilitate quantification.

Protrusions were identified as the actin tails extending from the cells surface. The area

of the cells as well as the length of the actin tails were measured using ImageJ

software. At least ten cells were scored per group.

1687

Single cell RNA-sequencing analysis

PBMCs isolated from mice were filtered using a 40-µm nylon cell strainer and red

blood cells were removed by red blood cell lysis solution. Dissociated cells were

washed with DPBS containing 2% FBS. Cells were stained with 0.4% Trypan blue to

check the viability. Sequencing libraries were prepared using randomly interrupted

whole-transcriptome amplification products to enrich the 3' end of the transcripts

linked with the cell barcode and unique molecular identifier. All the procedures

including the library construction were performed strictly according to the standard

manufacturer's protocol [10× Genomics Chromium Single-Cell 3′ kit (V3)].

Sequencing libraries were quantified using a High Sensitivity DNA Chip on a

Bioanalyzer 2100 and the Qubit High Sensitivity DNA Assay. Single cells were

sequenced using an Illumina NovaSeq 6000 sequencing system (paired-end

multiplexing run, 150 bp) by Majorbio Bio-pharm Technology Co., Ltd (Shanghai,

China). The sequencing and bioinformatic analysis were performed on platform of

Majorbio (Shanghai, China).

Reads were processed using the Cell Ranger (v7.1.0) pipeline with default and

recommended parameters. FASTQs generated from Illumina sequencing output were

aligned to the mouse genome, version GRCm38, using the STAR algorithm. Gene-

barcode matrices containing the barcoded cells and gene expression counts were

generated, and then imported into the Seurat (v3.2.0) R toolkit for quality control and

downstream analysis of our single cell RNAseq data.

Cells were clustered using graph-based clustering of the PCA reduced data with the

Louvain Method after computing a shared nearest neighbor graph. For sub-clustering,

we applied the same procedure of scaled, dimensionality reduction, and clustering to

the specific set of data. For each cluster, we used the Wilcoxon Rank-Sum Test to find

genes with significant deferential expression compared to the remaining clusters.

Differentially expression genes (DEGs) were identified with Seurat's FindMarkers,

using a likelihood ratio test. DEGs with $|\log_2$ (fold change)$| > 1$ and p value ≤ 0.05

were considered to be significant. In addition, GO functional-enrichment analysis was

performed to identify which DEGs were significantly enriched in GO terms and

metabolic pathways at Bonferroni-corrected P-value ≤ 0.05 compared with the whole-

transcriptome background.

1720

Lipid metabolomics

Metabolite extraction and mass spectrometry was performed using an Ultra-High

Performance Liquid Chromatography-mass spectrometry (UHPLC-MS) by Biomarker

Technologies (Beijing, China). The raw data was collected under the control of the

acquisition software (MassLynx V4.2, Waters), and then processed by Progenesis QI

software integrated with the METLIN database (Waters). The theoretical fragment

identification and mass deviation were within 100 ppm. Following normalization of

peak areas relative to the total peak area, principal component analysis (PCA) and

Spearman correlation analysis were performed to evaluate the repeatability of the

samples. A total of 4,625 named metabolites were identified for the datasets. The

identified compounds were classified, and pathway information was annotated by

using the KEGG, HMDB and LipidMaps databases. The enrichment significance of

differential metabolites in KEGG pathways was evaluated using a hypergeometric

distribution test.

1735

RNA-seq (RNA Sequencing)

RNA was extracted from mouse liver tissues using a Trizol extraction and Purelink

RNA kit (ThermoFisher Scientific). Bioanalyzer chips were used to assess RNA

quality prior to RNA sequencing. A total of 1 μg RNA per sample was used as input

material for RNA sample preparation. Sequencing libraries were generated using a

Hieff NGS Ultima Dual-mode mRNA Library Prep Kit for lumina following

manufacturer's recommendations, and index codes were added to attribute sequences

for each sample. The libraries were sequenced on an Illumina NovaSeq platform to

generate 150 bp paired-end reads, according to the manufacturer's instructions.

1745

Measurement of ECAR

ECAR was measured using the XF Glycolysis Stress Test kit. In brief, liver tissue was

inoculated into XFe24 Cell Culture Microplates and cultured overnight at 37℃ in an

incubator containing 5% CO_2. Three injections (10 mM glucose, 1.5 μM oligomycin

and 50 mM 2-DG) were performed at the indicated timepoints. ECAR values were

normalized to the total amount of proteins. ECAR parameters were calculated using

Wave software (v.2.4.3, Seahorse Bioscience, Agilent Technologies) according to

manufacturer instructions.

1754

Fura-4 Ca^{2+} imaging

Cells were washed with PBS and incubated with 5 μM Fluo-4 AM and 0.04%

Pluronic-127 for 60 minutes at 37℃. Then, cells were washed with PBS and

incubated with PBS for 30 minutes at 37℃. Fluorescence was detected using a

CytoFLEX Flow Cytometer (Beckman Coulter) equipped with a 488 nm excitation

wavelength. 30 seconds after loaded with Fluo-4 AM, cells were then treated with

BHB (10 mM) as indicated in the figures. Data was analyzed with Flow Jo to

determine the fluorescence intensity.

1763

Real-time qRT-PCR

Total RNA was isolated using a RNeasy Mini Kit (QIAGEN). The mRNA was

reverse-transcribed and quantified using a HiScript II 1st Strand cDNA Synthesis Kit

(Vazyme). qRT-PCR was performed on an ABI StepOnePlus PCR system (Thermo

Fisher), and results were normalized to GAPDH mRNA levels. Data were analyzed

using the $2^{-\Delta\Delta CT}$ method. All the primers used for qRT-PCR are listed in Table S3.

1770

Statistical analysis

Statistical analysis was performed using GraphPad Prism Software. Two-group

comparisons were made using two-tailed unpaired Student's t-test. One-way analysis

of variance (ANOVA) with appropriate multiple comparisons tests was used to

compare three independent groups. For comparisons of multiple factors, two-way

ANOVA with appropriate multiple comparisons tests was used. A value of $p < 0.05$

was considered significant. Specific details of statistical analysis are presented above

or in associated figure legends.

1779

1780

1781

References

1.   Forrester, J.V., McMenamin, P.G., and Dando, S.J. (2018). CNS infection and

immune privilege. Nat Rev Neurosci *19*, 655-671. 10.1038/s41583-018-0070-

8.

2.   Langen, U.H., Ayloo, S., and Gu, C. (2019). Development and Cell Biology of

the Blood-Brain Barrier. Annu Rev Cell Dev Biol *35*, 591-613.

10.1146/annurev-cellbio-100617-062608.

3.   Loh, L.N., Gao, G., and Tuomanen, E.I. (2017). Dissecting Bacterial Cell Wall

1790   Entry and Signaling in Eukaryotic Cells: an Actin-Dependent Pathway

1791   Parallels Platelet-Activating Factor Receptor-Mediated Endocytosis. mBio *8*.

1792   10.1128/mBio.02030-16.

1793   4.   Tabusi, M., Thorsdottir, S., Lysandrou, M., Narciso, A.R., Minoia, M.,

1794   Srambickal, C.V., Widengren, J., Henriques-Normark, B., and Iovino, F.

1795   (2021). Neuronal death in pneumococcal meningitis is triggered by

1796   pneumolysin and RrgA interactions with beta-actin. PLoS Pathog *17*,

1797   e1009432. 10.1371/journal.ppat.1009432.

1798   5.   Coureuil, M., Lecuyer, H., Bourdoulous, S., and Nassif, X. (2017). A journey

1799   into the brain: insight into how bacterial pathogens cross blood-brain barriers.

1800   Nat Rev Microbiol *15*, 149-159. 10.1038/nrmicro.2016.178.

1801   6.   Coureuil, M., Lecuyer, H., Scott, M.G., Boularan, C., Enslen, H., Soyer, M.,

1802   Mikaty, G., Bourdoulous, S., Nassif, X., and Marullo, S. (2010).

1803   Meningococcus Hijacks a beta2-adrenoceptor/beta-Arrestin pathway to cross

1804   brain microvasculature endothelium. Cell *143*, 1149-1160.

1805   10.1016/j.cell.2010.11.035.

1806   7.   Lopes-Luz, L., Mendonca, M., Bernardes Fogaca, M., Kipnis, A., Bhunia,

1807   A.K., and Buhrer-Sekula, S. (2021). Listeria monocytogenes: review of

1808   pathogenesis and virulence determinants-targeted immunological assays. Crit

1809   Rev Microbiol *47*, 647-666. 10.1080/1040841X.2021.1911930.

1810   8.   Ghosh, P., Halvorsen, E.M., Ammendolia, D.A., Mor-Vaknin, N., O'Riordan,

1811   M.X.D., Brumell, J.H., Markovitz, D.M., and Higgins, D.E. (2018). Invasion

of the Brain by Listeria monocytogenes Is Mediated by InlF and Host Cell

Vimentin. mBio *9*. 10.1128/mBio.00160-18.

9.   Pagelow, D., Chhatbar, C., Beineke, A., Liu, X., Nerlich, A., van Vorst, K.,

Rohde, M., Kalinke, U., Forster, R., Halle, S., et al. (2018). The olfactory

epithelium as a port of entry in neonatal neurolisteriosis. Nat Commun *9*, 4269.

10.1038/s41467-018-06668-2.

10.   Levin, B.R., Baquero, F., Ankomah, P.P., and McCall, I.C. (2017). Phagocytes,

Antibiotics, and Self-Limiting Bacterial Infections. Trends Microbiol *25*, 878-

892. 10.1016/j.tim.2017.07.005.

11.   Boada-Romero, E., Martinez, J., Heckmann, B.L., and Green, D.R. (2020).

The clearance of dead cells by efferocytosis. Nat Rev Mol Cell Biol *21*, 398-

414. 10.1038/s41580-020-0232-1.

12.   Locati, M., Curtale, G., and Mantovani, A. (2020). Diversity, Mechanisms,

and Significance of Macrophage Plasticity. Annu Rev Pathol *15*, 123-147.

10.1146/annurev-pathmechdis-012418-012718.

13.   Zhai, L., Huang, C., Ning, Z., Zhang, Y., Zhuang, M., Yang, W., Wang, X.,

Wang, J., Zhang, L., Xiao, H., et al. (2023). Ruminococcus gnavus plays a

pathogenic role in diarrhea-predominant irritable bowel syndrome by

increasing serotonin biosynthesis. Cell Host Microbe *31*, 33-44 e35.

10.1016/j.chom.2022.11.006.

14.   Van der Merwe, S., Chokshi, S., Bernsmeier, C., and Albillos, A. (2021). The

multifactorial mechanisms of bacterial infection in decompensated cirrhosis. J

Hepatol *75 Suppl 1*, S82-S100. 10.1016/j.jhep.2020.11.029.

15.	Maudet, C., Kheloufi, M., Levallois, S., Gaillard, J., Huang, L., Gaultier, C.,

Tsai, Y.H., Disson, O., and Lecuit, M. (2022). Bacterial inhibition of Fas-

mediated killing promotes neuroinvasion and persistence. Nature *603*, 900-906.

10.1038/s41586-022-04505-7.

16.	Pinho-Ribeiro, F.A., Deng, L., Neel, D.V., Erdogan, O., Basu, H., Yang, D.,

Choi, S., Walker, A.J., Carneiro-Nascimento, S., He, K., et al. (2023). Bacteria

hijack a meningeal neuroimmune axis to facilitate brain invasion. Nature *615*,

472-481. 10.1038/s41586-023-05753-x.

17.	Xu, K., Yin, N., Peng, M., Stamatiades, E.G., Shyu, A., Li, P., Zhang, X., Do,

M.H., Wang, Z., Capistrano, K.J., et al. (2021). Glycolysis fuels

phosphoinositide 3-kinase signaling to bolster T cell immunity. Science *371*,

405-410. 10.1126/science.abb2683.

18.	Escoll, P., Song, O.R., Viana, F., Steiner, B., Lagache, T., Olivo-Marin, J.C.,

Impens, F., Brodin, P., Hilbi, H., and Buchrieser, C. (2017). Legionella

pneumophila Modulates Mitochondrial Dynamics to Trigger Metabolic

Repurposing of Infected Macrophages. Cell Host Microbe *22*, 302-316 e307.

10.1016/j.chom.2017.07.020.

19.	Yang, D., Jacobson, A., Meerschaert, K.A., Sifakis, J.J., Wu, M., Chen, X.,

Yang, T., Zhou, Y., Anekal, P.V., Rucker, R.A., et al. (2022). Nociceptor

neurons direct goblet cells via a CGRP-RAMP1 axis to drive mucus

production and gut barrier protection. Cell *185*, 4190-4205 e4125.

10.1016/j.cell.2022.09.024.

20.   Loft, A., Schmidt, S.F., Caratti, G., Stifel, U., Havelund, J., Sekar, R., Kwon,

Y., Sulaj, A., Chow, K.K., Alfaro, A.J., et al. (2022). A macrophage-hepatocyte

glucocorticoid receptor axis coordinates fasting ketogenesis. Cell Metab *34*,

473-486 e479. 10.1016/j.cmet.2022.01.004.

21.   Lu, J.F., Zhu, M.Q., Xia, B., Zhang, N.N., Liu, X.P., Liu, H., Zhang, R.X.,

Xiao, J.Y., Yang, H., Zhang, Y.Q., et al. (2023). GDF15 is a major determinant

of ketogenic diet-induced weight loss. Cell Metab *35*, 2165-2182 e2167.

10.1016/j.cmet.2023.11.003.

22.   Tomlinson, K.L., Chen, Y.T., Junker, A., Urso, A., Wong Fok Lung, T., Ahn,

D., Hofstaedter, C.E., Baskota, S.U., Ernst, R.K., Prince, A., and Riquelme,

S.A. (2023). Ketogenesis promotes tolerance to Pseudomonas aeruginosa

pulmonary infection. Cell Metab *35*, 1767-1781 e1766.

10.1016/j.cmet.2023.09.001.

23.   Seufert, A.L., Hickman, J.W., Traxler, S.K., Peterson, R.M., Waugh, T.A.,

Lashley, S.J., Shulzhenko, N., Napier, R.J., and Napier, B.A. (2022). Enriched

dietary saturated fatty acids induce trained immunity via ceramide production

that enhances severity of endotoxemia and clearance of infection. Elife *11*.

10.7554/eLife.76744.

24.   Wilhelm, C., Surendar, J., and Karagiannis, F. (2021). Enemy or ally? Fasting

as an essential regulator of immune responses. Trends Immunol *42*, 389-400.

10.1016/j.it.2021.03.007.

25.   McFarland, A.P., Yalin, A., Wang, S.Y., Cortez, V.S., Landsberger, T., Sudan,

R., Peng, V., Miller, H.L., Ricci, B., David, E., et al. (2021). Multi-tissue

single-cell analysis deconstructs the complex programs of mouse natural killer

and type 1 innate lymphoid cells in tissues and circulation. Immunity *54*,

1320-1337 e1324. 10.1016/j.immuni.2021.03.024.

26.   Ahmad, F., Rani, A., Alam, A., Zarin, S., Pandey, S., Singh, H., Hasnain, S.E.,

and Ehtesham, N.Z. (2022). Macrophage: A Cell With Many Faces and

Functions in Tuberculosis. Front Immunol *13*, 747799.

10.3389/fimmu.2022.747799.

27.   Hill, D.A., Lim, H.W., Kim, Y.H., Ho, W.Y., Foong, Y.H., Nelson, V.L.,

Nguyen, H.C.B., Chegireddy, K., Kim, J., Habertheuer, A., et al. (2018).

Distinct macrophage populations direct inflammatory versus physiological

changes in adipose tissue. Proc Natl Acad Sci U S A *115*, E5096-E5105.

10.1073/pnas.1802611115.

28.   Jaitin, D.A., Adlung, L., Thaiss, C.A., Weiner, A., Li, B., Descamps, H.,

Lundgren, P., Bleriot, C., Liu, Z., Deczkowska, A., et al. (2019). Lipid-

Associated Macrophages Control Metabolic Homeostasis in a Trem2-

Dependent Manner. Cell *178*, 686-698 e614. 10.1016/j.cell.2019.05.054.

29.   Luis, G., Godfroid, A., Nishiumi, S., Cimino, J., Blacher, S., Maquoi, E., Wery,

C., Collignon, A., Longuespee, R., Montero-Ruiz, L., et al. (2021). Tumor

resistance to ferroptosis driven by Stearoyl-CoA Desaturase-1 (SCD1) in

cancer cells and Fatty Acid Biding Protein-4 (FABP4) in tumor

microenvironment promote tumor recurrence. Redox Biol *43*, 102006.

10.1016/j.redox.2021.102006.

30.   Victor, M.B., Leary, N., Luna, X., Meharena, H.S., Scannail, A.N., Bozzelli,

P.L., Samaan, G., Murdock, M.H., von Maydell, D., Effenberger, A.H., et al.

(2022). Lipid accumulation induced by APOE4 impairs microglial

surveillance of neuronal-network activity. Cell Stem Cell *29*, 1197-1212 e1198.

10.1016/j.stem.2022.07.005.

31.   Chen, Y., Zhang, J., Cui, W., and Silverstein, R.L. (2022). CD36, a signaling

receptor and fatty acid transporter that regulates immune cell metabolism and

fate. J Exp Med *219*. 10.1084/jem.20211314.

32.   Gudgeon, J., Marin-Rubio, J.L., and Trost, M. (2022). The role of macrophage

scavenger receptor 1 (MSR1) in inflammatory disorders and cancer. Front

Immunol *13*, 1012002. 10.3389/fimmu.2022.1012002.

33.   Botham, K.M., and Wheeler-Jones, C.P. (2013). Postprandial lipoproteins and

the molecular regulation of vascular homeostasis. Prog Lipid Res *52*, 446-464.

10.1016/j.plipres.2013.06.001.

34.   Xue, J., Zhao, Z., Zhang, L., Xue, L., Shen, S., Wen, Y., Wei, Z., Wang, L.,

Kong, L., Sun, H., et al. (2017). Neutrophil-mediated anticancer drug delivery

for suppression of postoperative malignant glioma recurrence. Nat

Nanotechnol *12*, 692-700. 10.1038/nnano.2017.54.

35.   Peyrusson, F., Varet, H., Nguyen, T.K., Legendre, R., Sismeiro, O., Coppee,

J.Y., Wolz, C., Tenson, T., and Van Bambeke, F. (2020). Intracellular

Staphylococcus aureus persisters upon antibiotic exposure. Nat Commun *11*,

2200. 10.1038/s41467-020-15966-7.

36.   Ding, Z., Pothineni, N.V.K., Goel, A., Luscher, T.F., and Mehta, J.L. (2020).

PCSK9 and inflammation: role of shear stress, pro-inflammatory cytokines,

and LOX-1. Cardiovasc Res *116*, 908-915. 10.1093/cvr/cvz313.

37.   Da Dalt, L., Castiglioni, L., Baragetti, A., Audano, M., Svecla, M., Bonacina,

F., Pedretti, S., Uboldi, P., Benzoni, P., Giannetti, F., et al. (2021). PCSK9

deficiency rewires heart metabolism and drives heart failure with preserved

ejection fraction. Eur Heart J *42*, 3078-3090. 10.1093/eurheartj/ehab431.

38.   Katsuki, S., P, K.J., Lupieri, A., Nakano, T., Passos, L.S.A., Rogers, M.A.,

Becker-Greene, D., Le, T.D., Decano, J.L., Ho Lee, L., et al. (2022).

Proprotein Convertase Subtilisin/Kexin 9 (PCSK9) Promotes Macrophage

Activation via LDL Receptor-Independent Mechanisms. Circ Res *131*, 873-

889. 10.1161/CIRCRESAHA.121.320056.

39.   Qi, Z., Hu, L., Zhang, J., Yang, W., Liu, X., Jia, D., Yao, Z., Chang, L., Pan, G.,

Zhong, H., et al. (2021). PCSK9 (Proprotein Convertase Subtilisin/Kexin 9)

Enhances Platelet Activation, Thrombosis, and Myocardial Infarct Expansion

by Binding to Platelet CD36. Circulation *143*, 45-61.

10.1161/CIRCULATIONAHA.120.046290.

40.   Yang, H., Li, N., Du, Y., Tong, C., Lu, S., Hu, J., Zhang, Y., and Long, M.

(2017). Neutrophil adhesion and crawling dynamics on liver sinusoidal

endothelial cells under shear flow. Exp Cell Res *351*, 91-99.

10.1016/j.yexcr.2017.01.002.

41.   Neupane, A.S., Willson, M., Chojnacki, A.K., Vargas, E.S.C.F., Morehouse, C.,

Carestia, A., Keller, A.E., Peiseler, M., DiGiandomenico, A., Kelly, M.M., et

al. (2020). Patrolling Alveolar Macrophages Conceal Bacteria from the

Immune System to Maintain Homeostasis. Cell *183*, 110-125 e111.

10.1016/j.cell.2020.08.020.

42.   Hernandez, G.E., and Iruela-Arispe, M.L. (2020). The many flavors of

monocyte/macrophage--endothelial cell interactions. Curr Opin Hematol *27*,

181-189. 10.1097/MOH.0000000000000573.

43.   Pang, X., He, X., Qiu, Z., Zhang, H., Xie, R., Liu, Z., Gu, Y., Zhao, N., Xiang,

Q., and Cui, Y. (2023). Targeting integrin pathways: mechanisms and advances

in therapy. Signal Transduct Target Ther *8*, 1. 10.1038/s41392-022-01259-6.

44.   Deng, L., Kersten, S., and Stienstra, R. (2023). Triacylglycerol uptake and

handling by macrophages: From fatty acids to lipoproteins. Prog Lipid Res *92*,

101250. 10.1016/j.plipres.2023.101250.

45.   Singh, V., Jamwal, S., Jain, R., Verma, P., Gokhale, R., and Rao, K.V. (2012).

Mycobacterium tuberculosis-driven targeted recalibration of macrophage lipid

homeostasis promotes the foamy phenotype. Cell Host Microbe *12*, 669-681.

10.1016/j.chom.2012.09.012.

46.   Patterson, M.T., Firulyova, M.M., Xu, Y., Hillman, H., Bishop, C., Zhu, A.,

Hickok, G.H., Schrank, P.R., Ronayne, C.E., Caillot, Z., et al. (2023). Trem2

promotes foamy macrophage lipid uptake and survival in atherosclerosis.

Nature Cardiovascular Research *2*, 1015-1031. 10.1038/s44161-023-00354-3.

47.   Andrews, N.W., and Corrotte, M. (2018). Plasma membrane repair. Curr Biol

28, R392-R397. 10.1016/j.cub.2017.12.034.

48.   Jimenez, A.J., Maiuri, P., Lafaurie-Janvore, J., Divoux, S., Piel, M., and Perez,

F. (2014). ESCRT machinery is required for plasma membrane repair. Science

343, 1247136. 10.1126/science.1247136.

49.   Cook, A.D., Braine, E.L., and Hamilton, J.A. (2003). The phenotype of

inflammatory macrophages is stimulus dependent: implications for the nature

of the inflammatory response. J Immunol *171*, 4816-4823.

10.4049/jimmunol.171.9.4816.

50.   Zhang, S., Peng, X., Yang, S., Li, X., Huang, M., Wei, S., Liu, J., He, G.,

Zheng, H., Yang, L., et al. (2022). The regulation, function, and role of

lipophagy, a form of selective autophagy, in metabolic disorders. Cell Death

Dis *13*, 132. 10.1038/s41419-022-04593-3.

51.   Newman, J.C., and Verdin, E. (2017). beta-Hydroxybutyrate: A Signaling

Metabolite. Annu Rev Nutr *37*, 51-76. 10.1146/annurev-nutr-071816-064916.

52.   Youm, Y.H., Nguyen, K.Y., Grant, R.W., Goldberg, E.L., Bodogai, M., Kim,

D., D'Agostino, D., Planavsky, N., Lupfer, C., Kanneganti, T.D., et al. (2015).

The ketone metabolite beta-hydroxybutyrate blocks NLRP3 inflammasome-

mediated inflammatory disease. Nat Med *21*, 263-269. 10.1038/nm.3804.

53.   Duking, T., Spieth, L., Berghoff, S.A., Piepkorn, L., Schmidke, A.M.,

Mitkovski, M., Kannaiyan, N., Hosang, L., Scholz, P., Shaib, A.H., et al.

(2022). Ketogenic diet uncovers differential metabolic plasticity of brain cells.

Sci Adv *8*, eabo7639. 10.1126/sciadv.abo7639.

54.   Matsuura, T.R., Puchalska, P., Crawford, P.A., and Kelly, D.P. (2023). Ketones

and the Heart: Metabolic Principles and Therapeutic Implications. Circ Res

132, 882-898. 10.1161/CIRCRESAHA.123.321872.

55.   Shah, U.A., and Iyengar, N.M. (2022). Plant-Based and Ketogenic Diets As

Diverging Paths to Address Cancer: A Review. JAMA Oncol *8*, 1201-1208.

10.1001/jamaoncol.2022.1769.

56.   Doran, K.S., Fulde, M., Gratz, N., Kim, B.J., Nau, R., Prasadarao, N.,

Schubert-Unkmeir, A., Tuomanen, E.I., and Valentin-Weigand, P. (2016).

Host-pathogen interactions in bacterial meningitis. Acta Neuropathol *131*,

185-209. 10.1007/s00401-015-1531-z.

57.   Steeb, B., Claudi, B., Burton, N.A., Tienz, P., Schmidt, A., Farhan, H., Maze,

A., and Bumann, D. (2013). Parallel exploitation of diverse host nutrients

enhances Salmonella virulence. PLoS Pathog *9*, e1003301.

10.1371/journal.ppat.1003301.

58.   Peyron, P., Vaubourgeix, J., Poquet, Y., Levillain, F., Botanch, C., Bardou, F.,

Daffe, M., Emile, J.F., Marchou, B., Cardona, P.J., et al. (2008). Foamy

macrophages from tuberculous patients' granulomas constitute a nutrient-rich

reservoir for M. tuberculosis persistence. PLoS Pathog *4*, e1000204.

10.1371/journal.ppat.1000204.

59.   Yang, Y., Fang, H., Xie, Z., Ren, F., Yan, L., Zhang, M., Xu, G., Song, Z.,

2010   Chen, Z., Sun, W., et al. (2024). Yersinia infection induces glucose depletion

2011   and AMPK-dependent inhibition of pyroptosis in mice. Nat Microbiol.

2012   10.1038/s41564-024-01734-6.

2013 60. Stubbs, B.J., Cox, P.J., Evans, R.D., Santer, P., Miller, J.J., Faull, O.K., Magor-

2014   Elliott, S., Hiyama, S., Stirling, M., and Clarke, K. (2017). On the Metabolism

2015   of Exogenous Ketones in Humans. Front Physiol *8*, 848.

2016   10.3389/fphys.2017.00848.

2017 61. Ari, C., Murdun, C., Koutnik, A.P., Goldhagen, C.R., Rogers, C., Park, C.,

2018   Bharwani, S., Diamond, D.M., Kindy, M.S., D'Agostino, D.P., and Kovacs, Z.

2019   (2019). Exogenous Ketones Lower Blood Glucose Level in Rested and

2020   Exercised Rodent Models. Nutrients *11*. 10.3390/nu11102330.

2021 62. Acar, R. (2021). Association between Beta-Hydroxybutyrate Levels and

2022   Survival in Sepsis Patients.

2023 63. Wang, A., Huen, S.C., Luan, H.H., Yu, S., Zhang, C., Gallezot, J.D., Booth,

2024   C.J., and Medzhitov, R. (2016). Opposing Effects of Fasting Metabolism on

2025   Tissue Tolerance in Bacterial and Viral Inflammation. Cell *166*, 1512-1525

2026   e1512. 10.1016/j.cell.2016.07.026.

2027 64. Pereira-Dutra, F.S., Souza, E.K., Souza, T.S., Goltara-Gomes, T.C., Ferraro-

2028   Moreira, F., Palhinha, L., Cunha-Fernandes, T., Rajao, M.A., Silva, A.R., and

2029   Bozza, P.T. (2024). Accumulation of lipid droplets induced by Listeria

2030   monocytogenes in macrophages: implications for survival and evasion of

2031   innate immunity. J Leukoc Biol *116*, 1364-1371. 10.1093/jleuko/qiae115.

2032 65. Koopmans, M.M., Engelen-Lee, J., Brouwer, M.C., Jaspers, V., Man, W.K.,

2033 Vall Seron, M., and van de Beek, D. (2018). Characterization of a Listeria

2034 monocytogenes meningitis mouse model. J Neuroinflammation *15*, 257.

2035 10.1186/s12974-018-1293-3.

2036 66. Ali, M.A., and Kravitz, A.V. (2018). Challenges in quantifying food intake in

2037 rodents. Brain Res *1693*, 188-191. 10.1016/j.brainres.2018.02.040.

2038 67. Young, K., and Morrison, H. (2018). Quantifying Microglia Morphology from

2039 Photomicrographs of Immunohistochemistry Prepared Tissue Using ImageJ. J

2040 Vis Exp. 10.3791/57648.

2041

Graphical Abstract

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 10

A
Risk factors for patients with bacterial CNS infection

	OR (95%CI)	adjust p
Ferritin	5.09 (1.55, 16.71)	0.006
Fibrinogen	3.57 (1.13, 11.25)	0.026
LDL	3.31 (1.01, 10.83)	0.043
Glucose	2.17 (1.04, 4.50)	0.028
AST	2.16 (1.01, 4.63)	0.035
Bilirubin	1.98 (0.86, 4.55)	0.077
Bicarbonate	1.73 (0.82, 3.63)	0.103
Lactate	1.53 (0.71, 3.32)	0.187
ALT	1.43 (0.68, 3.01)	0.225
Vitamin B12	1.41 (0.45, 4.46)	0.385
eAG	1.40 (0.41, 4.76)	0.409
Amylase	1.22 (0.35, 4.19)	0.500
CK	1.12 (0.47, 2.68)	0.489
BUN	1.10 (0.53, 2.26)	0.476
TSH	0.90 (0.36, 2.23)	0.500
Iron	0.85 (0.28, 2.58)	0.500
Chloride	0.83 (0.41, 1.69)	0.369
Magnesium	0.80 (0.39, 1.63)	0.332
Lipase	0.77 (0.31, 1.92)	0.375
Creatinine	0.65 (0.31, 1.36)	0.172
AG	0.65 (0.31, 1.35)	0.165
pH	0.63 (0.25, 1.56)	0.219
Triglycerides	0.63 (0.22, 1.78)	0.269
Calcium	0.55 (0.27, 1.13)	0.074
Sodium	0.52 (0.25, 1.07)	0.055
Potassium	0.42 (0.20, 0.87)	0.014
AP	0.38 (0.18, 0.83)	0.011
Phosphate	0.23 (0.11, 0.50)	0.000
HDL	0.19 (0.05, 0.66)	0.008
Transferrin	0.14 (0.04, 0.50)	0.002

$\text{Log}_2(\text{OR})$

Supplementary Figure 11