Title: Dysregulation of macrophage lipid metabolism underlies intracellular bacterial

neuroinvasion

Authors: Zhou Sha[1, #], Kun Yang[2, #], Shoupeng Fu[3, #], Xiaoyong Tong[4, #], Hui Yang[1],

Huiling Fang[1], Qianqian He[1], Ning Li[5, 6], Xinyu Shu[5, 6], Qi Liu[1], Yongliang Du[1],

Beibei Fu[1], Yan Xiong[1], Dong Guo[1], Jing Liu[4], Qian Li[7], Hao Zeng[8], Xiaokai Zhang[8],

Rui Yao[9], Xushuo Zhang[10], Wenjin Guo[3], Xuhu Mao[7], Mian Long[5, 6], Xiaoyuan Lin[7,

11, *], Quanming Zou[8, *], Haibo Wu[1, *]

1 School of Life Sciences, Chongqing University, Chongqing 401331, China

2 Chongqing Public Health Medical Center, Chongqing 400036, China

3 School of Veterinary Medicine, Jilin University, Changchun 130062, China

4 School of Pharmaceutical Sciences, Chongqing University, Chongqing 401331,

China

5 Center of Biomechanics and Bioengineering, Key Laboratory of Microgravity

(National Microgravity Laboratory), and Beijing Key Laboratory of Engineered

Construction and Mechanobiology, Institute of Mechanics, Chinese Academy of

Sciences, Beijing 100190, China

6 School of Engineering Sciences, University of Chinese Academy of Sciences,

Beijing 100049, China

7 Department of Clinical Microbiology and Immunology, College of Pharmacy and

Laboratory Medicine, Army Medical University (Third Military Medical University),

Chongqing 400038, China.

8 National Engineering Research Center of Immunological, Department of

Microbiology and Biochemical Pharmacy, College of Pharmacy and Laboratory

Medicine, Army Medical University (Third Military Medical University), Chongqing

400038, China.

9 Department of Pathology, Chongqing Hygeia Hospital, Chongqing 401331, China

10 Zhejiang Toyouvax Biopharming, Hangzhou 311100, China

11 Institut für Virologie, Freie Universität Berlin, Robert-von-Ostertag-Str. 7-13,

Berlin 14163, Germany.

# Equally Contributing Authors

* Corresponding authors

Xiaoyuan Lin, linxiaoyuan23@163.com

Quanming Zou, qmzou2007@163.com

Haibo Wu, hbwu023@cqu.edu.cn

Abstract

Acute infection of the central nervous system is one of the deadliest diseases, but the

mechanisms by which intracellular bacteria infiltrate the brain remain poorly

understood. Phagocytic cells are usually recognized as the battlefield on which war is

waged against intracellular bacteria; however, little is known about how the

intracellular bacteria take advantage of infected phagocytes to access the brain. In this

study, we find that a novel CD36$^+$ foamy macrophage subpopulation participates in

penetration of the brain by intracellular bacteria. Biomechanical analysis reveals that

abundant protrusions and adhesion molecules on foamy macrophages confer

significant resistance to the mechanical stress of blood flow, thereby providing more

opportunities for these macrophages to adhere to the vascular endothelial surface

during neuroinvasion. Through metabolomics analysis, we find that macrophage lipid

metabolism is recalibrated during bacterial neuroinvasion, and that β-hydroxybutyrate

promotes the formation and survival of CD36$^+$ foamy macrophages. Taken together,

our findings uncover a pathway by which intracellular bacteria hijack macrophages to

invade the brain, suggesting that lipid metabolism might play a role in the prevention

or resolution of bacterial neuroinvasion.

Keywords: intracellular bacteria; neuroinvasion; foamy macrophage; ketogenesis; β-

hydroxybutyrate

Introduction

Meningitis is an inflammation of the protective membranes covering the brain that caused by bacterial or viral infection.[1,2] The meninges consist of multiple layers of membranes that protect the central nervous system from injury and infection.[3,4] Some intracellular bacteria, including *Salmonella typhimurium* (*S. typhimurium*) and *Mycobacterium tuberculosis* (*M. tuberculosis*), can invade the meninges to cause severe brain pathology.[5-7] Bacterial meningitis is a life-threatening infection with extremely high mortality; however, the mechanism by which bacteria invade the brain remains unclear. Efforts to understand the factors that promote neuroinvasion and host responses to these pathogens are urgently needed.

Phagocytes, such as neutrophils and macrophages, play dominant roles in host response to intracellular bacteria.[8] The primary function of phagocytes is to uptake and destroy microorganisms and cell debris through tissue circulation and migration.[9,10] Via phagocytosis and antigen presentation, phagocytes act rapidly as the first line of defense against bacterial infections. However, intracellular bacteria can employ sophisticated strategies to replicate within phagocytes and evade host immune surveillance.[11-13] Recently, Marc Lecuit et al. showed that the surface protein internalin B of *Listeria monocytogenes* (*L. monocytogenes*) protects infected monocytes from immune killing by CD8$^+$ T cells, thus prolonging the lifespan of infected cells and facilitating the spread of infection throughout the central nervous system.[13] In this context, phagocytes, especially monocytes and macrophages, might

play important roles during neuroinvasion, but exactly how intracellular bacteria take

advantage of infected macrophages is not well understood.

Infectious diseases and host metabolism are closely linked. Metabolic signaling helps

awaken the immune system to resist pathogenic invasion. For example, glycolysis

bolsters T cell immunity to prevent *L. monocytogenes* infection.[14] In addition, many

studies have shown that bacteria can actively or passively manipulate host metabolites

to evade clearance by the host immune system.[11,15,16] *Ruminococcus gnavus* initiates

biosynthesis of phenethylamine and tryptamine, which exacerbates the progression of

diarrhea-predominant irritable bowel syndrome.[11] *Legionella pneumophila* induces

Warburg-like metabolism by secreting the type 4 secretion system effector MitF and

decreases oxidative phosphorylation (OXPHOS) in infected macrophages to increase

the pathogen's likelihood of survival.[15] A recent report indicates that $Nav1.8^+$

nociceptors send signals to meningeal macrophages via the calcitonin gene-related

peptide, and this neuroimmune axis suppresses host defenses and exacerbates

bacterial meningitis.[16] These findings imply a potentially close connection between

host metabolism and bacterial brain invasion.

Ketogenesis, induced by fasting or dietary supplementation, is a metabolic pathway

that provides an alternative form of energy for the body and is closely related to host

immune response.[17,18] A recent study reports that in pneumonia caused by

Pseudomonas aeruginosa, ketone bodies are recycled to the lungs via OXPHOS to

maintain airway homeostasis. [19] Consistently, a ketogenic diet is reported to activate

the expansion of protective γδ T cell responses against influenza virus infection. [20]

However, researchers have also found that pathogens can achieve intracellular

survival and infection progression by manipulating host ketogenesis. [21] Mice fed a

ketogenic diet develop a hyperinflammatory response to systemic lipopolysaccharide

and an increased mortality rate, phenomena that are independent of diet-induced

microbiota and hyperglycemia. [21] Therefore, the role of ketogenesis in the progression

and migration of infections is controversial. [22] β-hydroxybutyrate (BHB) produced by

ketogenesis is well known to promote the accumulation of intracellular lipid bodies,

which serve as both a source of nutrients and a secure niche for intracellular bacteria. [23]

However, how ketogenesis and BHB participate in regulating monocytic/macrophagic

homeostasis and their roles in bacterial neuroinvasion are poorly defined.

In this study, we discovered a novel subpopulation of $CD36^+$ macrophages that served

as immune 'Trojan horses', crossing the blood-brain barrier (BBB) and enabling

intracellular bacterial neuroinvasion. This macrophage subset has a foamy phenotype,

which improves the cells' ability to stay and migrate vertically on the cerebrovascular

endothelial surface under the mechanical stress of blood flow. Furthermore, we found

that serum BHB, which might be produced by a ketogenic diet, exacerbated bacterial

neuroinvasion by increasing the formation and promoting the survival of foamy

macrophages in peripheral blood. Our study outlines a new common strategy

employed by intracellular bacteria in brain invasion, which could be a potential

therapeutic target in the treatment of bacterial meningitis.

Results

CD36[+] foamy macrophages are potential carriers of bacteria during brain

invasion

To study the role of immune cells in bacterial neuroinvasion, we performed single-cell

RNA-sequencing (scRNA-seq) of peripheral blood mononuclear cells (PBMCs)

collected from mice in an acute meningitis model of *L. monocytogenes* infection

(Figure 1A). We conducted scRNA-seq 24 hours post infection, at which timepoint

bacteria had massively invaded the brain and mice began to die (Figure S1A and S1B).

Immune cell lineages were identified using canonical surface markers of dendritic

cells (DCs: *Xcr1, Flt3, CD209a*), NK cells (*Nkg7*), T cells (*CD3*),

monocyte/macrophages (Mo/Mφ: *Fcgr1, Adgre1*), B cells (*CD79*), neutrophils

(*S100a9, S100a8*; Figure 1B and 1C).[24] Among these lineages, Mo/Mφ are involved in

phagocytosis and intracellular killing of microorganisms; however, intracellular

bacteria can also live in these cells to achieve long-term latent infection.[25] In this case,

Mo/Mφ (n=3535 cells) were extracted and reclustered *in silico* to precisely delineate

subpopulations and their distributions before and after neuroinvasion. Based on

differentially expressed cell surface markers and transcripts, we annotated Mo/Mφ

into 6 subpopulations (Figure 1D, S1C). Results showed that the percentage of Cluster

2 cells was sharply increased in the neurolisteriosis group (Figure S1D). Further, we

found that Cluster 2 cells expressed high levels of genes related to lipid accumulation,

including *CD36*, peroxisome proliferator–activated receptor gamma (*Pparg*), *CD9*,

apolipoprotein E epsilon (*ApoE*), fatty-acid binding protein 4 (*Fabp4*), and triggering

receptor expressed on myeloid cells 2 (*Trem2*; Figure. 1E). [26-30] These data implied that

the functional role of Cluster 2 macrophages was closely related to lipid metabolism.

As previously reported, lipid accumulation in macrophages easily exceeds their

homeostatic capacity, triggering lipid droplet formation and causing the macrophages

to have a foamy appearance. [31,32] To understand the characteristics of Cluster 2

macrophages, $CD36^+$ and $CD36^-$ macrophages were isolated from peripheral blood by

fluorescence-activated cell sorting. Compared with $CD36^-$ macrophages, the $CD36^+$

subpopulation exhibited significantly higher granularity (SSC^{hi}; Figure S1E), which

indicated the number of proteins or organelles they contained and the protrusions on

their surfaces. Increased granularity is considered a characteristic of foamy

macrophages.[33] The sorted macrophages were cultured *in vitro* overnight and then

labeled with BODIPY FL, a lipophilic probe for polar and neutral lipids. Results

showed that the $CD36^+$ subpopulation accumulated significantly more lipid droplets

than the $CD36^-$ macrophages (Figure S1F). We also performed Oil Red O staining to

validate this finding (Figure S1F). To confirm the lipid uptake ability of the $CD36^+$

macrophages, we used fluorescently labeled 1,1′-dioctadecyl-3,3,3′,3′-

tetramethylindocarbocyanine perchlorate (Dil)–conjugated oxidized low-density

lipoprotein (oxLDL) particles and our findings showed that $CD36^+$ macrophages took

up more than twice as much ox-LDL as CD36$^-$ macrophages (Figure S1G). These

data confirmed the foamy appearance of the Cluster 2 macrophages during brain

invasion.

Next, we performed flow cytometry to detect percentage changes in foamy

macrophages over time during neurolisteriosis (Figure S1E). The results showed that

the percentage of CD36$^+$ foamy macrophages in peripheral blood steadily increased

over time during neuroinvasion (Figure 1F). The CD36$^+$ subgroup could exceed 52%

of all macrophages (Figure 1G). In addition, the percentage of foamy macrophages in

peripheral blood was positively associated with bacterial load in the brain (Figure 1H).

These data implied that CD36$^+$ foamy macrophages might be involved in bacterial

neuroinvasion. Furthermore, immunofluorescence co-localization analysis was used

to validate this hypothesis. The results showed that the number of CD36$^+$ foamy

macrophages attached to cerebrovascular endothelial cells (CVECs) increased as

neuroinvasion progressed in the acute meningitis model (Figure 1I). Taken together,

these results suggested that CD36$^+$ foamy macrophage might play an important role in

bacterial brain invasion.

CD36$^+$ foamy macrophages facilitate transport of intracellular bacteria across

the blood-brain barrier

Next, we sought to understand how necessary CD36$^+$ foamy macrophages were to

bacterial neuroinvasion. Using mouse primary CVECs and neurons in a Transwell

system, we constructed an *in vitro* BBB model (Figure 2A). Ox-LDL has been

reported to induce foamy cell formation by enhancing CD36 expression[34]; therefore,

we converted peritoneal macrophages isolated from mice into CD36$^+$ foamy

macrophages by treating them with different concentrations of ox-LDL for 24 hours

(Figure S2A). Subsequently, macrophages were infected with *L. monocytogenes*

(MOI=10) and then inoculated in the upper Transwell chamber of the BBB model. We

confirmed that intracellular bacterial load increased with the transformation of

macrophages into the foamy phenotype (Figure S2B). Of note, the assistance of ox-

LDL–stimulated macrophages significantly increased the number of *L.*

monocytogenes organisms crossing the BBB in a dose-dependent manner (Figure 2B).

Using crystal violet staining, we also demonstrated a dose-dependent increase in the

number of macrophages penetrating the endothelial monolayer upon stimulation with

ox-LDL (Figure 2C and 2D).

To understand whether the increase in CD36$^+$ foamy macrophages favored the transfer

of *L. monocytogenes* into the brain *in vivo*, we performed adoptive macrophage

transfer experiments through tail vein injection based on a cyclophosphamide (CTX)-

induced immunosuppressive mouse model (Figure 2E). Consistent with previous

studies, *L. monocytogene*-induced clinical phenotype and mortality were relieved in

the mice transferred with purified peritoneal macrophages (PMφ group; Figure 2F,

S2C and S2D).[35,36] However, body weight was reduced, comprehensive clinical scoring

was increased, and survival time was shortened in the mice transferred with CD36$^+$

foamy macrophages compared with the control (FMφ group; Figure 2F, S2C and

S2D). Furthermore, we showed that brain bacterial loads in mice transferred with

CD36$^+$ macrophages were significantly higher than those in mice transferred with

purified peritoneal macrophages (Figure 2G). Interestingly, bacterial loads did not

differ among the lung, spleen, and liver (Figure S2E). We also tested proinflammatory

factor levels in plasma and brain homogenate. The results showed that levels of

interleukin 6 (IL-6) and tumor necrosis factor alpha (TNF-α) were remarkably

increased (Figure 2H and 2I), indicating a more severe phenotype of meningitis in the

foamy macrophage transferred group. Pathological histology analysis confirmed the

existence of more severe inflammation and neuronal damage in the cerebral cortices

in mice transferred with foamy macrophages than in those of control (Figure S2F and

S2G). In addition, when mice were infected with a relatively low dose (5×10^7

CFU/mouse) of *L. monocytogenes*, the difference in bacterial load in brain tissues

between foamy macrophage transferred mice and the control group was even greater

than that under high dose condition (Figure S2H and S2I). These data suggested that

neuroinvasion mediated by CD36$^+$ foamy macrophages were not restricted to an acute

meningitis model induced by high dose bacteria.

We then wondered whether CD36$^+$ foamy macrophages might play the same role

during brain invasion by other intracellular bacteria. In the relevant experiment, a

positive correlation between CD36$^+$ foamy macrophages and neuroinvasion was

observed in meningitis models of the intracellular bacterium *S. typhimurium* (Figure

S3A and S3B). We observed the same result in neuroinvasion model of Methicillin-

resistant *Staphylococcus aureus* (MRSA; Figure S3C and S3D), which, although not a

classical intracellular pathogen, exhibits significant intracellular habitation.[37] However,

this finding was not validated in a neuroinvasion model of the extracellular bacterium

Streptococcus pneumoniae (*S. pneumoniae*; Figure S3E and S3F). Next, we

performed adoptive macrophage transfer experiments to further verify the carrier role

of CD36$^+$ foamy macrophages in intracellular bacterial invasion of the brain. We

again observed differences in brain tissue bacterial loads between the foamy

macrophage transferred and control groups in the meningitis models of MRSA (Fig.

S3G-S3K), but not in models using *S. pneumoniae* (Figure S3L-S3O). Taken together,

these data suggested that CD36$^+$ foamy macrophages greatly contributed to the

process of intracellular bacterial neuroinvasion.

Proprotein convertase subtilisin/kexin 9 (PCSK9) belongs to the proprotein

convertase family, whose members participate in cholesterol and fatty acid (FA)

metabolism. [38,39] Intracellular low-density lipoprotein cholesterol (LDL-C)

accumulates because LDL receptor located on the cell membrane cannot be degraded

normally in PCSK9$^{-/-}$ mice.[40,41] We found that PCSK9-deficient macrophages had

significantly higher lipid content compared with control (Figure S4A and S4B) and

that more *L. monocytogene* organisms parasitized PCSK9$^{-/-}$ macrophages than they

parasitized wild-type (WT) cells (Figure S4C). During neurolisteriosis, PCSK9$^{-/-}$ mice

had higher percentages of CD36$^+$ foamy macrophages in peritoneal blood than their

littermates (Figure S4D). Therefore, we used PCSK9$^{-/-}$ mice as a model of

spontaneously forming foamy macrophages, and verified the reliability of our

findings through adoptive macrophages transfer. After recovery from transfer for 12

hours, recipient mice were intraperitoneally injected with 2×10^8 CFU of *L.*

monocytogenes; 12 h later, samples were collected for further examinations (Figure

2J). We found that brain bacterial loads were significantly higher in PCSK9$^{+/+}$

recipients that had been transferred with PCSK9$^{-/-}$ macrophages than in those that had

received WT macrophages (Figure 2K). Similar results were observed in the PCSK9$^{-/-}$

recipients (Figure 2K). However, differences in bacterial load manifested in brain

tissues only, not in other organs such as the lung, spleen, and liver (Figure S4E).

Consistently, the mortality rate, and pro-inflammatory factors in the plasma and brain

homogenate were aggravated in both PCSK9$^{+/+}$ and PCSK9$^{-/-}$ recipients by transfer of

PCSK9$^{-/-}$ macrophages (Figure 2L, S4F and S4G). Since *L. monocytogene* is a

foodborne pathogen that can invade the body through the gastrointestinal tract, we

further confirmed the above-mentioned finding by oral administration of *L.*

monocytogenes in the same experimental model (Figure 2M and 2N, S4H-S4J). Taken

together, these data indicated that excess lipid accumulation contributed to the foamy

appearance of macrophages, thereby facilitating neuroinvasion by intracellular

bacteria.

Shear flow analysis demonstrates increased adhesion and resistance of foamy

macrophages during neuroinvasion

We next asked how CD36$^+$ foamy macrophages exploited their properties to

accumulate on the cerebral blood vessel wall and participate in the process of

intracellular bacterial neuroinvasion. Using a two-dimensional live-cell flow chamber

assay with a microfluidic chip[42], we deciphered the multistep process of how

macrophages adhered to and moved on the surface of vascular endothelium under

physiological flow (Figure 3A). Cellular adhesion ability was calculated by counting

total macrophages adhering to the endothelial monolayer under a given shear stress.

We found that as the percentage of CD36$^+$ foamy macrophages increased, so did the

number of cells adhering to the endothelial monolayer (Figure 3B and 3C). Because

adherent cells could be detached by an increase in shear stress, we tested the

dynamics of CD36$^+$ foamy macrophages' shear resistance. The results showed that

such resistance was significantly stronger than that of control macrophages: CD36$^+$

foamy macrophages outstandingly resisted high shear stress (Figure 3D). Further, we

investigated the effect of endothelial density on macrophagic adhesion. Results

showed that endothelial cell density, rather than cell type, greatly affected

macrophagic adhesion, and foamy macrophages tended to adhere to the surface of

dense endothelial cells (Figure S5A-S5D).

Once attached to the endothelial monolayer, the macrophage crawls along it and finds

a place to cross it in the presence of blood flow. [43,44] We next tested the performance of

CD36$^+$ foamy macrophages in post-arrest crawling on the endothelial monolayer. We

acquired sequential images of macrophages at a shear stress of 2 dyn/cm^2 and then

analyzed them by defining various parameters of crawling dynamics (Figure S5E).

The observations were presented by crawling trajectories of each individual cells

(Figure 3E), and we found that the crawling ability of adherent CD36$^+$ foamy

macrophages was dampened, reflecting in shortened crawling displacement and

slower crawling speed (Figure 3F and 3G). However, we saw no obvious differences

in crawling duration (Figure S5F). Directionality is also a critical aspect of cell

crawling dynamics, it is quantified as the ratio of displacement along shear flow to the

perpendicular. Our results showed that the directionality of macrophages was slightly

but significantly enhanced by ox-LDL stimulation, implying that macrophagic

foaminess tended to drive cell crawling in the vertical direction (Figure 3H). We also

compared a key dynamic parameter of transmigration duration, which is defined as

the interval from the timepoint of cell arrival at the site to the endpoint of completion

of transmigration beneath the endothelial monolayer (Figure S5G). The result showed

that transmigration duration was significantly shortened in CD36$^+$ foamy macrophage

(Figure 3I).

Considering that protrusion is critical for cell crawling and migration, we then

investigated whether intercellular lipid could affect the formation of protrusions. The

results showed that the numbers, as well as the lengths of protrusions were

significantly increased in CD36$^+$ foamy macrophages (Figure 3J and 3K). We further

quantified macrophagic cell area and found that the mean area of foamy macrophages

was significantly larger than the control (Figure 3L). Additionally, we examined

various integrins and adhesive ligands expressed on the cell surface, which have been

reported to be involved in adhesion and migration.[45] The results showed that the

expression levels of α-integrins, β-integrins and intercellular adhesion molecules were

significantly upregulated in the infected CD36$^+$ foamy macrophages (Figure 3M).

Collectively, these data suggested that the foaminess of macrophages greatly

enhanced their ability to adhere under mechanical stress, predisposing them to stay

and to migrate vertically on the surface of CVECs; this might be attributable to the

formation of protrusions as well as the expression of integrins and adhesion ligands.

β-hydroxybutyrate promotes foamy macrophage formation and exacerbates

neuroinvasion

Having shown foamy appearance to be closely linked to the excess lipid accumulation

in macrophages, we performed lipid metabolomics analysis of serum samples to

reveal how foamy macrophages were formed during bacterial neuroinvasion.

Principal component analysis (PCA) showed that distinct metabolite genotype clusters

were generated in the neurolisteriosis group (Figure 4A). The volcano plot indicated

that 70 FAs were upregulated and 71 FAs were downregulated ($p < 0.05$, |log$_2$(fold

change) | > 1) in the neurolisteriosis group (Figure 4B, Table S1). The 70 upregulated

FAs were classified into four subclusters by the number of carbon atoms in the alkyl

chain. We found that BHB, which has been reported to favor the foamy phenotype of

macrophages [46], was the only short-chain FA among all upregulated FAs (Figure 4C).

KEGG-based analysis revealed that metabolites related to the synthesis and

degradation of ketone bodies were enriched (Figure S6A), and metabolomics data

showed that BHB was significantly increased during neuroinvasion (Figure S6B).

Based on these data, we detected serum BHB level in the acute meningitis model of *L.*

monocytogenes infection. The results showed that BHB level climbed steadily over

time and then decreased slightly (Figure 4D). Consistently, an upregulation of genes

controlling ketogenic enzymes (HMGCS2 and CPT1A) in the liver was observed as

infection persisted (Figure S6C and S6D). Meanwhile, mice experienced weight loss

and decreased appetite during the experiments, which was partly related to enhanced

ketogenic ability after infection (Figure S6E and S6F). Furthermore, a steady increase

of serum BHB was also observed in the meningitis models established using *S.*

typhimurium and MRSA (Figure S6G and S6H). These data confirmed that production

of BHB was significantly induced during bacterial neuroinvasion.

As previously reported, BHB promotes the foamy phenotype of macrophages via the

hydroxycarboxylic acid receptor 2 (HCAR2).[23] Therefore, we next used HCAR2

knockout mice to study whether BHB could promote bacterial neuroinvasion. We

confirmed that lipid accumulation and percentage of CD36[+] macrophages all

increased dramatically upon BHB stimulation, and there was no significant difference

when HCAR2 was knocked out (Figure 4E-4G). BHB inhibits adenylyl cyclase

activity through the HCAR2 receptor and reduces cellular cyclic adenosine

monophosphate (cAMP) levels, thereby protecting lipid droplets from lipolysis by

hormone-sensitive lipases.[23] As a result, we found that cAMP pretreatment eliminated

the increase in lipid content and intercellular bacterial burden stimulated by BHB;

however, this phenomenon was not found in HCAR2-deficient macrophages (Figure

S6I and S6J). By using the *in vitro* BBB model (Figure 2A), the number of bacteria

crossing the barrier increased significantly in response to BHB stimulation in controls,

but not in HCAR2$^{-/-}$ macrophages (Figure 4H). Next, peritoneal macrophages from

HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice were isolated to further assess the effects of BHB on

the biophysical dynamics of macrophages. The data demonstrated that BHB

significantly altered the static or shear-resistant adhesion of control macrophages, but

not of HCAR2$^{-/-}$ macrophages (Figure 4I-4K). We also found that BHB greatly

impaired the crawling ability, including shortened crawling displacement, slowed

down crawling speed and unpredictable directionality of macrophages (Figure 4L-4O).

In addition, the transmigratory ability of macrophages was significantly increased

when cells were pretreated with BHB (Figure 4P). Consistently, the expression levels

of the α-integrin family were significantly upregulated by BHB treatment in controls,

but not in HCAR2-deficient macrophages (Figure 4Q). Taken together, these data

indicated that BHB facilitated the foamy phenotype of macrophages through the

HCAR2 receptor, thereby affecting the process of bacterial neuroinvasion.

β-hydroxybutyrate promotes the survival of macrophages via HCAR2-mediated

calcium influx

Because the foamy stage appears to be a terminal point of macrophage differentiation

[47], we wondered whether, in addition to converting peritoneal macrophages to the

foamy phenotype, cell survival also contributed to the increase in absolute quantity of

foamy macrophages induced by BHB treatment. The results showed that the lactate

dehydrogenase (LDH) activity in the culture medium was significantly reduced in

response to BHB treatment, indicating a decrease in cell death (Figure 5A). Propidium

iodide (PI) staining was used to evaluate the membrane integrity of infected

macrophages, and we found that the proportion of PI^+ cells decreased from 59.7% to

24.5% under BHB stimulation, with no significant difference in cell viability between

the BHB treatment and control groups when HCAR2 was knocked out (Figure 5B and

5C). Calcium influx is generally recognized as a signal that triggers cell membrane

repair.[48] To investigate the impact of BHB-HCAR2 signaling on cytosolic calcium

concentration, infected macrophages were loaded with Fluo-4 AM, a fluorescent Ca^{2+}

indicator. The results showed that BHB induced remarkable accumulation of

intracellular calcium, and HCAR2 knockdown eliminated calcium influx in

macrophages (Figure 5D). Pretreatment of macrophages with BAPTA, a Ca^{2+} chelator,

reversed the BHB-induced increase in cell survival (Figure 5E and 5F). Further, we

tested whether calcium flux was associated with foamy macrophage-mediated

intracellular bacterial penetration of the BBB. As a result, inhibition of calcium flux

had no effect on bacterial uptake by macrophages in the upper chamber (Figure S7A);

however, we observed differences in bacterial loads in the macrophages that had

crossed the barrier (Figure 5G).

We next investigated how BHB maintained the survival of foamy macrophages. As

previously reported, calcium influx initiates membrane repair mainly through

recruitment of the endosomal sorting complexes required for transport (ESCRT)

machinery.[49] Our data showed that BHB treatment significantly increased the number

of puncta that were positive for charged multivesicular body protein 4B (CHMP4B),

which indicated assembly of ESCRT in the infected macrophages; HCAR2

knockdown or BAPTA treatment could block this membrane repair process (Figure

5H and 5I, S7B and S7C).Cellular membrane repair also relies on lysosomes, which

can rapidly fuse with the plasma membrane when intracellular Ca^{2+} concentration

rises. Therefore, we examined lysosomal exocytosis in the infected macrophages

using exofacial lysosomal-associated membrane protein 1 (LAMP1) and the release of

lysosomal enzymes, such as acid phosphatase (ACP), β-galactosidase (β-gal) and β-

hexosaminidase (β-hex). The results confirmed that BHB promoted the lysosomal

exocytosis in infected macrophages through the HCAR2 receptor (Figure 5J-5L, S6D-

S6F). These data collectively indicated that BHB signaling activated Ca^{2+}-mediated

membrane repair, thereby maintaining the survival of foamy macrophages via the

HCAR2 receptor.

Ketogenesis exacerbates intracellular bacterial neuroinvasion

BHB can be synthesized via the metabolism of FAs and ketogenesis.[50] To understand

the relationship between ketogenesis and bacterial meningitis, we gave mice different

doses of ketone ester drink for 4 weeks, and then infected them with *L.*

monocytogenes via intraperitoneal injection (Figure 6A). The data showed that ketone

ester elevated the levels of serum BHB and the proportion of foamy macrophages in

peritoneal blood (Figure 6B and S8A), accompanied by remarkable increases in the

clinical scores, mortality rate and bacterial load in brain tissues (Figure 6C and 6D,

S8B). Dietary modulation can increase and maintain circulating metabolites in human

circulation, and a growing amount of evidence demonstrates that a ketogenic diet

elevates BHB levels.[51] The ketogenic diet is a low-carb, high-fat diet that causes

weight loss and provides numerous health benefits, including alleviating symptoms of

neurodegenerative diseases, cardiovascular diseases, and cancers.[52-55] We next asked

whether a ketogenic diet was associated with neurolisteriosis. HCAR2$^{-/-}$ mice and

their littermates were fed a normal or ketogenic diet for 4 weeks respectively, and

then infected with *L. monocytogenes* by intraperitoneally injection (Figure 6E). As a

result, the lifespan of HCAR2$^{+/+}$ mice fed a ketogenic diet was significantly shortened

under *L. monocytogenes* infection, and the survival disadvantage caused by dietary

modulation was abrogated by HCAR2 knockout (Figure 6F). Consistently, the clinical

scores (Figure. S8C), percentages of foamy macrophages in peritoneal blood (Figure

6G), the intracellular bacterial burden of macrophages (Figure 6H), and bacterial

loads in brain tissue (Figure 6I) were all greatly increased in HCAR2$^{+/+}$ mice on a

ketogenic diet. Interestingly, mice in this group lost less weight than mice in the

control group, and experienced greater spread of bacteria to other organs (Figure 6I

and S8D). We then examined the levels of pro-inflammatory factors in the plasma and

brain homogenate. The results showed that a ketogenic diet significantly increased the

levels of pro-inflammatory cytokines and induced more severe neuronal damage in

HCAR2$^{+/+}$ mice, while it had no effect on HCAR2$^{-/-}$ mice (Figure 6J-6M). These data

were confirmed by oral administration of *L. monocytogenes* in the same experimental

model (Figure S8E-S8H). We also used MRSA to further verify our findings. The data

indicated that, as expected, the ketogenic diet promoted brain invasion by MRSA and

exacerbated symptoms of meningitis (Figure S8I-S8K). Taken together, our data

showed that a ketogenic diet could exacerbate bacterial neuroinvasion and hasten the

demise of mice, primarily depending on elevations in serum BHB via the HCAR2

receptor.

Serum β-hydroxybutyrate fuels the symptoms of human bacterial meningitis

Our data thus far suggested that BHB-induced CD36^{+} foamy macrophages played an

important role in bacterial neuroinvasion in mice. Next, we sought to assess whether

the accumulation of foamy macrophages or BHB could also be identified and were

associated with bacterial meningitis in human samples. Gender- and age-matched

bacterial meningitis subjects (n=62) and healthy individuals (n=60) were recruited

and peripheral blood samples were collected for analysis (Figure 7A). We observed no

significant differences in blood glucose or insulin concentrations between the two

groups (Figure S9A and S9C), nor was there any correlation between these parameters

and BHB levels (Figure S9B and S9D), excluding the involvement of glucose and

insulin in the differential production of BHB. As a result, serum BHB levels were

significantly elevated in meningitis patients compared with healthy controls,

suggesting an infection-induced ketogenesis (Figure 7B). Interestingly, serum

cholesterol levels, including total cholesterol (TC) and LDL-C were increased in

meningitis patients compared with healthy controls, while high-density lipoprotein

cholesterol (HDL-C) was slightly but not significant reduced (Figure S9E-S9G). To

investigate whether BHB was associated the severity of infection, we further divided

patients into three groups, high BHB (in the top 1/3 among samples tested, n=21),

moderate (middle 1/3, n=20) and low (bottom 1/3, n=21; Figure 7A). Clearly, patients

with high BHB levels exhibited higher clinical scores than patients with low BHB

levels (Figure 7C). Consistently, high-BHB patients had higher cell counts and more

protein in the cerebrospinal fluid (CSF; Figure 7D and 7E); higher infection indices

such as white blood cell counts (WBC; Figure 7F) and procalcitonin (PCT; Figure

7G); and higher values for markers of inflammatory conditions such as erythrocyte

sedimentation rate (ESR; Figure 7H) and C-reactive protein (CRP; Figure 7I),

suggesting increased disease severity. These data implied that infection-induced

ketogenesis might provide important fuel for severe bacterial neuroinvasion.

To explore the dynamics of CD36$^+$ foamy macrophages and BHB in the pathogenesis

of bacterial meningitis, we followed-up with 7 individuals during their hospitalization.

We found that CD36$^+$ foamy macrophages and BHB levels in patients tended to

increase during disease progression, and decrease during the recovery stage (Figure 7J

and 7K). To further confirm the relationship between foamy macrophages and

bacterial meningitis pathology, we collected PBMCs from patients or healthy controls

for analysis. Compared with healthy controls, the expression levels of genes involved

in cholesterol uptake were significantly increased, while those of genes related to

cholesterol utilization and excretion were decreased in macrophages from meningitis

patients (Figure 7L). Taken together, these data indicated that BHB fueled human

bacterial meningitis by promoting the development of CD36$^+$ foamy macrophages.

Discussion

Our findings suggested that CD36$^+$ foamy macrophages might serve as carriers for

intracellular bacteria that colonize in the brain, facilitating the penetration of these

bacteria across the BBB. Also, we found that BHB level increased significantly in the

peripheral blood of meningitis patients. Mechanistically, BHB regulated the dynamics

of and morphological changes to foamy macrophages and promoted cell survival by

activating the HCAR2 receptor, which is shown here to be a key signaling molecule

for intracellular bacterial neuroinvasion. Furthermore, under ketogenic diet conditions,

elevated BHB levels promoted the formation of foamy macrophages, thereby

exacerbating bacterial transport to the brain.

Normally, the BBB prevents bacteria or viruses from crossing unless there is

significant trauma to the brain. However, specific bacteria can penetrate the BBB via

their unique mechanisms and cause severe brain infections.[1,13] For example, *S.*

pneumoniae co-localizes with polymeric immunoglobulin receptor (pIgR) and platelet

endothelial cell adhesion molecule (PECAM), leading to brain invasion.[56]

Streptococcus agalactiae enhances the production of pro-inflammatory cytokines and

intercellular adhesion molecule-1(ICAM-1) by endothelial cells via activation of the

inducible nitric oxide synthase (iNOS) pathway, resulting in direct BBB disruption.[1]

Among intracellular bacteria, internalin B of *L. monocytogenes* is reported to protect

infected monocytes from Fas-mediated cell killing by $CD8^+$ T cells and prolong their

life-span[13]; while *M. tuberculosis* might utilize leucine-responsive regulatory protein

(Lrp) to inhibit the production of pro-inflammatory cytokines, thereby promoting their

survival and neuroinvasion.[57] In this context, it is likely that intracellular bacteria use

the host's cells, especially immune cells, to achieve brain invasion. In this study, we

discovered that $CD36^+$ foamy macrophages could be hijacked by intracellular bacteria

and serve as vehicles for neuroinvasion. Compared with peritoneal macrophages,

foamy macrophages tend to have more protrusions and greater cell area, which make

them more conducive to colonization on CVECs under the pressure of blood flow.

Pathogens can exploit most of the host's nutrients, including glucose, glycerol, FAs,

N-acetylglucosamine (GlcNAc), and several other carbon substances, for invasive

transmission.[58] The T3SS effector SopE2 of *S. Typhimurium* utilizes accumulated

glycolytic intermediates as carbon sources, leading to intracellular replication and

pathogenicity.[59] Our findings demonstrated that intracellular bacteria can actively

influence host free FA metabolism (Figure 4A-4C) and reprogram macrophages,

thereby exacerbating neuroinvasion. Foamy macrophages are formed through

dysregulation of lipid metabolism, and foamy phenotype impairs the immune function

of macrophages and contributes to pathogenesis. For example, foamy macrophages

play a key role in the initiation, progression, and instability of atherosclerosis.[60,61] In

multiple sclerosis, myelin-laden foamy macrophages in brain lesions are associated

with demyelinating activity and chronic active lesions.[62] During tuberculosis, foamy

macrophage necrosis expands tuberculous lesions in lung parenchyma, causing

progressive destruction of lung tissue and loss of lung function in patients with active

disease.[63] In this study, we discovered that BHB produced from ketogenic diets

promoted the formation of $CD36^+$ foamy macrophages in peripheral blood, thereby

facilitating bacterial neuroinvasion. Interestingly, we found that serum cholesterol

levels, including TC and LDL-C were significantly higher in meningitis patients than

in healthy controls (Figure S9E and S9G). We hypothesize that cholesterol or other

lipid metabolites might also contribute to the formation of $CD36^+$ foamy macrophages.

However, this hypothesis should be further verified experimentally.

Our biomechanical data indicated that $CD36^+$ foamy macrophages adhered more

strongly to the endothelial surface than normal macrophages, and that this adhesion-

enhancing effect was more pronounced when endothelial cells were dense (Figure 3B

and 3C, S5A and S5B). This suggested that foamy macrophage-mediated bacterial

invasion affected the brain to a greater extent than other tissues. Accordingly, we

showed that the increase in $CD36^+$ foamy macrophages in peripheral blood led to the

transfer of bacteria mainly to the brain, rather than to other organs (Figure 2G and

S2E). However, we found that a ketogenic diet and ketone ester supplementation

significantly increased bacterial load not only in the brain, but also in other tissues

such as the liver and lung (Figure 6D and 6I). This suggested that ketogenesis, in

addition to promoting the generation of foamy macrophage, might play other

unknown roles in the intracellular bacterial infections. Therefore, the role of

ketogenesis in the formation of $CD36^+$ foamy macrophages and the contribution of

ketogenesis to bacterial neuroinvasion should be further determined.

Host basal metabolic status may also greatly influence bacterial neuroinvasion by

modifying immune cells. A previous study found that cardiac endothelial cells can

oxidize ketone bodies and promote cell proliferation, migration, and blood vessel

sprouting.[64] Increasing ketone body levels with a ketogenic diet increased endothelial

cell proliferation and prevented vascular rarefication in a hypertrophied mouse

model.[64] Whether ketogenesis directly affects the development or permeability of the

BBB, leading to an increase in bacterial invasion, is unclear. Meanwhile, our study

found that mice experienced weight loss and decreased appetite during the

experiments (Figure S6E and S6F). Therefore, we currently cannot determine whether

the elevated BHB levels were caused directly by pathogenic bacteria or by hepatic

ketogenesis.

This study demonstrated that a BHB-rich environment created by the ketogenic diet

rendered mice more susceptible to *L. monocytogene*-induced neuroinvasion.

Mechanistically, BHB protected $CD36^+$ foamy macrophages from senescence via

HCAR2-mediated membrane repair. However, inhibition of HCAR2 signaling or

calcium influx reversed the improvement in foamy macrophage survival but did not

completely reverse the exacerbation in bacterial neuroinvasion induced by BHB

(Figure 5E-5G). These data suggested that BHB-induced bacterial brain invasion was

not solely dependent on HCAR2 signaling but that other potential mechanisms might

also be involved in this process. However, this hypothesis requires further

experimental verification.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No.

92369115), the Fundamental Research Funds for the Central Universities (No.

2023CDJXY-009 and 2019CDYGZD009), Natural Science Foundation of Chongqing,

China (No. CSTB2023NSCQMSX0402), and Chongqing Talents: Exceptional Young

Talents Project (No. cstc2021ycjh-bgzxm0099). The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Author Contributions

H.W., Q.Z., X.L. and Z.S. conceived and designed the study. Z.S., H.Y., H.F., Q.H.,

B.F., Y.X. and D.G. performed the animal experiments. Z.S., K.Y., Q.H., Q.L., Y.D.

and Xushuo Z. carried out healthy volunteer and patient recruitment and sample

collection. Z.S., N.L. and X.S. carried out the live-cell flow chamber assays. H.W.,

Z.S., K.Y., H.Y. and Xiaokai Z. collected and analyzed the data. X.T., S.F., J. L., R.Y.,

W.G., X.M. and M.L. contributed to experimental material and insightful suggestions.

H.W., X.L., Xushuo Z. and Z.S. wrote the manuscript. H.W. and Q.Z. coordinated the

project. All authors read and approved the final manuscript.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1. Identification of CD36[+] foamy macrophage accumulation during

neurolisteriosis.

(A) Summary of experimental design (n=5 control mice and n=7 mice infected with

2×10^8 CFU *L. monocytogenes* for 24 hours).

(B) Uniform Manifold Approximation and Projection (UMAP) representation of

integrated transcriptome-based clustering. A total of 17351 peripheral blood

mononuclear cells (PBMCs) were clustered and major immune cell lineages were

identified.

(C) Dot plot showing the expression of selected transcripts in immune cell clusters.

All transcripts were significantly enriched in the pertinent clusters with adjusted p <

0.05.

(D) A t-distributed Stochastic Neighbor Embedding (tSNE) analysis of the re-

clustered cells corresponding to monocytes and macrophages (baseline, n=1804; neurolisteriosis, n=1731).

(E) Expression of the indicated transcripts (encoding genes associated with lipid accumulation) was projected onto a monocyte/macrophage tSNE plot.

(F-G) An acute mouse neurolisteriosis model was generated by intraperitoneal injection of 2×10^8 CFU *L. monocytogenes.* Quantification of CD36$^+$ foamy macrophages in PBMCs (F) and total macrophages (G) was detected by flow cytometry (n=5 for each group). Experiments were performed in duplicate. Data are presented as mean ± SEM. *p < 0.05, **p < 0.01.

(H) Correlation between CD36$^+$ foamy macrophages and bacterial burdens in brain during the neurolisteriosis. R, correlation coefficient.

(I) Immunofluorescence staining of CD36$^+$ foamy macrophages in cerebral vasculature of mice with neurolisteriosis. CD36$^+$ foamy macrophages were stained with CD11b (red) and CD36 (green). The boundary of vascular was depicted in dashed lines. Scale bar, 100 μm.

Abbreviations: PBMC, peripheral blood mononuclear cell; i.p., intraperitoneally; FMφ, foamy macrophages; CFU, colony forming units.

Figure 2. Infected foamy macrophages transfer bacteria into brain parenchyma *in vitro* and *in vivo*.

(A) Schematic diagram of an *in vitro* BBB model based on Transwell system.

(B) Quantification of bacterial burden in the macrophages penetrating into the lower

chamber of Transwell system under different concentrations of ox-LDL.

(C-D) (C) Representative image of macrophages that penetrated into the lower

chamber. Scale bar, 100 μm. (D) Quantification of the penetrating cells in each field

of view.

(E) Experimental diagram of adoptive macrophage transfer.

(F-I) After the adoptive macrophage transfer, recipient mice were intraperitoneally

infected with 2×10^8 CFU *L. monocytogenes*. (F) Survival of recipient mice (n=10 for

each group). (G-I) Bacterial burdens in brains (G), and cytokine levels in plasma (H)

and brain homogenate (I) of recipient mice on day 1.25 (n=5 for each group).

(J-N) (J) Schematic diagram of adoptive macrophage transfer and infection

experiments among PCSK9$^{+/+}$ (WT) and PCSK9$^{-/-}$ (KO) mice. Peritoneal

macrophages from PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice were injected into PCSK9$^{+/+}$ and

PCSK9$^{-/-}$ mice, respectively, followed by *L. monocytogenes* infection.

(K, M) Bacterial burdens in the brains of recipient mice after intraperitoneal injection

(K) or oral administration (M) of *L. monocytogenes* (n=5 for each group).

(L, N) Survival of recipient mice after intraperitoneal injection (L) or oral

administration (N) of *L. monocytogenes* (n=10 for each group).

Experiments were performed in duplicate (E-N) or triplicate (A-D). Data are

presented as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ox-LDL, oxidized

low density lipoprotein; FOV, field of view; NS, normal saline; PMφ, purified

peritoneal macrophages; FMφ, CD36$^+$ foamy macrophages; i.p., intraperitoneally; ns,

nonsignificant.

Figure. 3 Two-dimensional live-cell flow chamber assay of the migration

characteristics of infected foamy macrophages.

(A) Schematic diagram of live-cell flow chamber experiment. Primary peritoneal

macrophages were pretreated with ox-LDL for 24 hours and then infected with *L.*

monocytogenes (MOI=10) for 6 hours.

(B-C) (B) Optical images of ox-LDL pretreated macrophages adhering to the surface

of monolayer endothelial cells under shear flow. Scale bar, 200 μm. (C)

Quantification of adherent cells in each field of view.

(D) Resistance to shear flow after adhesion of ox-LDL pretreated macrophages to

endothelial cells. Upon 10-min adherence of equal macrophages to the monolayer

endothelial cells at 1 dyn/cm^2, the shear stress was stepwise increased to 2, 4, 8, 16,

and 32 dyn/cm^2 for 30 s sequentially to detach those adhered cells.

(E-H) Crawling dynamics of ox-LDL pretreated macrophages under shear flow. (E)

Typical trajectories of ox-LDL pretreated macrophages crawling on monolayer

endothelial cells. For each trajectory, the arrested sites of macrophages were overlaid

at the origin of the diagram (x/y=0/0) and the corresponding end points were indicated

with dots. Arrows indicate the flow direction. (F-H) Quantification of crawling

displacement (F), velocity (G) and directness (H).

(I) Transmigration duration of ox-LDL pretreated macrophages after adhesion to the

surface of monolayer endothelial cells.

(J-K) Quantification of the number (J) and mean length (K) of protrusions in ox-LDL

pretreated macrophages.

(L) Quantification of cell area in ox-LDL pretreated macrophages.

(M) qRT-PCR analysis of relative expression of genes associated with intercellular

adhesion molecules.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ox-LDL, oxidized low density lipoprotein.

Figure 4. β-hydroxybutyrate aggravates bacterial neuroinvasion by promoting

the formation of CD36$^+$ foamy macrophages via HCAR2 receptor.

(A-C) Lipid metabolomic profiling analyses of serum from the mice with or without

neurolisteriosis. The mice of neurolisteriosis group were infected with 2×10^8 CFU *L.*

monocytogenes for 24 hours (n=6 for each group). (A) Principal coordinates analysis

(PCA) analysis of differences between groups. (B) Expression of fatty acids (FAs) in

the serum was shown in the volcano plot. Differentially expressed FAs were selected

with |log2 fold change|>1 and adjusted P < 0.05. The dark blue dots represent down-

regulated FAs and the orange dots represent up-regulated FAs. (C) Classification of

the up-regulated FAs (SCFA, short chain fatty acids; MCFA, medium chain fatty acids;

LCFA, long chain fatty acids; VLCFA, very long chain fatty acids) shown in the pie

chart.

(D) Biochemical analysis of serum BHB levels in mice during the neurolisteriosis.

(E-Q) Primary peritoneal macrophages were extracted from 8-week-old HCAR2$^{+/+}$

and HCAR2$^{-/-}$ mice in preparation for further functional analysis. HCAR2$^{+/+}$ and

HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) for 24 hours and then

infected with *L. monocytogenes* (MOI=10) for 6 hours.

(E-F) (E) Flow cytometry analysis of BODYPY FL 493/503 in BHB-treated

macrophages (0, 5, 10 mM). (F) The histogram represents normalized MFI.

(G) Flow cytometry analysis of CD36$^+$ foamy macrophages in BHB-treated

macrophages (0, 5, 10 mM).

(H) Quantification of bacterial burden in the macrophages penetrating into the lower

chamber of Transwell system. The macrophages stimulated with ox-LDL (80 ng/ml)

were used as positive control.

(I-J) (I) Optical images of adherent macrophages on the surface of monolayer

endothelial cells. Scale bar, 200 μm. (J) Quantification of adherent cells in each field

of view.

(K) Resistance to shear flow after adhesion of macrophages to endothelial cells.

(L-P) Crawling and migrating dynamics of infected macrophages under shear flow. (L)

Typical trajectories of infected macrophages crawling on monolayer endothelial cells.

Quantification of crawling displacement (M), velocity (N), directness (O) and

transmigration duration (P).

(Q) mRNA expression of intercellular α-integrin in infected macrophages.

Experiments were performed in duplicate (D) or triplicate (E-Q). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-

hydroxybutyrate; MFI, mean value of fluorescence intensity.

Figure 5. β-hydroxybutyrate promotes the survival of macrophage via HCAR2-mediated calcium influx.

(A-L) Primary peritoneal macrophages were extracted from 8-week-old HCAR2$^{+/+}$ and HCAR2$^{-/-}$ mice, cells were then infected with *L. monocytogenes* (MOI=10) for 6 hours.

(A-C) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) before infection. (A) Quantification of the percentage of released lactate dehydrogenase (LDH) in cell culture medium. (B) Immunofluorescence of Propylene glycol (PI)-stained macrophages. DAPI (blue) labels the nucleus and PI (red) labels the cell membrane. (C) Quantification of the percentage of PI$^+$ cells after infection.

(D) Flow cytometry analysis of dynamics in intracellular calcium concentration. Infected macrophages were preloaded with Fluo-4 AM. BHB (10 mM) was added to the buffer after a span of 30 s.

(E-G) Macrophages were pretreated with BHB (10 mM) or BAPTA (10 mM) for 24 hours before infection. (E) Quantification of the percentage of released lactate dehydrogenase (LDH) in cell culture medium. (F) Quantification of the percentage of PI$^+$ cells after infection. (G) Quantification of bacterial burden in the macrophages penetrating into the lower chamber of Transwell system.

(H-I) (H) Infected macrophages were treated with BHB (10 mM). Time-lapse images of CHMP4B-EGFP in macrophages were collected at 0, 5, 10, 15, 20, 30 min post BHB treatment with confocal microscopy. (I) Quantification of CHMP4B-EGFP puncta.

(J-L) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) for

24 hours before infection. (J) Flow cytometry analysis of LAMP1 expression on the

cell surface. (K) The histogram representing normalized MFI. (L) Quantification of

the lysosomal enzyme (acid phosphatase [ACP], β-galactosidase [β-gal] and β-

hexosaminidase [β-hex]) activity released into the culture medium.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate.

Figure 6. Ketogenesis exacerbates neurolisteriosis through β-hydroxybutyrate-

HCAR2 signaling.

(A-D) (A) Schematic diagram of *in vivo* KE supplementation. C57BL/6 mice were

supplemented with KE (10, 20 mg/ml) in drinking water for 2 weeks and then

challenged with 2×10^8 CFU *L. monocytogenes*. Serum BHB levels (B) and bacterial

burdens in the parenchymal organs (D) were detected (n=5 for each group). (C)

Survival of mice exposed *L. monocytogenes* (n=10 for each group).

(E-M) (E) Schematic diagram of *in vivo* ketogenic diet. HCAR2$^{+/+}$ and HCAR2$^{-/-}$

mice were treated with ketogenic or control diets, respectively, for 4 weeks, followed

by intraperitoneal infection with 2×10^8 CFU *L. monocytogenes*. (F) Survival of mice

exposed to *L. monocytogenes* (n=10 for each group). (G) Flow cytometry analysis of

CD36$^+$ foamy macrophages before and after neurolisteriosis. (H-M) (H-I) Bacterial

burdens in peripheral blood macrophages (H) and parenchymal organs (I) of mice on

day 29. (J-K) Cytokine levels in the plasma (J) and brain homogenate (K). (L-M)

Representative Hematein & Eosin staining (L) and pathology scores (M) of cerebral

cortex tissue. Scale bar, 10 μm. (n=5 for each group).

Experiments were performed in duplicate. Data are presented as mean ± SEM. $*p <$

0.05, $**p < 0.01$. Abbreviations: ns, nonsignificant; KE, Ketone ester; CD, control

diet; KD, ketogenic diet; FMφ, foamy macrophages; i.p., intraperitoneally.

Figure 7. Levels of β-hydroxybutyrate are decreased in bacterial meningitis in

humans, which correlated with the severity of disease.

(A) Schematic diagram of HC (n=60) and BM (n=62) samples. The BM patients were

divided into three groups according to serum BHB levels.

(B) Serum BHB level in HC and BM patients.

(C-I) Clinical scores (C), cell counts (D) and protein levels (E) in cerebrospinal fluid,

white blood cells (WBC, F), procalcitonin (PCT, G), erythrocyte sedimentation rate

(ESR, H) and C-reactive protein (CRP, I) in BM patients with different BHB levels.

(J-K) Dynamics of $CD36^+$ foamy macrophages (J) and serum BHB (K) in 7 BM

donors after admission. The time after admission was divided into symptomatic and

convalescent periods according to clinical and laboratory indexes of infection. The

scatter plot on the right panel represents the statistics of AUC.

(L) Relative expression of genes involved in lipid metabolism in PBMCs from HC

and BM patients was detected by qRT-PCR.

Data are presented as mean ± SEM. $*p < 0.05$, $**p < 0.01$. Abbreviations: ns,

nonsignificant; HC, healthy control; BM, bacterial meningitis; AUC, area under the

curve; FMφ, foamy macrophages; BHB, β -hydroxybutyrate .

Figure S1. Additional results of identification of CD36⁺foamy macrophage

accumulation during the neurolisteriosis, related to Figure 1.

(A-B) C57BL/6 mice were intraperitoneally infected with 2×10^8 CFU *L.*

monocytogenes. (A) Bacterial burdens in brains. (B) Survival of the mice (n=10 for

each group). The yellow shading represents the range of confidence interval.

(C) Cell clusters identified in 10× scRNA-seq dataset. The top 10 enriched genes for

each cluster are highlighted. Normalized z-score is used to illustrate gene expression.

(D) Proportion of each cluster according to experimental conditions in Figure 1D.

(E) Gating strategy for analysis of CD36⁺ foamy macrophages in PBMCs.

(F) CD36⁺ and CD36⁻ macrophages were sorted from peripheral blood by flow

cytometry and then stained with BODIPY FL 493/503 and oil Red O. Scale bar, 10

μm.

(G) Normalized intake of Dil-oxLDL by CD36⁺ and CD36⁻ macrophages.

Experiments were performed in duplicate (A-B) or triplicate (F-G). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: FMφ, foamy macrophages.

Figure S2. Additional results of adoptive transfer of foamy macrophages *in vitro*

and *in vivo*, related to Figure 2.

(A) Conversion of mice peritoneal macrophages into CD36⁺ foamy macrophages by

treating them with different concentrations of ox-LDL for 24 hours.

(B) Intracellular bacterial burden was quantified in macrophages challenged with *L.*

monocytogenes (MOI=10) in the upper chamber of Transwell system under different

concentrations of ox-LDL.

(C-G) The adoptive macrophage transfer experiment was shown in Figure 2E-2I (n=5

for each group). Recipient mice were intraperitoneally infected with 2×10^8 CFU *L.*

monocytogenes. (C-D) Body weight changes (C) and clinical scores (D) of recipient

mice during neurolisteriosis. (E) Bacterial burdens in other tissues (lung, spleen, liver)

of recipient mice on day 1.25. (F-G) Representative Hematein & Eosin staining (F)

and the pathology scoring (G) of cerebral cortex tissue. Scale bar, 20 μm.

(H-I) After adoptive macrophage transfer, recipient mice were intraperitoneally

infected with 5×10^7 CFU *L. monocytogenes*. Bacterial burdens in brains (H) and other

tissues (I) of recipient mice on day 1.25 were detected.

Experiments were performed in duplicate (C-I) or triplicate (A-B). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; NS, normal

saline; PMφ, purified peritoneal macrophages; FMφ, CD36$^+$ foamy macrophages; i.p.,

intraperitoneally.

Figure S3. Accumulation of CD36$^+$ foamy macrophages exacerbate intracellular

bacteria neuroinvasion.

(A-F) Mice were intravenously infected with *S. Typhimurium* (A-B), MRSA (C-D) or

S. pneumoniae (E-F), respectively (n=5 for each group). (A, C, E) Flow cytometry

analysis of CD36$^+$ foamy macrophages during neuroinvasion. (B, D, F) Correlation

between CD36$^+$ foamy macrophages and bacterial burden in the brain during bacteria

neuroinvasion. R, correlation coefficient.

(G-K) After the adoptive macrophage transfer, recipient mice were intraperitoneally

infected with 2×10^8 CFU MRSA. (G) Survival of recipient mice (n=10 for each

group). (H-I) Bacterial burdens in the brains (H) and other tissues (lung, spleen, liver).

(J-K) Cytokine levels in plasma (J) and brain homogenate (K) of recipient mice on

day 1.5 (n=5 for each group).

(L-O) Mice were intraperitoneally infected with 1×10^8 CFU *S. pneumoniae* (n=5 for

each group). (L) Survival rate of mice (n=10). (M) Bacterial burdens in tissues. (N-O)

Cytokine levels in plasma (N) and brain homogenate (O) of recipient mice on day 2

(n=5 for each group).

Experiments were performed in duplicate. Data are presented as mean ± SEM. *$p <$

0.05, **$p < 0.01$. Abbreviations: ns, nonsignificant.

Figure S4. Additional results of bidirectional transfer of macrophages from

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice, related to Figure 2.

(A-B) (A) Flow cytometry analysis of lipid droplets in peritoneal macrophages of

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice. (B) The histogram represents normalized MFI.

(C) Peritoneal macrophages of PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice were infected with *L.*

monocytogenes (MOI=10) for 2 hours and then incubated in a fresh solution

containing 5 μg/mL lysostaphin for 2 or 12 hours. Intracellular bacterial burden of

infected macrophages was quantified

(D) Flow cytometry analysis of CD36$^+$ foamy macrophages during neurolisteriosis in

PCSK9$^{+/+}$ and PCSK9$^{-/-}$ mice, respectively (n=5 for each group).

(E-G) Recipient mice in Figure 2J were intraperitoneally infected with 2×10^8 CFU *L.*

monocytogenes. (E) Bacterial burdens in other tissues (lung, spleen, liver). (F-G)

Cytokine levels in plasma (F) and brain homogenate (G) (n=5 for each group).

(H-J) Recipient mice in Figure 2J were orally infected with 2×10^8 CFU *L.*

monocytogenes. (H) Bacterial burdens in other tissues (lung, spleen, liver). (I-J)

Cytokine levels in plasma (I) and brain homogenate (J) (n=5 for each group).

Experiments were performed in duplicate (D-J) or triplicate (A-C). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; i.p.,

intraperitoneally.

Figure S5. Supplemental data for the live-cell flow chamber assays, related to

Figure 3.

(A-D) Optical images of *L. monocytogenes* infected macrophages (with or without ox-

LDL stimulation) on the surface of endothelial monolayers with different densities (A)

or endothelial monolayers derived from different tissues (C). Scale bar, 200 μm. (B, D)

Quantification of adherent cells in each field of view.

(E) Schematic trajectory used for defining macrophage crawling parameters.

(F) Crawling duration of *L. monocytogene*-infected macrophages after adhesion to

endothelial monolayer surface.

(G) Relative fluorescence intensity of a macrophage on endothelial surface over time.

It is a random example to illustrate the definition of transmigration duration. A 70%

decrease in fluorescence intensity after a macrophage crosses a monolayer is

considered successful transmigration.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: CVEC, cerebrovascular endothelial cell; PVEC,

pulmonary vascular endothelial cell; LSEC, liver sinusoidal endothelial cell; TEM,

transendothelial migration; ns, nonsignificant.

Figure S6. Additional results from the influence of β-hydroxybutyrate on

macrophages during neuroinvasion, related to Figure 4.

(A) KEGG enrichment analysis of lipid metabolites.

(B) Mass spectrometry analysis of serum BHB levels.

(C-D) Relative expression of the gene transcripts regulating ketogenesis in liver

(HMGCS2, CPT1A) detected by qRT-PCR.

(E-F) Body weight changes (E) and food uptake (F) were monitored every 6 hours

during neuroinvasion.

(G-H) Biochemical analysis of serum BHB levels in mice during neuroinvasion by *S.*

typhimurium (G) and MRSA (H).

(I-J) HCAR2$^{+/+}$ and HCAR2$^{-/-}$ macrophages were pretreated with BHB (10 mM) and

cAMP (1 mM) for 24 hours and then infected with *L. monocytogenes* (MOI=10) for 6

hours. Quantification of the normalized MFI of BODIPY FL 493/503 (I) and the

bacterial burden in the macrophages penetrating into the lower chamber of Transwell

system (J).

Experiments were performed in duplicate (C-H) or triplicate (I-J). Data are presented

as mean ± SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; MFI, mean

fluorescence intensity.

Figure S7. Additional results on the role of β-hydroxybutyrate in regulating

macrophage survival, related to Figure 5.

(A-F) Primary peritoneal macrophages were extracted and then infected with *L.*

monocytogenes (MOI=10) for 6 hours.

(A) Macrophages were pretreated with BHB (10 mM) or BAPTA (10 mM) for 24

hours before infection. Intracellular bacterial burden of macrophages in the upper

chamber of Transwell system was quantified.

(B-C) (B) Infected macrophages were treated with BHB (10 mM). Time-lapse images

of CHMP4B-EGFP in macrophages were collected at 0, 5, 10, 15, 20, 30 min post

BHB treatment with confocal microscopy. (C) Quantification of CHMP4B-EGFP

puncta.

(D-F) Peritoneal macrophages were pretreated with BHB (10 mM) or BAPTA (10

mM) for 24 hours before infection. (D) Flow cytometry analysis of LAMP1

expression on the cell surface. (E) The histogram representing normalized MFI. (F)

Quantification of the lysosomal enzyme activity released into the culture medium.

Experiments were performed in triplicate. Data are presented as mean ± SEM. *p <

0.05, **p < 0.01. Abbreviations: ns, nonsignificant; BHB, β-hydroxybutyrate; MFI,

mean fluorescence intensity.

Figure S8. Additional results on the role of ketogenesis in bacterial neuroinvasion, related to Figure 6.

(A-B) The KE supplemental experiment was shown in Figure 6A-6D. (A) Flow cytometry analysis of $CD36^+$ foamy macrophages in mice supplemented with KE. (B) Clinical scores of mice supplemented with KE.

(C-D) The ketogenic diet experiment was shown in Figure 6E-6M. Body weight changes (C) and clinical scores (D) of mice during *L. monocytogenes* infection.

(E-H) $HCAR2^{+/+}$ and $HCAR2^{-/-}$ mice were treated with ketogenic and control diets, respectively, for 4 weeks, followed by oral infection with 2×10^8 CFU *L. monocytogenes*. (E) Survival of mice exposed to *L. monocytogenes* (n=10 for each group). (F) Bacterial burdens in organs (brain, lung, spleen, liver). Cytokine levels in plasma (G) and brain homogenate (H) (n=5 for each group).

(I-K) C57BL/6 mice were treated with ketogenic and control diets, respectively, for 4 weeks, followed by intravenous infection with 2×10^8 CFU MRSA. (I) Survival of mice exposed to MRSA (n=15 for each group). (J) Bacterial burdens in organs (brain, lung, spleen, liver). (K) Clinical scores of mice during neuroinvasion (n=5 for each group).

Experiments were performed in duplicate. Data are presented as mean $\pm$ SEM. *p < 0.05, **p < 0.01. Abbreviations: ns, nonsignificant; i.p., intraperitoneally.

Figure S9. Additional results of high levels of BHB and human tuberculous meningitis, related to Figure 7.

(A-D) Glucose (A) and insulin (C) levels in serum from HC and BM patients.

Correlation between serum glucose level and BHB (B), as well as serum insulin levels and BHB (D) in BM patients. R, correlation coefficient

(E-F) Serum triglyceride (TC, E), high density lipoprotein (HDL-C, F) and low density lipoprotein (LDL-C, G) from HC and BM patients.

Data are presented as mean ± SEM. $*p < 0.05$, $**p < 0.01$. Abbreviations: ns, nonsignificant, HC, health control, BM, bacterial meningitis.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Clinical samples

Bacterial meningitis (BM) disease subjects (n = 62, 32 males and 30 females, aged 41.62 ±15.68 years) and healthy donors (n = 60, 30 males and 30 females, aged 38.32±9.17 years) were recruited from Chongqing Public Health Medical Center. Isolation of PBMCs and single-cell suspensions was performed using PBMC Isolation Kit (P8680, Solarbio) by density gradient centrifugation. Plasma was collected from each patient and donor and then used for quantification of BHB. This study was carried out in strict accordance with the Guidelines for the Care and Use of Animals of Chongqing University, and was compliant with the "Guidance of the Ministry of Science and Technology (MOST) for the Review and Approval of Human Genetic

Resources". Healthy individuals and patients providing blood samples were given

informed consent. The ethics committee approved this consent procedure. The clinical

symptoms of the BM patients included burnout, headache, irritability, vomiting, fever,

neck stiffness, convulsion, and consciousness disorder. Each symptom is worth 1

point.

Animal

All animal experiments were performed following institutional guidelines at

Chongqing University. In all experiments, control and experimental mice were 6-8-

week-old littermates of male C57BL/6 mice. Littermates of the same sex were housed

together regardless of genotype. All mice were maintained under a 12-hour light-dark

cycle with ad libitum access to regular food and water.

Bacterial strains

L. monocytogenes (19115) and *S. pneumoniae* 49619 were grown in Brain Heart

Infusion Broth. Methicillin-resistant *Staphylococcus aureus* USA300 and *S.*

Typhimurium were grown in Luria-Bertani Broth.

Data and code availability

Single-cell RNA-seq data have been deposited at GEO and are publicly available as

of the date of publication.

METHOD DETAILS

Single cell RNA-sequencing analysis

Isolated PBMCs from the mice were transported in in sterile culture dish with 10 ml

Dulbecco's Phosphate-Buffered Saline (DPBS, Thermo Fisher Scientific) on ice to

remove the residual tissue storage solution, then minced on ice. We used dissociation

enzyme 0.25% Trypsin (Thermo Fisher Scientific) and 10 ug/mL DNase I (Sigma)

dissolved in PBS with 5% Fetal Bovine Serum (FBS, GIBCO) to digest the tissues.

PBMCs were dissociated at 37 ℃ under a shaking speed of 50 rpm for about 40 min.

We repeatedly collected the dissociated cells at interval of 20 min to increase cell

yield and viability. Cell suspensions were filtered using a 40-μm nylon cell strainer

and red blood cells were removed by red blood cell lysis solution (Thermo Fisher

Scientific). Dissociated cells were washed with DPBS containing 2% FBS.

10× beads were then subject to second-strand cDNA synthesis, adaptor ligation, and

universal amplification. Sequencing libraries were prepared using randomly

interrupted whole-transcriptome amplification products to enrich the 3' end of the

transcripts linked with the cell barcode and UMI. All the remaining procedures

including the library construction were performed according to the standard

manufacturer's protocol (Chromium Single Cell 3' v3). Sequencing libraries were

quantified using a High Sensitivity DNA Chip (Agilent) on a Bioanalyzer 2100 and

the Qubit High Sensitivity DNA Assay (Thermo Fisher Scientific). The libraries were

sequenced on NovaSeq6000 (Illumina) using 2×150 chemistry. The sequencing and

bioinformatic analysis were performed on platform of Majorbio Co., Ltd (Shanghai,

China).

Reads were processed using the Cell Ranger (v7.1.0) pipeline with default and

recommended parameters. FASTQs generated from Illumina sequencing output were

aligned to the mouse genome, version GRCm38, using the STAR algorithm. Gene-

barcode matrix containing the barcoded cells and gene expression counts were

generated. This output was then imported into the Seurat (v3.2.0) R toolkit for quality

control and downstream analysis of our single cell RNAseq data.

Cells were clustered using graph-based clustering of the PCA reduced data with the

Louvain Method after computing a shared nearest neighbor graph. For sub-clustering,

we applied the same procedure of scaled, dimensionality reduction, and clustering to

the specific set of data. For each cluster, we used the Wilcoxon Rank-Sum Test to find

significant deferentially expressed genes comparing to the remaining clusters.

To identify differential expression genes (DEGs) between two different samples or

clusters, was performed using the function FindMarkers in Seurat, using a likelihood

ratio test. Essentially, DEGs with $|log2FC| > 0.25$ and Q value ≤ 0.05 were

significantly different expressed genes. In addition, functional-enrichment analysis

GO were performed to identify which DEGs were significantly enriched in GO terms

and metabolic pathways at Bonferroni-corrected P-value ≤ 0.05 compared with the

whole-transcriptome background. GO functional enrichment analysis were carried out

by Goatools (https://github.com/tanghaibao/Goatools).

1055

Mouse model of acute bacteria brain neuroinvasion

Mice were challenged with pathogenic bacteria (2×10^8 CFU *L. monocytogenes* or

MRSA, 1×10^8 CFU *S. Typhimurium* or *S. pneumoniae*) and weighed and scored every

3 hours. The progression of acute bacterial meningitis was determined by clinical

scores as previously described with a few adaptations.[65] Clinical scoring was

performed based on the Table S2, including activity (0–2), time to return to upright

position (0–2), eyes (0–2) and neurological examination (0–2). Mice were deeply

anaesthetized by an intraperitoneal injection of tribromoethanol solution (Avertin, 500

mg/kg) and transcranial perfused with saline solution for brain tissue

pathomorphology. Blood was sampled immediately before perfusion. Tissue samples,

including brain, lung, liver, and spleen, were dissected and weighed.

1067

Analysis Bacterial load in cell or tissue

Tissue samples of mice were transferred into 2 ml tubes containing 4 mm stainless

steel beads (Qiagen) and 1 ml of ice-cold sterile saline. Homogenization was carried

out in TissueLyser II (Qiagen) for 5 min at 70 Hz. Serial dilutions of the homogenates

were plated on the plates incubated at 37℃ and CFU were counted.

The infected macrophages were incubated in a fresh solution containing 5 μg/mL

lysostaphin for the indicated periods and then lysed by 0.2% Triton to release

intracellular bacteria. The lysates were diluted to appropriate dilutions with sterile

PBS for CFU count on agar plates.

1077

Brain histopathology and Immunostaining

Brain samples were collected from infected mice 24 hours after injection. Mice were

deeply anaesthetized with tribromoethanol solution (Avertin, 500 mg/kg,

intravenously) and transcranial perfused with 30 ml saline solution followed by 10 ml

of 4% Paraformaldehyde (PFA) in saline. After dissection, brain samples were

maintained in 4% PFA at 4 ℃. Fixed brains were embedded in paraffin, sectioned and

half of the slides were stained using haematoxylin and eosin. Each field was assigned

a score from 0 to 3 based on neuronal morphology: grade 0 (not altered); grade 1 (no

vacuolation with small numbers of pyknotic cells); grade 2 (moderate vacuolation and

pyknosis); and grade 3 (extensive vacuolation, pyknosis and tissue loss or liquefactive

necrosis).

Fresh frozen brain tissue sections of 5-10 mm thickness were mounted on slides, air

dried for 10 minutes at room temperature and fixed in ice cold 10% neutral-buffered

formalin for 5-10 min. Slices were washed in PBS, then incubated for 2 hours in 3%

bovine serum albumin (BSA) and then 0.1% Triton X-100 in PBS. The slides were

labelled with the appropriate primary antibodies of CD11b and CD36 overnight at 4 ℃

in mild blocking conditions. Next day, the slices were washed in PBS, and then

incubated with secondary antibodies and DAPI. Tissues were washed in PBS and then

mounted on glass slides under coverslips in mounting medium. The slides were left in

darkness overnight before observation under a Zeiss LSM900 or LM710 microscope.

1098

1099 **Cell sorting and flow cytometry.**

1100 Isolated mouse PBMCs were cleaned twice with PBS. After calculating and

determining the appropriate concentration, appropriate amount of fluorescent-coupled

primary antibody (F4/80-APC, CD11b-FITC, as the marker of macrophage, CD36-

FITC) was added respectively. Incubate at 4 ℃ for 30 min away from light. The

living macrophages were collected according to the expression of CD36 and then

culture in vitro overnight. The sorting cells were detected the lipid content using

BODIPY FL and oil Red O. To evaluate the ability of lipid uptake, macrophages were

incubated with 40 µg/ml DiI-ox-LDL particles for 6h and then detect the fluorescence

intensity by flow cytometry.

To determine the proportion of foamy macrophages population, cells were stained

with fluorescent-coupled primary antibody (F4/80-APC, CD11b-PE, CD36-FITC) at

4 ℃ for 30 min away from light. To detect the expression of LAMP1 on the cell

surface, infected macrophages were cleaned twice with PBS and then stained with

LAMP1-FITC at 4 ℃ for 30 min away from light. The cells were washed three times

and then re-suspended in 200µl of the staining buffer for final FACS analysis. Data

was analyzed with FlowJo software.

Isolation and culture of primary cells

To generate peritoneal macrophages, 8-week-old mice were injected intraperitoneally

with 3% thioglycolate broth. After 72 hours, peritoneal cells were harvested, and

macrophages were enriched by quick adherence.

Primary endothelial cells were isolated from 8 to 12-week-old male C57BL/6 mice.

The cerebral cortex in the brains were separated and rolled on the filter paper to

remove the meninges and choroid plexuses. A mechanical reduction was performed

by mincing the brain pieces using surgical blades. The homogenate was centrifuged

(800 g, 8 min, 4℃) and the precipitate was digested in DMEM containing 1 mg/ml

type II collagenase and 58.5 U/ml type I DNase for 75 min on a benchtop orbital

shaker at 37 ℃. The supernatant was finally collected and centrifuged at 500 g for 8

min. The isolated cerebrovascular endothelial cells were purified by the anchoring

technique. The liver and lung from an anaesthetized mouse were perfused via the

portal vein sequentially with balanced salt solution containing 5 mM glucose, 0.01%

sodium heparin, and 5 mM EGTA and high glucose DMEM medium containing 4

mM $CaCl_2$, 2% FBS and 0.05% collagenase IV (Sigma-Aldrich, St. Louis, MO) at a

flow rate of 5 ml/min for 5 min, respectively. The following steps are similar in order

to separate the liver sinusoids endothelial cells and pulmonary vascular endothelial

cells.

Isolation of primary embryonic neurons and single-cell suspensions was performed

using Pierce Primary Neuron Isolation Kit (88280, Thermo Scientific) by density

gradient centrifugation.

1139

***In vitro* blood-brain barrier**

The *in vitro* BBB model was constructed with primary endothelial cells using a

Transwell cell culture system. Briefly, CVECs were seeded onto the upper chamber of

the Transwell pre-coated with gelatine, and cultured with the medium containing 10%

FBS. The integrity of the monolayer cell was evaluated by measuring the TEER

values using a Millicell-ERS voltohmmeter (Millipore). The cell monolayers with

TEER values higher than 300 $\Omega.cm^2$ were used as the BBB model for the

transmigration studies. The neurons cells were seeded onto the lower chamber of the

Transwell for bacterial load.

Macrophages were exposed to *L. monocytogenes* (MOI=10) for 6 hours. After the

BBB model has been constructed, the infected macrophages were digested for cell

counting and added into the upper chamber. The conventional incubation time was 12

hours and then the neurons cells in the bottom chamber were lysed for penetrating

bacterial quantification.

1154

Adoptive macrophages transfer in mouse bacteria brain neuroinvasion

Peritoneal macrophages from donor mice were collected via intraperitoneal injection

of thioglycolate. Following further cultivation, purified macrophages were obtained.

To collect CD36$^+$ foamy macrophages, purified macrophages were stimulated with 80

µg/ml ox-LDL over 24 hours. Recipient mice were pretreated with 3 days of CTX and

then intraperitoneally given 0.2 ml normal saline or cell suspension (1×10^6 per mouse)

on Day 0. After adoptive macrophages transfer, recipient mice were intraperitoneally

injected with 2×10^8 CFU of *L. monocytogenes* in 12 hours and euthanized after 18

hours.

Purified peritoneal macrophages were harvested from PCSK9$^{+/+}$ and PCSK9$^{-/-}$ donor

mice respectively, and then transferred to recipient mice as depicted in Figure 2J.

Recipient mice were infected with 2×10^8 CFU *L. monocytogenes* 12 hours after

macrophages transfer and euthanized after 24 hours.

1168

Live-cell flow chamber assay

A flow cell system was used to test macrophages adhesion, shear resistance, crawling,

and transmigration, on endothelial monolayer, which was performed as previously

described.[42] Shear flow was applied by a syringe pump (PHD22/2000, Harvard

Apparatus) into a microfluidic chip with a 100 μm × 1 mm× 15 mm main flowing

zone. Different shear stresses were acquired by setting corresponding volumetric flow

rates. Isolated macrophages were stimulated under different concentration of ox-LDL

for 24 hours and then infected with *L. monocytogenes* (MOI=10) for 6 hours.

Macrophages were re-suspended and stained with Hoechst. The cells were then

perfused over pre-formed endothelial monolayer for 8 min at a shear stress of 0.1

dyne/cm^2. Then, the adhesion and crawling dynamics of macrophages on endothelial

monolayer was recorded at 2 dyn/cm^2 for 10 min at 20× objective by a CCD camera.

Images were taken from the video every second and analyzed the crawling parameters

and transmigrating duration with Image J software. Here a macrophage crossing the

monolayer with a 70% fluorescence intensity reduction was deemed transmigrated.

For testing shear-resistant macrophage adhesion, primed macrophages were re-

suspended and counted. The same number of cells (2×10^4) were completely perfused

over pre-formed endothelial monolayer within 10 min. The shear stress was increased

stepwise from 1 to 2, 4, 8, 16, 32 dyn/cm^2 for 30s each. Remaining adherent

macrophages on endothelial monolayer were recorded at 20× objective by a CCD

camera and counted using Image J software. The normalized adhesion fraction was

calculated by dividing the number of adherent macrophages at the endpoint of each

shear phase by the number at the endpoint of 1 dyn/cm^2. For defining the macrophage

transmigration dynamics, the time-lapse images were analyzed offline frame-by-frame

every second.

Quantification of *L. monocytogenes* protrusion formation

Isolated peritoneal macrophages were stimulated under different concentration of ox-

LDL for 24 hours and then infected with CFSE-labeled *L. monocytogenes* (MOI=10)

for 6 hours. At 4 hours post-infection, cells were fixed, permeabilized, and labeled

with mouse anti-ezrin and phalloidin-Alexa 647. Samples were analyzed using

confocal microscopy. Protrusions were identified as bacteria with ezrin tails that

project from host cells into neighboring. In each experiment, ten cells were scored for

each condition. The lengths of ezrin comet tails were measured using ImageJ software.

Lipid metabolomics

Metabolite extraction and mass spectrometry detection was processed using Ultra-

High Performance Liquid Chromatography-mass spectrometry (UHPLC-MS) of

Biomarker Technologies (Beijing, China). The raw data collected using MassLynx

V4.2 is processed by Progenesis QI software for peak extraction, peak alignment, and

other data processing operations, based on the Progenesis QI software online

METLIN database and Biomark's self-built library for identification. The theoretical

fragment identification and mass deviation are within 100 ppm. After normalizing the

original peak area information with the total peak area, the follow-up analysis was

performed. Principal component analysis (PCA) and Spearman correlation analysis

were used to judge the repeatability of the samples within group and the quantity

control samples. The identified compounds are searched for classification and

pathway information in KEGG and HMDB databases.

A total of 4,625 named metabolites were retained for datasets. The R language

package ropls was used to perform OPLS-DA modeling, and 200 times permutation

tests was performed to verify the reliability of the model. The screening criteria are

FC>1, $p < 0.05$ and VIP>1. The difference metabolites of KEGG pathway enrichment

significance were calculated using hypergeometric distribution test.

1222

Fura-4 Ca^{2+} imaging

Cells were removed from incubator and washed with PBS and then incubated with 5

μM Fluo-4 AM (Beyotime) for 60 min at 37 ℃ with 0.04% Pluronic-127. After

incubation, cells were washed with PBS and incubated with PBS for 30 min at 37 ℃.

Fluorescence were detected with a CytoFLEX Flow Cytometer (Beckman Coulter)

equipped with 488 nm excitation wavelength. As indicated, BHB (10 mM) were

added into the cells pre-loaded with Fluo-4 30 s after the record. Data was analyzed

with Flow Jo to determine the fluorescence intensity.

1231

Diets for *in vivo* assays

When indicated, mice were administered a supplement of 20 mg/ml ketone ester (D-

β-hydroxybutyrate-(R)-1,3 butanediol monoester) in the drinking water for 4 weeks

before the infection. Ketogenic diets (# XTKD01, Xietong Pharma Co.) contained 10%

protein and 90% fat while the control diets contained 23.2% protein, 2.8% fat and 74%

sugar. Mice were consequently placed on either the ketogenic or control diet for an

additional 4 weeks before infection.

1239

Real-time qRT-PCR

Total RNA was isolated using an RNeasy Mini Kit (QIAGEN). The mRNA was

reverse-transcribed and quantified using a HiScript II 1st Strand cDNA Synthesis Kit

(Vazyme). qRT-PCR was performed on an ABI StepOnePlus PCR system (Thermo

Fisher Scientific), and results were normalized to GAPDH mRNA levels. Data were

analyzed using the 2-DDct method.

1246

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis was performed using GraphPad Prism Software. Two-group

comparisons were made using two-tailed unpaired Student's t-test. One-way analysis

of variance (ANOVA) with appropriate multiple comparisons tests was used to

compare three independent groups. For comparisons of multiple factors, two-way

ANOVA with appropriate multiple comparisons tests was used. Exact p values for

specific group comparisons in all the figures. A value of $p < 0.05$ was considered

significant. Specific details of statistical analysis are presented above or in associated

figure legends.

Reference

1. Doran, K.S., Fulde, M., Gratz, N., Kim, B.J., Nau, R., Prasadarao, N., Schubert-Unkmeir, A., Tuomanen, E.I., and Valentin-Weigand, P. (2016). Host-pathogen interactions in bacterial meningitis. Acta Neuropathol *131*, 185-209. 10.1007/s00401-015-1531-z.

2. Hasbun, R. (2022). Progress and Challenges in Bacterial Meningitis: A Review. JAMA *328*, 2147-2154. 10.1001/jama.2022.20521.

3. Rua, R., and McGavern, D.B. (2018). Advances in Meningeal Immunity. Trends Mol Med *24*, 542-559. 10.1016/j.molmed.2018.04.003.

4. Merlini, A., Haberl, M., Strauss, J., Hildebrand, L., Genc, N., Franz, J., Chilov, D., Alitalo, K., Flugel-Koch, C., Stadelmann, C., et al. (2022). Distinct roles of the meningeal layers in CNS autoimmunity. Nat Neurosci *25*, 887-899. 10.1038/s41593-022-01108-3.

5. Byrd, T.F., and Davis, L.E. (2007). Multidrug-resistant tuberculous meningitis. Curr Neurol Neurosci Rep *7*, 470-475. 10.1007/s11910-007-0073-8.

6. Huynh, J., Donovan, J., Phu, N.H., Nghia, H.D.T., Thuong, N.T.T., and Thwaites, G.E. (2022). Tuberculous meningitis: progress and remaining questions. Lancet Neurol *21*, 450-464. 10.1016/S1474-4422(21)00435-X.

7. van de Beek, D., Brouwer, M.C., Koedel, U., and Wall, E.C. (2021). Community-acquired bacterial meningitis. Lancet *398*, 1171-1183.

1277 10.1016/S0140-6736(21)00883-7.

1278 8. Levin, B.R., Baquero, F., Ankomah, P.P., and McCall, I.C. (2017). Phagocytes,

1279 Antibiotics, and Self-Limiting Bacterial Infections. Trends Microbiol *25*, 878-

1280 892. 10.1016/j.tim.2017.07.005.

1281 9. Boada-Romero, E., Martinez, J., Heckmann, B.L., and Green, D.R. (2020).

1282 The clearance of dead cells by efferocytosis. Nat Rev Mol Cell Biol *21*, 398-

1283 414. 10.1038/s41580-020-0232-1.

1284 10. Locati, M., Curtale, G., and Mantovani, A. (2020). Diversity, Mechanisms,

1285 and Significance of Macrophage Plasticity. Annu Rev Pathol *15*, 123-147.

1286 10.1146/annurev-pathmechdis-012418-012718.

1287 11. Zhai, L., Huang, C., Ning, Z., Zhang, Y., Zhuang, M., Yang, W., Wang, X.,

1288 Wang, J., Zhang, L., Xiao, H., et al. (2023). Ruminococcus gnavus plays a

1289 pathogenic role in diarrhea-predominant irritable bowel syndrome by

1290 increasing serotonin biosynthesis. Cell Host Microbe *31*, 33-44 e35.

1291 10.1016/j.chom.2022.11.006.

1292 12. Van der Merwe, S., Chokshi, S., Bernsmeier, C., and Albillos, A. (2021). The

1293 multifactorial mechanisms of bacterial infection in decompensated cirrhosis. J

1294 Hepatol *75 Suppl 1*, S82-S100. 10.1016/j.jhep.2020.11.029.

1295 13. Maudet, C., Kheloufi, M., Levallois, S., Gaillard, J., Huang, L., Gaultier, C.,

1296 Tsai, Y.H., Disson, O., and Lecuit, M. (2022). Bacterial inhibition of Fas-

1297 mediated killing promotes neuroinvasion and persistence. Nature *603*, 900-906.

1298 10.1038/s41586-022-04505-7.

1299   14.   Xu, K., Yin, N., Peng, M., Stamatiades, E.G., Shyu, A., Li, P., Zhang, X., Do,

1300   M.H., Wang, Z., Capistrano, K.J., et al. (2021). Glycolysis fuels

1301   phosphoinositide 3-kinase signaling to bolster T cell immunity. Science *371*,

1302   405-410. 10.1126/science.abb2683.

1303   15.   Escoll, P., Song, O.R., Viana, F., Steiner, B., Lagache, T., Olivo-Marin, J.C.,

1304   Impens, F., Brodin, P., Hilbi, H., and Buchrieser, C. (2017). Legionella

1305   pneumophila Modulates Mitochondrial Dynamics to Trigger Metabolic

1306   Repurposing of Infected Macrophages. Cell Host Microbe *22*, 302-316 e307.

1307   10.1016/j.chom.2017.07.020.

1308   16.   Yang, D., Jacobson, A., Meerschaert, K.A., Sifakis, J.J., Wu, M., Chen, X.,

1309   Yang, T., Zhou, Y., Anekal, P.V., Rucker, R.A., et al. (2022). Nociceptor

1310   neurons direct goblet cells via a CGRP-RAMP1 axis to drive mucus

1311   production and gut barrier protection. Cell *185*, 4190-4205 e4125.

1312   10.1016/j.cell.2022.09.024.

1313   17.   Loft, A., Schmidt, S.F., Caratti, G., Stifel, U., Havelund, J., Sekar, R., Kwon,

1314   Y., Sulaj, A., Chow, K.K., Alfaro, A.J., et al. (2022). A macrophage-hepatocyte

1315   glucocorticoid receptor axis coordinates fasting ketogenesis. Cell Metab *34*,

1316   473-486 e479. 10.1016/j.cmet.2022.01.004.

1317   18.   Puchalska, P., and Crawford, P.A. (2017). Multi-dimensional Roles of Ketone

1318   Bodies in Fuel Metabolism, Signaling, and Therapeutics. Cell Metab *25*, 262-

1319   284. 10.1016/j.cmet.2016.12.022.

19.   Tomlinson, K.L., Chen, Y.T., Junker, A., Urso, A., Wong Fok Lung, T., Ahn,

D., Hofstaedter, C.E., Baskota, S.U., Ernst, R.K., Prince, A., and Riquelme,

S.A. (2023). Ketogenesis promotes tolerance to Pseudomonas aeruginosa

pulmonary infection. Cell Metab *35*, 1767-1781 e1766.

10.1016/j.cmet.2023.09.001.

20.   Goldberg, E.L., Molony, R.D., Kudo, E., Sidorov, S., Kong, Y., Dixit, V.D.,

and Iwasaki, A. (2019). Ketogenic diet activates protective gammadelta T cell

responses against influenza virus infection. Sci Immunol *4*.

10.1126/sciimmunol.aav2026.

21.   Seufert, A.L., Hickman, J.W., Traxler, S.K., Peterson, R.M., Waugh, T.A.,

Lashley, S.J., Shulzhenko, N., Napier, R.J., and Napier, B.A. (2022). Enriched

dietary saturated fatty acids induce trained immunity via ceramide production

that enhances severity of endotoxemia and clearance of infection. Elife *11*.

10.7554/eLife.76744.

22.   Wilhelm, C., Surendar, J., and Karagiannis, F. (2021). Enemy or ally? Fasting

as an essential regulator of immune responses. Trends Immunol *42*, 389-400.

10.1016/j.it.2021.03.007.

23.   Singh, V., Jamwal, S., Jain, R., Verma, P., Gokhale, R., and Rao, K.V. (2012).

Mycobacterium tuberculosis-driven targeted recalibration of macrophage lipid

homeostasis promotes the foamy phenotype. Cell Host Microbe *12*, 669-681.

10.1016/j.chom.2012.09.012.

24.   McFarland, A.P., Yalin, A., Wang, S.Y., Cortez, V.S., Landsberger, T., Sudan,

R., Peng, V., Miller, H.L., Ricci, B., David, E., et al. (2021). Multi-tissue

single-cell analysis deconstructs the complex programs of mouse natural killer

and type 1 innate lymphoid cells in tissues and circulation. Immunity *54*,

1320-1337 e1324. 10.1016/j.immuni.2021.03.024.

25.   Ahmad, F., Rani, A., Alam, A., Zarin, S., Pandey, S., Singh, H., Hasnain, S.E.,

and Ehtesham, N.Z. (2022). Macrophage: A Cell With Many Faces and

Functions in Tuberculosis. Front Immunol *13*, 747799.

10.3389/fimmu.2022.747799.

26.   Hill, D.A., Lim, H.W., Kim, Y.H., Ho, W.Y., Foong, Y.H., Nelson, V.L.,

Nguyen, H.C.B., Chegireddy, K., Kim, J., Habertheuer, A., et al. (2018).

Distinct macrophage populations direct inflammatory versus physiological

changes in adipose tissue. Proc Natl Acad Sci U S A *115*, E5096-E5105.

10.1073/pnas.1802611115.

27.   Jaitin, D.A., Adlung, L., Thaiss, C.A., Weiner, A., Li, B., Descamps, H.,

Lundgren, P., Bleriot, C., Liu, Z., Deczkowska, A., et al. (2019). Lipid-

Associated Macrophages Control Metabolic Homeostasis in a Trem2-

Dependent Manner. Cell *178*, 686-698 e614. 10.1016/j.cell.2019.05.054.

28.   Luis, G., Godfroid, A., Nishiumi, S., Cimino, J., Blacher, S., Maquoi, E., Wery,

C., Collignon, A., Longuespee, R., Montero-Ruiz, L., et al. (2021). Tumor

resistance to ferroptosis driven by Stearoyl-CoA Desaturase-1 (SCD1) in

cancer cells and Fatty Acid Biding Protein-4 (FABP4) in tumor

microenvironment promote tumor recurrence. Redox Biol *43*, 102006.

10.1016/j.redox.2021.102006.

29. Victor, M.B., Leary, N., Luna, X., Meharena, H.S., Scannail, A.N., Bozzelli,

P.L., Samaan, G., Murdock, M.H., von Maydell, D., Effenberger, A.H., et al.

(2022). Lipid accumulation induced by APOE4 impairs microglial

surveillance of neuronal-network activity. Cell Stem Cell *29*, 1197-1212 e1198.

10.1016/j.stem.2022.07.005.

30. Xu, S., Chaudhary, O., Rodriguez-Morales, P., Sun, X., Chen, D., Zappasodi,

R., Xu, Z., Pinto, A.F.M., Williams, A., Schulze, I., et al. (2021). Uptake of

oxidized lipids by the scavenger receptor CD36 promotes lipid peroxidation

and dysfunction in CD8(+) T cells in tumors. Immunity *54*, 1561-1577 e1567.

10.1016/j.immuni.2021.05.003.

31. Guerrini, V., and Gennaro, M.L. (2019). Foam Cells: One Size Doesn't Fit All.

Trends Immunol *40*, 1163-1179. 10.1016/j.it.2019.10.002.

32. Luo, Y., Duan, H., Qian, Y., Feng, L., Wu, Z., Wang, F., Feng, J., Yang, D., Qin,

Z., and Yan, X. (2017). Macrophagic CD146 promotes foam cell formation

and retention during atherosclerosis. Cell Res *27*, 352-372. 10.1038/cr.2017.8.

33. Kim, K., Shim, D., Lee, J.S., Zaitsev, K., Williams, J.W., Kim, K.W., Jang,

M.Y., Seok Jang, H., Yun, T.J., Lee, S.H., et al. (2018). Transcriptome

Analysis Reveals Nonfoamy Rather Than Foamy Plaque Macrophages Are

Proinflammatory in Atherosclerotic Murine Models. Circ Res *123*, 1127-1142.

10.1161/CIRCRESAHA.118.312804.

34. Botham, K.M., and Wheeler-Jones, C.P. (2013). Postprandial lipoproteins and

the molecular regulation of vascular homeostasis. Prog Lipid Res *52*, 446-464.

10.1016/j.plipres.2013.06.001.

35.   Jaggi, U., Matundan, H.H., Yu, J., Hirose, S., Mueller, M., Wormley, F.L., Jr.,
and Ghiasi, H. (2021). Essential role of M1 macrophages in blocking cytokine
storm and pathology associated with murine HSV-1 infection. PLoS Pathog *17*,
e1009999. 10.1371/journal.ppat.1009999.

36.   Pasula, R., Martin, W.J., 2nd, Kesavalu, B.R., Abdalla, M.Y., and Britigan,
B.E. (2017). Passive transfer of interferon-gamma over-expressing
macrophages enhances resistance of SCID mice to Mycobacterium
tuberculosis infection. Cytokine *95*, 70-79. 10.1016/j.cyto.2017.02.009.

37.   Peyrusson, F., Varet, H., Nguyen, T.K., Legendre, R., Sismeiro, O., Coppee,
J.Y., Wolz, C., Tenson, T., and Van Bambeke, F. (2020). Intracellular
Staphylococcus aureus persisters upon antibiotic exposure. Nat Commun *11*,
2200. 10.1038/s41467-020-15966-7.

38.   Ding, Z., Pothineni, N.V.K., Goel, A., Luscher, T.F., and Mehta, J.L. (2020).
PCSK9 and inflammation: role of shear stress, pro-inflammatory cytokines,
and LOX-1. Cardiovasc Res *116*, 908-915. 10.1093/cvr/cvz313.

39.   Da Dalt, L., Castiglioni, L., Baragetti, A., Audano, M., Svecla, M., Bonacina,
F., Pedretti, S., Uboldi, P., Benzoni, P., Giannetti, F., et al. (2021). PCSK9
deficiency rewires heart metabolism and drives heart failure with preserved
ejection fraction. Eur Heart J *42*, 3078-3090. 10.1093/eurheartj/ehab431.

40.   Katsuki, S., P, K.J., Lupieri, A., Nakano, T., Passos, L.S.A., Rogers, M.A.,
Becker-Greene, D., Le, T.D., Decano, J.L., Ho Lee, L., et al. (2022).

Proprotein Convertase Subtilisin/Kexin 9 (PCSK9) Promotes Macrophage

Activation via LDL Receptor-Independent Mechanisms. Circ Res *131*, 873-

889. 10.1161/CIRCRESAHA.121.320056.

41.   Qi, Z., Hu, L., Zhang, J., Yang, W., Liu, X., Jia, D., Yao, Z., Chang, L., Pan, G.,

Zhong, H., et al. (2021). PCSK9 (Proprotein Convertase Subtilisin/Kexin 9)

Enhances Platelet Activation, Thrombosis, and Myocardial Infarct Expansion

by Binding to Platelet CD36. Circulation *143*, 45-61.

10.1161/CIRCULATIONAHA.120.046290.

42.   Yang, H., Li, N., Du, Y., Tong, C., Lu, S., Hu, J., Zhang, Y., and Long, M.

(2017). Neutrophil adhesion and crawling dynamics on liver sinusoidal

endothelial cells under shear flow. Exp Cell Res *351*, 91-99.

10.1016/j.yexcr.2017.01.002.

43.   Neupane, A.S., Willson, M., Chojnacki, A.K., Vargas, E.S.C.F., Morehouse, C.,

Carestia, A., Keller, A.E., Peiseler, M., DiGiandomenico, A., Kelly, M.M., et

al. (2020). Patrolling Alveolar Macrophages Conceal Bacteria from the

Immune System to Maintain Homeostasis. Cell *183*, 110-125 e111.

10.1016/j.cell.2020.08.020.

44.   Hernandez, G.E., and Iruela-Arispe, M.L. (2020). The many flavors of

monocyte/macrophage--endothelial cell interactions. Curr Opin Hematol *27*,

181-189. 10.1097/MOH.0000000000000573.

45.   Pang, X., He, X., Qiu, Z., Zhang, H., Xie, R., Liu, Z., Gu, Y., Zhao, N., Xiang,

Q., and Cui, Y. (2023). Targeting integrin pathways: mechanisms and advances

in therapy. Signal Transduct Target Ther *8*, 1. 10.1038/s41392-022-01259-6.

46.   Deng, L., Kersten, S., and Stienstra, R. (2023). Triacylglycerol uptake and

handling by macrophages: From fatty acids to lipoproteins. Prog Lipid Res *92*,

101250. 10.1016/j.plipres.2023.101250.

47.   Patterson, M.T., Firulyova, M.M., Xu, Y., Hillman, H., Bishop, C., Zhu, A.,

Hickok, G.H., Schrank, P.R., Ronayne, C.E., Caillot, Z., et al. (2023). Trem2

promotes foamy macrophage lipid uptake and survival in atherosclerosis.

Nature Cardiovascular Research *2*, 1015-1031. 10.1038/s44161-023-00354-3.

48.   Andrews, N.W., and Corrotte, M. (2018). Plasma membrane repair. Curr Biol

28, R392-R397. 10.1016/j.cub.2017.12.034.

49.   Jimenez, A.J., Maiuri, P., Lafaurie-Janvore, J., Divoux, S., Piel, M., and Perez,

F. (2014). ESCRT machinery is required for plasma membrane repair. Science

343, 1247136. 10.1126/science.1247136.

50.   Newman, J.C., and Verdin, E. (2017). beta-Hydroxybutyrate: A Signaling

Metabolite. Annu Rev Nutr *37*, 51-76. 10.1146/annurev-nutr-071816-064916.

51.   Youm, Y.H., Nguyen, K.Y., Grant, R.W., Goldberg, E.L., Bodogai, M., Kim,

D., D'Agostino, D., Planavsky, N., Lupfer, C., Kanneganti, T.D., et al. (2015).

The ketone metabolite beta-hydroxybutyrate blocks NLRP3 inflammasome-

mediated inflammatory disease. Nat Med *21*, 263-269. 10.1038/nm.3804.

52.   Duking, T., Spieth, L., Berghoff, S.A., Piepkorn, L., Schmidke, A.M.,

Mitkovski, M., Kannaiyan, N., Hosang, L., Scholz, P., Shaib, A.H., et al.

(2022). Ketogenic diet uncovers differential metabolic plasticity of brain cells.

Sci Adv *8*, eabo7639. 10.1126/sciadv.abo7639.

53.   Lu, J.F., Zhu, M.Q., Xia, B., Zhang, N.N., Liu, X.P., Liu, H., Zhang, R.X.,

Xiao, J.Y., Yang, H., Zhang, Y.Q., et al. (2023). GDF15 is a major determinant

of ketogenic diet-induced weight loss. Cell Metab *35*, 2165-2182 e2167.

10.1016/j.cmet.2023.11.003.

54.   Matsuura, T.R., Puchalska, P., Crawford, P.A., and Kelly, D.P. (2023). Ketones

and the Heart: Metabolic Principles and Therapeutic Implications. Circ Res

132, 882-898. 10.1161/CIRCRESAHA.123.321872.

55.   Shah, U.A., and Iyengar, N.M. (2022). Plant-Based and Ketogenic Diets As

Diverging Paths to Address Cancer: A Review. JAMA Oncol *8*, 1201-1208.

10.1001/jamaoncol.2022.1769.

56.   Iovino, F., Engelen-Lee, J.Y., Brouwer, M., van de Beek, D., van der Ende, A.,

Valls Seron, M., Mellroth, P., Muschiol, S., Bergstrand, J., Widengren, J., and

Henriques-Normark, B. (2017). pIgR and PECAM-1 bind to pneumococcal

adhesins RrgA and PspC mediating bacterial brain invasion. J Exp Med *214*,

1619-1630. 10.1084/jem.20161668.

57.   Liu, Y., Li, J.Y., Chen, S.T., Huang, H.R., and Cai, H. (2016). The rLrp of

Mycobacterium tuberculosis inhibits proinflammatory cytokine production

and downregulates APC function in mouse macrophages via a TLR2-mediated

PI3K/Akt pathway activation-dependent mechanism. Cell Mol Immunol *13*,

729-746. 10.1038/cmi.2015.58.

58.   Steeb, B., Claudi, B., Burton, N.A., Tienz, P., Schmidt, A., Farhan, H., Maze,

A., and Bumann, D. (2013). Parallel exploitation of diverse host nutrients

enhances Salmonella virulence. PLoS Pathog *9*, e1003301.

10.1371/journal.ppat.1003301.

59. Jiang, L., Wang, P., Song, X., Zhang, H., Ma, S., Wang, J., Li, W., Lv, R., Liu,

X., Ma, S., et al. (2021). Salmonella Typhimurium reprograms macrophage

metabolism via T3SS effector SopE2 to promote intracellular replication and

virulence. Nat Commun *12*, 879. 10.1038/s41467-021-21186-4.

60. Allahverdian, S., Chaabane, C., Boukais, K., Francis, G.A., and Bochaton-

Piallat, M.L. (2018). Smooth muscle cell fate and plasticity in atherosclerosis.

Cardiovasc Res *114*, 540-550. 10.1093/cvr/cvy022.

61. Robichaud, S., Rasheed, A., Pietrangelo, A., Doyoung Kim, A., Boucher, D.M.,

Emerton, C., Vijithakumar, V., Gharibeh, L., Fairman, G., Mak, E., et al.

(2022). Autophagy Is Differentially Regulated in Leukocyte and Nonleukocyte

Foam Cells During Atherosclerosis. Circ Res *130*, 831-847.

10.1161/CIRCRESAHA.121.320047.

62. Boven, L.A., Van Meurs, M., Van Zwam, M., Wierenga-Wolf, A., Hintzen,

R.Q., Boot, R.G., Aerts, J.M., Amor, S., Nieuwenhuis, E.E., and Laman, J.D.

(2006). Myelin-laden macrophages are anti-inflammatory, consistent with

foam cells in multiple sclerosis. Brain *129*, 517-526. 10.1093/brain/awh707.

63. Peyron, P., Vaubourgeix, J., Poquet, Y., Levillain, F., Botanch, C., Bardou, F.,

Daffe, M., Emile, J.F., Marchou, B., Cardona, P.J., et al. (2008). Foamy

macrophages from tuberculous patients' granulomas constitute a nutrient-rich

reservoir for M. tuberculosis persistence. PLoS Pathog *4*, e1000204.

10.1371/journal.ppat.1000204.

64.   Weis, E.M., Puchalska, P., Nelson, A.B., Taylor, J., Moll, I., Hasan, S.S.,

Dewenter, M., Hagenmuller, M., Fleming, T., Poschet, G., et al. (2022).

Ketone body oxidation increases cardiac endothelial cell proliferation. EMBO

Mol Med *14*, e14753. 10.15252/emmm.202114753.

65.   Koopmans, M.M., Engelen-Lee, J., Brouwer, M.C., Jaspers, V., Man, W.K.,

Vall Seron, M., and van de Beek, D. (2018). Characterization of a Listeria

monocytogenes meningitis mouse model. J Neuroinflammation *15*, 257.

10.1186/s12974-018-1293-3.

1507

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7