Title

Effect of translation enhancing nascent SKIK peptide on the arrest peptides containing consecutive Proline

Author List

Yuma Nishikawa, Riko Fujikawa, Hideo Nakano, Takashi Kanamori, Teruyo Ojima-Kato*

*, the person to whom correspondence

Abstract

Ribosome arrest peptides (RAPs) such as SecM arrest peptide (SecM AP) and WPPP with consecutive Pro residues, are known to induce translational stalling in *Escherichia coli*. We demonstrate that the translation enhancing SKIK peptide tag, which consisted of four amino acid residues Ser-Lys-Ile-Lys, effectively alleviate translational arrest caused by WPPP. Moreover, the proximity between SKIK and WPPP significantly influences the extent of this alleviation, observed in both PURE cell-free protein synthesis and in vivo protein production systems, resulting in a substantial increase in the yield of proteins containing such RAPs. Furthermore, we unveil that nascent SKIK peptide tag and translation elongation factor P (EF-P) which alleviate ribosome stalling in consecutive-Pro rich protein, synergistically promote translation. A kinetic analysis based on the generation of super folder green fluorescent protein under in vitro translation reaction reveals that the ribosome turnover is enhanced by more than 10-fold when the SKIK peptide tag is positioned immediately upstream of the SecM AP sequence. Our findings provide valuable insights into optimizing protein production processes, which are essential for advancing synthetic biology applications.

Keywords

polyproline, ribosome arrest peptides, SKIK, translation enhancement, EF-P

Introduction

Protein synthesis is a fundamental process in synthetic biology, and the translation by ribosomes is an essential step in protein synthesis. Nonetheless, achieving high productivity for all proteins of interest remains a challenge, often leading to encounters with difficult-to-express proteins for reasons yet to be fully elucidated. Addressing this issue necessitates considering various strategies. These include codon optimization[1–6], loosening the secondary structure of mRNA[7–9], utilizing a combination of host-vector systems comprising ribosome binding site (RBS), promoter, and terminator[10–13]. Furthermore,

co-expression of molecular chaperons[14], and the use of fusion protein tags (e. g., maltose

binding protein domain, glutathione S-transferase, small ubiquitin-related modifier) have

proven beneficial[15,16]. optimizing culture conditions plays a pivotal role in enhancing

productivityI[17–19].

Recently, some nascent polypeptides generated during the translation process have been

reported as ribosome arrest peptides (RAPs), which cause translational stalling by

interacting with internal components of the ribosome when they pass through the ribosome

exit tunnel in the nascent state. RAP-mediated translation arrest is directed by the amino

acid sequence of the nascent polypeptide chain instead of the mRNA nucleotide sequence.

It is independent of codon usages or the secondary structure of mRNA[21].

RAPs have been found in a variety of organisms, including SecM and CmlA leader in *E.*

coli[20–23], VemP in *Vibrio alginolyticus*[24,25], and MifM in *Bacillus subtilis*[26]. Among them, SecM

is the most studied, and the partial sequence $F^{150}XXXXWIXXXXGIRAGP^{166}$ (X is any amino

acid) at the C-terminus is required and sufficient for causing ribosomal stalling[27], where new

peptidyl bond formation is inhibited when the 166th Pro is going to be synthesized (P-site:

SecM$_{1-165}$-Gly-tRNA, A-site: Pro-tRNA) at the peptidyl-transferase center (PTC). Due to the

structural rigidity and unique status as an N-alkylamino acid, Pro-tRNA at the A site slows

down the peptidyl bond formation in the RAPs containing SecM and poly-Pro[28]. As such, the

WPPP (FXXYXIWPPP: X is any amino acid) sequence, artificially created RAP, efficiently

induces ribosomal stalling due to its consecutive Pro residues[29,30]. The universally

conserved bacterial elongation factor called elongation factor P (EF-P, PDB 1UEB) is

responsible for preventing the translational stalling caused by consecutive Pro in prokaryotic

cells[31–34]. The shape and size of EF-P are similar to tRNA and it interacts with the ribosome

via the E site in the PTC[35–37]. However, Woolstenhulme et al. showed that EF-P had little or

no effect on stalling by the full-length WPPP AP (FXXYXIWPPP) and RXPP motif[30].

We have previously reported that the insertion of an "SKIK peptide tag" composed of the

four amino acids Ser-Lys-Ile-Lys at the N-terminus of difficult-to-express proteins, enhances

protein production not only in both *E. coli* in vivo and in vitro systems but also in yeast[38].

More recently, we revealed that changing nucleotide sequences corresponding to these four

amino acids had no effect on protein production and that the nascent SKIK peptide

generated during the translation process could enhance the translation process, but not

transcription[39]. Surprisingly, ribosomal stalling by RAPs such as SecM AP

(FSTPVWISQAQGIRAGP) and chloramphenicol induced CmlA leader (KNAD) was

canceled by a nascent SKIK peptide generated immediately upstream of the RAPs.

Moreover, the combined investigation of the SKIK peptide tag and SecM AP suggested that

the SKIK peptide, even when positioned internally within the protein, could enhance

translation. Additionally, it was suggested that the distance between SKIK and the AP motif

might be crucial for SKIK to effectively alleviate translational stalling induced by RAPs.

However, it should be noted that in our previous study, the SKIK peptide tag showed no

significant impact on translation when it was positioned immediately upstream of the WPPP

AP (FQKYGIWPPP)[39].

Based on our recent findings, we hypothesized that ribosomal stalling induced by RAPs

containing a poly-Pro motif could be alleviated by optimizing the distance between the

nascent SKIK peptide and the poly-Pro motif. In this study, we investigated the positional

effects of the SKIK peptide tag on the ribosomal stalling caused by the poly-Pro AP motif in

both *E. coli* in vitro and in vivo systems. Additionally, we uncovered that EF-P and the SKIK

peptide tag synergistically enhance translation, resulting in a significant increase in protein

production. Translation promotion was observed only when the SKIK peptide was tagged,

not by adding the synthesized free SKIK peptide to the in vitro translation system,

emphasizing the importance of the nascent state of the SKIK peptide in counteracting arrest.

Furthermore, we performed a kinetic analysis using an in vitro translation system to

elucidate how the SKIK peptide tag influences on translation.

Results and Discussion

Influences of the distance between SKIK and WPPP arrest peptide

As described above, the ribosomal stalling by WPPP (FQKYGIWPPP), an artificial AP

containing consecutive Pro, was not affected by the SKIK peptide tagging while the

ribosomal stalling by SecM AP was more effectively reduced when the SKIK peptide tag

was positioned closer[39]. We therefore created DNA constructs corresponding to the amino

acid sequence depicted in Figure 1 to explore the hypothesis that an optimal position for the

SKIK peptide tag exists concerning the poly-Pro motif. Here, the SKIK tag positioned

immediately before the reported AP motif (FQKYGIWPPP) of WPPP was designated as the

standard (+0 aa). Minus and plus numbers indicate the count of deleted amino acid residues

from F in the WPPP AP motif and inserted G, respectively. To confirm that only the four

amino acids "WPPP" function as an arrestor, a version with -6 aa (SKIKWPPP), where the

SKIK moiety was substituted with GGGG or AAAA, was also constructed.

In the results of in vitro translation using mRNA, the fluorescence intensity, reflecting the

synthesized sfGFP quantity after 90 minutes of reaction, demonstrated a rising trend as the

distance between SKIK and WPPP decreased (Figure 2A). Particularly noteworthy was the

fluorescence intensity at -5 aa (28,860), where SKIK and WPPP AP were brought closer,

being approximately 5 times higher than at +0 aa (5,660), indicating the effective alleviation

of ribosomal stalling induced by poly-Pro by the SKIK peptide tag. Western blotting analysis

for His tag detection also revealed an increase in the full-length protein synthesis as the

distance between SKIK and poly-Pro decreased. Moreover, insertions of G at +4−7 aa and

+13 aa also exhibited approximately twice the fluorescence intensity of +0 aa, suggesting

the possibility of multiple conditions where SKIK can effectively mitigate the ribosomal

stalling caused by poly-Pro. It is worth noting that the fluorescence intensity for -6 aa

(SKIKWPPP) was 20,200, whereas GGGG (sequence GGGGWPPP) and AAAA

(AAAWPPP) exhibited significantly lower intensities. This indicates that the four amino acids

WPPP alone can induce translational stalling, and SKIK counteracts this stalling. In the RAP

employed in this study, the WPPP AP motif (FQKYGIWPPP), it was reported that the

presence of F, Y, I, and W upstream of PPP is essential for translational arrest. However,

Elgama et al. demonstrated that the strength of translation arrest induced by a poly-Pro

motif is influenced by the specific amino acid residues immediately upstream[40]. This may

support our findings that SKIK has the potential to alleviate translational arrest induced by

poly-Pro, and its effectiveness seems to depend on the distance between SKIK and

poly-Pro motif.

The translation rate, determined by monitoring the real-time synthesis of sfGFP from mRNA,

correlated with the amount of the protein obtained after 90 minutes of reaction (Figure 2A, B).

Hence, under the 90-minute CFPS conditions in this experiment, it can be inferred that the

final protein amount is indeed influenced by the translation rate. Nevertheless, as the

distance increased beyond "+8 aa," a declining trend in synthesis levels was observed.

Upon comparison with the protein production levels of "GGGG," a sequence excluding SKIK,

it can be inferred that translational arrest was alleviated in all sequences containing SKIK,

albeit with varying degrees.

The more prominent trend of translational arrest alleviation observed in vivo may be

attributed to the fact that in vitro reactions were only 90 minutes, while in vivo, we conducted

longer expression induction. Additionally, the presence of natural EF-P in the living cells

could contribute to this difference.

To the best of our knowledge, there have been no reports on function of the SKIK peptide to

alleviate the ribosomal stalling by poly-Pro or on the importance of their positional

relationship. Therefore, our findings will provide a critical and novel insight in the field of

synthetic biology.

Kinetic analysis of the protein synthesis

We then performed a kinetic analysis of translation to better understand the effect of the

SKIK peptide tag on translation. Here, SecM AP-sfGFP with and without the SKIK peptide

tag and simple translation reaction model were used (Figure 3A, B). For this analysis, we

designate mRNA as "Substrate (S)," sfGFP as "Product (P)," and ribosome as "Enzyme (E)"

for convenience. We applied the following three assumptions to simplify the analysis: 1)

there is no folding rate limitation for sfGFP, 2) there are sufficient elements necessary for

translation, and no molecules inhibit the translation reaction, and 3) mRNA and ribosome do

not dissociate until the translation of single protein molecule is completed.

The translation rate v was defined as follows.

$v = k_2[\text{ES}]$ $\qquad\qquad$ (1)

The following equation is given by the steady-state approximation.

$[\text{ES}] = \frac{k_1}{k_2}[\text{E}][\text{S}]$ $\qquad\qquad$ (2)

On the other hand, [E] can be expressed as follows:

$[\text{E}] = [\text{E}]_0 - [\text{ES}]$ $\qquad\qquad$ (3)

The following equation can be derived by substituting (3) into (2).

$[\text{ES}] = \frac{[\text{E}]_0[\text{S}]}{k_2/k_1 + [\text{S}]}$ $\qquad\qquad$ (4)

Based on the assumption that $k_2[\text{E}]_0 = V_{max}$, substituting equation (1) into (4) results in a

Michaelis-Menten-like equation as follows:

$v = \frac{V_{max}[\text{S}]}{k_2/k_1 + [\text{S}]}$ $\qquad\qquad$ (5)

Where $[\text{S}] = k_2/k_1$ when the translation rate is 1/2 V_{max}.

The translation rate was assessed by continuously monitoring the fluorescence intensity of

sfGFP generated by CFPS using mRNA templates ranging from 0.316 to 24.2 µM. As

shown in Figure 3C, the rate of sfGFP production escalated proportionally with the quantity

of mRNA template utilized. Based on these measurements, the translation rate was

calculated (Table 1). It was observed that the translation rate increased by over 10-fold

when the SKIK peptide was tagged, regardless of mRNA concentrations, peaking at a

maximum enhancement of 16.8-fold at 15.5 µM.

Utilizing the translation rates calculated for each mRNA concentration in Table 2, nonlinear

regression analysis was conducted using the R package "renz"[41](Figure 3D). The

regression curve aptly fits the experimental data, enabling the calculation of, V_{max} and k_2/k_1

based on Equation (5). As a result, the V_{max} values for SKIK peptide tag absence and

presence were determined to be 0.20 mM$_{GFP}$/M$_{Ribosome}$/s and 4.50 mM$_{GFP}$/M$_{Ribosome}$/s,

respectively. The addition of SKIK peptide tag led to a 22.5-fold increase in V_{max}.

The calculated k_2/k_1 values were 8.45 μM without SKIK and 21.9 μM with SKIK,

representing a 2.59-fold increase upon SKIK peptide tag addition (Table 3). $[E]_0$ = 1 μM and

$V_{max} = k_2 [E]_0$ were used to calculate k_2, and subsequently, k_1 was calculated from the

non-linear regression analysis-derived k_2/k_1. The results are also summarized in Table 3. As

evident from this, the addition of the SKIK peptide tag led to values of k_1 and k_2 that were

8.65-fold and 22.5-fold higher, respectively.

Since the RBS is the same in the two examined mRNA sequences, it seems unlikely that the

affinity between the ribosome and mRNA changes significantly. Therefore, the increase in k_1

resulting from the addition of SKIK peptide tag suggests an augmentation in the pool of

ribosomes capable of initiating translation. Similarly, the increase in k_2 implies an

acceleration in the rate of translation from initiation to termination. These kinetic analyses

indicate that the addition of the SKIK peptide tag could enhance the ribosomal turnover rate.

Regarding studies on kinetic analysis of translation, previous analyses have utilized the

incorporation of radiolabeled amino acids into synthesized proteins[42], and more recently,

elongation rates per codon have been calculated based on ribosome density observed on

mRNA through ribosome profiling [43,44]. To our knowledge, our analysis, based on the

production of sfGFP as an indicator, has not been reported previously and is a first attempt.

Elongation rate per codon is commonly used as an indicator of translation efficiency, with

reported values ranging from 1.66 codons/s[45] and in *E. coli* in vivo to 10 codons/s[42] in vitro,

4.2 codons/s for *S. cerevisiae,* and 5.2 codons/s for mouse stem cells[43]

In this study, SKIK-SecM AP-sfGFP gene used consists of 267 codons. The apparent

elongation rate per codon calculated from the estimated V_{max} of SKIK-SecM AP-sfGFP was

1.2 codons/s. Although this value is lower than those reported in previous studies,

considering that ribosomal stalling by SecM is not completely alleviated by SKIK peptide tag

[39], and our reaction model disregards the translation initiation and termination stages which

require time, the calculated elongation rate appears reasonable. Although it should be noted

that use of this model might be limited, we concluded that the simple translation rate

analysis method proposed in this study worked well for our purpose.

Effect of EF-P and SKIK peptide

EF-P is a translation factor reported to promote the synthesis of poly-Pro motif. Therefore,

we investigated how the SKIK peptide tag, capable of relieving ribosomal stalling induced by

poly-Pro, and EF-P, would affect the translation of sequences containing poly-Pro. We

examined using four sequences including -5 aa, +0 aa, GGGG, and AAAA. Among these,

the ribosomal stalling was most effectively alleviated by SKIK peptide tag alone in -5 aa

(Figure 2A). As a result of EF-P addition at a final concentration of 1 μM, the protein

production was increased to 1.51-fold and 4.40-fold in -5 aa and +0 aa, respectively,

compared with no additive. In particular, the protein production was highest for -5 aa with

EF-P, indicating a synergistic effect by SKIK peptide tag and EF-P (Figure 4A). It suggests

the possibly that EF-P and SKIK peptide tag individually promote translation in this case. By

contrast, GGGG and AAAA, which do not contain SKIK, did not show any change in protein

levels with addition of EF-P. To confirm the nascent state of SKIK peptide is essential for

translation promotion, CFPS of four mRNA templates was conducted with the addition of the

chemically synthesized free SKIK peptides at 100 μM which was 100 times higher

concentration of ribosome in CFPS reaction. As expected, no influence was observed

(Figure 4B). Huter et al. demonstrated that EF-P recognizes the P-site tRNA and the E-site

mRNA codon, stabilizing the conformation of the P-site tRNA and thereby promoting

peptidyl bond formation[46]. Taken together with the results of additional of free SKIK peptide

and our previous study[39], it is likely that nascent state of SKIK peptide tag is crucial for

translation promotion. Therefore, we expect that the mechanism of action of externally

functioning EF-P and SKIK peptide tag at the nascent state may be different. Since the

mechanism of translation enhancement by SKIK peptide tag has not yet been elucidated,

we believe that further structural analysis will be necessary.

Conclusions

In this study, we revealed that the translation enhancing SKIK peptide tag effectively

alleviates ribosomal stalling induced by the WPPP when the distance between them is

closer in both the PURE cell-free protein synthesis and in vivo protein production systems,

resulting in significant increment of production of proteins containing such poly-Pro motif. A

kinetic analysis based on the simple translation model showed that the ribosome turnover is

enhanced by more than 10-fold when the SKIK peptide tag. Furthermore, we found that the

use of both SKIK peptide tag and EF-P synergistically enhanced translation for proteins

containing poly-Pro. Our findings and simple strategy have practical significance in

enhancing protein production and contribute to the advancement of synthetic biology.

Figure Legends

Figure 1. DNA constructs used to study the distance between the SKIK peptide tag

and poly-Pro motif of WPPP.

The amino acid sequences including start M encoded immediately upstream of sfGFP are

shown. SKIK and inserted G are shown in red and gray, respectively.

Figure 2. Influences of the distance between SKIK peptide and WPPP poly-Pro motif

on protein production and translation rate.

The sample names and sequences are corresponding to those of Figure 1.

A) Fluorescence intensity and Western blotting analysis of sfGFP detecting His tag. In

vitro translation was carried out using mRNA templates (1.66 µM) for 90 min (n = 2).

B) Translation rate calculated from the real-time monitoring of the fluorescence intensity

of the generated sfGFP during in vitro translation (n = 2, mRNA concentration; 1.66

µM).

C) Fluorescence intensity of the produced sfGFP in *E. coli* BL21(DE3) living cell

expression system (n = 6). SDS-PAGE and CBB staining analysis of the sfGFP bands

are shown.

Figure 3. Kinetic analysis of translation by PURE cell-free protein synthesis.

A) mRNA constructs and amino acid sequences of SecM AP and SKIK-SecM AP.

B) Translation reaction model: E, S, and P represent the ribosome, mRNA, and sfGFP

(product), respectively. k_1 and k_2 denote the rate constants for the reactions indicated

by the respective arrows. The ES complex encompasses all stages from initiation to

termination of translation.

C) Real-time fluorescence intensity measurements. Generation of sfGFP of various

mRNA concentrations was monitored during 90min (5,400 s) CFPS reaction. Plus and

minus in the parentheses indicate the presence and absence of the SKIK peptide tag,

respectively.

D) Non-linear regression analysis of translation rate.

Figure 4. Effect of EF-P and free SKIK peptide on protein production in CFPS.

A) Fluorescence intensity of sfGFP by in vitro translation using mRNA (1.61 µM) after 90

min reaction. +/– indicates the presence/absence of EF-P (1 µM).

B) Fluorescence intensity of sfGFP by in vitro translation using mRNA after 90 min

reaction. The mRNA concentration was 1.19 µM except for SecM AP and SKIK-SecM

AP (2.37 μM). +/− indicates the presence/absence of synthesized free SKIK peptide

(100 μM).

Tables

Table 1. Translation rate

mRNA (µM)	v (mM$_{GFP}$/M$_{Ribosome}$/s)		$V_{SKIK}/V_{No\ tag}$
	No tag	SKIK	
24.2	0.157	2.25	14.3
15.5	0.121	2.02	16.8
10.7	9.73×10^{-2}	1.61	16.5
6.33	0.105	0.823	7.82
3.16	5.38×10^{-2}	0.514	9.56
1.58	2.69×10^{-2}	0.292	10.9
0.316	5.48×10^{-3}	5.72×10^{-2}	10.4
No template	9.52×10^{-4}		

287 The translation rate, v, calculated from the plot shown in Figure 3D, and the ratio of v with
288 and without SKIK are presented as $V_{SKIK}/V_{No\ tag}$. No tag and SKIK mean SecM AP-sfGF and
289 SKIK-SecM AP-sfGFP, respectively.
290
291

292 **Table 2. Parameters on translation calculated based on real-time translation**

293 **monitoring.**

	V_{max} (mM$_{GFP}\cdot$M$_{Ribosome}^{-1}\cdot$s^{-1})	k_2/k_1 (μM)	k_1	k_2
No tag	0.20	8.45	2.37×10^7	0.20×10^3
SKIK	4.50	21.9	20.5×10^7	4.50×10^3
SKIK/No tag (-fold)	22.5	2.59	8.65	22.5

V_{max} and k_2/k_1 were calculated using non-linear regression analysis with the renz software,

as shown in Figure 3D. The rate constants k_1 and k_2 were estimated from the values of V_{max}

and k_2/k_1 obtained through the non-linear regression analysis. The units of k_1 and k_2 are

M$_{GFP}\cdot$M$_{Ribosome}^{-2}\cdot$s$^{-1}\cdot$M$_{mRNA}^{-1}$ and M$_{GFP}\cdot$M$_{Ribosome}^{-2}\cdot$s^{-1}, respectively. M$_{Ribosome}$ and M$_{mRNA}$

represent the concentrations of ribosome and mRNA in mol/L, respectively.

Methods

DNA primer list

DNA primers used in this research are summarized in Supplementaly Table 1.

Plasmid construction

Total 21 plasmids were prepared by whole plasmid PCR using our previously constructed

pET22b-SKIK-WPPP-sfGFP-His as the template and DNA primers listed in Supplementary

Table 1. *Dpn*I treated PCR products were purified with silica column (FastGene Gel/PCR

Extraction Kit, Nippon Genetics Co., Ltd., Tokyo, Japan) and then HiFi assembly reaction

(New England Biolabs, Ipswich, MA) was carried out to connect each end of linear DNA

products. The plasmids were prepared using plasmid mini prep kit (FastGene Plasmid Mini

Kit, Nippon Genetics Co., Ltd.) from overnight cultured *E. coli* DH5alpha transformants. The

sequences of all plasmids confirmed by Sangar sequencing are available in Supporting

Information.

Cell-free protein synthesis (CFPS).

DNA fragments containing T7 promoter and terminator used for CFPS were prepared by

PCR with the primer pair F1 and R1, constructed plasmids as the template, and Tks Gflex

DNA polymerase (98°C 1 min; (98°C 10 sec; 50°C 5 sec; 68°C 30 sec) × 30 cycles; 68°C

1min; 12°C ∞, Takara, Kusatsu, Japan). Amplified DNA was purified using silica column and

used for in vitro mRNA synthesis. For in vitro mRNA synthesis, the T7 RiboMAX

Large-Scale RNA Production System (Promega, Madison, WI, USA) was used and the

synthesized mRNAs were purified with NucleoSpin RNA Plus (Takara). The concentration of

mRNA was measured by Nano Drop One (Thermo Fisher scientific, MA, USA) and the

molecular weight were calculated using RNA Molecular Weight Calculator on AAT Bioquest

(https://www.aatbio.com).

CFPS reactions were carried out with PURE*frex* 2.1 (GeneFrontier Corporation, Kashiwa,

Japan) under the following condition. Solution I, 4 µL; Solution II, 0.5 µL; Solution III (20 µM

ribosome), 0.5 µL; 10 mM cysteine, 0.5 µL; RNase Inhibitor (TOYOBO, Osaka, Japan), 1

µL; at final reaction volume of 10 µL; incubated at 37°C for 90 min. The final concentration of

the template mRNA depends on the examination. An elongation factor EF-P (GeneFrontier

Corporation) and synthesized SKIK peptide dissolved in nuclease-free water (94.6% purity,

GenScript, Tokyo) was added to achieve a final concentration of 1 µM and 100 µM,

respectively, as needed.

In vivo expression

Terrific broth (Bacto tryptone, 12 g; yeast extract, 24 g; KH_2PO_4, 6.8 g; $Na_2PO_4 \cdot 12H_2O$, 7.1

g; $MgSO_4$, 0.15 g; $(NH_4)_2SO_4$, 2.58 g; 10%(w/v) glucose, 5 mL; and 8%(w/v) lactose; 25 mL,

per 1 L, adjusted pH 7.4) was used as the autoinduction medium. The single colonies of

ECOS *E. coli* BL21(DE3) (Nippon Gene, Tokyo, Japan) transformed with each plasmid were

inoculated to 4 mL of LB liquid and grown with shake at 37°C for 19 h. Then the aliquots (50

µL) were transferred to the 1 mL of terrific broth and cultured at 30°C until the OD600

reached 2.3–6.9. All media were supplemented with 100 µg/mL ampicillin. After collecting

the bacterial cells by centrifugation and washing with phosphate buffer saline buffer (PBS,

137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4; pH 7.4), the OD600 values

of each sample were adjusted to the same. Protein was extracted from the cells using

E.coli/Yeast Protein Extraction Buffer (Tokyo Chemical Industry Co., Ltd., Tokyo, Japan),

and the lysate fraction was used for analysis.

Preparation of Standard sfGFP

ECOS *E. coli* BL21(DE3) was transformed with pET22b-sfGFP-His and cultured in 100 mL

of LB medium at 37°C with shake until the OD600 value reached 0.4~0.6. IPTG was then

added at a final concentration of 0.1 mM and cultured at 37°C for another 5 h. Bacterial cells

were collected by centrifugation (4°C, 9000 rpm, 10 min), resuspended in 10 mL of PBS,

and then the bacteria were crushed by sonication (30 min). The supernatant of the lysate

was applied to a Ni-NTA agarose HP column (Fujifilm Wako Pure Chemical Corporation,

Osaka, Japan) equilibrated with Binding buffer (20 mM sodium phosphate buffer, 0.5 M

NaCl, 20 mM imidazole, pH 8.0), and the column was washed 3 times with 6 mL of binding

buffer and eluted 6 times with 1 mL of Elution buffer (20 mM sodium phosphate buffer, 0.5 M

NaCl, 500 mM imidazole, pH 8.0). The eluate was added to the dialysis membrane and

dialyzed for a total of 24 h (buffer was changed twice during the dialysis). The concentration

of purified sfGFP was calculated from the A280 nm value (A.U.) measured by Nano Drop

One. Molecular weight (27788.3 Da) and molar extinction coefficient (19,035 $M^{-1}cm^{-1}$) of

sfGFP were calculated using ProtParam tool on Expasy (https://www.expasy.org)[47].

Fluorescence measurement

Each CFPS reaction solution was diluted 25-fold with water and dispensed 50 µL per well

into Flat Bottom Microfluor Plates Black (Thermo Fisher scientific, MA, USA), and the

fluorescence intensity of sfGFP was measured by a microplate reader (Infinite 200 PRO,

TECAN, ZH, Switzerland) at excitation wavelength 485 nm (bandwidth 9 nm)/emission

wavelength of 535 nm (bandwidth 20 nm).

Real-time monitoring of the translation in CFPS

The reaction was performed using a real-time PCR system (StepOnePlus Real-Time PCR

System, Life Technologies, CA, USA) by incubating at 37°C, and the fluorescence intensity

was measured every minute after 30 s. Fluorescence intensity per sfGFP molecule was

calculated using purified sfGFP (14.6 μM, 1.46 μM, 0.146 μM, 0.0146 μM, and 0.00146 μM)

as the standard. Translation rate defined as the generated sfGFP per ribosome per time

was calculated from the rate of the fluorescence increase where the slope was the greatest

and linear for all samples.

Western Blotting

The samples (1 μL) were subjected to SDS-PAGE and transferred to a nitro cellulose

membrane (Bio-Rad Laboratories, CA, USA). The membrane was blocked using 3% skim

milk in PBST (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.05 %

Tween-20; pH 7.4) for 1 h at room temperature. After washing three times with PBST, the

membrane was incubated with Anti-His-tag mAb-HRP-DirectT (MBL, Tokyo, Japan) at

1:5000 dilution with Can Get Signal Solution 2 (Toyobo, Osaka, Japan) for 1 h at room

temperature. After washing with PBST three times, blots were visualized with TMB solution

(Nacalai tesque, Kyoto, Japan).

Supporting Information

The Supporting Information including DNA sequences for plasmid with GenBank format and

additional experimental data (PDF) is available free of charge.

Author Information

Corresponding author

Teruyo Ojima-Kato - Laboratory of Molecular Biotechnology, Graduate School of

Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601,

Japan

https://orcid.org/0000-0001-5290-5176; E-mail: teruyo@agr.nagoya-u.ac.jp

Authors

Yuma Nishikawa - Laboratory of Molecular Biotechnology, Graduate School of

Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601,

Japan

Riko Fujikawa - Laboratory of Molecular Biotechnology, Graduate School of Bioagricultural

Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan

Hideo Nakano - Laboratory of Molecular Biotechnology, Graduate School of Bioagricultural

Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan

https://orcid.org/0000-0001-6133-9794

Takashi Kanamori - GeneFrontier Corporation, 273-1 Kashiwa, Kashiwa, Chiba 277-0005,

Japan

https://orcid.org/0000-0001-9150-5968

Author contributions

Y.N., H.N., T.K., and T.O.K. designed the study and analyzed the data. Y.N. performed

CFPS experiments. R.F. performed in vivo expression experiments. Y.N. and T.O.K. wrote

the manuscript. T.O.K. supervised the project.

Acknowledgment

This research was supported by Japan Science and Technology agency FOREST Program

(grant No. JPMJFR2204), KAKENHI 23K04989, and GteXProgram Japan Grant Number

JPMJGX23B6 and JPMJGX23B4. PURE*frex* 2.1 and supplemental components used in this

study were provided by GeneFrontier Corporation. We thank Mrs. Kana Yamauchi (Nagoya

University) for her help to the preparation of the purified sfGFP. We also thank Prof. Hisashi

Hemmi (Nagoya University) and Prof. Tomokazu Ito (Nagoya University) for their kind

advice and help on translation analysis.

Abbreviations

RAP, Ribosomal arrest peptide

RBS, ribosome binding site

PTC, peptidyl-transferase center

CFPS, cell-free protein synthesis

sfGFP, superfolder green fluorescent protein

References

(1)   Makrides, S. C. Strategies for Achieving High-Level Expression of Genes in
438   Escherichia Coli. *Microbiol Rev* **1996**, *60* (3).
439   https://doi.org/10.1128/mr.60.3.512-538.1996.

440   (2)   Gvritishvili, A. G.; Leung, K. W.; Tombran-Tink, J. Codon Preference Optimization
441   Increases Heterologous PEDF Expression. *PLoS One* **2010**, *5* (11).
442   https://doi.org/10.1371/journal.pone.0015056.

443   (3)   Bentele, K.; Saffert, P.; Rauscher, R.; Ignatova, Z.; Blüthgen, N. Efficient Translation
444   Initiation Dictates Codon Usage at Gene Start. *Mol Syst Biol* **2013**, *9*.
445   https://doi.org/10.1038/msb.2013.32.

446   (4)   Wu, X.; Jörnvall, H.; Berndt, K. D.; Oppermann, U. Codon Optimization Reveals
447   Critical Factors for High Level Expression of Two Rare Codon Genes in Escherichia
448   Coli: RNA Stability and Secondary Structure but Not TRNA Abundance. *Biochem*
449   *Biophys Res Commun* **2004**, *313* (1). https://doi.org/10.1016/j.bbrc.2003.11.091.

450   (5)   Saito, Y.; Kitagawa, W.; Kumagai, T.; Tajima, N.; Nishimiya, Y.; Tamano, K.; Yasutake,
451   Y.; Tamura, T.; Kameda, T. Developing a Codon Optimization Method for Improved
452   Expression of Recombinant Proteins in Actinobacteria. *Sci Rep* **2019**, *9* (1).
453   https://doi.org/10.1038/s41598-019-44500-z.

454   (6)   Gingold, H.; Dahan, O.; Pilpel, Y. Dynamic Changes in Translational Efficiency Are
455   Deduced from Codon Usage of the Transcriptome. *Nucleic Acids Res* **2012**, *40* (20),
456   10053–10063. https://doi.org/10.1093/nar/gks772.

457   (7)   Jana, S.; Deb, J. K. Strategies for Efficient Production of Heterologous Proteins in
458   Escherichia Coli. *Applied Microbiology and Biotechnology*. 2005.
459   https://doi.org/10.1007/s00253-004-1814-0.

460   (8)   Emory, S. A.; Bouvet, P.; Belasco, J. G. A 5′-Terminal Stem-Loop Structure Can
461   Stabilize MRNA in Escnerichia Coli. *Genes Dev* **1992**, *6* (1).
462   https://doi.org/10.1101/gad.6.1.135.

463   (9)   Frederick, M. I.; Heinemann, I. U. Regulation of RNA Stability at the 3′ End. *Biological*
464   *Chemistry*. 2021. https://doi.org/10.1515/hsz-2020-0325.

465   (10)   Gold, L. Posttranscriptional Regulatory Mechanisms in Escherichia Coli. *Annual*
466   *Review of Biochemistry*. 1988. https://doi.org/10.1146/annurev.bi.57.070188.001215.

467   (11)   De Smit, M. H.; Van Duin, J. Secondary Structure of the Ribosome Binding Site
468   Determines Translational Efficiency: A Quantitative Analysis. *Proc Natl Acad Sci U S*
469   *A* **1990**, *87* (19). https://doi.org/10.1073/pnas.87.19.7668.

(12)   Hall, M. N.; Gabay, J.; Débarbouillé, M.; Schwartz, M. A Role for MRNA Secondary
Structure in the Control of Translation Initiation. *Nature* **1982**, *295* (5850).
https://doi.org/10.1038/295616a0.
(13)   Kastelein, R. A.; Berkhout, B.; Van Duin, J. Opening the Closed Ribosome-Binding
Site of the Lysis Cistron of Bacteriophage MS2. *Nature* **1983**, *305* (5936).
https://doi.org/10.1038/305741a0.
(14)   Kyratsous, C. A.; Silverstein, S. J.; DeLong, C. R.; Panagiotidis, C. A.
Chaperone-Fusion Expression Plasmid Vectors for Improved Solubility of
Recombinant Proteins in Escherichia Coli. *Gene* **2009**, *440* (1–2).
https://doi.org/10.1016/j.gene.2009.03.011.
(15)   Marblestone, J. G. Comparison of SUMO Fusion Technology with Traditional Gene
Fusion Systems: Enhanced Expression and Solubility with SUMO. *Protein Science*
2006, *15* (1), 182–189. https://doi.org/10.1110/ps.051812706.
(16)   LaVallie, E. R.; McCoy, J. M. Gene Fusion Expression Systems in Escherichia Coli.
Curr Opin Biotechnol **1995**, *6* (5). https://doi.org/10.1016/0958-1669(95)80083-2.
(17)   Corisdeo, S.; Wang, B. Functional Expression and Display of an Antibody Fab
Fragment in Escherichia Coli: Study of Vector Designs and Culture Conditions.
Protein Expr Purif **2004**, *34* (2). https://doi.org/10.1016/j.pep.2003.11.020.
(18)   Gupta, P.; Sahai, V.; Bhatnagar, R. Enhanced Expression of the Recombinant Lethal
Factor of Bacillus Anthracis by Fed-Batch Culture. *Biochem Biophys Res Commun*
2001, *285* (4). https://doi.org/10.1006/bbrc.2001.5282.
(19)   Yee, L.; Blanch, H. W. Recombinant Protein Expression in High Cell Density
Fed-Batch Cultures of Escherichia Coli. *Bio/Technology* **1992**, *10* (12).
https://doi.org/10.1038/nbt1292-1550.
(20)   Muto, H.; Nakatogawa, H.; Ito, K. Genetically Encoded but Nonpolypeptide
Prolyl-TRNA Functions in the A Site for SecM-Mediated Ribosomal Stall. *Mol Cell*
2006, *22* (4), 545–552. https://doi.org/10.1016/j.molcel.2006.03.033.
(21)   Nakatogawa, H.; Ito, K. *Secretion Monitor, SecM, Undergoes Self-Translation Arrest*
in the Cytosol the Export Status of SecM Affects Translation of SecA Encoded in the
Same Messenger RNA Strand. The Cis-Specific Effect Suggests That the SecM
Translocation In; 2001; Vol. 7.
(22)   Lovett, P. S. Translation Attenuation Regulation of Chloramphenicol Resistance in
Bacteria - A Review. *Gene* **1996**, *179* (1).
https://doi.org/10.1016/S0378-1119(96)00420-9.
(23)   Ishii, E.; Chiba, S.; Hashimoto, N.; Kojima, S.; Homma, M.; Ito, K.; Akiyama, Y.; Mori,
H.; Randall, L. L. Nascent Chain-Monitored Remodeling of the Sec Machinery for

Salinity Adaptation of Marine Bacteria. *Proc Natl Acad Sci U S A* **2015**, *112* (40). https://doi.org/10.1073/pnas.1513001112.

(24)   Mori, H.; Sakashita, S.; Ito, J.; Ishii, E.; Akiyama, Y. Identification and Characterization of a Translation Arrest Motif in VemP by Systematic Mutational Analysis. *Journal of Biological Chemistry* **2018**, *293* (8), 2915–2926. https://doi.org/10.1074/jbc.M117.816561.

(25)   Kolář, M. H.; Nagy, G.; Kunkel, J.; Vaiana, S. M.; Bock, L. V.; Grubmüller, H. Folding of VemP into Translation-Arresting Secondary Structure Is Driven by the Ribosome Exit Tunnel. *Nucleic Acids Res* **2022**, *50* (4), 2258–2269. https://doi.org/10.1093/nar/gkac038.

(26)   Chiba, S.; Lamsa, A.; Pogliano, K. A Ribosome-Nascent Chain Sensor of Membrane Protein Biogenesis in Bacillus Subtilis. *EMBO Journal* **2009**, *28* (22). https://doi.org/10.1038/emboj.2009.280.

(27)   Nakatogawa, H.; Ito, K. The Ribosomal Exit Tunnel Functions as a Discriminating Gate. *Cell* **2002**, *108* (5), 629–636. https://doi.org/10.1016/S0092-8674(02)00649-9.

(28)   Pavlov, M. Y.; Watts, R. E.; Tan, Z.; Cornish, V. W.; Ehrenberg, M.; Forster, A. C. *Slow Peptide Bond Formation by Proline and Other N-Alkylamino Acids in Translation*; 2008. www.pnas.orgcgidoi10.1073pnas.0809211106.

(29)   Tanner, D. R.; Cariello, D. A.; Woolstenhulme, C. J.; Broadbent, M. A.; Buskirk, A. R. Genetic Identification of Nascent Peptides That Induce Ribosome Stalling. *Journal of Biological Chemistry* **2009**, *284* (50). https://doi.org/10.1074/jbc.M109.039040.

(30)   Woolstenhulme, C. J.; Parajuli, S.; Healey, D. W.; Valverde, D. P.; Petersen, E. N.; Starosta, A. L.; Guydosh, N. R.; Johnson, W. E.; Wilson, D. N.; Buskirk, A. R. Nascent Peptides That Block Protein Synthesis in Bacteria. *Proc Natl Acad Sci U S A* **2013**, *110* (10). https://doi.org/10.1073/pnas.1219536110.

(31)   Doerfel, L. K.; Wohlgemuth, I.; Kothe, C.; Peske, F.; Urlaub, H.; Rodnina, M. V. EF-P Is Essential for Rapid Synthesis of Proteins Containing Consecutive Proline Residues. *Science (1979)* **2013**, *339* (6115). https://doi.org/10.1126/science.1229017.

(32)   Ude, S.; Lassak, J.; Starosta, A. L.; Kraxenberger, T.; Wilson, D. N.; Jung, K. Translation Elongation Factor EF-P Alleviates Ribosome Stalling at Polyproline Stretches. *Science (1979)* **2013**, *339* (6115). https://doi.org/10.1126/science.1228985.

(33)   Peil, L.; Starosta, A. L.; Lassak, J.; Atkinson, G. C.; Virumäe, K.; Spitzer, M.; Tenson, T.; Jung, K.; Remme, J.; Wilson, D. N. Distinct XPPX Sequence Motifs Induce Ribosome Stalling, Which Is Rescued by the Translation Elongation Factor EF-P. *Proc Natl Acad Sci U S A* **2013**, *110* (38). https://doi.org/10.1073/pnas.1310642110.

(34) Hersch, S. J.; Wang, M.; Zou, S. B.; Moon, K. M.; Foster, L. J.; Ibba, M.; Navarre, W.
W. Divergent Protein Motifs Direct Elongation Factor P-Mediated Translational
Regulation in Salmonella Enterica and Escherichia Coli. *mBio* **2013**, *4* (2).
https://doi.org/10.1128/mBio.00180-13.
(35) Blaha, G.; Stanley, R. E.; Steitz, T. A. Formation of the First Peptide Bond: The
Structure of EF-P Bound to the 70S Ribosome. *Science (1979)* **2009**, *325* (5943).
https://doi.org/10.1126/science.1175800.
(36) Hanawa-Suetsugu, K.; Sekinet, S. I.; Sakai, H.; Hori-Takemoto, C.; Terada, T.; Unzai,
S.; Tame, J. R. H.; Kuramitsu, S.; Shirouzu, M.; Yokoyama, S. Crystal Structure of
Elongation Factor P from Thermus Thermophilus HB8. *Proc Natl Acad Sci U S A* **2004**,
101 (26). https://doi.org/10.1073/pnas.0308667101.
(37) Yanagisawa, T.; Sumida, T.; Ishii, R.; Takemoto, C.; Yokoyama, S. A Paralog of
Lysyl-TRNA Synthetase Aminoacylates a Conserved Lysine Residue in Translation
Elongation Factor P. *Nat Struct Mol Biol* **2010**, *17* (9).
https://doi.org/10.1038/nsmb.1889.
(38) Ojima-Kato, T.; Nagai, S.; Nakano, H. N-Terminal SKIK Peptide Tag Markedly
Improves Expression of Difficult-to-Express Proteins in Escherichia Coli and
Saccharomyces Cerevisiae. *J Biosci Bioeng* **2017**, *123* (5), 540–546.
https://doi.org/10.1016/j.jbiosc.2016.12.004.
(39) Ojima-Kato, T.; Nishikawa, Y.; Furukawa, Y.; Kojima, T.; Nakano, H. Nascent MSKIK
Peptide Cancels Ribosomal Stalling by Arrest Peptides in Escherichia Coli. *Journal of
Biological Chemistry* **2023**, *299* (5). https://doi.org/10.1016/j.jbc.2023.104676.
(40) Elgamal, S.; Katz, A.; Hersch, S. J.; Newsom, D.; White, P.; Navarre, W. W.; Ibba, M.
EF-P Dependent Pauses Integrate Proximal and Distal Signals during Translation.
PLoS Genet **2014**, *10* (8). https://doi.org/10.1371/journal.pgen.1004553.
(41) Aledo, J. C. Renz: An R Package for the Analysis of Enzyme Kinetic Data. *BMC
Bioinformatics* **2022**, *23* (1). https://doi.org/10.1186/s12859-022-04729-4.
(42) Pavlov, M. Y.; Ehrenberg, M. Rate of Translation of Natural MRNAs in an Optimized in
Vitro System. *Arch Biochem Biophys* **1996**, *328* (1).
https://doi.org/10.1006/abbi.1996.0136.
(43) Sharma, A. K.; Sormanni, P.; Ahmed, N.; Ciryam, P.; Friedrich, U. A.; Kramer, G.;
O'Brien, E. P. A Chemical Kinetic Basis for Measuring Translation Initiation and
Elongation Rates from Ribosome Profiling Data. *PLoS Comput Biol* **2019**, *15* (5).
https://doi.org/10.1371/journal.pcbi.1007070.
(44) Correction to: The Impact of Ribosomal Interference, Codon Usage, and Exit Tunnel
Interactions on Translation Elongation Rate Variation (PLOS Genetics, (2018), 14, 1,

(E1007166), 10.1371/Journal.Pgen.1007166). *PLoS Genetics.* 2018.

https://doi.org/10.1371/journal.pgen.1007620.

(45) Pedersen, S. Escherichia Coli Ribosomes Translate in Vivo with Variable Rate.

EMBO J **1984**, *3* (12). https://doi.org/10.1002/j.1460-2075.1984.tb02227.x.

(46) Huter, P.; Arenz, S.; Bock, L. V.; Graf, M.; Frister, J. O.; Heuer, A.; Peil, L.; Starosta, A.

L.; Wohlgemuth, I.; Peske, F.; Nováček, J.; Berninghausen, O.; Grubmüller, H.;

Tenson, T.; Beckmann, R.; Rodnina, M. V.; Vaiana, A. C.; Wilson, D. N. Structural

Basis for Polyproline-Mediated Ribosome Stalling and Rescue by the Translation

Elongation Factor EF-P. *Mol Cell* **2017**, *68* (3).

https://doi.org/10.1016/j.molcel.2017.10.014.

(47) Wilkins, M. R.; Gasteiger, E.; Bairoch, A.; Sanchez, J. C.; Williams, K. L.; Appel, R. D.;

Hochstrasser, D. F. Protein Identification and Analysis Tools in the ExPASy Server.

Methods in molecular biology (Clifton, N.J.). 1999.

https://doi.org/10.1385/1-59259-584-7:531.

Graphical abstract

Figure 1

Figure 2

Figure 3

A

B

Figure 4