Title: Disordered proteins interact with the chemical environment to tune their protective function during drying

Authors: Shraddha KC[1], Kenny Nguyen[1], Vincent Nicholson[1], Annie Walgren[1], Tony Trent[1], Edith Gollub[2], Sofia Romero[2], Alex S. Holehouse[3,4], Shahar Sukenik[2,*], Thomas C. Boothby[1,*]

Affiliations

1. University of Wyoming, Department of Molecular Biology. Laramie, WY

2. Department of Chemistry and Biochemistry, University of California Merced, Merced, CA, USA

3. Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, St. Louis, MO, USA

4. Center for Biomolecular Condensates, Washington University in St. Louis, St. Louis, MO, USA

Correspondence

*Thomas Boothby: tboothby@uwyo.edu

*Shahar Sukenik: ssukenik@ucmerced.edu

Abstract

The conformational ensemble and function of intrinsically disordered proteins (IDPs) are sensitive to their solution environment. The inherent malleability of disordered proteins combined with the exposure of their residues accounts for this sensitivity. One context in which IDPs play important roles that is concomitant with massive changes to the intracellular environment is during desiccation (extreme drying). The ability of organisms to survive desiccation has long been linked to the accumulation of high levels of cosolutes such as trehalose or sucrose as well as the enrichment of IDPs, such as late embryogenesis abundant (LEA) proteins or cytoplasmic abundant heat soluble (CAHS) proteins. Despite knowing that IDPs play important roles and are co-enriched alongside endogenous, species-specific cosolutes during desiccation, little is known mechanistically about how IDP-cosolute interactions influence desiccation tolerance. Here, we test the notion that the protective function of desiccation-related IDPs is enhanced through conformational changes induced by endogenous cosolutes. We find that desiccation-related IDPs derived from four different organisms spanning two LEA protein families and the CAHS protein family, synergize best with endogenous cosolutes during drying to promote desiccation protection. Yet the structural parameters of protective IDPs do not correlate with synergy for either CAHS or LEA proteins. We further demonstrate that for CAHS, but not LEA proteins, synergy is related to self-assembly and the formation of a gel. Our results demonstrate that functional synergy between IDPs and endogenous cosolutes is a convergent desiccation protection strategy seen among different IDP families and organisms, yet, the mechanisms underlying this synergy differ between IDP families.

Keywords: Intrinsically disordered proteins, cosolutes, desiccation tolerance, synergy, CAHS proteins, LEA proteins, oligomerization, sucrose, trehalose

Introduction

Intrinsically disordered proteins (IDPs) make up about 40% of the eukaryotic proteome [1,2]. Unlike typical well-folded proteins, IDPs are characterized by a lack of defined tertiary structure, and instead exist as an ensemble of dynamic, interconverting conformations [3,4]. Despite their disordered nature, IDPs are known to play important roles in many biological processes including regulation of transcription and translation, metabolic signaling, subcellular organization, molecular chaperoning and response and adaptation to environmental cues [3,5].

Despite lacking a stable three-dimensional structure, IDPs still follow a similar paradigm by which form begets function. Different from well-folded proteins, however, an IDP's sequence determines the ensemble of conformations it adopts, and this ensemble can be important for the IDP's function(s) [2,4,6]. However, sequence is not the only determinant of the conformations present in an IDP's ensemble [7–9]. This is because IDP ensembles have relatively few intramolecular bonds and a large solvent-accessible surface area, which makes their ensembles more sensitive to the physicochemical environment than the relatively rigid structures of well-folded proteins [8–10].

The sensitivity of IDP ensembles to their solution environment and their link to IDP function poses a fundamental question: how do sequence and solution combine to tune IDP ensemble and function? To explore this question, we focus on a biological phenomenon where the intracellular environment undergoes drastic physical chemical changes: desiccation.

Organisms across every biological kingdom can survive near-complete desiccation by entering a state of reversible suspended metabolism known as anhydrobiosis (from Greek for 'life without water') [11,12]. As water effluxes from the cell during drying, the concentration of cosolutes increases by orders of magnitude, dramatically changing the physicochemistry of the cell [13,14]. In addition to the decrease in water content and concomitant increase in cosolute concentrations, the composition of the intracellular environment changes massively because of a regulated metabolomic response to drying mounted by anhydrobiotic organisms [12,15].

The acquisition of desiccation tolerance has historically been linked to the intracellular buildup of cosolutes such as trehalose, sucrose, arabinose, stachyose, and raffinose in plants and trehalose in some animals, fungi, and bacteria [15–20]. More recently, the accumulation of high levels of IDPs has also been linked to desiccation tolerance in many organisms [12,21–24]. Common examples of desiccation-related IDPs include the late embryogenesis abundant (LEA) proteins, which are the most widely studied desiccation-related IDPs due to their early identification and widespread distribution among different species and kingdoms of life [21,23–25]. LEA proteins are classified into seven different families based on the presence of conserved motif sequences [25,26]. Another family of desiccation-related IDPs are the tardigrade-specific cytosolic abundant heat soluble (CAHS) proteins [12,22,27].

Simultaneous enrichment of disordered proteins and endogenous cosolutes during desiccation promotes an ideal setting in which to study IDP-cosolute interactions [19,28–30]. In the

desiccation field, these interactions have been observed to produce a functional synergy in promoting tolerance to drying. Trehalose, a cosolute enriched alongside IDPs in many desiccation tolerant systems, has previously been shown to enhance IDP protective function *in vitro* and *in vivo* [19,28–30]. These observations prompted us to ask about the specificity of these interactions and if desiccation-related IDPs may have coevolved to work synergistically alongside their endogenously enriched cosolutes to promote desiccation protection.

To test whether desiccation-related IDP sequences have evolved to work with their intracellular chemical environment, here we use representative proteins from three families of IDPs: one CAHS protein (CAHS D) and proteins from two LEA families. Our result shows that full-length CAHS D and LEA proteins derived from four different organisms synergize better with endogenous protective cosolutes compared to protective exogenous cosolutes from other organisms.

To reveal the underpinnings of cosolute:IDP synergy, we examine the secondary and tertiary structure of protective IDPs in the presence of two disaccharides that are similar in terms of size but distinct with respect to chemistry and use across taxa. In all cases, the secondary structure (residual helicity) and tertiary structure (radius of gyration) do not change significantly in the presence of synergistic cosolutes and thus cannot explain the enhancement in function observed with synergistic cosolutes. We next assessed quaternary structure as both CAHS and LEA proteins are known to oligomerize [23,31], and for CAHS proteins, oligomerization leads to gelation [31–35]. While synergistic cosolutes did not influence LEA oligomerization, CAHS D oligomerization and gelation were enhanced in the presence of synergistic cosolutes. We further show that CAHS D's synergy can be explained through direct repulsive interactions between cosolutes and CAHS D's sidechains. However, this explanation does not hold for LEA proteins, implying that synergy in different protein families occurs through distinct mechanisms.

Our study showcases that different families of protective IDPs can have orthogonal modes of action and different functions in divergent solution environments. Beyond expanding our understanding of desiccation tolerance, these findings shed light on the sensitivity of IDP ensemble and function to the chemical composition of their environment. This is important as IDPs are ubiquitous across biology and function in key developmental processes and disease states that are concomitant with large changes in intracellular chemistry. Understanding how disordered proteins interact and evolve with the solution environment will provide insights into these biological mechanisms and phenomena.

Results

Desiccation-related IDPs are enriched in organisms alongside specific cosolutes during drying

To test whether IDP sequences have evolved to be functionally-tuned by the composition of the intracellular environment during drying, we selected six desiccation-related IDPs. These IDPs come from two LEA families [25,36] as well as the CAHS family (Table 1). We selected four

LEA_4 proteins each from a different desiccation tolerant organism. These organisms include
the plant *Arabidopsis thaliana* (AtLEA3-3)*,* the nematode *Aphelenchus avenae* (AavLEA1), the
tardigrade *Hypsibius exemplaris* (HeLEA68614), and the rotifer *Adineta vaga* (AvLEA1C) (Table
1). To assess whether synergy extends across LEA families, we selected a LEA_1 protein from
A. thaliana (AtLEA4-2) (Table 1). Finally, CAHS D was selected from the tardigrade *H.*
exemplaris (Table 1). These organisms were selected not only because they all utilize LEA
proteins to survive desiccation, but also because they accumulate different disaccharides to
varying degrees during drying [16,17,19,37–41] (Table 1). The organisms we chose use one or
both of two disaccharides - trehalose and sucrose which are similar in size, but chemically
distinct.
161
162 **LEA motifs are not sufficient to mediate synergistic interaction with endogenous**
163 **cosolutes during desiccation**
164
To assess whether endogenous cosolutes induce functional changes in desiccation-related
IDPs, we began by testing peptides encoding LEA motifs derived from full-length LEA_4 and
LEA_1 proteins (Table 1). Family 1 LEA (LEA_1) proteins are characterized by a 20-mer
repeating motif, whereas Family 4 LEA (LEA_4) proteins are characterized by the repetition of
an 11-mer LEA_4 motif [25,36,42–44]. These motifs are often found in multiple linear or
nonlinear repeats across the length of a LEA protein [25,36,42]. Interestingly, LEA_4 motifs
have previously been suggested to be sufficient to confer desiccation protection to desiccation-
sensitive proteins and membranes to a degree similar to full-length LEA proteins, both *in vitro*
and *in vivo [44–47]*. With this in mind, we expected to observe synergy between LEA motif
repeats and their paired endogenous cosolute(s).
175
176 We generated 11-mer LEA_4 motif peptides (At11, Aav11, He11, and Av11) and a 20-mer
177 LEA_1 peptide (At20) and measured the ability of these motifs to protect lactate dehydrogenase
178 (LDH), a desiccation sensitive enzyme during drying. LDH assay is used to assess the function
179 of desiccation protectants to protect the activity of LDH which otherwise retains only
180 approximately 2% of its pre-desiccation activity when dried and rehydrated [19,22,28,48].
181
The protective capacity for each LEA motif and cosolute was assessed across a range of
concentrations with LDH (Fig. 1A). Most LEA_4 motifs displayed levels of protection so low that
a 50% level of protection could not be reached even at concentrations exceeding 1 mM (Fig.
1A&B, Fig. S1A). Additionally, higher concentrations of the LEA_4 motifs tend to inactivate the
enzyme when kept under control conditions (4 °C, see Methods), over a 16 hour incubation
period during the assay (Fig S1C). The LEA_1 20-mer motif At20, however, showed robust
concentration-dependent protection of LDH, demonstrating that LEA_4 and LEA_1 motifs are
functionally distinct (Fig. 1A&B, Fig. S1A). Concentration-dependent protection was also
observed for our cosolutes trehalose and sucrose (Fig. 1A, Fig. S1A).
191
Since LEA_4 motifs often exist in tandem repeats of 11-mers within a full-length LEA protein
[44,45,49], we wondered if the observed lack of protection was a result of their short length or
repeat number. We synthesized 2X (At22) and 4X (At44) tandem repeats of the *A. thaliana* 11-

mer LEA_4 motif (At11). Our results show minimal potency in preserving *in vitro* LDH function
during drying regardless of motif length (Fig. 1A, Fig. S1A).

Despite the low protection displayed by our LEA_4 peptides, we opted to use them in cosolute
synergy assays, as we reasoned that perhaps they would become functional when in solution
with trehalose or sucrose. We picked sub-optimal concentrations of protectants (Table S1) so
that under instances of synergistic protection, the additive protection of cosolute:peptide
mixtures would not exceed 100%. We then performed synergy assays where sucrose or
trehalose was combined with LEA motifs at molar ratios of 1:100, 1:10, 1:1, 10:1, and 100:1
(cosolute:protein). The upper limit (10:1 and 100:1) of these ratios closely align with known
cosolute:protein ratios that produce synergistic protection between the tardigrade disordered
protein CAHS D and trehalose [19]. Here we report on synergy by showing the individual
protective ability of the cosolute and IDP on its own, the sum of these protective values
(hypothetical additive effect), and the actual measured protection produced by combination of
the cosolute and peptide (Fig. 1C). We quantify synergy using the following equation:

% synergy = (% LDH protection)$_{IDP+cosolute}$ - (% LDH protection)$_{IDP}$ - (% LDH protection)$_{cosolute}$

In nearly all cases, synergy was not observed for 11-mer LEA_4 peptides with either sucrose or
trehalose (Fig. 1D&E, Fig.S1B). In fact, in several cases, mixing LEA_4 peptides with sucrose or
trehalose elicited antagonistic, rather than synergistic, effects (Fig. 1D&E, Fig. S1B). These
results suggest that LEA_4 motifs do not robustly preserve LDH function, nor do they interact
with cosolutes trehalose or sucrose in a functionally productive fashion. Similar to 11-mer
motifs, the *A. thaliana* 22- and 44-mer peptides did not synergize with either trehalose or
sucrose (Fig. 1D&E, Fig. S1B). Unlike LEA_4 motifs, the 20-mer LEA_1 motif, At20, was both
protective and synergized with both trehalose and sucrose (Fig. 1D&E, Fig. S1B).

Taken together, these experiments demonstrate a diversity in disordered protein/motif function,
where LEA_4 motifs largely are not protective to the enzyme LDH during drying, nor are they
synergistic with endogenous cosolutes. Conversely, the LEA_1 motif tested is highly protective
and synergizes with either sucrose or trehalose.

Desiccation-related IDPs synergize with endogenous cosolutes

While LEA proteins are identified through homology in conserved LEA motif repeats, they also
contain varying quantities of non-motif sequence (Table 1). Since we observed that LEA_4
motifs generally provide relatively little protection and tend not to synergize with endogenous
cosolutes in LDH assays, we wondered if full-length LEA proteins might. We also included
CAHS D in our analysis as it has been previously known to synergize with trehalose and, to a
lesser extent, with sucrose [19].

We began by testing the baseline protection of our proteins using the LDH assay. All full-length
LEA_1 and LEA_4 proteins confer protection for LDH activity up to the pre-desiccated value
(Fig. 2A, Fig. S2A). Likewise, CAHS D provided concentration-dependent protection to LDH as

previously observed [19,22,48] (Fig. 2A, Fig. S2A). We also included bovine serum albumin
(BSA) in these studies as a well-studied control [19,48]. Unlike the LEA motifs, most full-length
proteins protected 50% LDH at concentrations less than 1 mM (Fig. 2B).

Using data derived from the concentration range of LDH assays, we chose a sub-optimal
concentration that provides 15-45% protection for each protein to perform synergy experiments
with (Table S2). Our results show that nearly all full-length IDPs showed synergy with either
sucrose or trehalose or both (Fig. 2C&D). Exceptions to this are AvLEA1C, which is derived
from a rotifer that accumulates neither trehalose nor sucrose, and BSA, which comes from
cows, which of course have no capacity for anhydrobiosis. Remarkably, in cases where LEA
proteins displayed synergy, they were always more synergistic with endogenous compared to
exogenous cosolutes (Fig. 2E).

Taken together, these experiments demonstrate that synergistic interactions between IDPs and
cosolutes extend across multiple families of desiccation-protective IDPs found in a variety of
organisms. Furthermore, proteins derived from different organisms synergize best with their
endogenous cosolute(s) to promote desiccation tolerance. It is also of note that while many of
the LEA motifs tested at the beginning of this study did not display synergy with trehalose or
sucrose, corresponding full-length proteins did. Likewise, At20 showed synergy with both
trehalose and sucrose, while full-length AtLEA4-2 was only synergistic with sucrose. Thus, not
only do these experiments demonstrate that full-length LEA proteins synergize with their
endogenous cosolute(s) (Fig. 2E), but they demonstrate that this synergy, at least in part, is
driven by sequence features beyond conserved motifs.

Trehalose and sucrose do not elicit local ensemble changes to desiccation-related IDPs
in solution or in the dry state under the conditions tested

We next wondered what mechanism(s) drive the functional synergy observed between
desiccation-protective IDPs and endogenous cosolutes. We reasoned that functional synergy
might be driven by cosolute-induced changes to the IDP ensemble. To test this, we first
examined the secondary structure contained in the ensemble of LEA proteins and CAHS D
using circular dichroism (CD) spectroscopy.

Each full-length protein was first assessed using CD in an aqueous state by itself (Fig. 3A). All
full-length LEA proteins displayed a single minimum at ~200 nm (Fig. 3A, black) indicating that
LEA proteins are disordered in the aqueous state. CAHS D also displayed a minimum at ~200
nm and a slight minimum around ~220 nm, indicating that while disordered, it also has some
propensity for helical structure in solution (Fig. 3A, black), in line with previous studies [31,50].
This is in contrast to BSA, which showed a high propensity for helical structure as denoted by
the double minima at 222 and 210 nm (Fig. 3A, black). To see if the addition of cosolutes
induces secondary structural change in the aqueous state, we obtained CD spectra of
cosolute:protein mixtures at 100:1 molar ratios. Somewhat to our surprise, trehalose (Fig.
3A&C, blue) and sucrose (Fig. 3A&C, green) do not induce any significant structural changes to
any of the full-length IDPs tested here (Fig. S3A).

LEA proteins gain helical conformation upon drying or in response to low water availability.
Drying-induced helicity has been postulated to drive their protective function [23,30,51]. We
reasoned that while our cosolutes do not induce detectable changes to LEA/CAHS secondary
structure in solution, cosolutes could induce structural changes in proteins in a dry state. To test
this, we examined our proteins using CD in a desiccated state [52,53]. Our results show a
significant structural change for all LEA proteins and CAHS D in the dry state, indicated by a
shift from disordered spectra with a minimum at ~200 nm to a helical structure with two minima
at ~222 and ~210 nm (Fig. 3B). This is in contrast to BSA which started out helical and showed
little change in the spectrum (Fig. 3B). These changes manifest for pure proteins without any
addition of synergistic cosolutes. To quantify the influence of drying on secondary structure, we
examined the changes in the ratio of CD signal at 222 and 210 nm. This ratiometric value
reports on secondary structure in a concentration-independent way [54]. Using this metric, all
LEAs and CAHS D display a significant change in structure going from the aqueous to
dehydrated state, whereas BSA remains the same (Fig. 3C).

Next, we examined combinations of our proteins with trehalose or sucrose in a desiccated state.
As with aqueous samples, the addition of trehalose or sucrose did not induce significant
changes in the secondary structure of LEA and CAHS D proteins in the dry state (Fig. 3B&C,
Fig. S3B). To assess whether there is a link between the minimal structural changes we
observed and functional synergy in LDH assays, the change in the ratio of signal at 210 and 222
nm in desiccated and aqueous states was compared to synergy observed for that same mixture.
Synergistic protection observed in our LDH assays did not correlate with secondary structural
changes with the addition of trehalose (p=0.9863 for aqueous, p=0.1113 for desiccated, Fig.
S3C&D) or sucrose (p= 0.6673 for aqueous, p=0.9863 for desiccated, Fig. S3E&F).

Taken together, these results indicate that while LEA proteins and CAHS D undergo a structural
transition during desiccation, this phenomenon does not require, nor is it affected by, the
presence of trehalose and sucrose. Furthermore, synergistic protection observed in our LDH
assays is not mediated by local ensemble changes in these IDPs.

Trehalose and sucrose do not elicit changes in global ensemble dimensions for
desiccation-related IDPs, but promote oligomerization of CAHS D

We next wondered if the synergistic interactions observed between our IDPs and cosolutes
could instead be explained by a change in global dimensions, such as expansion or compaction
of the protein. To measure global dimensions, which cannot be detected using CD, we used
small angle X-ray scattering (SAXS), which allows a model-free estimation of the radius of
gyration (R_g) of an IDP as well as a prediction of the molecular weight [55]. Each protein was
measured with no cosolute and with different molar ratios of trehalose and sucrose. We
reasoned that a cosolute-dependent change in the radius of gyration (R_g) could indicate
changes in tertiary or quaternary structure, which may correlate with increased function or
synergy. We note that for SAXS experiments cosolutes were used at concentrations between

20 and 50 mM due to technical restrictions on how much protein can accurately be assayed and
a desire to maintain molar ratios used in other experiments.

We began by testing the R_g of BSA in different solution conditions. As a well-folded protein, we
expected that BSA would be relatively insensitive to changes in the solution environment. The
R_g values obtained via this approach match existing literature (Fig. 4A) [56]. Additionally, adding
cosolutes did not modulate R_g (Fig. 4A, Fig. S4A). While our molecular weight approximations
trended higher than expected for monomeric BSA, we reason that this may be due to the
propensity of BSA to form small populations of low-level oligomers [57,58].

We next measured the R_g of our LEA proteins in various solution conditions. We observed that
regardless of the solution environment, all of our LEA proteins have an R_g that falls within error
of readings in other solution environments (Fig. 4B-F). At the concentrations used here,
cosolutes do not induce significant changes in the global dimensions of LEA proteins. While the
predicted molecular weight (pMW) from SAXS for these proteins were somewhat variable, we
see no consistent trend between the presence of cosolutes and change in pMW for any LEA
protein (Fig. 4B-F).

Finally, we tested CAHS D in similar solution conditions. We obtained an R_g value for CAHS D
that lies between the values reported by other groups (see Methods) [31,33]. While this R_g was
consistent in 1.6:1 disaccharide solutions, 16:1 disaccharide solutions, and 160:1 sucrose
solution, we found that the 160:1 trehalose solution had a Guinier region that was sharply
curved upwards, even upon the protein's first exposure to the X-ray source (Fig. 4G, Fig. S4G).
This is consistent with the presence of large oligomerized species. A Bayesian approximation of
the molecular weight of these samples supported this result (Fig. 4G). Most of our CAHS D
samples showed a pMW that was only slightly elevated from the known value for the monomeric
protein. In contrast, the 160:1 sucrose sample had a pMW about 300% higher than other
samples, and the 160:1 trehalose sample had a pMW about 1000% higher (Fig. 4G).

While SAXS shows little change in Rg or pMW for LEA proteins, there is evidence that not only
CAHS D but also some LEA proteins tend to oligomerize [23,59]. However, LEA oligomerization
appears to be weak and transient, requiring extreme crowding and/or sensitive methods to
detect [23,59,60]. We therefore wanted to use a more sensitive method to assess LEA
oligomers, and how they might be affected by cosolutes.

To characterize oligomerization of LEA proteins in a more sensitive fashion, we used photo-
induced crosslinking of unmodified proteins (PICUP), a zero-length crosslinking method that
uses a light activatable crosslinking system and is known to capture transient oligomeric species
[61,62]. PICUP has previously been used to characterize oligomeric forms in LEA proteins *in*
vitro [60,63]. All of our LEA proteins showed a propensity to form oligomers, even at low
concentrations (Fig. S5). However, the presence of trehalose or sucrose did not elicit changes
in oligomeric populations (Fig. S5). These results confirm that LEA proteins are able to form
transient oligomers, but also demonstrate that at levels where sucrose and trehalose are
synergistic with these proteins oligomerization is unaffected.

Taken together, these results show a divergence in the behavior of LEA and CAHS proteins in
the presence of synergistic cosolutes. For CAHS D, the SAXS data suggests a relationship
between the presence of cosolutes and increased oligomerization. However, this dataset
showed no evidence of a cosolute inducible increase in molecular weight or R_g for LEA proteins.
This was further supported by PICUP, which despite detecting LEA oligomers did not show that
they were enhanced by cosolutes. Thus, our data suggests that CAHS D oligomerization is
promoted by the presence of synergistic cosolutes while LEA oligomerization is not.
378

379 **Synergistic cosolutes promote gelation of CAHS D but not LEA proteins**
380

381 CAHS D oligomers, as well as those of other CAHS proteins, have been reported to undergo
382 self-assembly to form a gel network [31–35,64]. We wondered if the cosolute induced
383 oligomerization of CAHS proteins in the presence of trehalose as seen in the SAXS experiments
384 could be attributed to the propensity of CAHS D to form gels.
385

To test this, we performed differential scanning calorimetry (DSC) on CAHS D to observe the
presence or absence of a gel melt. To begin, we tested CAHS D at 6 mg/mL (0.235 mM), which
has previously been established to be a non-gelling concentration [31]. Consistent with this, at
0.235 mM, we find that CAHS D does not undergo a characteristic gel melt indicating a lack of
gelation (Fig. 5A, black). Addition of trehalose and sucrose at increasing molar ratios (1:1, 10:1,
100:1, and 500:1) showed thermal features characteristic of endothermic phase transitions (e.g.,
a gel melting), indicating that the presence of these cosolutes induced gelation (Fig. 5A).
Measuring the area under these melt curves allows us to calculate the enthalpy of melting (Fig.
S6A). Change in enthalpy measurements for the cosolute:protein mixtures relative to the protein
provide us a quantification of how gelation is affected by different amounts of cosolutes.
Trehalose induced significant gelation at a 100:1 ratio, while 500:1 of sucrose was required to
induce a significant gel melt (Fig. 5B). This is consistent with trehalose producing larger
oligomeric species in our SAXS experiments (Fig. 4G), indicating that trehalose has a larger
influence than sucrose on the gelation of CAHS D (Fig. 5B).
400

Furthermore, to test whether synergistic cosolutes enhance gelation, we tested CAHS D at 12
mg/mL (0.47 mM), a concentration above CAHS D's gelation threshold [31]. Addition of
trehalose or sucrose at increasing molar concentrations promoted the formation of stronger gels
evident by the enthalpy of melting measurements (Fig. S6B-E). These experiments demonstrate
that not only do trehalose and sucrose induce subgelling concentrations of CAHS D to form
gels, but they also enhance the strength of gels formed by higher concentrations of the protein.
407

Unlike CAHS proteins, gelation of LEA proteins has not been commonly observed or reported.
The exception to this is AfrLEA6 (a LEA protein from the brine shrimp *A. franciscana*) that
appears to undergo phase separation, forming a hydrogel-like matrix upon desiccation [65]. To
see if our LEA proteins gel, we performed DSC experiments on our LEA proteins and cosolutes.
We analyzed similar molar concentrations of LEA proteins on their own and in mixtures with
cosolutes at equivalent ratios (1:1, 10:1, 100:1, and 500:1). None of our LEA proteins by

themselves or in mixtures with trehalose or sucrose show evidence of gelation (Fig. 5B-G).
Likewise, BSA also failed to form a gel (Fig. 5I).
416
417 Taken together, these results demonstrate that trehalose and sucrose affect the oligomerization
418 and phase state of different IDPs in distinct ways. The observation that trehalose induces more
419 synergy as well as more gelation of CAHS D relative to sucrose leads us to speculate that
420 gelation and synergistic protection of LDH may be linked.
421
422 **Direct cosolute:IDP interactions drive synergy for CAHS D, but not LEA proteins**
423
To explain the possible relationship between synergy and oligomerization-driven gelation of
CAHS D, we quantified the interactions between each IDP and its cosolute environment using
transfer free energies (TFEs). The TFE is a measure of the change in free energy undergone by
a macromolecule when transferring from water to a concentrated solution of some osmolyte
(typically 1 M) [66–68]. Using transfer free energy values derived from literature, we calculated
the effect of trehalose and sucrose on the ability of CAHS D to dimerize: $\Delta\Delta G_{tr}^{M\to D}$. This is
calculated by finding the free energy of CAHS D's monomeric state upon transfer to an
osmolyte solution (ΔG_{tr}^{M}), doing the same for the dimeric state (ΔG_{tr}^{D}), and then taking the
difference (Fig. 6A). A strong negative value for $\Delta\Delta G_{tr}^{M\to D}$ indicates that the presence of the
cosolute pushes the population towards dimers. A positive value indicates that the addition of
the cosolute pushes the population towards monomers.
435
In order to perform these calculations, we utilized AlphaFold2 and AlphaFold Multimer to
determine plausible conformations for both CAHS D's monomeric and dimeric state. These
structures have a helical linker region, which is consistent with our CD measurements (Fig.
3A&B) and is consistent with previous reports for this protein [31]. We reasoned that
dimerization is indicative of gelation since CAHS D dimers were especially prevalent in
crosslinking data [31], and previous research suggests that CAHS D dimer formation is a
necessary step toward gelation [31]. TFEs were then calculated based on the solvent
accessible surface area of different chemical groups in monomeric vs. dimeric state (see
methods). Our calculations reveal that trehalose has a negative $\Delta\Delta G_{tr}^{M\to D}$ with CAHS D,
meaning the dimeric state is favored in the presence of this cosolute. Sucrose's $\Delta\Delta G_{tr}^{M\to D}$ is
close to 0, neither stabilizing nor destabilizing the dimer (Fig. 6B). In addition to these cosolutes,
we wanted to explore the effect of a cosolute with a positive $\Delta\Delta G_{tr}^{M\to D}$. Glycine betaine, a
common stabilizing cosolute, but without known roles in tardigrade desiccation tolerance,
displayed a positive $\Delta\Delta G_{tr}^{M\to D}$ (Fig. 6B). Together these cosolutes span a range of
$\Delta\Delta G_{tr}^{M\to D}$ values that are expected to increase the dimer population (trehalose), have a minimal
impact on dimerization (sucrose), or increase the monomer population (glycine betaine) of
CAHS D [68,69]. While data for trehalose and sucrose are in line with this analysis, we next
sought to determine empirically the impact of glycine betaine on CAHS D gelation.
454
To test the effects of glycine betaine on CAHS D dimerization predicted by these TFE
calculations, we first repeated our DSC and SAXS experiments in the presence of glycine
betaine. Unlike trehalose and sucrose, below the protein's gelation threshold no increase in

enthalpy of melting was observed upon the addition of glycine betaine (Fig. 6C&D). To probe
whether glycine betaine inhibits CAHS D oligomerization, we conducted additional DSC
experiments above the protein's gelation threshold. While trehalose and sucrose enhanced
gelation of CAHS D (Fig S6B-E), we observed a decrease in enthalpy of melting when glycine
betaine was present at the 500:1 molar ratio, signifying inhibition of CAHS D gelation (Fig.
6E&F).

We then performed SAXS on CAHS D with 1000:1 molar ratios of each cosolute. The intent of
using such a high molar ratio was to mimic the high concentration of cosolutes that CAHS D
would experience during desiccation [13]. A non-gelling concentration of CAHS D in 1000:1
glycine betaine yielded a scattering profile similar to the protein with no cosolute indicating a
lack of gelation (Fig. 6G). Meanwhile, 1000:1 sucrose and trehalose yielded scattering profiles
consistent with gelation, similar to those previously reported [31]. This is made especially
evident by a peak at q=0.06 Å^{-1}, which reports on the width of CAHS D's gel fibers and matches
previously reported scattering profiles for gelled CAHS D [31]. Consistent with our hypothesis
that the more negative $\Delta\Delta G_{tr}^{M \to D}$ for trehalose will increase dimerization and subsequent
gelation, we observed an increased curvature in the Guinier region indicating increased
fibrillization (Fig. S7C&D).

Finally, we tested the impact of glycine betaine on CAHS D's protective capacity. If induction of
self-assembly of CAHS D is a mechanism underlying trehalose/sucrose induced synergy, then
one would expect that glycine betaine's inhibition of gelation would result in no synergy, or even
have an antagonistic effect. We found a significant antagonistic relationship between glycine
betaine and CAHS D on LDH protection (Fig. 6H). Furthermore, Pearson correlation between
the $\Delta\Delta G_{tr}^{M \to D}$ of a cosolute (scaled by concentration) and its ability to induce synergy in CAHS D
was statistically significant (Fig. 6I). This correlation however did not hold for LEA proteins or for
BSA (Fig. S7E-J).

Taken together, these results suggest that the driving force for gelation in CAHS D and its ability
to synergize with a given cosolute is rooted in the direct interaction between CAHS D and the
prevalent cosolute. This observation ruled out the possibility for the combined interactions of the
CAHS D:cosolute on the client protein, LDH. However, trying to apply this model to the LEAs
tested in this work failed to yield meaningful correlates. Thus, we propose that direct interactions
between cosolute and LEA proteins cannot explain the synergy observed with cosolutes.
Overall, our study demonstrates that while synergy between desiccation-related IDPs and
endogenous cosolutes appears to be a widespread and conserved behavior, the mechanisms
underlying this synergy vary between IDP families.

Discussion

499 In this study, we examined the interplay between IDP sequence, solution environment,
500 ensemble, and function. To do this, we have taken advantage of the dramatic changes to the
501 cosolute content of anhydrobiotic organisms brought on by desiccation, and compared the

effects of these cosolutes between three families of desiccation-related IDPs (LEA_4, LEA_1, and CAHS proteins). We demonstrate that endogenous cosolutes enriched during desiccation, enhance the protective capacity of CAHS D, full-length LEA_4, and LEA_1 proteins in an *in vitro* enzyme assay. Surprisingly, the functional changes were not accompanied by any detectable structural changes to the monomeric ensemble of these IDPs. However, synergistic cosolutes did induce oligomerization and gelation of CAHS D. Finally, we show that in the case of CAHS D, but not LEAs, oligomerization, gelation, and protective synergy can be traced to direct interactions between cosolute and protective protein. Our results suggest that while functional synergy between the solution environment spans multiple IDP families, different mechanisms can underlie synergistic interactions for different proteins.

Functional synergy for the full-length proteins from different organisms mirrored the endogenous cosolute environment in that organism. In all cases, an IDP protected LDH activity more with its endogenous cosolute compared to an exogenous cosolute. These differences in synergy appear to extend across even subtle variations in cosolute use. For example, nematodes and tardigrades both accumulate trehalose during desiccation, but tardigrades accumulate orders of magnitude less [19,37–39]. Consistent with this, both tardigrade proteins used in this study synergized with trehalose at an order of magnitude lower concentration than what was required to elicit synergy with the nematode LEA protein.

Overall, while our study found that synergy with endogenous cosolutes is observed across two families of LEA proteins as well as CAHS proteins, we observed a major difference in structural changes induced in these protein families. While in CAHS D synergy could be traced back to the interaction between the protective protein and the cosolute, no underlying mechanism was detected for LEAs. What then could be driving the synergy we observe for these proteins?

One possibility is that in most of our assays we do not consider the protein being protected. Our studies of molecular mechanisms for synergy do not consider the underlying effect of both protectant protein and cosolute on LDH directly. It is possible that the presence of both endogenous cosolute and protein create a solvation environment that becomes highly protective for e.g. rehydration. In line with this, recent studies have highlighted the ability of LEA proteins to stabilize sugar glasses in a dry state [49,70–72]. Glass formation is known to preserve labile biomolecules during desiccation, contributing to survival [22,73]. Different glasses vary significantly in their protective capacity, and studies have attempted to find structural properties that explain this difference [72,74]. Because trehalose and sucrose both form glasses when dried [75,76], it is possible that our LEA proteins are inducing a change in the glass's structural properties that leads to synergy.

Another possibility is a difference in the nature of TFE-induced oligomerization. While repulsive cosolutes drive homotypic interactions between CAHS D monomers, the same thermodynamic force can promote heterotypic interactions between LEAs and other proteins [77]. For example, trehalose may stabilize electrostatic interactions between LEA proteins and LDH during our *in vitro* synergy assays. If the protective capacity of LEA proteins is dependent on direct interactions between the protectant and the client protein, then this is a plausible explanation for synergy. However, it is currently unknown whether or not this is the case.

Another major question posed by this research is why sucrose was able to elicit synergy in CAHS D. Our computational approach predicted that sucrose should be agnostic to CAHS D's ability to form dimers, and yet we clearly see that, *in vitro*, sucrose is a moderately potent driver of gelation. We believe several factors could explain this effect. While sucrose does not drive dimerization through direct 'soft' repulsion, it may still do so by acting as a crowder eliciting an excluded volume effect [78]. Another possible manifestation of an excluded volume effect is the slight increase in the melting peak that is observed with all cosolutes used in this study at high molar ratios (Fig. S6F-I, Fig. S7K-L). Alternatively, given that CAHS D must polymerize beyond the dimeric state to form a gel, sucrose may stabilize a higher-level oligomer that was not captured in our $\Delta\Delta G_{tr}^{M\rightarrow D}$ analysis.

IDPs are known to play important regulatory functions during development and disease progression that often occurs together with changes to the chemical composition of the intracellular environment. For example, there are known links between type II diabetes and Alzhiemer's disease, and it has been shown that the intrinsically disordered neurodegenerative peptide Aβ42 undergoes pathological oligomerization in the presence of glucose whose levels mirror those found in diabetic patients [5]. Our study showcases how different cosolute environments can have a direct effect on the function of IDPs. By understanding the rules governing desiccation related IDP-cosolute interactions, we might better understand the influence of changing chemical environments on a host of other IDPs.

Acknowledgements

Support for this project came from NSF via the IntBio research program under awards 2128069 to TCB, 2128067 to SS, and 2128068 to ASH. SK and KN were supported in part by the USDA National Institute of Food and Agriculture, Hatch project #1012152. In addition, this work was made possible in part through support from an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health (Grant # 2P20GM103432). We thank members of the Water and Life Interface Institute (WALII), supported by NSF DBI grant #2213983, for helpful discussions. We thank Dr. Greg Hura and Kathryn Burnett for their correspondence and help in performing the SAXS experiments. SAXS experiments were conducted at the Advanced Light Source (ALS), operated by Lawrence Berkeley National Laboratory on behalf of the Department of Energy, Office of Basic Energy Sciences, through the Integrated Diffraction Analysis Technologies (IDAT) program, supported by DOE Office of Biological and Environmental Research.

Author Contributions

Conceptualization: **SK, KN, ASH, SS, TCB**
Data Curation: **SK, KN, VN, EG, SR**
Formal Analysis: **SK, KN, VN, EG, SR**
Funding Acquisition: **ASH, SS, TCB**

Investigation: **SK, KN, VN, AW, AT, EG, SR**
Protein expression and purification: **SK, KN, AW**
LDH functional experiments: **SK, KN**
Circular dichroism spectroscopy: **EG, SR**
Small angle X-ray scattering: **VN**
Cross-linking: **SK**
Differential scanning calorimetry/gelation: **SK, KN**, **AW**
TFE computation/analysis: **VN**, **AT**
Methodology: **SK, KN, VN, EG, SS, TCB**
Project Administration: **VN**
Software: **AT**
Supervision: **ASH, SS, TCB**
Visualization: **SK, KN, VN**
Writing – Original Draft: **SK, KN, VN, TCB**
Writing – Review & Editing: **SK, KN, VN, AW, AT, SR, EG, ASH, SS, TCB**
599
600   **Declarations of Interest**
601   A.S.H. is a scientific consultant with Dewpoint Therapeutics and is on the Scientific Advisory
602   Board of Prose Foods. The work reported here was not influenced by these affiliations. All other
603   authors declare no competing interests.
604
605   **Lead Contact**
606   Further information and request for resources and reagents should be directed to and will be
607   fulfilled by TCB (thomas.boothby@uwyo.edu)
608
609   **Data availability**
610   All data generated or analyzed during this study are included in this published article (File
611   S1.zip).
612
613   **Code availability statement**
614   All custom code used in this study is included in this published article (File S2.zip).
615
616
617   **Methods**
618
619   **Protein Sequences**
620   Sequences of all peptides and proteins used in this study are available in File S3.
621
Cloning
Inserts for full-length proteins- AavLEA1, AtLEA4-2, AvLEA1C, CAHS D, and HeLEA68614
were synthesized as codon optimized gBlocks (Integrated DNA Technologies) and cloned into
the pET28b expression vector using gibson assembly (New England Biosciences). AtLEA3-3

was cloned in pET28a vector by Twist Bioscience. Clones were propagated in DH5α cells (NEB,
Cat. #C2987H) and verified by Sanger sequencing (Eton Bioscience).
628

Protein Expression

Expression constructs were transformed into BL21 (DE3) cells (New England Biosciences, Cat.
#C2527H) and plated on Luria-Bertani (LB) agar plates with 50 µg/mL kanamycin. At least 3
single colonies were chosen for each construct and tested for expression. Constructs were
expressed in 1 L LB/kanamycin medium and grown at 37 °C while shaking at 180 rpm
(Eppendorf Innova S44i) until an OD600 of 0.6 was reached. The culture was induced with 1 mM
IPTG, and grown for the next 4 hours while shaking. AvLEA1C was grown for 1 hour following
IPTG addition. Cells were harvested by centrifugation at 4000 rpm for 30 minutes at 4 °C. Cell
pellets were resuspended in 5 mL of 20 mM Tris buffer, pH 7.5 supplemented with 30 µL of 1X
protease inhibitor [Sigma Aldrich, Cat. P2714]). Cell pellets were stored at -80 °C until further
use.
640

Protein Purification

Frozen pellets were thawed at room temperature, subjected to heat lysis in boiling water for 10
minutes, and cooled down for 15 minutes. These were then centrifuged at 10500 rpm at 10 °C
for 30 minutes, and the supernatant was later filter-sterilized through a 0.22 µm filter to remove
any insoluble particles (EZFlow Syringe Filter, Cat. 388-3416-OEM). The filtrate was diluted two
times the volume with buffer UA (8 M urea [Acros Organics, CAS No. 57-13-6], 50 mM sodium
acetate [Tocris CAS No. 127-09-3], pH 4). This was loaded onto a HiPrep SP HP 16/10 (Cytiva,
Cat. 29018183) cation exchange column and purified on an AKTA Pure (Cytiva, Cat. #29018224),
controlled using the UNICORN 7-9.1 Workstation pure-BP-exp (Cytiva, Cat. #29128116). CAHS D
was eluted using a 0-40% UB (8 M urea, 50 mM sodium acetate, and 1 M NaCl, pH 4) gradient
and fractionated over 15 column volumes. LEA proteins were eluted using the 0-70% UB
gradient over 15 column volumes. Protein fractions were assessed using SDS-PAGE and
selected fractions were dialysed in a 3.5 kDa tubing (SpectraPor 3 Dialysis Membrane, Part No.
132724) in 20 mM sodium phosphate buffer pH 7, followed by six rounds of Milli-Q water (18.2
MΩcm) at four hours interval each. Concentration of the dialyzed fractions were then quantified
using Qubit 4 fluorometer (Invitrogen, REF Q33226), flash frozen, then lyophilized (Labconco
FreeZone 6, Cat. 7752021) for 48 hours, and stored at -20 °C until further use.
658

LEA Motif Sequence Identification

660 LEA_4 and LEA_1 sequence motifs were identified in full-length LEA proteins using RADAR
661 (https://www.ebi.ac.uk/Tools/pfa/radar/). In cases where RADAR was unable to identify
662 repetitive motifs (e.g., in cases where a full-length LEA protein had only one or two motif
663 repeats), manual selection and alignment of motifs was performed.
664

Lactate Dehydrogenase (LDH) Protection Assay

LDH assay was adopted from previous studies [19,22,28,48]. Protectants were resuspended at
a final concentration range 20 mg/mL to 0.1 mg/mL in 25 mM Tris HCl pH 7. Rabbit muscle L-
lactate dehydrogenase (LDH), sourced from Sigma (Sigma-Aldrich, Cat 10127230001), was
added to each solution at a concentration of 0.1 mg/mL. Half of this sample was dried in a

vacuum desiccator (SAVANT Speed Vac Concentrator) for 16 hours, while the other half was
refrigerated at 4 °C for the same duration. Water was added to both desiccated and non-
desiccated samples to a final volume of 250 µL each. 10 µL sample was mixed with 980 µL
phosphate pyruvate buffer (100 mM sodium phosphate, 2 mM sodium pyruvate; pH 6.00)
supplemented with 10 µL of 10 mM NADH (Sigma-Aldrich NADH; disodium salt, grade II) in a
quartz cuvette. LDH activity was measured as the kinetics of the decrease in NADH absorption
at 340 nm for a minute in NanodropOne (Thermo Scientific). Percent protection was calculated
as a ratio of NADH absorbance for the desiccated samples normalized to non-desiccated
controls. Each sample was performed in triplicate.
679
Lactate Dehydrogenase (LDH) Synergy Assay
The protection data for individual protein or motif was used to select a suboptimal protective
concentration. Trehalose or sucrose was mixed in equal parts with proteins at 2X concentration
in 100 µL resuspension buffer (25 mM Tris HCl pH 7) at respective molar ratios. LDH assay was
performed for the mixtures as described previously. For each mixture, LDH protection was
assessed individually and as a mixture. The sum of the protection conferred by individual
protein and cosolute was determined, which would refer to the expected additive protection.
Synergy was determined by statistical comparison of this expected additive protection with the
experimental protection.
689
Circular Dichroism (CD) Spectroscopy
CD spectroscopy was adopted from Bremer et al. [52]. Lyophilized proteins were resuspended
in 25 mM NaPi pH 7 to a concentration of 200 µM. The resuspended protein was then mixed in
equal parts with the NaPi buffer, 20 mM trehalose, or 20 mM sucrose in separate samples to a
100:1 molar ratio, with a final protein concentration of 100 µM and cosolute concentration of 10
mM. Protein concentration was confirmed with either a UV-vis (Thermo Scientific, GENESYS 50
UV-visible spectrophotometer) or a Qubit (Life Technologies, Qubit 3.0 Fluorometer). 20 µL
aliquots of the samples were deposited on a 0.05 mm quartz cuvette and measured in a Circular
Dichroism (CD) spectrometer (JASCO, J-1500 model). New 20 µL aliquots were then deposited
on one half of a 0.05 mm quartz cuvette and spread across part of the cuvette with the tip of a
pipette, to a surface area of about 1 cm^2. The samples were then desiccated in a vacuum
chamber with drierite for 1 hour to create a dry film, and another CD measurement was taken
immediately after the vacuum was stopped. Each measurement was performed in triplicate.
703
Small-Angle X-ray Scattering (SAXS) – Sample Preparation
Lyophilized protein was resuspended at high concentration in a buffer containing 20 mM tris HCl
(pH = 7.0) and the correct amount of cosolute to reach the desired molar ratio. Protein samples
were then quantified with the Qubit Protein Assay from ThermoFisher Scientific (catalog#
Q33212). The proteins were then diluted into 8 mg/mL and 4 mg/mL stocks using the same
cosolute solution. Due to the necessity for each sample to have an identical buffer blank, the
concentration of cosolute in the 8 mg/mL sample had to be the same as in the 4 mg/mL sample,
meaning the molar ratio would be doubled in the 4 mg/mL sample. For each sample, a small
aliquot of buffer was saved and stored at 4 °C for use as a blank. Samples and buffer blanks
were filtered using 0.22 µm syringe filters and loaded into an Axygen 96-well polypropylene

PCR Microplate (Corning product# PCR-96-FS-C), which was then sealed with an AxyMat
Sealing Mat (product# AM-96-PCR-RD) and wrapped in parafilm. Plates were shipped to
Lawrence Berkeley National Labs in a styrofoam cooler filled with cold packs. All SAXS
measurements were performed by the SIBYLS group at the Lawrence Berkeley National
Laboratory HT-SAXS beamline (12.3.1) [79,80]. For technical restrictions, proteins were
measured at 20:1 and 200:1 molar ratios of trehalose and sucrose instead of 10:1 and 100:1.
The exception to this was CAHS D, which was tested with a wider range of molar ratios (1.6:1,
16:1, and 160:1).

We note that the radius of gyration that we calculated for CAHS D is approximately 5 angstroms
greater than previously reported by our group [31]. We believe that this difference can be
attributed to minor differences in our approach. While we used Size Exclusion Chromatography
(SEC)-coupled SAXS and a relatively dilute CAHS D sample in our previous study, the number
of SAXS experiments in this study necessitated a higher-throughput approach that omitted the
SEC step. We therefore believe that our control samples contained a higher fraction of transient
oligomeric species, which inflated the radius of gyration without significantly curving the Guinier
region. Given that this fact was consistent between all of our CAHS D samples, the inter-
environmental comparisons are still valid.

Small-Angle X-ray Scattering (SAXS) – Guinier and pMW Analysis
Notable aggregation, likely induced by exposure to X-rays, was present in some samples,
especially in solutions that contained cosolutes. This was controlled for by excluding scattering
data from samples that had already been exposed to large amounts of X-ray radiation and were
thus statistically different from the initial readings. Despite some readings having been
excluded, a Guinier analysis was able to be conducted for each protein-cosolute combination.
Buffer subtractions and Guinier analysis were performed using BioXTAS RAW v. 2.1.4 [81,82].
A $qMaxR_g$ of 1.1 was used to establish linear fits in the Guinier region [55,83]. Samples with
Guinier regions that could not be fit were excluded from the study. Molecular weight
approximations were performed using the method described in Hajizadeh et al. 2018, which is
programmed directly into BioXTAS RAW [84]. The 8 mg/mL samples tended to be far more
aggregation-prone than the 4 mg/mL samples, so only the 4 mg/mL data is reported here.

Photo-Induced Cross-Linking of Unmodified Proteins (PICUP)
PICUP crosslinking was performed as previously described [60,63]. Briefly, lyophilized protein,
Ru(II)bpy32+, and ammonium persulfate were resuspended in 20 mM Tris pH 7.5. Each
reaction mixture constituted protein at the desired concentration with 1.25 mM Tris (2,2'-
bipyridyl)dichlororuthenium(II)hexahydrate (Sigma, CAS No. 50525-27-4, and 2.5 mM
ammonium persulfate (Sigma, CAS No. 7727-54-0) to a final volume of 10 µL. For mixtures,
cosolutes and proteins were mixed at a 100:1 molar ratio at 2X molar concentration.
Photoreaction was triggered by flashing 72 W light through a 2.5 cm water filter for 10 seconds
in a dark room. The reaction was immediately quenched by adding 10 µL of 2X Laemmli buffer
containing 4 % SDS and 10 % β-mercaptoethanol. The reaction mixture was heated at 95 °C for
5 minutes. 8.5 µL of each sample was run in denaturing SDS-PAGE gels and stained with
coomassie blue to visualize the oligomeric states.

Differential Scanning Calorimetry (DSC) Measurements

Differential Scanning Calorimetry (DSC) Measurements
Samples were prepared in Eppendorf tubes at the desired molar ratios with cosolutes. Protein
mixtures were resuspended and incubated at 55 °C for 5 minutes to ensure proper solubility. 25
µL of the sample was hermetically sealed into a previously massed pair of DSC aluminum
hermetic pan and hermetic lid (Catalog 900793.901 and 901684.901, respectively, TA
instruments). The sample mass was determined after the sample was sealed within the pan and
lid. The sealed samples were then run on a TA DSC2500 instrument. The DSC method for
heating experiments is as follows:

Samples were equilibrated at 20 °C, heated to 60 °C at a 5 °C per minute ramp, and then
cooled to 20 °C at a 5 °C per minute ramp. Samples were held for a 10 minute isothermal hold
at 20 °C and heated to 60 °C at a 5 °C per minute ramp. Trios software (TRIOS version
#5.0.0.44608, TA Instruments) was used to analyze enthalpy for samples showing the melt
curves. The changes in enthalpy for the mixtures were calculated relative to the protein.

AlphaFold2 Structural Modelling

AlphaFold2 Structural Modelling
Protein structure predictions (both monomeric and multimeric) were performed using Google's
AlphaFold Colab notebook. The setting "relax_use_gpu" was checked to increase the speed of
individual predictions. The number of recycles was left at 3 [85]. Representative images of each
structure are provided (Fig. S8). All analysis was done in triplicate to account for variation in
AlphaFold's predictions. Representative images of each structure are provided (Fig. S8). All pdb
files can be found in supplementary data.

Transfer Free Energy (TFE) Calculations

782 **Transfer Free Energy (TFE) Calculations**
783 TFE values of each cosolute for each amino acid were pulled from existing literature [68,69,86].
784 We used experimentally derived TFE values for the transfer of a chemical group - amino acid
785 side chains or backbone - into 1 M solutions of trehalose, sucrose, or glycine betaine [67–69].
786 We then calculated the TFE for each conformation using the formula:

$$\Delta G_{tr} = \sum_{N} \sum_{i=1} \alpha_i g_N$$

Here, ΔG_{tr} is the TFE of a protein conformation from water to 1 M cosolute solution, *N* is the
chemical group, *i* is a numerical index for all instances of the chemical group, α is the surface
area of the specific instance of the chemical group in square angstrom, and *g* is the
experimental value of the transfer free energy for that chemical group per square angstrom of
exposed surface area [67]. By doing this for two conformations of a protein, one can find the
change in free energy of conformational change that can be attributed to the presence of an
osmolyte.

Importantly, two "end-state" protein conformations are required to calculate a $\Delta\Delta G_{tr}$ value,
which is typically done for well-folded proteins [67]. However, we believe the proteins used in
this study constitute an important exception. For CAHS, several different groups have noted the
propensity of CAHS proteins for oligomerization, and recent research has identified the dimer as

a particularly stable CAHS D conformer [31,33,35]. This notion is supported by the crosslinking
data, in which the CAHS D dimer is especially prominent [31]. We thus use the following
equation for the effects of cosolutes on CAHS dimerization:
802

$$\Delta\Delta G_{tr}^{M \to D} = \Delta G_{tr}^{D} - 2 \times \Delta G_{tr}^{M}$$

804 Where ΔG_{tr}^{M} and ΔG_{tr}^{D} is the TFE for the monomeric and dimeric state, respectively. The
805 monomer and dimer conformations were obtained from AlphaFold Multimer and AlphaFold 2
806 predictions, respectively (see AlphaFold method). Surface area for residues was calculated
807 using SOURSOP, a python package for protein structure analysis [87].
808

809 To calculate the $\Delta\Delta G_{tr}^{folding}$ for our LEA proteins and for BSA, we compared an AlphaFold
810 prediction of a monomeric protein with a theoretical conformation in which all residues are 100%
811 exposed. This is because (a) LEAs showed no tendency to form a gel, and (b) the CD spectra
812 switched between a primarily disordered conformation to a helical conformation upon
813 desiccation (Fig. 3). For all LEAs, we calculated

$$\Delta\Delta G_{tr}^{folding} = \Delta G_{tr}^{final} - \Delta G_{tr}^{initial}$$

Here, $\Delta G_{tr}^{initial}$ represents the free energy of a completely denatured protein chain where the
maximum theoretical accessibility is achieved (RASA = 1). $\Delta G_{tr}^{initial}$ represents the free energy
of the protein's "native" conformation (as determined by an AlphaFold prediction). AlphaFold
predictions of our LEA proteins were broadly helical (Fig. S8), and thus were used to represent
the disorder-to-helix transition commonly observed in LEA proteins.

Data Analysis and Visualisation
LDH protection was fitted into a sigmoidal curve by fitting a 5PL regression analysis using
GraphPad Prism v9.5.1 from which the resulting PD50 values were derived. Other plots were
plotted using R-Studio. Annotation for statistical significance include: p>0.05: NS, p= 0.01-0.05:
*, p= 0.001-0.01: **, p<0.001: ***
824
825
826

831
832
833

Figure 1: LEA motifs are not synergistic with endogenous cosolutes. A) Sigmoidal plot representing percent of LDH stabilization by LEA motifs and cosolutes as a function of the molar concentration. n=3, error bars=standard deviation. **B)** Protective dose 50 (PD50) for additives obtained by sigmoidal fitting of data in Fig. 1A. n=3, N/A represents instances where 50% protection was not achieved. **C)** Example plot showing possible outcomes (additive, synergistic, or antagonistic effect) from cosolute:IDP mixtures. "Cosolute" and "Protein" represent the percent LDH protection by cosolute and protein respectively. Experimental represents the experimental protection resulting from cosolute and protein mixtures. **D)** Synergy plots for trehalose:LEA peptide at 100:1 molar ratio. n=3, Welch's t-test was used for statistical comparison, error bars = standard deviation. **E)** Synergy plots for sucrose:LEA peptide at 100:1 molar ratio. **F)** Plot representing synergy with trehalose vs. sucrose from Fig. 1D and Fig. 1E. Dotted line represents instances where there is equal synergy with both trehalose and sucrose.

Figure 2: Full-length desiccation-related IDPs act synergistically with cosolutes. A)
Concentration dependence of LDH protection by full-length proteins and cosolutes used in this
study **B)** PD50 for additives obtained by sigmoidal fitting of concentration-dependent LDH
protection from Fig. 2A. n=3, N/A represents instances where 50% protection was not achieved.
C) Synergy plots for trehalose:protein at 100:1 molar ratio. n = 3, Welch's t-test was used for
statistical comparison, error bars = standard deviation. **D)** Synergy plots for sucrose: protein at
100:1 molar ratio **E)** Plot representing synergy with trehalose vs. sucrose from Fig. 2C and Fig.
2D.

861
862
Figure 3: Functional synergy is not mediated by secondary structural changes. Buffer
represents the CD analysis for the proteins without cosolutes. Trehalose and Sucrose represent
the CD analysis for trehalose:protein mixture and sucrose:protein mixture at 100:1 molar ratio
respectively **A, B)** CD spectra for protein and cosolute:protein mixtures at 100:1 molar ratio
under aqueous **(A)** and **(B)** desiccated conditions. Each plot represents the average of three
replicates, with the shaded region representing the standard deviation of the average **C)**
Changes in the ratio of CD signal at 222 and 210 nm for individual protein and cosolute:protein
mixtures under aqueous and desiccated conditions. n=3, error bars = standard deviation,
statistical analysis was done by Welch's t-test.
872

Figure 4: Cosolutes increase the global dimensions of CAHS D, but not LEA or BSA. Analyzed data from SAXS experiments of proteins at 4 mg/mL in 20 mM tris HCl pH 7. Proteins tested include **A)** BSA (0.0578 mM), **B)** AtLEA3-3 (0.22 mM), **C)** AavLEA1 (0.249 mM), **D)** HeLEA68614 (0.156 mM), **E)** AvLEA1C (0.163 mM), **F)** AtLEA4-2 (0.38mM), and **G)** CAHS D (0.156 mM). The left plot shows the radius of gyration of the protein in the presence of no cosolute (gray), increasing molar ratios of trehalose (blue shades), and increasing molar ratios of sucrose (green shades). Error bars represent uncertainty in the measurement, provided by BioXTAS RAW. The right plot shows molecular weight values derived from Guinier analysis (see Methods). The red dashed line indicates the monomeric protein's molecular weight. Color scheme is the same as in the left figure. Error bars represent >90% confidence interval, which is directly obtained from the analysis.

Figure 5: DSC thermograms show cosolutes promote gelation of CAHS D but not LEA proteins. A) DSC thermogram of 0.235 mM CAHS D with trehalose (left, blue lines) and sucrose (right, green lines) at increasing cosolute:protein molar ratios **B)** Change in enthalpy measurements for cosolute:CAHS D mixtures relative to CAHS D. Enthalpy measurements were done by taking the area of gel melt peaks represented by black dashes in (A) (see Methods). **C-H)** DSC thermogram of LEA proteins and BSA at 0.235 mM and with the addition of trehalose and sucrose at molar ratios of 1:1, 10:1, 100:1, and 500:1 (cosolute:protein). **I)** DSC thermogram for trehalose and sucrose in the absence of proteins at respective concentrations for different molar ratios.

Figure 6: Transfer free energy of CAHS D highlights the difference between synergistic
and non-synergistic cosolutes. A) Tanford's transfer model, depicting the effect of cosolutes
on the dimerization of two proteins. "M" represents the protein's monomeric state, "D"
represents the dimeric state, "aq" represents an aqueous solution, and "os" represents a
solution containing some osmolyte. **B)** The difference in $\Delta\Delta G_{tr}^{M\rightarrow D}$ between two CAHS D
monomers and a CAHS D dimer. All structures are predicted by AlphaFold2 [85,88]. **C)** DSC
thermograms of CAHS D at 6 mg/mL (0.235 mM) in varying molar ratios of glycine betaine **D)**
Change in enthalpy measurements for betaine:CAHS D mixtures relative to CAHS D at 6
mg/mL. Enthalpy measurements were done by taking the area of gel melt peaks represented by
black dashes in Fig. 6C. **E)** Same as (C) but with CAHS D at 12 mg/mL (0.47 mM) **F)** Change in
enthalpy measurements for betaine:CAHS D mixtures relative to CAHS D at 12 mg/mL. **G)**
SAXS data depicting scattering profiles of 4 mg/mL CAHS D in tris (black), 1000:1 trehalose
(blue), 1000:1 sucrose (green), and 1000:1 glycine betaine (red). **H)** LDH synergy assay for
glycine betaine and CAHS D at various molar ratios. **I)** A correlation of $\Delta\Delta G_{tr}^{M\rightarrow D}$ of a given
cosolute with CAHS D and its synergy in the LDH assay at different molar ratios. p-value is
given from a Pearson correlation.

Organism	Enriched disaccharide during desiccation	Protein Family	LEA Motif	Representative protein	Length (amino acid)	Motif (%)	Metapredict Disorder
Arabidopsis thaliana	Sucrose (high) / Trehalose (high) Angeles-Núñez and Tiessen 2010, Fait et al. 2006	LEA_4	AKSKADETLES	AtLEA3-3	169	19.5%	
Aphelenchus avenae	Trehalose (high) Higa and Womersley 1993, Browne et al. 2004	LEA_4	LRDTAAEKLHQ	AavLEA1	143	30.06%	
Hypsibius exemplaris	Trehalose (low) Hengherr et al. 2008, Nguyen et al. 2022	LEA_4	LKDKAGSAWNQ	HeLEA68614	238	26.85%	
Adineta vaga	X Tunacliffe and Lapinski 2003, Lapinski and Tunacliffe 2003	LEA_4	KDKAAEALDAI	AvLEA1C	231	25.54%	
Arabidopsis thaliana	Sucrose (high) / Trehalose (high) Angeles-Núñez and Tiessen 2010, Fait et al. 2006	LEA_1	AKEKNIGGA KAQGHAEKTM	AtLEA4-2	97	73.19%	
Hypsibius exemplaris	Trehalose (low) Hengherr et al. 2008, Nguyen et al. 2022	CAHS	X	CAHS D	227	▬	

Table 1. Summary of organisms, disaccharides, and proteins used in this study. Table displaying the organismal source of representative LEA_4, LEA_1 and CAHS proteins used in this study. In addition, the table displays endogenous cosolute reported in the literature to be co-enriched alongside LEA and CAHS proteins during desiccation in the given organism. The consensus sequence of 11-mer LEA_4 or 20-mer LEA_1 motifs as well as the length of the full-length proteins and predicted disorder using Metapredict [89] are shown. Shaded areas in the disorder plot correspond to the motif coordinates in the full-length LEA proteins. We note that the reason many of these profiles contain large folded regions is because the amphipathic LEA and CAHS proteins are predicted to form helices, which metapredict infers and incorrectly highlights these regions as 'folded' when really they are disordered in isolation.

956 **Supplemental Figures**

Figure S1. A) Sigmoidal plot representing concentration dependence of LDH protection by LEA motifs and cosolutes as a function of the weight/volume concentration **B)** Synergy plots for cosolutes with LEA motifs at 1:100, 1:10, 1:1, 10:1, and 100:1 molar ratios. n=3, Welch's t-test was used for statistical comparison, error bars = standard deviation. Numbers under the checkmarks represent the mean percentage synergy and standard deviation **C)** Change in NADH absorbance values for the refrigerated controls of Aav11 at different concentrations after 16 hours **D)** Percent synergy comparison for At20 with trehalose and sucrose at 100:1 ratio. Welch's t-test was used for statistical comparison.

Figure S2. A) Sigmoidal plot representing concentration dependence of LDH protection by full-length proteins and cosolutes as a function of the weight/volume concentration. **B)** Synergy plots for cosolute with full-length proteins at respective molar ratios. n=3, Welch's t-test was used for statistical comparison, error bars = standard deviation. Numbers under the checkmarks represent the mean percentage synergy and standard deviation **C-E)** Percent synergy comparison for **C)** AtLEA3-3 **D)** AavLEA1 **E)** CAHS D with trehalose and sucrose at 100:1 ratio. n=3, error bars= standard deviation. Welch's t-test was used for statistical comparison.

Figure S3. A) Plot comparing secondary structural changes in proteins with trehalose vs. sucrose at 100:1 molar ratio under hydrated conditions. Dotted line represents cases where there is equal structural shift in both trehalose and sucrose mixture. **B)** Plot comparing secondary structural alterations in proteins with trehalose vs. sucrose at 100:1 molar ratio under desiccated conditions. Dotted line represents cases where there is equal structural shift in both trehalose and sucrose mixture. **C)** Correlation plots for synergy vs. secondary structural shift in presence of trehalose in the hydrated state. **D)** Correlation plots for synergy vs. secondary structural shift in presence of trehalose in the desiccated state. **E)** Correlation plots for synergy vs. secondary structural shift in presence of sucrose in the hydrated state. **F)** Correlation plots for synergy vs. secondary structural shift in presence of sucrose in the desiccated state.

Figure S4. Raw SAXS data from the experiments shown in Fig. 4 for 4 mg/mL of **A)** BSA (0.0578 mM), **B)** AtLEA3-3 (0.22 mM), **C)** AavLEA1 (0.249 mM), **D)** HeLEA68614 (0.156 mM), **E)** AvLEA1C (0.163 mM), **F)** AtLEA4-2 (0.38 mM), and **G)** CAHS D (0.156 mM). Each plot displays raw SAXS data on a Guinier scale, with a zoomed-in portion in the top right displaying the Guinier region. The Guinier fit is displayed as a black trendline. In each plot, the protein's scattering profile in several solution environments is shown, including protein in the presence of no cosolute (black), a 20:1 molar ratio of trehalose (light blue), a 200:1 molar ratio of trehalose (dark blue), a 20:1 molar ratio of sucrose (light green), a 200:1 molar ratio of sucrose (dark green). CAHS D data contains slightly different molar ratios of 1.6:1 trehalose (very light blue), 16:1 trehalose (light blue), 160:1 trehalose (dark blue), 1.6:1 sucrose (very light green), 16:1 sucrose (light green), 160:1 sucrose (dark green). Unlike other samples, a Guinier analysis with a $qMaxR_g$ of 1.1 could not be obtained. A best attempt to establish the R_g of this sample ($qMaxR_g \sim 1.35$) revealed a significant increase in the R_g. Given the clear curvature of the Guinier region, this data was consistent with the presence of large oligomeric species.

1005

1006
1007
1008

Figure S5. Crosslinking gels for full-length LEA proteins and in mixtures with trehalose or
sucrose. A range of protein concentrations were used to cover the concentrations used in the
synergy assay. Protein concentrations used were **A)** 25 µM, **B)** 50 µM, **C)** 75 µM, **D)** 100 µM, **E)**
150 µM, and **F)** 200 µM. Each gel contains protein ladder (Precision Plus Protein™ Dual Xtra
Standards, Bio Rad, Catalog #161-0377) in the first lane, protein at desired concentration in the
second lane, crosslinked protein in the third lane, 100:1 trehalose:protein in the fourth lane,
crosslinked 100:1 trehalose:protein in the fifth lane, 100:1 sucrose:protein in the sixth lane, and
crosslinked 100:1 sucrose:protein in the last lane.

1017

1018
Figure S6. A) Sample enthalpy analysis using TRIOS software. First derivative analysis was
used to aid in selecting the onset and endset in the thermogram representative of protein
gelation. This is done by identifying observable spikes that correspond to where the curve
begins to dip (onset) or level out (endset). The area between the approximate onset and endset
is selected and TRIOS software is used to calculate defined onset and endset values, enthalpy
(J/g), and peak temperature (Tm, ˚C) by analyzing the area above this feature within the
identified range. **B)** DSC thermograms of CAHS D at 12 mg/mL (0.47 mM) in varying molar
ratios of trehalose (0:1, 1:1, 10:1, 100:1, and 500:1). **C)** DSC thermograms of CAHS D at 12
mg/mL (0.47 mM) in varying molar ratios of sucrose (0:1, 1:1, 10:1, 100:1, and 500:1). **D)**
Change in enthalpy measurements for trehalose:CAHS D mixtures relative to CAHS D at 12
mg/mL. Enthalpy measurements were done by taking the area of gel melt peaks represented by
black dashes in Fig. S6B **E)** Change in enthalpy measurements for sucrose:CAHS D mixtures
relative to CAHS D at 12 mg/mL. Enthalpy measurements were done by taking the area of gel
melt peaks represented by black dashes in Fig. S6C. **(F-I)** Melting temperature for the gel melts
as calculated from the DSC thermograms for CAHS D and cosolute:CAHS D mixtures at
respective ratios.
1035

Figure S7. Raw SAXS data from the experiments shown in Fig. 6 for 4 mg/mL of CAHS D in various solution conditions. **A)** Scattering curves for CAHS D in tris (gray), 10:1 glycine betaine (light red), 100:1 glycine betaine (medium red), 1000:1 glycine betaine (dark red). Guinier regions are displayed in the top left with Guinier fits drawn in black. **B)** Dimensionless Kratky analysis of the data from A. **C)** Zoomed in Guinier regions of CAHS D in tris (gray), 1000:1 glycine betaine (red), 1000:1 trehalose (blue), and 1000:1 sucrose (green). **D)** Scattering profiles for CAHS D in glycine betaine, trehalose, and sucrose normalized to the protein's scattering profile in tris. **E-J)** Scatterplots of $\Delta\Delta G_{tr}^{folding}$ and synergy for BSA and LEA proteins. All fits are done using linear regression. All p-values derived from Pearson correlation. **K-L)** Melting temperature for the gel melts as calculated from the DSC thermograms for CAHS D and Betaine:CAHS D mixtures at respective ratios.

1050
1051
Figure S8. Representative images of the pdb structures obtained from our AlphaFold predictions. All structures are colored by their pLDDT on a scale from 0 (red) to 100 (blue). **A**) Monomeric BSA. **B**) Monomeric AtLEA3-3. **C**) Monomeric AavLEA1. **D**) Monomeric HeLEA68614. **E**) Monomeric AvLEA1C. **F**) Monomeric AtLEA4-2. **G**) Monomeric CAHS D. **H**) Dimeric CAHS D. See supplementary data for pdb files of each prediction & their exact pLDDT.

Organism	Motif	Concentration (mM)
	At11	0.297
	At22	0.277
	At44	0.161
	Aav11	0.0562
	He11	0.287
	Av11	0.262
	At20	0.264

Table S1. Protein concentrations used in synergy assays for LEA motifs in this study.

Organism	Protein	Concentration (mM)
	AtLEA3-3	0.1215
	AavLEA1	0.112
	HeLEA68614	0.0392
	AvLEA1C	0.068
	AtLEA4-2	0.217
	CAHS D	0.0078
	BSA	0.054

Table S2. Protein concentrations used in synergy assays for full-length proteins in this study.

References

1.   Ward JJ, Sodhi JS, McGuffin LJ, Buxton BF, Jones DT. Prediction and functional analysis
of native disorder in proteins from the three kingdoms of life. J Mol Biol. 2004;337: 635–
645.

2.   Holehouse AS, Kragelund BB. The molecular basis for cellular function of intrinsically
disordered protein regions. Nat Rev Mol Cell Biol. 2023. doi:10.1038/s41580-023-00673-0

3.   van der Lee R, Buljan M, Lang B, Weatheritt RJ, Daughdrill GW, Dunker AK, et al.
Classification of intrinsically disordered regions and proteins. Chem Rev. 2014;114: 6589–
6631.

4.   Mao AH, Lyle N, Pappu RV. Describing sequence-ensemble relationships for intrinsically
disordered proteins. Biochem J. 2013;449: 307–318.

5.   Kedia N, Almisry M, Bieschke J. Glucose directs amyloid-beta into membrane-active
oligomers. Phys Chem Chem Phys. 2017;19: 18036–18046.

6.   Das RK, Ruff KM, Pappu RV. Relating sequence encoded information to form and function
of intrinsically disordered proteins. Curr Opin Struct Biol. 2015;32: 102–112.

7.   Soranno A, Koenig I, Borgia MB, Hofmann H, Zosel F, Nettels D, et al. Single-molecule
spectroscopy reveals polymer effects of disordered proteins in crowded environments. Proc
Natl Acad Sci U S A. 2014;111: 4874–4879.

8.   Moses D, Yu F, Ginell GM, Shamoon NM, Koenig PS, Holehouse AS, et al. Revealing the
Hidden Sensitivity of Intrinsically Disordered Proteins to their Chemical Environment. J
Phys Chem Lett. 2020;11: 10131–10136.

9.   Moses D, Ginell GM, Holehouse AS, Sukenik S. Intrinsically disordered regions are poised
to act as sensors of cellular chemistry. Trends Biochem Sci. 2023;48: 1019–1034.

10.   Theillet F-X, Binolfi A, Frembgen-Kesner T, Hingorani K, Sarkar M, Kyne C, et al.
Physicochemical properties of cells and their effects on intrinsically disordered proteins
(IDPs). Chem Rev. 2014;114: 6661–6714.

11.   Crowe JH. Anhydrobiosis: an unsolved problem. Plant, cell & environment. 2014. pp. 1491–
1493.

12.   Hesgrove C, Boothby TC. The biology of tardigrade disordered proteins in extreme stress
tolerance. Cell Commun Signal. 2020;18: 178.

13.   Romero-Perez PS, Dorone Y, Flores E, Sukenik S, Boeynaems S. When Phased without
Water: Biophysics of Cellular Desiccation, from Biomolecules to Condensates. Chem Rev.
2023;123: 9010–9035.

14.   Oliver MJ, Farrant JM, Hilhorst HWM, Mundree S, Williams B, Bewley JD. Desiccation
Tolerance: Avoiding Cellular Damage During Drying and Rehydration. Annu Rev Plant Biol.
2020;71: 435–460.

15.   Koster KL, Leopold AC. Sugars and desiccation tolerance in seeds. Plant Physiol. 1988;88:

829–832.

16. Angeles-Núñez JG, Tiessen A. Arabidopsis sucrose synthase 2 and 3 modulate metabolic
homeostasis and direct carbon towards starch synthesis in developing seeds. Planta.
2010;232: 701–718.

17. Fait A, Angelovici R, Less H, Ohad I, Urbanczyk-Wochniak E, Fernie AR, et al. Arabidopsis
seed development and germination is associated with temporally distinct metabolic
switches. Plant Physiol. 2006;142: 839–854.

18. Ryabova A, Cornette R, Cherkasov A, Watanabe M, Okuda T, Shagimardanova E, et al.
Combined metabolome and transcriptome analysis reveals key components of complete
desiccation tolerance in an anhydrobiotic insect. Proc Natl Acad Sci U S A. 2020;117:
19209–19220.

19. Nguyen K, Kc S, Gonzalez T, Tapia H, Boothby TC. Trehalose and tardigrade CAHS
proteins work synergistically to promote desiccation tolerance. Commun Biol. 2022;5: 1046.

20. Tapia H, Koshland DE. Trehalose is a versatile and long-lived chaperone for desiccation
tolerance. Curr Biol. 2014;24: 2758–2766.

21. Hincha DK, Thalhammer A. LEA proteins: IDPs with versatile functions in cellular
dehydration tolerance. Biochem Soc Trans. 2012;40: 1000–1003.

22. Boothby TC, Tapia H, Brozena AH, Piszkiewicz S, Smith AE, Giovannini I, et al.
Tardigrades Use Intrinsically Disordered Proteins to Survive Desiccation. Mol Cell. 2017;65:
975–984.e5.

23. Hernández-Sánchez IE, Maruri-López I, Martinez-Martinez C, Janis B, Jiménez-Bremont
JF, Covarrubias AA, et al. LEAfing through literature: late embryogenesis abundant proteins
coming of age—achievements and perspectives. J Exp Bot. 2022;73: 6525–6546.

24. Hand SC, Menze MA, Toner M, Boswell L, Moore D. LEA proteins during water stress: not
just for plants anymore. Annu Rev Physiol. 2011;73: 115–134.

25. Hundertmark M, Hincha DK. LEA (late embryogenesis abundant) proteins and their
encoding genes in Arabidopsis thaliana. BMC Genomics. 2008;9: 118.

26. Artur MAS, Zhao T, Ligterink W, Schranz E, Hilhorst HWM. Dissecting the Genomic
Diversification of Late Embryogenesis Abundant (LEA) Protein Gene Families in Plants.
Genome Biol Evol. 2019;11: 459–471.

27. Yamaguchi A, Tanaka S, Yamaguchi S, Kuwahara H, Takamura C, Imajoh-Ohmi S, et al.
Two novel heat-soluble protein families abundantly expressed in an anhydrobiotic
tardigrade. PLoS One. 2012;7: e44209.

28. Goyal K, Walton LJ, Tunnacliffe A. LEA proteins prevent protein aggregation due to water
stress. Biochem J. 2005;388: 151–157.

29. Kim SX, Çamdere G, Hu X, Koshland D, Tapia H. Synergy between the small intrinsically
disordered protein Hsp12 and trehalose sustain viability after severe desiccation. Elife.
2018;7. doi:10.7554/eLife.38337

30. LeBlanc BM, Hand SC. Target enzymes are stabilized by AfrLEA6 and a gain of α-helix
coincides with protection by a group 3 LEA protein during incremental drying. Biochim
Biophys Acta: Proteins Proteomics. 2021;1869: 140642.

31. Sanchez-Martinez S, Nguyen K, Biswas S, Nicholson V, Romanyuk AV, Ramirez J, et al.
Labile assembly of a tardigrade protein induces biostasis. bioRxiv. 2023. p.
2023.06.30.547219. doi:10.1101/2023.06.30.547219

32. Eicher JE, Brom JA, Wang S, Sheiko SS, Atkin JM, Pielak GJ. Secondary structure and
stability of a gel-forming tardigrade desiccation-tolerance protein. Protein Sci. 2022;31:
e4495.

33. Malki A, Teulon J-M, Camacho-Zarco AR, Chen S-WW, Adamski W, Maurin D, et al.
Intrinsically Disordered Tardigrade Proteins Self-Assemble into Fibrous Gels in Response
to Environmental Stress. Angew Chem Int Ed Engl. 2022;61: e202109961.

34. Yagi-Utsumi M, Aoki K, Watanabe H, Song C, Nishimura S, Satoh T, et al. Desiccation-
induced fibrous condensation of CAHS protein from an anhydrobiotic tardigrade. Sci Rep.
2021;11: 21328.

35. Tanaka A, Nakano T, Watanabe K, Masuda K, Honda G, Kamata S, et al. Stress-
dependent cell stiffening by tardigrade tolerance proteins that reversibly form a filamentous
network and gel. PLoS Biol. 2022;20: e3001780.

36. Battaglia M, Olvera-Carrillo Y, Garciarrubio A, Campos F, Covarrubias AA. The enigmatic
LEA proteins and other hydrophilins. Plant Physiol. 2008;148: 6–24.

37. Higa LM, Womersley CZ. New insights into the anhydrobiotic phenomenon: The effects of
trehalose content and differential rates of evaporative water loss on the survival of
Aphelenchus avenae. J Exp Zool. 1993;267: 120–129.

38. Browne JA, Dolan KM, Tyson T, Goyal K, Tunnacliffe A, Burnell AM. Dehydration-specific
induction of hydrophilic protein genes in the anhydrobiotic nematode Aphelenchus avenae.
Eukaryot Cell. 2004;3: 966–975.

39. Hengherr S, Heyer AG, Köhler H-R, Schill RO. Trehalose and anhydrobiosis in tardigrades-
-evidence for divergence in responses to dehydration. FEBS J. 2008;275: 281–288.

40. Tunnacliffe A, Lapinski J. Resurrecting Van Leeuwenhoek's rotifers: a reappraisal of the
role of disaccharides in anhydrobiosis. Philos Trans R Soc Lond B Biol Sci. 2003;358:
1755–1771.

41. Lapinski J, Tunnacliffe A. Anhydrobiosis without trehalose in bdelloid rotifers. FEBS Lett.
2003;553: 387–390.

42. Dure L 3rd. A repeating 11-mer amino acid motif and plant desiccation. Plant J. 1993;3:
363–369.

43. Knox-Brown P, Rindfleisch T, Günther A, Balow K, Bremer A, Walther D, et al. Similar Yet
Different-Structural and Functional Diversity among LEA_4 Proteins. Int J Mol Sci. 2020;21.
doi:10.3390/ijms21082794

44. Furuki T, Sakurai M. Group 3 LEA protein model peptides protect enzymes against

desiccation stress. Biochim Biophys Acta. 2016;1864: 1237–1243.

45. Furuki T, Shimizu T, Chakrabortee S, Yamakawa K, Hatanaka R, Takahashi T, et al.
Effects of Group 3 LEA protein model peptides on desiccation-induced protein aggregation.
Biochim Biophys Acta. 2012;1824: 891–897.

46. Furuki T, Sakurai M. Group 3 LEA protein model peptides protect liposomes during
desiccation. Biochim Biophys Acta. 2014;1838: 2757–2766.

47. Hibshman JD, Goldstein B. LEA motifs promote desiccation tolerance in vivo. BMC Biol.
2021;19: 263.

48. Piszkiewicz S, Gunn KH, Warmuth O, Propst A, Mehta A, Nguyen KH, et al. Protecting
activity of desiccated enzymes. Protein Sci. 2019;28: 941–951.

49. Shimizu T, Kanamori Y, Furuki T, Kikawada T, Okuda T, Takahashi T, et al. Desiccation-
induced structuralization and glass formation of group 3 late embryogenesis abundant
protein model peptides. Biochemistry. 2010;49: 1093–1104.

50. Wang S, Eicher J, Pielak GJ. Trifluoroethanol and the behavior of a tardigrade desiccation-
tolerance protein. Protein Sci. 2023;32: e4716.

51. Cuevas-Velazquez CL, Saab-Rincón G, Reyes JL, Covarrubias AA. The Unstructured N-
terminal Region of Arabidopsis Group 4 Late Embryogenesis Abundant (LEA) Proteins Is
Required for Folding and for Chaperone-like Activity under Water Deficit. J Biol Chem.
2016;291: 10893–10903.

52. Bremer A, Wolff M, Thalhammer A, Hincha DK. Folding of intrinsically disordered plant LEA
proteins is driven by glycerol-induced crowding and the presence of membranes. FEBS J.
2017;284: 919–936.

53. Navarro-Retamal C, Bremer A, Alzate-Morales J, Caballero J, Hincha DK, González W, et
al. Molecular dynamics simulations and CD spectroscopy reveal hydration-induced
unfolding of the intrinsically disordered LEA proteins COR15A and COR15B from
Arabidopsis thaliana. Phys Chem Chem Phys. 2016;18: 25806–25816.

54. Greenfield NJ. Using circular dichroism spectra to estimate protein secondary structure. Nat
Protoc. 2006;1: 2876–2890.

55. Kachala M, Valentini E, Svergun DI. Application of SAXS for the Structural Characterization
of IDPs. Adv Exp Med Biol. 2015;870: 261–289.

56. Graewert MA, Franke D, Jeffries CM, Blanchet CE, Ruskule D, Kuhle K, et al. Automated
pipeline for purification, biophysical and x-ray analysis of biomacromolecular solutions. Sci
Rep. 2015;5: 10734.

57. Babcock JJ, Brancaleon L. Bovine serum albumin oligomers in the E- and B-forms at low
protein concentration and ionic strength. Int J Biol Macromol. 2013;53: 42–53.

58. Levi V, González Flecha FL. Reversible fast-dimerization of bovine serum albumin detected
by fluorescence resonance energy transfer. Biochim Biophys Acta. 2002;1599: 141–148.

59. Shou K, Bremer A, Rindfleisch T, Knox-Brown P, Hirai M, Rekas A, et al. Conformational

selection of the intrinsically disordered plant stress protein COR15A in response to solution
osmolarity – an X-ray and light scattering study. Phys Chem Chem Phys. 2019;21: 18727–
18740.

60. Rivera-Najera LY, Saab-Rincón G, Battaglia M, Amero C, Pulido NO, García-Hernández E,
et al. A group 6 late embryogenesis abundant protein from common bean is a disordered
protein with extended helical structure and oligomer-forming properties. J Biol Chem.
2014;289: 31995–32009.

61. Fancy DA, Kodadek T. Chemistry for the analysis of protein-protein interactions: rapid and
efficient cross-linking triggered by long wavelength light. Proc Natl Acad Sci U S A.
1999;96: 6020–6024.

62. Preston GW, Wilson AJ. Photo-induced covalent cross-linking for the analysis of
biomolecular interactions. Chem Soc Rev. 2013;42: 3289–3301.

63. Liu G, Liu K, Gao Y, Zheng Y. Involvement of C-Terminal Histidines in Soybean PM1
Protein Oligomerization and Cu2+ Binding. Plant Cell Physiol. 2017;58: 1018–1029.

64. Eicher J, Hutcheson BO, Pielak GJ. Properties of a tardigrade desiccation-tolerance protein
aerogel. Biophys J. 2023;122: 2500–2505.

65. Belott C, Janis B, Menze MA. Liquid-liquid phase separation promotes animal desiccation
tolerance. Proc Natl Acad Sci U S A. 2020;117: 27676–27684.

66. Tanford C. Protein denaturation. C. Theoretical models for the mechanism of denaturation.
Adv Protein Chem. 1970;24: 1–95.

67. Auton M, Bolen DW. Application of the transfer model to understand how naturally
occurring osmolytes affect protein stability. Methods Enzymol. 2007;428: 397–418.

68. Auton M, Bolen DW. Predicting the energetics of osmolyte-induced protein
folding/unfolding. Proc Natl Acad Sci U S A. 2005;102: 15065–15068.

69. Hong J, Gierasch LM, Liu Z. Its preferential interactions with biopolymers account for
diverse observed effects of trehalose. Biophys J. 2015;109: 144–153.

70. Wolkers WF, McCready S, Brandt WF, Lindsey GG, Hoekstra FA. Isolation and
characterization of a D-7 LEA protein from pollen that stabilizes glasses in vitro. Biochim
Biophys Acta. 2001;1544: 196–206.

71. Shih M-D, Hsieh T-Y, Jian W-T, Wu M-T, Yang S-J, Hoekstra FA, et al. Functional studies
of soybean (Glycine max L.) seed LEA proteins GmPM6, GmPM11, and GmPM30 by CD
and FTIR spectroscopy. Plant Sci. 2012;196: 152–159.

72. Hincha DK, Zuther E, Popova AV. Stabilization of Dry Sucrose Glasses by Four LEA_4
Proteins from. Biomolecules. 2021;11. doi:10.3390/biom11050615

73. Sakurai M, Furuki T, Akao K-I, Tanaka D, Nakahara Y, Kikawada T, et al. Vitrification is
essential for anhydrobiosis in an African chironomid, Polypedilum vanderplanki. Proc Natl
Acad Sci U S A. 2008;105: 5093–5098.

74. Water content, transition temperature and fragility influence protection and anhydrobiotic

capacity. BBA Advances. 2024; 100115.

75. Crowe JH, Carpenter JF, Crowe LM. The role of vitrification in anhydrobiosis. Annu Rev
Physiol. 1998;60: 73–103.

76. Simperler A, Kornherr A, Chopra R, Bonnet PA, Jones W, Motherwell WDS, et al. Glass
transition temperature of glucose, sucrose, and trehalose: an experimental and in silico
study. J Phys Chem B. 2006;110: 19678–19684.

77. Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, et al. Composition-
dependent thermodynamics of intracellular phase separation. Nature. 2020;581: 209–214.

78. Sarkar M, Li C, Pielak GJ. Soft interactions and crowding. Biophys Rev. 2013;5: 187–194.

79. Dyer KN, Hammel M, Rambo RP, Tsutakawa SE, Rodic I, Classen S, et al. High-
throughput SAXS for the characterization of biomolecules in solution: a practical approach.
Methods Mol Biol. 2014;1091: 245–258.

80. Trame C, MacDowell AA, Celestre RS, Padmore HA, Cambie D, Domning EE, et al.
SIBYLS—a SAXS and Protein Crystallography Beamline at the ALS. AIP Conference
Proceedings. American Institute of Physics; 2004. pp. 502–505.

81. Nielsen SS, Toft KN, Snakenborg D, Jeppesen MG, Jacobsen JK, Vestergaard B, et al.
BioXTAS RAW, a software program for high-throughput automated small-angle X-ray
scattering data reduction and preliminary analysis. J Appl Crystallogr. 2009;42: 959–964.

82. Hopkins JB, Gillilan RE, Skou S. BioXTAS RAW: a free open-source program for small-
angle X-ray scattering data reduction. Foundations of Crystallography. 2018;74: a219.

83. Rieloff E, Skepö M. Determining Rg of IDPs from SAXS Data. Methods Mol Biol.
2020;2141: 271–283.

84. Hajizadeh NR, Franke D, Jeffries CM, Svergun DI. Consensus Bayesian assessment of
protein molecular mass from solution X-ray scattering data. Sci Rep. 2018;8: 7204.

85. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596: 583–589.

86. Auton M, Bolen DW. Additive transfer free energies of the peptide backbone unit that are
independent of the model compound and the choice of concentration scale. Biochemistry.
2004;43: 1329–1342.

87. Lalmansingh JM, Keeley AT, Ruff KM, Pappu RV, Holehouse AS. SOURSOP: A Python
package for the analysis of simulations of intrinsically disordered proteins. bioRxiv. 2023.
doi:10.1101/2023.02.16.528879

88. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Tunyasuvunakool K, et al. AlphaFold 2.
Fourteenth Critical Assessment of Techniques for Protein Structure Prediction; DeepMind:
London, UK. 2020. Available:
https://predictioncenter.org/casp14/doc/presentations/2020_12_01_TS_predictor_AlphaFol
d2.pdf

89. Emenecker RJ, Griffith D, Holehouse AS. Metapredict: a fast, accurate, and easy-to-use

predictor of consensus disorder and structure. Biophys J. 2021;120: 4312–4319.

1311