1    **Title:** Choices to landscapes: Mechanisms of animal movement scale to landscape patterns
2

3    **Authors and affiliations:**
4    Will Rogers[1], will.rogers@yale.edu https://orcid.org/0000-0001-9899-8064
5    Scott Yanco[1,2] scott.yanco@yale.edu https://orcid.org/0000-0003-4717-9370
6    Walter Jetz[1,2], walter.jetz@yale.edu https://orcid.org/0000-0002-1971-7277
7
8    [1] Department of Ecology & Evolutionary Biology, Yale University, P. O. Box 208106, New Haven, CT
9    06520-8106
10   [2] Yale Center for Biodiversity and Global Change

11
12   **Running title:** Movement processes resolve spatial patterns of occurrence
13
14   **Keywords:** Species distribution, Habitat Selection, Occurrence, Movement, Step-selection, Ergodicity
15
16   **Statement of authorship:** WR and SY conceptualized the paper. WR wrote the code and carried out the
17   analyses supporting the manuscript. WR wrote the initial draft with contributions from SY. All authors
18   provided feedback on the concept and contributed to the final draft.
19

**Corresponding author:**
Name: Will Rogers
Email: will.rogers@yale.edu, rogerswill47@gmail.com
Mail: Department of Ecology & Evolutionary Biology, Yale University, P. O. Box 208106, New
Haven, CT 06520-8106
Address: 165 Prospect Street, New Haven, CT, US 06511
Phone: (203) 432-6138
Fax: (203) 432-5176

**Abstract:**

Understanding the geographic distributions of animals is central to ecological inquiry and

conservation planning. Movement-based habitat selection models, like discrete-time step-selection

functions, are useful for identifying key environmental attributes which animals select upon and can

nearly perfectly capture preference patterns between environmental conditions. However, predictions

based only on selection can often fail to accurately describe true geographic distributions. We show that

by integrating local selection patterns, movement models, and explicit landscape constraints within the

same framework, we can better predict simulated distributions than occurrence-based frameworks of

selection alone. Using three case studies, we show that this framework can better predict distributions

of organisms across increasing scales of out-of-sample prediction: within individuals, between

individuals, and even between regional contexts. Movement modeling is a powerful tool to describe

both the selection paradigm that organisms apply to the environment and the movement patterns that

enable them to apply those selection paradigms. By understanding movement and selection at the same

time, habitats that species "prefer" may not be ultimately occupied, as many "preferred" patches may

be unreachable given movement strategies and landscape context, often leading to smaller geographic

distributions than estimated by selection or occurrence alone.

**Introduction:**

Interactions between organisms and their environment shape fundamental ecological

phenomena such as species ranges (Kearney & Porter 2009), population dynamics (Roughgarden 1975;

Kareiva *et al.* 1990), community composition (Menge & Olson 1990), and evolutionary feedbacks (Post &

Palkovacs 2009). These relationships also provide critical applied insights about, for example,

spatiotemporal patterns of biological diversity (Buckley *et al.* 2011; Woodin *et al.* 2013; Evans *et al.*

2015), population trends (Bozinovic *et al.* 2011; Madin *et al.* 2012), or genetic connectivity (Cushman &

Lewis 2010; Creel *et al.* 2019; Fletcher Jr *et al.* 2022) – insights necessary to address the twin planetary-

scale ecological crises of climate change and biodiversity loss (Powers & Jetz 2019). Therefore, there is

significant value in frameworks that can link the relationship between organisms and their environments

to their niches (Kearney 2006; Matthiopoulos 2022), and make predictions from such relationships

about the distribution of individuals and species geographically

Inferring relationships between individuals and their environment has traditionally relied on

observations of occurrence patterns over some set of environmental contexts which vary geographically

(Boyce & McDonald 1999; Boyce *et al.* 2002), typically by using environmental dimensions to contrast

observations from locations where organisms are putatively absent (Mackey & Lindenmayer 2001;

Johnson *et al.* 2006; Moorcroft & Barnett 2008). For example, species distribution models (SDMs)

compare the patterns of presences as compared to possible null distributions over large geographic

extents (Elith *et al.* 2011; Merow *et al.* 2014). Resource selection functions (RSFs) - an methodologically

similar framework to SDMs (Fieberg *et al.* 2021; Northrup *et al.* 2022) - use logistic regression to model

the probability of individual occurrences within some domain of predefined availability, like home

ranges (Boyce & McDonald 1999; Manly *et al.* 2007). Both SDMs and RSFs are occurrence-based

frameworks, capture information about multiple biological mechanisms that lead to species or individual

occurrences while accounting for elements of the observation process itself (Keating & Cherry 2004;

Phillips *et al.* 2009; Sastre & Lobo 2009; Wisz & Guisan 2009; Newbold *et al.* 2010). Notably, SDMs and

RSFs assume that all environmental conditions within the landscape, an animal's home range, etc. are

available for that animal to have selected upon, irrespective of the geographic configuration of these

environments.

In reality, not all environmental conditions within some geographic extent may be available to

an individual, either because of physiological constraints on animal movement or the spatial

arrangement of habitats (Box 1). For example, geographically isolated patches of preferred

environmental conditions may be unreachable by an organism and, therefore, go unused (Soberon &

Peterson 2005; Matthiopoulos *et al.* 2020; Van Moorter *et al.* 2023). Thus, patterns of animal movement

interacting with underlying environmental preferences may strongly influence realized patterns of

spatial occurrence if movement constraints or geographic heterogeneity (or both) are severe (Fig. 1).

This is an important problem, as both SDMs and RSFs are unequipped to describe the movement as a

force driving patterns of environmental selection. Unlike these occurrence-based frameworks, Step-

selection functions (SSFs) use regularly-spaced GPS locations to estimate patterns of selection between

environmental conditions that are available based on the movement constraints of the organism,

comparing patterns of selection with each "step" in a GPS track (Avgar *et al.* 2016). As a result, SSFs may

capture more realistic relationships between organisms and their environment by more accurately

describing availability in local geographic extents.

While SSFs incorporate movement in selection processes and can even integrate selection with

movement characteristics (Avgar *et al.* 2016), there is currently not analytical method to use SSFs for

predicting geographic distributions of occurrence. SSFs are a form of conditional logistic regression,

meaning that model inference about preference for or against environments at the end of a step is

conditioned on the environments at the beginning of the step.  Therefore, spatial predictions can only

be made in comparison to some baseline conditions. In practice, researchers typically select some

arbitrary set of environmental conditions (frequently the averager or median conditions within the

prediction domain) to use as the basis for prediction. This projection of SSFs ignores the movement

constraints inherent in model parametrization and instead generates patterns of "selection" across a

landscape which may not be equivalent to actual relative use (Avgar *et al.* 2017). The inability of SSFs to

directly return geographical predictions remains a hurdle for the applicability of movement models for

ecological inference and conservation planning. To overcome this, some authors have suggested agent-

based simulations based on parameterized SSF Field (Potts & Borger 2023), which provide the ability to

truly interrogate SSF fit but may invoke computational constraints with no guarantee of ergodicity.

Similarly, multi-stage frameworks have used separate resource selection (to assign quality to locations)

and step selection components (to assign connectivity between locations) to estimate geographic

distributions based (Van Moorter *et al.* 2023), which provides scalability but discretizes mechanisms of

movement and selection.  Effectively capturing both the patterns of selection in environmental space

and then reprojecting those environmental conditions onto geographic space while considering

movement constraints is critical for our ability to link relationships between individuals and

environments to geographic distributions useful for ecological inferences (Schick *et al.* 2008;

Matthiopoulos 2022).

Here, we outline a conceptual approach to use SSFs as a model of concurrent movement and

selection processes to return geographical output about the probability of use that should (i)

parametrize organismal-environmental interactions more mechanistically than occurrence-based

frameworks and (ii) generate more accurate predictions of habitat distribution than occurrence or

selection frameworks that ignore movement when defining habitat over geographic domains.

Specifically, we use fitted SSFs to generate stochastic matrices of movement at local scales informed by

concurrent selection and movement limitations, and we retrieve stable-state distributions (SSDs) within

an ergodic framework. Because SSFs measure organismal-environment interactions at finer scales than

occurrence-based frameworks, we sought to test how occurrence-based (RSF), selection alone (SSF), and

concurrent selection and movement (SSD) predicted the distributions of organisms across the

landscape. We hypothesized that our framework incorporating both selection and movement would be

more accurate than frameworks based on occurrence or selection alone. We tested this hypothesis in a

simulation with known movement and selection constraints, as well as three biological datasets.

**Box 1: Individual fundamental preferences are not equivalent to realized use**

The suite of various ecological and environmental conditions necessary for species to persist is often conceptualized as a species' niche. This concept of the niche is divorced from spatial variation such that positions in niche space may correspond to multiple geographic locations (or none), a concept known as Hutchinson's duality (see Colwell & Rangel 2009). Because of this duality, it is possible to project a niche on a map, charting the areas that meet the conditions to support a species. While a tantalizing prospect, biogeographers were quick to note alarming inconsistencies between observed distributions of species and those expected from geographic niche projections (e.g., Pearson & Dawson 2003). Among a myriad of factors affecting the actual distributions of species, one nearly universal phenomenon is that the conditions species need to persist are discontinuous over real landscapes. To access these patches, species must disperse and cross regions of relatively low quality. Therefore, factors like dispersal and landscape heterogeneity are core constraints on a species' distribution (Pulliam 2000; Soberon & Peterson 2005; Soberón 2007), alongside factors like species interaction (e.g., competition, predation, parasitism) identified decades prior (Hutchinson 1978).

The role of dispersal in the geographic projection of species' niches is extremely relevant to individual- and population-level spatial inferences in movement ecology. While the individual niche concept is hotly contested, individuals within populations certainly vary physiologically and behaviorally. This variation should affect the distributions of individuals in space, potentially fueled by some latent difference in underlying needs or constraints. For example, male ungulates commonly occupy different environments than females, often with larger home ranges and occasionally more predation risk. This spatial segregation is hypothesized to arise from sex-based differences in reproductive constraints or resource requirements (reviewed in Main *et al.* 1996). Clearly, individual differences go far beyond sex,

with highly individual traits like personality likely factoring into space use (Spiegel *et al.* 2017). While quantifying individual variation in space use is an exciting endeavor, these differences should have real-world consequences for where individuals and their populations can persist, where they will go, and how they will get there – objectives central to our ability to inform conservation action.

Habitat selection modeling is a promising avenue to understand individual- and population-level space use and directly parallels niche models occurring at a species level. It is common to map the patterns of habitat selection of an individual, disregarding the spatial context of the habitat selection. For species niche modeling, an understanding of dispersal and heterogeneity was necessary to reconcile measured niches with the geographic distributions of species. Just as dispersal and landscape heterogeneity limit the ability of species to occupy all suitable conditions, movement traits, and landscape heterogeneity may limit the ability of individuals to occupy all suitable conditions in a landscape. Consequently, the underlying environmental preferences of an individual should not equal its real distribution geographically under movement constraints in heterogeneous landscapes (Fig. 1).

**Methods:**

*Landscapes as transition matrices*

We first considered a landscape, *G*, made up of *n* cells along with environmental data (Fig. 2A).

We treated the occupancy of these cells as states, between which individuals might transition according

to rates described by a stochastic transition matrix, *A*, of *n* rows and columns. We defined each value in

this matrix, $A_{i,j}$, as the probability of moving from one cell ($G_i$) to another ($G_j$) in the landscape. For an

animal in $G_i$, the sum of all transition probabilities to all other cells in *G* (including $G_i$) must sum to one

(we isolate the probability of transitioning from $G_i$ to $G_i$ for clarity later):

$$P(G_i|G_i) + \sum_{k=1,k\neq i}^n \quad P(G_{k=j}|G_i) = 1.$$ Eq. 1

In this way, Matrix *A* contains the directional probabilities of movement between all cells in *G*

based on the occupancy of any given cell $G_i$, parametrized by later movement modeling, creating a row-

stochastic matrix. The inverse of this matrix, $A^{-1}$, defines a column-stochastic matrix with a first principal

eigenvector (associated with the unit eigenvalue in a stochastic matrix) defining the probabilities of the

stable-state distribution across *G*.

*SSFs can parametrize transition matrices*

To gather these transition probabilities, we used predicted relative risk informed by fitted SSFs,

a quantity derived by comparing baseline conditions (at $G_i$) to a set of available conditions (*G*) and often

described as a relative strength of selection (Avgar et al. 2017). Relative risks are not equivalent to

probabilities but can be kernelized when the number of available choices are finite. To do this, we divide

Eq. 1 by the probability of moving from $G_i$ to $G_i$:

$$\frac{P(G_i|G_i)}{P(G_i|G_i)} + \sum_{k=1,k\neq i}^n \quad \frac{P(G_{k=j}|G_i)}{P(G_i|G_i)} = \frac{1}{P(G_i|G_i)}.$$ Eq. 2A

146    Therefore, Eq. 2A shows that the sum of relative risks for moving to all cells in *G* is equal to the inverse

147    probability of remaining in the current cell. Or, simplified, the inverse sum of all relative selection

148    strengths is equal to the probability of remaining in $G_i$ (Eq. 2B):

$$\frac{1}{1 + \sum_{k=1,k\neq i}^n \frac{P(G_{k=j}|G_i)}{P(G_i|G_i)}} = P(G_i|G_i).$$ Eq. 2B

More importantly, Eq. 2B now describes how to link a limited set of relative selection strengths to the

probabilities of remaining in place. Therefore, we could solve for the probabilities of moving to all

comparison cells by multiplying the relative selection strength returned by the SSF predictions by the

probability of staying in place.

*Extension to Integrated Step Selection Functions*

SSFs can also explicitly incorporate behavioral elements of the movement model independently

or (more commonly) interactively with the habitat selection model. For example, integrated SSFs (iSSFs)

frequently include a term to describe step distances, resulting in a distributional description of the likely

movement distances around a focal position (Avgar *et al.* 2016). Functionally, this forces the probability

of selection for very long-distance movements to zero even when habitat conditions are otherwise

suitable. In this way, iSSF models can define local neighborhoods for animal movement. By incorporating

distance into SSF models, we can confidently compare $G_i$ to cells in $G$ within a biologically-informed limit

on distance (herein the 95th percentile of observed step distances, Fig. 2G). This substantially reduces

the information burden carried by the transition matrix $A$ by treating it as a sparse matrix containing

only transitions to cells within a reasonable biological limit of movement (Fig. 2H). This dramatically

increases the speed of computing stable-state distributions (Fig. 2K), though users should explore the

tradeoff between the question at hand and computational demand.

We tested the performance of our method in describing animal distributions using one

simulated dataset and three real datasets. We compared the predictive performance of these

mechanistic models to models lacking individual behavioral mechanisms: RSFs and SSFs. While SSF

predictions capture patterns of selection, movement is ignored by assuming that local patterns of

selection rules can be generalized across entire landscapes (Fig. 2F). Similarly, RSF predictions can

capture patterns of occurrence; however, patterns are only observable over narrow spatial extents and

do not capture patterns of movement in predictions to landscapes. SSFs and RSFs therefore are

juxtaposed to the more mechanistic SSD approach.

*Simulation*

We defined a simulated animal mover based on an SSF framework (Potts & Borger 2023). We

defined "true" environmental suitability by combining four rasters of random, spatially-autocorrelated

variables using four random adjustment parameters sampled from a uniform distribution (Fig. 3A, SI Fig.

1, SI Fig. 2). The movement model also included negative selection for longer step lengths and positive

selection for similar turn angles across steps. We simulated 10,000 initial steps (in-sample) and then a

set of 500,000 steps with random restarts every 1,000 locations (out-of-sample), with initial locations

sampled based on the environmental suitability surface. The dataset of 500,000 locations generated a

map of spatial intensity of use (Fig. 3B). We converted the dataset of 10,000 steps into a movement

track, jittered locations very slightly to create a continuous distribution of step lengths, and sampled 40

random steps per observed step using the "amt" package (Fig. 3C, SI Fig. 3) (Signer *et al.* 2019). We

converted the observed gamma distribution of step lengths into a wider gamma distribution with a

larger mean; specifically, we increased the shape and decreased the scale of the gamma distribution by

a factor of two. This increased random sampling of step distances that were both very close and very far

based on the animal mover (SI Fig. 4). Accordingly, we fit an SSF treating the step-terminus values of the

four input rasters as predictors in the step selection function, and we included a polynomial for step

length as a predictor using the "amt" package (SI Table 2) (Signer *et al.* 2019). For simplicity, in all

analyses below, we did not model turn angle in all SSF models. We used the fitted SSF model to

parametrize *A* and generate a vector of stable-state distributions as described above (SI Fig. 5, SI Fig. 6).

We used "amt" package to fit an RSF using package defaults and the four input rasters as

covariates (SI Table 2) (Signer *et al.* 2019). We also used the mean step length to predict the selection

probability for all cells in the landscape using a random cell in the landscape as a baseline from the fitted

SSF above. We then compared the error in predictions in two ways. First, we calculated the root-mean-

squared error (RMSE) between the input environmental suitability and the simulated out-of-sample

intensity of use, the SSD probability, the RSF kernelized probability of use, and the SSF kernelized

relative selection strength. Second, we calculated the RMSE between the simulated out-of-sample

intensity of use and the SSD, RSF, and SSF predictions.

*Case Study #1 - Intra-individual prediction*

To test predictive accuracy within individuals, we used a red deer dataset for a single individual

in the "amt" package to generate SSD, RSF, and SSF predictions across a raster of distance to forest

(calculated from a binary forest raster provided in the "amt" package, aggregated to a resolution of

100m by 100m). We separated the first (chronologically) 75% of observed data as an in-sample training

dataset and retained the latter 25% of observed data as an out-of-sample testing dataset. As above, we

sampled random steps by widening the observed gamma distribution of step lengths by a factor of two

with 40 random points per observed step. We included distance to forest as a covariate in models along

with a polynomial of step length in SSF models (SI Table 2, SI Fig. 7). We generated the stable-state

distribution as above using a neighborhood defined by the 95$^{th}$ percentile of observed step lengths. We

then generated the RSF and SSF probabilities as above and aggregated each of the three predicted

surfaces into centiles (100 bins). We tallied the proportion of out-of-sample testing locations that fell in

each centile of the predicted surfaces, and calculated Spearman's rank correlation coefficient, rho, for

the relationship between stable-state distribution, RSF, and SSF predictions vs. real use intensity. This is

a common model check for RSF predictions (Morris *et al.* 2016), and it can be interpreted as the stronger

the monotonic relationship between the percentiles of predicted use and use intensity, the better a

given prediction captures the biological intensity of use (even if not linear). To ensure that comparisons

were not sensitive to the underlying out-of-sample datasets, we bootstrapped the observed set of GPS

points (sampling with replacement) 100 times, generating 100 estimates of Spearman's rho per

predicted surface.

*Case Study #2 - Inter-individual predictions*

To demonstrate the predictive utility of SSD across individuals, we used a previously published

roe deer dataset for five individuals in Northern Italy (Cagnacci *et al.* 2011) to assemble the same SSD,

SSF, and RSF predictions. We resampled tracks to 4 hours (the median interval in GPS transmissions),

and as above, we isolated the first 75% of individual movements as in-sample, and the next 25% as out-

of-sample testing datasets. Because the dataset contained migratory movements, we decided to model

only summer movements of deer (defined as Julian day 120-290, SI Fig. 11). We included distance to

forest (calculated from 2006 CORINE landcover classes of conifer, mixed, and broadleaf forests) and

slope (calculated from STRM elevation models, aggregated at a scale of 129m by 185.2m) as covariates

in our models and predictions (SI Fig. 12) with a polynomial of step length, while the RSF only included

the square-root of distance to forest and slope (SI Table 2). We performed the same out-of-sample

evaluations as above with 100 bootstrapped Spearman rank correlations between surface centiles and

observed use intensity in 25% out-of-sample for individuals. Additionally, we used individual-level

models (fit using 75% of individual data) to predict the distributions of all other individuals' spatial use

intensity in the dataset and assessed the quality of using the same bootstrapping method (referred to as

"between individual" predictions).

*Case Study #3 - Prediction to novel contexts*

Lastly, to explore the predictive capacity of the SSD method to novel landscapes, we used a dataset of

eight fisher from Northern New York, USA (SI Fig. 16) (LaPoint *et al.* 2013) to generate stable-state

distribution, RSF, and SSF probabilities considering a landscape with square-root distance to forests or

wetlands and square-root human population density (Falcone 2016)(SI Table 1). Seven fisher inhabited a

similar environmental extent with a mix of forested and human landscapes (approximately 10km by

15km, SI Fig. 17), whereas one fisher occupied a completely different landscape approximately 30km

away from the others and with far less human infrastructure and population density (SI Fig. 22). We

regarded the seven local fisher as focal individuals for modeling, and we retained the non-local fisher as

a unique opportunity to evaluate individual-level predictive performance among environmental

contexts. We calculated distance to forest using the National Landcover Database from 2006 to identify

land classes which were wetland and deciduous, mixed, and evergreen forest. We dissolved the

resulting raster to generate a distance to forest or wetland using QGIS (v. 3.22) and GDAL proximity

analysis for computational efficiency before aggregating spatially to 91.2m by 123.2m resolution.

Because individuals in the fisher dataset had different GPS fix-rates, we resampled tracks to the median

fix rate of each individual (range: 2-15 minutes). We performed the same out-of-sample evaluations for

within individual (75/25 split) and between individual comparisons (one individual predicting the

distribution of all others) for all seven focal fishers using RSF, SSF, and SSD surfaces. We then performed

a novel comparison of each of the seven focal fisher movement models to predict the spatial

distribution of the non-local fisher in the novel environmental context, again using 100 bootstraps of the

test dataset to assess sensitivity of surface predictions.

The code used to generate this paper is available online (https://github.com/will-

rogers/StableStateSSF.git) and is provided as compiled document in an online supplement to the main

text. All code will be archived on Zenodo upon acceptance.

**Results:**

*Simulation*

We found that the kernelized surface of the relative strength of selection was almost perfectly

linearly related to the input environmental suitability surface (Pearson's $r$ = 0.997), a pattern supported

by the fact that the SSF surface most closely matched (by an order of magnitude) the input spatial

patterns of environmental suitability (RMSE = $7.57 \times 10^{-06}$) as compared to SSD (RMSE = $9.57 \times 10^{-05}$, $r$ =

0.934) and RSF surfaces (RMSE = 6.33x10$^{-05}$, $r$ = 0.931, Fig. 3G). Observed patterns of out-of-sample

space use intensity had a relatively large error in predictions of input environmental suitability (RMSE

=7.84x10$^{-05}$, $r$ = 0.950). These patterns of error were also non-linear (Fig. 3G), with the SSD surface and

the simulated intensity of use strongly departing from a 1:1 relationship. The SSD predictions (RMSE =

4.15x10$^{-05}$,  Fig. 3H) were almost twice as accurate as the SSF surface (RMSE = 8.13x10$^{-05}$) in predicting

the true simulated distribution of use intensity. The RSF surface lagged behind the best predicting model

(RMSE = 8.28x10$^{-05}$). We found that the SSF predictions strongly departed from a 1:1 relationship, while

the SSD predictions were now linear with the simulated intensity of use surface (Fig. 3H).

*Case Study #1 - Intra-individual prediction*

As in the simulation, the general patterns between surfaces appeared similar, but there was far

greater variation in spatial intensity of predicted use in the SSD surface (Figs. 4A-C, SI Figs. 8-10). We

found that the SSD surface for the red deer case study was best correlated with the observed use of

intensity for modeled data and out-of-sample predictions for the last 25% of the individual's movement

(Fig. 5A-B). Using only a single predictor in the model (distance to forest), the SSF and RSF surfaces were

similar after kernelizing with weaker correlations with modeled and the out-of-sample locations (Fig. 5A-

B).

*Case Study #2 - Inter-individual predictions*

We compared the RSF, SSF, and SSD surfaces in their ability to predict the future distributions of

individual roe deer as well as the distributions of other individuals in the dataset (Figs. 4D-F, SI Figs. 13-

15). For the predictions on modeled and out-of-sample predictions on the next 25% of chronological

observations, we found that two individuals were best predicted by SSF surfaces, while the majority

(three of five) were best predicted by SSD surfaces (Fig. 5C-D). This pattern was even more stark when

comparing the ability for individuals to predict the distribution of others in the dataset. Only one

individual was better predicted by the RSF surface, while four of five were best predicted by the SSD

surface (Fig. 5E).

*Case Study #3 - Prediction to novel contexts*

We used the same frameworks as above to assess the ability for RSF, SSF, and SSD surfaces to

predict modeled data, the next 25% of individual data, and other individuals in the dataset (SI Figs. 18-

20). We also compared the ability for individual models to make predictions in a novel environmental

context (SI Figs. 23-25). We found that, as in previous examples, most individual movements of modeled

data and out-of-sample predictions within individuals were best predicted by SSD surfaces (six of seven)

as compared to RSF (zero of seven) and SSF surfaces (one of seven, Fig. 5F-G). We also found that the

SSD surface of all seven individuals was better able to predict both local (Fig. 5H) and non-local space

use by other individuals in the dataset, as compared to the RSF and SSF surfaces (Fig. 5I). In fact, the RSF

and SSF surfaces were uniformly *negatively* correlated with the non-focal individual's space use in the

non-local context, while the SSD surface was positively correlated with observed space use in a novel

environmental context (Fig. 5I).

**Discussion**

Predicting animal distribution at landscape scales on the basis of organism-environment

interactions is an active area of research that informs both basic ecological inquiry as well as applied

work informing conservation, management, and global policy. We highlight the simultaneous integration

of environmental preference, movement capacity, and spatial arrangement of habitats as an important

outstanding challenge for current frameworks. In simulations and analyses of three biological datasets,

we found that the SSD approach was able to recover patterns in animal distribution that were not

captured by occurrence (RSF) or SSF frameworks. We found that occurrence-based methods, though

commonly used, can generate poor inferences about how animals are distributed in landscapes, despite

accurately capturing then nature of environment-organism relationships. Conversely, movement models

like SSFs have been proposed as a more mechanistic approach to understand these interactions, yet we

found that SSFs can indeed fail to capture patterns of organism distribution in landscapes when used

improperly (i.e., making unconditional or arbitrarily conditioned predictions across landscapes). While

SSFs can accurately define fine-scale selection paradigms (e.g., SSFs can nearly perfectly capture habitat

preference parameters, Fig. 3G), we found that the incorporation of movement mechanisms was

necessary to generate effective predictions of animal distribution (Fig. 3H). We propose that by

combining the fine-scale applicability of SSFs with ergodic frameworks, such as SSD, to capture scale-

dependent movement, we can generate high-resolution and accurate depictions of organismal

distributions in space (Fig. 1). In sum, these findings suggest that local mechanisms of organismal

interactions with their environment can be better generalized to novel spatial extents and unobserved

organisms than occurrence-based frameworks or frameworks that rely on the application of local

movement rules divorced of movement mechanisms.

Our approach builds upon previous frameworks and methods, but we introduce a new way of

using SSFs to parametrize transition matrices and incorporate distance-weighted movement

probabilities. For example, transition matrices are the currency underlying frameworks like *circuitscape*

(McRae *et al.* 2008), and are fundamental to more recent movement frameworks (Fletcher *et al.* 2019).

Similarly, it is recognized that SSF output can capitulate animal distributions through simulations from

SSF models, not from direct predictions of SSF models over an entire landscape (Potts & Borger 2023).

The advance presented here is that ergodic frameworks provide a convenient way to generate

simulations of SSF output assuming random starts in all possible landscape cells. Similarly, by using SSFs

to parametrize transition matrices and incorporate distance-weighted movement probabilities, animal

movement models can generate sparse transition matrices that are less assumptive than previous

methods used to generate transition matrices (McRae *et al.* 2008; Fletcher *et al.* 2019), and are

quantitatively more specific, directional, and biologically reasonable.

Mechanistic models can produce predictions of patterns that are possibly less affected by bias

than inferences and predictions based on patterns. Animal movement and step-selection can be

effective mechanistic models of animal-environment interactions, and we demonstrate how these

mechanisms can generate strong predictions of emergent animal distributions in landscapes.

Conceptually, we propose that SSF and movement modeling in general is one of the best available ways

to use remotely-sensed behavior to understand organism-centric mechanisms - like organism-

environment interactions - that empower predictions of emergent phenomena.

Movement ecology may serve as an effective link between mechanism-focused subfields of

behavior and physiology with pattern-focused subfields like biogeography. Harnessing the mechanisms

implicit in remotely-sensed information offers a tantalizing opportunity to merge within-organism

processes with their manifestation at grander scales in space. Importantly, this framework also removes

potentially fraught questions about how long to run SSF simulations and from where in landscapes to

initiate simulations.

Still, there remain limitations to the framework proposed above that would be fruitful avenues

for future work. One that may be immediately apparent to movement modelers is the lack of turn angle

inclusion in our ergodic framework. This is principally a computational issue. The transition matrix

specifies movement between states (cells) in the environment. If we were to also include how the turn

angle of entrance into a cell as a state dependency, it is not clear how to incorporate such a continuous

variable in state definitions. This is because incorporating turn angles (no matter how biologically

relevant) breaks the Markovian character of SSFs and, in doing so, offers conceptual and computational

complications. We suggest that SSFs be fit so that each step is only dependent on the current conditions

of the step in the above framework. We note that step distance is often heavily correlated with turn

angle (it is less possible to move large distances in consecutive steps with a high turn angle), and step

distance may effectively capture much of the turn angle signal. However, this is certainly an over-

generalization and this phenomenon would benefit from a more quantitative investigation. We

recognized that the landscape prediction from SSFs was an effective diagnostic to test the predictions

from an SSF in comparison to our biological expectations of animal distribution. We argue that this

framework may offer an important ability to assess SSF fit, a currently nebulous endeavor. There are

also many extensions of stochastic matrices for time-dependent predictions, connectivity modeling, and

use with environmentally variable survival (Fletcher 2019). Treating SSFs as ways to parametrize

transition matrices offers exciting opportunities for extensions in movement ecology and applications of

organismal processes to larger scales of predictions.

Acknowledgments:

We thank Johannes Signer, Francesca Cagnacci, and Scott LaPoint for their generosity in providing the

red deer, roe deer, and fisher datasets (respectively) as open-access resources for the scientific

community, and for their permission to use those datasets in our study. We also thank John Fieberg and

members of the Jetz and Ezenwa labs for their generous feedback on early versions of the project.

**Figures**

**Figure 1: Fundamental preferences do not equal realized use**

(**A**) The theoretical frequency of use based on the preference of a single-dimensional

environment (red) vs. the realized frequency of use given some movement constraint (blue). (**B**)

The visualization of the underlying preference surface used in simulations minus the realized

frequency of use from simulations with selection against longer steps. (**C**) Theoretical and (**D**)

empirical depictions of heterogeneity for the frequency of use of individual patches or locations

within a landscape with (blue) and without movement constraints (red). (**E**) In effect, movement

constraints should be predicted to amplify underlying patterns of landscape heterogeneity.

**Figure 2:** (**A**) Animal movements accompanied by environmental contexts, distilled into a

landscape made of cells, $G$. (**B**) Animal locations can be distilled into use vs. available

frameworks conditional on the landscape, suitable for resource selection analyses (RSFs). (**C**)

Alternatively, tracking data can be converted into use vs. available frameworks conditional on

steps, suitable for step selection analyses (SSFs). (**D**) RSFs predict the probability of use over a

landscape, which can be kernelized to return (**I**) probabilities of use over $G$. (**E**) SSFs predict the

relative probability between comparisons within a context, a distinct metric from the probability

of use. (**F**) It is common to make an arbitrary comparison between some either observed or

unobserved environmental contexts in $G$ to all possible contexts in $G$. (**J**) These spatial relative

risks are then kernelized, representing something akin to RSF output, though mathematically

inequivalent. (**G**) Harnessing the ability of SSFs to parametrize local movement patterns, we

can algebraically solve the probability of moving within a set of local environmental contexts ($G_j$)

by defining that the relative selection strength for a focal cell ($G_i$) compared to itself must equal

1. (**H**) Aggregating patterns of in- and out-selection over all neighborhoods of all $G_i$ resolves a

sparse stochastic matrix for all reasonable movements. (**K**) Solving for the principal eigenvector

of the stochastic matrix returns the stable state distribution - the probability of ending in some $G_i$

given all possible starts over the landscape, $G$.

**Figure 3:** (**A**) Combining four random, spatially-autocorrelated rasters to generate a landscape

of "suitable" environmental contexts ($G_i$). (**B**) Simulating 500 random animal tracks (with

constraints on step length and turn angle) over this landscape for 1,000 steps returns intensity

of use with similar patterns as the input "suitability", but with different intensities of use. (**C**) We

simulated 10,000 steps for a single animal using the same input surface to fit a resource

selection function (RSF) and a step-selection function (SSF). (**D**) Kernelized RSF probability of

use and (**E**) kernelized SSF relative selection strength mapped over the landscape. (**F**) Using

the SSF model, we estimated local movements, compiled those probabilities into a stochastic

matrix, and solved the matrix for the principal eigenvector, resolving the stable state distribution

of use over the landscape (SSD). (**G**) Comparison of the input "suitability" vs. observed use

intensity from simulations, SSD, SSF, and RSF outputs, with a sampling of observed points (for

plotting efficiency), a generalized additive model fit (GAM, with all data), and the root-mean

square error (RMSE) between the input and four other surfaces. (**H**) Comparison of the

observed use intensity from simulations vs. SSD, SSF, and RSF outputs, with a sampling of

observed points (for plotting efficiency), GAM fit (with all data), and RMSE between the input

and three other surfaces. We highlighted the minimum error surface in each scenario and

underlined the error in the observed surface to highlight how divergent expectations are from

simulated use. Spatial predictions are visualized untransformed and with the same color scale

to highlight differences in intensity.

**Figure 4:** (**A-C**) Spatial predictions of "use" (as a kernel over the landscape) for red deer, (**D-F**)

summer use of roe deer, (**G-I**) and fisher datasets based on (**A,D,G**) output from a resource

selection function (RSF), (**B,E,H**) output from a step-selection function (SSF), and (**C,F,I**) the

stable state distribution (SSD) from local movement patterns estimated from the SSF. Spatial

predictions for each species are visualized untransformed and with the same color scale to

highlight differences in intensity.

**Figure 5:** (**A,B**) Median (point) and 95% mass distributions (ranges) for 100 bootstraps of

Spearman's rho coefficients between centiles of predicted use surfaces and proportion of true

GPS fixes, where bootstraps are performed by resampling modeled and out-of-sample data with

replacement for red deer (*Cervus elaphus*), (**C-E**) roe deer (*Capreolus capreolus*), and (**F-I**)

fisher (*Pekania pennanti*) datasets. In four set of tests, we used the first (chronologically) 75% of

observed locations for individuals to predict (**A,C,F**) the modeled data ("In-Sample"), (**B,D,G**) the

next (chronologically) 25% of observed locations per individual ("Within Individual"), (**E,H**) all

observed locations for the other individuals in a dataset within the same geographic context

("Between Individuals"), and (**I**) to predict all observed locations for an individual in a completely

different environmental context than the individuals modeled ("Between Contexts").

References

Alston, J. M., C. H. Fleming, R. Kays, J. P. Streicher, C. T. Downs, T. Ramesh, B. Reineking, and J. M.

Calabrese. 2023. Mitigating pseudoreplication and bias in resource selection functions with

autocorrelation-informed weighting. Methods in Ecology and Evolution **14**:643-654.

Avgar, T., S. R. Lele, J. L. Keim, and M. S. Boyce. 2017. Relative Selection Strength: Quantifying effect size

in habitat- and step-selection inference. Ecol Evol **7**:5322-5330.

Avgar, T., J. R. Potts, M. A. Lewis, and M. S. Boyce. 2016. Integrated step selection analysis: Bridging the

gap between resource selection and animal movement. Methods in Ecology and Evolution

**7**:619-630.

Boyce, M. S., and L. L. McDonald. 1999. Relating populations to habitats using resource selection

functions. Trends in ecology & evolution **14**:268-272.

Boyce, M. S., P. R. Vernier, S. E. Nielsen, and F. K. A. Schmiegelow. 2002. Evaluating resource selection

functions. Ecological Modelling **157**:281-300.

Bozinovic, F., D. A. Bastías, F. Boher, S. Clavijo-Baquet, S. A. Estay, and M. J. A. Jr. 2011. The Mean and

Variance of Environmental Temperature Interact to Determine Physiological Tolerance and

Fitness. Physiological and Biochemical Zoology **84**:543-552.

Buckley, L. B., S. A. Waaser, H. J. MacLean, and R. Fox. 2011. Does including physiology improve species

distribution model predictions of responses to recent climate change? Ecology **92**:2214-2221.

Calabrese, J. M., G. Certain, C. Kraan, and C. F. Dormann. 2014. Stacking species distribution models and

adjusting bias by linking them to macroecological models. Global Ecology and Biogeography

**23**:99-112.

Calabrese, J. M., C. H. Fleming, and E. Gurarie. 2016. ctmm: an r package for analyzing animal relocation

data as a continuous-time stochastic process. Methods in Ecology and Evolution **7**:1124-1132.

Creel, S., G. Spong, M. Becker, C. Simukonda, A. Norman, B. Schiffthaler, and C. Chifunte. 2019.

Carnivores, competition and genetic connectivity in the Anthropocene. Pages 1-8  Scientific

reports.

Cushman, S. A., and J. S. Lewis. 2010. Movement behavior explains genetic differentiation in American

black bears. Landscape Ecology **25**:1613-1625.

Dingemanse, N. J., A. J. N. Kazem, D. Réale, and J. Wright. 2010. Behavioural reaction norms: animal

personality meets individual plasticity. Trends in ecology & evolution **25**:81-89.

Evans, T. G., S. E. Diamond, and M. W. Kelly. 2015. Mechanistic species distribution modelling as a link

between physiology and conservation. Conservation Physiology **3**.

Fieberg, J., J. Signer, B. Smith, and T. Avgar. 2021. A 'How to' guide for interpreting parameters in

habitat-selection analyses. J Anim Ecol **90**:1027-1043.

Fletcher Jr, R. J., J. A. Sefair, N. Kortessis, R. Jaffe, R. D. Holt, E. P. Robertson, S. I. Duncan, A. J. Marx, and

J. D. Austin. 2022. Extending isolation by resistance to predict genetic connectivity. Methods in

Ecology and Evolution **13**:2463-2477.

Grimm, V., and S. F. Railsback. 2005. Princeton University Press, Princeton.

JOHNSON, C. J., S. E. NIELSEN, E. H. MERRILL, T. L. McDONALD, and M. S. BOYCE. 2006. Resource

Selection Functions Based on Use-Availability Data: Theoretical Motivation and Evaluation

Methods. The Journal of Wildlife Management **70**:347-357.

Kareiva, P., A. Mullen, S. R. Southwood, M. P. Hassell, and R. M. May. 1990. Population dynamics in

spatially complex environments: theory and data. Philosophical Transactions of the Royal

Society of London. Series B: Biological Sciences **330**:175-190.

Kays, R., M. C. Crofoot, W. Jetz, and M. Wikelski. 2015. Terrestrial Animal Tracking as an Eye on Life and

Planet. Science.

Kearney, M., and W. Porter. 2009. Mechanistic niche modelling: combining physiological and spatial

503           data to predict species' ranges. Ecology Letters **12**:334-350.

504    Keating, K. A., and S. Cherry. 2004. USE AND INTERPRETATION OF LOGISTIC REGRESSION IN HABITAT-

505           SELECTION STUDIES. The Journal of Wildlife Management **68**:774-789.

506    Mackey, B. G., and D. B. Lindenmayer. 2001. Towards a hierarchical framework for modelling the spatial

507           distribution of animals. Journal of Biogeography **28**:1147-1166.

508    Madin, J. S., M. O. Hoogenboom, and S. R. Connolly. 2012. Integrating physiological and biomechanical

509           drivers of population growth over environmental gradients on coral reefs. Journal of

510           Experimental Biology **215**:968-976.

511    Manly, B., L. McDonald, D. L. Thomas, T. L. McDonald, and W. P. Erickson. 2007. Resource selection by

512           animals: statistical design and analysis for field studies. Springer Science & Business Media.

513    McRae, B. H., B. G. Dickson, T. H. Keitt, and V. B. Shah. 2008. Using circuit theory to model connectivity

514           in ecology, evolution, and conservation. Ecology **89**:2712-2724.

515    Menge, B. A., and A. M. Olson. 1990. Role of scale and environmental factors in regulation of community

516           structure. Trends in ecology & evolution **5**:52-57.

517    Merow, C., M. J. Smith, T. C. Edwards Jr, A. Guisan, S. M. McMahon, S. Normand, W. Thuiller, R. O.

518           Wüest, N. E. Zimmermann, and J. Elith. 2014. What do we gain from simplicity versus complexity

519           in species distribution models? Ecography **37**:1267-1281.

520    Moorcroft, P. R., and A. Barnett. 2008. Mechanistic home range models and resource selection analysis:

521           A reconciliation and unification. Ecology **89**:1112-1119.

522    Morris, L. R., K. M. Proffitt, and J. K. Blackburn. 2016. Mapping Resource Selection Functions in Wildlife

523           Studies: Concerns and Recommendations. Appl Geogr **76**:173-183.

524    Mueller, T., K. A. Olson, G. Dressler, P. Leimgruber, T. K. Fuller, C. Nicolson, A. J. Novaro, M. J. Bolgeri, D.

525           Wattles, S. DeStefano, J. M. Calabrese, and W. F. Fagan. 2011. How landscape dynamics link

individual- to population-level movement patterns: a multispecies comparison of ungulate

relocation data. Global Ecology and Biogeography **20**:683-694.

Newbold, T., T. Reader, A. El-Gabbas, W. Berg, W. M. Shohdi, S. Zalat, S. B. El Din, and F. Gilbert. 2010.

Testing the accuracy of species distribution models using species records from a new field

survey. Oikos **119**:1326-1334.

Northrup, J. M., E. Vander Wal, M. Bonar, J. Fieberg, M. P. Laforge, M. Leclerc, C. M. Prokopenko, and B.

D. Gerber. 2022. Conceptual and methodological advances in habitat-selection modeling:

guidelines for ecology and evolution. Ecological Applications **32**.

Phillips, S. J., M. Dudík, J. Elith, C. H. Graham, A. Lehmann, J. Leathwick, and S. Ferrier. 2009. Sample

selection bias and presence-only distribution models: implications for background and pseudo-

absence data. Ecological Applications **19**:181-197.

Post, D. M., and E. P. Palkovacs. 2009. Eco-evolutionary feedbacks in community and ecosystem ecology:

interactions between the ecological theatre and the evolutionary play. Philosophical

Transactions of the Royal Society B: Biological Sciences **364**:1629-1640.

Potts, J. R., and L. Borger. 2023. How to scale up from animal movement decisions to spatiotemporal

patterns: An approach via step selection. J Anim Ecol **92**:16-29.

Powers, R. P., and W. Jetz. 2019. Global habitat loss and extinction risk of terrestrial vertebrates under

future land-use-change scenarios. Nature Climate Change **9**:323-329.

Roughgarden, J. 1975. A Simple Model for Population Dynamics in Stochastic Environments. The

American Naturalist **109**:713-736.

Sastre, P., and J. M. Lobo. 2009. Taxonomist survey biases and the unveiling of biodiversity patterns.

Biological Conservation **142**:462-467.

Scharf, H. R., M. B. Hooten, D. S. Johnson, and J. W. Durban. 2018. Process convolution approaches for

modeling interacting trajectories. Environmetrics **29**.

Schick, R. S., S. R. Loarie, F. Colchero, B. D. Best, A. Boustany, D. A. Conde, P. N. Halpin, L. N. Joppa, C. M.

McClellan, and J. S. Clark. 2008. Understanding movement data and movement processes:

Current and emerging directions. Ecology Letters **11**:1338-1350.

Signer, J., J. Fieberg, and T. Avgar. 2019. Animal movement tools (amt): R package for managing tracking

data and conducting habitat selection analyses. Ecology and evolution **9**:880-890.

Sih, A. 2013. Understanding variation in behavioural responses to human-induced rapid environmental

change: a conceptual overview. Animal Behaviour **85**:1077-1088.

Spiegel, O., S. T. Leu, C. M. Bull, and A. Sih. 2017. What's your move? Movement as a link between

personality and spatial dynamics in animal populations. Ecology Letters **20**:3-18.

Stillman, R. A., S. F. Railsback, J. Giske, U. Berger, and V. Grimm. 2014. Making Predictions in a Changing

World: The Benefits of Individual-Based Ecology. BioScience **65**:140-150.

van der Vaart, E., A. S. A. Johnston, and R. M. Sibly. 2016. Predicting how many animals will be where:

How to build, calibrate and evaluate individual-based models. Ecological Modelling **326**:113-123.

Van Moorter, B., C. M. Rolandsen, M. Basille, and J.-M. Gaillard. 2016. Movement is the glue connecting

home ranges and habitat selection. Journal of Animal Ecology **85**:21-31.

Wisz, M. S., and A. Guisan. 2009. Do pseudo-absence selection strategies influence species distribution

models and their predictions? An information-theoretic approach based on simulated data. BMC

Ecology **9**:8.

Woodin, S. A., T. J. Hilbish, B. Helmuth, S. J. Jones, and D. S. Wethey. 2013. Climate change, species

distribution models, and physiological performance metrics: predicting when biogeographic

models are likely to fail. Ecology and evolution **3**:3334-3346.

1.

Avgar, T., Lele, S.R., Keim, J.L. & Boyce, M.S. (2017). Relative Selection Strength: Quantifying effect size in habitat- and step-selection inference. *Ecol Evol*, 7, 5322-5330.

2.

Avgar, T., Potts, J.R., Lewis, M.A. & Boyce, M.S. (2016). Integrated step selection analysis: Bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, 7, 619-630.

3.

Boyce, M.S. & McDonald, L.L. (1999). Relating populations to habitats using resource selection functions. *Trends in Ecology & Evolution*, 14, 268-272.

4.

Boyce, M.S., Vernier, P.R., Nielsen, S.E. & Schmiegelow, F.K.A. (2002). Evaluating resource selection functions. *Ecological Modelling*, 157, 281-300.

5.

Bozinovic, F., Bastías, D.A., Boher, F., Clavijo-Baquet, S., Estay, S.A. & Jr., M.J.A. (2011). The Mean and Variance of Environmental Temperature Interact to Determine Physiological Tolerance and Fitness. *Physiological and Biochemical Zoology*, 84, 543-552.

6.

Buckley, L.B., Waaser, S.A., MacLean, H.J. & Fox, R. (2011). Does including physiology improve species distribution model predictions of responses to recent climate change? *Ecology*, 92, 2214-2221.

7.

Cagnacci, F., Focardi, S., Heurich, M., Stache, A., Hewison, A.J.M., Morellet, N. *et al.* (2011). Partial migration in roe deer: migratory and resident tactics are end points of a behavioural gradient determined by ecological factors. *Oikos*, 120, 1790-1802.

8.

Colwell, R.K. & Rangel, T.F. (2009). Hutchinson's duality: The once and future niche. *Proceedings of the National Academy of Sciences*, 106, 19651-19658.

9.

Creel, S., Spong, G., Becker, M., Simukonda, C., Norman, A., Schiffthaler, B. & Chifunte, C. (2019). Carnivores, competition and genetic connectivity in the Anthropocene. *Scientific Reports*, 9, 1-8.

Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J. & Ferrier, S. (2009).

    Sample selection bias and presence-only distribution models: implications for

    background and pseudo-absence data. *Ecological Applications*, 19, 181-197.

38.

Post, D.M. & Palkovacs, E.P. (2009). Eco-evolutionary feedbacks in community and ecosystem

    ecology: interactions between the ecological theatre and the evolutionary play.

    *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364, 1629-1640.

39.

Potts, J.R. & Borger, L. (2023). How to scale up from animal movement decisions to

    spatiotemporal patterns: An approach via step selection. *J Anim Ecol*, 92, 16-29.

40.

Powers, R.P. & Jetz, W. (2019). Global habitat loss and extinction risk of terrestrial vertebrates

    under future land-use-change scenarios. *Nature Climate Change*, 9, 323-329.

41.

Pulliam, H.R. (2000). On the relationship between niche and distribution. *Ecology Letters*, 3,

    349-361.

42.

Roughgarden, J. (1975). A Simple Model for Population Dynamics in Stochastic Environments.

    *The American Naturalist*, 109, 713-736.

43.

Sastre, P. & Lobo, J.M. (2009). Taxonomist survey biases and the unveiling of biodiversity

    patterns. *Biological Conservation*, 142, 462-467.

44.

Schick, R.S., Loarie, S.R., Colchero, F., Best, B.D., Boustany, A., Conde, D.A. *et al.* (2008).

    Understanding movement data and movement processes: Current and emerging

    directions. *Ecology Letters*, 11, 1338-1350.

45.

Signer, J., Fieberg, J. & Avgar, T. (2019). Animal movement tools (amt): R package for

    managing tracking data and conducting habitat selection analyses. *Ecology and*

    *Evolution*, 9, 880-890.

46.

Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species