1 **Title**
2 Statistical design of a synthetic microbiome that suppresses diverse gut pathogens
3
4 **Authors**

5 Rita A. Oliveira[1,7], Emma McSpadden[1], Bipul Pandey[1,9], Kiseok Lee[2], Mahmoud Yousef[2], Robert
6 Y. Chen[3], Conrad Triebold[4], Fidel Haro[5], Valeryia Aksianiuk[6], Riya Patel[8], Kaavya Shriram[9],
7 Ramaswamy Ramanujam[1], Seppe Kuehn[2,10], Arjun S. Raman[1,10,11*]
8
9
Affiliations
[1]Duchossois Family Institute, University of Chicago, Chicago, IL, 60637
[2]Department of Ecology and Evolution, University of Chicago, Chicago, IL, 60637
[3]Department of Psychiatry, University of Washington, Seattle, WA, 98195
[4]Section of Genetic Medicine, University of Chicago, Chicago, IL, 60637
[5]Pritzker School of Medicine, University of Chicago, Chicago, IL, 60637
[6]Institute for Cell Biology, University of Bern, Bern, Switzerland
[7]Department of Medicine, Section of Infectious Disease & Global Health, University of Chicago
Medicine, Chicago, IL, 60637
[8]University of Illinois at Urbana Champagne, Urbana, IL, 61825
[9]Grinnell College, Grinnell, IA, 50112
[10]Center for Physics of Evolving Systems, University of Chicago, Chicago, IL, 60637
[11]Department of Pathology, University of Chicago, Chicago, IL, 60637
23
24 *Correspondence to: araman@bsd.uchicago.edu

Abstract

Engineering functional microbiomes is challenging due to complex interactions between bacteria and their environments[1–6]. Using a set of 848 gut commensal strains and clearance of multi-drug resistant *Klebsiella pneumoniae* (*Kp*-MH258) as a target function, we engineered a functional 15-member synthetic microbiome—SynCom15—through a statistical approach agnostic to strain phenotype, mechanism of action, bacterial interactions, or composition of natural microbiomes. Our approach involved designing, building, and testing 96 metagenomically diverse consortia, learning a generative model using community strain presence/absence as input, and distilling model constraints through statistical inference. SynCom15 cleared *Kp*-MH258 across *in vitro*, *ex vivo*, and *in vivo* environments, matching the efficacy of a fecal microbiome transplant in a clinically relevant murine model of infection. The mechanism of suppression by SynCom15 was related to fatty acid production coupled with environmental acidification. SynCom15 also suppressed other pathogens—*Clostridioides difficile*, *Escherichia coli*, and other *K. pneumoniae* strains—but through different mechanisms. Sensitivity analysis revealed models trained on strain presence/absence captured the statistical structure of pathogen suppression, illustrating that community representation was key to our approach succeeding. Our framework, 'Constraint Distillation', could be a general and efficient strategy for building emergent complex systems, offering a path towards synthetic ecology more broadly.

Main

Engineering communities of microbes for desired functions ('synthetic ecology') is of fundamental importance and holds great practical promise for addressing many problems facing humanity[7–9]. Across environmental, agricultural, and human health applications, designed communities have previously been created for specific functions. Yet despite these case-by-case successes, synthetic ecology remains largely empirical, driven by phenotypic screens, substantial prior knowledge, and context-specific information. In application to the human gut microbiome, such empiricism has typically entered design efforts in three ways. First, by mining large human cohorts or the microbiotas of laboratory mice to extract the natural statistics of existing gut communities (species prevalence and abundance, covariation structure, modules/guilds, compositional structures linked to phenotypes of interest)[10–15]. Second, by using intact or minimally processed natural communities (fecal microbial transplants, donor enrichments, fractionation/serial dilution) to transfer function (i.e. 'top-down' approach)[16–20]. Or third, by assembling defined consortia from individual bacterial strains but with the inclusion of rich biological priors (strain growth characteristics, functional screens of strains, pairwise interaction maps, metabolic capabilities, resource-interaction networks)[21–27].

Why is such knowledge typically used and broadly deemed necessary for engineering consortia? A central reason is complexity. Microbiomes comprise many species whose interactions are non-linear, redundant, and context-dependent[2,10,28,29]. Higher-order effects often defeat logic learned about system function derived from single strains or pairwise interactions between strains[3,6,30]. Thus, the combinatorial complexity of N strains is not additive, but scales as 2^N thereby becoming prohibitively expansive with respect to design efforts: a hypothetical consortium comprised of 100 strains encodes 10^{30} possible strain combinations. In physics and chemistry, engineering principles are scaffolded by 'forward models'—generative theory that maps parts to function in a deterministic fashion[31]. However in biology, such forward models are absent and therefore functional properties cannot be inferred from components alone. Collective

biological behaviors—so-called 'emergent' properties—arise from interaction patterns that are

difficult to observe, let alone encode *a priori*. Thus, lacking a physics of emergence and confronted

with daunting complexity, current bacterial consortia design strategies default to embedding

statistical, ecological, or mechanistic priors. Inclusion of these priors leads to two key limitations

with respect to consortia design: (i) biasing outcomes towards what nature already makes feasible

rather than exploring a truly synthetic space and (ii) resulting in bespoke solutions that exist on a

case-by-case basis rather than transferable design principles that can be broadly employed. Both

limitations place a ceiling on the potential of synthetic ecology with respect to future application

towards desired target functions.

There has been a recent substantial rise in the capacity to culture bacteria and create

strain banks[11,16,21,27,32]. This trend along with the observations regarding consortia design noted

above motivate a natural question of both fundamental and practical importance: are microbiomes

programmable? By 'programmable', we specifically mean two nested capabilities. First, the

capacity to create a reliable map relating microbiome composition to function learned (e.g.

predictive capacity) without the need to specify context-dependent mechanistic or functional

priors[33]. But also second, going beyond prediction, the ability to use that map to build microbiomes

that satisfy a desired target function with performance that holds across environments (i.e.

generative capacity and functional robustness)[10,34]. Taking inspiration from engineering, in circuit

theory existing principles enable the construction of apparently complex, functional, and robust

circuits from a list of parts. Analogously, can design principles be derived that enable construction

of synthetic, functionally robust microbiomes from strains alone?

When forward models are unavailable or limiting, statistical physics has addressed design

by solving the so-called 'inverse problem': inferring constraints that reflect statistical patterns

within observational data alone, then using only these constraints to synthesize new

systems[31,35,36]. Classic examples include the approach of maximum-entropy to learn energy

models from correlation structures in data; more recent examples include using sophisticated

statistical networks (e.g. artificial intelligence-based models) to represent relationships and

generate complex systems like proteins and materials[37–40]. These examples suggest a promising

strategy forward with respect to consortia design. Rather than encode any priors involving strain

phenotype, strain interactions, or mechanism of desired target function, solving the inverse

problem would involve inferring a statistical model of latent constraints residing within an

ensemble of synthetic consortia that is predictive of the target function and is generative. If

successful, this strategy would enable the construction of truly synthetic microbiomes—consortia

that do not resemble natural microbiomes—through establishing design principles that are

statistical in nature (e.g. 'statistical design') and thereby reframe consortia design as a data-driven

engineering problem.

In this work, we engineered a complex 15-member synthetic microbiome—SynCom15—

that suppressed a multi-drug resistant (MDR) pathogenic strain of *Klebsiella pneumoniae* (*Kp*-

MH258) across *in vitro*, *ex vivo*, and *in vivo* conditions, matching the rapidity, efficacy, and

robustness of a whole fecal microbial transplant (FMT) in a specific pathogen-free (SPF) mouse

model of infection. Rather than relying on biological knowledge involving strain phenotype, strain

interactions, natural microbiome composition, or mechanism of suppression to accomplish this

task, we engineered SynCom15 by first solving the inverse problem through statistically learning

a generative model of consortia design *de novo* using an ensemble of just 96 synthetic consortia

spanning various sizes that we evaluated for pathogen suppression, then distilling taxonomic

constraints learned by the model. Critically, the only biological information used in our process

was strain genome sequence for characterizing genomic diversity amongst our strain collection.

Through a series of experiments, we found that SynCom15 (i) suppressed *Kp*-MH258 through

fatty acid production coupled with environmental acidification, (ii) suppressed other pathogenic

strains of *K. pneumoniae*, *Escherichia coli*, and *Clostridioides difficile*, (iii) promoted recovery of

antibiotic-induced dysbiosis *in vivo*, and (iv) was not identifiable from sequencing statistics of

healthy human gut microbiomes. We pose that our statistical approach—'Constraint Distillation'—

may be generally useful for engineering bacterial consortia towards executing desired target

functions.

Results

Suppression of K. pneumoniae by engineered, diverse microbiomes does not translate across

environments

To design a synthetic bacterial consortium that suppresses MDR *K. pneumoniae*, we first

created a strain bank of 848 gut commensals from a set of 28 healthy human volunteers and

whole-genome sequenced all strains (Methods). A subset of these strains have been described

and characterized previously[32,41]. These commensals collectively reflected a breadth of

phylogenetic diversity typically found in human gut microbiomes (**Extended Data Fig. 1**,

Supplementary Table 1A). Without incorporating more information, the total possible design

space encompassed by this strain bank spanning all individual strains, pairs of strains, and higher-

order combinations was $\sim 10^{250}$—a space that is impractical from the standpoint of design.

Moreover, constructing consortia that constitute greater than 50 strains is markedly challenging

from the perspective of experimental culture. We therefore sought to reduce the number of

possible strains we could use to design consortia.

We performed a UMAP-based (Uniform Manifold Approximation and Projection)

dimension-reduction on an alignment of gene content across all strains (Methods). This defined

a latent embedding of strain relatedness that clustered strains by their phylum-level phylogenetic

designation, thereby providing a coarse-grained view of genomic diversity across our strain bank

(**Extended Data Fig. 2**, **Supplementary Table 1B**). We next sought to select a subset of strains

from our original 848 strains that comprised a manageable set from the perspective of engineering

synthetic consortia in the laboratory yet retained the structure of phylogenetic diversity captured

by the UMAP embedding. We therefore selected 46 strains that retained the phylogenetic diversity

of our strain bank—a number of strains that was within the range of previously published designed

consortia (**Fig. 1A**, **Supplementary Table 2**) (Methods)[16,27].

Despite down-sampling our commensal strain bank from 848 strains to 46 strains, the

resulting combinatorial complexity (~35.2 billion possibilities) was still unmanageable. Therefore,

we sought to impose a constraint on the consortia we designed—construct metagenomically

diverse consortia. The rationale behind this choice was to capture a large possible functional

capacity of each designed consortium. Previous studies have constructed diverse consortia by

simply combining phylogenetically diverse strains[11,27]. However, no algorithmic process currently

exists for creating ensembles of diverse consortia spanning various membership sizes from strain

banks. Primarily this is because the hierarchical nature of phylogeny makes defining differences

between strains in a quantitative manner challenging[32]. We therefore developed an algorithm that

constructs consortia directly off the two-dimensional UMAP coordinates shown in **Fig. 1A**

(**Supplemental Discussion**) (Methods). Briefly, for a consortium consisting of N members, the

algorithm proceeds as follows. First, a random seed bacterium is chosen; next, a second

bacterium is chosen such that it is distant from the seed in the UMAP coordinate space, a third

bacterium is chosen such that it is distant from the first two bacteria, and so on until N members

are chosen (**Extended Data Fig. 3A**). This strategy of design contains no biological information

outside of strain genomes for engineering microbial consortia nor any information regarding strain

phenotype or pathogen suppression *a priori*. We employed our design strategy and engineered

96 'Designed Microbial Communities' (DMCs) (**Fig. 1B,C**). The 96 DMCs spanned a size of 2 to

all 46 strains (**Supplementary Table 3A**). We found that the communities comprising greater than

10 strains asymptotically reached a degree of taxonomic diversity represented by the community

containing all 46 strains ('DMC46') (**Extended Data Fig. 3B**). Thus, our algorithm succeeded in

creating an ensemble of metagenomically diverse consortia.

We then tested all DMCs for their capacity to suppress an MDR strain of *K. pneumoniae*—

Kp-MH258—after 120 hours of co-culture in Brain Heart Infusion Media supplemented with

cysteine (BHIS) (**Extended Data Fig. 4A**) (Methods). The MH258 strain was isolated from a

patient at Memorial Sloan Kettering Hospital and was representative of the epidemic multilocus

sequence type (ST) 258 clone harboring the bla_{KPC}-encoded carbapenemase. We chose this

strain to use as our target for suppression because it is amongst the most multi-drug resistant

strains that have been previously characterized, exhibiting resistance against a diversity of

antibiotics[42]. BHIS media was chosen because all 46 gut commensal strains robustly grew in this

environment (**Extended Data Fig. 4B**, **Supplementary Table 3B**). We found that the set of 96

DMCs exhibited a continuous range of suppressive capacities in a reproducibly robust fashion

(**Fig. 1C**, **Extended Data Fig. 4C,D**, **Supplementary Table 3C,D**). During the first 48 hours of

co-culture, the abundance of *Kp*-MH258 remained uniformly high, after which community effects

began to manifest—some consortia suppressed *Kp*-MH258 greater than four orders of magnitude

while others had minimal impact on its fitness (**Extended Data Fig. 4E**, **Supplementary Table**

3D).

We evaluated three different compositional qualities of our DMCs that could be associated

with suppression of *Kp*-MH258. First, previous studies have suggested the importance of

including up to ten strains that are phylogenetically diverse in consortia designed to suppress *K.*

pneumoniae[27]. We found that the suppressive capacity of DMCs was not associated with the

number of species within our diverse consortia (**Extended Data Fig. 5A**). As one example,

DMC17 contained 12 strains and suppressed *Kp*-MH258 four orders of magnitude while DMC92

contained 19 strains and suppressed *Kp*-MH258 less than one order of magnitude (**Extended**

Data Fig. 5B, left). As another example, there were a total of 16 DMCs that contained 14 strains,

but their observed capacity to suppress *Kp*-MH258 varied from one to four orders of magnitude

(**Extended Data Fig. 5B**, right). Second, previous studies have suggested the importance of

including *Escherichia coli* strains in the background of large diverse consortia for suppressing *K.*

pneumoniae due to hypothesized overlap in metabolic niche[27]. Our set of 46 strains contained

one *E. coli* strain. We found no relationship between the membership size of consortia that

contained our *E. coli* strain and the capacity of a consortium to suppress *Kp*-MH258 (**Extended**

Data Fig. 5C). As an example, DMC92 contained an *E. coli* strain plus 18 other strains and

DMC03 contained an *E. coli* strain and two other strains. Despite the difference in consortia size,

both DMCs exhibited similar suppressive capacity of less than one order of magnitude (**Extended**

Data Fig. 5C,D). Finally, we tested whether the presence of any single strain was uniformly

responsible for the suppressive capacity of DMCs. Our results showed that the presence of 14

individual strains was significantly associated with the suppression of *Kp*-MH258 (**Extended Data**

Fig. 6). However, we also found that the presence of any one of these strains on its own did not

guarantee suppression by the DMC. DMCs containing these strains showed a range of

suppression that overlapped extensively with DMCs not containing these strains (**Extended Data**

Fig. 7). Collectively, these results illustrated that previously described taxonomic heuristics—

diversity/richness of consortium, presence/absence *of E.* coli—and presence/absence of single

strains were insufficient to describe our results.

Recent work has suggested that simplistic statistical models that do not consider higher-

order interactions between strains can be successful at predicting community phenotype from the

presence or absence of individual strains. A study that fit simple regression models across a large

number of previously published studies found that additive models alone that do not consider

even pairwise interactions between species can predict community phenotype[33]. Inspired by this

work, we next combined all 14 strains whose individual presence was significantly associated with

Kp-MH258 suppression (see **Extended Data Fig. 6**, yellow box). This consortium was not within

the 96 DMCs we had engineered in **Fig. 1C** and was designed using only the results from our

functional screen. We therefore termed this consortium 'DMC-Rational'. Notably, DMC-Rational

included strains displaying the most extreme regression coefficients derived from a linear

regression model that successfully related *Kp*-MH258 abundance with the additive effect of

individual strains alone (r^2 of ~0.8) (**Supplementary Table 3E**) (Methods). We found that DMC-

Rational suppressed *Kp*-MH258 greater than four orders of magnitude and was the best

performing DMC of all the DMCs comprised of 14 strains (**Fig. 1D**). This result demonstrated that

taking a simplistic statistical approach—constructing a consortium based on the suppressive

effect of each strain considered individually without interaction effects—enabled designing a

consortium that suppressed *Kp*-MH258 in BHIS.

Previous attempts at translating the suppression of *K. pneumoniae* by engineered

bacterial consortia from *in vitro* to *in vivo* settings have demonstrated, at best, limited success. As

an example, recently published work engineering a large diverse consortium of gut symbionts

showed robust *K. pneumoniae* suppression *in vitro* but only mild suppression *in vivo* with

suppression lasting only one day after infection[27]. Moreover, tests of pathogen suppression by

engineered consortium *in vivo* have primarily involved germ-free mouse models of infection—a

model system unrelated to clinically relevant courses of infection by *K. pneumoniae* where

treatment by broad-spectrum antibiotics creates an open niche for pathogen colonization while

patients retain a background microbiome and are subject to colonization by bacteria in the

environment[11,16,27].

Given these observations, we next tested a select group of DMCs for their capacity to

suppress *Kp*-MH258 in a more clinically realistic *in vivo* model of infection. Using a specific

pathogen-free (SPF) mouse model of *Kp*-MH258 infection, we administered several DMCs

spanning a range of suppressive capacities observed in BHIS media. Singly-housed mice were

treated with broad-spectrum antibiotics (metronidazole, neomycin, vancomycin) for four days,

colonized with *Kp*-MH258 on the day after stopping antibiotics, and subsequently gavaged two

days later with either phosphate buffered saline (PBS), a whole-stool fecal microbiota transplant

from mice not given antibiotic treatment (FMT), or a specific DMC (**Fig. 2A**) (Methods). We chose

to evaluate four DMCs—DMC46 (comprised of all 46 strains), DMC96 (comprised of 28 strains),

DMC94 (comprised of 24 strains), and DMC-Rational (comprised of 14 strains)—because they

suppressed *Kp*-MH258 in BHIS media at least four orders of magnitude. We also chose to

evaluate DMC86 (comprised of 20 strains) because this DMC was amongst the worst performing

with respect to pathogen suppression in BHIS media—suppressing *Kp*-MH258 within one order

of magnitude—while containing a similar number of strains as DMC94.

First, our results showed that no consortium suppressed *Kp*-MH258 to the same extent as

an FMT. By three days after gavage of the FMT, fecal abundance of *Kp*-MH258 dropped six orders

of magnitude to 10^4 colony forming units (CFUs) per gram of feces (**Fig. 2B**, **Supplementary**

Table 4A). In contrast, the best performing consortium in BHIS media—DMC46—was three

orders of magnitude worse than the FMT intervention three days after the last gavage. Second,

we found that seven days after the last gavage of the intervention (Day 11 of the experiment), the

suppressive effect of all consortia were within 1.5 orders of magnitude of each other, yielding a

fecal abundance of *Kp*-MH258 between 10^6 and 10^8 CFUs/gram of feces. This result illustrated

that all tested consortia performed in a roughly similar manner to each other within our *in vivo*

model (**Fig. 2B**, **Supplementary Table 4A**). Third, despite suppressing *Kp*-MH258 several orders

of magnitude in BHIS, DMC-Rational most closely resembled the effect of saline intervention on

Kp-MH258 colonization in mice. Thus DMC-Rational lost its suppressive potential when evaluated

in our *in vivo* setting. A systematic analysis of our results showed that the suppressive capacity of

a DMC in BHIS media was not associated with the capacity to suppress *Kp*-MH258 in our *in vivo*

model (**Fig. 2C**, **Supplementary Table 4B**).

Overall, our results showed that using simple heuristics—size/diversity of consortium,

presence of *E. coli*, or the cumulative effect of statistically important individual strains—was

insufficient for engineering a microbiome that suppressed *Kp*-MH258 across different

environments. A parsimonious explanation for these results is that environmental idiosyncrasies

demand bespoke consortia designs, making generally effective consortia inherently difficult to

engineer. However, prior work suggests that translatability can indeed be achieved by identifying

modules of interacting taxa. As a notable example, statistical analysis of human developing

microbiomes led to the identification of a module of interacting species that, when combined into

a consortium, recapitulated key taxonomic dynamics in a porcine model of microbiome

development[10]. More broadly, the idea of modularity has been proposed as a construct for

enhancing robustness across changing contexts in network science, protein biophysics, and

gene-regulatory circuits[34,43–48]. We therefore posed an alternative hypothesis: the capacity to

suppress *Kp*-MH258 across different environments may reside in groups of strains whose joint

effects form a collective unit such that translatable suppression is a property of the interaction

architecture rather than any single taxon or simple rule. Because such groups are not directly

observable from composition alone and because there does not exist obvious priors for encoding

translatability of function into consortia design, we turned to statistical learning as a path forward.

Discovering a 15-member bacterial consortium through machine learning and distillation of

model constraints

Machine-learning (ML) used towards designing synthetic bacterial consortia has typically

incorporated a number of detailed strain qualities including (i) individual growth rates, (ii) growth

rates in co-culture, and (iii) metabolic capacities that were either experimentally determined or

inferred from gene content[22–24,33]. On the other hand, using patterns of strain presence/absence

with regression-based statistics including interactions have accurately predicted ecosystem

structure-function relationships but is limited in detecting greater than pairwise interactions

between strains in the limit of finite sampling that may be critical for going beyond prediction and

towards generative design[33]. As ML models are capable of modeling non-linear interactions, we

reasoned that using ML but considering only the pattern of strain presence/absence across our

DMCs could aid in creating a statistical generative model of consortia design for suppressing *Kp*-

MH258 without requiring the need to consider biological priors.

We trained two models—a Random Forests (RF) ML model and a deep learning-based

neural network (DNN)—using the pattern of strain presence/absence across our 96 DMCs as

input and the CFUs of *Kp*-MH258 after 120 hours of co-culture with a DMC as output (Methods).

Notably, neither model included any information regarding (i) survival of microbes in culture, (ii)

dynamics of bacterial strains, (iii) phenotypic qualities of strains, (iv) nature of interactions

between microbes, or (v) interaction with *Kp*-MH258. Instead, the input to the model was a 46-

dimensional vector defined by the 46 strains that could be used to design a DMC. Each DMC was

characterized by '1's and '0's where a '1' reflected that a specific strain was included in the DMC

and a '0' reflected that a specific strain was left out of the DMC. We observed an in-sample

validation r^2 value of 0.98 for the RF model and 0.74 for the DNN model, indicating well-trained

models (**Supplementary Table 5**).

We next evaluated the capacity of the RF and DNN models to predict 60 new DMCs that

were not included in the original 96 DMCs we constructed. These 60 new DMCs spanned a range

of membership sizes and predicted capacities for *Kp*-MH258 suppression (**Supplementary Table**

6A). We found that both the RF and DNN models accurately predicted the suppressive capacity

of the 60 new DMCs with the RF model mildly outperforming the DNN model (r^2 of 0.6 versus

0.52) where variability in model performance was predictably large at low values of *Kp*-MH258

abundance (**Fig. 3A**, top; **Supplementary Table 6B**). Thus, considering only the pattern of

included strain presence/absence for a given DMC was sufficient to yield generative models of

consortia engineered to suppress *Kp*-MH258 in BHIS media.

We therefore next sought to use these models to design a suppressive consortium.

However, we noted this would be challenging: our ML models exhibited substantial variation in

the prediction of *Kp*-MH258 abundance below 10^5 CFUs because the input data was naturally

limited within this regime (**Fig. 3A**). Moreover, studying the models themselves was not a viable

solution either. While ML models like RF and DNN-based algorithms can be useful for prediction,

their internal logic is opaque, making it difficult to extract actionable design principles or

understand compositionally why particular consortia were effective. With RF models, feature

importance scores are useful for discriminatory tasks but do not capture combinatorial, redundant,

and context-dependent relationships necessary for revealing statistical interactions underlying the

success of model prediction and therefore necessary for generation of a system. In the case of

DNNs, no mapping exists between the architecture of the model and the input dataset since DNNs

are inherently non-linear, opaque transformations.

Given the above considerations, one option would have been to collect data on more

consortia such that the suppressive regime would be better sampled. However, it was not clear

how this could be done. First, simply creating more consortia would likely not have led to sampling

a greater number of suppressive communities. Second, the high variance of the ML predictions

in consortia performance below Kp-MH258 values of 10^5 would likely necessitate testing lots more

consortia. Given these issues, we desired a way to turn noisy ML predictions into actionable

guidance without conducting a large new experimental screen. We therefore posed an alternative

hypothesis: a large ensemble of outputs from our ML models could be leveraged for extracting

new taxonomic information about Kp-MH258 suppression. Our rationale for this hypothesis was

as follows. If predictions are unreliable in the extreme low-abundance regime but are globally

reasonable, then what can be trusted is the relative prediction of the model across many consortia.

Thus, we could use the ML models as rankers—assigning relative rankings such that certain

consortia are placed consistently lower than others—rather than relying on the precise numerical

prediction for a specific consortium. Relatedly, we recognized that any single 'top-ranked'

consortium could be inaccurately predicted. However, aggregation across many independently

top-ranked consortia could potentially reveal taxonomic regularities that are shared. Patterns that

recur across many such consortia would be far more likely to reflect true design signals than

model idiosyncrasies. This approach—(i) broadly query a complex and opaque model, (ii) keep

only the consistent answers, and (iii) summarize those answers into a compact rule set—is

analogous to the logic of 'distillation' in the field of artificial intelligence (AI)—a framework used to

make 'black-box' AI models understandable[49–51]. We therefore adopted the ethos of distillation by

generating a large *in silico* ensemble of consortia, selecting the consortia that were ranked as

strong-suppressors, and performing statistical inference via spectral analysis to extract

compositional constraints (which strains co-occur in association with Kp-MH258 suppression,

which strains are allowable substitutions, which strain combinations should be avoided) (see

Supplementary Discussion for further details). We chose to use our RF model for this approach

as it performed better than the DNN model.

We generated 10^5 DMCs by randomly sampling strain combinations from the 46-strain

pool where the totality of DMCs spanned a range of predicted *Kp*-MH258 suppression values as

determined by the RF model. We chose 10^5 communities to sample greater than 10^3 of the

complexity of the design space as defined by each strain independently (46 total strains

corresponds to a space of 4.6 x 10^4 considering 10^3 oversampling). Each DMC was represented

as a binary vector of strain presence/absence and scored using the RF model, resulting in a matrix

of 10^5 DMCs (rows) by 46 strains (columns) where each DMC was annotated with a predicted

abundance of *Kp*-MH258. We then filtered this design space by DMCs predicted to suppress *Kp*-

MH258 to a value below 10^5 CFUs, yielding a subset of 5,752 DMCs that were predicted to be

high-performing (**Fig. 3B**, **Supplementary Table 7A**). Next, to uncover model constraints on

consortia composition, we performed Singular Value Decomposition (SVD) of this set of DMCs

defined by their patterns of strain presence/absence. This resulted in a spectrum of eigenvectors

that defined axes of variation across DMCs predicted to function particularly well. We associated

the pattern of loadings of DMCs onto each eigenvector with the predicted capacity to suppress

Kp-MH258 for all 5,752 DMCs (Methods) (**Supplementary Table 7B**,**C**). This analysis revealed

that eigenvector 46—encoding <1% of data variance—defined taxonomic variation amongst *in*

silico generated DMCs that was most strongly associated with *Kp*-MH258 suppression (**Extended**

Data Fig. 8). This axis therefore represented a functionally relevant statistical constraint latent

within our RF model.

Two groups of strains contributed similarly onto eigenvector 46: one group of five strains

('Block 1') comprising *Clostridium innocuum*, *Clostridium symbiosum*, *Colinsella aerofaciens*, *E.*

coli, and *Bacteroides xylanisolvens*, and another group of ten strains ('Block 5') comprising

Lacrimispora celerecrescens, Bacteroides caccae, *Blautia faecis*, *Blautia obeum*, *Clostridium*

scindens, a *Bifidobacterium* species, *Megasphaera massiliensis*, *Coprococcus comes*,

Mitsuokella jalaludinii, and *Blautia producta* (**Fig. 3C**) (Methods). Three other blocks of strains

contributed relatively less towards eigenvector 46 ('Block 2', 'Block 3', and 'Block 4') (**Fig. 3C**).

We further observed that the strains belonging to Blocks 1 and 5 co-contributed to eigenvector 46

with each other more than with strains defining Blocks 2, 3, and 4 (**Fig. 3C**, **Supplementary Table**

7D-F). Thus, the 15 strains collectively comprising Blocks 1 and 5 formed the statistically-inferred

putative core within DMCs predicted to suppress *Kp*-MH258—the basis of a designed consortium

that we termed SynCom15 (**Fig. 3C**).

SynCom15 suppresses Kp-MH258 across diverse contexts

391   We evaluated the capacity for SynCom15, as well as Blocks 1 through 5, to suppress *Kp*-

MH258 when co-cultured in BHIS media as well as two *ex-vivo* contexts—media created from the

cecal contents of germ-free mice ('GF cecal content media') and media created from the cecal

contents of SPF mice that were treated with antibiotics for four days then given normal drinking

water for three days ('Ab-SPF cecal content media')—a scenario mimicking our *in vivo* infection

model (**Fig. 4A**) (Methods). The two *ex-vivo* conditions contained information regarding the host,

host diet, and host microbiome, and thus the collection of *in vitro* and *ex-vivo* contexts reflected

vastly different environments for evaluating the suppressive capacity of a consortium on *Kp*-

MH258. We found that of all the consortia tested, SynCom15 was the most suppressive

community across all three environments (**Fig. 4B**, **Supplementary Table 7G**).

401   We next tested the efficacy of SynCom15 in our SPF mouse model of infection (**Fig 4C**).

We found SynCom15 suppressed *Kp*-MH258 five orders of magnitude on the first day after the

third gavage, six orders of magnitude the third day after the third gavage, seven orders of

magnitude the fifth day after the third gavage and maintained a level of *Kp*-MH258 below detection

the seventh day after the third gavage (**Fig. 4D**). Notably, *Kp*-MH258 suppression due to

SynCom15 was more rapid and a greater magnitude than that of an FMT. We compared the

suppressive effect of SynCom15 with that of saline and three other consortia: Block 1, Block 2,

and DMC46. We evaluated Block 1 because it effectively suppressed *Kp*-MH258 in BHIS media

and was identified as consistently comprising a group of strains associated with a suppressive

phenotype towards DMCs by three independent statistical descriptions—(i) the effect of strains

considered individually (see **Extended Data Fig. 6**, yellow box), (ii) the feature importance scores

resulting from the RF model, and (iii) the results of our distillation analysis (**Extended Data Fig.**

9, **Supplementary Table 7H**). We evaluated Block 2 as a plausible negative control as it failed to

suppress *Kp*-MH258 in BHIS, GF cecal content media, and Ab-SPF cecal content media, and this

group of strains were associated with poor suppressive capacity from our distillation analysis. We

re-evaluated DMC46 because it contained all SynCom15 strains, and it was the consortia that

exhibited the highest suppressive capacity of all DMCs tested from **Fig. 1C** in our *in vivo* infection

model. We found that Blocks 1 and 2 did not suppress *Kp*-MH258 greater than saline intervention

(**Fig. 4D**, **Supplementary Table 7I**). Additionally, we found that DMC46 suppressed *Kp*-MH258

several orders of magnitude less on days 5 through 11 after infection by *Kp*-MH258 and was more

gradual than SynCom15, illustrating that removing 31 strains from DMC46 was beneficial for

achieving a heightened suppressive capacity of *Kp*-MH258 *in vivo* (**Fig. 4D**, **Supplementary**

Table 7I). We note that the increased suppressive capacity of DMC46 observed between days 11

and 16 compared to the results shown in **Fig. 2B** was also observed with the PBS intervention.

Thus, pathogen suppression during this time period was likely due to the natural course of

microbiota repair as our murine model was not germ-free but was open to the environment.

Collectively these results illustrated that (i) SynCom15 suppressed *Kp*-MH258 across

diverse, unrelated conditions and (ii) SynCom15 intervention resulted in a marked, rapid, and

sustained clearance of *Kp*-MH258 in a pre-clinically relevant *in vivo* mouse model of infection.

These findings motivated us to understand the mechanism of action of SynCom15 and to evaluate

its effect on gut pathogens other than *Kp*-MH258.

 SynCom15 ameliorates antibiotic-induced dysbiosis, suppresses Kp-MH258 by acidification and

metabolic remodeling, and suppresses other gut pathogens through various mechanisms

To understand how SynCom15 suppresses *Kp*-MH258, we started by characterizing the

suppressive capacity of SynCom15 from a taxonomic perspective. First, we evaluated the

suppressive capacity of each single strain on *Kp*-MH258 in BHIS media. We found that only four

strains suppressed *Kp*-MH258 more than two orders of magnitude in BHIS while the remaining

11 strains suppressed *Kp*-MH258 less than one order of magnitude (**Extended Data Fig. 10A**).

This result illustrated that interactions between strains were key for specifying the suppressive

capacity of SynCom15, consistent with prior literature demonstrating that the capacity to

decolonize *K. pneumoniae* in *in vitro* culture was not a property of single strains[27]. Additionally,

the Block 1 consortium comprised all four of the strains that suppressed *Kp*-MH258 more than

two orders of magnitude yet failed to suppress *Kp*-MH257 *in vivo*, suggesting that the translatable

capacity of SynCom15 could not be explained simply by the effects of individual strains *in vitro*.

Second, as SynCom15 includes a strain of *E. coli*—a species whose inclusion in diverse consortia

was previously shown to be important for designing consortia that suppress *K. pneumoniae*—we

evaluated the effect of removing *E. coli* from SynCom15 on pathogen suppression **(Extended**

Data Fig. 10B)[27] · We found that SynCom15 without *E. coli* matched the suppressive capacity of

SynCom15 in both BHIS and GF cecal content media (**Extended Data Fig 10C**,**D**). This result

illustrated that the effect of SynCom15 was not merely due to the inclusion of *E. coli* in the

consortium. Third, we evaluated whether the suppressive capacity of SynCom15 could be inferred

via taxonomic information gleaned from our mouse experiments. Taxonomic profiling of fecal

samples collected through the course of our experiment revealed that 10 of the 15 strains in

SynCom15 engrafted in at least one of the mice within the cohort with one such strain being *E.*

coli. Additionally, dynamics of SynCom15 strains showed that 5 of the 10 strains were present at

detectable fractional abundances throughout the course of the experiment—*C. symbiosum*, *B.*

xylanisolvens, *C. innocuum*, *B. obeum*, and *B. caccae* (**Extended Data Fig. 10B**, inset)

(Methods). We therefore built three communities—(i) a community constituting strains that

consistently engrafted the mice (10 species), (ii) a community constituting strains that consistently

engrafted the mice but without *E. coli*, and (iii) a community constituting strains that were detected

in mice across all timepoints (5 species). The 10-member community and the 10-member

community without *E. coli* suppressed *Kp*-MH258 two orders of magnitude in BHIS and did not

exhibit suppression in GF cecal content media. The 5-member community suppressed *Kp*-MH258

one order of magnitude in BHIS and did not exhibit suppression in GF cecal content media

(**Extended Data Fig. 10C,D**). Together these results suggested that elucidating a mechanism of

action of SynCom15 from the perspective of taxonomic composition alone would be challenging.

We therefore turned to a more function-centric view for understanding how SynCom15

suppresses *Kp*-MH258.

We leveraged the results of our mouse experiments and analyzed the longitudinal

dynamics of fecal microbiota structure to evaluate the effect of administering SynCom15 after

antibiotic exposure and infection by *Kp*-MH258 (Methods). We found that the effect of

administering SynCom15 mirrored the FMT intervention, returning the state of the microbiota to

that observed prior to antibiotic treatment (**Fig. 5A**, **Extended Data Fig. 11A,B**, **Supplementary**

Table 8A,B). Moreover, we noted that while treatment with FMT resulted in the return of mouse-

specific gut bacteria (e.g. *Duncaniella* and *Paramuribaculum*), treatment with SynCom15 resulted

in the detectable presence of human-specific gut bacteria (e.g. genera belonging to the genera

Bacteroides and a bloom of *Bifidobacterium*) (**Extended Data Fig. 11C**, **Supplementary Table**

8C). Histology of the mouse colon showed that SynCom15 was well tolerated as an intervention

showing no evidence of inflammation or tissue insult (**Extended Data Fig. 11D**). Together, these

results demonstrated that SynCom15 restores community diversity of the gut microbiota as

measured by coarse phylogenetic composition but results in a compositional signature distinct

from natural murine microbiota. Our findings motivated a deeper investigation into the functional

mechanisms by which SynCom15 suppresses pathogens.

To begin interrogating the functional output of SynCom15, we performed targeted

metabolic profiling on supernatants from the five most suppressive and least suppressive DMCs,

as well as SynCom15, when co-cultured with *Kp*-MH258 in BHIS media (Methods). This metabolic

panel encompassed metabolites central to microbiome functional capacities—fatty acids, indoles,

amines, and amino acids—and has been used previously to describe human gut microbiomes

(Methods)[32,41,52,53]. Our analysis revealed two distinguishing signatures of suppressive

consortia—select amino acid depletion and fatty acid production (**Fig. 5B**, **Extended Data Fig.**

12A, **Supplementary Table 9A,B**). Relative to *Kp*-MH258 monoculture, the most suppressive

DMCs and SynCom15 exhibited elevated levels of phenylacetic acid, valeric acid, hexanoic acid,

and 5-aminovaleric acid while SynCom15 uniquely showed additional increases in lactic and

succinic acid (**Fig. 5B**, **Supplementary Table 9A,C**). Additionally, multiple amino acids including

those with aromatic and non-polar sidechains—phenylalanine, alanine, isoleucine, leucine, and

valine—plus glutamate and 5-oxoproline were significantly depleted in the suppressive DMCs as

well as SynCom15 (**Fig. 5B**, **Supplementary Table 9A**,**C**). Thus, elevated levels of medium-chain

fatty acids (valeric acid, hexanoic acid) with aromatic and amino-acid derived acids (phenylacetic

acid and 5-aminovaleric acid) suggested environmental acidogenesis consistent with the

previously described mechanism involving pH-dependent suppression of gram-negative

Enterobacteriaceae[54]. Additionally, depletion of branched-chain (isoleucine and leucine), aromatic

(phenylalanine), and nitrogen handling (glutamate, 5-oxoproline, alanine) amino acids suggested

nutrient competition that could potentially limit *K. pneumoniae* growth, in accord with previous

studies illustrating the importance of branched-chain, aromatic, and glutamate family amino acids

(glutamate and proline) for *K. pneumoniae* fitness[55–57]. Consistent with these results, metabolic

profiling of fecal samples collected from our mouse model of *Kp*-MH258 infection showed a

significant increase in fatty acids at day 10 of the experiment and depletion of amino acids at day

12 of the experiment when mice were given the SynCom15 intervention compared to intervention

with saline (**Extended Data Fig. 12B**). Together, our findings suggested that metabolic

remodeling of the environment by SynCom15 was important for suppression of *Kp*-MH258.

Following our metabolomic analysis, we performed a series of experiments to evaluate

the mechanistic basis of suppression by SynCom15 in BHIS and GF cecal content media. First,

we evaluated whether the metabolic environment created by SynCom15, as opposed to contact-

dependent direct physical interaction, was the driver of *Kp*-MH258 suppression. To test this idea,

we co-cultured SynCom15 with *Kp*-MH258 in a transwell system where SynCom15 was physically

separated from *Kp*-MH258 but could still interact with *Kp*-MH258 in a contact-independent

manner via metabolites (Methods). We found that in both BHIS media and GF cecal content

media, SynCom15 suppressed *Kp*-MH258 to nearly the same degree in the transwell system as

when in co-culture (**Fig. 5C**, left; **Supplementary Table 10A**). This result illustrated that the

mechanism of *Kp*-MH258 suppression by SynCom15 was through contact-independent

interaction with the pathogen.

Previous literature has demonstrated that acidification of the environment can

substantially augment the effect of fatty acid production on suppression of gram negative

Enterobacteriaceae[54]. Because suppression in the GF cecal content media condition was

substantially higher than that of BHIS, we evaluated the pH of each media condition with and

without bacterial culture. Both media conditions exhibited a baseline pH of 7.0. However,

SynCom15 co-cultured with *Kp*-MH258 in GF cecal content media resulted in a drop in pH to 5.0

whereas the pH of the co-culture in BHIS media remained at 7.0. We next evaluated whether a

pH drop by itself was sufficient to suppress *Kp*-MH258. We grew *Kp*-MH258 alone in BHIS and

GF cecal content media and found that the pH of the culture in BHIS was maintained at 7.0 while

the pH of the culture in GF cecal content media dropped to 5.0. However, despite the acidification

of the culture in GF cecal content media, the fitness of *Kp*-MH258 remained high (**Fig. 5C**, left

and middle; **Supplementary Table 10B**). This result illustrated that environmental acidification

alone was not sufficient to explain suppression of *Kp*-MH258.

Our results so far showed that both SynCom15/*Kp*-MH258 co-culture and *Kp*-MH258

monoculture drop environmental pH, but that SynCom15/*Kp*-MH258 co-culture also produces

fatty acids. We therefore hypothesized that SynCom15 suppresses *Kp*-MH258 through a

combination of fatty acid production and acidification of the media—a mechanism consistent with

previous studies interrogating gut microbiome-mediated decolonization of gram negative

Enterobacteriaceae[54]. To test this hypothesis, we first collected the supernatant from SynCom15

co-cultured for 120 hours with *Kp*-MH258 in BHIS and GF cecal content media. Second, we

acidified the supernatant derived from BHIS media to a pH of 5.0 and buffered the supernatant

derived from GF cecal content media to a pH of 7.0 (Methods). Finally, we inoculated the acidified

and buffered supernatants with *Kp*-MH258 for 120 hours and compared the resulting pathogen

abundance in supernatants that were not acidified and buffered. We found that acidification of the

supernatant in BHIS suppressed *Kp*-MH258 greater than five orders of magnitude; buffering of

the supernatant in GF cecal content media increased the abundance of *Kp*-MH258 by greater

than four orders of magnitude (**Fig. 5C**, right; **Supplementary Table 10C**). To evaluate whether

this effect was specific to SynCom15, we collected the supernatant of *Kp*-MH258 monocultured

for 120 hours in BHIS and in GF cecal content media. We then followed the same protocol as

above. We found that *Kp*-MH258 cultured in its own supernatant in BHIS and GF cecal content

media maintained at their native pH retained its fitness, decreasing less than an order of

magnitude in each condition (**Fig. 5C**, right; **Supplementary Table 10C**). We also found that

changing the pH in either condition affected the fitness of *Kp*-MH258 but not to the same degree

as the supernatant collected from SynCom15/*Kp*-MH258 co-culture (**Fig. 5C**, right;

Supplementary Table 10C). We next evaluated whether the suppressive effect of the

supernatant collected from the co-culture of SynCom15/*Kp*-MH258 was because of the inclusion

of *Kp*-MH258. We collected the supernatant of SynCom15 cultured on its own for 120 hours in

both BHIS and GF cecal content media, followed by the same protocol of acidification and

buffering with inoculation of *Kp*-MH258. We found that the presence of *Kp*-MH258 in the co-

culture with SynCom15 did not affect the suppressive phenotype of the supernatant nor the effect

of pH on the suppressive phenotype (**Extended Data Fig. 12C**, **Supplementary Table 10D**).

Collectively, these results suggest that environmental acidification with fatty acid production was

a plausible mechanism by which SynCom15 suppresses *Kp*-MH258 across different

environments.

Several previous studies have illustrated the connection between fatty acid production,

nutrient depletion, and environmental acidification in naturally occurring microbiomes with

suppression of various pathogens[54,58–61]. We therefore sought to evaluate the suppressive effect

of SynCom15 against a library of phylogenetically distinct gut pathogens. These included different

strains of MDR *K. pneumoniae*, *E. coli*, and *C. difficile* as well as a single strain of vancomycin

resistant *Enterococcus* (VRE). We chose these pathogens because of (i) their clinical prevalence

with respect to disease, (ii) the difficulty associated with their treatment, and (iii) because they

cause disease through different biological mechanisms—*C. difficile* colitis through toxin

production and *K. pneumoniae*, VRE, and *E. coli* infection through induction of colitis and/or gut

barrier translocation leading to sepsis. We evaluated the co-culture of SynCom15 with these

pathogens in GF cecal content media. We note that due to antibiotic selection overlap, we

evaluated the fitness of *Kp*-MH1867 and MDR *E.coli* strains when co-cultured with SynCom15

without its *E. coli* strain. With respect to strains of *Enterobacteriaceae*, we found that co-culture

with SynCom15 suppressed all strains of *K. pneumoniae* and *E. coli* three to four orders of

magnitude (**Fig. 5D**, **Supplementary Table 11A**). We found that SynCom15 suppressed two *C.*

difficile strains—VPI and ST1-49—by two orders of magnitude, two other strains—R020291 and

ST1-75—greater than one order of magnitude; and one strain—ST1-6—less than one order of

magnitude (**Fig. 5D**, **Supplementary Table 11A**). Finally, we found there to be minimal effect of

SynCom15 on suppressing VRE (**Fig. 5D**, **Supplementary Table 11A**). Across all pathogens, we

found that the Block 1 consortium was generally not suppressive, resulting in the same level of

pathogens and the pathogen cultured alone (**Fig. 5D**, **Supplementary Table 11A**). Thus,

SynCom15 was able to suppress several strains of phylogenetically diverse gut pathogens.

We next evaluated mechanisms by which SynCom15 suppressed a selection of *K.*

pneumoniae, *E. coli*, and *C. difficile* strains. We co-cultured SynCom15 with *Kp*-MH225, *E. coli*

X43, and *C. difficile* VPI and ST1-49 in GF cecal content media using a transwell culture system

for 120 hours. We found that separating SynCom15 from *Kp*-MH225 abrogated the suppression

of the pathogen: SynCom15 suppressed *Kp*-MH225 three orders of magnitude in co-culture but

less than one order of magnitude in the transwell culture system (**Extended Data Fig. 13**). In

contrast, the transwell co-cultures of SynCom15 with *E. coli* X43 and *C. difficile* VPI and ST1-49

strains recapitulated pathogen suppression observed in co-culture. Using the same protocol of

buffering the GF cecal content media condition used in **Fig. 5C**, we found that suppression of *E.*

coli X43 and *C. difficile* VPI and ST1-49 were pH dependent (**Extended Data Fig. 13**,

Supplementary Table 11B).

Collectively, our interrogation of pathogen suppression by SynCom15 revealed several

findings. First, environmental acidification coupled with fatty acid production appeared to be a

mechanism by which SynCom15 suppressed *Kp*-MH258, possibly augmented by depletion of

specific amino acids. Second, though designed to suppress *Kp*-MH258, SynCom15 was able to

suppress gut pathogens belonging to diverse phylogenies that manifest pathology in humans by

different biological mechanisms. Third, the capacity to suppress these pathogens was dependent

on different mechanisms—through indirect, pH-dependent suppression for *E. coli* X43, *C. difficile*

VPI, and *C. difficile* ST1-49 strains versus through direct physical interaction for *Kp*-MH225—

illustrating the degeneracy of mechanisms for pathogen suppression encoded by SynCom15.

Lastly, beyond only pathogen suppression, we found that SynCom15 repairs the microbiota after

antibiotic-induced dysbiosis, promoting its return to a healthy and diverse state.

Strains of SynCom15 are not prevalent or abundant in gut microbiomes of healthy humans

We explored the extent to which SynCom15 was represented across healthy humans who

provided stool samples from which we created our strain bank. We first interrogated the

prevalence of the genera constituting SynCom15 strains in these fecal samples. We found that

SynCom15 was composed of a diversity of genera observed across the set of healthy gut

microbiomes (**Fig. 6A**). Next, we interrogated the prevalence of the SynCom15 species across

the fecal samples of the healthy donors (Methods). We found that no healthy human microbiome

contained more than eleven of the SynCom15 species above a fractional abundance of 0.1%

(**Fig. 6B**; **Supplementary Table 12**). Moreover, we found certain SynCom15 species to be sparse

in their prevalence across donors as detected by metagenomic sequencing. *M. jalaludinii* was not

detectable in any donor; *M. massiliensis* was detectable in two donors; *C. symbiosum* in three

donors; and *C. scindens* in four donors. Amongst strains that were most prevalent, *B. obeum* and

B. faecis were detectable in 20 donors; *L. celerecrescens* in 14 donors; *C. comes* in 13 donors;

B. caccae in 12 donors. Finally, we interrogated the fractional abundance of SynCom15 species

across the fecal samples. We found SynCom15 species were present at a relative abundance of

less than 5% across all donors, with most species being found at a relative abundance of less

than 0.5% (**Fig. 6C**).

Together, these findings illustrated two conclusions. First, the composition of SynCom15

was distinct from that found across healthy human gut microbiotas determined by metagenomic

sequencing. This was either because SynCom15, as a community, did not exist in the healthy

samples from our cohort or because several of SynCom15 strains were undetectable by our

sequencing methods due to their low abundance. Second, the strains comprising SynCom15

were low prevalence and abundance amongst fecal samples of healthy donors. This result

highlighted the power of generating and using broadly diverse strain banks for engineering

synthetic bacterial communities as compared to strain banks reflecting the compositional

abundance and prevalence distributions gleaned from analysis of natural human microbiomes—

an approach previously used to design synthetic microbiomes[10–12].

Describing communities by strain presence/absence was important for learning a generative

model of pathogen suppression

Our overall approach to deriving SynCom15 followed six sequential steps: (1) Compress the design space and (2) Design consortia implementing an algorithmic constraint based on selection for genomic diversity, (3) Build and (4) Test consortia for function, (5) Learn a statistical model of function from patterns of strain presence/absence of designed consortia, and (6) Infer constraints learned by the model. As this approach is distinct from the typical 'Design-Build-Test-Learn' (DBTL) approach used in systems theory engineering, we term this approach 'Constraint Distillation'. We note that for the specific function of Kp-MH258 suppression, Constraint Distillation was not iterative: SynCom15 was directly derived from just one round of Compress/Design-Build/Test-Learn-Infer. Thus, we next explored two facets of Constraint Distillation: (i) its experimental efficiency and (ii) key choices we made that enabled our framework to converge on SynCom15.

To quantify the experimental efficiency of Constraint Distillation in generating SynCom15, we computed an effective compression score that considers three measurements: (i) the total complexity of information prior to implementing each step of Constraint Distillation, (ii) the total complexity of information remaining after implementing each step of Constraint Distillation, and (iii) the information needed to converge on the output of each step of Constraint Distillation (**Fig. 7A**) (**Supplementary Discussion**). Starting with a strain bank of 848 gut commensal strains and reducing the possible space of strains to 46 required whole genome sequencing of all strains—a total of nearly 10^{12} base pairs of information—and subsequently implementing a UMAP-based dimension-reduction which required no additional information. This step reduced the space of design from 848 to 46 strains resulting in a ~10^{230} compression of information yet left greater than 35 billion strain combinations as the space of design. To arrive at SynCom15 required performing 179 experiments where each experiment evaluated a single DMC for its capacity to suppress Kp-

MH258 in BHIS media (96 experiments to train and validate the predictive model; 23 experiments

to evaluate reproducibility of assay; 60 experiments to demonstrate the generative capacity of the

predictive model). This yielded a compressive power of ~10^{11} (**Fig. 7B**). These findings illustrated

that Constraint Distillation enabled navigation of a remarkably high-dimensional combinatorial

space to ultimately arrive at SynCom15 using only the genome sequences of the individual strains

as prior biological knowledge.

We next evaluated the choices that were critical for the efficiency of Constraint Distillation.

The Compress step resulted in 46 strains that recapitulated the diversity of the 848 strains in our

strain bank. However, our results showed that DMC46—the consortium containing all 46 strains—

was not effective at suppressing *Kp*-MH258 *in vivo*. Thus, simply performing the Compress step

was insufficient for engineering a consortium that translated pathogen suppression across

environments. The Design step of leveraging genetic diversity to construct 96 DMCs from 46

strains resulted in a substantial reduction in complexity. However, our results from screening 96

DMCs illustrated that merely designing for genetic diversity proved insufficient for engineering

consortia that predictably suppressed *Kp*-MH258. Regarding the Test function itself, we found that

suppression of *Kp*-MH258 was not explainable by simple heuristics—(i) richness and diversity of

consortia, (ii) the presence/absence of individual strains, (iii) or the combination of

richness/diversity with the presence of strains previously shown to metabolically overlap with *K.*

pneumoniae like *E. coli*. Moreover, the result of inference on our RF model could have yielded a

phylogenetically constrained group of strains, e.g. all strains of the same genera. However, this

was not the case as SynCom15 remained phylogenetically diverse. Thus, the test function we

chose to target—suppressing *Kp*-MH258—was complex enough to require a specific set of

diverse, interacting microbes to satisfy suggesting that the experimental efficiency of our process

was likely not merely due to a simplicity of test function. These observations shifted our focus to

the final two steps of Constraint Distillation: Learning and Inference.

With respect to the Learning step, different types of ML models—RF and deep learning-

based neural networks—proved to be useful in statistically learning an input-output relationship

with our data. Thus, the specific class of ML model was not critical for the efficiency of Constraint

Distillation. However, a specific choice we made in the Learning step was to describe each of the

DMCs by their pattern of strain presence/absence. Notably, this choice neglected information

about dynamics of the co-culture, physiologic state of the microbes in the culture, interaction with

Kp-MH258 throughout the time-course of the co-culture, or mechanistic information regarding

pathogen suppression. Additionally, our efforts at characterizing the mechanism of action of

SynCom15 showed that the supernatant of the co-culture after 120 hours was sufficient for

suppressing *K. pneumoniae*, illustrating the importance of the metabolic environment created by

SynCom15 towards pathogen suppression. As such, we sought to understand whether a

generative model of consortia design could be created using the metabolic profiles of DMCs as

defined at 120 hours of co-culture with *Kp*-MH258. Our rationale was that pursuing this goal would

enable comparing two types of information—strain presence/absence of DMCs versus metabolite

content at 120 hours of co-culture—that were leveraged by us to (i) design SynCom15 or (ii)

explain how SynCom15 works respectively.

We performed targeted metabolic profiling of DMCs described in **Fig. 1C** in addition to the

60 DMCs that collectively served as the out-of-sample set when co-cultured with *Kp*-MH258 for

120 hours (**Supplementary Table 13A,B**). The metabolic panel used was the same used for

characterizing consortia in **Extended Data Fig. 12A**. We trained and validated an RF model and

DNN model on the metabolic profile of DMCs to predict *Kp*-MH258 abundance after 120 hours of

co-culture (Methods). We found that these models were well trained, exhibiting an r^2 value of 0.9

and 0.7 respectively with the validation set of data (**Extended Data Fig. 14A**, **Supplementary**

Table 13C). However, these models failed to generalize to out-of-sample communities (**Extended**

Data Fig. 14B, **Supplementary Table 13D**). Our finding that the models performed well on the

validation dataset but were unable to generalize out-of-sample pointed to model overfitting. We

emphasize that this result does not suggest that metabolomics is uninformative. Indeed, multiple

factors could limit its generative utility in our data, including incomplete coverage of relevant

metabolites, the use of a single endpoint (e.g., 120 hours of co-culture) rather than a time-course

based input, and missing important variables like supernatant pH—a trait we found was crucial

for explaining the mechanism of pathogen suppression by SynCom15. However, these results

provided us the unique opportunity to further understand Constraint Distillation as a process: what

statistical characteristics of an input space that describes communities enable a generative design

model to extract beyond the training ensemble?

We therefore examined the eigenspectrum of variation in the two input spaces used: strain

presence/absence and metabolites. Examination of each eigenspectrum with respect to the

capacity to suppress *Kp*-MH258 revealed a stark difference between the way information was

encoded in the two spaces. In the strain presence/absence space, the top eigenvectors were the

most strongly associated with *Kp*-MH258 suppression; additionally, this association decayed with

eigenvector rank (**Extended Data Fig. 14C**, **Supplementary Table 13E**). This alignment between

the statistical structure of variation and our target function concentrated in a few high-variance,

low-noise directions, yielded a low-dimensional coordinate along which suppression is ordered

thereby defining conditions where ML models can generalize. In the metabolite space, this

alignment was absent. A high-variance mode (the second eigenvector harboring >20% variance)

showed little association with suppression while multiple low-variance modes (harboring <5%

variance each) were more associated with suppression (**Extended Data Fig. 14D**,

Supplementary Table 13F). This observation provided an explanation as to why overfitting was

occurring: models fit the training data by over-indexing on fluctuant, low-variance modes but these

modes are unstable across new communities thereby degrading out-of-sample performance.

Thus, our results shed light on specific critical qualities of Constraint Distillation—a series

of sequential steps that took as input the genome sequences of strains and output a statistical

model of targeted, function-specific consortia design that directly led to the construction of

SynCom15 (**Fig. 7C**). Namely, the capacity to create a useful generative model depended on

describing the 96 DMCs in a way that enabled informational resonance between the statistical

structure of DMCs and variation in the target function. When statistical signal associated with

function is concentrated in the top variance modes of the input representation, learning models

can become generative. When statistical signal associated with function resides in low-variance

modes, models may fit but fail to generalize.

Discussion

We have engineered a defined, synthetic microbiome, SynCom15, that (i) suppresses *Kp*-

MH258 (a clinical MDR strain of *K. pneumoniae*) across diverse environments, (ii) matches the

efficacy of a whole stool transplant in a pre-clinically relevant *in vivo* mouse model of infection,

and (iii) is compositionally distinct from natural human gut microbiomes. We achieved this using

Constraint Distillation—a process that is agnostic to bacterial interactions, strain phenotypes, data

from cohort studies, qualities of natural microbiomes, or any mechanism of action. Instead,

Constraint Distillation comprised learning a map of taxonomic structure to desired function from

observational data alone to yield SynCom15. A plausible mechanism by which SynCom15

suppressed *Kp*-MH258 was through fatty acid production and acidification of the environment.

This mechanism was shared across suppression of multiple, phylogenetically distinct gut

pathogens including strains of *E. coli* and *C. difficile.* A sensitivity analysis of the Constraint

Distillation framework demonstrated that describing consortia by their pattern of strain

presence/absence was critical for achieving an accurate generative model of design. This was

because of informational overlap between the structure of taxonomic variation in the 96

engineered consortia that were functionally screened and the target function of suppressing *Kp*-

MH258.

We acknowledge clear limitations regarding (i) the effect of SynCom15 and (ii) Constraint

Distillation as an approach for engineering consortia. Regarding SynCom15, we note that our *in*

vivo model considered only one dietary background—normal mouse chow. However, it is known

that dietary background can have a substantial effect on the composition of the microbiome as

well as the capacity for probiotic interventions to be efficacious[62,63]. Normal mouse chow is fiber

rich; recent studies have demonstrated the limitations of more westernized diets with respect to

ameliorating antibiotic-induced dysbiosis[64]. Additionally, our *in vivo* model also only spanned a

single genetic background—C5J/Bl6. These experimental choices limit the extent to which

SynCom15 can be considered a generally useful consortia for rescuing dysbiosis of the gut

microbiome. It will be important for future efforts to vary such environmental variables that likely

play an important role in determining the efficacy of a live biotherapeutic intervention. Regarding

Constraint Distillation, our results showed that we could learn a generative model of consortia

design towards *K. pneumoniae* suppression using 96 synthetic consortia as an initial ensemble.

Importantly, this test function could be evaluated using *in vitro* culture. For test functions that

involve extra-enteric effects in the host (e.g. the capacity of the gut microbiome to influence the

tumor microenvironment, gut brain axis, immune system, or other organs), we note that assay

throughput may become a significantly limiting factor. To demonstrate that Constraint Distillation

can engineer synthetic microbiomes functions that, at baseline, require murine models, it will be

important to further develop theoretical frameworks that enable designing consortia in a more

informed manner such that the subsequent Build-Test/Learn-Infer steps can be rapidly iterated

upon if necessary.

As synthetic ecology efforts continue to evolve within application to the human gut

microbiome, we contrast the composition and functionality of our consortium with other consortia

that have previously been created as surrogates for 'healthy' microbiomes. Recent studies have

emphasized the power of large consortia, comprising greater than 40 strains, for pathogen

suppression and other functions associated with a healthy, diverse gut microbiome[11,27]. One

rationale for such an approach is to maximize the functional potential of a consortium by simply

adding more diverse strains. Our data demonstrate that it is possible to create a synthetic

consortium that is sparse—comprising only 15 members—yet can clear *K. pneumoniae* from the

gut as well as rescue the effects of antibiotic-induced dysbiosis. Moreover, our data illustrate that

a larger consortium comprising more gut commensals that is inclusive of SynCom15—DMC46—

was in fact less efficacious with respect to pathogen clearance, thereby illustrating the limitation

of using larger commensal consortia as therapeutics. Instead of directly inoculating the gut with

the large diversity of strains that may be present in a whole stool sample, the intervention of

SynCom15 appears to be more indirect—allowing the rescue of bacterial diversity through

creating an environmental milieu for commensal growth. We reason this is the property underlying

the sustained pathogen resistance we see well beyond gavage of SynCom15. In this sense,

SynCom15 represents a qualitatively different strategy for repopulating the gut microbiome after

perturbation relative to a full stool transplant: by creating an appropriate gut environment,

SynCom15 promotes the re-diversification of the gut ecosystem. From a more practical

perspective, to realize the value of synthetic ecology it will be important to attempt synthesizing

consortia as live biotherapeutics and establishing clinical trials for their efficacy. Given the

challenges associated with the regulatory landscape and manufacturing of live biotherapeutics,

having a defined mixture of 15 strains constitute the investigational new drug (IND) as opposed

to much larger consortia or other types of live biotherapeutic constructs that are undefined (e.g.

stool transplants or purified contents from healthy donor stool) will be a substantial benefit in

ultimately transitioning synthetic consortia to clinical settings.

Due to the complexity of microbiomes, there have been many recent efforts that augment

design with statistical formalism[21–24,65,66]. These efforts have used some level of biological

behavior to inform the statistical approach. Inclusion of these behaviors—metabolic qualities,

growth qualities, presumed architecture of interactions via consumer-resource models, interaction

characterization, functional screens—constitute the equivalent of a forward model imposed by the

experimenter. We note, this is not statistical design. Statistical design involves solving the inverse

problem—learning a model from observation alone. In other fields of biology and science

spanning protein design, material science, neuroscience, and complex systems-based macro-

ecology, implementation of statistical design involves (i) creating an ensemble of systems

reflecting extant diversity, (ii) learning a probabilistic model of constraint, and (iii) using the model

to generate synthetic systems[37–40,67–71]. In our work, we closely mirrored this approach with the

difference being that instead of collecting microbiome data from the wild and therefore

representing extant microbiome diversity, we created an ensemble of microbiomes using genetic

diversity as merely a sampling strategy. Thus, our approach fell directly within the purview of

statistical design. What are benefits of not including priors and only employing the strategy of

statistical design? First, this approach filters priors that are not general. For instance, our data

show that the previously published prior of consortia diversity and presence of *E. coli* to specify

K. pneumoniae suppression could not explain our results. Second, statistical design makes

possible the discovery of mechanism rather than anchoring on prior mechanisms as routes of

design. In our work, we first constructed SynCom15 then evaluated its mechanism of action rather

than the reverse. Third, learning a statistical model of design opens the possibility of truly synthetic

ecology—the creation of products that are unnatural and thereby expand the functional space

beyond what nature has created. We showed that SynCom15 was not detectable within natural

human gut microbiomes and encoded different mechanisms by which it achieved pathogen

suppression across phylogenetically diverse gut pathogens. While future work will test the limits

of generality with respect to using statistical design for engineering consortia against other

functions, our work has demonstrated that such a strategy is viable, efficient, and can lead to the

creation of novel synthetic products.

As increasingly more phenotypes are associated with the gut microbiome, we pose that a

useful test for judging the strength of association is through design of synthetic consortia that

affect the phenotype. We note that Constraint Distillation is agnostic to the specific function being

optimized. The 'Test' module ('T') can be adapted to any phenotype for which there is an assay.

Because implementing Constraint Distillation to engineer synthetic ecological systems does not

depend on natural community composition, known mechanisms, or pre-existing pathway

annotations, this approach offers a potentially general route to the statistical design of synthetic

microbiomes. More broadly however, Constraint Distillation is, in theory, also agnostic to the

specific type of 'part' used to construct a collective whole. That is, given an ensemble of parts

(e.g. bacterial strains) described by degrees of freedom (bacterial genomes), the process of

Constraint Distillation can be employed to create an emergent system. In this sense, Constraint

Distillation offers more than a procedure for specifically designing synthetic microbiomes but could

be a more abstract strategy for using collections of heterogeneous parts to engineer synthetic

emergent systems—systems where higher-order, unintuitive interactions between parts create a

functional collective. As physics-based principles for building emergent systems *de novo* do not

currently exist, testing this idea across both ecological and non-ecological domains represents an

exciting future direction.

Our results challenge prevailing assumptions about what is required to design functional

microbial communities for therapeutic effect. Prior studies have emphasized the need for high-

dimensional measurements, detailed mechanistic understanding, or complex dynamical models

to capture structure-function relationships that govern microbial ecosystems. Community design

efforts have typically drawn on a wide array of information sources including: (i) quantitative

modeling of interspecies interactions, (ii) knowledge of molecular mechanisms underlying target

functions, (iii) pathway-level genome annotations, (iv) human microbiome composition as a

blueprint, or (v) down-selection from natural communities exhibiting desirable traits such as

pathogen resistance (**Supplementary Table 14**, **Supplemental Discussion**)[10,11,13,16,22–25,27,72].

Moreover, strategies for translating microbiome function across unrelated, diverse contexts have

involved learning even more information about environmental differences and context-specific

bacterial interactions for consortia optimization[3,73]. Our findings paint a substantially different

picture. We find that generative principles of consortia design can emerge from far simpler inputs.

A binary encoding of just 96 metagenomically diverse synthetic communities by strain

presence/absence coupled with empirical functional measurements was sufficient to achieve two

key results: the capacity to train a statistical model mapping community structure with function

and guide the construction of a sparse community (SynCom15) that translated successfully to

diverse environments. Constraint Distillation proved substantially compressive, navigating a large

design space with minimal information relative to the full combinatorial complexity of design. In

computational terms, this workflow achieved an efficient search over a high-dimensional space

using a low-dimensional, statistically learnable representation. We hypothesize that the success

of this approach stems from an underappreciated property of many biological systems: that

despite their apparent complexity, biological systems are often governed by low-dimensional rules

that can be successfully inferred directly from observational data. Our results regarding the

informational resonance between patterns of strain presence/absence and pathogen suppression

supports this idea. Placing learned statistical patterns before biological phenomenology or

mechanistic understanding has found growing support across biology, including recent

breakthroughs in data-driven protein design and the adoption of AI-driven approaches at the scale

of genomes and tissues[74–81]. Our results suggest that the same principle may hold for microbial

ecosystems.

Figures

Oliveira et al., Figure 1

889
890
Figure 1. Designing, building, and testing synthetic microbiomes for suppression of *Kp*-
MH258. (A) Subset of commensal strain bank used to engineer synthetic consortia; strains are
dots (n = 46) within Uniform Manifold Approximation Plot (UMAP) axes. **(B)** Schematic of
algorithm used to create Designed Microbial Communities (DMCs). **(C)** Matrix of 96 engineered
DMCs (rows) as well as *Kp*-MH258 in monoculture (bottom row) by strains (columns). Blue pixel
indicates strain was included in the DMC, white pixel indicates strain was not included in the DMC.
Each row is labeled by 'Community richness'—the number of strains included in the design of the
DMC—and '*Kp*-MH258 abundance'—the colony forming units (CFUs) per milliliter of *Kp*-MH258
remaining after 120 hours of co-culture with the DMC in Brain Heart Infusion media supplemented
with cysteine (BHIS media). **(D)** Abundance of *Kp*-MH258 (y-axis) across time (x-axis) in
monoculture (black dots), in co-culture with all DMCs from panel C comprised of 14 strains (grey
distribution), or in co-culture with DMC-Rational (14 strains) in BHIS media.

A

B

903
904

Figure 2. Suppression of *Kp*-MH258 by DMCs *in vitro* does not translate *in vivo*. (A) Experimental workflow for *in vivo* model of *Kp*-MH258 infection. Specific-pathogen-free (SPF) mice were treated with metronidazole, neomycin, and vancomycin ('MNV') in drinking water for four days, gavaged with *Kp*-MH258 ('*Kp*-MH258 gavage'), then gavaged with either phosphate buffered saline (PBS), fecal microbial transplant (FMT), or one of several DMCs three days in a row. **(B)** Dynamics of fecal *Kp*-MH258 abundance through time for mice gavaged with either PBS, FMT, or one of five DMCs (see color key). Lines represent median values and error bars represent interquartile range. **(C)** Suppressive capacity of DMCs (colored dots; color scheme follows panel B) observed in BHIS at 120 hours of co-culture (x-axis) versus SPF-infected mice at day 16 post *Kp*-MH258 infection (y-axis). PBS gavage in mice and monoculture of *Kp*-MH258 alone in BHIS is represented by the black dot. Error bars represent interquartile range.

916
917
Figure 3. Distillation of constraints learned by generative model yields SynCom15. (A)
Predicted abundance of *Kp*-MH258 at 120 hours of co-culture with DMCs (x-axis) from a Random-
Forests model (RF, left) and a deep-learning based neural network (DNN, right) trained on data
from **Fig. 1C** versus observed abundance (y-axis). Each dot is one of 60 possible out-of-sample
DMCs not represented in the 96 DMCs shown in **Fig. 1C**. **(B)** 100,000 DMCs were generated *in*
silico and scored by the trained RF model. Histogram of 100,000 DMCs is shown as a function of
predicted *Kp*-MH258 abundance for each DMC after 120 hours of co-culture. DMCs for which the
RF model predicts less than 10^5 CFUs/mL of *Kp*-MH258 remaining after 120 hours of co-culture
are selected and converted into a matrix (inset). Rows are each of 5,752 possible DMCs ('*In silico*
DMC'); columns are one of 46 strains used to create DMCs in **Fig. 1C**; entries reflect whether a
strain is present (blue pixel) or absent (white pixel) in the DMC. **(C)** Loadings of each of the 46
strains onto eigenvector 46 of the matrix defined in panel B (left). Similarity between all pairs of

strains along eigenvector 46 was computed and resulting matrix was hierarchically clustered
resulting in distinct blocks of strains (right). Block 1 (orange) plus Block 5 (green) is defined as
SynCom15.

933
934
Figure 4. SynCom15 suppresses MDR *Kp*-MH258 across diverse environments. (A)
Workflow for creating cecal content media from germ-free (GF) or antibiotic-treated specific
pathogen free (Ab-treated SPF) mice (box). **(B)** Abundance of *Kp*-MH258 at 120 hours of co-
culture (y-axis) with each Block defined in **Fig. 3C**, SynCom15, and *Kp*-MH258 in monoculture
(x-axis) across BHIS media and medias defined in panel A. **(C)** Workflow for assaying suppressive
capacity of Blocks 1, 2, DMC46, SynCom15, and controls (PBS and FMT) in *in vivo* model of *Kp*-
MH258 infection (similar scheme as **Fig. 1D**). **(D)** *Kp*-MH258 fecal abundance versus time for
mice gavaged with various interventions (color key). Lines represent median values; error bars
represent interquartile range.

Figure 5. SynCom15 rescues antibiotic-induced dysbiosis, suppresses *Kp*-MH258 through metabolic remodeling plus environmental acidification, and suppresses a diversity of other gut pathogens. (A) PCoA of fecal microbiota for SPF mice on day 0, 7, and 16 of experiment shown in **Fig. 4C**; colored shape is centroid for indicated cohort. **(B)** Bar plots show distribution of normalized peak areas (y-axis) for each metabolite feature distinguishing the most suppressive from least suppressive DMCs (individual panels) across three different culture settings at the 72-, 96-, and 120-hour culture timepoint: *Kp*-MH258 in monoculture (black), *Kp*-

MH258 in co-culture with the most suppressive DMCs (dark blue), *Kp*-MH258 in co-culture with
the least suppressive DMCs (light blue), and *Kp*-MH258 in co-culture with SynCom15 (maroon).
*p < 0.05; **p < 0.01; ****p < 0.0001. **(C)** (Left) Abundance of *Kp*-MH258 in BHIS and GF cecal
content media in monoculture, co-culture with SynCom15, or co-culture with SynCom15 in a
transwell culture setup (SynCom15 TW) after 120 hours. (Middle) pH of BHIS remains constant
at 7.0 with and without bacterial culture; pH of GF cecal content media drops from 7.0 to 5.0 with
bacterial culture. (Right) Abundance of *Kp*-MH258 in BHIS and GF cecal content media when co-
cultured for 120 hours with supernatants collected from *Kp*-MH258 cultured alone (*Kp*-MH258
sup) or SynCom15/*Kp*-MH258 co-culture (SynCom15 sup). Both supernatants in BHIS media
exhibited a natural pH of 7.0; supernatants were then acidified to a pH of 5.0 (ΔpH). Both
supernatants in GF cecal content media exhibited a natural pH of 5.0; pH of supernatants was
then increased to 7.0 (ΔpH). **(D)** Abundance of various pathogens (y-axis) in monoculture (grey),
co-culture with SynCom15 (red), co-culture with SynCom15 without *E. coli* strain ('-*E. coli*') (red),
or co-culture with Block 1 (defined in **Fig. 3C**). Experiments were performed in GF cecal extract
media. In panels C and D, bars represent median values and error bars represent interquartile
range. **(E)** Summary of suppressive effects of SynCom15 on abundance of all tested pathogens.

Figure 6. Comparison of SynCom15 composition with composition of healthy human microbiomes. **(A)** Phylogenetic tree of genera present across fecal microbiomes of human donors. Brown genera are those found across SynCom15 strains. **(B)** Prevalence pattern for species of SynCom15 (rows) across donor fecal microbiomes (DFI is Duchossois Family Institute; columns). **(C)** Histogram of relative abundance for SynCom15 species (x-axis) across all fecal samples from population of healthy human donors.

A.

$$C = \frac{\left(\dfrac{A}{B}\right)}{I}$$

Complexity prior to compression

Complexity after compression

Effective compressive power

Information needed for compression

B.

Bank of 46 diverse commensal strains

Strain bank of 848 commensals

1   2   SynCom15

① Compress

② Design, Build, Test, Learn, and Infer

A: $2^{848}/2$
B: $2^{46}/2$
I: 8.65×10^{11}
(BPs of genomes)

A: $2^{46}/2$
B: 1
I: 179
(Experiments involving DMCs in BHIS)

$C_1 \sim 10^{230}$

$C_2 \sim 1.5 \times 10^{11}$

C.

Compress & Design

Genome

Strain bank

848 strains   46 strains

UMAP 2 / UMAP 1

Compression

Design consortia based on diversity

UMAP 2 / UMAP 1

Build & Test

96 Designed Bacterial Consortia (DMCs)

Strain presence/absence   Kp

DMC01
DMC02
DMC03
DMC04
DMC05
DMC06
...

Learn

RF model   DNN model

Prediction with strains presence/absence

Predicted Kp abundance

Observed Kp abundance

Infer

In silico DMCs

Distill constraints

SynCom15

978
979
Figure 7. Constraint Distillation: A strategy for the statistical design of synthetic functional
consortia from sequenced bacterial strains. (A) Computing effective compressive power. **(B)**
Compression from starting point—bank of 848 gut commensal strains—to SynCom15 required
two sets of information. The first was the genomes of all strains to compress the design space
from 848 to 46 strains (a total of information content of 8.65×10^{11}). The second was 179
experiments comprised of evaluating DMCs in BHIS media used in the Design-Build-Test-Learn-
Infer steps to engineer SynCom15. **(C)** Schematic of Constraint Distillation defining inputs—
genome-sequenced strain bank—and outputs—statistically designed consortium for target
function.
989

Genera present in a healthy cohort

Genera of 848 strains in commensal strain bank

990
991 **Extended Data Fig. 1.** Fecal samples collected from 28 healthy human donors were subject to
992 shotgun metagenomic sequencing. Tree of the genera comprising all fecal microbiomes is shown
993 here. Colored in blue are the distribution of genera observed in our bank of 848 commensal
994 strains.

Extended Data Fig. 2. Matrix of strain (rows) by Prokka feature (columns) for 848 gut commensal strains in our strain bank was created where entries are a '1' if the Prokka feature is present in the strain proteome, and '0' if Prokka feature is absent in the strain proteome (top). Matrix was subject to UMAP decomposition resulting in two axes (UMAP1 and UMAP2) used to define clusters of bacteria; each dot in this plot is an individual bacterial strain (bottom).

A

B

1003

Extended Data Fig. 3. (A) Workflow for algorithm used to design synthetic consortia.
Communities with three bacterial strains are shown as an example. Given three possible
communities that could be created, the first step is to choose the mean dispersal of the lowest
third of pairwise distances between strains for each community. In the example shown here, the
lowest third is equivalent to the minimum pairwise distance for each community due to the
communities being comprised of only three strains (gray box). The second step is to choose the
community with the maximal dispersion per Step 1. In the case shown here, 'Community 1' would
be chosen as a DMC to create. **(B)** Phylogenetic diversity (y-axis) versus number of species in
DMCs (x-axis). Each dot is the averaged prevalence of a specific phylogenetic category spanning
Phylum, Class, Order, Family, and Genus (see color key) across all DMCs of a specific
membership size. The maximum possible prevalence of each phylogenetic category is set by the
DMC containing all 46 strains used to engineer DMCs (the right-most set of dots).

Extended Data Fig. 4. (A) Workflow for evaluating clearance capacity of DMCs for *Kp*-MH258 *in*
vitro ('BHIS' is Brain Heart Infusion media supplemented with cysteine, 'Kan' is kanamycin). **(B)**
Bacterial growth (OD$_{600}$, y-axis) of each of the 46 strains comprising the columns of the matrix in
Fig. 1C grown in BHIS media for 48 hours. Bars represent average values, error bars represent

standard deviation. **(C)** *Kp*-MH258 abundance (y-axis) after co-culture with each DMC (x-axis) for
120 hours where the x-axis is ordered by the most suppressive DMCs (left) to the least
suppressive (right) DMCs. **(D)** Correlation between the suppressive capacity of several different
DMCs across two experimental replicates in BHIS media. **(E)** Time course of *Kp*-MH258
abundance (y-axis) when co-cultured with each DMC described in the rows of **Fig. 1C**. Solid line
represents the median value of *Kp*-MH258 abundance in co-culture across all 96 DMCs at a given
timepoint; grey shade represents the distribution of *Kp*-MH258 abundance in co-culture across all
96 DMCs at a given timepoint. 'Below detection level' delineates limit of the assay to detect
suppressive capacity of a DMC.

Extended Data Fig. 5. (A) Communities ordered from least rich to most rich (x-axis) versus *Kp*-MH258 abundance after 120 hours of co-culture (y-axis). Solid line indicates detection level of the assay; first entry on the x-axis is *Kp*-MH258 abundance in monoculture (black bar). **(B)** Time course of *Kp*-MH258 abundance when co-cultured with DMC92 and DMC17 (left); Time course of *Kp*-MH258 abundance when co-cultured with DMCs containing 14 strains (right). **(C)** DMCs containing *E. coli* as a bacterial strain (dots) are separated by their capacity to suppress *Kp*-MH258 in co-culture (x-axis) and the number of species they contain in their design (y-axis). Highlighted are two DMCs (DMC92 and DMC03) with similar suppressive capacities that represent markedly different degrees of community richness. **(D)** Time course of *Kp*-MH258

abundance when co-cultured with DMC92 and DMC03. In panels B and D, Time course of *Kp-*
MH258 is represented by median values and error bars are represented by interquartile range.

Extended Data Fig. 6. The effect of the presence ('+') or absence ('-') (x-axis) of a given strain
on the capacity of DMCs (dots) to suppress *Kp*-MH258 (y-axis). Each panel is stratified by each
of the 46 strains comprising all DMCs; solid black line indicates median value of *Kp*-MH258
abundance. Yellow box distinguishes set of strains associated with a statistically significant
relationship between their respective presence/absence and the capacity for DMCs to suppress
Kp-MH258. Statistical significance determined by Mann-Whitney test.

Extended Data Fig. 7. For each of the 14 strains whose individual presence (red line)/absence (black line) was associated with a statistically significant difference in the capacity of DMCs to suppress *Kp*-MH258, histograms of the number of DMCs (y-axis) across the measured range of suppressive capacity (x-axis) are plotted.

1057
1058
1059 **Extended Data Fig. 8.** Loading of each of the 5,752 DMCs generated from the RF model (blue
1060 dots) onto eigenvector 46 (x-axis) versus predicted abundance of *Kp*-MH258 at 120 hours of co-
1061 culture (y-axis).

1062
1063 **Extended Data Fig. 9.** Strains (columns) that are defined by three different types of descriptions:
1064 present in SynCom15, present in the top 15 strains as determined by feature importance score of
1065 our trained Random Forests (RF) model, present in the DMC-Rational consortium.

1066
1067
Extended Data Fig. 10. (A) *Kp*-MH258 abundance (y-axis) after co-culture with each SynCom15
strain (x-axis) individually for 120 hours in BHIS media. **(B)** Construction of several variants of
SynCom15 (rows) defined by the presence (yellow dot) or absence (white dot) of SynCom15
strains (columns). Inset illustrates strains that were detected in mice. First row indicates of inset
that the strain was found in any mouse at any given timepoint after gavage of SynCom15 post
antibiotic treatment and pathogen infection; second row of inset indicates percent of mice in which
strain was observed; third row of inset indicates percent of timepoints in which strain was
observed. **(C,D)** *Kp*-MH258 abundance (y-axis) after 120 hours of co-culture with each of the
SynCom15 variants shown in panel B in BHIS media (panel C) and GF cecal content media (panel
D).

Extended Data Fig. 11. (A) Chao and Shannon diversity indices (y-axes) versus time (x-axes) for SPF mice treated with MNV, infected with *Kp*-MH258 ('*Kp*') and given PBS, FMT, or SynCom15. Lines represent media values, error bars indicate interquartile range. **(B)** Distribution of average relative abundance for major bacterial phylogenies represented in the fecal microbiota through time (x-axis) for infected mice treated with FMT (left panel), PBS (middle panel), or SynCom15 (right panel). Distributions are defined spanning kingdom to genera-level descriptions. **(C)** Relative abundance of *Bacteroides*, *Duncaniella, Paramuribaculum* and *Bifidobacterium* genera that are differentially abundant amongst infected mice treated with FMT, saline, or SynCom15 prior to antibiotic treatment (day 0) and at day 16 after treatment (equivalent to 11

1089 days after infection with *K. pneumoniae*). Statistical tests performed are two-way ANOVA; **$p <$
1090 0.01; ***$p < 0.001$; ****$p < 0.0001$. **(D)** Hematoxylin and eosin stain of colon of infected mice given
1091 SynCom15.

A

B

C

1092
1093
Extended Data Fig. 12. (A) p-values for differential enrichment of metabolites were computed for
the five DMCs that were the most and least suppressive of *Kp*-MH258 compared to *Kp*-MH258 in
monoculture in BHIS media (two-way ANOVA). Each dot in the plot is a metabolite; the x-axis
defines the range of p-values observed in the most suppressive DMCs; the y-axis defines the
range of p-values observed in the least suppressive DMCs. The red box delineates set of
metabolites that are most significantly differentially enriched in the most suppressive DMCs but
are not significantly enriched in the least suppressive DMCs. **(B)** Distributions of normalized peak
areas (y-axes) of fatty acids and amino acids from fecal samples collected on Day 10 and Day 12
of mouse experiment shown in **Fig. 4C** for mice gavaged with saline (PBS) or SynCom15
(maroon). ***p<0.001; Mann-Whitney. **(C)** Abundance of *Kp*-MH258 in BHIS and GF cecal content
media when co-cultured for 120 hours with supernatants collected from SynCom15 cultured alone

for 120 hours. Supernatant in BHIS media exhibited a natural pH of 7.0 and was then acidified to
a pH of 5.0 (ΔpH). Supernatant in GF cecal content media exhibited a natural pH of 5.0 and was
buffered to a pH of 7.0 (ΔpH). Bars represent median values, error bars represent interquartile
range.

1109
1110 **Extended Data Fig. 13**. Abundance of various pathogens (y-axis) when monocultured ('-alone'),
1111 or co-cultured in a transwell setup with SynCom15 ('+SynCom15 TW') in GF cecal content media
1112 at both natural pH and pH adjusted to 7.0. Bars represent media values, error bars represent
1113 interquartile range

A

Metabolites

DMCs

Train

RF model

$R^2 = 0.9589$

Observed *Kp*-MH258
120 hours co-culture
(log_{10} CFUs/mL)

Predicted *Kp*-MH258,
in training, 120 hours co-culture
(log_{10} CFUs/mL)

DNN model

$R^2 = 0.735$

Observed *Kp*-MH258
120 hours co-culture
(log_{10} CFUs/mL)

Predicted *Kp*-MH258,
in training, 120 hours co-culture
(log_{10} CFUs/mL)

B

Metabolites

DMCs
(OOS)

Predict

RF model

$R^2 = 0.3449$

Observed *Kp*-MH258
120 hours co-culture
(log_{10} CFUs/mL)

Predicted *Kp*-MH258,
120 hours co-culture
(log_{10} CFUs/mL)

DNN model

Observed *Kp*-MH258
120 hours co-culture
(log_{10} CFUs/mL)

$R^2 = 0.01405$

Predicted *Kp*-MH258,
120 hours co-culture
(log_{10} CFUs/mL)

C

Strains

|Correlation| with Suppression

Variance explained by PC
(PCs of DMCs described by strain presence/absence)

D

Metabolites

|Correlation| with Suppression

Variance explained by PC
(PCs of DMCs described by metabolite content
at 120 h of co-culture with Kp-MH258)

1114

Extended Data Fig. 14. **(A)** Training RF and DNN models of *Kp*-MH258 abundance from
metabolite content of DMCs when co-cultured with *Kp*-MH258 for 120 hours. **(B)** Prediction of *Kp*-
MH258 abundance from metabolite content of 60 out-of-sample DMCs (see **Fig. 3A**) using trained
RF and DNN models from panel A. **(C,D)** Information associated with *Kp*-MH258 suppression (y-
axis) versus principal component ('PC') (x-axis; ordered from right to left by variance explained
by each principal component).

Methods

Collection and whole genome sequencing of 848 gut commensals

Fecal samples were obtained from 28 human donors that fell within the age range of 18 to 63 with a median age of 35. Donors were selected as those with no antibiotic use in the past year, no known history of diabetes, colitis, autoimmune disease, cancer, pneumonia, dysentery, or cellulitis at time of consent. Institutions that approved protocols of fecal sample collection were Memorial Sloan Kettering (MSK) and the University of Chicago. Fresh fecal samples were immediately reduced in an anaerobic chamber upon collection and diluted and cultured on various growth media. Agar media types vary, but include any of the following: Columbia Blood Agar, Brain Heart Infusion + Yeast, Brain Heart Infusion + Mucin, Brain Heart Infusion + Yeast + Acetate or N-acetylglucosamine, reinforced Clostridial Agar, Peptone Yeast Glucose, Yeast Casitone Fatty Acids, Defined media M5. Colonies were selected and grown to be sufficiently turbid, 20% glycerol/PBS stocks were created and stored in a -80°C freezer.

Colonies were selected for whole-genome based on pyro-sequencing of the 16S region which provides a rough estimate of genus level designation. For each donor, only colonies that had a sequence identity threshold of less than 99% from CD-Hit (v. 4.8.1) were selected for whole-genome sequencing[82,83]. Bacterial genomic DNA was extracted using QIAamp DNA Mini Kit (QIAGEN) according to manufacturer's manual. The purified DNA was quantified using a Qubit 2.0 fluorometer. 1000ng of each sample was prepared for sequencing using the QIAseq FX DNA Library Kit (QIAGEN). The protocol was carried out for a targeted fragment size of 550bp. Sequencing was performed on the MiSeq or NextSeq platform (Illumina) with a paired-end (PE) kit in pools designed to provide 1-3 million PE reads per sample with read length of 250 or 150 bp.

Adapters were trimmed off with Trimmomatic (v0.39) with following parameters: the leading and trailing 3 bp of the sequences were trimmed off, quality was controlled by a sliding window of 4, with an average quality score of 15 (default parameters of Trimmomatic)[84]. Moreover,

any read that was less than 50 bp long after trimming and quality control were discarded. The

remaining high-quality reads were assembled into contigs using SPAdes (v3.15.4)[85].

Taxonomic classification of the assembled contigs was performed with the following

methods: (a) Kraken2 (v2.1.2; (b) full/partial length 16S rRNA gene from each isolated colony's

assembled contigs is extracted and input into BLASTn (v2.10.1+) to query against NCBI's RNA

RefSeq database[86–88]. Top five hits for each query are manually curated to determine an isolate's

identity, with identity and coverage cutoff both at 95%; (c) GTDB-Tk (v1.5.1)[89]. The final taxonomy

is determined by the consensus of the three methods. Any colony that did not match initial pyro-

sequencing taxonomy or lacked consensus was excluded from the commensal strain bank.

1156

Dimension-reduction of commensal strain bank

All gut commensal strains were annotated by their Prokka annotations and an alignment

was created (848 rows comprising commensal strains, 150181 columns comprising Prokka

annotated features)[90]. Each entry in the alignment is a '1' or a '0' indicating the presence or

absence of a specific feature in a particular bacterial proteome. Uniform Manifold Approximation

and Projection (UMAP) using default parameters (top 10 Principal Components with standard

clustering) was used to visualize relative position of each gut commensal strain (see **Extended**

Data Fig. 2).

1165

1166   *Algorithm for designing metagenomically diverse consortia from UMAP-based embedding*

1167   To design a bacterial consortium comprised of N strains, we perform the following steps.

1168   Given a UMAP plot of strains, the following algorithm was used to engineer synthetic microbial

1169   consortia.

1170

1171   *Step 1*: Create 10,000 communities randomly of size N. The ensemble of all 10,000

1172   communities of size N is represented as

$$C_{size\ N} = \{c_1, \dots, c_{10,000}\} \qquad (1)$$

1173

1174   *Step 2*: Each community, c_i, is defined by a set of *N* bacterial strains:

$$c_i = \{s_1, \dots, s_N\} \qquad (2)$$

1175

1176   where s_j is strain *j* in c_i. Compute all pairwise distances in the UMAP space for all strains

1177   in C_i. For instance, the pairwise distance between strain 1 and 2 is:

$$pd_{1,2} = dist(s_1, s_2) \qquad (3)$$

1178

1179   where 'dist' is the function that computes the distance between s_1 and s_2 in the UMAP

1180   space. We define the distribution of all pairwise distances for c_i as

$$PD_i = \{pd_{1,2}, pd_{1,3}, \dots, pd_{N-1,N}\} \qquad (4)$$

1181

1182   *Step 3*: Order PD_i for a given c_i from largest to smallest values, then compute the mean

1183   pairwise distance across the lower 30% of values comprising PD_i. We term this value the

1184   'mean adjusted dispersal'.

1185

1186   *Step 4*: Compute the mean adjusted dispersal for all communities in $C_{size\ N}$.

1187

1188   *Step 5*: Identify the community within the 10,000 communities comprising $C_{size\ N}$ with the

1189   maximum mean adjusted dispersal. This community is the designed community

1190   comprising *N* strains.

1191

For downsampling the 848 strains in our commensal strain bank to the 46 strains used to design

DMCs (see **Fig. 1A**), the above approach was used where *N* was set to 46 and the UMAP plot

used is shown in **Extended Data Fig. 2**. 46 was chosen as the maximum number of strains we

were confidently able to manipulate to create bacterial cultures. For creating the DMCs shown in

Fig. 1C, the above approach was used where *N* spanned a range from 2 to 29 strains and the

UMAP plot used is shown in **Fig. 1A**.

A schematic of the process is outlined for a community comprised of three strains in **Extended**

Data Fig. 3A.

1200

1201   _Creating the Klebsiella pneumoniae MH258 strain used in experiments_

1202   The _Kp_-MH258 isolate was previously described elsewhere[42]. For better in vitro and in

1203   vivo selection of this strain, _Kp_-MH258 was transformed by electroporation with pmCherry-sfGFP

1204   (86441; addgene).

1205

Evaluating the effect of DMCs on the fitness of MDR Kp-MH258

The 46 bacterial strains described in **Supplementary Table 1B** were individually

inoculated from a frozen stock into 900μL of BHI supplemented with cysteine 0.1% (BHIS)

previously reduced. Strains were incubated at 37°C in static conditions for 48h in anaerobiosis to

ensure that the most fastidious strains reach stationary phase. _K. pneumoniae_-MH258 sGFP was

also inoculated in the same conditions, but only 24h after commensal isolates inoculation due to

the fast growth capacity of this species and was incubated for 24h. All strain densities were

assessed by taking 100 μL of each culture and measuring OD_{600} in a Biotek Cytation 5. To build

all DMCs, isolates were inoculated in 900 μL of BHIS previously reduced in different combinations

with an initial OD_{600} of 0.001, so that the densest community reaches a maximum total initial OD_{600}

of approximately 0.05. _K. pneumoniae_ was added at the same initial OD_{600} of 0.001 (approximately

5 x 10^6 CFUs/mL) to all DMCs. Cultures were incubated at 37°C in static conditions and

anaerobiosis for 5 days. To _assess K._ pneumoniae abundance, 10 μL of each culture were

collected daily and homogenized in 90 μL of PBS and serially diluted. Diluted samples were plated

in BHIS with kanamycin (50μg/mL). Plates were incubated at 37°C overnight in aerobiosis. GFP

expressing _K. pneumoniae-MH258_ colony forming units (CFUs) were enumerated. In parallel,

100uL of each culture was also collected to recover the cell phase and the supernatants at 72h,

96h, and 120h. These samples were stored at -80°C to be later processed for shotgun

metagenomics and metabolomics.

1225

Training additive linear regression model on strain presence/absence for suppressing Kp-MH258

We used the LinearRegression module, available under scikit learn (sklearn.linear) python

package[91]. The intercept option was set to true which allowed the model to have a non-zero linear

intercept if necessary for quality fit. The input data (independent variables) was a vector of 46 1's

and 0's as shown in the matrix displayed in **Fig. 1C** corresponding to the pattern of presence-

absence for each DMC. The dependent variable for model training was the *Kp-MH258* log

abundance value for each of these 96 experiments. Both independent and dependent variables

associated with the original 96 experiments were used to train the linear model's coefficients and

intercept. After training, we predicated the *Kp*-MH258 log abundances for these 96 experiments.

The R^2 value and the Pearson correlation between the prediction and actual abundance values

were 0.844 and 0.92 respectively. We extracted and recorded this model's regression coefficients

for each strain. The larger magnitude of this value means a larger effect on the *Kp-MH258* log

abundance.

1239

Animal protocol and husbandry

All mouse experiments were performed in accordance with and approved by the

Institutional Animal Care and Use Committee of the University of Chicago under protocol 72599.

Male specific-pathogen-free C57BL/6J mice, aged 8 weeks to 10 weeks, from Jackson

Laboratories were used for all experiments. Mice were kept within a facility that maintained a

12 hour light and 12 hour dark cycle and controlled humidity (30–70%) and temperature (68–

79 °F). Mice were housed in sterile, autoclaved cages with irradiated feed (LabDiets 5K67) and

acidified, autoclaved water upon arriving at the on-site mouse facility. Mouse handling and cage

changes were performed by investigators wearing sterile gowns, masks and gloves in a sterile

biosafety hood. Mice were cohoused with their original shipment group until starting the

experiment.

1251

Preparation of mice stool samples for fecal microbiota transplant (FMT)

Fecal samples from 15-20 mice SPF mice from different cages (to increase sample

diversity) were collected to a 50 mL tube. Samples were transferred immediately to the anaerobic

chamber (anaerobic exposure was kept under 30 min). Samples were dissolved in 1 mL of PBS

20% glycerol 0.1% cysteine (previously filtered and reduced) per fecal pellet (1mL per ~20 mg of

fecal sample) using a mechanical pestle and vortexing. Samples were aliquoted in cryovials and

stored -80°C until use.

1259

An in-vivo mouse model of MDR Kp-MH258 infection

C57BL/6J male at 8-10 weeks of age were singly housed and placed under an antibiotic

regime (0.25g MNV – metronidazole, neomycin, vancomycin) in the drinking water (day 0). Four

days later, antibiotic treatment was halted and mice were placed on normal acidified water (day

4). Cages and food were also changed. On day 5 all mice were gavaged with 100μL of PBS

containing 500 CFUs of Kp-MH258, prepared as previously explained. On days 7, 8, and 9 mice

were gavaged with 100uL of either selected defined bacterial consortia, a fecal microbiota

transplant from naïve healthy mice, or PBS. Fecal samples were collected on days 0, 4, 7, 10,

12, 14, 16, and 21 (final day of the experiment) for 16s rRNA sequencing and on day 10 and 12

for metabolomics. These were immediately place on dry ice after collection and later stored at -

80°C. To assess for Kp-MH258 levels, fecal samples were collected on days 7, 10, 12, 14, 16,

and 21. Fecal samples were homogenized in 1mL of PBS and serially diluted. Undiluted and

diluted samples were plated in BHIS and kanamycin (50μg/mL).

1273

 Training an RF model and a deep learning-based model on DMC strain presence/absence as

 input and Kp-MH258 abundance at 120 hours of co-culture with DMC as output

For training the RF model, we used a RandomForestRegressor, available with scikit-learn

python package[91]. Tree Depth was set to 12 levels per tree, the number of trees was set to 100,

and the maximum number of features was set to "sqrt" (square-root of the number of strains total).

Out-of-bag error was measured by a combination of R^2 (where numbers less than 1 indicate

more error) and Mean Squared Error (where larger numbers indicate more error). To train and

validate our model, we randomly split our dataset into 90% training and 10% true-out-of-sample

100 times. The input data was a vector of 46 1's and 0's as shown in the matrix displayed in **Fig.**

1C corresponding to the pattern of presence-absence for each DMC. In each iteration, the

RandomForestRegressor was fit to the training set via 6-fold cross validation. Cross-validation

accuracy was measured through Pearson Correlation. The true out of sample set was then

predicted, and prediction accuracy was measured by computing Mean Squared Error and

Pearson Correlation of the predicted versus measured *K. pneumoniae* abundances after 120

hours of co-culture with the DMC. Feature Importance Scores for all features were observed and

stored. This process was repeated 100 times, and prediction accuracies and feature significance

scores were averaged. An additional RandomForestRegressor model was then trained on the

entirety of the dataset with 6-fold cross-validation. Cross-validation accuracy was measured by

calculating Mean Squared Error, Pearson Correlation, and R^2. Averaged prediction accuracies

and feature significance scores were used to estimate prediction error.

For training the deep learning-based model ('deep neural network', DNN), we used the

same input data format as described above. We next created 5000 Neural networks using

PyTorch each with two hidden layers (50 neurons going to 10 neurons) and a single output neuron

tasked with predicting the *Kp*-MH258 log abundance values. For each neural network, we first

performed a random 80% -20% training-validation split of the original 96 experiments. This neural

network was trained on its training split using a dynamic momenta-dependent backpropagation

algorithm where the gradient is calculated on the training split's loss alone while the momentum

parameter is reduced when the loss function on the validation split stagnates. This momentum-

based scheme allows for smooth (non-oscillatory) and faster training and reduces the chance of

our neural networks getting stuck in local optima.

1304

Defining SynCom15 through distillation of the RF model

Isolating eigenvector most associated with Kp-MH258 clearance.

The matrix in **Fig. 3B** was subject to Singular Value Decomposition (SVD) resulting in 46

principal components of data-variance (eigenvectors). We found that the first principal component

(PC1) was significantly associated with community complexity (**Supplementary Table 7B**). To

isolate the effect of *Kp*-MH258 clearance from community size, we first performed a series of

steps to 'regress out' the effect of community size. First, let x_i be the community size of DMC i.

Let y_i be the predicted *Kp*-MH258 clearance from the RF model for DMC i. A linear model is then

created regressing community size against *Kp*-MH258 clearance taking the form:

$$y_i = \beta_1 x_i + \beta_0 + \varepsilon \quad (5)$$

$$\hat{y}_i = \beta_1 x_i + \beta_0 \quad (6)$$

where $\hat{y}_i$ is the *Kp*-MH258 clearance of DMC i as a function of its size. The residuals of this

linear model are given by

$$\varepsilon_i = y_i - \hat{y}_i \quad (7)$$

where ε_i is the degree of *Kp*-MH258 clearance of DMC i after removing linearly modeled

information related to the size of the DMC. All principal components were regressed against r_i

and principal component 46 (PC46) was found to be the most significantly associated with

predicted, residualized *Kp*-MH258 clearance (**Supplementary Table 7C**).

1323

*Defining the matrix in **Fig. 3C***

Let $u \in \mathbb{R}^{46}$ be the vector of projections of each strain on eigenvector 46 of the matrix

defined in **Fig. 3B**.

Let s be the scalar value denoting the maximum value of u

$$A^{46x46} = \left(a_{ij}\right) \quad (8)$$

$$a_{ij} = d_{ij}^2 = ||u_i - u_j||^2 \quad (9)$$

Where $|| \cdot ||$ denotes the Euclidian norm on $\mathbb{R}^{46}$

$$A_{i,j} = \begin{bmatrix} 0 & \cdots & d_{46,1}^2 \\ \vdots & \ddots & \vdots \\ d_{1,46}^2 & \cdots & 0 \end{bmatrix} \quad (10)$$

1332

$$S_{i,j} = s - A_{i,j} = \begin{bmatrix} s & \cdots & s - d_{46,1}^2 \\ \vdots & \ddots & \vdots \\ s - d_{1,46}^2 & \cdots & s \end{bmatrix} \quad (11)$$

The resulting symmetric similarity matrix, $S_{i,j}$, with rows and columns indicating each strain and

each element representing the similarity between strain i and strain j describes how strains are

related to one another based on their projections along eigenvector 46. Hierarchical clustering

on the resulting similarity matrix was then performed to identify groups of strains. Strains that

are more similar are often found in communities that suppress *Kp-MH258* and those that are

more distant are rarely found in communities that suppress *Kp-MH258*.

1340

Creation of ex-vivo media

8–10-week-old wild-type male C57BL/6J mice were used for all studies. Mice were initially

obtained from The Jackson Laboratory and subsequently bred and raised in a GF isolator. After

removal from the GF isolator, mice were handled in a sterile manner and individually housed in

sealed negative pressure bio-containment unit isolators. Throughout breeding, mice were housed

within the University of Chicago Gnotobiotic Research Animal Facility (GRAF) and maintained at

a 12 hour light and 12 hour dark cycle and controlled humidity (30–70%) and temperature (68–

79 °F). Gnotobiotic mice were fed an ad libitum diet of autoclaved Teklad Global 18% Protein

Rodent Diet (Sterilizable) (2018S/2018SC). To create 'GF cecal content media', mice were

euthanized and cecal contents were collected, weighted, and homogenized in 10mL of sterile

distilled water on a of per gram of content. Cecal suspension was centrifuged, and supernatants

were filtered through a 0.22 mm filter. GF cecal extract media was stored at -80°C.

To create 'Ab-SPF cecal content media' C57BL/6J SPF male mice at 8-10 weeks of age

were singly housed and placed under an antibiotic regime (0.25g MNV – metronidazole,

neomycin, vancomycin) in the drinking water (day 0). Four days later, antibiotic treatment was

halted and mice were placed on normal acidified water (day 4). Cages and food were also

changed. On day 7 were euthanized and cecal contents were collected, weighted, and

homogenized in 10mL of sterile distilled water on a of per gram of content. Cecal suspension was

centrifuged, and supernatants were filtered through a 0.22 mm filter. Ab-SPF cecal extract media

was stored at -80°C.

1361

Determining engraftment of SynCom15 strains in SPF mice

To determine SynCom15 strain engraftment, 16s rRNA sequences from all 15 strains were

blasted against 16S rRNA sequences derived from fecal samples of antibiotic-treated SPF mice

gavaged with SynCom15 consortium. Fecal-derived sequences were assigned to a SynCom15

strain if their 16s rRNA percentage sequence identity was 100% with a minimum of a 95%

coverage.

1368

Determining structure of microbiota in infected SPF mice given saline, FMT, or SynCom15

DNA was extracted using the QIAamp PowerFecal Pro DNA kit (Qiagen). Before

extraction, samples were subjected to mechanical disruption using a bead beating method.

Briefly, samples were suspended in a bead tube (Qiagen) along with lysis buffer and loaded on a

bead mill homogenizer (Fisherbrand). Samples were then centrifuged, and supernatant was

resuspended in a reagent that effectively removed inhibitors. DNA was then purified routinely

using a spin column filter membrane and quantified using Qubit.

16S sequencing was performed for murine studies, where V4–V5 region within 16S rRNA

gene was amplified using universal bacterial primers—563F (5′-nnnnnnnn-NNNNNNNNNNNNN-

AYTGGGYDTAAA-GNG-3′) and 926R (5′-nnnnnnnn-NNNNNNNNNNNNN-CCGTCAATTYHT-

TTRAGT-3′), where 'N' represents the barcodes and 'n' are additional nucleotides added to offset

primer sequencing. Approximately 412-bp region amplicons were then purified using a spin

column-based method (Minelute, Qiagen), quantified and pooled at equimolar concentrations.

Illumina sequencing-compatible Unique Dual Index adapters were ligated onto the pools using

the QIAseq 1-step amplicon library kit (Qiagen). Library quality control was performed using Qubit

and TapeStation and sequenced on Illumina MiSeq platform to generate 2 × 250 bp reads.

Raw V4–V5 16S rRNA gene sequence data were demultiplexed and processed through

the dada2 pipeline (v1.18.0) into amplicon sequence variants (ASVs) with minor modifications in

R (v4.0.3)[92]. Specifically, reads were first trimmed at 190 bp for both forward and reverse reads

to remove low-quality nucleotides. Chimeras were detected and removed using the default

consensus method in the dada2 pipeline. Then, ASVs with length between 320 bp and 365 bp

were kept and deemed as high-quality ASVs. Taxonomy of the resultant ASVs was assigned to

the genus level using the RDP Classifier (v2.13) with a minimum bootstrap confidence score of

80[93].

1393

Metabolic profiling of DMCs

For metabolite extraction from liquid cultures, samples were incubated at −80 °C between

1 h and 12 h. Four volumes of methanol spiked with internal standards were added to each culture

supernatant. Samples were then centrifuged at −10 °C and 20,000 × g for 15 min followed by the

transfer of 100 µL of supernatant to pre-labelled mass spectrometer autosampler vials (MicroLiter,

09-1200).

For metabolite extraction from fecal samples, extraction solvent (80% methanol spiked

with internal standards and stored at -80 °C) was added at a ratio of 100 mg of material/mL of

extraction solvent in beadruptor tubes (Fisherbrand; 15-340-154). Samples were homogenized at

4 °C on a Bead Mill 24 Homogenizer (Fisher; 15-340-163), set at 1.6 m/s with 6 thirty-second

cycles, 5 seconds off per cycle. Samples were then centrifuged at -10 °C, 20,000 x g for 15 min

and the supernatant was used for subsequent metabolomic analysis.

Short chain fatty acids were derivatized as described by Haak *et al*. with the following

modifications[94]. The metabolite extract (100 µL) was added to 100 µL of 100 mM borate buffer

(pH 10) (Thermo Fisher, 28341), 400 µL of 100 mM pentafluorobenzyl bromide (Millipore Sigma;

90257) in Acetonitrile (Fisher;A955-4), and 400 µL of n-hexane (Acros Organics; 160780010) in

a capped mass spec autosampler vial (Microliter; 09-1200). Samples were heated in a

thermomixer C (Eppendorf) to 65 °C for 1 hour while shaking at 1300 rpm. After cooling to RT,

samples were centrifuged at 4 °C, 2000 x g for 5 min, allowing phase separation. The hexanes

phase (100 µL) (top layer) was transferred to an autosampler vial containing a glass insert and

the vial was sealed. Another 100 µL of the hexanes phase was diluted with 900 µL of nhexane

in an autosampler vial. Concentrated and dilute samples were analyzed using a GC-MS (Agilent

7890A GC system, Agilent 5975C MS detector) operating in negative chemical ionization mode,

using a HP-5MSUI column (30 m x 0.25 mm, 0.25 µm; Agilent Technologies 19091S-433UI),

methane as the reagent gas (99.999% pure) and 1 µL split injection (1:10 split ratio). Oven ramp

parameters: 1 min hold at 60 °C, 25 °C per min up to 300 °C with a 2.5 min hold at 300 °C. Inlet

temperature was 280 °C and transfer line was 310 °C. A 10-point calibration curve was prepared

with acetate (100 mM), propionate (25 mM), butyrate (12.5 mM), and succinate (50 mM), with 9

subsequent 2x serial dilutions.

Metabolites were also analyzed using GC-MS with electron impact ionization. The

metabolite extract (100 µL) mass spec autosampler vials (Microliter; 09-1200) and dried down

completely under nitrogen stream at 30 L/min (top) 1 L/min (bottom) at 30 °C (Biotage SPE Dry

96 Dual; 3579M). To dried samples, 50 µL of freshly prepared 20 mg/mL methoxyamine (Sigma;

226904) in pyridine (Sigma; 270970) was added and incubated in a thermomixer C (Eppendorf)

for 90 min at 30 °C and 1400 rpm. After samples are cooled to room temperature, 80 µL of

derivatizing reagent (BSTFA + 1% TMCS; Sigma; B-023) and 70 µL of ethyl acetate (Sigma;

439169) were added and samples were incubated in a thermomixer at 70 °C for 1 hour and 1400

rpm. Samples were cooled to RT and 400 µL of Ethyl Acetate was added to dilute samples. Turbid

samples were transferred to microcentrifuge tubes and centrifuged at 4 °C, 20,000 x g for 15 min.

Supernatants were then added to mass spec vials for GCMS analysis. Samples were analyzed

using a GC-MS (Agilent 7890A GC system, Agilent 5975C MS detector) operating in electron

impact ionization mode, using a HP-5MSUI column (30 m x 0.25 mm, 0.25 µm; Agilent

Technologies 19091S- 433UI) and 1 µL injection. Oven ramp parameters: 1 min hold at 60 °C, 16

°C per min up to 300 °C with a 7 min hold at 300 °C. Inlet temperature was 280 °C and transfer

line was 300 °C.

Data analysis was performed using MassHunter Quantitative Analysis software (version

B.10, Agilent Technologies) and confirmed by comparison to authentic standards. Normalized

peak areas were calculated by dividing raw peak areas of targeted analytes by averaged raw

peak areas of internal standards.

1443

Evaluating suppressive capacity of SynCom15 on various strains of K. pneumoniae, E. coli, and

C. difficile

The 15 bacterial strains that compose SynCom15 were individually inoculated from a

frozen stock into 900 µL of BHIS previously reduced. Strains were incubated at 37°C in static

conditions for 48h in anaerobiosis to ensure that the most fastidious strains reach stationary

phase. Pathogen strains of *K. pneumoniae* (MH258-sGFP, MH225, MH189, MH1867, and

Kp43816), *E. coli* (X43, T18002), VRE, and *C. difficile* (VPI, ST-49, R20291, ST-6, ST-75) were

also inoculated in the same conditions, but only 24h after commensal isolates inoculation due to

the fast growth capacity of these species and was incubated for 24h. All strain densities were

assessed by taking 100 μL of each culture and measuring OD_{600} in a Biotek Cytation 5. To build

SynCom15, SynCom15 minus commensal *E. coli* MSK19.28, Block 1, and Block 1 minus *E. coli*

MSK19.28, isolates were inoculated in 900 μL of GF cecal content media previously reduced with

an initial OD_{600} of 0.001, as described above. *K. pneumoniae, E. coli, VRE, or C. difficile* strains

were added at the same initial OD_{600} of 0.001 to the different DMCs or to media alone. Cultures

were incubated at 37°C in static conditions and anaerobiosis for 5 days. To *assess K.*

pneumoniae, VRE, and *C. difficile* abundance, 10 μL of each culture were collected at 24h, 96,

and 120h of culture and homogenized in 90 μL of PBS and serially diluted. Diluted samples were

plated in: BHIS with kanamycin (50mg/mL) for the *Kp*-MH258-sfGFP strain; BHIS with ampicillin

(100mg/mL) for *Kp*-MH225, *Kp*-MH189, *Kp*-MH1867, Kp43816, *E. coli* X43, and T18002; BHIS

with Vancomycin (50mg/mL)/ Streptomycin (100mg/mL) for the VRE strain; and BHIS with

norfloxacin [6g/L], moxalactam [16g/L], D-Cycloserine [62.5g/L], and Cefoxitin [16g/L] for all *C.*

difficile strains. Plates were incubated at 37°C overnight in aerobiosis for all strains except for the

C. difficile strains, which dilutions, plating and incubation of the plates were done under

anaerobiosis. CFUs were enumerated 24h later for the *K. pneumoniae* and *E. coli* strains, and

48h later for VRE and *C. difficile* strains. In parallel, 100 μL of each culture was also collected to

recover the cell phase and the supernatants at 24h, 96h, and 120h. These samples were stored

at -80°C to be later processed for shotgun metagenomics and metabolomics.

1471

Elucidating indirect, pH dependent mechanism of suppression

For testing SynCom15 dependency on contact for pathogen suppression, the 15 bacterial

strains that compose SynCom15 were individually inoculated from a frozen stock into 900 μL of

BHIS previously reduced. Strains were incubated at 37°C in static conditions for 48h in

anaerobiosis to ensure that the most fastidious strains reach stationary phase. Pathogen strains

of *K. pneumoniae* (MH258-sGFP and MH225), *E. coli* (X43), and *C. difficile* (VPI and ST-49) were

also inoculated in the same conditions, but only 24h after commensal isolates inoculation due to

the fast growth capacity of these species and was incubated for 24h. Pathogen cultures were

diluted into BHIS or GF cecal content media in 24-well plates, and DMCs cultures were diluted

into upper chambers of Transwells inserts (0.4 μ m ThinCert for 24-well plates, greiner bio-one)

such that each strain started at an OD_{600} of 0.001. To *assess K. pneumoniae*, *E. coli*, and *C.*

difficile abundance, 10 μL of each culture were collected at 24h, 96h, and 120h of culture and

homogenized in 90 μL of PBS and serially diluted. Samples were diluted, plated, incubated, and

stored as described above.

DMCs plus pathogen or pathogen alone cultures were also collected to measure pH after

120h (pH probe Fisherbrand). When indicated culture's pH was adjusted to 5.0 using HCl or 7.0

using NaOH. Cell-free spent medium from these cultures (with or without adjusted pH) were

obtained using 0.22 μ m Millex-GP syringe filters (Millipore). Supernatants were used as culture

media to inoculate the different pathogens. A supernatant derived from a culture that contains a

specific pathogen was only used to inoculate the same pathogen. As before, 10 μL of each culture

were collected at 24h, 96h, and 120h of culture and homogenized in 90 μL of PBS and serially

diluted for plating the appropriated plates and conditions for the different pathogens, as indicated

before. In parallel, 100 μL of each culture was also collected to recover the cell phase and the

supernatants at 24h, 96h, and 120h. These samples were stored at -80°C to be later processed

for shotgun metagenomics and metabolomics.

1497

Comparison of SynCom15 with microbiotas of healthy human donors

To investigate the presence of SynCom15 strains in samples from healthy human donors,

SynCom15 strains taxonomic names were searched in the 22 fecal samples obtained from the

DFI 22 human donors. For SynCom15 strain unclassified to species level *Bifidobacterium* sp., the

most closely related species annotated by GTDB with an 98.21% ANI (*Bifidobacterium*

pseudocatenulatum) was used[89,95].

1504

Training an RF- and DNN-based model for predicting Kp-MH258 suppression from metabolite

content

Data Handling:

Our metabolite data consists of 118 metabolites measured as well as the Kp-MH258 CFUs

at the end of 120 hours. The Kp-MH258 CFU measurement has a resolution of 1000 CFU. or any

CFU value recorded as 0, we regularized it by adding a 1000. The log of this regularized CFU as

well as the log fold changes are then unit norm-centered and used to train both Random Forest

(RF) and Deep Neural Network (DNN) Models. Since norm-centering is an invertible

transformation, we undo this transformation on the final model prediction value to get the

predicted log suppression.

1515

1516   *Training RF model:*
1517
For training the RF model, we used a RandomForestRegressor, available with scikit-learn

python package. Tree Depth was set to 12 levels per tree, the number of trees was set to 5000,

and the maximum number of features was set to "sqrt" (square-root of the number of strains total).

For any tree in a RF model, a randomly chosen 1/3rd of the training datapoints is internally withheld

as validation set. Since the RF model internally performs this cross validation using decisions

from trees that don't use a given datapoint, and the number of trees we have is large compared

to the number of datapoints, the out-of-bag (OOB) accuracy score—the average accuracy of

prediction on a datapoint across these trees—is our metric for generality of the model. For RF

model for predicting Kp-MH258 suppression_built on metabolites, the OOB score is **0.20** on a

scale of 0 to 1 strongly suggesting that metabolites are a poor choice of basis. The score was

stable to any additional trees added to the model. With the model trained, we noted the R^2 score

on the training set. Using this RF model, we predicted and recorded the log suppression on the

60 true-out-of-sample experiments.

1531

Training DNN based model:

For training the metabolite based deep learning-based model DNN, we used the same

input data format as described above. We next created 5000 Neural networks using PyTorch each

with two hidden layers (50 neurons going to 10 neurons) and a single output neuron tasked with

predicting the *Kp-MH258* log abundance values. For each neural network, we first performed a

random 80% -20% training-validation splits. This neural network was trained on its training split

using a dynamic momenta-dependent backpropagation algorithm where the gradient is calculated

on the training split's loss alone while the momentum parameter is reduced when the loss function

on the validation split stagnates. This momentum-based scheme allows for smooth (non-

oscillatory) and faster training and reduces the chance of our neural networks getting stuck in

local optima. Once trained, we defined the generalization capacity of each neural network as the

R^2 value on the prediction of the validation split with floor set at 0. Higher R^2 means that the neural

network has properly generalized over the validation split. These 5000 trained neural networks

were used to predict log suppression on the 60 true-out-of-sample experiments. We aggregated

the result of this neural network ensemble experiment by weighing each prediction with the

generalization capacity of the respective neural network to calculate the weighted mean and

standard deviation of the predicted *Kp-MH258* log abundance values for all DMCs included

(training/validation set as well as out-of-sample set).

Supplementary Tables

1551

Supplementary Table 1. Data pertaining to **Extended Data Figures 1** and **2**.

Supplementary Table 2. Phylogenetic description of 46 strains used to construct bacterial

communities.

Supplementary Table 3. Data pertaining to design of DMCs.

Supplementary Table 4. Data pertaining to **Fig. 2B** and **2C**.

Supplementary Table 5. Validation data for ML models.

Supplementary Table 6. Data regarding validating ML models on out-of-sample DMCs.

Supplementary Table 7. Defining SynCom15 using machine learning and statistical inference

of model constraints; data pertaining to **Fig. 4B** and **4D**.

Supplementary Table 8. Taxonomic compositional analysis of mouse experiments.

Supplementary Table 9. Metabolomic data associated with DMCs.

Supplementary Table 10. Mechanism of *Kp*-MH258 suppression by SynCom15.

Supplementary Table 11. SynCom15 suppression of phylogenetically distinct pathogens.

Supplementary Table 12. Fractional abundance of SynCom15 strain-associated species in

fecal samples of healthy donors.

Supplementary Table 13. Data and analysis associated with **Extended Data Fig. 14**.

Supplementary Table 14. A summary of select previous microbiome design efforts.

Supplementary Discussion

On constructing consortia based on phylogenetic diversity using the UMAP-based embedding

As our goal was to design phylogenetically diverse consortia, we sought to do so in an algorithmic, quantitative manner. Typically, phylogeny is reflected as a hierarchy of relatedness between bacterial species[89,95–97]. While such a view provides a reasonable way to classify bacteria, hierarchies are intrinsically difficult to use as a parametrization of defining distance[32]. Conveyed using intuition: what does 'different' mean when objects are hierarchically related to each other? By virtue of being hierarchically related, two objects can be different but be relatively more similar than two other bacteria. As such, ensuring diverse sampling off a tree is a challenging concept to encode into an algorithm.

As an alternative to hierarchical parametrizations of ensembles, non-linear embeddings compress hierarchical relationships into an *N*-dimensional space where *N* is determined by the user[98–100]. In our case, we compressed the phylogeny of our strain bank into two UMAP-based dimensions, yielding a dimension-reduced space. Typically, such spaces are used purely for visualization. The reason for this is that distance in the space cannot be ascribed to hamming distance between objects due to the non-linear compression intrinsic to UMAP. What does this consideration practically mean for downstream analyses? First, new data cannot be projected into an existing UMAP space without a reparameterization of the space. Second, two UMAP plots comprised of distinct data cannot be compared to each other from a geometric standpoint. Third, computing distance within the UMAP space is meaningless with respect to the relative similarity between two objects.

However, in our case, the UMAP-based embedding directly captured phylogenetic differences between groups of strains. Note, this did not mean that distance in the UMAP scaled with phylogenetic distance, but that, by definition, large distances in the UMAP space corresponded to sampling of different phylogenies. As such, using dispersion in the UMAP space was a data-driven way to create diverse consortia. We note that this approach is likely imperfect and future work will be needed to create diverse ensembles directly off the inferred phylogenetic tree.

Model Distillation:

The motivation behind distillation in our work stems from a broader challenge in machine learning (ML): predictive models often succeed in fitting data but remain opaque with respect to the principles, if any, they have internalized. In fields such as computer vision and natural language processing, this has spurred the development of 'model distillation' frameworks which attempt to extract simplified rules, constraints, or low-dimensional representation from complex black-box models[49]. The underlying ethos is to bridge the gap between prediction and understanding. That is, move from models that merely output correct answers and towards models that reveal generative rules that can be reused across contexts.

We adapted this philosophy in our work to understand how the RF model was learning to predict the suppressive capacity of DMCs. A key difference in our approach from the typical approaches of distillation used in AI is that we did not focus on the architecture of the models themselves, but rather inferred constraints that the models learned within their latent representation of the data. The motivation behind this choice was inspired by previous studies that have interrogated the structure of biological systems using statistical inference spanning the fields of protein science, neuroscience, genome science, microbiome science, and cancer genomics[10,47,68,74,80,81,101–103]. The essence of the idea has been the following. Evolution creates diversity through the process of iterative selection and variation. However, the constraints governing this process are opaque and unintuitive from the perspective of a researcher and therefore we are not informed about a 'forward model' of biology. We therefore collect the outputs of this process—extant diversity—create an alignment to compare these outputs, and statistically

learn constraints describing the outputs[104].
The result of this analysis provides an
understanding of what component parts in
complex biological systems that arise from
the process of evolution are relatively
important versus unimportant for the
survival of the system (**Fig. SD1A**). In
analogy to this approach, we derived a
synthetic generating function in the form of
a machine-learning model that can be used
to engineer consortia with desired *Kp-*
MH258 suppression. As such, we could
mirror the process of creating extant
diversity from the evolutionary process by
creating lots of consortia that were
predicted to achieve pathogen suppression
by running the ML model. While the logic of
suppression would not be revealed
observing the composition of any single
consortia, the collection of suppressive
consortia would contain conserved
taxonomic signatures reflecting that all
consortia are predicted to suppress *Kp-*
MH258. Thus, statistical decomposition of
the consortia reflecting the 'extant diversity'
of communities that suppress *Kp*-MH258
would reveal key taxonomic features
learned by the synthetic generating function (**Fig. SD1B**).

Figure SD1. (A) Evolution creates extant diversity. The ensemble of extant diversity is subject to statistical decomposition to learn constraints about the evolutionary process. **(B)** In our work, a synthetic generating function was created in the form of an ML model. This ML model then created >5,000 consortia that were predicted to achieve pathogen suppression of greater than 10^5 CFUs/ml. The resulting consortia are analogous to 'extant diversity'—synthetic communities that survived the selection criteria of needing to achieve a high degree of pathogen suppression. The resulting consortia are subject to statistical decomposition to learn what constraints, if any, are latent within the statistical generating function.

1648
Assessing the compressive power of our approach
The process by which we converged on SynCom15 as a community that clears *K.*
pneumoniae involved a six-step process: <u>C</u>ompress/<u>D</u>esign-<u>B</u>uild-<u>T</u>est-<u>L</u>earn-<u>I</u>nfer. Within this
process there were two points at which biological and experimental information was used for
progressing through steps: (i) execution of the <u>C</u>ompress step utilized whole genomes of strains
to reduce the size of the design space from spanning 848 possible strains to 46 possible strains
and (ii) execution of the <u>D</u>esign-<u>B</u>uild-<u>T</u>est-<u>L</u>earn-<u>I</u>nfer steps required screening 96 DMCs. We
therefore sought to compute the equivalent of a 'compression ratio' for converging on a single
complex community from a bank of 848 strains. In evaluating computational algorithms, the
compression ratio is a measure of data complexity prior to compression relative to after
compression. As our process considered biological information in the form of bacterial genome
sequences and experiments, we normalized the compression ratio by the amount of information
needed to perform the compression. We therefore defined an 'effective' compressive power as

1662
$$C = \frac{\left(\frac{A}{B}\right)}{I} \qquad (1)$$

where C is the effective compressive power of a process, A is the complexity of data prior to
compression, B is the complexity of data after compression, and I is the information needed for
compression from A to B.
For our first step, we reduced the strain bank from 848 strains to 46 strains representative
of the full phylogenetic diversity by genome sequencing each of the 848 strains, annotating each
genome by their gene content, and performing dimension-reduction via a UMAP analysis.
Therefore, the total complexity prior to compression was $2^{848}/2$, the total complexity after

compression was $2^{46}/2$, and the information needed to be collected for compression were all base
pairs of the 848 commensal strains (8.5×10^{11} basepairs). Considering these values, the effective
compression of our first step was $\sim 10^{230}$—a substantial compression driven by the sizeable drop
in complexity of the strain bank. For our second step, we used the diversity of the 46 strains to
create 96 DMCs, 23 experiments to demonstrate the reproducibility of our assay, and 60
experiments to evaluate the 'out-of-sample' DMCs—a total of 179 experiments. Therefore, the
total complexity prior to compression was $2^{46}/2$, the total complexity after compression was 1
(SynCom15), and the information needed to be collected for compression 179 experiments.
Considering these values, the effective compression for our second step was $\sim 10^{11}$. Collectively,
this analysis showed that despite the apparently immense amount of data reflected in the whole
genome sequences of 848 bacterial strains, this complexity is offset by many orders of magnitude
through Constraint Distillation.
1682
Contextualizing Constraint Distillation amongst prior microbiome design strategies
A broad range of strategies have been proposed for designing synthetic microbial
communities with desired functions. As summarized in **Supplementary Table 14**, most efforts fall
along distinct categories of design approaches. In one category are approaches that treat
communities as reaction networks whose collective behavior can be captured by dynamical
systems models. These studies typically use extensive information about individual strains—
growth in mono- and co-culture, inferred or measured metabolic qualities, curated genome-scale
metabolic models, or experimentally parametrized interaction matrices—to simulate community
dynamics and identify candidate consortia for a target function. Such models can provide
mechanistic insight and can enable *in silico* exploration of design space, but in practice require
substantial prior knowledge, are difficult to parametrize at scale, and are often limited to specific
conditions or considering small numbers of strains. Typically, newer approaches of statistical
modeling—machine learning and artificial intelligence approaches like deep-learning—have been
used within this category to augment Design-Build-Test-Learn paradigms. Although powerful,
these methods are typically data hungry, requiring hundreds to thousands of tested communities,
and are often optimized for prediction within a single experimental context. As a result,
generalization of rules across environments, particularly regarding synthetic design of consortia
towards gut microbiome-related applications, has remained largely untested using this strategy.
A second category is rooted in empirical reduction of natural communities. Here, complex
microbiotas (e.g. whole fecal communities) are iteratively diluted, fractionated, or rationally
trimmed to yield a defined consortium. This strategy implicitly treats the natural ecosystem as a
blueprint: composition and structure of the starting community constrain which taxa are even
considered, and down-sampling tends to preserve high-abundance or strongly interacting
members. While this has yielded designed consortia, this strategy is predicated on the existence
of natural microbiomes that have already solved the problem. Thus, design towards possibly novel
target functions is not possible through using only this approach. Moreover, this strategy is not
meant to create principles of design—the capacity to design a new, functional microbiome in a
predictive manner.

A third category is leveraging existing sequencing efforts between human or mouse
cohorts to perform comparative analysis and define bacteria that differentiate the cohorts from
each other. A typical example would be to compare a 'diseased' cohort with a 'healthy' cohort then
create a consortium based on these differences. This strategy has also been historically effective
at creating consortia but requires (i) the presence of taxonomic differences between cohorts, (ii)
the taxonomic differences faithfully map to differences in phenotype, and (iii) the method of
comparative analysis is accurately capturing taxonomic differences from a statistical perspective
without compressive loss of information. Like the second category, this strategy is also not meant
to create principles of consortia design such that new consortia can be engineered to function in
a predictive manner.

Figure SD2. Previous efforts at community design can be conceptualized into three categories with respect to use of prior knowledge: (i) Use of natural communities to subsequently perform down-sampling, (ii) Statistical characterization of natural communities that reflect phenotypes of interests from cohorts, and (iii) Use of strain phenotypes/strain-strain interactions. Our manuscript illustrates the utility of using genome sequences of strains to Compress the space of design and Design metagenomically diverse consortia without any behavioral or functional knowledge of the constituent strains or communities. These communities are then Built, Tested for function, a model is Learned between strain presence/absence and output function, Inference is performed to distill taxonomic constraints learned by the model. Together these steps define the process we term 'Constraint Distillation'.

Constraint Distillation occupies a distinct point in this landscape (**Fig. SD2**). Our approach
deliberately dispenses with several underlying foundations of prior work. We do not (i) model
dynamical interactions between community members, (ii) encode mechanism-specific features
related to *K. pneumoniae* suppression, (iii) rely on pathway-level annotations or curated metabolic
networks at the scale of individual strains or designed communities, (iv) use human microbiome
composition as a template, or (v) start from naturally occurring communities that already display
the desired phenotype. Instead, we represent each designed community only by a binary vector
of strain presence/absence, where the communities were initially designed solely based on
genome-encoded diversity. From this sparse and functionally agnostic description along with a
modestly sized screen of 96 consortia, we trained an ML model and then inferred functionally
relevant statistical constraints learned by the model that defined SynCom15. Conceptually, our
work complements rather than replaces prior efforts. Mechanistic, dynamical, and approaches
embracing '-omics' remain essential for elucidating how specific consortia function and for de-
risking translation of live biotherapeutics into clinical settings. However, the SynCom15 case study
demonstrates that generative design of a sparse, broadly effective community does not require
these ingredients as prerequisites. Thus, Constraint Distillation offers a distinct mode of
microbiome engineering: one that treats community design primarily as an inference problem on
compositional space with mechanistic understanding as an output rather than an input of the
design process.

References

1740

1.   Lawson, C. E. *et al.* Common principles and best practices for engineering microbiomes.
Nat. Rev. Microbiol. **17**, 725–741 (2019).
2.   Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. *Science* **361**,
469–474 (2018).
3.   Faith, J. J., Ahern, P. P., Ridaura, V. K., Cheng, J. & Gordon, J. I. Identifying gut microbe-
host phenotype relationships using combinatorial communities in gnotobiotic mice. *Sci.*
Transl. Med. **6**, 220ra11 (2014).
4.   Inda, M. E., Broset, E., Lu, T. K. & de la Fuente-Nunez, C. Emerging frontiers in microbiome
engineering. *Trends Immunol.* **40**, 952–973 (2019).
5.   Venturelli, O. S. *et al.* What is the key challenge in engineering microbiomes? *Cell Syst.* **14**,
85–90 (2023).
6.   Faust, K. & Raes, J. Microbial interactions: from networks to models. *Nat. Rev. Microbiol.*
10, 538–550 (2012).
7.   Zomorrodi, A. R. & Segrè, D. Synthetic Ecology of Microbes: Mathematical Models and
Applications. *J. Mol. Biol.* **428**, 837–861 (2016).
8.   Alivisatos, A. P. *et al.* MICROBIOME. A unified initiative to harness Earth's microbiomes.
Science **350**, 507–508 (2015).
9.   Dubilier, N., McFall-Ngai, M. & Zhao, L. Microbiology: Create a global microbiome effort.
Nature Publishing Group UK http://dx.doi.org/10.1038/526631a (2015)
doi:10.1038/526631a.
10. Raman, A. S. *et al.* A sparse covarying unit that describes healthy and impaired human gut
microbiota development. *Science* **365**, (2019).
11. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut
microbiome. *Cell* **185**, 3617-3636.e19 (2022).
12. Tian, S. *et al.* A designed synthetic microbiota provides insight to community function in
Clostridioides difficile resistance. *Cell Host Microbe* **33**, 373-387.e9 (2025).
13. Brugiroux, S. *et al.* Genome-guided design of a defined mouse microbiota that confers
colonization resistance against Salmonella enterica serovar Typhimurium. *Nat. Microbiol.* **2**,
16215 (2016).
14. Osbelt, L. *et al.* Klebsiella oxytoca causes colonization resistance against multidrug-
resistant K. pneumoniae in the gut via cooperative carbohydrate competition. *Cell Host*
Microbe **29**, 1663-1679.e7 (2021).
15. Oliveira, R. A. *et al.* Klebsiella michiganensis transmission enhances resistance to
Enterobacteriaceae gut invasion by nutrition competition. *Nat Microbiol* **5**, 630–641 (2020).
16. Furuichi, M. *et al.* Commensal consortia decolonize Enterobacteriaceae via ecological
control. *Nature* **633**, 878–886 (2024).
17. Honda, K. *et al.* Rationally-defined microbial consortia suppress multidrug-resistant
proinflammatory Enterobacteriaceae via ecological control. *Res. Sq.* (2023)
doi:10.21203/rs.3.rs-3462622/v1.
18. Auchtung, J. M., Preisner, E. C., Collins, J., Lerma, A. I. & Britton, R. A. Identification of
simplified microbial communities that inhibit Clostridioides difficile infection through
dilution/extinction. *mSphere* **5**, (2020).
19. Lawley, T. D. *et al.* Targeted restoration of the intestinal microbiota with a simple, defined
bacteriotherapy resolves relapsing Clostridium difficile disease in mice. *PLoS Pathog.* **8**,
e1002995 (2012).
20. Petrof, E. O. *et al.* Stool substitute transplant therapy for the eradication of Clostridium
difficile infection: "RePOOPulating" the gut. *Microbiome* **1**, 3 (2013).
21. Gowda, K., Ping, D., Mani, M. & Kuehn, S. Genomic structure predicts metabolite dynamics
in microbial communities. *Cell* **185**, 530-546.e25 (2022).

22. Clark, R. L. *et al.* Design of synthetic human gut microbiome assembly and butyrate
production. *Nat. Commun.* **12**, 3254 (2021).
23. Baranwal, M. *et al.* Recurrent neural networks enable design of multifunctional synthetic
human gut microbiome dynamics. *Elife* **11**, (2022).
24. Stein, R. R. *et al.* Computer-guided design of optimal microbial consortia for immune
system modulation. *Elife* **7**, (2018).
25. Atarashi, K. *et al.* Treg induction by a rationally selected mixture of Clostridia strains from
the human microbiota. *Nature* **500**, 232–236 (2013).
26. Dsouza, M. *et al.* Colonization of the live biotherapeutic product VE303 and modulation of
the microbiota and metabolites in healthy volunteers. *Cell Host Microbe* **30**, 583-598.e8
(2022).
27. Spragge, F. *et al.* Microbiome diversity protects against pathogens by nutrient blocking.
Science **382**, eadj3502 (2023).
28. Morin, M. A., Morrison, A. J., Harms, M. J. & Dutton, R. J. Higher-order interactions shape
microbial interactions as microbial community complexity increases. *Sci. Rep.* **12**, 22640
(2022).
29. Gallardo-Navarro, O., Aguilar-Salinas, B., Rocha, J. & Olmedo-Álvarez, G. Higher-order
interactions and emergent properties of microbial communities: The power of synthetic
ecology. *Heliyon* **10**, e33896 (2024).
30. Feng, L. *et al.* Identifying determinants of bacterial fitness in a model of human gut
microbial succession. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 2622–2633 (2020).
31. Front Matter. *Inverse Problem Theory and Methods for Model Parameter Estimation* i–xii
Preprint at https://doi.org/10.1137/1.9780898717921.fm (2005).
32. Doran, B. A. *et al.* Subspecies phylogeny in the human gut revealed by co-evolutionary
constraints across the bacterial kingdom. *Cell Syst.* (2025) doi:10.1016/j.cels.2024.12.008.
33. Skwara, A. *et al.* Statistically learning the functional landscape of microbial communities.
Nat Ecol Evol **7**, 1823–1833 (2023).
34. Lee, K. K., Park, Y. & Kuehn, S. Robustness of microbiome function. *Curr. Opin. Syst. Biol.*
36, 100479 (2023).
35. Gubbins, D. Book reviews. *Geophys. J. Int.* **94**, 167–168 (1988).
36. Clermont, G. & Zenker, S. The inverse problem in mathematical biology. *Math. Biosci.* **260**,
11–15 (2015).
37. Cocco, S., Feinauer, C., Figliuzzi, M., Monasson, R. & Weigt, M. Inverse statistical physics
of protein sequences: a key issues review. *Rep. Prog. Phys.* **81**, 032601 (2018).
38. Sanchez-Lengeling, B. & Aspuru-Guzik, A. Inverse molecular design using machine
learning: Generative models for matter engineering. *Science* **361**, 360–365 (2018).
39. Mora, T. & Bialek, W. Are biological systems poised at criticality? *J. Stat. Phys.* **144**, 268–
302 (2011).
40. Park, H., Li, Z. & Walsh, A. Has generative artificial intelligence solved inverse materials
design? *Matter* **7**, 2355–2367 (2024).
41. Odenwald, M. A. *et al.* Bifidobacteria metabolize lactulose to optimize gut metabolites and
prevent systemic infection in patients with liver disease. *Nat Microbiol* **8**, 2033–2049 (2023).
42. Xiong, H. *et al.* Distinct Contributions of Neutrophils and CCR2+ Monocytes to Pulmonary
Clearance of Different Klebsiella pneumoniae Strains. *Infect. Immun.* **83**, 3418–3427
(2015).
43. Hernández, U., Posadas-Vidales, L. & Espinosa-Soto, C. On the effects of the modularity of
gene regulatory networks on phenotypic variability and its association with robustness.
Biosystems. **212**, 104586 (2022).
44. Hartwell, L. H., Hopfield, J. J., Leibler, S. & Murray, A. W. From molecular to modular cell
biology. *Nature* **402**, C47-52 (1999).

45. Newman, M. E. J. Modularity and community structure in networks. *Proc. Natl. Acad. Sci.*
U. S. A. **103**, 8577–8582 (2006).

46. Halabi, N., Rivoire, O., Leibler, S. & Ranganathan, R. Protein sectors: evolutionary units of
three-dimensional structure. *Cell* **138**, 774–786 (2009).

47. McLaughlin, R. N., Jr, Poelwijk, F. J., Raman, A., Gosal, W. S. & Ranganathan, R. The
spatial architecture of protein function and adaptation. *Nature* **491**, 138–142 (2012).

48. Espinosa-Soto, C. & Wagner, A. Specialization can drive the evolution of modularity. *PLoS*
Comput. Biol. **6**, e1000719 (2010).

49. Frosst, N. & Hinton, G. Distilling a neural network into a soft decision tree. *arXiv [cs.LG]*
(2017) doi:10.48550/arXiv.1711.09784.

50. Boix-Adsera, E. Towards a theory of model distillation. *arXiv [cs.LG]* (2024).

51. Arrieta, A. B. *et al.* Explainable Artificial Intelligence (XAI): Concepts, taxonomies,
opportunities and challenges toward responsible AI. *arXiv [cs.AI]* (2019)
doi:10.48550/arXiv.1910.10045.

52. Stutz, M. R. *et al.* Immunomodulatory fecal metabolites are associated with mortality in
COVID-19 patients with respiratory failure. *Nat. Commun.* **13**, 6615 (2022).

53. de Porto, A. P. *et al.* Fecal metabolite profiling identifies critically ill patients with increased
30-day mortality. *Sci. Adv.* **11**, eadt1466 (2025).

54. Sorbara, M. T. *et al.* Inhibiting antibiotic-resistant Enterobacteriaceae by microbiota-
mediated intracellular acidification. *J. Exp. Med.* **216**, 84–98 (2019).

55. Bachman, M. A. *et al.* Genome-Wide Identification of Klebsiella pneumoniae Fitness Genes
during Lung Infection. *MBio* **6**, e00775 (2015).

56. Silver, R. J. *et al.* Amino acid biosynthetic pathways are required for Klebsiella pneumoniae
growth in immunocompromised lungs and are druggable targets during infection.
Antimicrob. Agents Chemother. **63**, e02674-18 (2019).

57. Vornhagen, J. *et al.* The Klebsiella pneumoniae citrate synthase gene, gltA, influences site
specific fitness during infection. *PLoS Pathog.* **15**, e1008010 (2019).

58. Pickard, J. M., Zeng, M. Y., Caruso, R. & Núñez, G. Gut microbiota: Role in pathogen
colonization, immune responses, and inflammatory disease. *Immunol. Rev.* **279**, 70–89
(2017).

59. Oliphant, K. & Allen-Vercoe, E. Macronutrient metabolism by the human gut microbiome:
major fermentation by-products and their impact on host health. *Microbiome* **7**, 91 (2019).

60. Horrocks, V., King, O. G., Yip, A. Y. G., Marques, I. M. & McDonald, J. A. K. Role of the gut
microbiota in nutrient competition and protection against intestinal pathogen colonization.
Microbiology **169**, 001377 (2023).

61. Baktash, A. *et al.* Mechanistic insights in the success of fecal Microbiota transplants for the
treatment of Clostridium difficile infections. *Front. Microbiol.* **9**, 1242 (2018).

62. Gehrig, J. L. *et al.* Effects of microbiota-directed foods in gnotobiotic animals and
undernourished children. *Science* **365**, (2019).

63. Delannoy-Bruno, O. *et al.* Evaluating microbiome-directed fibre snacks in gnotobiotic mice
and humans. *Nature* **595**, 91–95 (2021).

64. Kennedy, M. S. *et al.* Diet outperforms microbial transplant to drive microbiome recovery in
mice. *Nature* 1–9 (2025).

65. Plata, G., Srinivasan, K., Krishnamurthy, M., Herron, L. & Dixit, P. Designing host-
associated microbiomes using the consumer/resource model. *mSystems* **10**, e0106824
(2025).

66. Thompson, J. C., Zavala, V. M. & Venturelli, O. S. Integrating a tailored recurrent neural
network with Bayesian experimental design to optimize microbial community functions.
PLoS Comput. Biol. **19**, e1011436 (2023).

67. Russ, W. P., Lowery, D. M., Mishra, P., Yaffe, M. B. & Ranganathan, R. Natural-like function
in artificial WW domains. *Nature* **437**, 579–583 (2005).

68. Socolich, M. *et al.* Evolutionary information for specifying a protein fold. *Nature* **437**, 512–
518 (2005).
69. Bialek, W. & Ranganathan, R. Rediscovering the power of pairwise interactions. *arXiv [q-*
bio.QM] (2007) doi:10.48550/arXiv.0712.4397.
70. Cavagna, A. *et al.* New statistical tools for analyzing the structure of animal groups. *Math.*
Biosci. **214**, 32–37 (2008).
71. Cavagna, A. *et al.* Scale-free correlations in starling flocks. *Proc. Natl. Acad. Sci. U. S. A.*
107, 11865–11870 (2010).
72. Caballero, S. *et al.* Cooperating Commensals Restore Colonization Resistance to
Vancomycin-Resistant Enterococcus faecium. *Cell Host Microbe* **21**, 592-602.e4 (2017).
73. Sulaiman, J. E. *et al.* Elucidating human gut microbiota interactions that robustly inhibit
diverse Clostridioides difficile strains across different nutrient landscapes. *Nat. Commun.*
15, 7416 (2024).
74. Russ, W. P. *et al.* An evolution-based model for designing chorismate mutase enzymes.
Science **369**, 440–445 (2020).
75. Yeh, A. H.-W. *et al.* De novo design of luciferases using deep learning. *Nature* **614**, 774–
780 (2023).
76. Anishchenko, I. *et al.* De novo protein design by deep network hallucination. *Nature* **600**,
547–552 (2021).
77. Watson, J. L. *et al.* De novo design of protein structure and function with RFdiffusion.
Nature **620**, 1089–1100 (2023).
78. Wang, J. *et al.* Scaffolding protein functional sites using deep learning. *Science* **377**, 387–
394 (2022).
79. Brixi, G. *et al.* Genome modeling and design across all domains of life with Evo 2. *bioRxiv*
2025.02.18.638918 (2025) doi:10.1101/2025.02.18.638918.
80. Zaydman, M. A. *et al.* Defining hierarchical protein interaction networks from spectral
analysis of bacterial proteomes. *Elife* **11**, (2022).
81. Behera, V. *et al.* Conserved principles of spatial biology define tumor heterogeneity and
response to immunotherapy. *bioRxiv* 2024.10.18.619136 (2024)
doi:10.1101/2024.10.18.619136.
82. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).
83. Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. CD-HIT Suite: a web server for clustering and
comparing biological sequences. *Bioinformatics* **26**, 680–682 (2010).
84. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).
85. Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A. & Korobeynikov, A. Using SPAdes De
Novo Assembler. *Curr. Protoc. Bioinformatics* **70**, e102 (2020).
86. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol. **20**, 257 (2019).
87. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).
88. O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current status,
taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733-45 (2016).
89. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).
90. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069
(2014).
91. Liaw, A. & Wiener, M. Classification and Regression by randomForest. (2007).
92. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.
Nat. Methods **13**, 581–583 (2016).

93. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid
1943   assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*
1944   **73**, 5261–5267 (2007).

1945   94. Haak, B. W. *et al.* Impact of gut colonization with butyrate-producing microbiota on
1946   respiratory viral infection following allo-HCT. *Blood* **131**, 2978–2986 (2018).

1947   95. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a
1948   phylogenetically consistent, rank normalized and complete genome-based taxonomy.
1949   *Nucleic Acids Res.* **50**, D785–D794 (2022).

1950   96. Guindon, S., Delsuc, F., Dufayard, J.-F. & Gascuel, O. Estimating Maximum Likelihood
1951   Phylogenies with PhyML. in *Bioinformatics for DNA Sequence Analysis* (ed. Posada, D.)
1952   113–137 (Humana Press, Totowa, NJ, 2009).

1953   97. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
1954   phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

1955   98. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection
1956   for Dimension Reduction. *arXiv [stat.ML]* (2018).

1957   99. Kobak, D. & Berens, P. The art of using t-SNE for single-cell transcriptomics. *Nat. Commun.*
1958   **10**, 5416 (2019).

1959   100.Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat.*
1960   *Biotechnol.* **37**, 38–44 (2018).

1961   101.Stringer, C., Pachitariu, M., Steinmetz, N., Carandini, M. & Harris, K. D. High-dimensional
1962   geometry of population responses in visual cortex. *Nature* **571**, 361–365 (2019).

1963   102.Eisen, J. A. Phylogenomics: improving functional predictions for uncharacterized genes by
1964   evolutionary analysis. *Genome Res.* **8**, 163–167 (1998).

1965   103.Pellegrini, M., Marcotte, E. M., Thompson, M. J., Eisenberg, D. & Yeates, T. O. Assigning
1966   protein functions by comparative genome analysis: protein phylogenetic profiles. *Proc. Natl.*
1967   *Acad. Sci. U. S. A.* **96**, 4285–4288 (1999).

1968   104.Stein-O'Brien, G. L. *et al.* Enter the Matrix: Factorization Uncovers Knowledge from Omics.
1969   *Trends Genet.* **34**, 790–805 (2018).

Data Availability. The datasets generated in our study are available within Supplementary Tables. Metagenomic data generated from profiling human fecal microbiomes used in this study are publicly available on NCBI under BioProject ID PRJNA838648. 16S data generated from mouse experiments used in this study will be publicly available on NCBI under BioProject ID PRJNA1074807. Raw data files associated with metabolomic data used in this study will be found on MassIVEW repository MSV000094183.

Code availability. All code was written in either Python or R; code for all analysis will be found on Github (https://github.com/aramanlab/Oliveira_et_al_2024).

Figures. Figure panels associated with data were generated using either the Prism software (v10.2.0), various available packages in R, or Python. Figure schemes were generated using BioRender (BioRender.com) or Adobe Illustrator.

Acknowledgements. We thank members of the Pamer, Kuehn, and Raman Laboratories for helpful discussion. We thank D. Pincus, M. Mani, A. Murugan, R. Ranganathan, and E. Pamer for helpful discussion. We thank members of the biobank, genomics, and metabolomics core services within the Duchossois Family Institute (DFI) at the University of Chicago and E. Pamer for their help in isolate collection, sequencing, and metabolomic profiling of all samples described in this manuscript. We thank the late E. Littman for his bioinformatic contributions. This work was supported by the Duchossois Family Institute (DFI) at the University of Chicago, the Dr. Ralph and Marian Falk Medical Research Trust, and NIH grant RM35GM146702.

Author Information. E.M. aided in many of the experiments involving pathogen suppression and elucidating mechanism of SynCom15. B.P. aided in modeling of strain presence/absence with linear regression and deep-learning models as well as metabolomic-based modeling of consortia.

K.L. wrote the code to implement the UMAP-based design strategy. M.Y. wrote the code to create the RF model based on strain presence/absence and scored 100,000 DMCs using the resulting RF model. R.Y.C. performed statistical analysis of the 100,000 DMCs to ultimately yield SynCom15; C.T. aided in the assay of all DMCs for evaluating their capacity to clear *Kp*-MH258 as well as the effect of selected communities in an *in vivo* setting. F.H. and V.A. aided with establishing the plate-based assay to evaluate clearing *Kp*-MH258; R.P. and K.S. aided in evaluating the suppressive capacity of SynCom15 against pathogen strains of *K. pneumoniae*, *E. coli*, and *C. difficile*. R.R. aided in analysis of taxonomic composition of fecal pellets procured from mice and analysis of fecal samples collected from healthy human donors. R.O. conducted all experiments involving DMCs across different conditions, all *in vivo* experiments, all experiments involving characterization of SynCom15, provided material for metabolomic and genomic analysis; S.K. and A.S.R. conceived of the statistical approach for community design involving creating a statistical model from an ensemble of diverse consortia; R.Y.C. and A.S.R. conceived of performing statistical inference on the learned model to distill model constraints. R.O. and A.S.R. conceived of the *in vitro* and *in vivo* experiments, experiments involved in understanding mechanism of SynCom15 action, the analysis of healthy human samples, metabolic profiling of DMCs, and sensitivity analysis of Constraint Distillation as a process. R.O. and A.S.R. wrote the manuscript.

Ethics Declarations. Patents (63/543,XXX & 63/543XXX) related to this research have been filed by the University of Chicago with S.K., R.O., and A.S.R. as inventors.

Materials and Correspondence. Author to whom correspondence and materials request should be addressed is A.S.R.