Widespread evolution of poricidal flowers: A striking example of morphological convergence across flowering plants

Avery Leigh Russell[1†*], Rosana Zenil-Ferguson[2*], Stephen L. Buchmann[3], Diana D. Jolles[4], Ricardo Kriebel[5], Mario Vallejo-Marín[6]

[1] Department of Biology, Missouri State University, Springfield, MO, United States

[2] Biology Department, University of Kentucky, Lexington, KY, United States

[3] Department of Ecology and Evolutionary Biology and Department of Entomology, University of Arizona, Tucson, AZ, United States

[4] Department of Biological Sciences, Plymouth State University, Plymouth, NH, United States

[5] Department of Botany, Institute of Biodiversity Science and Sustainability, California Academy of Sciences, San Francisco, CA, United States

[6] Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

*co-first authors

†Corresponding author

Department of Biology, 910 S John Q Hammons Pkwy, Temple Hall, Missouri State University

Springfield, MO 65897, USA

607 220 4432

†AveryRussell@MissouriState.edu

Total word count for main text: 3834; total tables: 3; total figures: 6 (all color)

Running title: Convergent evolution of poricidal flowers

Summary

- The evolution of tube-like floral morphologies that control pollen release via small openings (functionally poricidal flowers) represents a taxonomically and geographically widespread instance of repeated and independent evolution of a functionally similar morphology. Poricidal flowers are often closely associated with buzz pollination by bees, but we lack an updated angiosperm-wide survey of their phylogenetic distribution.

- We identify all presently known angiosperm genera containing poricidal flowers via a literature survey. We determined the phylogenetic distribution of poricidal flowers and minimum number of independent gains and losses via a genus-level and species level angiosperm-wide phylogeny. We estimated if evolution of poricidal flowers is associated with changes in speciation/extinction via diversification rate analyses.

- Poricidal flowers occur across at least 87 angiosperm families and 635 genera containing > 28,000 species. At the genus level, more than 200 independent gains and 145 independent losses of poricidal flowers occurred. Across angiosperms, genus-level analyses suggest poricidal flower evolution is associated with lower net diversification rates (origination-extinction). Species-level analyses argue no diversification linked to poricidal morphology but within the two focal families we found family-specific diversification effects.

- We present different hypotheses for the functional significance of poricidal flowers, including its association with buzz pollination, and suggest future directions for studies elucidating the extent to which morphological and functional convergence of poricidal flowers are correlated.

Keywords: Bees, buzz pollination, convergent evolution, diversification analysis, pollen, poricidal anthers, poricidal flowers.

Introduction

Patterns of convergent evolution are of fundamental interest in evolutionary biology. Broad convergence on functionally equivalent morphology across disparate groups is particularly common in plants, as in the case of succulent morphology or carnivorous pitcher plant traps, and has helped elucidate shared selective pressures, mechanisms, and developmental biases (Bennici, 2003; Ellison & Gotelli, 2009; Thorogood *et al.*, 2018). Likely the most geographically and taxonomically widespread pattern of convergent evolution is tubular floral morphology that restricts pollen dispersal via small openings, i.e., functionally poricidal floral morphology (hereafter 'poricidal flowers'; Fig. 1)(Vallejo-Marin & Russell, 2024). Previous studies estimate that approximately 22,000 plant species possess poricidal anthers ('poricidal species'), spread across 72 families and 544 genera (Buchmann, 1983). Additionally, flower traits that control pollen dispersal are key to angiosperm reproductive success (Hargreaves *et al.*, 2009) and many poricidal plant species are also agriculturally important, such as tomatoes, cranberries, blueberries, and kiwis (Cooley & Vallejo-Marín, 2021). Yet the phylogenetic distribution and evolutionary patterns of poricidal flowers has not been formally characterized and the latest published taxonomic survey of poricidal flowers is now nearly 45 years old (Buchman, 1978; Buchmann, 1983).

Poricidal flowers share a common functional morphology in which pollen is held within tubular floral structures with small openings (e.g., pores, valves, or slits; hereafter 'pores') through which the pollen is dispersed. In most cases, the tubular structure is composed of modified enlarged anthers, but may alternatively consist of modifications to the corolla or even the entire flower (Fig. 1) (Harris, 1905; Michener, 1962; Buchmann, 1983; De Luca & Vallejo-Marin, 2013; Russell *et al.*, 2017; Dellinger *et al.*, 2019b; Vallejo-Marin, 2019a). Poricidal flowers have been studied for over a century (Harris, 1905) and are frequently associated with a suite of floral characteristics (Buchmann, 1983; Renner, 1989; De Luca & Vallejo-Marin, 2013). Most poricidal plant species are animal pollinated, nectarless, and have dry, loose pollen (reduced pollenkitt lipids) and many are associated with enantiostyly (style handedness) and heteranthery (multiple kinds of stamens) (Buchmann, 1983; Vallejo-Marin *et al.*, 2010). Furthermore, poricidal flowers are often closely associated with the buzz pollination interaction (although not all poricidal flowers are buzz pollinated; see Vallejo-Marin and Russell (2024) for some exceptions), in which pollination is nearly exclusively performed by bee species capable of rapidly contracting their indirect flight muscles while on the flower, generating strong vibrations that expel pollen onto the bee's body (Macior, 1964; Buchmann, 1983; Proenca, 1992; Vallejo-Marin, 2019a; Vallejo-Marin & Russell, 2024). Indeed, poricidal flowers are thought to be a key pollen resource for these buzzing bees, particularly in tropical regions (Delgado *et al.*, 2023). Poricidal floral morphologies may facilitate speciation by filtering out less effective

pollinators (Cardinal *et al.*, 2018). Yet because of specialization on a subset of pollinators, poricidal

morphology may make these taxa more vulnerable to extinction (Renner, 1989; Pacheco Filho *et al.*,

2015; Dellinger *et al.*, 2019a; Vallejo-Marin, 2019b). Consequently, poricidal floral morphologies are

consistently suggested to be adaptations to pollinators that might be key traits affecting patterns of

speciation and extinction.

Previous attempts at comprehensively studying poricidal flowers were restricted to species

with pored anthers presumed to be buzz pollinated (Harris, 1905; Buchmann, 1983). However,

functionally poricidal floral morphologies are more diverse (Fig. 1). Additionally, clades of poricidal

species can comprise thousands of species (e.g., *Solanum*, Melastomataceae) and the rate of gain

and loss of poricidal floral morphologies has yet to be quantified. Although poricidal species are

extremely widespread, clades are often all poricidal or not and thus studies of small phylogenetic

clades suffer from the limitations of rarely seeing evolutionary transitions to and from poricidal

flowers. Consequently, large-scale analyses are needed to allow generalization and provide sufficient

statistical replication to test the effect of poricidal flower morphologies on diversification (Beaulieu

& O'Meara, 2018; Zenil-Ferguson *et al.*, 2019; Joly & Schoen, 2021). However, large-scale

phylogenetic analyses bring their own challenges (Helmstetter *et al.*, 2022). First, it is challenging to

obtain information on multiple traits for thousands of species (i.e., presence or absence of poricidal

flowers). Second, well resolved mega-phylogenies are required. Fortunately, recent angiosperm-

wide phylogenies have become available (Sauquet & Magallón, 2018; Jin & Qian, 2019; Janssens *et*

al., 2020). Third, running complex diversification models in large phylogenies is computationally

demanding and proven methods continue to be developed to speed up large-scale and complex

analyses. Finally, the effects of a single trait in a large tree can be confounded with heterogeneity in

the process of diversification by other unmeasured factors (Beaulieu & O'Meara, 2018; Beaulieu &

O'Meara, 2019; Donoghue & Edwards, 2019).

Here, we investigate the evolutionary history of poricidal flowers across angiosperms to

address the following questions. **1)** What proportion of angiosperms possesses poricidal flowers? **2)**

How many times have poricidal flowers been gained and lost over the evolutionary history of

flowering plants at the genus level? **3)** Is the presence of poricidal flowers associated with changes in

origination of genera and diversification rates (origination-extinction)? And **4)** Is the presence of

poricidal flowers associated with diversification rates (speciation-extinction) across all angiosperms

and within some key families? We therefore conducted an extensive literature survey to identify all

presently known angiosperm genera containing species with poricidal flowers. We then used an

angiosperm-wide phylogeny at the genus level (as recently done in Meseguer *et al.*, 2022) to

determine the phylogenetic distribution of poricidal flowers, the geologic ages of poricidal lineages,

and the number of independent gains and losses of functionally poricidal floral morphology before

the origination of genera. We also conducted diversification rate analyses at the species level to

estimate if the evolution of poricidal flowers is associated with changes in speciation/extinction.

Finally, we present an overview of both classic and novel hypotheses for why poricidal flowers

evolve and suggest directions for future research about the extent to which convergence in poricidal

flower morphology is accompanied by convergence in function.

Material and Methods

Literature survey

We defined "poricidal flowers" as flowers with tube-like floral structures (typically formed by

individual anthers or groups of anthers, but sometimes by staminodes, sepals or corolla), restricting

pollen release via an opening that can be described as a pore, small slit, or valve. We conducted our

literature search between 2015 and November 2020 for English, German, and Portuguese language

articles describing all presently known genera of functionally poricidal plant species. Sources

included online books, online search engines (Web of Science, Google Scholar), dissertations, and

primary research papers. Search terms included "anther", "stamen", or "flower" and "apical pore",

"dehisce", "dehiscence", "dehiscent", "introrse", "longitudinal", "porandrous", "porate", "pore",

"pored", "poricidal", "porose", "porous", "short slit", "thecal pores", "valve", or "valvular". The

search was performed at the genus level because numerous well characterized genera are

frequently monomorphic for the presence/absence of poricidal morphology (e.g., *Solanum*, *Senna*,

Miconia) and information for all species in rarer and/or tropical genera is frequently sparse

(although we also recognize that monomorphism within genera should not be taken for granted and

there is evidence of potential exceptions, e.g., Renner, 1989; Gavrutenko *et al.*, 2020). For each

genus, we documented the flower parts involved in forming the tubular structures that comprise

poricidal flower morphology, as well as how pollen is released from poricidal anthers (Fig. 2, Table 1;

Supporting Information Table S1). We also confirmed non-poricidal morphology for monomorphic

genera when morphological information was available, but generally did not catalog the

accompanying literature.

Angiosperm phylogeny and estimates of species number per genus

To estimate the incidence of poricidal taxa across angiosperms we cross-referenced our 638

poricidal genera against three angiosperm-wide databases. First, we used The Plant List (2013),

which contained 271,405 species, 12,645 genera, and 396 families of angiosperms, after filtering out

infraspecific classifications (i.e., subspecies and varieties). Next, we obtained two recent angiosperm

time-calibrated phylogenies from the literature. The first phylogeny is the Nelsen et al. (2018) genus-

level phylogeny of angiosperms, which contains 10,785 genera in 409 families. Second, we used the

Jin and Qian (2019) species-level, time-calibrated phylogeny of vascular plants, available through the

R package *PhyloMaker* (Jin & Qian, 2019). This phylogeny contains 417 families, 10,363 genera and

74,531 species of angiosperms.

Phylogenetic comparative methods

State dependent diversification and origination analyses

We used two time-tree phylogenies to perform a hidden-state dependent speciation and extinction

analysis at the genus level. First, we used the R package V.PhyloMaker by Jin and Qian (2019) that

relies on the Smith and Brown (2018) angiosperm phylogenetic tree to obtain a phylogenetic tree at

the genus level. The Smith and Brown (2018) tree was primarily created using mega tree

methodology, that is, matching different large angiosperm phylogenies to generate a larger and

more complete tree for genus-level data, but cannot be used for species-level data due to lack of

resolution at that level. Therefore, for the angiosperm wide genus-level analysis, we pruned the tree

to one species per genus (for a similar approach see Meseguer *et al.*, 2022). The second tree we

used was the Meseguer et al. (2022) genus level-phylogeny, also created from the Smith and Brown

(2018) tree. The published Meseguer et al. (2022) phylogenetic tree was presented as a bifurcating

tree. However, we noted that when we pruned the Smith and Brown (2018) phylogeny we had to

resolve polytomies in the genus-level tree, and we did this at random using the function multid2di()

in R package ape (Paradis *et al.*, 2004). This is an important detail, because the algorithm used to

resolve polytomies might change the estimation of diversification rates, and this is the main

difference between the Meseguer *et al.* 2022 and our pruned tree. For both genus-level phylogenies

we matched a total of 10,328 genera. Each tree was matched to binary states of poricidal status

(0=not poricidal, 1=poricidal) using treeplyr (Uyeda & Harmon, 2020), using the genus-level

character state obtained from the literature search. In addition, we chose four families from the

Smith and Brown (2018) phylogenetic tree (Solanaceae, Melastomataceae, Ericaceae, and

Fabaceae), and for each of the four families we obtained the species-level phylogeny and matched it

with poricidal status.

153   The matching trees and states on the tips of the tree were entered as inputs for state-

dependent diversification models. Using RevBayes (Höhna *et al.*, 2016), we wrote a binary hidden

state speciation and extinction model HiSSE (Beaulieu & O'Meara, 2016) using a graphical model

approach (Jordan, 2004) to estimate rates of net origination (origination of a genus minus extinction

rate) for poricidal and not poricidal states associated with two hidden states A and B. We fitted this

model for the two genus level phylogenies, ours and the Meseguer *et al.* (2022) phylogeny for all

angiosperms, and for the family level phylogenies.

The HiSSE model is used to identify both differences in net diversification between poricidal

and not poricidal states, but also differences given by hidden states A and B. The difference between

hidden states is of particular interest, especially in large phylogenies, because it represents the

potential for diversification due to unmeasured factors (Beaulieu & O'Meara, 2016). Thus, estimating

this difference disentangles the effect of poricidal morphology on diversification from other

potential sources of diversification. We only fitted HiSSE and not the simplest version without the

hidden states (BiSSE, Maddison *et al.*, 2007), because in a large tree our null hypothesis should

always be one that has heterogeneity in the rates of diversification due to unobserved factors, and

not one where the trait of interest has all the signal. As discussed in the recent synthesis of SSE

(state-dependent speciation and extinction) models, phylogenies with large samples always find that

HiSSE is a preferred model over BiSSE (Helmstetter *et al.*, 2023).

We performed a second HiSSE analysis matching our information of poricidal states to the

species-level phylogeny of Janssens et al. 2020. The Janssens et al. (2020) time-calibrated phylogeny

was specifically designed to address species-level macroevolution analyses, which is the main

difference with the Brown and Smith (2018) tree. Although the latter tree has more species

sampling, the uncertainty of the placement of the species is much larger than the Janssens et al.

(2020) tree, hence our preference for Janssens et al. (2020) for our species-level analysis. Based on

matK (incl. trnK) and rbcL. plastids this phylogeny has 36,101 taxa representing 8,399 genera and is

dated using 56 angiosperm fossils. We matched this phylogeny to our species-level poricidal data

and managed to match 28,979 taxa. In addition, we found 267 small genera where all species had

the same tip value, but none of them were present in the tree. Hence, we added the trait value as

representative of the genus to maximize the information when using the R package phyndr (Pennell

et al., 2016), resulting in a phylogeny of 29,246 tips with poricidal binary states. This represents the

largest plant phylogeny to date where a HiSSE analysis has been performed.

A Bayesian framework was used to obtain posterior probability densities via Markov chain

Monte Carlo (MCMC) for all HiSSE model parameters for genus-level and species-level phylogenies.

We checked for convergence of our MCMC algorithm by running our inference twice and observing

stationarity in the posterior distributions, as well as matching of convergence values in software

TRACER (Rambaut *et al.*, 2018). These two runs were not combined, and we retained only one run

for the results. Additionally, we verified that the effective sample size (ESS) for each parameter was

higher than 200. In the species-level analyses, convergence of the MCMC is very difficult due to

numerical precision at calculating likelihoods of the model. In this case we reached an ESS of more

than 200 for the posterior distribution of the HiSSE after an 11-month long MCMC run,

acknowledging that there are many local maxima for the posterior distribution that are very close,

which makes estimation of diversification rates difficult. However, all these local maxima have the

same consistent result in diversification, which is discussed in the Results section. The proportion of

genera or species sampled was calculated according to the number of genera listed in The Plant List

(2013), and the state of root was fixed to non-poricidal in the angiosperm-wide analysis and

unknown for the family-level ones.

Ancestral state reconstruction and estimation of minimum number of transitions

We implemented ancestral state reconstruction in RevBayes using a HiSSE model, as proposed by

Freyman and Höhna (Freyman & Höhna, 2018) for the genus level phylogenetic tree from Jin and

Qian (2019) matched with our poricidal status data. To infer which of four states (non-poricidal A,

non-poricidal B, poricidal A, poricidal B) is an internal node, we calculated the marginal posterior

probability of each internal node and chose the state that maximized that probability as shown in

Fig. 3 (Zenil-Ferguson *et al.*, 2019). Given these estimates for the internal states we calculated from

node to node whether there has been a transition from not poricidal to poricidal or vice versa. This

approach allows us to estimate the minimum number of expected transitions. An approximation to

the exact number would require stochastic mapping that is numerically challenging when scaling a

HiSSE model to large phylogenetic trees (Freyman & Höhna, 2019; Zenil-Ferguson *et al.*, 2023). All

results and ancestral reconstructions are illustrated using the R package RevGadgets (Tribble *et al.*,

2021). Ancestral reconstructions were not computationally feasible for the species-level

phylogenetic tree HiSSE analysis due to the large phylogeny size.

Results

Distribution of poricidal flowers across taxonomic levels

In a seminal paper, Buchmann (1983) estimated that 15,000-20,000 species of angiosperms in 72

families and 544 genera have anthers that dehisce through apical pores or slits (poricidal anthers).

Here, we found that poricidal flowers, including species with poricidal anthers, are reported from 87

families and 638 genera (Fig. 2, Table 1; Supporting Information Table S1). These values correspond

to approximately 21% of plant families and 5% of genera included in three different data sets of

flowering plants analyzed here (Table 1). Using the 271,405 species of angiosperms in the largest

database (The Plant List), which includes 635 of 638 genera identified in our study, yields an upper

estimate of 28,218 poricidal species (5.02% of genera and 10.39% of species) (Table 1). We obtained

a similar value from the other database that includes species-level information (Jin & Qian, 2019),

with an upper estimate of 10.93% (8,149 of 74,531) poricidal species and 5.06% (525 of 10363)

poricidal genera. Our results therefore suggest that poricidal flower morphologies have evolved in a

fifth of families and up to one in ten angiosperm species.

Different types of poricidal flowers across taxonomic levels

Flower parts forming the tubular structures that comprise poricidal flower morphology include

poricidal anthers (an estimated 97% of poricidal species), corollas, or other functionally poricidal

floral parts (i.e., staminodes or sepals) (Table 2). Additionally, how pollen is released from poricidal

anthers is particularly diverse, although such classifications are morphological, rather than

developmental, and may therefore be quite inaccurate. Morphological variation includes pores (60

families and 441 genera), short terminal slits (46 families and 125 genera), valves (15 families and 74

genera), and introrse dehiscence forming terminal pores or short slits (16 families and 53 genera)

(Supporting Information Table S1). A given poricidal flower morphology is rarely restricted to a

particular family or genus and even multiple kinds of poricidal anthers are known within a genus

(Endress, 1996).

Diversification rates and transitions between poricidal and non-poricidal flowers

Ancestral character state reconstruction on the angiosperm wide HiSSE analysis at the genus level

with our matched phylogeny from Jin and Qian (2019) indicated a minimum of 200 transitions from

non-poricidal to poricidal flowers and 145 reverse transitions (poricidal to non-poricidal), when

aggregated across hidden states (Fig. 3). This HiSSE analysis also suggests differences in rates of net

origination (origination rate of genera is calculated as the birth rate minus the death rate per state in

the HiSSE model) with and without poricidal flowers, depending on the state of the hidden character

(Fig. 4). Specifically, the net diversification rate of poricidal genera is on average 26% lower than for

non- poricidal taxa but only for the B state (credible intervals: 0.218 ± 0.0122 vs. 0.295 ± 0.0074,

respectively), and similarly 31% lower when in combination with hidden state A (credible intervals:

0.0119 ± 0.0013 vs. 0.0172 ± 0.0011, for non-poricidal vs. poricidal, respectively) (Fig. 4).

Interestingly, poricidal state B has a faster net diversification than non-poricidal state A, indicating

that some, but not all poricidal lineages can have faster net diversification than some non-poricidal

lineages (Fig. 3). Furthermore, poricidal genera appear early in angiosperm evolution and are

substantially more common in older nodes versus more recent nodes (Fig. 5). However, both of

these results should be interpreted with caution (for instance, because we do not test time-

dependent diversification; Louca & Pennell, 2020).

Yet the relationship between poricidal flowers and origination rates varies when analyzed at

lower taxonomic levels (Supporting Information Fig. S1). In the species-level Solanaceae HiSSE

analysis, poricidal flowers diversified at a higher rate compared to non-poricidal flowers, but again,

this relationship varied with the state of the hidden character. In the HiSSE analysis of Ericaceae, we

found a higher diversification rate of poricidal flowers in combination with hidden character state A,

but lower when combined with character state B, indicating a strong context-dependence effect of

poricidal morphologies (Fig. S1). For the two other families analyzed at the species level

(Melastomataceae and Fabaceae), models did not converge after running the analyses for several

months due to high uncertainty on the branch lengths in the Melastomataceae and Fabaceae trees

and thus we lack results for these groups (additional discussion in Supporting Information). Using the

Meseguer et al. (2022) phylogeny, we found that poricidal states have faster net diversification for

poricidal in states A and B (Supporting Information Fig. S2). This difference might be due to

differences in how polytomies were resolved in the Meseguer et al. phylogeny. When using this

phylogeny, many local maxima were identified in the output of the MCMC algorithm, and when we

took the global maximum, we found that the ancestral state reconstruction at many nodes is highly

uncertain, indicating that the inference using this tree is highly uncertain as well.

For the species-level HiSSE analyses using the Jin and Qian (2019) phylogenetic tree with our

data, we found that there are no differences in net diversification (speciation minus extinction rates)

between non-poricidal and poricidal states (Fig. 6A). Instead, we only found differences between

hidden state net diversification rates with probability 0.99 (Fig. 6A), indicating that diversification

patterns come from unobserved factors and not our poricidal states. Traditionally, a model selection

procedure is used to compare HiSSE models against character independent models (CID) to test if

differences in diversification stem from hidden states. However, this is not necessary when using a

Bayesian framework, as the posterior distribution of HiSSE can be transformed to test the character

independent model hypothesis (see supporting information for a demonstration in Figs. S3 and S4).

This result is consistent with the synthesis on state-dependent diversification presented in

Helmstetter et al. (2023), where traits fitted to phylogenies with large sampling typically have no

correlation with diversification rates. Our diversification results in the species-level analysis along

with our genus-level HiSSE analysis indicate that the evolution of poricidal state might be key in the

generation of new angiosperm genera, but once those genera are established, poricidal state is not

as critical in diversification at the species level. Furthermore, we found that transitions from non-

poricidal to poricidal states are on average 33x and 6.6x slower than from poricidal back to non-

poricidal in A and B hidden states, respectively (95% credible intervals: non-poricidal to poricidal:

$q_{01}^A = (5.28x10^{-5}, 1.13x10^{-4})$, and $q_{01}^B = (2.65x10^{-4}, 5.88x10^{-4})$; poricidal back to non-

poricidal: $q_{10}^A = (1.75 \times 10^{-3}, 3.81 \times 10^{-3})$, and $q_{10}^B = (1.42 \times 10^{-3}, 4.52 \times 10^{-3})$), (Fig. 6B). This

pattern of asymmetry in transition rates from a "specialized" poricidal state back to a "generalized"

non-poricidal state was also found in recent state-dependent diversification analyses (Day *et al.*,

2016; Zenil-Ferguson *et al.*, 2023), where more evidence of evolving out of ecological specialization

has been found.

295 Discussion

Taxa with functionally poricidal floral morphology comprise an estimated 10% of angiosperm species

and this morphology has at least 200 independent origins at the generic level (Table 1, S1). Evidently,

there is relatively strong selective pressure to evolve floral poricidal morphology. Given the diversity

of angiosperm clades and ecosystems in which poricidal flowers have arisen, likely multiple non-

mutually exclusive hypotheses explain their overall evolution (Vallejo-Marin & Russell, 2024). For

instance, because poricidal flowers hold their pollen within tube-like morphology, their pollen is

likely much better protected from environmental stressors such as ultraviolet radiation and rain,

relative to the pollen of non-poricidal lineages (Table 3; c.f., Gottsberger & Silbauer-Gottsberger,

1988). Similarly, pollen is a nutrient-rich substrate for yeast and bacteria and poricidal morphology

might limit microbial antagonists gaining access to the pollen (Table 3). Poricidal morphology is also

frequently physically toughened, such as the endothecial thickenings and calcium oxalate crystal

druses common to poricidal anthers (Gerenday & French, 1988; Carrizo García *et al.*, 2008; Burrieza

et al., 2010). Such modifications may conceivably reduce the theft of protein-rich pollen by chewing

herbivores (Table 3; Coté & Gibernau, 2012; but see Paiva, 2021). Accordingly, we might expect the

evolution and occurrence of poricidal flowers to be more common among angiosperm taxa living in

regions experiencing greater pollen-related biotic and abiotic stressors.

Classical hypotheses for the evolution of poricidal morphology have highlighted the role of

pollinators (Vallejo-Marin & Russell, 2024). Most, but not all species with poricidal flowers are

pollinated by floral buzzing bees, which are prolific consumers of pollen (Michener, 1962;

Buchmann, 1983; Hargreaves *et al.*, 2009; Danforth et al. 2019; Vallejo-Marin, 2022). Poricidal

morphology might control the rate of pollen release, and thus reduce the tempo of pollen removal

by pollen-consuming pollinators (Table 3; e.g., Larson & Barrett, 1999; De Luca *et al.*, 2014; Brito *et*

al., 2016). Likewise, poricidal morphology may result in more effective dispersal of pollen to

conspecifics by pollinators, by enhancing the precision of pollen placement on the pollinator body

and/or by filtering more generalized and less effective pollinators (Table 3; e.g., Pacheco Filho *et al.*,

2015; Mesquita-Neto *et al.*, 2018; Kemp & Vallejo-Marin, 2021; Vallejo-Marin & Russell, 2024).

However, pollinator-mediated selection alone appears insufficient to explain the evolution of

poricidal morphology across diverse angiosperm clades. Poricidal species have diverse mating

systems, from selfing to outcrossing, and many poricidal flowers are wind pollinated, not animal

pollinated (e.g., in Poaceae and Halophytaceae;Pozner & Cocucci, 2006; Table S1). Although

vibration can be a key mechanism of wind-mediated pollen release (Timerman & Barrett, 2018), and

could function analogously to floral buzzing by bees for these poricidal species, the functional

benefits of poricidal morphology likely go well beyond pollen release, as discussed above and in

Vallejo-Marin and Russell (2024). While future work will be required to investigate the relative

contribution of each hypothesis to mediating the evolution of poricidal floral morphology across

diverse clades, we provide a list of key predictions and citations to facilitate the development of

these research directions (Table 3).

333   We also catalogue substantial diversity and bias in how poricidal morphology is achieved

(Fig. 1; Table 2). While floral parts such as anthers, corollas, sepals, and/or staminodes can be

involved, anther modifications are particularly common (~97% of poricidal genera Table 2, S1).

Within taxa, variation in which floral parts are used to achieve poricidal morphology is uncommon.

Additionally, there is substantial morphological diversity in poricidal anther among taxa, especially

with respect to anatomical differences in dehiscence (e.g., via slits, pores, or valves; Harris, 1905;

Buchmann & Hurley, 1978; Buchmann, 1983; Endress, 1996). These patterns strongly suggest

developmental biases, but the developmental pathways of poricidal morphology remain unexplored

to our knowledge. Finally, our study sets the stage for future work to investigate the evolutionary

history of other floral traits that have been repeatedly proposed to be potentially correlated with

poricidal floral morphology, such as loss of nectar, pore size, pollen size and exine ornamentation,

pollen quantity, or dry and loose pollen (Vogel, 1978; Dukas & Dafni, 1990; Cruden, 2000; Roulston

et al., 2000; Freitas & Sazima, 2003; Brito *et al.*, 2016).

346   Despite the apparent adaptive benefits of poricidal morphology, we find significantly lower

origination rates for poricidal genera when considering angiosperms as a whole (31% - 26% lower,

for hidden states A and B respectively), suggesting that this trait is frequently costly. One obvious

potential cost is that poricidal morphology increases pollination specialization, reducing the capacity

for these plant taxa to persist or diversify (Renner, 1989; Dellinger *et al.*, 2019b). Specialization in a

variety of ecological systems (e.g., parasitism, herbivory, pollination) can be associated with reduced

diversification rates (Moran, 1988; Tripp & Manos, 2008; Day *et al.*, 2016; Raia *et al.*, 2016) although

this relationship may not be straightforward (Forister *et al.*, 2012; Armbruster, 2014; Hardy & Otto,

2014; Day *et al.*, 2016; Zenil-Ferguson *et al.*, 2023). As we described in the species-level analyses, we

found that poricidal specialization is not linked to diversification. However, it might be important in

the origination of some genera of angiosperms, as our genus-level state-dependent model shows

fast net diversification for poricidal state B. That this difference in diversification is present at both

the genus- and species-level indicates that evolving poricidal flowers is complex, but once the trait

has evolved and persists (either by more speciation or less extinction), a poricidal flower ancestor

can then establish a new genus (Fig. 4). At the same time, specialization is easy to reverse to a

generalist non-poricidal state (Figure 6B) and poricidal flowers might be costly to reacquire, which

could explain why poricidal states are not linked to recent diversifications for the majority of

angiosperms (Fig 6A). A main reason for the difficulty in reacquiring the poricidal state is that animal

pollination in poricidal plant species is frequently restricted to a subset of possible pollinators. In

fact, many poricidal plant taxa are pollinated by bees capable of floral buzzing or, rarely, by birds or

bees that activate a bellows-like mechanism in the stamen (Sazima *et al.*, 1993; Dellinger *et al.*,

2019b) and is likely often more specialized. Furthermore, the regional diversity of poricidal taxa and

buzzing bees can be strongly positively correlated (Pacheco Filho *et al.*, 2015), suggesting an even

greater degree of poricidal plant-bee specialization (Mesquita-Neto *et al.*, 2018).

The costs of poricidal morphology may also be related to pollen being the only reward

offered by the majority of poricidal species (Vogel, 1978; Buchmann, 1983; Endress, 1994; Vallejo-

Marin *et al.*, 2010). In addition to resulting in specialization on pollen-foraging pollinators, pollen

offered as a food reward might reduce its availability for export to conspecific flowers (Hargreaves *et*

al., 2009). Furthermore, while separating male and female phases in flowers (dichogamy) reduces

selfing and/or interference with cross-pollinating mechanisms, dichogamy in pollen-only rewarding

species also potentially reduces visitation to the rewardless female phase flowers. If the costs of

dichogamy outweigh its benefits for pollen-only rewarding species, this may explain why the

simultaneous presentation of pollen and stigmas (homogamy) in pollen-only flowers is common

(Lloyd & Webb, 1986; Webb & Lloyd, 1986; Renner, 1989).

Notwithstanding the various potential costs of poricidal morphology and its associated

traits, some plant families have higher diversification rates associated with poricidal flowers,

although this is contingent on the state of hidden characters (e.g., Solanaceae, Ericaceae).

Accordingly, other factors that vary across angiosperm groups may be involved in the diversification

of poricidal taxa, such as flower modularity (Dellinger *et al.*, 2019a), abiotic and biotic selective

forces described in Table 3, the relative availability of potential pollinators (Pacheco Filho *et al.*,

2015; Mesquita-Neto *et al.*, 2018), or the presence of other pollen sources in the community (Kemp

et al., 2022).

Poricidal morphology is thought to be an adaptive plateau and appears to constrain floral

evolution (Renner, 1989; Dellinger *et al.*, 2019b). Yet poricidal floral morphology is not necessarily a

dead end. We estimate 145 independent losses of this morphology on average, suggesting this trait

is evolutionarily labile. Multiple non-mutually exclusive hypotheses might explain a return to

loculicidal anther dehiscence. For example, the loss of key pollinators, a reduction in herbivory,

and/or reduced selection by microbial pathogens could all reduce the potential benefits of poricidal

versus non-poricidal morphology (Schwartz-Tzachor *et al.*, 2006; Brito *et al.*, 2017). A detailed

understanding of the biogeography of poricidal taxa and of pollinators and other selective agents

will likely be required to distinguish among these hypotheses. Given differences in how functional

poricidal morphology is achieved, certain poricidal types may also be more evolutionarily labile than

others. For example, the length of the distal slits in poricidal anthers often varies among taxa or even

over the lifetime of a given flower (e.g., in *Actinidia*; Goodwin, 1986), and thus evolutionary

transitions from partial to more and more complete longitudinal dehiscence for some floral forms

may be developmentally relatively simple to achieve.

In conclusion, given the hundreds of thousands of non-poricidal taxa and the tens of

thousands of poricidal taxa, transitions to poricidal morphology are relatively rare, but once evolved,

the probability of losing poricidal morphology is relatively high. Other possibilities for this pattern

include early bursts of speciation associated with poricidal morphology or greater chances for the

poricidal state to go extinct later, given that poricidal genera are proportionally more common early

in angiosperm evolution, but later become rarer relative to non-poricidal genera. Our results thus

suggest that selective forces driving the evolution and maintenance of poricidal morphology may

vary substantially over time and space. Given our coding of genera as poricidal or not, our estimates

for the independent loss and gain of poricidal morphology are also likely highly conservative and

should be interpreted cautiously, although we also recover this pattern of faster poricidal loss in the

species-level analysis with a much larger tree. Despite extensive documentation of poricidal

morphology (Supporting Information Table S1), detailed data across multiple species has only been

collected for some of the major lineages (Brito *et al.*, 2016; Dellinger *et al.*, 2019a; Dellinger *et al.*,

2019b). Furthermore, the pollinators or even pollination systems of poricidal species are frequently

incompletely documented. Additionally, poricidal flowers are associated with diverse pollination

systems, including many wind-pollinated Poaceae, fly and/or non-buzzing bee pollinated Araceae,

Rafflesiaceae (taxa with toothpaste-like pollen extrusion), and *Rhododendron* (taxa with clumped

pollen attached via sticky viscin threads), and bird pollinated *Agarista* and *Axinaea* (taxa with

bellows flowers). Considering such diversity, we believe that future studies investigating functions

beyond only the control of how and when pollen is released (see Table 3) will be invaluable in

understanding the selective pressures driving the gain and loss of poricidal morphology.

Acknowledgements

MVM was partially supported by grants RPG-2018-235 from The Leverhulme Trust and RGP0043/2022 from the Human Frontier Science Program. RZF supported by NSF-DEB 2323170. MVM and RZF greatly benefited from discussions supported by DiveRS-CESAB project AS/CESAB181003. SLB was partially supported by NSF-DEB 1929499. This paper is dedicated to the memories of James A. Harris, Charles D. Michener, and Stefan Vogel for their foundational efforts in elucidating the ecology of poricidal flowers and their bee pollinators.

Author Contributions

SLB, DDJ, RK, and ALR initially conceived and designed the study, with subsequent contributions from MVM on study design and framing. ALR & SLB conducted the literature review, and DDJ, MVM and RK collected taxonomic and phylogenetic data. MVM and RZF performed the diversification analyses and prepared figures. ALR and MVM wrote the first draft of the manuscript; other authors provided editorial advice.

Data Accessibility

The datasets and code supporting this article will be uploaded to Dryad and made publicly available upon publication.

References

Armbruster WS. 2014. Floral specialization and angiosperm diversity: phenotypic divergence, fitness trade-offs and realized pollination accuracy. *Aob Plants* **6**.

Beaulieu JM, O'Meara BC. 2018. Can we build it? Yes we can, but should we use it? Assessing the quality and value of a very large phylogeny of campanulid angiosperms. *American Journal of Botany* **105**(3): 417-432.

Beaulieu JM, O'Meara BC. 2019. Diversity and skepticism are vital for comparative biology: a response to Donoghue and Edwards (2019). *American Journal of Botany* **106**(5): 613-617.

Beaulieu JM, O'Meara BC. 2016. Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. *Systematic Biology* **65**(4): 583-601.

Bennici A. 2003. The convergent evolution in plants. *Rivista di biologia* **96**(3): 485-489.

Brito VL, Rech AR, Ollerton J, Sazima M. 2017. Nectar production, reproductive success and the evolution of generalised pollination within a specialised pollen-rewarding plant family: a case study using Miconia theizans. *Plant Systematics and Evolution* **303**(6): 709-718.

Brito VLG, Fendrich TG, Smidt EC, Varassin IG, Goldenberg R. 2016. Shifts from specialised to
generalised pollination systems in Miconieae (Melastomataceae) and their relation with
anther morphology and seed number. *Plant Biology* **18**(4): 585-593.

Brito VLG, Leite FB, Jorge LR, Sazima M. 2021. Distinct pollen release dynamics between stamens
generate division of labour in pollen flowers of two *Pleroma* species (Melastomataceae).
Flora **285**: 151961.

Buchman SL. 1978. *Vibratile ("Buzz") Pollination in Angiosperms with Poricidally Dehiscent Anthers.*
PhD, University of California at Davis.

Buchmann SL 1983. Buzz pollination in angiosperms. In: Jones CE, Little RJ eds. *Handbook of
Experimental Pollination Biology.* NY: Scientific and Academic Editions, 73-113.

Buchmann SL, Hurley JP. 1978. Biophysical model for buzz pollination in Angiosperms. *Journal of
Theoretical Biology* **72**(4): 639-657.

Burrieza HP, López-Fernández MP, Láinez V, Montenegro T, Maldonado S. 2010. On the nature and
origin of the oxalate package in Solanum sisymbriifolium anthers. *Protoplasma* **247**(1): 45-
56.

Cardinal S, Buchmann SL, Russell AL. 2018. The evolution of floral sonication, a pollen foraging
behavior used by bees (Anthophila). *Evolution* **72**(3): 590-600.

Carrizo García C, Matesevach M, Barboza G. 2008. Features related to anther opening in *Solanum*
species (Solanaceae). *Botanical Journal of the Linnean Society* **158**(2): 344-354.

Christensen SM, Munkres I, Vannette RL. 2021. Nectar bacteria stimulate pollen germination and
bursting to enhance microbial fitness. *Current Biology* **31**(19): 4373-4380. e4376.

Cooley H, Vallejo-Marín M. 2021. Buzz-pollinated crops: A global review and meta-analysis of the
effects of supplemental bee pollination in tomato. *Journal of Economic Entomology.*

Coté GG, Gibernau M. 2012. Distribution of calcium oxalate crystals in floral organs of Araceae in
relation to pollination strategy. *American Journal of Botany* **99**(7): 1231-1242.

Cruden RW. 2000. Pollen grains: why so many? *Plant Systematics and Evolution* **222**: 143-165.

Cullen NP, Fetters AM, Ashman T-L. 2021. Integrating microbes into pollination. *Current opinion in
insect science* **44**: 48-54.

Danforth, B.N., Minckley, R.L, Neff, J.L. 2019. The Solitary Bees: Biology, Evolution, Conservation,
Princeton University Press, Princeton, 472 pp.

Day E, Hua X, Bromham L. 2016. Is specialization an evolutionary dead end? Testing for differences
in speciation, extinction and trait transition rates across diverse phylogenies of specialists
and generalists. *Journal of Evolutionary Biology* **29**(6): 1257-1267.

De Luca PA, Bussiere LF, Souto-Vilaros D, Goulson D, Mason AC, Vallejo-Marin M. 2013. Variability
in bumblebee pollination buzzes affects the quantity of pollen released from flowers.
Oecologia **172**(3): 805-816.

De Luca PA, Cox DA, Vallejo-Marin M. 2014. Comparison of pollination and defensive buzzes in
bumblebees indicates species-specific and context-dependent vibrations.
Naturwissenschaften **101**(4): 331-338.

De Luca PA, Vallejo-Marin M. 2013. What's the 'buzz' about? The ecology and evolutionary
significance of buzz-pollination. *Current Opinion in Plant Biology* **16**(4): 429-435.

Delgado T, Leal LC, El Ottra JHL, Brito VLG, Nogueira A. 2023. Flower size affects bee species
visitation pattern on flowers with poricidal anthers across pollination studies. *Flora* **299**:
152198.

**Dellinger AS, Artuso S, Pamperl S, Michelangeli FA, Penneys DS, Fernández-Fernández DM, Alvear
M, Almeda F, Armbruster WS, Staeder Y. 2019a.** Modularity increases rate of floral
evolution and adaptive success for functionally specialized pollination systems.
Communications biology **2**(1): 1-11.

**Dellinger AS, Chartier M, Fernández-Fernández D, Penneys DS, Alvear M, Almeda F, Michelangeli
FA, Staedler Y, Armbruster WS, Schönenberger J. 2019b.** Beyond buzz-pollination–
departures from an adaptive plateau lead to new pollination syndromes. *New Phytologist*
221(2): 1136-1149.

Dellinger AS, Pöllabauer L, Loreti M, Czurda J, Schönenberger J. 2019c. Testing functional
hypotheses on poricidal anther dehiscence and heteranthery in buzz-pollinated flowers.
Acta ZooBot Austria **156**: 197-214.

Donoghue MJ, Edwards EJ. 2019. Model clades are vital for comparative biology, and ascertainment
bias is not a problem in practice: a response to Beaulieu and O'Meara (2018). *American
Journal of Botany* **106**(3): 327-330.

Dukas R, Dafni A. 1990. Buzz-pollination in three nectariferous Boraginaceae and possible evolution
of buzz-pollinated flowers. *Plant Systematics and Evolution* **169**(1-2): 65-68.

Ellison AM, Gotelli NJ. 2009. Energetics and the evolution of carnivorous plants—Darwin's 'most
wonderful plants in the world'. *Journal of experimental botany* **60**(1): 19-42.

Endress PK. 1994. *Diversity and Evolutionary Biology of Tropical Flowers.* Cambdridge: Cambridge
Universtity Press.

Endress PK 1996. Diversity and evolutionary trends in angiosperm anthers. In: D'Arcy WG, Keating
RC eds. *The Anther: Form, Function and Phylogeny.* Cambridge: Cambridge University Press,
92-110.

Forister M, Dyer LA, Singer M, Stireman III JO, Lill J. 2012. Revisiting the evolution of ecological
specialization, with emphasis on insect–plant interactions. *Ecology* **93**(5): 981-991.

Freitas L, Sazima M. 2003. Floral biology and pollination mechanisms in two viola species-from
nectar to pollen flowers? *Annals of Botany* **91**(3): 311-317.

Freyman WA, Höhna S. 2018. Cladogenetic and anagenetic models of chromosome number
evolution: a Bayesian model averaging approach. *Systematic Biology* **67**(2): 195-215.

Freyman WA, Höhna S. 2019. Stochastic character mapping of state-dependent diversification
reveals the tempo of evolutionary decline in self-compatible Onagraceae lineages.
Systematic Biology **68**(3): 505-519.

Gavrutenko M, Reginato M, Kriebel R, Nicolas AN, Michelangeli FA. 2020. Evolution of floral
morphology and symmetry in the Miconieae (Melastomataceae): multiple generalization
trends within a specialized family. *International Journal of Plant Sciences* **181**(7): 732-747.

Gerenday A, French J. 1988. Endothecial thickenings in anthers of porate monocotyledons.
American Journal of Botany **75**(1): 22-25.

Goodwin R. 1986. Kiwifruit flowers: anther dehiscence and daily collection of pollen by honey bees.
New Zealand journal of experimental agriculture **14**(4): 449-452.

Gottsberger G, Silbauer-Gottsberger I. 1988. Evolution of flower structures and pollination in
Neotropical Cassiinae (Caesalpinaceae) species. *Phyton* **28**: 293-320.

Harder LD, Barclay MR. 1994. The functional significance of poricidal anthers and buzz pollination:
controlled pollen removal from *Dodecatheon*. *Functional Ecology* **8**(4): 509-517.

Hardy NB, Otto SP. 2014. Specialization and generalization in the diversification of phytophagous
insects: tests of the musical chairs and oscillation hypotheses. *Proceedings of the Royal*
Society B: Biological Sciences **281**(1795): 20132960.

Hargreaves AL, Harder LD, Johnson SD. 2009. Consumptive emasculation: the ecological and
evolutionary consequences of pollen theft. *Biological Reviews* **84**(2): 259-276.

Harris JA. 1905. The dehiscence of anthers by apical pores. *Missouri Botanical Garden Annual Report*
1905: 167-257.

Helmstetter AJ, Glemin S, Käfer J, Zenil-Ferguson R, Sauquet H, de Boer H, Dagallier L-PMJ, Mazet
N, Reboud EL, Couvreur TLP, Condamine FL. 2022. Pulled diversification rates, lineages-
through-time plots, and modern macroevolutionary modeling. *Systematic Biology* **71**(3):
758-773.

Helmstetter AJ, Zenil-Ferguson R, Sauquet H, Otto SP, Mendez M, Vallejo-Marin M, Schonenberger
J, Burgarella C, Anderson B, de Boer H, et al. 2023. Trait-dependent diversification in
angiosperms: Patterns, models and data. *Ecology Letters* **26**(4): 640-657.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP, Ronquist F. 2016.
RevBayes: Bayesian phylogenetic inference using graphical models and an interactive model-
specification language. *Systematic Biology* **65**(4): 726-736.

Janssens SB, Couvreur TLP, Mertens A, Dauby G, Dagallier LMJ, Vanden Abeele S, Vandelook F,
Mascarello M, Beeckman H, Sosef M, et al. 2020. A large-scale species level dated
angiosperm phylogeny for evolutionary and ecological analyses. *Biodiversity Data Journal* **8**:
e39677.

Jin Y, Qian H. 2019. V.PhyloMaker: an R package that can generate very large phylogenies for
vascular plants. *Ecography* **42**(8): 1353-1359.

Joly S, Schoen DJ. 2021. Repeated evolution of a reproductive polyphenism in plants is strongly
associated with bilateral flower symmetry. *Current Biology.*

Jordan MI. 2004. Graphical models. *Statistical science* **19**(1): 140-155.

Kemp JE, Telles FJ, Vallejo-Marín M. 2022. Reduced visitation to buzz-pollinated *Cyanella*
hyacinthoides in the presence of other pollen sources in the hyperdiverse Cape Floristic
Region. *Ecology and Evolution* **12**(4): e8784.

Kemp JE, Vallejo-Marin M. 2021. Pollen dispensing schedules in buzz-pollinated plants:
Experimental comparison of species with contrasting floral morphologies. *American Journal*
of Botany **108**(6): 1-13.

Koski MH, Ashman T-L. 2015. Floral pigmentation patterns provide an example of Gloger's rule in
plants. *Nature Plants* **1**(1): 1-5.

Koski MH, Galloway LF. 2018. Geographic variation in pollen color is associated with temperature
stress. *New Phytologist* **218**(1): 370-379.

Larson BMH, Barrett SCH. 1999. The ecology of pollen limitation in buzz-pollinated *Rhexia virginica*
(Melastomataceae). *Journal of Ecology* **87**(3): 371-381.

Lloyd DG, Webb CJ. 1986. The avoidance of interference between the presentation of pollen and
stigmas in Angiosperms. I. Dichogamy. *New Zealand Journal of Botany* **24**(1): 135-162.

Louca S, Pennell MW. 2020. A general and efficient algorithm for the likelihood of diversification
and discrete-trait evolutionary models. *Systematic Biology* **69**(3): 545-556.

Macior LW. 1964. Experimental study of floral ecology of *Dodecatheon meadia*. *American Journal of*
Botany **51**(1): 96-108.

Maddison WP, Midford PE, Otto SP. 2007. Estimating a binary character's effect on speciation and
extinction. *Systematic Biology* **56**(5): 701-710.

Meseguer AS, Carrillo R, Graham SW, Sanmartín I. 2022. Macroevolutionary dynamics in the
transition of angiosperms to aquatic environments. *New Phytologist.*

Mesquita-Neto JN, Blüthgen N, Schlindwein C. 2018. Flowers with poricidal anthers and their

complex interaction networks—Disentangling legitimate pollinators and illegitimate visitors.

Functional Ecology **32**(10): 2321-2332.

Michener CD. 1962. An interesting method of pollen collecting by bees from flowers with tubular

anthers. *Revista de biologia tropical* **10**(2): 167-175.

Moran NA. 1988. The evolution of host-plant alternation in aphids: evidence for specialization as a

dead end. *The American Naturalist* **132**(5): 681-706.

Nelsen MP, Ree RH, Moreau CS. 2018. Ant–plant interactions evolved through increasing

interdependence. *Proceedings of the National Academy of Sciences* **115**(48): 12253-12258.

Pacheco Filho AJdS, Verola CF, Lima Verde LW, Freitas BM. 2015. Bee-flower association in the

Neotropics: implications to bee conservation and plant pollination. *Apidologie* **46**(4): 530-

541.

Paiva ÉAS. 2021. Do calcium oxalate crystals protect against herbivory? *The Science of Nature*

108(3): 1-7.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of phylogenetics and evolution in R language.

Bioinformatics **20**(2): 289-290.

Pennell MW, FitzJohn RG, Cornwell WK. 2016. A simple approach for maximizing the overlap of

phylogenetic and comparative data. *Methods in ecology and evolution* **7**(6): 751-758.

Pereira Nunes CE, Vallejo-Marín M. 2022. How much pollen do beelike floral vibrations remove

from different types of anthers? *International Journal of Plant Sciences* **183**(9): 768-776.

Pozner R, Cocucci A. 2006. Floral structure, anther development, and pollen dispersal of

Halophytum ameghinoi (Halophytaceae). *International Journal of Plant Sciences* **167**(6):

1091-1098.

Proenca CEB. 1992. Buzz pollination: Older and more widespread than we think? *Journal of Tropical*

Ecology **8**: 115-120.

Raia P, Carotenuto F, Mondanaro A, Castiglione S, Passaro F, Saggese F, Melchionna M, Serio C,

Alessio L, Silvestro D. 2016. Progress to extinction: increased specialisation causes the

demise of animal clades. *Scientific Reports* **6**(1): 1-10.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior summarization in Bayesian

phylogenetics using Tracer 1.7. *Systematic Biology* **67**(5): 901-904.

Renner SS. 1989. A survey of reproductive biology in Neotropical Melastomataceae and

Memecylaceae. *Annals of the Missouri Botanical Garden* **76**(2): 496-518.

Roulston TH, Cane JH, Buchmann SL. 2000. What governs protein content of pollen: Pollinator

preferences, pollen-pistil interactions, or phylogeny? *Ecological Monographs* **70**(4): 617-643.

Russell AL, Buchmann SL, Papaj DR. 2017. How a generalist bee achieves high efficiency of pollen
collection on diverse floral resources. *Behavioral Ecology* **28**(4): 991-1003.
Russell AL, Fetters AM, James EI, Ashman T-L. 2021. Pollinator effectiveness is affected by
intraindividual behavioral variation. *Oecologia*: 1-12.
Sauquet H, Magallón S. 2018. Key questions and challenges in angiosperm macroevolution. *New*
Phytologist **219**(4): 1170-1187.
Sazima M, Vogel S, Cocucci A, Hausner G. 1993. The perfume flowers of *Cyphomandra* (Solanaceae).
Pollination by Euglossine bees, bellows mechanism, osmophores, and volatiles. *Plant*
Systematics and Evolution **187**(1-4): 51-88.
Schwartz-Tzachor R, Dafni A, Potts SG, Eisikowitch D. 2006. An ancient pollinator of a contemporary
plant (Cyclamen persicum): When pollination syndromes break down. *Flora-Morphology,*
Distribution, Functional Ecology of Plants **201**(5): 370-373.
The-Plant-List 2013. http://www.theplantlist.org/.
Thorogood CJ, Bauer U, Hiscock SJ. 2018. Convergent and divergent evolution in carnivorous pitcher
plant traps. *New Phytologist* **217**(3): 1035-1041.
Timerman D, Barrett SCH. 2018. Divergent selection on the biomechanical properties of stamens
under wind and insect pollination. *Proceedings of the Royal Society B* **285**(1893): 20182251.
Tribble CM, Freyman WA, Landis MJ, Lim JY, Barido-Sottani J, Kopperud BT, Höhna S, May MR.
2021. RevGadgets: an R Package for visualizing Bayesian phylogenetic analyses from
RevBayes. *Methods in ecology and evolution*.
Tripp EA, Manos PS. 2008. Is floral specialization an evolutionary dead-end? Pollination system
transitions in Ruellia (Acanthaceae). *Evolution: International Journal of Organic Evolution*
62(7): 1712-1737.
Uyeda J, Harmon L. 2020. treeplyr:'dplyr'functionality for matched tree and data objects. *R package*
version 0.1 **9**.
Vallejo-Marin M. 2019a. Buzz pollination: studying bee vibrations on flowers. *New Phytologist*
224(3): 1068-1074.
Vallejo-Marin M. 2019b. Evolutionary tinkering allows buzz pollinated plants to escape from an
adaptive dead-end. *New Phytologist* **221**(2): 618-620.
Vallejo-Marin M. 2022. How and why do bees buzz? Implications for buzz pollination. *Journal of*
experimental botany **73**(4): 1080-1092.
Vallejo-Marin M, Da Silva EM, Sargent RD, Barrett SC. 2010. Trait correlates and functional
significance of heteranthery in flowering plants. *New Phytologist* **188**(2): 418-425.

Vallejo-Marin M, Russell AL. 2024. Harvesting pollen with vibrations: Towards an integrative
understanding of the proximate and ultimate reasons for buzz pollination. *Annals of Botany*:
mcad189.

Vogel S 1978. Evolutionary shifts from reward to deception in pollen flowers. In: Richards AJ ed. *The*
Pollination of Flowers by Insects. London: Academic Press, 89-96.

Webb CJ, Lloyd DG. 1986. The avoidance of interference between the presentation of pollen and
stigmas in angiosperms. II. Herkogamy. *New Zealand Journal of Botany* **24**(1): 163-178.

Zenil-Ferguson R, McEntee JP, Burleigh JG, Duckworth RA. 2023. Linking ecological specialization to
its macroevolutionary consequences: An example with passerine nest type. *Systematic*
Biology **72**(2): 294-306.

Zenil-Ferguson R, Burleigh JG, Freyman WA, Igić B, Mayrose I, Goldberg EE. 2019. Interaction
among ploidy, breeding system and lineage diversification. *New Phytologist* **224**(3): 1252-
1265.

Zhang C, Yang Y-P, Duan Y-W. 2014. Pollen sensitivity to ultraviolet-B (UV-B) suggests floral
structure evolution in alpine plants. *Scientific Reports* **4**(1): 1-4.

Tables and figures

Table 1. Incidence of poricidal flowers at the species, genus, and family levels, as estimated from three separate angiosperm-wide databases. Notice that the proportions of poricidal flowers at each taxonomic level is very similar, irrespective of the database analyzed. Poricidal = number of taxa in which at least one species or genus is reported as poricidal (genera treated as monomorphic). Non-poricidal = Number of taxa with no reported poricidal species or genera.

Database	Taxonomic level	Poricidal	Non-poricidal	Total	% Poricidal
The Plant List (2013)	Family	87	309	396	21.96%
	Genus	635	12010	12645	5.02%
	Species	28218	243187	271405	10.39%
Nelsen et al. (2018)	Family	86	323	409	21.02%
	Genus	521	10264	10785	4.83%
	Species	--	--	--	--
Jian and Qin (2019)	Family	87	330	417	20.86%
	Genus	525	9838	10363	5.06%
	Species	8149	66382	74531	10.93%

681   **Table 2.** Type of floral structure involved in forming a tube-like structure with a small opening

682   containing pollen grains (poricidal flower) exhibited by 639 genera. Two genera (*Tyleria* and

683   *Testulea*; Ochnaceae) contain species forming poricidal flowers via pores and staminodes and thus

684   are included more than once in the table. Data from Supporting Information Table S1.

Floral structure forming a tube	Number of genera (%)	Number of families	Example taxa
Anthers	**633 (99.6%)**	86	*Solanum* (Solanaceae) *Rhexia* (Melastomataceae)
Staminodes	**4 (0.62%)**	1	*Adrenarake, Sauvagesia, Testulea, Tyleria* (Ochnaceae)
Corolla	**3 (0.47%)**	2	*Pedicularis, Rhinanthus* (Orobanchaceae) *Ternstroemia* (Pentaphylacaceae)
Sepals	**1**	1	*Dalechampia* (Euphorbiaceae)

687 **Table 3.** Hypotheses for the evolution of poricidal flowers discussed in the text (see also Vallejo-

688 Marin & Russell, 2024). The second column presents predictions arising from these hypotheses. The

689 different hypotheses are not mutually exclusive.

Hypotheses for the evolution of poricidal flowers	Predictions	References
Escalating response to pollen consumption by greedy pollinators	Poricidal species lose less pollen to pollinator consumption	No direct support, but see: (Dukas & Dafni, 1990; Harder & Barclay, 1994; Hargreaves *et al.*, 2009; Russell *et al.*, 2017; Russell *et al.*, 2021)
Filtering mechanism selecting subset of effective pollinators	Fewer, but more effective pollinators able to access pollen of poricidal species	(Pacheco Filho *et al.*, 2015; Mesquita-Neto *et al.*, 2018; Kemp & Vallejo-Marin, 2021)
Pollen dispensing mechanism	Pollen of poricidal species dispensed more gradually and/or more precisely to pollinators and on particular pollinator body parts	(Larson & Barrett, 1999; De Luca *et al.*, 2013; Dellinger *et al.*, 2019c; Brito *et al.*, 2021; Kemp & Vallejo-Marin, 2021; Pereira Nunes & Vallejo-Marín, 2022)

Protection of pollen from environmental stress (UV, rain, desiccation, etc.)	Pollen of poricidal species germinates more often; adaptations of pollen to environmental stress are reduced in poricidal species	No direct support, but see: (Buchmann, 1983; Gottsberger & Silbauer-Gottsberger, 1988; Zhang *et al.*, 2014; Koski & Ashman, 2015; Koski & Galloway, 2018)
Protection of pollen from herbivory	Poricidal morphology reduces pollen consumption	No direct support, but see: (Burrieza *et al.*, 2010; Coté & Gibernau, 2012; Paiva, 2021)
Protection from microbial antagonists	Adaptations of pollen to microbial antagonists are reduced in poricidal species	No direct support, but see: (Christensen *et al.*, 2021; Cullen *et al.*, 2021)

Figure 1. Diversity of poricidal flower structures illustrated across 11 taxa. The images depict stamens or whole flowers of taxa with poricidal flowers. For illustration of the diverse ways to build a poricidal flower, false colors indicate the structure or tissue involved: Corolla, Filament and accessory tissue (Filament), and Anther. The phylogeny at the center indicates relationships among taxa. Species as follows: 1 = *Chimaphila umbellata* (Ericaceae), 2 = *Demosthenesia cordifolia* (Ericaceae), 3 = *Solanum elaeagnifolium* (Solanaceae), 4 = *Pedicularis groenlandica* (Orobanchaceae), 5 = *Berberis sp.* (Berberidaceae), 6 = *Tibouchina urvilleana* (Melastomataceae), 7 = *Apeiba tibourbou* (Malvaceae), 8 = *Sloanea terniflora* (Elaeocarpaceae), 9 = *Sauvagesia capillaris* (Ochnaceae), 10 = *Senna* sp. (Fabaceae), 11 = *Arctostaphylos* sp. (Ericaceae). Redrawn from original 3D models by William Singleton and Daniel Hornung.

708   **Figure 2**. Proportion of taxa with poricidal flowers at the family, genus, and species level, estimated
709   from The Plant List (2013) database.

Figure 3. Ancestral character state reconstruction from the hidden state-dependent diversification
model (HiSSE) on a genus-wide 10,328 tip phylogeny across angiosperms (see Methods). Non-
poricidal states in hidden state B (light grey) diversify equally or more quickly than poricidal states in
hidden state B (yellow). However, poricidal states in B (yellow) have diversified faster than other
poricidal states and non-poricidal states in hidden state A (red and blue respectively). Even when
non-poricidal states (light gray) represent the majority of the diversification in angiosperms, some
yellow "bursts" of poricidal state originate genera in large families like Solanaceae, Ericaceae,
Fabaceae, and Melastomataceae.

Figure 4. Posterior probability density of estimated diversification rates (origination − extinction) of poricidal vs. non-poricidal taxa (genus) in a hidden state dependent speciation and extinction model with two hidden states (A, B) in an angiosperm-wide genus-level phylogeny. States A and B represent other potential factors unmeasured by this study, but which affect diversification (Beaulieu and O'Meara, 2016, Zenil-Ferguson et al. 2019, Zenil-Ferguson et al. 2023). The states capture different paces of diversification, during which lineages can be poricidal or non-poricidal.

Figure 5. Proportion (top panel) and logged (bottom panel) estimated number of nodes over time

(millions of years before present) of poricidal vs. non-poricidal taxa (genus) in a hidden state

dependent speciation and extinction model with two hidden states (A, B) in an angiosperm-wide

phylogeny.

Figure 6. Hidden state dependent speciation and extinction model fitted to non-poricidal and

poricidal states in a species-level phylogeny with 29,246 tips. (A) No differences in net diversification

were found between poricidal and non-poricidal states in both hidden state A and B. (B) Transitions

to poricidal states from non-poricidal are slower (NP   P) than transitions from poricidal back to

non-poricidal (P   NP) for both hidden states A and B.