Widespread evolution of poricidal flowers: A striking example of

morphological convergence across flowering plants

Avery Leigh Russell[1][†][*], Rosana Zenil-Ferguson[2][*], Stephen L. Buchmann[3], Diana D. Jolles[4],

Ricardo Kriebel[5], Mario Vallejo-Marín[6]

[1] Department of Biology, Missouri State University, Springfield, MO, United States

[2] Biology Department, University of Kentucky, Lexington, KY, United States

[3] Department of Ecology and Evolutionary Biology and Department of Entomology, University of

Arizona, Tucson, AZ, United States

[4] Department of Biological Sciences, Plymouth State University, Plymouth, NH, United States

[5] Department of Botany, Institute of Biodiversity Science and Sustainability, California Academy of

Sciences, San Francisco, CA, United States

[6] Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

*co-first authors

†Corresponding author

Department of Biology, 910 S John Q Hammons Pkwy, Temple Hall, Missouri State University

Springfield, MO 65897, USA

607 220 4432

†AveryRussell@MissouriState.edu

Author Contributions

SLB, DDJ, RK, and ALR initially conceived and designed the study, with subsequent contributions from MVM on study design and framing. ALR & SLB conducted the literature review, and DDJ, MVM and RK collected taxonomic and phylogenetic data. MVM and RZF performed the diversification analyses and prepared figures. ALR and MVM wrote the first draft of the manuscript; other authors provided editorial advice.

Funding

MVM was partially supported by grants RPG-2018-235 from The Leverhulme Trust and RGP0043/2022 from the Human Frontier Science Program. RZF supported by NSF-DEB 2323170. MVM and RZF greatly benefited from discussions supported by DiveRS-CESAB project AS/CESAB181003. SLB was partially supported by NSF-DEB 1929499.

Conflicts of Interest

No conflicts of interest are declared.

Data Accessibility

The datasets and code supporting this article will be uploaded to Dryad and made publicly available upon publication.

Acknowledgements

This paper is dedicated to the memories of James A. Harris, Charles D. Michener, and Stefan Vogel for their foundational efforts in elucidating the ecology of poricidal flowers and their bee pollinators.

Abstract

The evolution of tube-like floral morphologies that control pollen release via small openings (functionally poricidal flowers) represents a taxonomically and geographically widespread instance of repeated and independent evolution of a functionally similar morphology. Poricidal flowers are also often closely associated with buzz pollination by bees. Yet we lack an updated angiosperm-wide survey of their phylogenetic distribution. We identify all known angiosperm genera containing poricidal flowers via a literature survey. We determined their phylogenetic distribution and minimum number of independent gains and losses via a species-level angiosperm-wide phylogeny. We estimated if evolution of poricidal flowers is associated with changes in speciation/extinction via diversification rate analyses. Poricidal flowers occur across 87 angiosperm families and 639 genera containing >28,000 species. At the species level, an average of 205 independent gains and 215 losses of poricidal flowers occurred. Angiosperm-wide analyses suggest an early burst in poricidal evolution, but no differences in net diversification (origination-extinction) between non-poricidal and poricidal taxa. Analyses for two focal families however indicate strong context-dependent effects of poricidal flowers on diversification. Poricidal evolution thus represents a large-scale example of convergent evolution in floral form, but effects on diversification appear to be strongly contingent on phylogenetic and ecological background.

Keywords: Bees, buzz pollination, convergent evolution, diversification analysis, pollen, poricidal anthers, poricidal flowers.

Introduction

Patterns of convergent evolution are of fundamental interest in evolutionary biology. Broad convergence on functionally equivalent morphology across disparate groups is particularly common in plants, as in the case of succulent morphology or carnivorous pitcher plant traps, and has helped elucidate shared selective pressures, mechanisms, and developmental biases (Bennici, 2003; Ellison & Gotelli, 2009; Thorogood *et al.*, 2018). A very geographically and taxonomically widespread pattern of convergent evolution is tubular floral morphology that restricts pollen dispersal via small openings, i.e., functionally poricidal floral morphology (hereafter 'poricidal flowers'; Fig. 1) (Vallejo-Marin & Russell, 2024). Previous studies estimate that approximately 22,000 plant species possess poricidal anthers ('poricidal species'), spread across 72 families and 544 genera (Buchmann, 1983). Additionally, flower traits that control pollen dispersal are key to angiosperm reproductive success (Hargreaves *et al.*, 2009; Minnaar *et al.* 2019) and many poricidal plant species are also agriculturally important, such as tomatoes, cranberries, blueberries, and kiwifruit (Cooley & Vallejo-Marín, 2021). Yet the phylogenetic distribution and evolutionary patterns of poricidal flowers has not been formally characterized and the latest published taxonomic survey of poricidal flowers is now over 40 years old (Buchman, 1978; Buchmann, 1983).

Poricidal flowers share a common functional morphology in which pollen is held within tubular floral structures with small openings (e.g., pores, valves, or slits; hereafter 'pores') through which the pollen is dispersed. In most cases, the tubular structure is composed of modified enlarged anthers, but may alternatively consist of modifications to the corolla or even the entire flower (Fig. 1) (Harris, 1905; Michener, 1962; Buchmann, 1983; De Luca & Vallejo-Marin, 2013; Russell *et al.*, 2017; Dellinger *et al.*, 2019b; Vallejo-Marin, 2019a). Poricidal flowers have been studied for over a century (Harris, 1905) and are

frequently associated with a suite of floral characteristics (Buchmann, 1983; Renner, 1989;

De Luca & Vallejo-Marin, 2013). Most poricidal plant species are animal pollinated,

nectarless, and have dry, loose pollen (reduced pollenkitt lipids) and many are associated

with enantiostyly (style handedness) and heteranthery (multiple kinds of stamens)

(Buchmann, 1983; Vallejo-Marin *et al.*, 2010). Furthermore, poricidal flowers are often

closely associated with the buzz pollination interaction (although not all poricidal flowers are

buzz pollinated; see Vallejo-Marin and Russell (2024) for some exceptions), in which

pollination is nearly exclusively performed by bee species capable of rapidly contracting their

indirect flight muscles while on the flower, generating strong vibrations that expel pollen

onto the bee's body (Macior, 1964; Buchmann, 1983; Proenca, 1992; Vallejo-Marin, 2019a;

Vallejo-Marin & Russell, 2024). Indeed, poricidal flowers are thought to be a key pollen

resource for these buzzing bees, particularly in tropical regions (Delgado *et al.*, 2023).

Poricidal floral morphologies may facilitate speciation by filtering out less effective

pollinators (Cardinal *et al.*, 2018). Yet because of specialization on a subset of pollinators,

poricidal morphology may make these angiosperm taxa more vulnerable to extinction

(Renner, 1989; Pacheco Filho *et al.*, 2015; Dellinger *et al.*, 2019a; Vallejo-Marin, 2019b).

Consequently, poricidal floral morphologies are consistently suggested to be adaptations to

pollinators and thus they might be key traits affecting patterns of speciation and extinction.

Previous attempts at comprehensively studying poricidal flowers were restricted to

species with pored anthers presumed to be buzz pollinated (Harris, 1905; Buchmann, 1983).

However, functionally poricidal floral morphologies are more diverse (Fig. 1). Additionally,

clades of poricidal species can comprise thousands of species (e.g., *Solanum*,

Melastomataceae) and the rate of gain and loss of poricidal floral morphologies has yet to be

quantified. Although poricidal species are extremely widespread, clades are often all

poricidal or not and thus studies of small phylogenetic clades suffer from the limitations of

rarely seeing evolutionary transitions to and from poricidal flowers. Consequently, large-scale analyses are needed to allow generalization and provide sufficient statistical replication to test the effect of poricidal flower morphologies on diversification (Beaulieu & O'Meara, 2018; Zenil Ferguson *et al.*, 2019; Joly & Schoen, 2021). However, large-scale phylogenetic analyses bring their own challenges (Helmstetter *et al.*, 2022). First, it is challenging to obtain information on multiple traits for thousands of species (i.e., presence or absence of poricidal flowers). Second, well resolved mega-phylogenies are required. Fortunately, recent angiosperm-wide phylogenies have become available (Sauquet & Magallón, 2018; Jin & Qian, 2019; Janssens *et al.*, 2020). Third, running complex diversification models in large phylogenies is computationally demanding and proven methods continue to be developed to speed up large-scale and complex analyses. Finally, the effects of a single trait in a large tree can be confounded with heterogeneity in the process of diversification by other unmeasured factors (Beaulieu & O'Meara, 2018; Beaulieu & O'Meara, 2019; Donoghue & Edwards, 2019).

Here, we investigate the evolutionary history of poricidal flowers across angiosperms to address the following questions. **1)** What proportion of angiosperms possesses poricidal flowers? **2)** How many times have poricidal flowers been gained and lost over the evolutionary history of flowering plants? **3)** Is the presence of poricidal flowers associated with changes in net diversification (speciation minus extinction) and diversification rates across all angiosperms? And **4)** Is the presence of poricidal flowers associated with diversification rates (speciation-extinction) within some key families? We therefore conducted an extensive literature survey to identify all presently known angiosperm genera containing species with poricidal flowers. We then used a species-level angiosperm-wide phylogeny to determine the phylogenetic distribution of poricidal flowers, the geologic ages of poricidal lineages, and the number of independent gains and losses of functionally

poricidal floral morphology. We also conducted diversification rate analyses to estimate if the

evolution of poricidal flowers is associated with changes in speciation/extinction. Finally, we

present an overview of both classic and novel hypotheses for why poricidal flowers evolve

and suggest directions for future research about the extent to which convergence in poricidal

flower morphology is accompanied by convergence in function.

Figure 1. Diversity of poricidal flower structures illustrated across 11 taxa. The images

depict stamens or whole flowers of taxa with poricidal flowers. For illustration of the diverse

ways to build a poricidal flower, false colors indicate the structure or tissue involved: Corolla,

Filament and accessory tissue (Filament), and Anther. The phylogeny at the center indicates

relationships among taxa. Species as follows: 1 = *Chimaphila umbellata* (Ericaceae), 2 =

Demosthenesia cordifolia (Ericaceae), 3 = *Solanum elaeagnifolium* (Solanaceae), 4 =

Pedicularis groenlandica (Orobanchaceae), 5 = *Berberis sp.* (Berberidaceae), 6 = *Pleroma*

urvilleanum (Melastomataceae), 7 = *Apeiba tibourbou* (Malvaceae), 8 = *Sloanea terniflora*

(Elaeocarpaceae), 9 = *Sauvagesia capillaris* (Ochnaceae), 10 = *Senna* sp. (Fabaceae), 11 =

Arctostaphylos sp. (Ericaceae). Redrawn from original 3D models by William Singleton and

Daniel Hornung.

Material and Methods

Literature survey

We defined "poricidal flowers" as flowers with tube-like floral structures (typically formed

by individual anthers or groups of anthers, but sometimes by staminodes, sepals or corolla),

restricting pollen release via an opening that can be described as a pore, small slit, or valve.

We conducted our literature search between 2015 and November 2020 for English, German,

and Portuguese language articles describing all presently known genera of functionally

poricidal plant species. Sources included online books, online search engines (Web of

Science, Google Scholar), dissertations, and primary research papers. Search terms included

"anther", "stamen", or "flower" and "apical pore", "dehisce", "dehiscence", "dehiscent",

"introrse", "longitudinal", "porandrous", "porate", "pore", "pored", "poricidal", "porose",

"porous", "short slit", "thecal pores", "valve", or "valvular". The search was performed at the

genus level because numerous well-characterized genera are monomorphic for the

presence/absence of poricidal morphology (e.g., *Solanum*, *Senna*, *Miconia*) and

comprehensive species-level information for rarer and/or tropical genera is frequently sparse

(although we also recognize that monomorphism within genera should not be taken for

granted and there are exceptions in some genera, e.g., Renner, 1989; Gavrutenko *et al.*,

2020). For each genus, we documented the flower parts involved in forming the tubular

structures that comprise poricidal flower morphology, as well as how pollen is released from

poricidal anthers (Fig. S2, Table 1; Supplementary Material Table S1). We also confirmed

non-poricidal morphology for monomorphic genera when morphological information was

available, but generally did not catalog the accompanying literature.

Table 1. Incidence of poricidal flowers at the species, genus, and family levels, as estimated

from an angiosperm-wide database and within the Janssens et al. (2020) phylogeny used for

the angiosperm-wide analysis. The proportions of poricidal flowers at each taxonomic level

are similar. Poricidal = number of taxa in which at least one species or genus is reported as

poricidal (genera treated as monomorphic). Non-poricidal = Number of taxa with no reported

poricidal species or genera.

Source	Taxonomic level	Poricidal	Non-poricidal	Total	% Poricidal
The Plant List (2013)	Family	87	309	396	21.96%
	Genus	635	12,010	12,645	5.02%
	Species	28,218	243,187	271,405	10.39%
Janssens et al. (2020)	Family	79	376	455	17.36%
	Genus	423	6,849	7,272	5.82%
	Species	2,607	26,639	29,246	8.91%

Angiosperm phylogeny and estimates of species number per genus

To estimate the incidence of poricidal and non-poricidal species across the angiosperms we

cross-referenced our 639 poricidal genera against The Plant List (2013), an angiosperm-wide

database, which contained 271,405 species, 12,645 genera, and 396 families of angiosperms,

after filtering out infraspecific classifications (i.e., subspecies and varieties). We then

matched the Janssens et al. (2020) phylogenetic tree to our species-level data using the R

package *treeplyr* (Uyeda and Harmon 2020) and found direct matches for 28,980 species.

The Janssens et al. (2020) phylogenetic tree was specifically developed to address species-

level macroevolutionary analyses like those performed in the present study. The Janssens et

al. (2020) phylogenetic tree is based on matK (incl. trnK) and rbcL. plastids and has 36,101

taxa representing 8,399 genera and was dated using 56 angiosperm fossils. Additionally,

there were 266 species in the Janssens et al. (2020) tree (each representing a different genus)

that were absent in our species-level data, but were of genera represented in our species-level

data. To maximize information for the phylogeny, we assigned the trait value of each genus

in our species-level data to these 266 species in the Janssens et al. (2020) tree, via the R

package *phyndr* (Pennell *et al.*, 2016). This resulted in an ultrametric and bifurcating

phylogenetic tree with a total sample size of 29,246 taxa (tips) with poricidal binary states

that we used for the phylogenetic comparative analyses below.

Phylogenetic comparative methods

State dependent diversification and origination analyses

The 29,246 taxa in the resulting time-calibrated bifurcating tree have tips coded as binary

states ('non-poricidal' or 'poricidal') as the input of a hidden state dependent speciation and

extinction model (HiSSE, Beaulieu and O'Meara 2016), with the goal of testing whether

differences in diversification between poricidal and non-poricidal states exist. For every

HiSSE model we identify the differences in net diversification (speciation minus extinction)

between poricidal and non-poricidal states, but also the diversification differences given by

hidden states A and B. The difference between hidden states is of particular interest,

especially in large phylogenies, because it represents the potential for diversification due to

unmeasured factors or heterogeneity, but not the trait of interest (Beaulieu & O'Meara, 2016).

Thus, estimating the differences between the main and hidden states disentangles the effect of

poricidal state on diversification from other potential sources of diversification. We only

fitted HiSSE and not a simpler model without hidden states (BiSSE, Maddison *et al.*, 2007)

for two main reasons. First, in large phylogenetic trees, many unobserved traits likely

contribute to diversification, and thus our null hypothesis should always have heterogeneity

in diversification rates due to these unobserved traits (instead of the trait of interest having all

the signal). Second, in a Bayesian framework the posterior distribution of the HiSSE model

can be re-written to represent the null hypothesis determined by the character independent

model with two hidden states (CID-2), which describes a model of diversification where the

rates change only due to hidden states and not the trait of interest, as described below.

The HiSSE model was fitted under a Bayesian framework within the RevBayes

environment (Höhna *et al.*, 2016). We obtained the posterior probability densities for every

parameter in the model via a Markov chain Monte Carlo (MCMC) algorithm. We checked for

convergence of our MCMC algorithm by running two chains and observing that the posterior

distributions reached the same range of values, and that stationarity was achieved (see

Supplementary Material for details). Convergence via Gelman and Rubin's convergence

diagnostic (Brooks and Gelman 1998) was verified using the software TRACER (Rambaut *et

al.*, 2018) (Fig. S4-S6). We assumed the sampling fraction for the HiSSE model was 10%,

given the current estimate of ~300,000 extant flowering plant species (The Angiosperm

Phylogeny Group 2016). We do not add a sampling correction for states, because our tree

reflects our literature survey at the species, genus, and family level (Table 1).

To examine whether there might be family specific patterns in how poricidal state

might influence diversification rates, a second round of HiSSE models were performed for

four large angiosperm families matched to the Smith and Brown (2018) phylogenetic tree:

Solanaceae, Melastomataceae, Ericaceae, and Fabaceae. These families were selected

because they have well-resolved trees, are large, and are variable for poricidal state. For each

of the four families we pruned the species-level phylogenies from the Smith and Brown

(2018) tree and matched them with poricidal status. The matching trees and states on the tips

were the inputs for HiSSE. All these analyses were accomplished via RevBayes (Höhna *et*

al., 2016) and convergence was checked similarly to our angiosperm-wide phylogenetic tree.

We reached an ESS of more than 200 for all parameters.

Testing state-dependent diversification in a Bayesian framework

The character independent models (Beaulieu and O'Meara, 2016) are at the core of testing if

the trait of interest is linked to diversification. In the approach used by the R package *hisse*

(Beaulieu *et al.*, 2023) multiple models of state-dependent diversification are fitted to data

and then a model selection procedure using the Akaike Information Criterion (AIC)

determines if a model that includes diversification rates linked to the main states is preferable

over a model with diversification rate differences given only by the hidden states. When

fitting state-dependent diversification models in a Bayesian framework, as done in this study,

the model selection part of the process is unnecessary since the samples of the posterior

distributions for the parameters of HiSSE can be transformed to test the hypotheses laid out

by a CID-2 model as we describe here. The null hypothesis in CID-2 assumes that the two

main states (non-poricidal and poricidal) are equal within each hidden state (A or B), but a

given main state is different between hidden states. In statistical terms, the null hypothesis is:

$$H_0: \quad r_{Non-Poricidal\ A} = r_{Poricidal\ A}$$

$$r_{Non-Poricidal\ B} = r_{PoricidalB}$$

$$r_{Non-Poricidal\ A} \neq r_{Non-Poricidal\ B}$$

$$r_{Poricidal\ A} \neq r_{Poricidal\ B}$$

Where r_i refers to the net diversification (the difference between speciation and extinction, in

mathematical terms, $\lambda_i - \mu_i$) in state i. Using our species-level HiSSE results (Fig. 2b H_0),

we created two test statistics to determine the probability of the differences between non-

poricidal and poricidal states within each hidden state. These test statistics represent the

differences of net diversification between non-poricidal and poricidal states within each

hidden state. Mathematically, these can be written as:

$\quad T_A = r_{Non-Poricidal\ A} - r_{Poricidal\ A}$ and $T_B = r_{Non-Poricidal\ B} - r_{Poricidal\ B}$

and their posterior distributions are calculated by taking the differences in the speciation and

extinction sampling estimates in each iteration of MCMC (Gelman *et al.*, 1995).

For hypothesis testing, we calculate the quantile that the value of 0 represents ($T_A = 0$): the

probability of the absolute value of T_A being greater than zero (in mathematical notation

$P(|T_A| > 0)$). If $P(|T_A| > 0) > 0.05$, non-poricidal and poricidal states in hidden states A are

the same, similarly for $T_B = 0$. This simple test statistic transformation negates the need for

model selection, while allowing us to test for the same hypothesis delineated by CID-2.

Models without diversification. Estimation of number of poricidal transitions

and their timing

As discussed in the Results section, we found that poricidal status does not inform the process

of diversification. For this reason, we fitted a simpler model: a Markov model with two states

(poricidal or not poricidal) and hidden states (A and B), excluding diversification rates. We

estimated the transition rates of this model using the R package *corhmm* (Beaulieu *et al.*,

2022). Under this model, we estimated 1000 stochastic maps using R package *phytools*

(Revell, 2012) to extract the distribution of the expected number of transitions between

poricidal and non-poricidal states in both hidden categories, and to characterize the

distribution of waiting times where the transitions between states happened. The distribution

of the number of transitions allows us to assess which transitions are more common, and the

distribution of waiting times shows whether these transitions are more common at specific

times in the evolution of angiosperms. We calculate the relative lineage through time plots

per state by calculating the lineages through time possessing a given state, and at a given time

interval, by dividing them by the total lineages. This relative LTT allows us to compare

differences in the distributions of lineages at a given time point that are difficult to observe in

a 29K-tip phylogenetic tree (e.g., two lineages at 199 mya vs. 29K lineages at 0 mya) (Fig.

5).

Results

296   Distribution of poricidal flowers across taxonomic levels

297   In a seminal paper, Buchmann (1983) estimated that 15,000-20,000 species of angiosperms in

298   72 families and 544 genera have anthers that dehisce through apical pores or slits (poricidal

anthers). Here, we found that poricidal flowers, including species with poricidal anthers, are

reported from 87 families and 639 genera (Fig. S2, Table 1; Table S1). Using the 271,405

species of angiosperms in the The Plant List database, which includes 635 of 639 genera

identified in our study, yields an upper estimate of 28,218 poricidal species (21.96% of plant

families, 5.02% of genera, and 10.39% of species) (Table 1). Our results therefore suggest

that poricidal flower morphologies have evolved in a fifth of families and up to one in ten

angiosperm species.

Different types of poricidal flowers across taxonomic levels

Flower parts forming the tubular structures that comprise poricidal flower morphology

include poricidal anthers (an estimated 97% of poricidal species), corollas, or other

functionally poricidal floral parts (i.e., staminodes or sepals) (Table 2). Additionally, how

pollen is released from poricidal anthers is particularly diverse, although such classifications

are morphological, rather than developmental, and may therefore be quite inaccurate.

Morphological variation includes pores (60 families and 441 genera), short terminal slits (46

families and 125 genera), valves (15 families and 74 genera), and introrse dehiscence forming

terminal pores or short slits (16 families and 53 genera) (Supplementary Material Table S1).

A given poricidal flower morphology is rarely restricted to a particular family or genus and

even multiple kinds of poricidal anthers are known within a genus (Endress, 1996).

Table 2. Type of floral structure involved in forming a tube-like structure with a small

opening containing pollen grains (poricidal flower) exhibited by 639 genera. Two genera

(*Tyleria* and *Testulea*; Ochnaceae) contain species forming poricidal flowers via pores and

staminodes and thus are included more than once in the table. Data from Supplementary

Material Table S1.

Floral structure	Number of	Number of	Example taxa

forming a tube	genera (%)	families	
Anthers	**633 (99.6%)**	86	*Solanum* (Solanaceae) *Rhexia* (Melastomataceae)
Staminodes	**4 (0.62%)**	1	*Adrenarake, Sauvagesia, Testulea, Tyleria* (Ochnaceae)
Corolla	**3 (0.47%)**	2	*Pedicularis, Rhinanthus* (Orobanchaceae) *Ternstroemia* (Pentaphylacaceae)
Sepals	**1**	1	*Dalechampia* (Euphorbiaceae)

Diversification rates and transitions between poricidal and non-poricidal flowers

For the angiosperm-wide species-level HiSSE analyses using the (Janssens et al. 2020) phylogenetic tree with our data, we found that there were no differences in net diversification (speciation minus extinction rates) between non-poricidal and poricidal states (Fig. 2). Instead, we only found differences between hidden state net diversification rates with probability ($P(|T_A| > 0) = 0$ and $P(|T_B| > 0) = 0$) (Fig. 2B). This result is consistent with the synthesis on state-dependent diversification presented in Helmstetter et al. (2023), where discrete traits modeled under SSE approaches in trees with large numbers of tips typically have no correlation with diversification rates. In the species-level Solanaceae HiSSE analysis, poricidal flowers diversified at a higher rate compared to non-poricidal flowers, but again, this relationship varied with the state of the hidden character. In the HiSSE analysis of Ericaceae, we found a higher diversification rate of poricidal flowers in combination with hidden character state A, but lower when combined with character state B, indicating a strong context-dependence effect of poricidal morphologies (Fig. S2). We lack results for the two other families analyzed at the species level (Melastomataceae and Fabaceae), because models

did not converge after running the analyses for several months, due to high uncertainty on the

branch lengths (additional discussion in Supplementary Material).

Figure 2. A) Posterior distribution of net diversifications under the hidden state dependent

speciation and extinction model (HiSSE) for non-poricidal and poricidal states. No

differences in diversification are found between main states, but only in hidden states. B) Test

statistics   and   that calculate the difference between poricidal and non-poricidal states in

each hidden state show that it is highly probable that there are no differences in

diversification due to poricidality.

Our diversification results for the angiosperm-wide species-level HiSSE analysis

indicate that the evolution of poricidality is faster for certain lineages (in hidden state B), but

not for all angiosperms. Furthermore, we found that transitions from non-poricidal to

poricidal states are on average 33x and 6.6x slower than from poricidal back to non-poricidal

in A and B hidden states, respectively (95% credible intervals: non-poricidal to poricidal:

353   ), and   ); poricidal back to non-

poricidal:), and   )), (Fig. 5B).

This pattern of asymmetry in transition rates from a "specialized" poricidal state back to a

"generalized" non-poricidal state was also found in recent state-dependent diversification

analyses (Day *et al.*, 2016; Zenil-Ferguson *et al.*, 2023), where more evidence of evolving out

of ecological specialization has been found.

Figure 3. Posterior distribution of rates of transition between non-poricidal and poricidal

states under hidden states A and B using the hidden state dependent speciation and extinction

model (HiSSE). For both hidden states it is easier to transition out of the poricidal state than

into the poricidal state (   ).

Discrete trait evolution recovers asymmetry of transitions outside of poricidal

state

After fitting a discrete state model with hidden states, we also recover asymmetry in

maximum likelihood estimates ($q_{01}^{A} = 4.68x10^{-3}$, $q_{01}^{B} = 2.34x10^{-6}$, $q_{10}^{A} = 1.28x10^{-2}$, and

$q_{10}^{B} = 2.19x10^{-4}$). The software would not permit recovering likelihood or confidence

intervals for these maximum likelihood estimates and thus we cannot test if the estimates are

significantly different. However, it is worth noting that point estimates differ by at least 100x.

Using the 1000 stochastic maps simulated from the discrete and hidden states Markov model

(Mkn), we calculated the distribution of the expected number of changes between states. In

Table 3 we summarize the statistics of these transitions, and found the same pattern, with

fewer transitions into poricidal states than transitions out of poricidal states.

Type of transition	Mean	Median	Credible Intervals (95%)
Non-Poricidal $_A$ → Poricidal $_A$	202.22	202	(184, 220)
Poricidal $_A$ → Non-Poricidal $_A$	214.79	215	(189, 239)
Non-Poricidal $_B$ → Poricidal $_B$	3.75	4	(1, 7)
Poricidal $_B$ → Non-Poricidal $_B$	22.85	23	(15, 31)

Table 3. Summary statistics of transitions obtained from the 1000 stochastic maps. These

statistics were obtained by following each lineage and counting the number of transitions

from one state to the next.

Timing of poricidality: A fifty million year wait and establishment

Under the discrete and hidden states Markov model (Mkn) we simulated 1000 stochastic

maps, but we present just one (Fig. 4). In this figure, we observed that poricidal states

(Poricidal$_A$ (red) and Poricidal$_B$ (yellow)) evolved around 150 million years ago (~ 50 million

years after the origin of angiosperms; Fig. 5) and remain in that state until the present

(especially in state Poricidal$_B$, Fig. 4). This observation is reinforced by the relative lineage

through time plots (Fig. 5c and 5f), where state Poricidal$_A$ suddenly increases in relative

numbers, whereas Non-Poricidal$_A$ decreases. Later, state Non-Poricidal$_A$ mostly transitions

into Non-Poricidal$_B$, and Poricidal$_A$ into Poricidal$_B$ and grows in numbers without switching

further states, as seen in the single stochastic map (Fig. 4).

392
Figure 4. A single representative stochastic map calculated via an angiosperm-wide

phylogenetic tree with 29,246 taxa under a discrete-state and hidden state model using R

packages *corhmm* (Beaulieu et al. 2022) and *phytools* for visualization (Revell 2012). Non-

poricidal states in blue (0A) and grey (0B). Poricidal states in red (1A) and yellow (1B). Even

when non-poricidal states (light gray) represent the majority of the diversification in

angiosperms, some yellow "bursts" of poricidal state happened, such as within large families

like Solanaceae, Ericaceae, Fabaceae, and Melastomataceae. Digital art of illustrative

poricidal flowers by Moth Castagna.

Figure 5. Relative lineage through time (RLTT) plots from the 1000 stochastic maps. The relative lineage through time plots are calculated by counting the number of lineages through time in each state and dividing them by the total number of lineages. A) The RLTT for Non-Poricidal$_A$ state indicates a decline of lineages in this state starting around 180 million years ago, with this state mostly transitioning into B) Non-Poricidal$_B$ state and D) Poricidal$_A$. F) Poricidal$_A$ state transitions later into Poricidal$_B$. C) and F) have the total in each main state shown independently from the hidden state.

Discussion

Our study shows that the repeated evolution of functionally poricidal floral morphology represents a major case of convergent evolution in flower form across angiosperms. Although

we found no overall effect on speciation-extinction rates, effects of evolving poricidal

flowers on species diversification likely strongly depend on the phylogenetic, and probably

ecological, context in which they occur, as suggested by our focal family-level analyses. Our

survey updates previous estimates of the incidence of taxa with functionally poricidal floral

morphology, showing that about 10% of angiosperm species have this morphology. We can

also provide a quantitative estimate of its repeated evolution, with an average minimum of

205 independent origins across the angiosperms (Table 3). Although poricidal flowers have

repeatedly independently evolved, which selective pressures drive their evolution are

currently unknown. Given the diversity of angiosperm clades and ecosystems in which

poricidal flowers have arisen (Buchmann 1983; Russell *et al.*, 2024), and the multiple non-

mutually exclusive hypotheses explaining their overall evolution (Vallejo-Marin & Russell,

2024), comparative studies across multiple independent origins of poricidal flowers are

needed to identify the role of different selective pressures on the evolution of this convergent

floral morphology.

When considering angiosperms as a whole, we find significantly lower origination

rates for poricidal taxa (33x - 6.6x lower), suggesting that this trait is frequently costly. One

obvious potential cost is that poricidal morphology increases pollination specialization,

reducing the capacity for these plant taxa to persist or diversify (Renner, 1989; Dellinger *et*

al., 2019b). Specialization in a variety of ecological systems (e.g., parasitism, herbivory,

pollination) can be associated with reduced diversification rates (Moran, 1988; Tripp &

Manos, 2008; Day *et al.*, 2016; Raia *et al.*, 2016) although this relationship may not be

straightforward (Forister *et al.*, 2012; Armbruster, 2014; Hardy & Otto, 2014; Day *et al.*,

2016; Zenil-Ferguson *et al.*, 2023). At the same time, specialization is easy to reverse to a

generalist non-poricidal state (Fig. 3) and poricidal flowers might be costly to reacquire,

which could explain why poricidal states are not linked to recent diversifications for most

angiosperms (Fig. 5D). A main reason for the difficulty in reacquiring the poricidal state is

that animal pollination in poricidal plant species is frequently restricted to a subset of possible

pollinators. In fact, many poricidal plant taxa are pollinated by bees capable of floral buzzing

or, rarely, by birds or bees that activate a bellows-like mechanism in the stamen (Sazima *et*

al., 1993; Dellinger *et al.*, 2019b) and is likely often more specialized. Furthermore, the

regional diversity of poricidal taxa and buzzing bees can be strongly positively correlated

(Pacheco Filho *et al.*, 2015), suggesting an even greater degree of poricidal plant-bee

specialization (Mesquita⬚Neto *et al.*, 2018).

The costs of poricidal morphology may also be related to pollen being the only reward

offered by the majority of poricidal species (Vogel, 1978; Buchmann, 1983; Endress, 1994;

Vallejo-Marin *et al.*, 2010). In addition to resulting in specialization on pollen-foraging

pollinators, pollen offered as a food reward might reduce its availability for export to

conspecific flowers (Hargreaves *et al.*, 2009). Furthermore, while separating male and female

phases in flowers (dichogamy) reduces selfing and/or interference with cross-pollinating

mechanisms, dichogamy in pollen-only rewarding species also potentially reduces visitation

to the rewardless female phase flowers. If the costs of dichogamy outweigh its benefits for

pollen-only rewarding species, this may explain why the simultaneous presentation of pollen

and stigmas (homogamy) in pollen-only flowers is common (Lloyd & Webb, 1986; Webb &

Lloyd, 1986; Renner, 1989).

Notwithstanding the various potential costs of poricidal morphology and its associated

traits, some plant families have higher diversification rates associated with poricidal flowers,

although this is contingent on the state of hidden characters (e.g., Solanaceae, Ericaceae).

These hidden characters might for instance include the availability of the right species of

buzz-pollinating bees or optimal environmental conditions for facilitating pollen while being

vibrated (e.g., low aridity and low wind; see Russell *et al.*, 2024; Vallejo-Marin & Russell,

2024). Other factors that have been suggested to modulate the diversification of poricidal taxa

include flower modularity (Dellinger *et al.*, 2019a), the relative availability of potential

pollinators (Pacheco Filho *et al.*, 2015; Mesquita⬜Neto *et al.*, 2018), or the presence of other

pollen sources in the community (Kemp *et al.*, 2022).

Poricidal morphology is thought to be an adaptive plateau and appears to constrain

floral evolution (Renner, 1989; Dellinger *et al.*, 2019b). Yet poricidal floral morphology is

not necessarily a dead end. We estimate 215 independent losses of this morphology on

average, suggesting this trait is evolutionarily labile. Multiple non-mutually exclusive

hypotheses might explain a return to loculicidal anther dehiscence. For example, the loss of

key pollinators, a reduction in herbivory, and/or reduced selection by microbial pathogens

could all reduce the potential benefits of poricidal versus non-poricidal morphology

(Schwartz-Tzachor *et al.*, 2006; Brito *et al.*, 2017). A detailed understanding of the

biogeography of poricidal taxa and of pollinators and other selective agents will likely be

required to distinguish among these hypotheses (see Russell *et al.*, 2024), which the present

dataset will facilitate. Given differences in how functional poricidal morphology is achieved,

certain poricidal types may also be more evolutionarily labile than others. For example, the

length of the distal slits in poricidal anthers often varies among taxa or even over the lifetime

of a given flower (e.g., in *Actinidia*; Goodwin, 1986), and thus evolutionary transitions from

partial to more and more complete longitudinal dehiscence for some floral forms may be

developmentally relatively simple to achieve.

Finally, we catalogue substantial diversity and bias in how poricidal morphology is

achieved (Fig. 1; Table 2). While floral parts such as anthers, corollas, sepals, and/or

staminodes can be involved, anther modifications are particularly common (~99% of

poricidal genera Table 2, S1). Within taxa, variation in which floral parts are used to achieve

poricidal morphology is uncommon. Additionally, there is substantial morphological

diversity in poricidal anther among taxa, especially with respect to anatomical differences in

dehiscence (e.g., via slits, pores, or valves; Harris, 1905; Buchmann & Hurley, 1978;

Buchmann, 1983; Endress, 1996). These patterns strongly suggest developmental biases, but

the developmental pathways of poricidal morphology remain unexplored to our knowledge.

Our study sets the stage for future work to investigate the evolutionary history of floral traits

that repeatedly proposed to be potentially correlated with poricidal floral morphology, such

as loss of nectar, pore size, pollen size and exine ornamentation, pollen quantity, or dry and

loose pollen (Vogel, 1978; Dukas & Dafni, 1990; Cruden, 2000; Roulston *et al.*, 2000;

Freitas & Sazima, 2003; Brito *et al.*, 2016).

Classical hypotheses for the evolution of poricidal morphology have highlighted the

role of pollinators (Vallejo-Marin & Russell, 2024). Most, but not all species with poricidal

flowers are pollinated by floral buzzing bees, which are prolific consumers of pollen

(Michener, 1962; Buchmann, 1983; Hargreaves *et al.*, 2009; Danforth et al. 2019; Vallejo-

Marin, 2022). Poricidal morphology might control the rate of pollen release, and thus reduce

the tempo of pollen removal by pollen-consuming pollinators (e.g., Larson & Barrett, 1999;

De Luca *et al.*, 2014; Brito *et al.*, 2016). Likewise, poricidal morphology may result in more

effective dispersal of pollen to conspecifics by pollinators, by enhancing the precision of

pollen placement on the pollinator body and/or by filtering more generalized and less

effective pollinators (e.g., Pacheco Filho *et al.*, 2015; Mesquita☐Neto *et al.*, 2018; Kemp &

Vallejo-Marin, 2021; Vallejo-Marin & Russell, 2024). However, pollinator-mediated

selection alone appears insufficient to explain the evolution of poricidal morphology across

diverse angiosperm clades. Poricidal species have diverse mating systems, from selfing to

outcrossing, and many poricidal flowers are wind pollinated, not animal pollinated (e.g., in

Poaceae and Halophytaceae; Pozner & Cocucci, 2006; Table S1). Indeed, some poricidal taxa

arose as early as 150 million years ago, potentially ~26 million years before buzz pollinating

bees (Cardinal *et al.*, 2018; Almeida *et al.*, 2023). Although vibration can be a key

mechanism of wind-mediated pollen release (Timerman & Barrett, 2018), and could function

analogously to floral buzzing by bees for these poricidal species, the functional benefits of

poricidal morphology likely go well beyond pollen release, as discussed in Vallejo-Marin and

Russell (2024).

In conclusion, given the hundreds of thousands of non-poricidal taxa and the tens of

thousands of poricidal taxa, transitions to poricidal morphology are relatively rare, but once

evolved, the probability of losing poricidal morphology is relatively high. Other possibilities

for this pattern include early bursts of speciation associated with poricidal morphology or

greater chances for the poricidal state to go extinct later, given that poricidal genera are

proportionally more common early in angiosperm evolution, but later become rarer relative to

non-poricidal genera. Our results thus suggest that selective forces driving the evolution and

maintenance of poricidal morphology may vary substantially over time and space. Given our

coding of genera as poricidal or not, our estimates for the independent loss and gain of

poricidal morphology are also likely highly conservative and should be interpreted

cautiously. Despite extensive documentation of poricidal morphology (Supplementary

Material Table S1), detailed data across multiple species has only been collected for some of

the major lineages (Brito *et al.*, 2016; Dellinger *et al.*, 2019a; Dellinger *et al.*, 2019b).

Furthermore, the pollinators or even pollination systems of poricidal species are frequently

incompletely documented. Additionally, poricidal flowers are associated with diverse

pollination systems, including many wind-pollinated Poaceae, fly and/or non-buzzing bee

pollinated Araceae, Rafflesiaceae (taxa with toothpaste-like pollen extrusion), and

Rhododendron (taxa with clumped pollen attached via sticky viscin threads), and bird

pollinated *Agarista* and *Axinaea* (taxa with bellows flowers). Considering such diversity,

future studies investigating functions beyond only the control of how and when pollen is

released will be invaluable in understanding the selective pressures driving the gain and loss

of poricidal morphology.

Data Accessibility

542 The datasets and code supporting this article will be uploaded to Dryad and made publicly

543 available upon publication.

References

Almeida EAB, Bossert S, Danforth BN, Porto DS, Freitas FV, Davis CC, Murray EA,

Blaimer BB, Spasojevic T, Ströher PR, Orr MC, Packer L, Brady SG,

Kuhlmann M, Branstetter MG, Pie MR. 2023. The evolutionary history of bees in

time and space. *Current Biology* **33**(16): 3409-3422.e6.

Armbruster WS. 2014. Floral specialization and angiosperm diversity: phenotypic

divergence, fitness trade-offs and realized pollination accuracy. *Aob Plants* **6**.

Beaulieu JM, O'Meara BC. 2018. Can we build it? Yes we can, but should we use it?

Assessingthe quality and value of a very large phylogeny ofcampanulid angiosperms.

American Journal of Botany **105**(3): 417-432.

Beaulieu JM, O'Meara BC. 2019. Diversity and skepticism are vital for comparative

biology: a response to Donoghue and Edwards (2019). *American Journal of Botany*

106(5): 613-617.

Beaulieu JM, O'Meara BC. 2016. Detecting hidden diversification shifts in models of trait-

dependent speciation and extinction. *Systematic Biology* **65**(4): 583-601.

Beaulieu J, O'Meara B, Oliver J, Boyko J. 2022. corHMM: Hidden Markov Models of

Character Evolution. *R package version 2.8*.

Beaulieu J, O'Meara B, Caetano D., Boyko J. Vasconcelos T. 2023. hisse: Hidden State

Speciation and Extinction. *R package version 2.1.11*.

Bennici A. 2003. The convergent evolution in plants. *Rivista di biologia* **96**(3): 485-489.

Brito VL, Rech AR, Ollerton J, Sazima M. 2017. Nectar production, reproductive success

and the evolution of generalised pollination within a specialised pollen-rewarding

plant family: a case study using Miconia theizans. *Plant Systematics and Evolution*

303(6): 709-718.

Brito VLG, Fendrich TG, Smidt EC, Varassin IG, Goldenberg R. 2016. Shifts from

specialised to generalised pollination systems in Miconieae (Melastomataceae) and

their relation with anther morphology and seed number. *Plant Biology* **18**(4): 585-

593.

Brito VLG, Leite FB, Jorge LR, Sazima M. 2021. Distinct pollen release dynamics

between stamens generate division of labour in pollen flowers of two *Pleroma* species

(Melastomataceae). *Flora* **285**: 151961.

Brooks SP., Gelman A. 1998. General methods for monitoring convergence of iterative

simulations. *Journal of Computational and Graphical Statistics* **7**(4): 434-455.

Buchman SL. 1978. *Vibratile ("Buzz") Pollination in Angiosperms with Poricidally*

Dehiscent Anthers. PhD, University of California at Davis.

Buchmann SL 1983. Buzz pollination in angiosperms. In: Jones CE, Little RJ eds.

Handbook of Experimental Pollination Biology. NY: Scientific and Academic

Editions, 73-113.

Buchmann SL, Hurley JP. 1978. Biophysical model for buzz pollination in Angiosperms.

Journal of Theoretical Biology **72**(4): 639-657.

Burrieza HP, López-Fernández MP, Láinez V, Montenegro T, Maldonado S. 2010. On

the nature and origin of the oxalate package in Solanum sisymbriifolium anthers.

Protoplasma **247**(1): 45-56.

Cardinal S, Buchmann SL, Russell AL. 2018. The evolution of floral sonication, a pollen

foraging behavior used by bees (Anthophila). *Evolution* **72**(3): 590-600.

Carrizo García C, Matesevach M, Barboza G. 2008. Features related to anther opening in

Solanum species (Solanaceae). *Botanical Journal of the Linnean Society* **158**(2): 344-

354.

Christensen SM, Munkres I, Vannette RL. 2021. Nectar bacteria stimulate pollen

germination and bursting to enhance microbial fitness. *Current Biology* **31**(19): 4373-

4380. e4376.

Cooley H, Vallejo-Marín M. 2021. Buzz-pollinated crops: A global review and meta-

analysis of the effects of supplemental bee pollination in tomato. *Journal of Economic*

Entomology.

Coté GG, Gibernau M. 2012. Distribution of calcium oxalate crystals in floral organs of

Araceae in relation to pollination strategy. *American Journal of Botany* **99**(7): 1231-

1242.

Cruden RW. 2000. Pollen grains: why so many? *Plant Systematics and Evolution* **222**: 143-

165.

Cullen NP, Fetters AM, Ashman T-L. 2021. Integrating microbes into pollination. *Current*

opinion in insect science **44**: 48-54.

Danforth, B.N., Minckley, R.L, Neff, J.L. 2019. The Solitary Bees: Biology, Evolution,

Conservation, Princeton University Press, Princeton, 472 pp.

Day E, Hua X, Bromham L. 2016. Is specialization an evolutionary dead end? Testing for

differences in speciation, extinction and trait transition rates across diverse

phylogenies of specialists and generalists. *Journal of Evolutionary Biology* **29**(6): 1257-1267.

De Luca PA, Bussiere LF, Souto-Vilaros D, Goulson D, Mason AC, Vallejo-Marin M. 2013. Variability in bumblebee pollination buzzes affects the quantity of pollen released from flowers. *Oecologia* **172**(3): 805-816.

De Luca PA, Cox DA, Vallejo-Marin M. 2014. Comparison of pollination and defensive buzzes in bumblebees indicates species-specific and context-dependent vibrations. *Naturwissenschaften* **101**(4): 331-338.

De Luca PA, Vallejo-Marin M. 2013. What's the 'buzz' about? The ecology and evolutionary significance of buzz-pollination. *Current Opinion in Plant Biology* **16**(4): 429-435.

Delgado T, Leal LC, El Ottra JHL, Brito VLG, Nogueira A. 2023. Flower size affects bee species visitation pattern on flowers with poricidal anthers across pollination studies. *Flora* **299**: 152198.

Dellinger AS, Artuso S, Pamperl S, Michelangeli FA, Penneys DS, Fernández-Fernández DM, Alvear M, Almeda F, Armbruster WS, Staeder Y. 2019a. Modularity increases rate of floral evolution and adaptive success for functionally specialized pollination systems. *Communications biology* **2**(1): 1-11.

Dellinger AS, Chartier M, Fernández☐Fernández D, Penneys DS, Alvear M, Almeda F, Michelangeli FA, Staedler Y, Armbruster WS, Schönenberger J. 2019b. Beyond buzz☐pollination–departures from an adaptive plateau lead to new pollination syndromes. *New Phytologist* **221**(2): 1136-1149.

Dellinger AS, Pöllabauer L, Loreti M, Czurda J, Schönenberger J. 2019c. Testing functional hypotheses on poricidal anther dehiscence and heteranthery in buzz-pollinated flowers. *Acta ZooBot Austria* **156**: 197-214.

Donoghue MJ, Edwards EJ. 2019. Model clades are vital for comparative biology, and

ascertainment bias is not a problem in practice: a response to Beaulieu and O'Meara

(2018). *American Journal of Botany* **106**(3): 327-330.

Dukas R, Dafni A. 1990. Buzz-pollination in three nectariferous Boraginaceae and possible

evolution of buzz-pollinated flowers. *Plant Systematics and Evolution* **169**(1-2): 65-

68.

Ellison AM, Gotelli NJ. 2009. Energetics and the evolution of carnivorous plants—Darwin's

'most wonderful plants in the world'. *Journal of experimental botany* **60**(1): 19-42.

Endress PK. 1994. *Diversity and Evolutionary Biology of Tropical Flowers*. Cambdridge:

Cambridge Universtity Press.

Endress PK 1996. Diversity and evolutionary trends in angiosperm anthers. In: D'Arcy WG,

Keating RC eds. *The Anther: Form, Function and Phylogeny*. Cambridge: Cambridge

University Press, 92-110.

Forister M, Dyer LA, Singer M, Stireman III JO, Lill J. 2012. Revisiting the evolution of

ecological specialization, with emphasis on insect–plant interactions. *Ecology* **93**(5):

981-991.

Freitas L, Sazima M. 2003. Floral biology and pollination mechanisms in two viola species-

from nectar to pollen flowers? *Annals of Botany* **91**(3): 311-317.

Freyman WA, Höhna S. 2018. Cladogenetic and anagenetic models of chromosome number

evolution: a Bayesian model averaging approach. *Systematic Biology* **67**(2): 195-215.

Freyman WA, Höhna S. 2019. Stochastic character mapping of state-dependent

diversification reveals the tempo of evolutionary decline in self-compatible

Onagraceae lineages. *Systematic Biology* **68**(3): 505-519.

Gavrutenko M, Reginato M, Kriebel R, Nicolas AN, Michelangeli FA. 2020. Evolution of

floral morphology and symmetry in the Miconieae (Melastomataceae): multiple

generalization trends within a specialized family. *International Journal of Plant*

Sciences **181**(7): 732-747.

Gelman A, Carlin JB, Stern HS, Rubin DB. 1995. Bayesian data analysis. Chapman and

Hall/CRC.

Gerenday A, French J. 1988. Endothecial thickenings in anthers of porate monocotyledons.

American Journal of Botany **75**(1): 22-25.

Goodwin R. 1986. Kiwifruit flowers: anther dehiscence and daily collection of pollen by

honey bees. *New Zealand journal of experimental agriculture* **14**(4): 449-452.

Gottsberger G, Silbauer-Gottsberger I. 1988. Evolution of flower structures and

pollination in Neotropical Cassiinae (Caesalpinaceae) species. *Phyton* **28**: 293-320.

Harder LD, Barclay MR. 1994. The functional significance of poricidal anthers and buzz

pollination: controlled pollen removal from *Dodecatheon*. *Functional Ecology* **8**(4):

509-517.

Hardy NB, Otto SP. 2014. Specialization and generalization in the diversification of

phytophagous insects: tests of the musical chairs and oscillation hypotheses.

Proceedings of the Royal Society B: Biological Sciences **281**(1795): 20132960.

Hargreaves AL, Harder LD, Johnson SD. 2009. Consumptive emasculation: the ecological

and evolutionary consequences of pollen theft. *Biological Reviews* **84**(2): 259-276.

Harris JA. 1905. The dehiscence of anthers by apical pores. *Missouri Botanical Garden*

Annual Report **1905**: 167-257.

Helmstetter AJ, Glemin S, Käfer J, Zenil-Ferguson R, Sauquet H, de Boer H, Dagallier

L-PMJ, Mazet N, Reboud EL, Couvreur TLP, Condamine FL. 2022. Pulled

diversification rates, lineages-through-time plots, and modern macroevolutionary

modeling. *Systematic Biology* **71**(3): 758-773.

Helmstetter AJ, Zenil-Ferguson R, Sauquet H, Otto SP, Mendez M, Vallejo-Marin M,

Schonenberger J, Burgarella C, Anderson B, de Boer H, et al. 2023. Trait-

dependent diversification in angiosperms: Patterns, models and data. *Ecology Letters*

26(4): 640-657.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP,

Ronquist F. 2016. RevBayes: Bayesian phylogenetic inference using graphical

models and an interactive model-specification language. *Systematic Biology* **65**(4):

726-736.

Janssens SB, Couvreur TLP, Mertens A, Dauby G, Dagallier LMJ, Vanden Abeele S,

Vandelook F, Mascarello M, Beeckman H, Sosef M, et al. 2020. A large-scale

species level dated angiosperm phylogeny for evolutionary and ecological analyses.

Biodiversity Data Journal **8**: e39677.

Jin Y, Qian H. 2019. V.PhyloMaker: an R package that can generate very large phylogenies

for vascular plants. *Ecography* **42**(8): 1353-1359.

Joly S, Schoen DJ. 2021. Repeated evolution of a reproductive polyphenism in plants is

strongly associated with bilateral flower symmetry. *Current Biology*.

Jordan MI. 2004. Graphical models. *Statistical science* **19**(1): 140-155.

Kemp JE, Telles FJ, Vallejo-Marín M. 2022. Reduced visitation to buzz-pollinated

Cyanella hyacinthoides in the presence of other pollen sources in the hyperdiverse

Cape Floristic Region. *Ecology and Evolution* **12**(4): e8784.

Kemp JE, Vallejo-Marin M. 2021. Pollen dispensing schedules in buzz-pollinated plants:

Experimental comparison of species with contrasting floral morphologies. *American*

Journal of Botany **108**(6): 1-13.

Koski MH, Ashman T-L. 2015. Floral pigmentation patterns provide an example of Gloger's

rule in plants. *Nature Plants* **1**(1): 1-5.

Koski MH, Galloway LF. 2018. Geographic variation in pollen color is associated with temperature stress. *New Phytologist* **218**(1): 370-379.

Larson BMH, Barrett SCH. 1999. The ecology of pollen limitation in buzz-pollinated *Rhexia virginica* (Melastomataceae). *Journal of Ecology* **87**(3): 371-381.

Lloyd DG, Webb CJ. 1986. The avoidance of interference between the presentation of pollen and stigmas in Angiosperms. I. Dichogamy. *New Zealand Journal of Botany* **24**(1): 135-162.

Louca S, Pennell MW. 2020. A general and efficient algorithm for the likelihood of diversification and discrete-trait evolutionary models. *Systematic Biology* **69**(3): 545-556.

Macior LW. 1964. Experimental study of floral ecology of *Dodecatheon meadia*. *American Journal of Botany* **51**(1): 96-108.

Maddison WP, Midford PE, Otto SP. 2007. Estimating a binary character's effect on speciation and extinction. *Systematic Biology* **56**(5): 701-710.

Meseguer AS, Carrillo R, Graham SW, Sanmartín I. 2022. Macroevolutionary dynamics in the transition of angiosperms to aquatic environments. *New Phytologist*.

Mesquita⬜Neto JN, Blüthgen N, Schlindwein C. 2018. Flowers with poricidal anthers and their complex interaction networks—Disentangling legitimate pollinators and illegitimate visitors. *Functional Ecology* **32**(10): 2321-2332.

Michener CD. 1962. An interesting method of pollen collecting by bees from flowers with tubular anthers. *Revista de biologia tropical* **10**(2): 167-175.

Minnaar C, Anderson B, de Jager ML, Karron JD. 2019. Plant–pollinator interactions along the pathway to paternity. *Annals of Botany* **123**:225-245

Moran NA. 1988. The evolution of host-plant alternation in aphids: evidence for specialization as a dead end. *The American Naturalist* **132**(5): 681-706.

Pacheco Filho AJdS, Verola CF, Lima Verde LW, Freitas BM. 2015. Bee-flower

association in the Neotropics: implications to bee conservation and plant pollination.

Apidologie **46**(4): 530-541.

Paiva ÉAS. 2021. Do calcium oxalate crystals protect against herbivory? *The Science of*

Nature **108**(3): 1-7.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of phylogenetics and evolution in

R language. *Bioinformatics* **20**(2): 289-290.

Pennell MW, FitzJohn RG, Cornwell WK. 2016. A simple approach for maximizing the

overlap of phylogenetic and comparative data. *Methods in ecology and evolution* **7**(6):

751-758.

Pereira Nunes CE, Vallejo-Marín M. 2022. How much pollen do beelike floral vibrations

remove from different types of anthers? *International Journal of Plant Sciences*

183(9): 768-776.

Pozner R, Cocucci A. 2006. Floral structure, anther development, and pollen dispersal of

Halophytum ameghinoi (Halophytaceae). *International Journal of Plant Sciences*

167(6): 1091-1098.

Proenca CEB. 1992. Buzz pollination: Older and more widespread than we think? *Journal of*

Tropical Ecology **8**: 115-120.

Raia P, Carotenuto F, Mondanaro A, Castiglione S, Passaro F, Saggese F, Melchionna

M, Serio C, Alessio L, Silvestro D. 2016. Progress to extinction: increased

specialisation causes the demise of animal clades. *Scientific Reports* **6**(1): 1-10.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior summarization

in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**(5): 901-904.

Renner SS. 1989. A survey of reproductive biology in Neotropical Melastomataceae and

Memecylaceae. *Annals of the Missouri Botanical Garden* **76**(2): 496-518.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution *PeerJ* **12**: e16505.

Roulston TH, Cane JH, Buchmann SL. 2000. What governs protein content of pollen:

Pollinator preferences, pollen-pistil interactions, or phylogeny? *Ecological*

Monographs **70**(4): 617-643.

Russell AL, Buchmann SL, Papaj DR. 2017. How a generalist bee achieves high efficiency

of pollen collection on diverse floral resources. *Behavioral Ecology* **28**(4): 991-1003.

Russell AL, Fetters AM, James EI, Ashman T-L. 2021. Pollinator effectiveness is affected

by intraindividual behavioral variation. *Oecologia*: 1-12.

Russell AL, Buchmann SL, Ascher JS, Wang Z, Kriebel R, Jolles DD, Orr MC, Hughes

AC. 2024. Global patterns and drivers of buzzing bees and poricidal plants. *Current*

Biology **34**(14): 3055-3063.e5.

Sauquet H, Magallón S. 2018. Key questions and challenges in angiosperm macroevolution.

New Phytologist **219**(4): 1170-1187.

Sazima M, Vogel S, Cocucci A, Hausner G. 1993. The perfume flowers of *Cyphomandra*

(Solanaceae). Pollination by Euglossine bees, bellows mechanism, osmophores, and

volatiles. *Plant Systematics and Evolution* **187**(1-4): 51-88.

Schwartz-Tzachor R, Dafni A, Potts SG, Eisikowitch D. 2006. An ancient pollinator of a

contemporary plant (Cyclamen persicum): When pollination syndromes break down.

Flora-Morphology, Distribution, Functional Ecology of Plants **201**(5): 370-373.

Smith SA, Brown JW. 2018. Constructing a broadly inclusive seed plant phylogeny.

American Journal of Botany **105**(3): 302–314.

The-Plant-List 2013. http://www.theplantlist.org/.

Thorogood CJ, Bauer U, Hiscock SJ. 2018. Convergent and divergent evolution in

carnivorous pitcher plant traps. *New Phytologist* **217**(3): 1035-1041.

Timerman D, Barrett SCH. 2018. Divergent selection on the biomechanical properties of

stamens under wind and insect pollination. *Proceedings of the Royal Society B*

285(1893): 20182251.

Tribble CM, Freyman WA, Landis MJ, Lim JY, Barido☐Sottani J, Kopperud BT,

H☐hna S, May MR. 2021. RevGadgets: an R Package for visualizing Bayesian

phylogenetic analyses from RevBayes. *Methods in ecology and evolution.*

Tripp EA, Manos PS. 2008. Is floral specialization an evolutionary dead☐end? Pollination

system transitions in Ruellia (Acanthaceae). *Evolution: International Journal of*

Organic Evolution **62**(7): 1712-1737.

Uyeda J, Harmon L. 2020. treeplyr:'dplyr'functionality for matched tree and data objects. *R*

package version 0.1 **9**.

Vallejo-Marin M. 2019a. Buzz pollination: studying bee vibrations on flowers. *New*

Phytologist **224**(3): 1068-1074.

Vallejo-Marin M. 2019b. Evolutionary tinkering allows buzz pollinated plants to escape

from an adaptive dead-end. *New Phytologist* **221**(2): 618-620.

Vallejo-Marin M. 2022. How and why do bees buzz? Implications for buzz pollination.

Journal of experimental botany **73**(4): 1080-1092.

Vallejo-Marin M, Da Silva EM, Sargent RD, Barrett SC. 2010. Trait correlates and

functional significance of heteranthery in flowering plants. *New Phytologist* **188**(2):

418-425.

Vallejo-Marin M, Russell AL. 2024. Harvesting pollen with vibrations: Towards an

integrative understanding of the proximate and ultimate reasons for buzz pollination.

Annals of Botany: mcad189.

Vogel S 1978. Evolutionary shifts from reward to deception in pollen flowers. In: Richards

AJ ed. *The Pollination of Flowers by Insects*. London: Academic Press, 89-96.

Webb CJ, Lloyd DG. 1986. The avoidance of interference between the presentation of

pollen and stigmas in angiosperms. II. Herkogamy. *New Zealand Journal of Botany*

24(1): 163-178.

Zenil-Ferguson R, McEntee JP, Burleigh JG, Duckworth RA. 2023. Linking ecological

specialization to its macroevolutionary consequences: An example with passerine nest

type. *Systematic Biology* **72**(2): 294-306.

Zenil☐Ferguson R, Burleigh JG, Freyman WA, Igić B, Mayrose I, Goldberg EE. 2019.

Interaction among ploidy, breeding system and lineage diversification. *New*

Phytologist **224**(3): 1252-1265.

Zhang C, Yang Y-P, Duan Y-W. 2014. Pollen sensitivity to ultraviolet-B (UV-B) suggests

floral structure evolution in alpine plants. *Scientific Reports* **4**(1): 1-4.

820