CORTICOTHALAMIC MODELLING OF SLEEP NEUROPHYSIOLOGY WITH APPLICATIONS TO MOBILE EEG

Taha Morshedzadeh[1,2,*], Kevin Kadak[1,2], Sorenza P. Bastiaens[1,2], M. Parsa Oveisi[1,7], Davide Momi[1], Zheng Wang[1], Shreyas Harita[1,2], Maurice Abou Jaude[3], Christopher A. Aimone[3], Steve Mann[5], Sean L. Hill[1,4,6], and John D. Griffiths[1,2,4,7,*]

[1]Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto (*primary affiliation*).

[2]Institute of Medical Science (IMS), University of Toronto

[3]Interaxon Inc., Toronto, ON, Canada

[4]Department of Psychiatry, University of Toronto

[5]Department of Electrical & Computer Engineering, University of Toronto

[6]Department of Physiology, University of Toronto

[7]Institute of Biomedical Engineering, University of Toronto

[*]*Corresponding author* – Email: taha.morshedzadeh@gmail.com; john.griffiths@utoronto.ca;

February 29, 2024

ABSTRACT

Recent developments in mathematical modelling of EEG data enable estimation and tracking of otherwise-inaccessible neurophysiological parameters over the course of a night's sleep. Likewise, advancements in wearable electronics have enabled easier & more affordable at-home collection of sleep EEG data. The convergence of these two advances, namely neurophysiological modelling for mobile sleep EEG, has the potential to significantly improve sleep assessments in research and the clinic. However, this subject area has received limited attention in existing literature. To address this, we used an established mathematical model of the corticothalamic system to analyze EEG power spectra from 5 datasets, spanning from in-lab, research-grade systems to at-home mobile EEG devices. In the present work, we compare the convergent and divergent features of the data and the estimated physiological model parameters. While data quality and characteristics differ considerably, several key patterns consistent with previous theoretical and empirical work are observed. During the transition from lighter to deeper NREM stages, i) the exponent of the aperiodic ($1/f$) spectral component is increased, ii) bottom-up thalamocortical drive is reduced, iii) corticocortical connection strengths are increased. This effect, which we observe in healthy individuals across all 5 datasets, is interestingly absent in individuals taking SSRI antidepressants, suggesting possible effects of ascending neuromodulatory systems on corticothalamic oscillations. Our results provide a proof-of-principle for the utility and feasibility of this physiological modelling-based approach to analyzing data from mobile EEG devices, providing a mechanistic measure of brain physiology during sleep at home or in the lab.

Keywords Neural Field Modelling · Electroencephalography · Power Spectral Density · Corticothalamic System · Excitation & Inhibition

1   Introduction

Sleep neurophysiology and EEG

Sleep is a vital and near-universal physiological function, manifested in most of the animal kingdom in a regular circadian pattern [1, 2]. It is far more than just a state of rest and reduced energy expenditure. States of sleep serve as a period during which the brain undergoes significant changes, including metabolic homeostasis and recovery [3, 4], synaptic regulation, and memory consolidation [1, 5]. Disturbances in sleep rhythms can also increase susceptibility to various types of psychiatric and neurological conditions such as mood disorders, epilepsy, and dementia [6–9]. Despite these associations, sleep disorders remain highly underdiagnosed clinically, or misdiagnosed as other neurological ailments [10].

The physiological state of the brain moves through a complex trajectory of dynamical regimes during a night's sleep. These changes evolve on the timescale of tens of minutes, and their electrical footprints are reflected in (and are indeed defined by) electroencephalography (EEG) recordings. Polysomnography (PSG), is one of the most widely-used methods for evaluating sleep in the clinical setting. It involves the concurrent monitoring of EEG, electromyography (EMG), electrooculography (EOG), movement, and respiration. Sleep stages are defined in terms of the properties of EEG time series data over standard (30s) windows, and the time series of the stages for the successive windows forms the hypnogram. This data is typically evaluated over a single channel, following the –mostly correct– assumption that brain activity changes similarly across all EEG channels during sleep [11]. Each of these stages has characteristic phenomenological definitions defined by the common sleep staging standards [12, 13]. Although it has strong diagnostic and prognostic utility [14, 7, 15–17], classical sleep staging is highly constrained as it is limited to only 5 values (the 5 stages W, N1, N2, N3, REM) to capture the vast continuum of brain states in sleep. This problem is further exacerbated by the highly subjective interpretation of different stages by human scorers, which has led to considerable inter-expert variability [18, 19]. Therefore, it is crucial to augment this information with more detailed quantitative approaches for evaluating brain activity trajectories in sleep.

Power spectral estimation is one of the fundamental methods for studying the characteristics of a time series signal across different frequencies. Studying the EEG power spectral density (PSD) in the same 30-second windows used for sleep staging can provide us with a more high-dimensional evaluation of brain states over these intervals. EEG PSDs can be reliably described in terms of two main components: i) A background $1/f^n$ trend, understood to be non-oscillatory or 'aperiodic', and defined by its exponent and offset, and ii) An oscillatory component which is highly periodic, featuring well-defined attributes such as frequency, amplitude, and bandwidth. The aperiodic component is an intrinsic feature of many natural processes, and is believed in the neuroscientific context to reflect variable excitatory/inhibitory balance [20]. It has also been linked to cognitive decline in ageing [21], cognitive speed [22], and movement [23]. Periodic activity is traditionally examined in the frequency bands delta (0.5-4 Hz), theta (4-7.5 Hz), alpha (7.5-12 Hz), beta (16-30 Hz), and gamma (>30 Hz). During wakefulness, the brain exhibits high-frequency low-amplitude activity, and as the subject transitions to NREM sleep, the activity transitions into a low-frequency high-amplitude

pattern.[24, 25]. The transition from lighter to deeper NREM sleep is associated with increased slope of the EEG power spectrum, along with increases in the amplitude of the delta band and a decrease in the amplitude the alpha band [26]. Wakefulness also is signified by a presence of alpha and gamma peaks, and REM (rapid eye movement) sleep is correlated with increases in gamma and theta but not alpha peaks [27].

The principal brain structure that drives brain state changes during sleep, including our measurement of them with EEG, is the corticothalamic system [28–30]. Different stages of sleep have been linked to changes in corticothalamic activity [29, 31–34] and to changes in the periodic and aperiodic components of the EEG signals over those changes [35, 36, 34, 37]. For instance, the transition from wakefulness to N1 sleep is also characterized by an increase in the slope of the $1/f$ component and the low-frequency band powers [35, 36], which is itself observed to be associated with corticothalamic communication [30].

Sleep stage N3, also known as Slow-wave sleep (SWS), is understood as the deepest stage of NREM sleep, showing strongly synchronized cortical activity in the infra-slow (<1 Hz) and delta (1-4 Hz) frequency bands. This synchronized cortical activity has been shown to be driven locally through corticocortical connections, and with reduced thalamocortical input [31, 38–40]. Interestingly, 1 Hz transcranial magnetic stimulation (TMS) in the cortex can effectively entrain this 1Hz cortical oscillation around the stimulation site [41], indicating that cortical activation is the primary source driving this oscillation. Synaptic homeostasis and long-term potentiation (LTP) have also been found to occur strongly in SWS [5, 33], and brain stimulation at this stage can trigger memory replays and improve memory recall [42].

Mathematical modelling of sleep-wake dynamics

This deep foundation of experimental knowledge in neuroscience across multiple species, spatial scales, and observable phenomena, provides a strong motivation for the development and use of mathematical models that explain sleep EEG in terms of their underlying neurophysiological processes across the units of the corticothalamic circuitry. One of the most widely used and extensively studied models of this kind to date was introduced by Robinson et al. [30], which describes, at the mesoscopic spatial scale, a four-node corticothalamic network containing the thalamic relay, thalamic reticular, cortical excitatory, and cortical inhibitory neural populations. With this structure, the Robinson model has proved highly capable of replicating measured EEG time series and power spectra [30, 43–45], with applications including evoked potentials [46, 47], alpha rhythms [48], and sleep & arousal [49, 50], to name only a few. In a 2015 paper, Abeysuriya et al. demonstrate the use of this model to study the trajectories of physiological brain states expressed through the EEG, across a night of sleep [51]. By fitting the model-generated power spectra to those observed in empirical EEG, circuit mechanisms such as corticocortical, corticothalamic, and intrathalamic connection strengths can be estimated from 30-second windows rolling throughout the night, and their changes compared against separately-scored PSG classifications. In this way, mathematical modelling of corticothalamic system dynamics can be used to enrich the observations made via classical sleep stages and conventional power spectral analysis.

Emerging mobile neurotechnologies for sleep EEG measurements

One of the most significant technical developments in the field of EEG over the past decade has been a suite of hardware, software, and commercial innovations leading to the widespread availability of low-cost ("consumer-grade") wireless mobile EEG devices. The lower price tag, smaller footprint, use of flexible components such as conductive rubber and conductive fabric, and more streamlined setup of these systems hold great promise for scientists and clinicians needing to access larger samples of subjects and over many more nights than is possible with traditional in-lab sleep EEG assessments. Two of the most established mobile sleep EEG headsets on the market today are Muse S by Interaxon [52] and Dreem by Beacon Biosignals [53]. Although, these products face stiff competition from other startups that with smaller but increasing market share, such as Cerebra [54], URGOnight [55] and IDUN, along with major consumer electronics companies such as LG Electronics (sleepwave.ai) that are looking to enter the mobile sleep EEG market.

This approach can enable an easier and more affordable overnight recording of sleep EEG at home or in the research lab. The easier setup and reduced cost can readily enable the researchers to make recordings over more repeated nights and for a larger population.

Present work

The recent advances cited above in our fundamental understanding of sleep neurophysiology, our ability to formulate and model it mathematically, and in the emergence of new technologies promising to radically up-scale the accessibility of EEG-based sleep monitoring, prompt a series of important research questions at the intersection of these topics. This was the focus of the present study. Selecting several widely used open-access research- and consumer-grade sleep EEG datasets, we used power spectral analysis to first evaluate the changes to periodic and aperiodic spectrum components across sleep stages during the night of sleep, assessing the performance of different EEG systems in capturing variations in physiological brain states. We then fit the Robinson corticothalamic model to these EEG power spectra, with a view to studying mechanisms underlying these physiological states over sleep stages, and evaluating their correlations to the depth of NREM sleep. Lastly, we used health data from one of the analyzed cohorts to investigate the correlations of model-estimated neurophysiological parameters with specific mental & physical health biomarkers.

2 Methods

2.1 EEG Datasets

We used EEG data from multiple sources, described in the following. All datasets were acquired according to the ethics board regulation at the hosting institutions. They were accessed and used in accordance with their relevant licences and data-sharing agreements. The left frontal, central, or temporoparietal channels were used in each dataset, specifically F3 or adjacent 10/20 system locations, subject to availability and data quality.

2.1.1   Sleep European Data Format - Extended (Sleep-EDFX)

We used 197 recordings from 185 subjects (97 female / 78 male, mean age 54.7), which were recorded in the time span of 1987-1991 and 1994 using portable Walkman-style cassettes at home [56, 57]. The accessed data had been digitized from the analog signal at the sampling rate of 100 Hz. In this dataset, 153 of the subjects had no previous health conditions and 44 were generally healthy but had trouble sleeping. We accessed the dataset through PhysioNet [58], acquiring the version last updated in 2018. We selected the data from the Fpz-Cz electrode channel for this work. The sleep stages were originally marked according to the Rechtschaffen & Kales (R&K) method [13], and were transformed into the AASM standard for further use in this project. In this paper, we will refer to this dataset as EDF-X for brevity.

2.1.2   Dreem Open Datasets (DOD)

This dataset includes 80 PSGs collected using a research-grade PSG setup from 80 subjects (54 male / 26 female, mean age 42.39). The dataset was curated by Dreem, a manufacturer of sleep EEG headsets, to benchmark automatic sleep staging methods [59, 60]. The data was sampled at 250 Hz and scored by sleep professionals based on the 2007 AASM manual. This dataset is comprised of two sections: 25 healthy subjects recorded in Bretigny-Sur-Orge, France, over 12 mastoid-referenced EEG channels; and 55 at Redwood City, CA, USA, from subjects with Obstructive Sleep Apnea (OSA), with 8 mastoid-referenced EEG channels. We used the Fp2-O2 channel data from this dataset for the current project.

2.1.3   Nap-EEG Dataset

This dataset was acquired from a healthy cohort of 22 individuals (16 male / 6 female; mean age 25.5 ± 7.03). The data was recorded between 10:00 and 17:00 over 1 to 2 consecutive days at the City College of New York, totalling 41 recordings. Subjects took a high- or low-load cognitive task and then took a nap for 30 minutes, where their EEG would be recorded over 64 channels of EEG and 2 channels of EOG, sampled at 1.5 KHz [61]. The data includes sleep stages and 2528 sleep spindles manually annotated, according to the AASM system. The dataset was accessed through the Open Science Framework (OSF) [62]. The PO8 channel data was chosen for the purpose of this project.

2.1.4   Wisconsin Sleep Cohort (WSC) Dataset

The WSC dataset is recorded from a large cohort of state employees in Wisconsin, United States. We requested and accessed the standardized dataset through the National Sleep Research Resource (NSRR) [63, 64]. This set includes 2570 recordings from 1123 subjects. This is a longitudinal data set in which the same subjects came to the sleep lab every 4 to 5 years for a PSG recording. Each subject has 1 to 5 EEG recordings through the years, each approximately 4.5 years apart on average (mean 4.54 ± 1.50 years) between 2000 and 2015. The percentage of subjects with 1, 2, 3, 4, and 5 recordings was respectively 32.5%, 17.1%, 39.6%, 10.6%, and 0.2%. The subjects were 37 to 85 years old (mean age 59.82 ± 8.49, 1385 male / 1185 female). The 6 recording EEG electrodes were referenced to the ipsilateral mastoid electrodes and sampled at 100 Hz for the data from 2000 - 2009, and at 200 Hz from 2009 to 2015. This dataset also

includes a large variety of mental and physical health information, such as the Zung Depression Scale, anxiety scales, caffeine consumption, number of recent nights with insomnia, blood pressure disorders, current medications, etc. We used the 556 first recordings in this dataset from 248 subjects, 39 to 81 years of age (mean age 60.29 ± 8.62, 137 male / 111 female). The C3 channel data was chosen for the purpose of this project.

2.1.5   Muse Sleep Dataset (MSD)

Muse S is a wireless sleep EEG headband manufactured by Interaxon (Toronto, ON, Canada). The data is sampled at 256 Hz, recorded from dry electrodes TP9, TP10, Fp1 and Fp2, and referenced at the FpZ electrode. We used 10 recordings made between 2020 and 2022, provided by Interaxon, that were selected from Muse's Sleep Dataset (MSD). MSD is an internal dataset of overnight at-home sleep recordings collected with the Muse S EEG headset [52]. This dataset was collected in accordance with the privacy policy that users agree to when using the Muse headband and ensures their informed consent concerning the use of EEG data for scientific research purposes. The subjects were 26 to 68 years old (mean age 38.70 ± 13.07, 7 male / 3 female) at the time of recordings. Sleep stages for these recordings were produced by Muse's proprietary automated sleep staging algorithm [52]. Due to the increased presence of artifacts in the data recorded from dry EEG electrodes [65], for each subject, we marked the 30s epochs with a standard deviation larger than that of the whole recording, thus dropping an approximate 7.48% of all epochs from power spectral analysis & model fitting across the entire dataset. In this work, we use the EEG data in the TP7 channel from the MSD dataset, as it uses a frontal reference, thus quantifying the differential trace between the temporoparietal and frontal electrodes.

2.2   EEG & Hypnogram Data Analysis

2.2.1   Pre-processing

The data were organized and processed using MNE-Python [66]. The power spectral density from the data was calculated in 30s segments using Welch's method [67] in 4s Hamming windows with a 1s overlap. Choice of window lengths in EEG signal processing should be optimized for the analysis objectives in question [68]. In the present case, this choice of window sizes was made to balance the sharpness of the peak frequencies –due to noise-driven changes in the power of those bands– with a physiologically-plausible level of specificity in key rhythms such as alpha (7.5-12 Hz). And the segment length here was chosen as it is the segment length over which the sleep stages are labelled. The sleep staging system used here is the 2007 standard, issued by the American Academy of Sleep Medicine (AASM) [12]. The epochs with stages marked as *unknown* were omitted for all datasets.

2.2.2   Aperiodic component estimation using FOOOF

We used the Python library `FOOOF v1.0.0` to separate the periodic and aperiodic components of the empirical and fitted power spectra [69]. The algorithm fit a Gaussian power spectrum corresponding to the aperiodic component to each EEG PSD without a knee, in the range of 0-45 Hz, with bins of the size 0.25 Hz. The Gaussian spectrum was then deducted from the EEG power spectrum to separate the periodic (oscillatory) components. This process was

Table 1: Abbreviations and descriptions of sleep the hypnogram-based sleep metrics calculated using YASA

Metric	Description	Unit
TIB	Total duration of the hypnogram.	min
SPT	Total duration from first to last period of sleep.	min
WASO	Duration of wake periods between the first and last periods of sleep	min
TST	Total duration of N1 + N2 + N3 + REM sleep between the first and last periods of sleep.	min
SE	(TST / TIB) × 100	%
SME	(TST / SPT) × 100	%
SOL	Latency to first epoch of sleep	min
<Stage> Latency	Latency to the first instance of the sleep stage *<Stage>*	min

implemented iteratively and optimized to get maximum 4 oscillatory peaks, each between 1 to 4 Hz in bandwidth, and with at least 1 V^2 / Hz amplitude. We used the extracted exponent (slope) & offset of the fitted aperiodic ($1/f$-like) component and the frequency & power of the periodic components to study the phenomenological properties of the power spectra.

2.2.3 Calculating sleep metrics and statistics using YASA

We used the Python library YASA v0.6.3 to estimate some common sleep metrics based on the presented hypnograms [70]. This library calculated values for sleep architecture and quality metrics using the subjects' hypnograms according to the AASM guidelines [12]. Table 1 includes these evaluated metrics.

2.3 Neurophysiological Model of Thalamocortical Activity

This work used a *neural field model* of thalamocortical dynamics to simulate plausible activity observed in the EEG data [30, 45]. In this physiological wave equation model, we model activity across these units of the thalamocortical circuitry, generated by: cortical excitatory (pyramidal) neurons (e), cortical inhibitory interneurons (i), thalamic reticular nucleus (r) and thalamic relay nuclei (s).

In each of these populations, a mean firing rate (i.e., pulse density) of each population denoted as Q_a is calculated based on the mean somatic voltage (V_a). Henceforth in this document, a & b will represent any of the modelled populations, $a, b \in \{e, i, r, s\}$.

With θ_a as the mean firing threshold and σ'_a as the standard deviation of the somatic voltage for the population a, we can calculate Q_a as:

$$Q_a = S(V_a) = \frac{Q_a^{max}}{1 + e^{-\frac{(V_a - \theta_a)}{\sigma'_a}}}. \tag{1}$$

Using Q, the number of outgoing axonal spikes from the population (ϕ_a) can be determined via this equation:

$$D_a \phi_a = Q_a \tag{2}$$

Figure 1: ***Method for fitting the sleep EEG power spectra to the model.*** **A)** Schematic of the neurophysiological model of the thalamocortical system, which simulates each channel of EEG as an independent active unit spanning the thalamic and cortical components. **B)** With this in mind, we accessed 5 datasets (EDF-X, Nap-EEG, DOD, WSC, MSD) which included sleep EEG and hypnograms. **C)** The *empirical* power spectra over 30-second epochs calculated using Welch's method and the *fitted* power spectra generated using Braintrak to yield the fitted parameters. **D)** Time series of the fitted gain (G) parameters for subject 11, recording no. 1, from the Nap-EEG dataset. Each fitted epoch yields a set of 5 gains. The gain parameters are then used to calculate the circuit gain parameters x, y, z. **E)** Distribution of the parameters x, y, z across the entire Nap-EEG dataset is shown. Different sleep stages are denoted by the colour of the dots in the scatter plot. Different stages are clustered in different areas of the subspace. The dashed line in the 2D x, y plot marks the stability boundary at $x + y = 1$.

$$D_a = \frac{1}{\gamma_a^2}\frac{\partial^2}{\partial t^2} + \frac{2}{\gamma_a}\frac{\partial}{\partial t} + 1 - r_a^2\nabla^2, \tag{3}$$

where D_a is a nonlinear term that dampens the incoming spike rate into a *field* equation (ϕ), and the temporal damping rate is $\gamma_a = v_a/r_a$. For each population a, v_a is the axonal conduction velocity, which is approximated to 10 m/s for myelinated axons that form the thalamocortical projections. r_a is the total range of axons of type a. This model assumes that long-range connections are myelinated and hence have a higher v_a. Shorter-range connections are not myelinated and will only have negligible values, of v, rendering their effects on cortical activity insignificant. Among the thalamocortical connections, only the thalamic relay-excitatory cortical $(r \leftrightarrow e)$ connection has a non-negligible distance, and as a result, we can take r_e as the only significant r value affecting the propagation and assume the other r values to be 1.

In the subcortical units (r and s), we also assume spatial uniformity such that $\nabla^2 = 0$. Given the large value of γ_a for the thalamocortical connection, the damper term (D_a) converges to 1. As such, we can approximate that in the physiological states, $Q_a = \phi_a$.

To account for the delays introduced by the synapses, we introduce $1/\beta$ and $1/\alpha$ which are respectively the *rise* and *decay* time constants of the postsynaptic soma activation, in the response of a population to a spike. We can write the dendritic impulse response function as:

$$L(u) = \frac{\alpha\beta}{\beta - \alpha}(e^{-\alpha u} - e^{-\beta u}) \tag{4}$$

So, if $\alpha \neq \beta$, the dendritic activation function can be written as:

$$D_{\alpha\beta} = \frac{1}{\alpha\beta}\frac{d^2}{dt^2} + \left(\frac{1}{\alpha} + \frac{1}{\beta}\right)\frac{d}{dt} + 1 \tag{5}$$

The Fourier transform of $L(u)$ yields a function in which the dendritic impulse response acts as a low-pass filter with the cut-off frequency at α, and exhibiting a more steep attenuation at β Hz:

$$L(\omega) = (1 - \frac{i\omega}{\alpha})^{-1}(1 - \frac{i\omega}{\beta})^{-1}, \tag{6}$$

where $\omega = 2\pi f$ is the angular frequency and f is the frequency in Hz.

We take $V_a(r, t)$ to be the electrical field (voltage) in population a, influenced by: 1) ϕ_{ab} which is the incoming activation by the presynaptic populations from population b, 2) N_{ab} the mean number of synapses between a and b, and 3) s_{ab} the strength of each synapse between these two populations (the time-integrated response strength in the soma for each incoming spike):

$$V(r, t) = \sum_b V_{ab}(r, t) \tag{7}$$

$$D_\alpha(t)V_{ab}(r, t) = N_{ab}s_{ab}\phi_b(r, t - \tau_{ab}) \tag{8}$$

We further define $\nu_{ab} = N_{ab}s_{ab}$ as the strength of all incoming synapses. The value of s (and hence ν) are considered positive for excitatory neurons and negative for inhibitory neurons. In this work, we assume *random connectivity* in the excitatory and inhibitory populations in the cortex, which means that $N_{ia} = N_{ea}$ for any population a. Therefore, we simplify the ν values as follows: $\nu_{ee} = \nu_{ie}$, $\nu_{ei} = \nu_{ii}$, and $\nu_{es} = \nu_{is}$. Hence, we are left with the 8 independent ν values: $\nu_{ee}, \nu_{ei}, \nu_{es}, \nu_{se}, \nu_{sr}, \nu_{rs}, \nu_{re}$, and ν_{sn}.

By taking the Fourier transform of Eqn. (8), we can represent the cortical excitatory field (ϕ_e) in terms of the external sensory input field (ϕ_n) in the Fourier domain:

$$\frac{\phi_e(k,\omega)}{\phi_n(k,\omega)} = \frac{G_{es}G_{sn}L^2e^{i\omega t_0/2}}{(1 - G_{srs}L^2)(1 - G_{ei}L)(k^2r_e^2 + q^2r_e^2)} \tag{9}$$

$$q^2(\omega)r_e^2 = \left(1 - \frac{i\omega}{\gamma_e}\right)^2 - \frac{G_{ee}L(\omega) + G_{es}L(\omega)S}{1 - G_{ei}L(\omega)}, \tag{10}$$

$$\tag{11}$$

$$S = \frac{(LG_{se} + LG_{sr}LG_{re})e^{i\omega t_0/2}}{1 - LG_{sr}LG_{rs}}, \tag{12}$$

where $k = 2\pi/\lambda$ is the wave vector with wavelength λ.

In a steady state, can assume V_a to be the only the perturbations to the function and take a linearized approximation of Eqn. (1), around the first term of the Taylor expansion. We define the parameter ρ as the derivative of the first term in this expansion. Hence we can reinterpret Eqn. (1) as:

$$Q_a(r,t) = \rho_a V_a(r,t) \tag{13}$$

$$\rho_a = S'(V_a^{(0)}) \tag{14}$$

G_{ab} is defined as the gain value between populations a and b, describing the strength of the response in population a as the result of the unit input of population b, determined by all the scalars that affect the activation of a synaptic population:

$$G_{ab} = \rho_a \nu_{ab} = \rho_a N_{ab} s_{ab} \tag{15}$$

These gain variables are multiplicative, so that $G_{abc} = G_{ab}G_{bc}$. In this manner, the gains in functionally significant loops can be simplified as $G_{ese} = G_{es}G_{se}$ representing the excitatory cortico-thalamo-cortical loop directed by the thalamic relay nuclei, $G_{esre} = G_{es}G_{sr}G_{re}$ representing the inhibitory cortico-thalamo-cortical loop directed by both the thalamic relay and reticular nuclei, and $G_{srs} = G_{sr}G_{rs}$ representing the gain in the inhibitory intrathalamic feedback loop.

In this case, we assume the uniform distribution of the cortical excitatory units (that is, spatially-uniform values of the wave vector k). If we approximate the brain as a finite-sized rectangular sheet with dimensions $lx \times ly$, we can calculate the EEG power spectrum $P(\omega)$ as the integration of $\phi_e(k,\omega)$ over the wave vector k:

$$P_{EEG}(\omega) = P_0 \left| \frac{L(\omega)T/G_{sn}}{1 - G_{ie}L(\omega)} \right|^2 \frac{(2\pi)^2}{l_x l_y} \times \sum_{m,n=-\infty}^{\infty} \frac{e^{-k_{m,n}^2 k_0^2}}{|k_{m,n}^2 r_e^2 + q^2(\omega)r_e^2|^2}, \tag{16}$$

where:

$$T = \frac{LG_{sn}e^{i\omega t_0/2}}{1 - LG_{sr}LG_{rs}}, \tag{17}$$

$$P_0 = \frac{\pi|\phi_n|^2}{r_e^2}G_{es}G_{sn} \tag{18}$$

$$k_{m,n}^2 r_e^2 = (2\pi m r_e/l_x)^2 + (2\pi n r_e/l_y)^2 \tag{19}$$

In Eqn. (16), the term $e^{-k_{m,n}^2/k_0^2}$ represents the low-pass spatial filtering induced by the dispersion of EEG electrical fields through the scalp and the cerebrospinal fluid between the cortex and the EEG sensor. This dispersion will also be spatially uniform given the uniformity of the vector k. The low-pass cutoff k_0 is set at $10\,\mathrm{m}^{-1}$ based on prior empirical observations by Srinivasan et al. [71].

To mitigate the effects of high-frequency EMG artifacts introduced by pericranial, cervical, and extraocular muscles on the higher frequencies in of the power spectra [72, 73], an additional EMG power spectrum is added to that of the EEG [74, 75]:

$$P_{total}(\omega) = P_{EEG}(\omega) + A_{EMG}\left(\frac{\omega/2\pi f_{EMG}}{1 + (\omega/2\pi f_{EMG})}\right)^2, \tag{20}$$

where the A_{EMG} term is fitted together with the other parameters for f_{EMG} in the range of 10 to 50 Hz.

It is worth noting regarding Eqn. (16) that by fitting this power spectrum function, we only get 5 gain (G) values (G_{ee}, G_{ei}, G_{ese}, G_{esre}, G_{srs}) along with α, β, t_0, and the fitted artifact term A_{EMG}. There are infinite solutions to find ν_a with the power spectrum model fitting approach, since $\nu_a = N_a s_a$. Similar to ν, the values of the gain parameters are negative for inhibitory synapses and positive for excitatory synapses, making G_{ei} and G_{sr} negative and all the other gains positive.

2.3.1 *xyz* space

In the stable regions of the 9-dimensional parameter space at low frequencies, a reduced 3-dimensional space could be defined to represent the model parameters, in which: 1) x is the cortical loop gain and represents the corticocortical connection strength, 2) y is the corticothalamic gain and represents how effectively the thalamus can drive cortical activity, and 3) z is the intrathalamic gain. These three parameters are calculated via the following equations:

$$x = \frac{G_{ee}}{(1 - G_{ei})} \tag{21}$$

$$y = \frac{G_{ese} + G_{esre}}{(1 - G_{srs})(1 - G_{ei})} \tag{22}$$

$$z = \frac{-G_{srs}\alpha\beta}{(\alpha + \beta)^2} \tag{23}$$

Each underlying state of brain activity gives more or less unique combinations of xyz. This system can be used to represent thalamocortical activity in many brain states with fewer dimensions than the entire fitted parameter set.

Eqn (22) asserts that the balance between cortical excitatory versus inhibitory activity determines the positive or negative sign of y. Excitation brings y toward more positive values, and inhibition will shift it to negative values. Eqns (21) and (23), respectively, indicate that the values of x and z will always be positive.

2.4 Simulation and model fitting

Simulations and model fitting were performed in MATLAB using the *Braintrak* library [75, 51, 76]. This toolbox implements a Markov Chain Monte Carlo (MCMC) method, using the Metropolis-Hastings algorithm for model fitting. The analytic power spectrum of the model, as defined in Eqns. (16) & 20, was fitted to the empirical power spectra from 30-second windows in the data. The parameters implemented were restricted to the stability limits defined in the previous literature [51] to ensure the biological feasibility of the attained fitted parameters. Furthermore, the value of the gain parameters for all connections was limited to 20 ($|G_{ab}| < 20$) to reduce the sensitivity of the model to noise in the input.

2.4.1 Fitting metrics

Using the method described above, we fitted the parameters of the corticothalamic model to empirical data. We used chi-squared (χ^2) error for model optimization, calculated between the 1 and 45 Hz frequency bins. The optimisation target function aims to reduce the error χ^2 between the empirical and model-generated power spectra. In this model fitting approach, the parameter space is firstly explored during a random walk with a length of 50,000 and with large step sizes, accepting the top points which get us to a region close to the target values. After this "burn-in" period, Braintrak takes smaller steps to approximate the ground truth more closely.

$$\chi^2 = \sum_j W_j |\frac{P_j(x) - P_j^{exp}}{P_j^{exp}}|, \tag{24}$$

where j corresponds to each unique Fourier transform frequency bin of the power spectra. P_j^{exp} is the expected model power spectrum and $P_j(x)$ is the empirical power spectrum for that bin. The term W_j is a scaling factor to increase the effect of the lower frequency bands compared to the higher frequency bands ($W_j \propto f^{-1}$), thus increasing the sensitivity of the optimizer to the high frequency bins of the power spectrum and minimizing its sensitivity to lower frequencies. This can be valuable in reducing artifacts observed in the EEG data, since the main artifacts affecting our 1-45 Hz window include the glossokinetic, movement, eye blink, and sweat artifacts, all of which produce low-frequency artifacts that must be mitigated [77, 78].

The complexity of the model was also calculated using the Akaike Information Criterion (AIC) [79]. AIC denotes the complexity of the combinations of parameters that together yield the model power spectrum. The lower the value of AIC, the simpler (or more parsimonious) the model. High values of AIC may denote overfitting of power spectra by fitting complicated combinations of parameters. AIC is described by this formula:

$$AIC = 2k - 2\ln(\mathcal{L}), \tag{25}$$

where k is the number of model parameters (9 in this case), and $\mathcal{L}$ is the maximum of the likelihood function for this model.

3　Results

Our analyses in this study evaluated the methodology for neurophysiological modelling-based brain state estimation described above [51, 75] across several datasets recorded from research-grade and consumer-grade devices. In the following, we first summarize several key characteristics of the sleep EEG and hypnogram data used, and then turn to our model fitting results and their physiological interpretation.

3.1　Comparison of EEG features across sleep datasets

3.1.1　Hypnogram-based sleep stage compositions

As can be seen in the group-averaged hypnogram summaries (Fig. 2E), sleep stages N1-N3 and W (wake) are well-sampled across all five of the studied datasets. REM sleep is also present in all datasets except Nap-EEG, since the 30-minute recordings used in that study are much shorter than the average 80-100 minute mark at which the first episode of REM appears [80]. The other four datasets all include several dozen minutes of REM-labelled sleep periods on average, although for the EDF-X dataset the average percentage of time spent in REM across the subject group is only 2.99% and the REM sleep latency (as seen in Fig. 2A), is unusually long (average of around 515 minutes).

Improved sleep quality with mobile EEG

Sleep efficiency (SE, the percentage of time from the whole recording spent in sleep) between all datasets is comparable, averaging between 66.61% and 92.86% — although we note that for three of the five datasets, this value is below the recommended healthy range of 80-100% [81]. This value is the highest for the Muse S dataset with 92.86%. Sleep maintenance efficiency (SME, the percentage of time in sleep between the first and last stages of sleep) is also comparable for all datasets, with Nap-EEG performing the best among all (98.23%), and Muse S performing best for the whole-night recordings. For Muse S, this is most likely due to the improved comfort factor associated with the light and non-intrusive nature of mobile EEG headsets and the fact that the recordings are done at the subjects' home and in their familiar and comfortable beds. This interpretation is further corroborated by comparing the subject-averaged total minutes of "Wake After Sleep Onset (WASO)" between the different datasets. Muse S subjects spend an average of 20.50 minutes awake after sleep onset, which is also the lowest among the whole-night recordings, further demonstrating that the subjects have less interrupted sleep when using mobile EEG equipment. This value is comparable to the range of 54-88 minutes for the three research-grade whole-night recordings (DOD, EDF-X, and WSC). Subjects also fall asleep faster with Muse S (average of 11.30 minutes Sleep Latency) than all other whole-night recordings, with the exception of EDF-X, for which an accurate sleep latency could not be calculated (see Supplementary Material section 3.1). Thus, mobile EEG can contain a more naturalistic and representative sample of physiological states and sleep stages in a full night of sleep than conventional research-grade EEG, and help us evaluate the normative trajectories of their changes in health and disease.

Figure 2: ***Comparison of sleep architecture metrics and power spectral features across datasets.*** **A)** Average metrics evaluating sleep architecture and sleep quality in a recording, compared across the datasets. Error bars represent the standard deviation of the values per subject. **B)** Distribution of $1/f$ exponents in each dataset, separated across the various sleep stages. **C)** *bottom:* Subject-averaged area under curve (AUC) of the EEG spectral power for each frequency band. *top:* Example power spectra from the dataset noted in the bar plot, with each under-curve band highlighted. **D)** Subject-averaged FOOOF-calculated peak power for each frequency band. **E)** Composition of the sleep stages forming the hypnograms in each dataset.

3.1.2 EEG power spectral features across sleep stages

In the next stage, we used two different prevailing approaches for a quantitative comparison of the power spectra across datasets:

I) In one approach, the background scale-free $1/f$ activity is separated from the oscillatory activity with defined frequencies, and then the properties of each periodic & aperiodic component is examined. We used the Python library FOOOF for this task, as described in section 2.2.3.

II) The second approach is to compare the areas under curve (AUC) for the power spectral plots in each of those frequency bands. We used the trapezoidal integration method to calculate the AUC.

Aperiodic components vary across datasets and track sleep stages robustly

We separated the periodic and aperiodic components of the power spectra using FOOOF and calculated the exponent of the $1/f$ component for each power spectrum. The range of the $1/f$ exponents from different datasets are vastly different. Regardless of this variance, moving from lighter to deeper NREM sleep is generally associated with an increase in the value of the exponent and the value is then reduced again in the transition to REM sleep (Fig. 2A).

The scale-free changes in the slope of the $1/f$ component, which are thought to be results of background physiological processes and general brain activity patterns [82] can disproportionately increase the AUC in low-frequency domains, including delta (slow) and sub-delta (infra-slow) frequencies. We see this by comparing the power of each band as compared using FOOOF vs. AUC methods in Fig 2: The datasets Dreem and WSC, which possess the highest average $1/f$ exponents (as noted in Fig. 2A) demonstrate highest AUCs in the delta band (Fig. 2B), but by using FOOOF to separate the periodic & aperiodic components and examine the height of each unique peak apart from the contributions of $1/f$, we observe that the delta band is the least dominant of all peaks 2C.

In fact, we see in that comparison that delta is the highest-powered band compared to all other bands in each dataset if we only use the AUCs, but separating the $1/f$ component using FOOOF relegates the rank of the delta peaks amplitudes to the last place in all cases. We further note that if we rely only on the AUCs, the alpha peaks are either the highest or the second-highest calculated peaks in all datasets with the exception of the Muse S. This demonstrates the importance of the contributions of the aperiodic component to the power of each frequency band in the power spectra and how it affects the traditional AUC methods for calculating band power. Solely observing the AUCs for each of those bands without this separation would have concluded a domination of low-frequency activity for all datasets, with minimal difference across the datasets, but FOOOF allows us to make that distinction between the footprints of each recording setup on band peak amplitudes.

In the MSD data, the peak height is lower than the other datasets for most bands (Fig. 2C). Separating the $1/f$ components from the raw power spectra in for this dataset almost completely reverses the order of the peaks with regards to frequency: the $1/f$-separated peaks are highest for gamma and the lowest for delta, but the the AUCs are

highest for delta and lowest for gamma. This points to the fact that for this dataset, the $1/f$ component is more dominant than the periodic component and that the $1/f$ exponent and is a robust feature separating the sleep stages and the physiological state of the subject, as evidenced by its strong variation across different sleep stages for Muse S (Fig. 2A), consistent with prior literature showing that most of the variation in individual sleep stages can be explained by $1/f$ components [36]. The $1/f$ exponents are one of the canonical features of our physiological model's power spectra as well, as described in Robinson et al. [30], where the power law of the power spectra is a defining feature of the system, directly correlating with cortical gains and the primary oscillation frequencies, thus also making it a suitable criterion for tracking the activity of the corticothalamic circuitry.

3.2   Physiological modelling

Using the neurophysiological model of thalamic circuitry described in Abeysuriya et al. [51], Abeysuriya and Robinson [75], we fit the EEG power spectra across the various datasets. Despite the considerable difference in the properties of the EEG power spectra across these datasets, the model still performs well in fitting to all data. Goodness of fit was satisfactory, with all models demonstrating similar distributions of error (χ^2) and model complexity (AIC), meaning that the model is fitting closely to power spectra without over-complicating the model parameters. The error distribution is marginally higher for Dreem than for other datasets which could be explained by the wide distribution of $1/f$ exponents in this dataset (Supplementary Fig. S1). The resulting fitted model parameters from all datasets exhibit patterns of change across sleep stages that are similar and in line with prior literature on thalamocortical communication in sleep. Corticothalamic communication is reduced in sleep and decreases further as the subject goes into deeper NREM sleep stages.

3.2.1   Physiological transitions from light to deep NREM

The regular progression of sleep stages throughout a night of sleep commences with wakefulness (W), then transitioning to light NREM sleep (N1), followed by deeper stages of sleep (N2 and N3), eventually reaching REM sleep. Each individual cycles through the REM/NREM stages multiple times, with the cycles taking an average of 90 minutes [80]. In this transition within NREM sleep from N1 to N2 and to N3, the physiological properties of the functional brain circuits change along a clear trajectory [51], which we quantify using our neurophysiological modelling approach.

We first demonstrate this characteristic trajectory [51] and the differences between sleep stages using power spectra and fits from the Nap-EEG dataset. Similar overall results are obtained with the other four datasets, which are detailed further in the Supplementary Material Fig. S4. During the transition from W to N3, the $1/f$ exponent becomes larger and peaks in the alpha frequency band are reduced as deeper stages are reached (Fig. 3A). Two major patterns are observed in the neurophysiological model in conjunction with this change in the power spectra:

Figure 3: *Tracking the changes in the parameters in different sleep stages in the Nap-EEG dataset.* **A**) High-level schematic of the corticothalamic model, as described by the corticocortical (x), corticothalamic (y), and thalamothalamic (z) connection strengths. **B**) Average empirical vs. fitted spectra in the entire dataset, separated by sleep stages. **C**) Average power spectra in each sleep stage, separated between empirical and fitted. **D**) Box plots comparing the distribution of x, y, and z parameters across different sleep stages. Stages with significant difference in mean parameters are denoted by (*). **E**) Visualization of the xyz time series in conjunction with the hypnogram for one complete recording. **F**) Comparison of fitted and empirical power spectra at notable points in the whole-recording xyz time series with extreme x or y values, noting the associated alterations in the power spectra. **G**) Schematic demonstrating the change observed in the following panels. As the subject transitions from light to deep NREM, the connection strength in the corticocortical circuit is increased and the connection strength of the corticothalamic circuitry approaches zero.

Corticocortical amplification is increased from W to N3

Figure 3C shows the distribution of the corticocortical (x), corticothalamic (y), and intrathalamic (z) loop gain parameters across all epochs in the Nap-EEG dataset. The parameter x, calculated according to Eqn. 21, represents the net corticocortical excitatory connection strength. This parameter takes values between 0 and 1, with values close to 1 demonstrating highest degree of excitatory corticocortical amplification via the the excitatory projections connecting the various cortical regions and lowest corticocortical inhibition.

As can be seen, the progression from lighter to deeper sleep stages (N1 → N2 → N3) is associated with an increase in the average value of x, with estimates of this parameter in N3 approaching its maximum value of 1. This observation was confirmed statistically with an independent-samples *t*-test, which showed a significant increase in x from W to N3 ($t = 17.29$, $p < 0.001$). The corresponding comparison was also statistically significant in the other four datasets

(Supplementary Fig. S4). This points to an association between deep NREM sleep and reduced thalamocortical drive of cortical dynamics. The reduction in the absolute value means that the bottom-up thalamocortical drive, in either the inhibitory or excitatory modes, is reduced in NREM sleep.

Bottom-up thalamocortical modulation is reduced from W to N3

During the transition from lighter to deeper stages of NREM, the distribution of y parameters–which indicates how strongly the thalamus drives cortical dynamics via thalamocortical projections–becomes narrower and more leptokurtic, with the absolute value of y decreasing and approaching 0. Based on the properties of the y circuit parameter, this signifies a reduction in the influence of thalamocortical gains (both inhibitory or excitatory). Per Eqn (22), positive values of y would denote the dominance of the excitatory part of the corticothalamic loop (G_{ese}) over the inhibitory part (G_{esre}), and hence net excitatory bottom-up stimulation thalamo-cortically. In contrast, negative values of y signify the dominance of the term G_{esre}, where the inhibitory effect is due to GABAergic projections from the thalamic reticular nucleus. The negative value y therefore denotes a net inhibition applied to the cortex by the thalamus. As can be seen in the middle panel of Figure 3C the absolute value of y in N3 sleep approaches 0, signifying the absence of excitation or inhibition driven from the thalamus towards the cortex. Similarly to the previous section, we used the independent-samples t-test to compare the $|y|$ values between stages W and N3, demonstrating a significant reduction in the parameter ($t = -11.48$, $p < 0.001$). Again, this effect was replicated across the other four datasets (Supplementary Fig. S4).

To further confirm that this relationship constitutes an ordinal trend across all four sleep stages, we assigned NREM sleep depth values of 0-3 to stages W-N3, and performed Pearson's r-test with these and the lumped circuit gain parameters. This returned significant correlations between sleep depth and both the absolute corticothalamic circuit gain $|y|$ ($r = -0.226$, $p < 0.001$) and corticocortical circuit gain x ($r = 0.301$, $p = 9.83 \times 10^{-45}$).

3.2.2 Relationship of physiological circuit parameters to periodic and aperiodic EEG power spectrum features

We have demonstrated physiological model-based extraction of information on corticothalamic system state from windowed EEG power spectra across sleep stages and in multiple datasets. A key question that this analysis raises is "what features of the computed spectra contribute to the estimated physiological parameters"? As noted in Figure 2, different sleep stages have characteristic fingerprints across the periodic and aperiodic ($1/f$) components of the EEG power spectrum, that are generally consistent across all five datasets studied here. Given the evident associations of each sleep stages with the corticothalamic circuit activity parameters (Fig. 3) as well as the aperiodic components of the power spectra (Fig. 2A), we aimed to directly determine the interplay between the strength of various thalamocortical sub-circuits (gain (G) parameters) and the broadband power and $1/f$ exponents of the power spectra, along with a comparison of how each of these parameters relate to the changes in the power spectra.

To determine more precisely how the aperiodic components of the empirical power spectra give rise to the fitted physiological model parameters, we studied the correlation of these parameters across the entire Nap-EEG dataset with the slope and offset of the $1/f$ (Fig. 4A). This step was repeated for all datasets and all gain parameters in the Supplementary Figs. S5 – S9. We then examined further the contributions of isolated individual parameters to the power spectrum structure, by first initializing models at the estimated parameter values from a typical fitted epoch, and then systematically manipulating each gain (G) parameter, per Eqns. (16) –(19), observing changes in the model power spectra (Fig. 4B).

Lower thalamic and corticothalamic gains generate a steeper $1/f$

In all datasets, G_{ese}—the circuit gain related to the positive feedback loop between the cortex and the thalamic relay nuclei—has a significant negative correlation with the exponent and the offset of the $1/f$ component (Figure 4A). That is, stronger excitatory corticothalamic feedback, signified by a higher G_{ese}, results in a flatter $1/f$ component and a reduced area under the curve in the low-frequency domain. In general, we observe in the model fit results that epochs from sleep stage N3 tend to cluster in regions of parameter space with lower G_{ese} values and higher $1/f$ exponents. Thus, through the progression from wake to light and into deeper sleep, the exponent of the $1/f$ components increases, due to the progressive weakening of the excitatory corticothalamic feedback loop, as predicted analytically in Robinson et al. [30]. Concurrently with this, a positive correlation is observed between $1/f$ exponents and the negative-valued (inhibitory) gains of the loops associated with the thalamic reticular nucleus (G_{esre} and G_{srs}). As can be seen clearly in Figure 4B), a decrease in the $1/f$ exponent and offset (flatter aperiodic component) is observed as both of these negative inhibitory gains become more pronounced. This is in line with observations by Abeysuriya et al. [51], in which a significant reduction in the strength of thalamothalamic inhibitory connections–represented by thalamo-thalamic circuit gain (z)–was observed in deeper sleep stages, which possess power spectra with higher $1/f$ exponents (Fig. 2A).

Greater corticocortical excitation can yield steeper $1/f$ components

The analyses in Figure 4B also show that amplifying the gains in the cortical excitatory (G_{ee}) or inhibitory (G_{ei}) connections results in a broadband increase in spectral power. The increase in G_{ee} is slightly more effective at increasing the lower frequency components. However, unlike the effects noted above for G_{ese}, G_{esre} and G_{srs}, modulation of G_{ee} or G_{ei} was not found to influence the observed spectra in this way in the datasets studied, and the correlations between these gain parameters and the $1/f$ components are not strongly correlated (Supplementary Fig. S7).

To further evaluate this effect of the gains on $1/f$ exponents, the analysis above was conducted for all the other datasets in addition to Nap-EEG (Supplementary Figs. S5–S9) which confirms this effect for most of the datasets. All of the datasets, with the exception of WSC, demonstrate significant moderate correlations between $1/f$ exponents and the corticothalamic gains (G_{ese} and G_{esre}) and insignificant or significant mild correlations between cortical gains and the $1/f$ exponents. Abeysuriya et al. [51] report G_{ee} to be the most variable gain among sleep stages. Our findings demonstrate that this is not mediated by the $1/f$ exponents at least in these datasets. In the case of WSC, the distribution

Figure 4: *Correlation between model parameters and the shape of the power spectra.* **A)** The correlation between the exponent of the aperiodic component with each thalamic gain parameter and x, y, z, calculated using Pearson's r-test. p- and r-values reported in each legend. **B)** The effects of the incremental increase of each gain parameter on the shape of the power spectra. The baseline power spectra (in black) was taken from the fit to a real 30-second epoch (in Nap-EEG in stage N2 with no prominent peak). The absolute value of each gain value was increased in steps of 0.05 and the power spectrum generated from those parameters was generated (deducting 0.05 from negative gains and adding 0.05 to negative gains in each step. The colour bar denotes the values specified, starting from the darker color as the baseline and changing the gain parameters successively towards the lighter copper colour. **C)** Correlation of the gain parameters with the EEG Band power and the $1/f$ exponent in the Nap-EEG dataset **D)** Correlation of the gains with the power spectra resulting from the incremental sweep of the gain parameters.

of the $1/f$ exponents does not differ greatly between sleep stages as seen in Fig. 2A, which suggests that it might not be as adequate an indicator of the brain's physiological state as it is for other datasets.

Fitted model parameters represent the power spectra robustly

In the next step, we examined the correlations between the fitted gain parameters and the power of each band and the $1/f$ exponent in the EEG power spectra, for each dataset and for the power spectra attained through the manual modification of single connection strengths. Pearson's r-tests were implemented between the power of each band for each power spectrum and the gain parameters corresponding to it. The statistical significance of each correlation was corrected for the repeated hypothesis testing using the Bonferroni method [83] to prevent the false detection of patterns. It is observed that the offset & the exponents of the aperiodic components correlate with fitted parameters in the same direction and with similar strengths, as evidenced by Figs. S5 to S9. This analysis was completed for the Nap–EEG dataset in Fig. 3C and was repeated for all other datasets and all fitted and calculated parameters in the Supplementary Material Figs. S10 to S14.

It is notable that the gain parameters G_{ee} and G_{ese} are excitatory and hence have a positive sign. Meanwhile, the gains G_{ei}, G_{esre}, and G_{srs} are negative, and hence have a negative sign. Stronger connection strengths correspond to larger absolute values of these gains (more positive for the excitatory and more negative for the inhibitory). This must be taken into account when interpreting the correlations in Fig. 4.

Comparing the correlations for the fitted parameters (Fig. 4C) and the power spectra generated by changing the parameters (Fig. 4D) reveals that the positive gains are correlated and the negative gains are anti-correlated with the exponent of the $1/f$ component. The direction of the correlations is similar for both instances (the sign of the r-value). However, the intensity of the correlations ($|r|$) is larger for the manually-set parameters, which could be explained by the fact that all model parameters can also change for the fitted parameters, while in the manually-set instance, all parameters are fixed other than the changed parameter. The correlations between the changed parameter and the exponent in the set parameter instance is very strong ($|r| > 0.86$) for all parameters except for that of the cortical excitatory self-connection (G_{ee}). Examining the parameters fitted to the Nap-EEG dataset, those correlations are moderate ($0.5 < r < 0.6$) for all parameters except for the cortical excitatory connection strength (G_{ee}) and also the cortical inhibitory connection strength (G_{ei}).

Physiological model captures changes in the spectra driven by both aperiodic and periodic components

As seen in Fig 4D, in examples such as the direct modification of the cortical inhibitory connection strength (G_{ei}) alone, we observe a very strong correlation of the gain with the EEG band power calculated via the "area under curve" (AUC) method. But by separating the periodic and aperiodic components using FOOOF, we observe a different phenomenon; the aperiodic ($1/f$) exponent is increased in the same direction and approximately with the same intensity as the AUC band powers, but we only see one correlation with the FOOOF-discerned peaks in the alpha band, and no peaks were generated in any of the other EEG frequency bands. The case for the cortical excitatory connection strength (G_{ee}) is somewhat different, where the AUCs of almost all bands except for delta have a very strong anti-correlation with the gain values ($r > 0.84$), but the FOOOF-discerned $1/f$ exponent does not show a notable correlation ($r = 0.07$) and the only correlating FOOOF-detected peak is alpha, with a weak correlation ($r = 0.17$).

As seen in Fig 4C, Comparing the FOOOF and AUC-detected phenomena in the parameters fitted to the Nap-EEG dataset demonstrates this common thread as well; all the connection strengths involving thalamus (G_{ese}, G_{esre}, and G_{srs}) exhibit weak to moderate correlations with the AUCs of EEG bands, and they similarly correlate moderately with the $1/f$ exponents. Comparing the r-values of the correlations between FOOOF and AUC-measured peaks with these three gains demonstrates a strong disagreement between these two common metrics of power band estimation, where only two of the peaks are strongly detected by both methods, and in the case of theta band activity, they change in the opposite directions. This is due to the effects of the change in the $1/f$ component, where the exponent of this aperiodic component is correlated with the area under the low-frequency bands (e.g., delta and theta), and anti-correlated with the high-frequency bands. In other words, the shape of the power spectra and how it follows power law can change the detected values for those power bands.

Despite the variations, several consistent phenomena can be identified that are associated with higher absolute values of any of the three thalamic gains ($|G_{\{ese,esre,srs\}}|$:

- The $1/f$ component is moderately decreased ($-0.51 < r < -0.56$).

- The AUC for the delta, theta, and alpha bands is decreased with a weak-to-moderate correlation (respectively, $-0.43 < r < -0.61$, $-0.18 < r < -0.35$, and $-0.18 < r < -0.47$).

- The AUC for beta and gamma band is weakly decreased (respectively, $-0.34 < r < -0.38$, $-0.18 < r < -0.28$).

- FOOOF peaks in theta are increased with a moderate correlation ($-0.44 < r < -0.68$).

.

We observe that the gain parameters attained by fitting the power spectrum can strongly capture the changes in both the periodic and aperiodic components of the power spectra. This model can describe both the broadband changes in the power spectra as described by the AUC measurements and the $1/f$ components, and the sharp peaks in narrower frequency bands, as measured via the FOOOF peaks. Among the fitted datasets, the $1/f$ exponents and the changes in the AUCs tend to describe the fitted gain parameters better than the sharp FOOOF-detected peaks in most of the datasets, as evidenced by Figs. 4C & 4D, pointing to the power of this model of EEG power spectra to represent the different power spectral features arising from the physiological properties of canonical brain circuitry.

3.2.3 Model parameters are associated with mental health markers

In this step, the fitted parameters of the Wisconsin Sleep Cohort (WSC) were analyzed in conjunction with the biomarkers included in this dataset. Using Pearson's r test, the correlations between each of the 227 biomarkers and the average value per night for each of the 9 fitted model parameters were examined. To correct for the repeated pairwise correlation analysis, we used the False Discovery Rate (FDR) method introduced by Benjamini and Hochberg to correct the p-values in a ranked manner, taking into account the probable false positives in repeated testing [84, 85], bringing the p-value threshold for rejecting the null hypothesis from 0.05 to approximately 0.0396. There were 46

significant correlations between the parameters, but they were all weak–with the highest $|r|$-value for any correlation being 0.16. The table containing all these correlations is included in the Supplementary Material, Table S1. Despite the weak correlations, discernible patterns arise when observing which specific parameters correlate with which biomarkers. For example, the connection strengths in the inhibitory thalamothalamic feedback loop (G_{srs}) and the the full thalamothalamic circuit (z) exhibit correlations with the administration of various medication groups and neurochemicals, such as alpha blockers, selective serotonin reuptake inhibitors (SSRIs), diabetes medication, and alcohol. This could be attributed to the various efferent cholinergic [86, 87] and serotoninergic [88] synapses that TRN receives, along with the complex calcium-dependent dynamics underlying its firing state and frequency [29], which would potentially be altered with the administration of these medications. The average thalamocortical circuit (y) exhibits significant correlations with biomarkers related to sleep quality and sleep debt, such as waking through sleep or daytime sleepiness. This is corroborated by evidence linking sleep deprivation to increased hyperexcitability and reduced specificity and functional connectivity in the thalamocortical connections [89, 90].

SSRI medication alters thalamocortical connectivity during deep NREM

Given the importance of changes in sleep as a comorbidity of many mental health disorders, we next focused on studying the correlations of the average physiological circuit parameters fitted over one night with the WSC variables related to mental health–namely trait and state anxiety, scores from the Zung self-rating depression scale [91], and antidepressant medication. The Zung index is a normalized integer score value between 25 to 100, wherein the scores between 50 and 59 are scored as mild depression, between 60 and 69 as moderate depression, and any value higher than 70 as severe depression. Across our physiological model parameters we observed a weak but significant correlation ($r = 0.10$, $p = 0.01$) between the consumption of SSRI antidepressant medication and the gain of the thalamothalamic circuit (z).

We first tested whether the SSRI medication successfully reduces the severity of depression. In total, in 437 (78.60%) of the fitted recordings, the subject reported being on SSRI medication, and in the other 119 (21.40%), they were off SSRI medications. An independent-samples two-sided t-test demonstrates that the on-medication group has a significantly lower Zung index ($t = -9.850$, $p < 0.001$) than the off-medication group.

Next, we compared the composition of the parameters in different stages of sleep between the on- and off-medication group. We repeated the analysis in Fig. 3 on the on-SSRIs and off-SSRIs groups separately as well, to see if the transition from W to N3 yields the same reduction in $|y|$ and increase in x in both subgroups. Independent t-tests were used to compare the means of x and $|y|$ between the two stages W and N3. In the off-medication group, we see a significant *depth of sleep* effect–where the reduction in $|y|$ is significant and negative ($t = -7.778$, $p < 0.001$) as is the increase in x ($t = 2.004$, $p < 0.001$). In the on-medication group, both of these effects are significantly reduced, with no significant reduction in N3 thalamocortical circuit gains ($t = -7.778$, $p = 0.997$) and very slight increase in the x values ($t = 2.045$, $p = 0.020$).

Interactions between depression or SSRI biomarkers with parameters x,y,z are nonlinear

Finally, we explored the ability of our estimated neurophysiological circuit parameters to predict disease outcomes using ML-based biomarker stratification. To this end, we separated the data into training and test groups, with 80% of the fitted recordings in the training group and the other 20% in the testing group. We then trained a linear kernel Support Vector Machine (SVM) to test if xyz in the on and off-medication groups are consistently separated using this support vector. The task is to predict the on- or off-medication group (binary label, $\in [0, 1]$) from the average parameters ($[x, y, z]$) from each whole-night recording. The training and testing data separation and model implementation was done using the Python scientific computing library `scikit-learn` v. 1.1.2 [92]. The algorithm performed poorly at predicting the SSRI medication outcome. In contrast, the model only performs very well at predicting the on-medication class, and not the off-medication class (true positive=88.40%, false positive=11.6%), and both true and false negatives 0%. The model achieves an accuracy of 88.40%, but it is only the baseline that is using the load-imbalance between the positive and negative labels, and the precision is 0.000%. Up-sampling the on-medication group, such that they comprise 50% of all the labels, still does not help with improving the precision of this model. In summary, these results demonstrate that whole-night average parameter values cannot linearly predict whether or not the subject is on SSRI medication. For this task, we must utilize algorithms that accept added dimensions for the input matrix, such as convolutional neural networks (CNNs), or apply dimensionality-reduction techniques such as singular value decomposition (SVD) or Principal Component Analysis (PCA).

4   Discussion

In this work we aimed to study how brain activity changes across different sleep stages, in health vs. disease, and as a function of recording technology (research-grade vs. mobile EEG). EEG power spectral density was calculated in 30s windows matching those of the hypnogram, delineating the frequency-domain characteristics of oscillatory and aperiodic background brain activity. Then, we used a neurophysiological modelling method introduced by Robinson et al. [45, 51, 75] to estimate various physiological parameters of corticothalamic brain circuits, and observe how these parameters change over sleep stages. Multiple sleep EEG datasets were employed to replicate our principal findings and to demonstrate the usage of this approach in various research and non-research scenarios, including most importantly, using at-home sleep EEG recordings from the consumer-grade sleep EEG headset Muse S. Changes in the $1/f$-parameterization of the power spectra was shown to be significantly correlated with the corticothalamic gain parameters linked to bottom-up thalamocortical drive of the cortical activity, with the exponents becoming larger with depth of sleep (Fig. 4 and Figs. S5 to S9). Deeper NREM sleep stages were also observed to undergo a severance of effective bottom-up thalamocortical control, signified by reduced thalamocortical circuit gains ($|y|$) and increased cortical excitability, signified by elevated corticocortical circuit gain values (x) (Fig. 3 and Fig. S4). Administration of SSRI medication was observed to block this disintegration of corticothalamic connections in deep sleep. Finally, it was demonstrated that this physiological modelling approach can effectively integrate the periodic & aperiodic

components of the EEG power spectra more robustly than common PSD analysis techniques and provide a reliable and physiologically explainable parameterization of those spectra.

4.1   Key Results

Thalamic relay excitation increases the $1/f$ ***slope***

A central result that was consistent across most of the analyzed datasets was that whereas connectivity strengths for cortico-cortical connections (G_{ee} and G_{ei}, x) had negligible associations with the fitted $1/f$ offsets and exponents, strong correlations with $1/f$ features were seen for connections involving thalamic units (G_{ese}, G_{esre}, and G_{srs}). The pattern is such that the higher the value of the gains (either excitatory or inhibitory), the bigger the exponent of the $1/f$ component, and so the more steep the background trend.

Previous literature points to the importance of the thalamic reticular nucleus as a regulator of excitatory thalamic nuclei activity, including relay nuclei [93, 29, 28]. This thalamic control loop attenuates the excitatory drive from thalamic relay nuclei to the cortex, thereby regulating the activity of the cortical neural populations underlying measured EEG signals.

As described in the 2 section, in the Robinson model, the thalamic relay nuclei (s) constitutes the main excitatory output of the thalamus, whose influence is balanced by the thalamic reticular nucleus (r) that implements a negative feedback loop, extinguishing the thalamic relay excitatory output. The loop gain G_{srs} gain parameter summarizes how the balance between these two units ($s \to r \to s$) enables this inhibitory feedback. One implication of our modelling results is that increased thalamo-thalamic inhibitory activity, signified by increases in inhibitory thalamic gains, flattens the $1/f$ spectrum by inhibiting the thalamocortical circuit driving the cortex. When this inhibition is removed, the network-level disinhibition in the cortex leads to a more steep $1/f$ slope. Previous authors (Gao et al. [20], Lombardi et al. [82] have suggested that higher $1/f$ exponents can be regarded as a criterion for higher cortical inhibition, driven not by thalamic but by cortical inhibitory populations, whereas our findings concentrate on the bottom-up thalamo-cortical axis of communication and how increases in its absolute gains lead to increased $1/f$ exponents.

A caveat for this model, as noted by Abeysuriya and Robinson [75], is that the gain (G) parameters are dependent on both the steady-state neural field and the synaptic strength of each population (per Eqn.(15)) and changes in either parameter can lead to a rise in the gain parameters, yielding infinite solutions for the exact delineation of these two parameters. Furthermore, another simplification in this model is the assumption of random outgoing synaptic connectivity, leading to $G_{ei} = G_{ii}$ and $G_{ie} = G_{ii}$, which may imbalance this cortical E/I balance estimation.

Thalamo-cortical disinhibition during the progression from wake to deep sleep

Using the neurophysiological model of thalamocortical system, we demonstrated that with the transition from wake to sleep, concurrent with an increase in the $1/f$ exponents, the values of the corticothalamic circuit (x) increase and the absolute values of the thalamo-cortical circuit gain (y) approach 0 (Fig. 3C). As we move from lighter sleep to deeper

NREM sleep (from wakefulness to sleep stage N1, and then to N2, and then N3), the values of x increase further, such that in N3, their values are distributed very narrowly, close to the maximum value of 1.0. Concurrently, the value of $|y|$ decreases, such that in N3, it has a narrow distribution close to 0 (both in figure 4C. As noted, the values of y depend not only on the existence of thalamocortical activity, but also on the nature of its contribution (i.e. whether it influences inhibitory or excitatory activity from the thalamus to the cortex) [51]. In this sense, the deeper stages of NREM sleep, especially N3, involve increased cortical excitability, but at the same time, the thalamocortical population is insensitive to activity propagated through the thalamus. This highlights prior work by Nir et al. [40] demonstrating that EEG slow wave activity during deep NREM sleep is regionally and not globally synchronized, and that the oscillatory phases vary spatially over the cortical surface. Additionally, Massimini et al. [41] demonstrate that these slow waves can be locally interrupted and entrained using transcranial magnetic stimulation (TMS) at 1 Hz, suggesting a cortically-generated dynamic where local stimulation has the capacity to disrupt them. This pattern of dis-facilitation and dis-inhibition in N3 are in line with previous work using biophysical models denoting a reduction in thalamic excitation or inhibition in the brain in slow wave sleep [32] despite an increase in cortical synaptic strength [38, 94, 95].

The sign of y depends on the dataset, rather than the immediate power spectra

We found the mean and the mode of the fitted values for the circuit gains (x, y, and z) to be slightly different in various datasets. This could be justified by the different amount of time spent in different sleep stages (Fig. 2A) and thus the different oscillatory regimes dominating the data (Fig. 2C).

Despite the observation in Abeysuriya et al. [51] reporting excitatory thalamocortical regimes noted by positive y values in wakefulness decreases to negative values with the transition from wake to sleep, we observed that the numerical sign of y depends more on the datasets used than the stages of sleep. We noted the frequent "sign" of y to changes across datasets, irrelevant of wakefulness vs. sleep. For instance, 72.38% of all y values among 120,855 wake epochs in the EDF-X dataset were negative and 66.20% of all 2,119 epochs during sleep in the Nap-EEG dataset were positive. Hence, our work suggests a more nuanced take where the transition from wake to sleep shifts to a more inhibitory regime, marked by reductions in y along the depth of sleep axis, but that does not reflect an overall domination of bottom up inhibition as soon as sleep is initiated. Future work comparing the topography of these effects can shine light on the network-level variations in such changes.

4.2   Limitations and Next steps

In this work we have focused primarily on changes in model parameters associated with transitions between sleep stages. However, these stages are far from the only physiologically-significant features we can extract from sleep EEG datasets. Other phenomena of interest could for instance be the dynamics of alpha activity in the final minutes of transitioning from wakefulness to sleep, which prior work has found to be associated with pathologies such as insomnia and sleep deprivation [96–98].

Studying sleep spindles

Another area of interest for future work that the framework presented here can be well suited to studying is transient oscillatory events in EEG traces such as sleep spindles and k-complexes. Abeysuriya et al. [49] utilize this model of the EEG power spectra to generate the power spectral density resulting from spindle generation. In this work the authors use stability analysis of the corticothalamic system to predict the nonlinear harmonic frequencies of the spindle peaks. In a follow-up study [50], they then demonstrated the existence of these spindle harmonics in empirical EEG data, as well robust fitting of an extended version of the model. In this case, the nonlinear harmonic frequencies of the spindle are resulting from the thalamo-thalamic feedback loop, and differ from the linear harmonic frequencies associated with the primary alpha and beta peaks. This study of the predicted harmonics in EEG power spectra can be useful for comparing the specific underlying corticothalamic connections generating such activity.

A long line of work on murine sleep EEG has demonstrated the importance of the corticothalamic system in generating sleep spindles. Empirically, shifts in the excitatory and inhibitory firing rates of various neural populations have been implicated in the generation of sleep spindles, initiated by a transition in the thalamic reticular nucleus [28, 99]. The circuit mechanisms and underlying mathematical structure of spindle generation in these detailed thalamic models [100] and the coarser-grained corticothalamic models [50, 49] may be related, but are not identical. They could be best understood as either complementary or competing candidate theories of this prominent phenomenon observed in human sleep EEG. An important direction for future work should be to compare and characterize the relevant parallels between these two frameworks, such as relating the models' excitation and inhibition parameters to the $1/f$ EEG features across sleep stages described in this study.

The current work does not include any in-depth assessment of sleep spindles, as only one of the datasets used here (Nap-EEG) contained consistent spindle events with expert labelling in the EEG data. Separation of the spindles in the other datasets requires expert or machine learning-based detection of the spindles, which was out of the scope of the present work. Furthermore, sleep spindles start and end in short spans of approximately 2 seconds, which is much shorter than our standard power spectrum epochs of 30 seconds. The window sizes would therefore need to be substantially shortened to accommodate spindle-oriented analyses, which would in turn deleteriously increase the proportion of noise-driven peaks in the spectra, making model fitting less stable and consistent. In future work, we will study spindles as they appear in mobile EEG data specifically, and characterize changes in their occurrence, frequency, length, amplitude, etc ove sleep stages. These data can then be used to inform fitting of the corticothalamic model to the spindle PSDs, per [50, 49], thereby mapping these empirically-observed changes to transitions in the model parameters.

As we have indicated, sleep spindles are oscillatory events which begin, rise, decay, and then conclude in a well-characterized and parameterizable fashion. It is also notable that the phase and frequency of the spindles can vary spatially. In this study, due to computational limitations and variation in the data sets, we were restricted to fitting the power spectra to only one EEG channel. A logical next step would be to individually fit *all* EEG channels from the datasets, and analyze the parameters in the channel space, or to implement the spatial modes in the analytic power

spectrum (k). The topography and spatial modes of these trajectories are topics of active interest in the field [101], and observing their changes in wakefulness and sleep in health and disease, especially in the context of mobile EEG, has clear scientific and clinical value.

In the future we will also consider transform-based machine learning, in which the transform is not merely a pre-processor but is also itself part of a neural network [102, 103], as well as phase-based methods [104]. Indeed, much of the important information in oscillatory activity during sleep is arguably better represented in terms of phase space, as well as scale space, phase scale, and the chirplet transform [102, 103], because sleep stages are often characterized by changes in frequency (acceleration of phase) [105–107]. Chirplet-based analytical approaches potentially offer a more biologically sympathetic perspective on neural signal analysis, which can aid corticothalamic modelling of sleep neurophysiology by better capturing time-varying frequency modulations in the EEG [108].

Implementation of this pipeline in large cohorts

Combining the above-outlined strategy with *de-novo* at-home sleep recordings using the Muse S headset, with a larger sample size than the N=10 studied here, is a promising extension of the present work. In particular, this has major potential for studying sleep EEG features and the physiological underpinnings at-scale - both in terms of number of subjects (hundreds to thousands) and number of sleep sessions per subject (dozens or more). Adding biomarkers related to sleep quality and general health, for example through surveys or integration with other wearable biometric devices, would also be of great utility in delineating the physiological basis of those biomarkers.

In the present work, simple features of the distribution of 9 fitted corticothalamic model parameters across a night of sleep (such as mean, mode, and standard deviation) were used. In future work, using data-driven dimensionality reduction techniques to identify underlying sub-structures within these parameter values may prove an effective use of the physiological model outputs to help predict the health status and outcomes, both in extant datasets such as WSC [64], as well as new Muse S recordings with surveys described above.

4.3   Conclusions

In summary, our work has showcased the adaptability and reliability of this neurophysiological model [51, 75] for generating the trajectory of brain states during a sleep recording, utilizing a range of EEG data with various setups and recorded in various locales. This method adds another degree of physiological interpretability to the observations made based on EEG time series and power spectra. A robust interplay was observed between the aperiodic and periodic power spectral components, fitted model parameters, and their stage-dependent dynamics. This method can be effective for comparing sleep EEG between and among subjects and inferring latent health or disease states.

5   Data and Resource Availability

1. The Sleep EDF–Extended (EDF-X) dataset version 1.0.0 used in this work is available on PhysioNet at `https://physionet.org/content/sleep-edfx/`, and described in Kemp et al. [56].

2. Dreem-Open-Datasets (DOD-H & DOD-O) and their annotations are available openly online. Full instructions on acquiring the data are included at `https://github.com/Dreem-Organization/dreem-learning-open`.

3. The Wisconsin Sleep Cohort (WSC) dataset is available on the National Sleep Research Resource (NSRR) [63] at `https://sleepdata.org/datasets/wsc` and can be accessed openly for academic research.

4. The Nap-EEG dataset is available via the Open Science Foundation (OSF) at `https://osf.io/chav7/`. Further information about the data is included by the authors at `https://github.com/nmningmei/Get_Sleep_data`.

5. The MCMC model fitting algorithm implemented on MATLAB is available at `https://github.com/BrainDynamicsUSYD/braintrak`.

6. The analysis and visualization code used in this paper is included in this GitHub Repository: `https://github.com/GriffithsLab/MorshedzadehEtAl2024_sleep-eeg-nft`.

6   Disclosure Statement

MAJ and CAA are full-time employees of Interaxon Inc. at the time of writing.

7   Acknowledgements

Computing: The computing infrastructure for this work included the CAMH Specialized Computing Cluster (SCC) and the cluster Narval at Calcul Québec, part of the Digital Research Alliance of Canada. The SCC is funded by the Canada Foundation for Innovation's (CFI) Research Hospital Fund. These platforms allowed for quick and robust large-scale model fitting to entire datasets and whole-dataset analyses.

Grants: This work was completed as a part of T.M.'s MSc thesis at the Institute of Medical Science (IMS), Temerty Faculty of Medicine, University of Toronto. Project funding and stipend for this degree were provided via UofT EMHSeed. The project was further supported by the CAMH Discovery Fund and the Krembil Foundation. The Wisconsin Sleep Cohort (WSC) Study and data was supported by the U.S. National Institutes of Health, National Heart, Lung, and Blood Institute (R01HL62252), National Institute on Aging (R01AG036838, R01AG058680), and the National Center for Research Resources (1UL1RR025011). The National Sleep Research Resource (NSRR) was supported by the U.S. National Institutes of Health, National Heart Lung and Blood Institute (R24 HL114473, 75N92019R002).

Author contributions *(in alphabetic order)*: **CA**: Methodology, Resources, Data Curation, **DM**: Writing - Original Draft, Visualization, **JDG**: Conceptualization, Methodology, Software, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration, Funding acquisition, **KK**: Writing - Review & Editing, **MAJ**: Methodology, Data Curation, **MPO**: Visualization, Writing - Review & Editing, **SB**: Writing - Review & Editing, **SH**: Writing - Original Draft, **SLH**: Supervision, Project administration, Funding acquisition, Writing - Review & Editing, **SM**: Funding acquisition, Writing - Review & Editing, **TM**: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, **ZW**: Writing - Original Draft.

References

[1] Chiara Cirelli and Giulio Tononi. Is sleep essential? *PLoS Biol*, 6(8):e216, 2008. doi: 10.1371/journal.pbio. 0060216.

[2] Serge Daan, DG Beersma, and Alexander A Borbély. Timing of human sleep: recovery process gated by a circadian pacemaker. *Am J Physiol Regul Integr Comp Physiol*, 246(2):R161–R183, 1984. doi: 10.1152/ajpregu. 1984.246.2.r161.

[3] Lulu Xie, Hongyi Kang, Qiwu Xu, Michael J. Chen, Yonghong Liao, Meenakshisundaram Thiyagarajan, John O'Donnell, Daniel J. Christensen, Charles Nicholson, Jeffrey J. Iliff, Takahiro Takano, Rashid Deane, and Maiken Nedergaard. Sleep Drives Metabolite Clearance from the Adult Brain. *Science*, 342(6156):373–377, October 2013. ISSN 0036-8075. doi: 10.1126/science.1241224.

[4] Emmanuel Mignot. Why we sleep: the temporal organization of recovery. *PloS Biol*, 6(4):e106, 2008. doi: 10.1371/journal.pbio.0060106.

[5] Wilfredo Blanco, Catia M. Pereira, Vinicius R. Cota, Annie C. Souza, César Rennó-Costa, Sharlene Santos, Gabriella Dias, Ana M. G. Guerreiro, Adriano B. L. Tort, Adrião D. Neto, and Sidarta Ribeiro. Synaptic Homeostasis and Restructuring across the Sleep-Wake Cycle. *PLoS Comput Biol*, 11(5):e1004241, May 2015. ISSN 1553-7358. doi: 10.1371/journal.pcbi.1004241.

[6] Le Shi, Si-Jing Chen, Meng-Ying Ma, Yan-Ping Bao, Ying Han, Yu-Mei Wang, Jie Shi, Michael V. Vitiello, and Lin Lu. Sleep disturbances increase the risk of dementia: A systematic review and meta-analysis. *Sleep Med Rev*, 40:4–16, August 2018. ISSN 1087-0792. doi: 10.1016/j.smrv.2017.06.010.

[7] Yo-El S Ju, Aleksandar Videnovic, and Bradley V Vaughn. Comorbid sleep disturbances in neurologic disorders. *Continuum (Minneap Minn)*, 23(4):1117–1131, 2017. doi: 10.1212/con.0000000000000501.

[8] Daniela Minecan, Alamelu Natarajan, Mary Marzec, and Beth Malow. Relationship of epileptic seizures to sleep stage and sleep depth. *Sleep*, 25(8):899–904, December 2002. ISSN 0161-8105. doi: 10.1093/sleep/25.8.56. URL `https://pubmed.ncbi.nlm.nih.gov/12489898`.

[9] David Nutt, Sue Wilson, and Louise Paterson. Sleep disorders as core symptoms of depression. *Dialogues Clin Neurosci*, 10(3):329, September 2008. doi: 10.31887/DCNS.2008.10.3/dnutt.

[10] G. Stores. Clinical diagnosis and misdiagnosis of sleep disorders. *J Neurol Neurosurg Psychiatry*, 78(12):1293, December 2007. doi: 10.1136/jnnp.2006.111179.

[11] J Zeitlhofer, P Anderer, S Obergottsberger, P Schimicek, S Lurger, E Marschnigg, B Saletu, and L Deecke. Topographic mapping of EEG during sleep. *Brain Topogr*, 6:123–129, 1993. doi: 10.1007/bf01191077.

[12] Conrad Iber and American Academy of Sleep Medicine. *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications*. American Academy of Sleep Medicine, Westchester, IL, 2007. OCLC: 438914652.

[13] Anthony Kales, Allan Rechtschaffen, Los Angeles University of California, and National Institute of Neurological Diseases and Blindness (U.S.). *A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects*. NIH publication. United States Government Printing Office, Washington, DC, 1968. doi: 10.1016/0013-4694(69)90021-2.

[14] Charles Bae and Nancy Foldvary-Schaefer. The use of sleep studies in neurological practice. *Semin Neurol*, 24 (3):237–248, September 2004. ISSN 0271-8235. doi: 10.1055/s-2004-835072.

[15] Axel Steiger and Mayumi Kimura. Wake and sleep EEG provide biomarkers in depression. *J Psychiatr Res*, 44 (4):242–252, March 2010. ISSN 1879-1379. doi: 10.1016/j.jpsychires.2009.08.013.

[16] Adam Wichniak, Aleksandra Wierzbicka, and Wojciech Jernajczyk. Sleep as a biomarker for depression. *Int Rev Psychiatry*, October 2013. URL https://www.tandfonline.com/doi/full/10.3109/09540261.2013.812067.

[17] José Haba-Rubio, Pedro Marques-Vidal, Daniela Andries, Nadia Tobback, Martin Preisig, Peter Vollenweider, Gérard Waeber, Gianina Luca, Mehdi Tafti, and Raphaël Heinzer. Objective sleep structure and cardiovascular risk factors in the general population: the HypnoLaus Study. *Sleep*, 38(3):391–400, March 2015. ISSN 1550-9109. doi: 10.5665/sleep.4496.

[18] Daphne Chylinski, Christian Berthomier, Eric Lambot, Sonia Frenette, Marie Brandewinder, Julie Carrier, Gilles Vandewalle, and Vincenzo Muto. Variability of sleep stage scoring in late midlife and early old age. *J Sleep Res*, 31(1):e13424, 2022.

[19] Magdy Younes, Jill Raneri, and Patrick Hanly. Staging sleep in polysomnograms: analysis of inter-scorer variability. *J Clin Sleep Med*, 12(6):885–894, 2016. doi: 10.5664/jcsm.5894.

[20] Richard Gao, Erik J Peterson, and Bradley Voytek. Inferring synaptic excitation/inhibition balance from field potentials. *NeuroImage*, 158:70–78, 2017. doi: 10.1016/j.neuroimage.2017.06.078.

[21] Kusum Thuwal, Arpan Banerjee, and Dipanjan Roy. Aperiodic and Periodic Components of Ongoing Oscillatory Brain Dynamics Link Distinct Functional Aspects of Cognition across Adult Lifespan. *eNeuro*, 8(5): ENEURO.0224–21.2021, September 2021. doi: 10.1523/eneuro.0224-21.2021.

[22] Guang Ouyang, Andrea Hildebrandt, Florian Schmitz, and Christoph S. Herrmann. Decomposing alpha and 1/f brain activities reveals their differential associations with cognitive processing speed. *NeuroImage*, 205:116304, January 2020. doi: 10.1016/j.neuroimage.2019.116304.

[23] Gert Pfurtscheller and Christa Neuper. *Movement and ERD/ERS*, pages 191–206. Springer US, 2003. doi: 10.1007/978-1-4615-0189-3_12.

[24] Thomas E Scammell, Elda Arrigoni, and Jonathan O Lipton. Neural circuitry of wakefulness and sleep. *Neuron*, 93(4):747–765, 2017. doi: 10.1016/j.neuron.2017.01.014.

[25] Eric R. Kandel, James H. Schwartz, Thomas M. Jessell, Steven A. Siegelbaum, and A. J. Hudspeth. *Principles of Neural Science, Fifth Edition*. McGraw Hill Professional, October 2012. ISBN 978-0-07-181001-2. Google-Books-ID: Z2yVUTnlIQsC.

[26] Jae Myeong Kang, Seo-Eun Cho, Jong Youn Moon, Soo In Kim, Jong Won Kim, and Seung-Gul Kang. Difference in spectral power density of sleep electroencephalography between individuals without insomnia and frequent hypnotic users with insomnia complaints. *Sci Rep*, 12(1):2117, 2022. doi: 10.1038/s41598-022-05378-6.

[27] Yuval Nir, Marcello Massimini, Melanie Boly, and Giulio Tononi. Sleep and Consciousness. In Andrea Eugenio Cavanna, Andrea Nani, Hal Blumenfeld, and Steven Laureys, editors, *Neuroimaging of Consciousness*, pages 133–182. Springer, Berlin, Heidelberg, 2013. ISBN 978-3-642-37580-4. doi: 10.1007/978-3-642-37580-4_9. URL https://doi.org/10.1007/978-3-642-37580-4%5F9.

[28] Mircea Steriade, David A. McCormick, and Terrence J. Sejnowski. Thalamocortical Oscillations in the Sleeping and Aroused Brain. *Science*, 262(5134):679–685, oct 29 1993. doi: 10.1126/science.8235588. URL http://dx.doi.org/10.1126/SCIENCE.8235588.

[29] David A. McCormick and Thierry Bal. SLEEP AND AROUSAL: Thalamocortical Mechanisms. *Annu Rev Neurosci*, 20(1):185–215, March 1997. doi: 10.1146/annurev.neuro.20.1.185. URL http://dx.doi.org/10.1146/ANNUREV.NEURO.20.1.185.

[30] PA Robinson, CJ Rennie, JJ Wright, H Bahramali, E Gordon, and DL Rowe. Prediction of electroencephalographic spectra from neurophysiology. *Phys Rev E*, 63(2):021903, 2001. doi: 10.1103/physreve.63.021903.

[31] Igor Timofeev, François Grenier, and Mircea Steriade. Disfacilitation and active inhibition in the neocortex during the natural sleep-wake cycle: An intracellular study. *Proc. Natl. Acad. Sci. U.S.A.*, 98(4):1924–1929, February 2001. doi: 10.1073/pnas.98.4.1924.

[32] Sean Hill and Giulio Tononi. Modeling sleep and wakefulness in the thalamocortical system. *J Neurophysiol*, 93(3):1671–1698, 2005. doi: 10.1152/jn.00915.2004.

[33] Giulio Tononi and Chiara Cirelli. Sleep function and synaptic homeostasis. *Sleep Med Rev*, 10(1):49–62, 2006. doi: 10.1016/j.smrv.2005.05.002.

[34] Antoine R. Adamantidis, Carolina Gutierrez Herrera, and Thomas C. Gent. Oscillating circuitries in the sleeping brain. *Nat Rev Neurosci*, 20(12):746–762, December 2019. ISSN 1471-0048. doi: 10.1038/s41583-019-0223-4.

[35] Tadao Hori. Spatiotemporal Changes of EEG Activity During Waking-Sleeping Transition Period. *Int J Neurosci*, January 1985. ISSN 1129-3354. doi: 10.3109/00207458509149139. URL https://www.tandfonline.com/doi/abs/10.3109/00207458509149139.

[36] Bence Schneider, Orsolya Szalárdy, Péter P. Ujma, Péter Simor, Ferenc Gombos, Ilona Kovács, Martin Dresler, and Róbert Bódizs. Scale-free and oscillatory spectral measures of sleep stages in humans. *Front Neuroinf*, 16:989262, October 2022. ISSN 1662-5196. doi: 10.3389/fninf.2022.989262.

[37] Leslie D Claar, Irene Rembado, Jacqulyn R Kuyat, Simone Russo, Lydia C Marks, Shawn R Olsen, and Christof Koch. Cortico-thalamo-cortical interactions modulate electrically evoked EEG responses in mice. *eLife*, 12: RP84630, jun 2023. ISSN 2050-084X. doi: 10.7554/eLife.84630. URL `https://doi.org/10.7554/eLife.84630`.

[38] Steve K. Esser, Sean Hill, and Giulio Tononi. Breakdown of Effective Connectivity During Slow Wave Sleep: Investigating the Mechanism Underlying a Cortical Gate Using Large-Scale Modeling. *J Neurophysiol*, October 2009. doi: 10.1152/jn.00059.2009. URL `https://journals.physiology.org/doi/full/10.1152/jn.00059.2009`.

[39] Thien Thanh Dang-Vu, Manuel Schabus, Martin Desseilles, Geneviève Albouy, Mélanie Boly, Annabelle Darsaud, Steffen Gais, Géraldine Rauchs, Virginie Sterpenich, Gilles Vandewalle, Julie Carrier, Gustave Moonen, Evelyne Balteau, Christian Degueldre, André Luxen, Christophe Phillips, and Pierre Maquet. Spontaneous neural activity during human slow wave sleep. *Proc Natl Acad Sci*, 105(39):15160–15165, September 2008. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.0801819105. URL `https://pnas.org/doi/full/10.1073/pnas.0801819105`.

[40] Yuval Nir, Richard J. Staba, Thomas Andrillon, Vladyslav V. Vyazovskiy, Chiara Cirelli, Itzhak Fried, and Giulio Tononi. Regional Slow Waves and Spindles in Human Sleep. *Neuron*, 70(1):153–169, April 2011. ISSN 0896-6273. doi: 10.1016/j.neuron.2011.02.043.

[41] Marcello Massimini, Fabio Ferrarelli, Steve K. Esser, Brady A. Riedner, Reto Huber, Michael Murphy, Michael J. Peterson, and Giulio Tononi. Triggering sleep slow waves by transcranial magnetic stimulation. *Proc Natl Acad Sci*, 104(20):8496–8501, May 2007. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.0702495104. URL `https://pnas.org/doi/full/10.1073/pnas.0702495104`.

[42] Yina Wei, Giri P. Krishnan, Lisa Marshall, Thomas Martinetz, and Maxim Bazhenov. Stimulation Augments Spike Sequence Replay and Memory Consolidation during Slow-Wave Sleep. *J Neurosci*, 40(4):811–824, January 2020. ISSN 0270-6474. doi: 10.1523/JNEUROSCI.1427-19.2019.

[43] PA Robinson, CJ Rennie, and DL Rowe. Dynamics of large-scale brain activity in normal arousal states and epileptic seizures. *Phys Rev E*, 65(4):041924, 2002. doi: 10.1103/physreve.65.041924.

[44] Christopher J Rennie, Peter A Robinson, and James J Wright. Unified neurophysical model of EEG spectra and evoked potentials. *Biol Cybern*, 86(6):457–471, 2002. doi: 10.1007/s00422-002-0310-9.

[45] PA Robinson, CJ Rennie, DL Rowe, SC O'Connor, JJ Wright, E Gordon, and RW Whitehouse. Neurophysical modeling of brain dynamics. *Neuropsychopharmacology*, 28(1):S74–S79, 2003. doi: 10.1038/sj.npp.1300143.

[46] Tara Babaie-Janvier and Peter A Robinson. Neural field theory of evoked response potentials with attentional gain dynamics. *Front Hum Neurosci*, 14:293, 2020. doi: 10.3389/fnhum.2020.00293.

[47] Peter A Robinson, Natasha C Gabay, and Tara Babaie-Janvier. Neural Field Theory of Evoked Response Sequences and Mismatch Negativity With Adaptation. *Front Hum Neurosci*, 15:655505, 2021. doi: 10.3389/fnhum.2021.655505.

[48] PA Robinson, RW Whitehouse, and CJ Rennie. Nonuniform corticothalamic continuum model of electroencephalographic spectra with application to split-alpha peaks. *Phys Rev E*, 68(2):021922, 2003. doi: 10.1103/physreve.68.021922.

[49] Romesh G Abeysuriya, Chris J Rennie, and Peter A Robinson. Prediction and verification of nonlinear sleep spindle harmonic oscillations. *J Theor Biol*, 344:70–77, 2014. doi: 10.1016/j.jtbi.2013.11.013.

[50] RG Abeysuriya, CJ Rennie, PA Robinson, and JW Kim. Experimental observation of a theoretically predicted nonlinear sleep spindle harmonic in human EEG. *Clin Neurophysiol*, 125(10):2016–2023, 2014. doi: 10.1016/j.clinph.2014.01.025.

[51] Romesh G. Abeysuriya, Christopher J. Rennie, and Peter A. Robinson. Physiologically based arousal state estimation and dynamics. *J Neurosci Methods*, 253, 06 2015. doi: 10.1016/j.jneumeth.2015.06.002.

[52] Maurice Abou Jaoude, Aravind Ravi, Jiansheng Niu, Hubert Banville, Nicolas Florez Torres, and Christopher Aimone. Automated Sleep Staging on Wearable EEG Enables Sleep Analysis at Scale. In *2023 11th International IEEE/EMBS Conference on Neural Engineering (NER)*, pages 1–4. IEEE, 2023. doi: 10.1109/NER52421.2023.10123829. URL https://ieeexplore.ieee.org/abstract/document/10123829.

[53] Pierrick J Arnal, Valentin Thorey, Eden Debellemaniere, Michael E Ballard, Albert Bou Hernandez, Antoine Guillot, Hugo Jourde, Mason Harris, Mathias Guillard, Pascal Van Beers, et al. The Dreem Headband compared to polysomnography for electroencephalographic signal acquisition and sleep staging. *Sleep*, 43(11):zsaa097, 2020. doi: 10.1093/sleep/zsaa097.

[54] Magdy Younes, Marc Soiferman, Wayne Thompson, and Eleni Giannouli. Performance of a new portable wireless sleep monitor. *J Clin Sleep Med*, 13(2):245–258, February 2017. ISSN 1550-9397. doi: 10.5664/jcsm.6456.

[55] Noralie Krepel, Tommy Egtberts, Emma Touré-Cuq, Pierre Bouny, and Martijn Arns. Evaluation of the URGOnight Tele-neurofeedback Device: An Open-label Feasibility Study with Follow-up. *Appl Psychophysiol Biofeedback*, 47(1):43–51, March 2022. ISSN 1573-3270. doi: 10.1007/s10484-021-09525-z.

[56] Bob Kemp, Aeilko H Zwinderman, Bert Tuk, Hilbert AC Kamphuisen, and Josefien JL Oberye. Analysis of a sleep-dependent neuronal feedback loop: the slow-wave microcontinuity of the EEG. *IEEE Trans Biomed Eng*, 47(9):1185–1194, 2000. doi: 10.1109/10.867928.

[57] Bob Kemp, A Zwinderman, B Tuk, H Kamphuisen, and J Oberyé. Sleep-EDF Database Expanded, 2018. URL https://www.physionet.org/content/sleep-edfx/1.0.0/.

[58] Ary L Goldberger, Luis AN Amaral, Leon Glass, Jeffrey M Hausdorff, Plamen Ch Ivanov, Roger G Mark, Joseph E Mietus, George B Moody, Chung-Kang Peng, and H Eugene Stanley. PhysioBank, PhysioToolkit,

and PhysioNet: components of a new research resource for complex physiologic signals. *Circulation*, 101(23): e215–e220, 2000. doi: 10.1161/01.cir.101.23.e215.

[59] Antoine Guillot, Fabien Sauvet, Emmanuel H During, and Valentin Thorey. Dreem open datasets: Multi-scored sleep datasets to compare human and automated sleep staging. *IEEE Trans Neural Syst Rehabilitation Eng*, 28 (9):1955–1965, 2020. doi: 10.1109/tnsre.2020.3011181.

[60] Valentin Thorey, Simon Kerns, and Antoine Guillot. Dreem-Organization/dreem-learning-open, February 2021. URL `https://github.com/Dreem-Organization/dreem-learning-open`. original-date: 2019-11-05T16:44:38Z.

[61] Ning Mei, Michael D Grossberg, Kenneth Ng, Karen T Navarro, and Timothy M Ellmore. A high-density scalp EEG dataset acquired during brief naps after a visual working memory task. *Data Brief*, 18:1513–1519, 2018. doi: 10.1016/j.dib.2018.04.073.

[62] Timothy Ellmore, Ning Mei, Kenneth Ng, and Karen Tatiana Navarro. Nap EEG, July 2016. URL `https://osf.io/chav7/`. Publisher: OSF.

[63] Guo-Qiang Zhang, Licong Cui, Remo Mueller, Shiqiang Tao, Matthew Kim, Michael Rueschman, Sara Mariani, Daniel Mobley, and Susan Redline. The National Sleep Research Resource: towards a sleep data commons. *J Am Med Inform Assoc*, 25(10):1351–1358, 2018. doi: 10.1145/3233547.3233725.

[64] Terry Young, Mari Palta, Jerome Dempsey, Paul E Peppard, F Javier Nieto, and K Mae Hla. Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study. *WMJ*, 108(5):246, 2009.

[65] Christoph Guger, Gunther Krausz, Brendan Z. Allison, and Guenter Edlinger. Comparison of Dry and Gel Based Electrodes for P300 Brain–Computer Interfaces. *Front Neurosci*, 6:25155, May 2012. ISSN 1662-453X. doi: 10.3389/fnins.2012.00060.

[66] Alexandre Gramfort, Martin Luessi, Eric Larson, Denis A Engemann, Daniel Strohmeier, Christian Brodbeck, Roman Goj, Mainak Jas, Teon Brooks, Lauri Parkkonen, et al. MEG and EEG data analysis with MNE-Python. *Front Neurosci*, 7:267, 2013. doi: 10.3389/fnins.2013.00267.

[67] Peter Welch. The use of fast Fourier transform for the estimation of power spectra: a method based on time averaging over short, modified periodograms. *IEEE Trans Audio Electroacoust*, 15(2):70–73, 1967. doi: 10.1109/tau.1967.1161901.

[68] Jie Cui and Willy Wong. Optimal window length in the windowed adaptive chirplet analysis of visual evoked potentials. In *2006 International Conference of the IEEE Engineering in Medicine and Biology Society*, pages 4580–4583. IEEE, 2006.

[69] Thomas Donoghue, Matar Haller, Erik J Peterson, Paroma Varma, Priyadarshini Sebastian, Richard Gao, Torben Noto, Antonio H Lara, Joni D Wallis, Robert T Knight, et al. Parameterizing neural power spectra into periodic and aperiodic components. *Nat Neuro*, 23(12):1655–1665, 2020. doi: 10.1038/s41593-020-00744-x.

[70] Raphael Vallat and Matthew P. Walker. An open-source, high-performance tool for automated sleep staging. *eLife*, October 2021. doi: 10.7554/eLife.70092.

[71] Ramesh Srinivasan, Don M Tucker, and Michael Murias. Estimating the spatial Nyquist of the human EEG. *Beh Res Meth Instr Comp*, 30(1):8–19, 1998. doi: 10.3758/bf03209412.

[72] Suresh Muthukumaraswamy. High-frequency brain activity and muscle artifacts in MEG/EEG: A review and recommendations. *Front Hum Neurosci*, 7:47068, April 2013. ISSN 1662-5161. doi: 10.3389/fnhum.2013.00138.

[73] Federico Barban, Michela Chiappalone, Gaia Bonassi, Dante Mantini, and Marianna Semprini. Yet another artefact rejection study: an exploration of cleaning methods for biological and neuromodulatory noise. *J Neural Eng*, 18(4):0460c2, August 2021. ISSN 1741-2552. doi: 10.1088/1741-2552/ac01fe.

[74] Donald L Rowe, Peter A Robinson, and Christopher J Rennie. Estimation of neurophysiological parameters from the waking EEG using a biophysical model of brain dynamics. *Journal of theoretical biology*, 231(3):413–433, 2004. doi: 10.1016/j.jtbi.2004.07.004.

[75] RG Abeysuriya and PA Robinson. Real-time automated EEG tracking of brain states using neural field theory. *J Neurosci Methods*, 258:28–45, 2016. doi: 10.1016/j.jneumeth.2015.09.026.

[76] Romesh Abeysuriya. BrainDynamicsUSYD/braintrak, January 2019. URL https://github.com/BrainDynamicsUSYD/braintrak. original-date: 2015-04-01T22:37:43Z.

[77] Ushtar Amin, Fábio A. Nascimento, Ioannis Karakis, Donald Schomer, and Selim R. Benbadis. Normal variants and artifacts: Importance in EEG interpretation. *Epileptic Disord*, 25(5):591–648, jul 2023.

[78] Jose Antonio Urigüen and Begoña Garcia-Zapirain. EEG artifact removal—state-of-the-art and guidelines. 12 (3):031001, April 2015. ISSN 1741-2552. doi: 10.1088/1741-2560/12/3/031001.

[79] Scott I Vrieze. Model selection and psychological theory: a discussion of the differences between the Akaike information criterion (AIC) and the Bayesian information criterion (BIC). *Psychol Methods*, 17(2):228, 2012. doi: 10.1037/a0027127.

[80] Mary A. Carskadon and William C. Dement. Normal Human Sleep: An Overview. In *Principles and Practice of Sleep Medicine (Sixth Edition)*, pages 15–24.e3. Elsevier, Walthm, MA, USA, January 2017. ISBN 978-0-323-24288-2. doi: 10.1016/B978-0-323-24288-2.00002-7.

[81] Sophie Desjardins, Sylvie Lapierre, Carol Hudon, and Alain Desgagné. Factors involved in sleep efficiency: a population-based study of community-dwelling elderly persons. *Sleep*, 42(5):zsz038, 2019. doi: 10.1093/sleep/zsz038.

[82] Fabrizio Lombardi, Hans J Herrmann, and Lucilla de Arcangelis. Balance of excitation and inhibition determines 1/f power spectrum in neuronal networks. *Chaos*, 27(4):047402, 2017. doi: 10.1063/1.4979043.

[83] Winston Haynes. *Bonferroni Correction*, pages 154–154. Springer New York, New York, NY, 2013. ISBN 978-1-4419-9863-7. doi: 10.1007/978-1-4419-9863-7_1213. URL https://doi.org/10.1007/978-1-4419-9863-7_1213.

[84] Yoav Benjamini and Yosef Hochberg. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Series B Stat Methodol*, 57(1):289–300, 1995. ISSN 00359246. doi: 10.1111/j.2517-6161.1995.tb02031.x. URL http://www.jstor.org/stable/2346101.

[85] Winston Haynes. *Bonferroni Correction*, pages 154–154. Springer New York, New York, NY, 2013. ISBN 978-1-4419-9863-7. doi: 10.1007/978-1-4419-9863-7_1213. URL https://doi.org/10.1007/978-1-4419-9863-7_1213.

[86] Qiang Li, Jiang-Ling Song, Si-Hui Li, M. Brandon Westover, and Rui Zhang. Effects of Cholinergic Neuromodulation on Thalamocortical Rhythms During NREM Sleep: A Model Study. *Front Comp Neurosci*, 13, 2020. ISSN 1662-5188.

[87] David A. McCormick. Chapter 36: Actions of acetylcholine in the cerebral cortex and thalamus and implications for function. In *Progress in Brain Research*, volume 98, pages 303–308. Elsevier, Walthm, MA, USA, January 1993. doi: 10.1016/S0079-6123(08)62412-7.

[88] José J. Rodríguez, Harun N. Noristani, Walter B. Hoover, Stephanie B. Linley, and Robert P. Vertes. Serotonergic projections and serotonin receptor expression in the reticular nucleus of the thalamus in the rat. *Synapse*, 65(9): 919–928, September 2011. ISSN 1098-2396. doi: 10.1002/syn.20920.

[89] Margaret N. Shouse. Sleep deprivation increases thalamocortical excitability in the somatomotor pathway, especially during seizure-prone sleep or awakening states in feline seizure models. *Exp. Neurol.*, 99(3):664–677, March 1988. ISSN 0014-4886. doi: 10.1016/0014-4886(88)90183-5.

[90] Yongcong Shao, Lubin Wang, Enmao Ye, Xiao Jin, Wei Ni, Yue Yang, Bo Wen, Dewen Hu, and Zheng Yang. Decreased Thalamocortical Functional Connectivity after 36 Hours of Total Sleep Deprivation: Evidence from Resting State fMRI. *PLoS One*, 8(10), 2013. doi: 10.1371/journal.pone.0078830.

[91] William WK Zung. A self-rating depression scale. *Arch Gen Psychiatry*, 12(1):63–70, 1965. doi: 10.1037/t04095-000.

[92] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay. Scikit-learn: Machine Learning in Python. *J Mach Learn Res*, 12:2825–2830, 2011.

[93] RW Guillery, SL Feig, and DA Lozsadi. Paying attention to the thalamic reticular nucleus. *Trends Neurosci*, 21 (1):28–32, 1998. doi: 10.1016/s0166-2236(97)01157-0.

[94] Brady A. Riedner, Vladyslav V. Vyazovskiy, Reto Huber, Marcello Massimini, Steve Esser, Michael Murphy, and Giulio Tononi. Sleep Homeostasis and Cortical Synchronization: III. A High-Density EEG Study of Sleep Slow Waves in Humans. *Sleep*, 30(12):1643–1657, December 2007. ISSN 0161-8105. doi: 10.1093/sleep/30.12.1643.

[95] Vladyslav V. Vyazovskiy, Brady A. Riedner, Chiara Cirelli, and Giulio Tononi. Sleep Homeostasis and Cortical Synchronization: II. A Local Field Potential Study of Sleep Slow Waves in the Rat. *Sleep*, 30(12):1631–1642, December 2007. ISSN 0161-8105. doi: 10.1093/sleep/30.12.1631.

[96] Nazanin Biabani, Adam Birdseye, Sean Higgins, Alessio Delogu, Jan Rosenzweig, Zoran Cvetkovic, Alexander Nesbitt, Panagis Drakatos, Joerg Steier, Veena Kumari, David O'Regan, and Ivana Rosenzweig. The neurophysiologic landscape of the sleep onset: a systematic review. *Journal of Thoracic Disease*, 15(8), August 2023. ISSN 2077-6624. doi: 10.21037/jtd-23-325.

[97] Justus T. C. Schwabedal, Maik Riedl, Thomas Penzel, and Niels Wessel. Alpha-wave frequency characteristics in health and insomnia during sleep. *J Sleep Res*, 25(3):278–286, June 2016. ISSN 1365-2869. doi: 10.1111/jsr. 12372.

[98] Jintao Wu, Qianxiang Zhou, Jiaxuan Li, Yang Chen, Shuyu Shao, and Yi Xiao. Decreased resting-state alpha-band activation and functional connectivity after sleep deprivation. *Sci Rep*, 11(1):1–10, January 2021. ISSN 2045-2322. doi: 10.1038/s41598-020-79816-8.

[99] Pablo Fuentealba, Igor Timofeev, and Mircea Steriade. Prolonged hyperpolarizing potentials precede spindle oscillations in the thalamic reticular nucleus. *Proc Natl Acad Sci*, 101(26):9816–9821, 2004. doi: 10.1073/pnas. 0402761101.

[100] Elisabetta Iavarone, Jane Simko, Ying Shi, Marine Bertschy, María García-Amado, Polina Litvak, Anna-Kristin Kaufmann, Christian O'Reilly, Oren Amsalem, Marwan Abdellah, et al. Thalamic control of sensory processing and spindles in a biophysical somatosensory thalamoreticular circuit model of wakefulness and sleep. *Cell Rep*, 42(3), 2023. doi: 10.1016/j.celrep.2023.112200.

[101] James C. Pang, Kevin M. Aquino, Marianne Oldehinkel, Peter A. Robinson, Ben D. Fulcher, Michael Breakspear, and Alex Fornito. Geometric constraints on human brain function. *Nature*, 618(7965):566–574, June 2023. ISSN 1476-4687. doi: 10.1038/s41586-023-06098-1.

[102] Jie Cui and Willy Wong. Investigation of short-term changes in visual evoked potentials with windowed adaptive chirplet transform. *IEEE Transactions on Biomedical Engineering*, 55(4):1449–1454, 2008.

[103] Steve Mann and Simon Haykin. Adaptive chirplet: an adaptive generalized wavelet-like transform. In *Adaptive Signal Processing*, volume 1565, pages 402–413. SPIE, 1991.

[104] Gerald B Folland. *Harmonic analysis in phase space*. Number 122. Princeton university press, 1989.

[105] Diego Z Carvalho, Günther JL Gerhardt, Guilherme Dellagustin, Emerson L de Santa-Helena, Ney Lemke, Alan Z Segal, and Suzana V Schönwald. Loss of sleep spindle frequency deceleration in obstructive sleep apnea. *Clinical Neurophysiology*, 125(2):306–312, 2014.

[106] Diego Carvalho, Suzana Schönwald, Guilherme Dellagustin, Emerson de Santa-Helena, Ney Lemke, and Gunther Gerhardt. Sleep spindle frequency modulation (chirp) in obstructive sleep apnea (p04. 025), 2012.

[107] Suzana V Schönwald, Diego Z Carvalho, Guilherme Dellagustin, Emerson L de Santa-Helena, and Günther JL Gerhardt. Quantifying chirp in sleep spindles. *Journal of Neuroscience Methods*, 197(1):158–164, 2011.

[108] Rafael Toledo Fernandes de Souza, Günther Johannes Lewczuk Gerhardt, Suzana Veiga Schönwald, José Luiz Rybarczyk-Filho, and Ney Lemke. Synchronization and propagation of global sleep spindles. *PLoS One*, 11(3): e0151369, 2016.

List of Figures