The Interplay of Ecological Opportunity and Adult Foraging Strategies

in Juvenile Sparrow Individual Specialization

Natalia Ricote[1], Constanza Weinberger[1], Natalia Ramírez-Otarola[2], Sara Bustamante[1,3],

María Lucía Málaga[1,3], Gonzalo Barceló[4], Pablo Sabat[3,5,7], Seth D. Newsome[6] and Karin

Maldonado[1*]

[1]Departamento de Ciencias, Facultad de Artes Liberales, Universidad Adolfo Ibáñez,

Santiago, Chile

[2]Escuela de Medicina Veterinaria, Facultad de Medicina y Ciencias de la Salud,

Universidad Mayor, Camino La Pirámide 5750, Huechuraba, Santiago, Chile.

[3]Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile,

Santiago, Chile

[4]Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, 1630

Linden Dr., Madison, Wisconsin 53706, USA

[5]Center of Applied Ecology and Sustainability (CAPES), Santiago, Chile

[6]Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA

[7] Millenium Nucleus of Patagonian Limit of Life (LiLi)

Correspondence:

corresponding author:

karin.maldonado@uai.cl

Abstract

Temporal variation in resource availability, amplified by global change, may have strong

impacts on species breeding at temperate and high latitudes that cue their reproduction to

exploit seasonal resource pulses. This study examines how resource availability

(ecological opportunity) and parental care influence niche partitioning between and within

age classes in the rufous-collared sparrow, which provides extensive parental care. We

hypothesized juveniles would exhibit narrower niches focused on high-quality resources

compared to adults, regardless of ecological opportunity. We used stable isotope analysis

to quantify individual and population niches in juveniles and adults across the breeding

season in two cohorts experiencing contrasting resource landscapes. Contrary to our initial

hypothesis, juveniles exhibited greater individual specialization and smaller total niche

widths during periods when resource availability was high in comparison to periods of low

ecological opportunity. Interestingly, niche characteristics of adults remained stable across

seasons despite a shift in trophic level, highlighting their potential role in providing a

consistent diet for their young. These findings reveal a dynamic interplay between

ecological opportunity, parental care, and individual specialization, with important

implications for understanding population resilience under variable resource scenarios.

The study also suggests that adult sparrows adapt their provisioning strategies based on

ecological opportunity, potentially buffering offspring from environmental fluctuations.

Understanding age-specific responses to resource variation is crucial for predicting

species responses to ecological conditions, particularly in regions like central Chile where

seasonal resource limitation is expected to become more variable in response to climate

change.

Introduction

Global change is expected to increase variation in the availability of resources (Jia et al.,

2019), and within-population variability in resource use may increase its resilience to

environmental change (Ramos et al., 2020; Riverón et al., 2021). Trophic niche differences

between age classes have been a key focus of population-level research, while more

recently, individual specialization, where individuals utilize narrower resource ranges than

their population, has gained interest (Costa-Pereira et al., 2018). These kinds of variation

may be driven by the same mechanisms that underlie resource partitioning among species

to reduce competition (Keast, 1977; Maiorana, 1978; Svanbäck and Bolnick, 2005). For

example, individual specialization can occur within a single age class, reducing

intraspecific competition and enhancing a population's resilience to fluctuations in resource

availability. Furthermore, the diversification of ecological roles promoted by individual

niche differentiation can contribute to ecosystem stability (Araújo et al., 2011; Bolnick et

al., 2011). Additionally, individual specialization can mitigate the costs of transitioning

between different resources due to functional trade-offs in which generalist individuals

experience decreased exploitation efficiency when using a wide range of resources.

Consequently, high levels of individual specialization would enable individuals to allocate

energy towards other essential functions that enhance fitness such as reproduction

(Bolnick et al., 2003; Araújo et al., 2011; Maldonado et al., 2019).

In general, it can often be assumed that parents feed their offspring a diet as diverse as

what they consume, leading to similar resource use and no niche partitioning between adults

and juveniles. In birds, parental decisions about how to allocate food to their nestlings are

strongly influenced by nestling behavior, parental sex, life history traits, and the nutritional

needs of their offspring, which require high-quality food to rapidly grow and enhance their

probability of survival (Alonso et al., 2012; Raya Rey et al., 2012; Beaulieu and Sockman,

2014). It has been suggested that protein intake is higher at early ages due to the

predominance of anabolic reactions caused by the formation of new tissues (Kyriazakis and

Oldham, 1993). This is consistent with studies on the overlap of population niches among

different age classes, which show that juveniles occupy higher trophic levels compared to

adults (Hobson, 1993; Hammerschlag-Peyer et al., 2011). Several studies have reported

niche partitioning between adults and juveniles, as well as differences in individual

specialization among age classes due to disparities in foraging abilities, morphology (*i.e.*

body mass, bill length), or social status (Durell, 2003; Hall et al., 2021). (Sword and Dopman,

1999; Frédérich et al., 2010; Vander Zanden et al., 2013; Hall et al., 2021). For example, in

western sandpipers (*Calidris mauri*) adults and juveniles differed in individual specialization

levels because juveniles have less experience in handling prey, shorter bills, and lower body

mass (Hall et al., 2021). A common feature in these studies, however, is that they report

differences between adults and juveniles in species that have little or no parental care or in

juveniles that can forage independently (Durell, 2003; Hall et al., 2021). Moreover, these

studies have not considered resource variation as a potential explanatory variable, despite

its well-documented influence on both niche breadth and niche partitioning (Finke and

Snyder, 2008; Araújo et al., 2011; Stephens et al., 2019).

Our hypothesis is that ecological opportunity modulates niche partitioning in both adults

and juveniles through the influence of seasonal resource availability on parental behavior

(prey selection). In species with significant parental care, we expect parents to select high-

quality food to ensure their brood's survival even if it results in more competition with other

adults during periods of low food availability (Beaulieu and Sockman, 2014). Thus, juveniles

should exhibit narrower niches focused on high-quality prey occupying higher trophic levels

that contain high protein contents, and lower degrees of individual specialization compared

to adults independent of environmental shifts in resource availability. On the contrary, adults

are expected to have higher levels of individual specialization during conditions of high

resource availability when ecological opportunity is enhanced (Hall et al., 2021). No studies

have examined how parental care, specifically the provisioning of food from parents to

offspring, affects niche partitioning both between and within age classes under contrasting

levels of ecological opportunity.

To assess these hypotheses, we studied the rufous-collared sparrow (*Zonotrichia*

capensis) that provides parental care while chicks are in the nest and during the post-

fledging period as juveniles learn to forage independently (Miller and Miller, 1968; Tubaro,

1990). To compare niche partitioning between and within age classes, we exploited the

species' capability to generate multiple broods within a single breeding season, allowing for

two distinct cohorts of nestlings and adults to experience contrasting ecological opportunity.

We used isotope-based estimates of dietary variation, an effective approach to quantify

individual specialization via estimates of individual and population isotopic variance by

analyzing consumer tissues that reflect diet over different timescales (Matich et al., 2011;

Hanson et al., 2015; Bond et al., 2016; Maldonado et al., 2017). Nitrogen isotope (δ^{15}N)

values have been widely used to characterize trophic level and food-chain length (Post,

2002) and here we measured δ^{15}N values of blood plasma, red blood cells, and claw tips

collected from each individual as a proxy of dietary variation to quantify individual and total

trophic niches of juveniles and adults throughout the breeding season (Bearhop et al., 2004;

Vander Zanden et al., 2010).

Materials and Methods

Fieldwork and Tissue Collection. We captured rufous-collared sparrows (*Zonotrichia*

capensis) with mist nets in Quebrada de la Plata (33°31'S 70°50'W) in central Chile in

spring (November 2020) and summer (January 2021). We categorized captured

individuals into adults (N_{spring} =16, N_{summer}= 20) or juveniles (fledglings; N_{spring}= 12,

N_{summer}=8) according to plumage characteristics described in (Miller and Miller, 1968). The

study site is characterized by a Mediterranean climate with cool wet winters and relatively

hot dry summers (Di Castri and Hajek, 1976). In addition, at this site the availability of

insects and plants varies seasonally being highest in spring and lowest in winter; summer

values also are lower than in spring (Lopez-Calleja, 1995).

Parental care in this species extends from hatching until complete independence at 40

days old, with both parents actively feeding the chicks. Fledglings start self-feeding at 18

days old and achieve independence by four weeks, although they may still receive food

from parents until day 40 (Miller and Miller, 1968; Tubaro, 1990). Individuals captured

during November in spring would correspond to late September broods, while those

captured in late January (summer) would correspond to early December broods, ensuring

that both broods were raised under markedly different ecological conditions.

Immediately after capture, individuals were placed in a bird-holding bag (Ecotone,

Poland), weighed to the nearest 0.1 g on a digital scale, and a ~50µl blood sample was

collected from the brachial vein using heparinized microhematocrit capillary tubes. We

then sampled claw tips, which were placed in an Eppendorf tube. All samples were stored

frozen in liquid nitrogen prior to preparation for stable isotope analysis. All tissue samples

were dried at 80°C, homogenized, and ~0.5–0.6 mg of each sample was loaded into tin

capsules for stable isotope analysis.

Resource Availability. The rufous-collared sparrow is an omnivorous species that routinely

consumes both plants and insects (Lopez-Calleja, 1995). We first examined seasonal

variation in available resources in our study site. To measure plant availability, we

established five permanent 1m^2 quadrats distributed across the study area in which we

identified cover percentage of living herbaceous species during both seasons: mid-

September and mid-November for spring and mid-December and mid-January for

summer. We also collected leaf and seed samples of 12 plant species during the spring;

seed samples of 6 species were collected during the summer since live plant cover was

absent in our study quadrats. This collection allowed us to characterize the (baseline) δ^{15}N

composition required to estimate relative trophic level (see below).

We sampled terrestrial arthropods with Barber pitfall traps filled with water and

biodegradable detergent (Southwick, 1978) placed every 1 m along five linear transects of

10 m distributed across the study area and collected after 24 h (Jaksic and Lazo, 1999).

We counted and identified the collected arthropods to the lowest taxonomic resolution in

the laboratory.

We measured the density of arthropods for each transect and sample day by

quantifying the number of individuals collected divided by the number of Barber pitfall

traps. Density of plant resources was measured by the total percent cover of living plants

at each sample quadrant. We then estimated the mean density of each assemblage for

each season and its confidence interval by bootstrap procedure with 10,000 iterations.

Resource diversity was estimated by quantifying the Hill numbers for the plant and

arthropod assemblages (Chao et al., 2014). Seasonal species richness and Shannon

diversity index were calculated by using sample-size-based rarefaction and extrapolation

estimation of the Hill numbers of order q equal to 0 and 1, respectively. The estimates

were based on species abundance, measured as the percent cover of living plant species

and number of individuals within each arthropod morphotype (Chao et al., 2014; Hseih et

al., 2019). All estimations were performed with the package iNEXT (Hseih et al., 2019).

Stable Isotope Analysis. We measured δ^{15}N values in blood plasma, red blood cells, and

claw tips of juvenile and adult individuals to quantify niche characteristics in seasons

associated with high (spring) and low (summer) resource availability. Although many

tissues can be compared to construct a time series of isotopic variation, blood plasma, red

blood cells, and claws can be easily collected in the field without sacrificing animals

(Podlesak et al., 2005; Carleton et al., 2008). δ^{15}N values of claw tips integrate dietary

inputs over ~6 weeks prior to sample collection, while blood plasma and red blood cells

integrate diet over ~1-2 and ~3-4 weeks prior to collection, respectively (Hobson and

Clark, 1992; Carleton et al., 2008; Hahn et al., 2014). Young birds initiate the post-juvenile

molt either at the time of independence or within about 10 days thereafter (Miller and

Miller, 1968). Therefore, fledglings had not achieved total independence or had only

recently done so at the time of capture. This implies that the variance in diet composition

based on analysis of these three tissues, with the most recent one reflecting diet

integrated over 7 days prior to capture, represents the diet of juvenile individuals during

the parental care period when they had not yet achieved foraging independence.

Nitrogen isotope (δ^{15}N) values of sparrow tissues were measured on a Costech

4010 elemental analyzer coupled to a Thermo Finnigan Delta Plus XP isotope ratio mass

spectrometer at the University of New Mexico Center for Stable Isotopes (Albuquerque,

NM). Stable isotope data are expressed in delta (δ) notation as parts per thousand (‰)

according to the equation, $\delta^{15}N = (R_{sample} / R_{standard} - 1) \times 1000$, where R represents the

ratio heavy to light isotopes (^{15}N/^{14}N) of the sample relative to the international reference

standard, respectively. Measured δ^{15}N values were calibrated relative to AIR with certified

reference materials. Within-run estimates of analytical precision were obtained by

measurements of an internal laboratory reference material and yielded a precision (SD) ≤

0.2‰.

Total Niche Width and Individual Diet Specialization. To quantify the total niche width and

degree of individual diet specialization between age groups and seasons, we modified the

Roughgarden index for use with tissue δ^{15}N data (Roughgarden, 1972; Maldonado et al.,

2017, 2019). Roughgarden used a simple framework to quantify individual diet

specialization: the total niche width (TNW) of the (adult or juvenile) population is the sum

of two components: (1) the within-individual component (WIC) which reflects the average

of the variability of resources used by individuals, and (2) the between individual

component (BIC) that represents the between-individual variation in the average use of

resources. The degree of individual diet specialization is reflected in the WIC/TNW ratio

where low ratios close to zero indicate a higher prevalence of specialization within an age

class, indicating a generalist population composed of individual specialists that each use a

small subset of the total niche width.

Trophic Level. We calculated the relative trophic level (TL) of each individual as follows

(Post, 2002): $TL = (1 + (\delta^{15}N_{bird} - \delta^{15}N_{producers}) / \Delta^{15}N)$, where the $\delta^{15}N_{bird}$ represents the mean

isotopic signature of the three tissues, $\delta^{15}N_{producers}$ correspond to the baseline isotopic

signature of the primary producers and $\Delta^{15}N$ is the trophic discrimination enrichment factor.

We used a mean trophic discrimination of 2.48‰, which is the mean value of collagen

(claws), blood, and plasma tissues for birds (Caut et al., 2009), and a baseline $\delta^{15}N$ value

of 3.47‰ estimated for plant samples from the study site.

Statistical Analysis. To assess for differences in individual diet specialization between

juveniles and adults against a null model, we employed a non-parametric Monte Carlo

procedure with 10,000 iterations to derive p-values. Notably, the null model characterizes

the trophic niche of either juveniles or adults as being comprised of generalist individuals

who forage randomly across the entire niche of their respective age classes (Bolnick et al.,

2002). This Monte Carlo procedure is implemented in the R package RInSp (Zaccarelli et

al., 2013). To evaluate whether significant differences exist in TNW, BIC, WIC, and

WIC/TNW between juveniles and adults, we utilized a Monte Carlo permutation procedure.

To calculate a p-value, the observed mean difference in WIC/TNW between juveniles and

adults was compared against the mean differences generated from 10,000 random

permutations of juvenile-adult groups. We then used a one-way Analysis of Variance

(ANOVA) followed by Tukey's post hoc test to assess potential differences in trophic levels

between the two age classes across seasons. Finally, we estimated the isotopic niche

overlap between juveniles and adults in both seasons by calculating the overlapped areas

relative to each pair of δ^{15}N kernel density estimations using the overlapping R package

(Pastore, 2018; Pastore and Calcagnì, 2019). To determine if the estimated area of

overlap was significantly different from what would be expected by chance, we conducted

a permutation test with 10,000 random combinations of the δ^{15}N values and compared the

estimated overlap area with the $CI_{0.95}$ of the permuted data.

Results

The percent of living plant cover decreased from 86% (CI: 82.1–90.1%) in spring to 0%

during the summer. A total of 27 herbaceous plant species were available during spring,

with an estimated Shannon diversity ($D^{q=1}$) of 4.5 (CI: 4.3–5.3). Conversely, no significant

differences were observed in arthropod density (d) between seasons; d= 20.7 (CI: 9.7–

37.7) in spring and d = 15.2 (CI: 9.4–22.6) in summer. Shannon diversity was also non-

significant: $D^{q=1}$ = 11.255 (CI: 10.89 – 11.97) for spring and $D^{q=1}$ = 12.43 (CI: 11.94 – 13.64)

for summer (See Supplementary Materials). However, the taxonomic richness of

arthropods was significantly higher in spring ($D^{q=0}$ = 132.00, CI: 120.94 – 143.06) than in

summer ($D^{q=0}$ = 81.00, CI: 73.60 – 88.40).

Individual diet specialization and total niche width (TNW) varied significantly between

seasons for juveniles but not for adults (Figure 1, Table 1). Juveniles exhibited higher

degrees of individual diet specialization in spring than in summer, while TNW was higher in

summer than in spring. In contrast, adults maintained a consistent degree of individual diet

specialization and TNW across seasons. Notably, adults showed a decrease in trophic

level in summer relative to spring, while juveniles maintained a consistent trophic level

between seasons (Table 2). The amount of niche overlap between juveniles and adults

varied from 0.595 in spring to 0.390 in summer (Table 3).

Discussion

This study provides valuable insights into the trophic niche structure and individual diet

specialization of juveniles and adults in a species with substantial parental care during

periods that differ in ecological opportunity. Contrary to our expectations, we found that

ecological opportunity influences both total niche width (TNW) and the degree of individual

diet specialization in juveniles (Table 1). However, adults show no change in these niche

characteristics over the breeding season despite a shift in their trophic level (Table 2).

Previous studies (Lopez-Calleja, 1995) and our findings show that ecological

opportunity is highest in spring because of higher arthropod richness and higher plant

cover and diversity. The greater availability of resources enables adults to specialize their

resource use for them and their offspring. Consequently, both age groups were primarily

composed of individual specialists exhibiting nearly identical degrees of individual diet

specialization and trophic levels. The high overlap found between juveniles and adults

further indicates the similarities in niche characteristics between these two groups during

this season (Table 3).

Following the same rationale and according to the niche variation hypothesis, we

expected that faced with a reduction in food resources, the total niche width and degree of

individual diet specialization should decrease in both age groups. In contrast to this

prediction, the prevalence of individual diet specialization only decreased in juveniles in

summer by means of a slight increase of WIC and a broadening of their total niche width

(Table 1). Our explanation is that juveniles require a protein-rich diet for development,

which in turn compels adults to provide food to maintain juvenile requirements under

conditions of low resource availability, even at the expense of their own ability to

specialize. This explanation aligns with optimal foraging theory, suggesting that parents

feed their offspring with additional less preferred food items when resources are scarce,

which results in a wider TNW and more generalist foraging behavior observed in juveniles

during the summer relative to spring. The seasonal shift from specialists to generalists

observed for juveniles can also result from parents becoming less selective. Under

reduced resource availability, parents are likely compelled to be less selective in their prey

choice to minimize predation risk associated with an increase in search time. This change

in selectivity would allow them to reduce provisioning rates without compromising

nutritional delivery to their offspring (Skutch., 1949).

The TNW and WIC of adults were similar across seasons, but this age class occupied

a lower trophic level in summer relative to spring. This resulted in a significantly lower

amount of dietary overlap between juveniles and adults during the summer (Table 3). This

suggests that adults may be inherently more flexible in resource selection because they

are not constrained by the same nutritional requirements as developing nestlings. These

conditions enable adults to persist during periods of relative resource scarcity while

prioritizing the delivery of higher trophic level prey items to their nestlings during critical

periods of their growth and development.

In conclusion, the results of this study highlight a dynamic interplay between

ecological opportunity and individual diet specialization, with significant implications for

understanding species resilience to environmental variation (Bolnick et al., 2003; Araújo et

al., 2011). Notably, our study reveals that adult sparrows adapt their parental care

strategies in response to seasonal fluctuations in resource availability. The dichotomy in

resource use between age groups illustrates the complex strategies species employ to

ensure survival and reproductive success. For juveniles, the shift towards generalism in

times of resource scarcity may be imperative to survival, ensuring adequate nutrient intake

for growth and development. In contrast, an adult population's ability to maintain niche

width but alter trophic position underlines a strategic flexibility that likely minimizes

intraspecific competition and optimizes resource utilization across the population.

Ultimately, this study underscores the importance of considering age-specific responses to

fully understand the ecological and evolutionary mechanisms governing niche partitioning

and individual diet specialization within populations. In Mediterranean ecosystems like

central Chile, climate change projections predict an increase in temperatures and a

decrease in annual precipitation (Bozkurt et al., 2017, 2018), therefore the relatively dry

resource-limited conditions associated with summer may extend across a larger portion of

the reproductive season. Studies like ours that focus on characterizing dietary flexibility

and understanding how resource use may shift under these conditions are essential for

understanding species response(s) to ongoing environmental change.

Acknowledgments: This work was funded by NR Postdoctoral Fellowship FONDECYT

3210704, KM Project FONDECYT REGULAR 1200513, and PS ANID PIA/BASAL center

FB0002.

References

Alonso, H., Granadeiro, J. P., Paiva, V. H., Dias, A. S., Ramos, J. A., and Catry, P. (2012).

Parent-offspring dietary segregation of Cory's shearwaters breeding in contrasting

environments. *Marine Biology* 159, 1197–1207. doi: 10.1007/s00227-012-1900-2

Araújo, M. S., Bolnick, D. I., and Layman, C. A. (2011). The ecological causes of individual

specialisation. *Ecology Letters* 14, 948–958. doi: 10.1111/j.1461-0248.2011.01662.x

Bearhop, S., Adams, C. E., Waldron, S., Fuller, R. A., and Macleod, H. (2004).

Determining trophic niche width: A novel approach using stable isotope analysis. *Journal*

of Animal Ecology 73, 1007–1012. doi: 10.1111/j.0021-8790.2004.00861.x

Beaulieu, M., and Sockman, K. W. (2014). Comparison of optimal foraging versus life-

history decisions during nestling care in Lincoln's Sparrows Melospiza lincolnii through

stable isotope analysis. *Ibis* 156, 424–432. doi: 10.1111/ibi.12146

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., et al.

(2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology*

and Evolution 26, 183–192. doi: 10.1016/j.tree.2011.01.009

Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., et al.

(2003). The ecology of individuals: Incidence and implications of individual specialization.

American Naturalist 161, 1–28. doi: 10.1086/343878

Bolnick, D. I., Yang, L. H., Fordyce, J. A., Davis, J. M., and Svanbäck, R. (2002).

Measuring individual-level resource specialization. *Ecology* 83, 2936–2941.

Bond, A. L., Jardine, T. D., and Hobson, K. A. (2016). Multi-tissue stable-isotope analyses

can identify dietary specialization. *Methods in Ecology and Evolution* 7, 1428–1437.

Bozkurt, D., Rojas, M., Boisier, J. P., and Valdivieso, J. (2017). Climate change impacts on

hydroclimatic regimes and extremes over Andean basins in central Chile. *Hydrology and*

Earth System Sciences Discussions, 1–29. doi: 10.5194/hess-2016-690

Bozkurt, D., Rojas, M., Boisier, J. P., and Valdivieso, J. (2018). Projected hydroclimate

changes over Andean basins in central Chile from downscaled CMIP5 models under the

low and high emission scenarios. *Climatic Change* 150, 131–147. doi: 10.1007/s10584-

018-2246-7

Carleton, S. A., Kelly, L., Anderson-Sprecher, R., and del Rio, C. M. (2008). Should we

use one-, or multi-compartment models to describe 13C incorporation into animal tissues?

Rapid Communications in Mass Spectrometry: An International Journal Devoted to the

Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry 22, 3008–3014.

Caut, S., Angulo, E., and Courchamp, F. (2009). Variation in discrimination factors (Δ15N

and Δ13C): the effect of diet isotopic values and applications for diet reconstruction.

Journal of Applied Ecology 46, 443–453.

Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., et al. (2014).

Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation

in species diversity studies. *Ecological Monographs* 84, 45–67. doi: 10.1890/13-0133.1

Costa-Pereira, R., Rudolf, V. H. W., Souza, F. L., and Araújo, M. S. (2018). Drivers of

individual niche variation in coexisting species. *Journal of Animal Ecology* 87, 1452–1464.

doi: 10.1111/1365-2656.12879

Di Castri, F., and Hajek, E. R. (1976). Bioclimatología de chile.

Durell, S. leV Dit. (2003). The implications for conservation of age-and sex-related feeding

specializations in shorebirds. *Wader Study Group Bulletin*, 35–39.

Finke, D. L., and Snyder, W. E. (2008). Niche Partitioning Increases Resource Exploitation

by Diverse Communities. *Science* 321, 1488–1490. doi: 10.1126/science.1160854

Frédérich, B., Lehanse, O., Vandewalle, P., and Lepoint, G. (2010). Trophic niche width,

shift, and specialization of dascyllus aruanus in Toliara Lagoon, Madagascar. *Copeia*,

218–226. doi: 10.1643/CE-09-031

Hahn, S., Dimitrov, D., Rehse, S., Yohannes, E., and Jenni, L. (2014). Avian claw

morphometry and growth determine the temporal pattern of archived stable isotopes.

Journal of Avian Biology 45, 202–207. doi: 10.1111/j.1600-048X.2013.00324.x

Hall, L. A., De La Cruz, S. E. W., Woo, I., Kuwae, T., and Takekawa, J. Y. (2021). Age-

and sex-related dietary specialization facilitate seasonal resource partitioning in a

migratory shorebird. *Ecology and Evolution* 11, 1866–1876. doi: 10.1002/ece3.7175

Hammerschlag-Peyer, C. M., Yeager, L. A., Araújo, M. S., and Layman, C. A. (2011). A

hypothesis-testing framework for studies investigating ontogenetic niche shifts using stable

isotope ratios. *PLoS ONE* 6. doi: 10.1371/journal.pone.0027104

Hanson, P. C., Pace, M. L., Carpenter, S. R., Cole, J. J., and Stanley, E. H. (2015).

Integrating landscape carbon cycling: research needs for resolving organic carbon budgets

of lakes. *Ecosystems* 18, 363–375.

Hobson, K. A. (1993). Trophic relationships among high Arctic seabirds: insights from

tissue-dependent stable-isotope models. *Marine Ecology Progress Series* 95, 7–18. doi:

10.3354/meps095007

Hobson, K. A., and Clark, R. G. (1992). Assessing avian diets using stable isotopes I:

Turnover of 13C in tissues. *Condor* 94, 181–188. doi: 10.2307/1368807

Hseih, T., Ma, K., and Chao, A. (2019). iNEXT: iNterpolation and EXTrapolation for

species diversity. *R package* 2, 19.

Jaksic, F. M., and Lazo, I. (1999). Response of a bird assemblage in semiarid Chile to the

1997-1998 El Niño. *The Wilson Bulletin*, 527–535.

Jia, G., Shevliakova, E., Artaxo, P., Noblet-Ducoudré, N. D., Houghton, R., House, J., et

al. (2019). "Land–climate interactions," in *Climate Change and Land: an IPCC special*

report on climate change, desertification, land degradation, sustainable land management,

food security, and greenhouse gas fluxes in terrestrial ecosystems.

Keast, A. (1977). Mechanisms Expanding Niche Width and Minimizing Intraspecific

Competition in Two Centrarchid Fishes. *Evolutionary Biology*, 333–395. doi: 10.1007/978-

1-4615-6953-4_7

Kyriazakis, I., and Oldham, J. D. (1993). Diet selection in sheep: the ability of growing

lambs to select a diet that meets their crude protein (nitrogen × 6.25) requirements. *British*

Journal of Nutrition 69, 617–629. doi: 10.1079/bjn19930064

Lopez-Calleja, M. V. (1995). Dieta de Zonotrichia capensis (Emberizidae) Y Diuca diuca

(Fringillidae): efecto de la variación estacional de los recursos tróficos y la riqueza de aves

granívoras en Chile central. *Revista Chilena de Historia Natural* 68, 321–331.

Maiorana, V. C. (1978). Difference in diet as an epiphenomenon: space regulates

salamanders. *Canadian Journal of Zoology* 56, 1017–1025. doi: 10.1139/z78-144

Maldonado, K., Bozinovic, F., Newsome, S. D., and Sabat, P. (2017). Testing the niche

variation hypothesis in a community of passerine birds. *Ecology* 98, 903–908. doi:

10.1002/ecy.1769

Maldonado, K., Newsome, S. D., Razeto-Barry, P., Ríos, J. M., Piriz, G., and Sabat, P.

(2019). Individual diet specialisation in sparrows is driven by phenotypic plasticity in traits

related to trade-offs in animal performance. *Ecology Letters* 22, 128–137. doi:

10.1111/ele.13174

Matich, P., Heithaus, M. R., and Layman, C. A. (2011). Contrasting patterns of individual

specialization and trophic coupling in two marine apex predators. *Journal of Animal*

Ecology 80, 294–305. doi: 10.1111/j.1365-2656.2010.01753.x

Miller, A. H., and Miller, V. D. (1968). the Behavioral Ecology and Breeding Biology of the

Andean Sparrow, Zonotrichia Capensis. *Source: Caldasia* 10, 83–154.

Pastore, M. (2018). Overlapping: a R package for Estimating Overlapping in Empirical

Distributions. *Journal of Open Source Software* 3, 1023. doi: 10.21105/joss.01023

Pastore, M., and Calcagnì, A. (2019). Measuring Distribution Similarities Between

Samples: A Distribution-Free Overlapping Index. *Frontiers in Psychology* 10. Available at:

https://www.frontiersin.org/articles/10.3389/fpsyg.2019.01089 (Accessed January 31,

2024).

Podlesak, D. W., McWilliams, S. R., and Hatch, K. A. (2005). Stable isotopes in breath,

blood, feces and feathers can indicate intra-individual changes in the diet of migratory

songbirds. *Oecologia* 142, 501–510.

Post, D. M. (2002). Using stable isotopes to estimate trophic position: models, methods,

and assumptions. *Ecology* 83, 703–718.

Ramos, R., Reyes-González, J. M., Morera-Pujol, V., Zajková, Z., Militão, T., and

González-Solís, J. (2020). Disentangling environmental from individual factors in isotopic

ecology: A 17-year longitudinal study in a long-lived seabird exploiting the Canary Current.

Ecological Indicators 111, 105963. doi: 10.1016/j.ecolind.2019.105963

Raya Rey, A., Polito, M., Archuby, D., and Coria, N. (2012). Stable isotopes identify age-

and sex-specific dietary partitioning and foraging habitat segregation in southern giant

petrels breeding in Antarctica and southern Patagonia. *Marine Biology* 159, 1317–1326.

doi: 10.1007/s00227-012-1912-y

Riverón, S., Raoult, V., Baylis, A. M. M., Jones, K. A., Slip, D. J., and Harcourt, R. G.

(2021). Pelagic and benthic ecosystems drive differences in population and individual

specializations in marine predators. *Oecologia* 196, 891–904. doi: 10.1007/s00442-021-

04974-z

Roughgarden, J. (1972). Evolution of niche width. *The American Naturalist* 106, 683–718.

Skutch., A. F. (1949). Do Tropical Birds Rear as Many Young as They Can Nourish ? *Ibis*

91, 430–455. doi: 10.1111/j.1474-919X.1949.tb02293.x

Southwick, L. H. (1978). *Ecological methods: with particular reference to the study of*

insect populations. Chapman and Hall.

Stephens, R. B., Hobbie, E. A., Lee, T. D., and Rowe, R. J. (2019). Pulsed resource

availability changes dietary niche breadth and partitioning between generalist rodent

consumers. *Ecology and Evolution* 9, 10681–10693. doi: 10.1002/ece3.5587

Svanbäck, R., and Bolnick, D. I. (2005). Intraspecific competition affects the strength of

individual specialization: An optimal diet theory method. *Evolutionary Ecology Research* 7,

993–1012.

Sword, G. A., and Dopman, E. B. (1999). Developmental specialization and geographic

structure of host plant use in a polyphagous grasshopper, Schistocerca emarginata

(=lineata) (Orthoptera: Acrididae). *Oecologia* 120, 437–445. doi: 10.1007/s004420050876

Tubaro, P. L. (1990). Aspectos causales y funcionales de los patrones de variación del

canto del chingolo (Zonotrichia capensis). Universidad de Buenos Aires. Facultad de

Ciencias Exactas y Naturales. Available at:

https://bibliotecadigital.exactas.uba.ar/download/tesis/tesis_n2382_Tubaro.pdf (Accessed

February 8, 2024).

Vander Zanden, H. B., Bjorndal, K. A., and Bolten, A. B. (2013). Temporal consistency and

individual specialization in resource use by green turtles in successive life stages.

Oecologia 173, 767–777. doi: 10.1007/s00442-013-2655-2

Vander Zanden, H. B., Bjorndal, K. A., Reich, K. J., and Bolten, A. B. (2010). Individual

specialists in a generalist population: Results from a long-term stable isotope series.

Biology Letters 6, 711–714. doi: 10.1098/rsbl.2010.0124

Zaccarelli, N., Bolnick, D. I., and Mancinelli, G. (2013). RI n S p: an r package for the

analysis of individual specialization in resource use. *Methods in Ecology and Evolution* 4,

1018–1023.

Table 1: Estimated within-individual component (WIC), between-individual component

(BIC), total niche width (TNW), and individual specialization (WIC/TNW) of adults and

juveniles individual rufous-collared sparrows captured during summer and spring. Units of

WIC, BIC, and TNW are ‰^{2}; sample sizes are noted in parentheses.

	Spring		Summer	
	Juveniles (11)	**Adults** (16)	**Juveniles** (8)	**Adults** (20)
WIC	0.252[a]	0.436[a]	1.924[b]	0.566[a]
BIC	0.958[a]	1.665[a]	1.482[a]	2.980[a]
TNW	1.210[a]	2.101[a]	3.405[b]	3.546[a]
WIC/TNW	0.208[a]	0.207[a]	0.565[b]	0.160[a]
Trophic Level	2.36[a]	2.57[a]	2.54[a]	2.0[b]

473 Different superscript letters denote significant differences ($P < 0.05$) among groups.

475 **Table 2**. Results of Analysis of Variance (ANOVA) assessing changes in trophic level among

476 sparrow age (juveniles and adults) and season of tissue collection (spring and summer).

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
factor(age)	1	0.22	0.22	0.64	0.43
factor(season)	1	1.45	1.45	4.20	0.04*
factor(age) : factor(season)	1	1.67	1.67	4.83	0.03*
Residuals	51	17.66	0.35		

477 *P<0.05

478 **Table 3:** Estimated isotopic niche overlap between juveniles and adults in both seasons

479 using $\delta^{15}N$ kernel density estimates. Permuted quantiles were calculated with a confidence

480 interval of 0.95.

Group 1	Group 2	Overlap	Permuted Quantiles	
			Low 95%	High 95%
J_{Spring}	J_{Summer}	0.593	0.534	0.884
J_{Spring}	A_{Spring}	0.595	0.572	0.866
J_{Spring}	A_{Summer}	0.308	0.574	0.894
J_{Summer}	A_{Spring}	0.489	0.511	0.851
J_{Summer}	A_{Summer}	0.390	0.492	0.890
A_{Spring}	A_{Summer}	0.206	0.641	0.910

482 **Figure 1:** Distributions with thin lines correspond to individual niches for sparrows

483 captured in the spring (blue) and summer (yellow) of juveniles (A and B) and adults (C and

484 D) generated from δ^{15}N values of blood plasma, red blood cells, and claw tips.

485 Distributions with thick lines represent the corresponding total population niche width

486 (TNW), which were scaled equally for the four populations enabling direct comparison.