The role of many-to-one mapping of vertebral form to function in Psittaciform tripedal locomotion

Anna R. Stuart[1*], Michael C. Granatosky[2,3], Ryan N. Felice[1,5,6] and Ryan D. Marek[1]

[1]Centre for Integrative Anatomy, Department of Cell and Developmental Biology, University College London, London, UK.

[2]Department of Anatomy, New York Institute of Technology College of Osteopathic Medicine, Old Westbury, New York, NY, USA

[3]Center for Biomedical Innovation, New York Institute of Technology College of Osteopathic Medicine, Old Westbury, New York, NY, USA

[4]Duke Lemur Center, Duke University, Durham, NC, USA

[5]Department of Life Sciences, Natural History Museum, London, UK

[6]Department of Genetics, Evolution, and Environment, University College London, London, UK

[*] corresponding author

Correspondence (email): anna.stuart.22@alumni.ucl.ac.uk

Acknowledgements:

This work was funded by a Leverhulme Trust Research Project Grant (award number RPG-2021-088) to RNF. We would like to thank Judith White at the Natural History Museum, Tring for access to specimens.

Abstract (197/200 words)

Parrots highlight the functional diversity of the avian neck by contributing to a range of behaviors,
including arboreal locomotion. The parrot neck is used alongside the beak and hindlimb to allow
them to successfully navigate arboreal habitats via tripedal locomotion. Whether specific
morphological characteristics of the neck enable this behavior are currently unknown. By combining
geometric morphometrics with phylogenetic comparative methods we investigate the factors
correlate with shape variation in the cervical vertebrae of parrots. We find that phylogeny, allometry,
integration, diet and tripedal locomotion all have a significant influence on the morphology of
psittaciform cervical vertebrae. However, the influence of diet and tripedal locomotion is weak, with
a high degree of morphospace overlap existing between dietary and neck use groups. Additionally,
we find no evidence of convergence in parrot neck morphology due to the incidence of tripedal
locomotion or dietary specialization. We thus conclude that changes to the neuromuscular control of
the neck, not morphological adaptations, are primarily responsible for tripedal locomotion in
parrots. We argue that many-to-one mapping of form to function allows parrots with similar neck
morphologies to participate in a range of behaviors, and this may be a common feature amongst all
birds.

Introduction

The avian forelimb is characterised by its highly specialised anatomy that allows for powered flight.
As a consequence of this derived structure and function, it is generally maladapted for many
functions outside of this key innovation (1–3). As such a large number of functions that were
ancestrally performed by the forelimb are instead shifted to the craniocervical system (i.e., beak,
head, and neck) in birds. Thus, the neck of birds functions across a broad range of activities as a
'surrogate forelimb' (3–5). Psittaciformes (parrots) exemplify the functional diversity of the avian
cervical column as the necks of this group participate in feeding, preening, tool use and most
spectacularly, locomotion (6–8). By utilising the neck and head as a third 'propulsive limb' parrots
are able to ascend vertical and traverse horizontal substrates by co-opting the craniocervical system
to function within a cyclical tripedal gait pattern (6,8,9). This adaptation of the craniocervical system
to function as a propulsive limb appears to be an evolutionary novelty unique to parrots, and
investigating this phenomenon may offer insight into how the head and neck can be exapted to
actively participate in locomotion.

The use of non-appendicular appendages in locomotion is well-documented across tetrapods. If one
of these appendages displays an interaction between the animal's mass and the substrate then it
can be defined as 'effective limb' (10). There are numerous examples of portions of the axial column
(particularly the tail) acting as an effective limb, however many of these are interacting with the
substrate in an incidental manner (e.g. tail dragging (11–13)), or acting as a stabiliser/brace (6).
Propulsive limbs (i.e., effective limbs that are used for propulsion (6)) are much rarer in tetrapods
and are often limited to the involvement of a tail in powered pentapedal gaits (14,15). Recent work
has highlighted the extraordinary ability of Psittaciformes to use the craniocervical system as a
propulsive limb, with the beak, neck and hindlimbs generating more relative tangential substrate
reaction forces than the forelimbs of humans and primates during vertical climbing (6,8,16). A large
proportion of this propulsive force must be driven by the neck as the thoracolumbar spine of birds is
adapted for stiffness and stability, not force production (6). This has led to the hypothesis that either
the neuronal (6,16) or musculoskeletal system of the psittaciform neck has undergone radical
changes to accommodate tripedal locomotion. If the adaptation is neuronal then many-to-one
mapping of form to function may be a key component of neck evolution in parrots, allowing novel
forms of locomotion to evolve rapidly without the need for large shifts in neck morphology across
the entire cervical spine. Parrots may have utilised many-to-one mapping over the course of their

evolutionary history to allow them to use their neck in a locomotory capacity without sacrificing the
diverse functionality that the 'surrogate forelimb' provides (4). If the adaptation is musculoskeletal,
then these changes can be observed by quantifying relationships between vertebral morphology and
neck usage in parrots.

Many-to-one mapping is a vital concept within evolutionary biology as it can result in a decoupling of
functional diversity from morphological diversity within a clade (17,18). This decoupling allows for
one morphological trait to accommodate multiple functions, and highlights that the morphological
form does not always directly correlate with function (17,19). Many-to-one mapping may be
exemplified by the avian neck as it performs a wide range of functions despite a similarity in its
overall structure between species, and this is evidenced by recent work which observed that the
regional modularity of the avian neck is conserved across many species, only adapting to highly
specialised functions such as carnivory (3,20). Here we investigate whether the psittaciform neck
breaks this pattern of conservatism in avian neck morphology in order to adapt to function during
locomotion, or if many-to-one mapping and changes to neuronal control of parrot neck musculature
are responsible for this novel behaviour.

Parrots have long been known to engage in climbing behaviour and tripedal locomotion during
climbing is thought to be a universal feature of Psittaciformes (21–23). Yet this ubiquity of tripedal
locomotion in parrots is anecdotal (21) and the extent of beak-assisted climbing in this clade has
never been quantified. By using a framework set out by recent research into avian foot use (24) we
can leverage enormous photographic databases of parrots identified to species level in order to
quantitatively determine the incidence of beak-assisted climbing across a broad range of extant
Psittaciformes. We then use the incidence of climbing data to assess any potential form-function
relationships between neck vertebral morphology and beak-assisted climbing. As the psittaciform
neck provides much of the propulsive force associated with climbing (6) we hypothesize that a
significant portion of vertebral shape variation will be governed by the incidence of beak-assisted
climbing. We may further speculate that the selective pressures potentially imposed by beak-
assisted climbing upon the neck leads to morphological convergence of the cervical spine in groups
that are frequently observed carrying out this behaviour. Prior work can be used as a basis for this
hypothesis, and has observed that non-psittaciform birds that share specialised neck kinematics (e.g.
carnivorous birds) display similar patterns of gross neck morphology (3). Phenotypic Integration, the
co-evolution of anatomical traits, is an important facilitator of morphological diversity and has been
found to be an important component of avian skull evolution (25,26). Integration often occurs when
multiple anatomical systems function together during a particular behaviour (27–29), and recent
data suggests it is a commonplace amongst extant avians (30). As tripedal locomotion involves the
neck and hindlimb working in tandem, we also hypothesize that neck-hindlimb integration will
significantly influence morphological variation of neck vertebrae across Psittaciformes.

Here we use a combination of geometric morphometrics, phylogenetic comparative methods and
behaviour quantification to investigate the relationship between neck usage (with a focus on beak-
assisted climbing) and neck vertebral morphology across a phylogenetically broad selection of extant
parrots. We also assess the relative contributions of other factors that have previously been found to
significantly influence the morphological variability of neck vertebral morphology, such as body size,
phylogeny, ecology, and morphological integration. Finally, we will use recently established
convergence metrics to test for convergence in the vertebral morphology parrots that frequently
undertake beak-assisted climbing.

Methods

Specimen details

We analysed the morphology of the cervical vertebrae, head, forelimb and hindlimb of 48 species of
extant parrots (Supplementary Table 1). Neck use behavior was also studied for 44 of 48 of these
species. Six of the 48 species were scanned at UCL with a Nikon XT H 225 microCT scanner. Scan data
for the remaining 42 species were downloaded from MorphoSource (see Supplementary Table 1 for
MorphoSource ID numbers). Dietary and foraging guilds for these species were taken from the
AVONET database (31). Phylogenetic trees were acquired from www.birdtree.org and pruned to
include only the 48 species included in this study.

Digitisation and 3D geometric morphometrics

Scans for all 48 species were segmented in Amira 3D (version 2021.1, Visualization Science Group,
Thermo Fisher Scientific) and the digital models outputted were cleaned and further processed in
MeshLab (32). To account for the variation in total cervical vertebral counts between species we
analysed 2 homologous vertebrae (the second cervical vertebrae, C2, and the last cervical vertebrae)
as well as 3 'functionally homologous' vertebrae (vertebrae at 25%, 50% and 75% along the cervical
column) (20,33,34). A vertebral landmark scheme consisting of 22 fixed landmarks (Supplementary
Figure 1, Supplementary Table 2) was constructed based on schemes from prior avian vertebral
morphometric studies (3,20,35,36). We next calculated head volumes by subjecting digital skull
models for each species to an α-shape fitting algorithm that is part of an in-house modified version
of the 'alphavol' package for MatLab (3,37). These volumes were then multiplied by the weighted
mean densities of soft tissues within the skull (approximated to the density of water, 997 kg/m^3).
This method may overestimate head mass as it does not model the degree of pneumatic bone and
soft tissue within each skull, but such granularity was outside the scope of this study. Head mass was
preferred over head shape as head shape often poorly correlates with ecology in birds (25,26,38).
We then measured limb element lengths digitally in Geomagic Wrap (Geomagic, United States).
Forelimb elements measured include the coracoid, scapula, humerus, radius, ulna and
carpometacarpus. Hindlimb elements measured include the femur, tibiotarsus and tarsometatarsus.
We size-corrected limb measurements using the following formula: limb element length/body
mass$^{0.33}$. We also size-corrected head mass by calculating the percentage of total body mass. Scaling
equations based on the humeral articulation facet of the coracoid were used to estimate body
masses (39).

Assessment of neck use behavior

To assess the extent to which the neck was being used by parrots, we devised a scoring system that
noted the presence or absence of four neck functions assumed to put relatively high loads on the
neck [i.e., beak assisted climbing (6,9,16), forceful flexion and extension (40–42), object carrying)].
We used the Macaulay Library (https://www.macaulaylibrary.org) to search for images of parrots
performing these neck use behaviors following protocols from prior literature (24). Up to 2000
images were searched for each of the 48 species in the study (24) (see Supplementary Table 3 for
behavior scorings and incidence data). We assessed the relationship between number of photos
searched and neck use score for each behavior and found it to be significant for each of the four neck
behaviors, as well as for the overall presence or absence of absolute neck use. To accommodate for
this we use an incidence of behavior metric that accounts for the number of photos searched.
Initially we observed a significant relationship between this incidence metric and the number of
photos searched, but after the removal of 4 outliers (species with an unusually high incidence of
beak assisted climbing despite < 50 photos searched per species) there was no significant correlation
($p < 0.05$).

It should be noted that our method for scoring neck function was a necessity owing to the lack of
available quantitative information on neck use in parrots (43). Generally, this approach is
synonymous with scan sampling, which provides an unbiased assessment of the activity budget of an
animal, but suffers from missing rare or uncommon behaviors (44,45). Indeed, this is confirmed with
the relatively low occurrence of neck use observed in parrots in this study (see below). For rare
behaviours, focal animal or specific behaviour sampling would be most appropriate (44,45).
However, no studies have yet to quantify the positional behaviour of parrots, let alone any non-
mammalian species, utilizing such methodology.

Statistical analysis

Morphological analysis

We first subjected landmark data to Procrustes superimposition. We then performed morphological
analyses on a pooled dataset that contained all vertebrae (C2, C25%, C50%, C75% and the last
cervical vertebrae) for all species, as well as separate analyses for each vertebral level. This allowed
us to study morphological evolution across the entire parrot neck and at the level of individual
vertebrae across the cervical spine (20). We calculated multivariate Blomberg's K (K_{mult}) using the
function physig() within the 'geomorph' R package to estimate the impact of phylogeny on the
morphological variation of parrot cervical vertebrae (46). To assess the relationship between
vertebral shape variation and ecological parameters, neck use behaviors and body mass we utilized
phylogenetic multivariate ANOVAs (pMANOVAs). A reduced dataset was used in the neck use
behavior pMANOVAs as we identified some outlier species (see prior methods section, 'Assessment
of neck use behavior'). We determined significant differences in morphology between ecological
groups using post-hoc pairwise tests with the 'pairwise' function in the 'RRPP' package in R.

We identified the degree of head-neck, neck-forelimb and neck-hindlimb integration using a
phylogenetic two-block partial least-squares (2BPLS) analysis in the R package 'geomorph' (46). We
then used the 'compare.pls' function (as part of the 'geomorph' R package) to search for significant
differences in effect sizes across integration tests. Next, we combined all measurements of individual
forelimb and hindlimb elements into a single forelimb or hindlimb matrix prior to inclusion in any
2BPLS analyses. Following this, we carried out pMANOVAs and 2BPLS tests for a pooled vertebral
dataset and for each individual vertebral level. In order to assess differences in morphological shape
change across the entire neck we applied Phenotypic Trajectory Analysis (PTA) (47). PTA plots a
trajectory through shape space for a particular group (in this case for a particular dietary or foraging
category) by connecting the mean shape of a particular vertebrae in a sequential chain from C2 to
the last cervical vertebrae. This allows for the statistical assessment of differences between certain
groups in the pattern of sequential vertebral shape change across the entire neck.

We produced phylomorphospace plots of PC1 and PC2 for each individual cervical vertebrae to
visualize differences in morphospace occupation between ecological groups and between disparate
incidences of neck use behavior values. We focused on exploring these axes because for each
vertebra, the subsequent PC axes each explained < 30% of the overall variation in each ordination.
We used these phylomorphospace plots to inform our hypotheses for morphological convergence
analyses. These visualizations allowed us to identify focal taxa that share the same ecology or neck
use traits and also occupied a distinct area of morphospace and were thus candidates for convergent
evolution. We then tested for morphological convergence using the Ct1-Ct4 metrics using the
'calcConvCt' and 'calcSigCt' functions in the R package 'convevol' (48,49).

Behavior analysis

Following the protocol of previous work (24) we first tested for any correlation between the number
of photos studied versus the overall occurrence of neck use behavior with a PGLS using the 'gls'
function of the R package 'nlme'. The absolute occurrence of neck use displayed a significant
correlation with the number of media analysed, and as such we accounted for the number of media
by using an incidence of neck use metric. We used Phylogenetic ANOVAs to assess potential
correlations between the incidence of neck use behaviors and ecological categories, body mass and
head mass. We visualized the distribution of the incidence of neck use behavior across our sample by
mapping this trait onto a phylogenetic tree using the 'contMap' function in 'phytools' (50).

Results

Morphological variation of parrot neck vertebrae

Three out of five of the studied cervical vertebrae occupy distinct areas of morphospace when all
vertebrae are projected onto the same morphospace, with only C50% and C75% displaying any level
of overlap (Supplementary Figure 4). Principal component (PC) 1 accounts for 38.15% of the
morphological variation and higher PC1 scores are associated with a lengthening of the centrum and
vertebral arch, a decrease in size of the neural spine, a widening of the neural canal and vertebral
arch, and an increase in length of the costal processes (Supplementary Figure 4). PC2 accounts for
27.63% of the morphological variation, with higher PC2 scores corresponding to a decrease in
centrum and vertebral arch length, an increase in neural spine height, a narrower neural canal, more
robust prezygopophyses and transverse processes, a widening of pre- and post-articular facets and in
increase in the size of the ventral spine (Supplementary Figure 4).

We then ordinated morphological variation for each vertebrae individually, revealing two distinct
patterns for either proximal or distal vertebrae. Proximal vertebrae (C50%, C75% and the last cervical
vertebrae) tended to display elongated and deeper centrums along the main axis of variation (PC1),
whilst centrum length shortened across PC1 for more distal vertebrae (C2 and C25%). PC1
corresponds to between 21.4% (C25%) and 43.78% (C75%) of the vertebral morphological variation,
and PC2 corresponds to between 11.27% (C2) and 16.28% (C50%) of vertebral morphological
variation (Figure 1). Increases to values of PC1 of C2 vertebrate correspond to a shortening of the
centrum and vertebral arch, an increase in height and width of the neural spine, more robust
zygapophyses and a reduction in size of both articular facets (Figure 1). An increase in PC2 scores of
C2 vertebrae is associated with a shortening and narrowing of both the centrum and vertebral arch
alongside an increase in neural spine height (Figure 1). Increases to PC1 scores of C25% vertebrae
correspond to a widening and shortening of the centrum and vertebral arch, a small increase in
height of both the neural and ventral spines, and a reduction in the height of the postarticular facet
(Figure 1). Increases to PC2 values correspond to subtle changes in C25% morphology such as a less
posteriorly pronounced ventral lip of the postarticular facet and a slight reduction in costal process
length (Figure 1). For middle (C50%) vertebrae, increases in PC1 scores are associated with a slight
increase in the length and depth of the centrum, a posterior shift in the position of the neural spine,
a narrowing of the transverse processes, shorter postzygapophyses and elongated costal processes
(Figure 1). Increases to PC2 scores in middle vertebrae are associated with an increased neural spine
height and enlarged transverse processes (Figure 1). Changes in vertebral shape across PC1 in C75%
vertebrae correspond to an increased centrum depth, a slight increase in neural spine height, a
minor elongation of the costal processes, and a more concave postarticular surface (Figure 1).
Increases in PC2 scores in C75% vertebrae are associated with increases to the width of both the
transverse processes and the postzygapophyses (Figure 1). For the last cervical vertebrae changes

associated with increased PC1 scores correspond to an increase in centrum length and depth,
increases to neural and ventral spine height, narrower transverse processes, more dorsally
positioned prezygapophyses and narrower postzygapophyses (Figure 1). Shape changes associated
with an increase in PC2 scores of the last cervical vertebrae correspond to a more anteriorly
positioned postzygapophysis and an enlarged postarticular facet (Figure 1).

Allometric and phylogenetic signal in parrot neck morphology

There is a significant relationship between body mass and vertebral shape in the pooled dataset ($p =$
0.004) as well as in C2 ($p = 0.001$), C25% ($p = 0.002$) and the last cervical vertebrae ($p = 0.035$)
(Supplementary Table 4). However, the proportion of morphological variation that can be explained
by body mass alone is small and varies between 0.044 (last cervical vertebrae) and 0.075 (C2)
(Supplementary Table 4). We calculated K_{mult} to determine the strength and significance of the
phylogenetic signal on the morphological variation of parrot cervical vertebrae. A significant ($P <$
0.05) and weak-to-moderate (lowest K_{mult} C2 = 0.496, highest K_{mult} last = 0.672) phylogenetic signal
can be observed in the morphological variation of the pooled vertebral dataset, as well as for all but
one (C75%, $p = 0.111$, $K_{mult} = 0.479$) of the individual cervical regions.

Ecological signal in parrot neck morphology

Results from the pMANOVAs suggest significant differences exist in morphology between dietary
groups within the pooled dataset ($p = 0.004$), as well as within C2 ($p = 0.004$), C25% ($p = 0.014$) and
C75% ($p = 0.014$) vertebrae (Supplementary Table 4). Diet displayed a weak correlation with vertebral
shape across all vertebrae studied (R^2 between 0.121 in C25% and 0.151 in C75%), yet displayed a
comparatively higher coefficient of correlation than body mass (R^2 $0.045 - 0.075$) (Supplementary
Table 4). Foraging guild displayed no significant correlation ($P > 0.05$) with vertebral morphology
across any of the vertebral regions studied (Supplementary Table 4). This weak ecological and
phylogenetic signal in parrot cervical morphology is reflected in the lack of distinction between
ecological and family-level groups in morphospace across all vertebral regions studied.

A post-hoc pairwise test revealed that between dietary groups, terrestrial herbivores and granivores
often displayed significantly different vertebral morphologies compared to most other dietary groups
(Supplementary Table 5). Terrestrial herbivores often displayed significant differences in vertebral
morphology compared to other dietary groups (8 comparisons across C2, C25% and C75%)
(Supplementary Table 5). Vertebral morphology was significantly different between herbivores and
frugivores in C2 ($p = 0.043$) and C25% ($p = 0.049$), between herbivores and generalists in C2 ($p =$
0.04), C25% ($p = 0.04$) and C75% ($p = 0.017$) and between herbivores and nectarivores in C2 ($p -$
0.012), C25% ($p = 0.015$) and C75% ($p = 0.02$) (Supplementary Table 5). Multiple dietary groups also
displayed significant differences in vertebral morphology when compared to granivores, including
nectarivores (C2 $p = 0.026$ and C25% $p = 0.044$) and generalists (C75%, $p = 0.032$) (Supplementary
Table 5). Terrestrial herbivores had predominantly taller neural spines than other dietary groups
across C2, C25% and C75%, a shortened centrum in C25% and a deeper centrum in C75%. Granivores
also display a taller neural spine across C2, C25% and C75%, as well as comparatively deeper centra
in C2 and C75%.

PTA detected that across all dietary, foraging and neck-use groupings, patterns of whole-neck
morphological variation were only significantly different between a select few dietary groups
(Supplementary Table 6). We tested for pairwise differences between dietary, foraging and
phylogenetic groupings separately and found only 4 dietary pairwise comparisons to be significant
(Supplementary Table 6, Supplementary Figure 4). Trajectory shape was significantly different

between terrestrial herbivores and generalists (p = 0.02), as well as between terrestrial herbivores
and granivores (p = 0.035) (Supplementary Table 6). The magnitude of differences between
trajectories was significant between generalists and frugivores (p = 0.035), as well as between
generalists and granivores (p = 0.005) (Supplementary Table 6).

308 The influence of neck use on parrot neck morphology

Our survey of neck use across 44 species of parrots (excluding outliers, see Methods) revealed that
across 40,893 images studied, the incidence of absolute neck use was 0.638%. Across each individual
neck use behavior, average incidence varied between 0.450% for beak-assisted climbing and 0.032%
for object carrying (Supplementary Table 3). Numerous species (out of 48 total) were never observed
performing each behavior and this value ranged between 13 and 14 in absolute incidence of neck
use and beak assisted climbing and 40 in forceful extension (Supplementary Table 3). Before
evaluating the relationship between the incidence of neck use behaviors and cervical morphology,
we assessed the relationship between behavior, body mass, head mass and ecology (diet and
foraging guild). Across all behaviors only the overall incidence of absolute neck use was significantly
correlated with body mass (p < 0.001) and head mass (p = 0.002), beak assisted climbing was also
significantly correlated with head mass only (p = 0.016) (Supplementary Table 7).

We investigated potential correlations between the incidence of each neck use behavior and cervical
vertebral morphology for the pooled dataset as well as for each individual cervical region.
Consequently, we found a significant, weak correlation between the incidence of total neck use and
the morphology of C2 vertebrae (p = 0.034, R^2 = 0.042), as well as similarly weak correlation between
incidence of beak assisted climbing and C2 vertebral morphology (p = 0.043, R^2 = 0.039)
(Supplementary Table 7). Species that have a high incidence of beak assisted climbing rarely occupy
distinct areas of morphospace (except for C2), and often coincide with species with minimal or no
occurrences of beak assisted climbing (Figure 2, Supplementary Figure 4). These high incidence
climbers also do not occupy similar areas of morphospace, often appearing far apart from each other
in all studied morphospaces (Figure 2, Supplementary Figure 4). Convergence tests revealed that
species with a high incidence of beak assisted climbing and total neck use were not convergent (P >>
0.05). We then visually identified smaller subsets (or pairs) of species with high incidence of neck use
behaviors or the same dietary guild that occupied similar areas of phylomorphospace and repeated
the convergence tests. None of these further convergence tests were significant (P >> 0.05).

335 The effects of integration on parrot cervical morphology

Parrots display significant integration across all of three of the studied regimes (P< 0.05 for head-
neck, neck-forelimb and neck-hindlimb integration, Supplementary Table 8). R PLS values were
generally high and ranged from 0.847 for head-neck integration across all vertebrae to 0.522 for
neck-forelimb integration in the last cervical vertebrae (Supplementary Table 8). Whereas R PLS
values and Z scores seem to decrease towards the proximal end of the cervical column, there are no
significant differences between Z scores for integration tests for different vertebrae. Many integrative
relationships are not significant when head, forelimb or hindlimb measures are adjusted for body
mass (Supplementary Table 8). Only the last cervical vertebrae retains a significant pattern of neck-
forelimb integration when adjusted for body mass (Supplementary Table 8). Significance is retained
in the pooled dataset, C50% and the last cervical vertebrae for head-neck integration, and no
vertebral region retains significance after adjusting hindlimb measurements for body mass
(Supplementary Table 8). Across all integration regimes studied, a gradient of body mass can be

observed whereby parrots with high body masses displayed extreme values of PLS1 and PLS2 and
parrots with low body mass clustered together at the other extreme of PLS1 and PLS2 (Figure 3).

Discussion

Here we present the results of the first study to investigate factors that have influenced the
morphological variation of psittaciform cervical vertebrae, as well as results from the first study to
quantify neck use across Psittaciformes. We find that variation in the morphology of parrot neck
vertebrae is governed by a variety of factors including phylogeny, allometry, ecology and integration
with both the head and forelimb. Contrary to prior anecdotal evidence, we find that beak-assisted
climbing is not a ubiquitous behavior across Psittaciformes and was not observed in 14 of the 48
species studied. Furthermore, we find that a weak but significant relationship exists between the
incidence of beak-assisted climbing and the morphological variability of psittaciform C2 vertebrae
(Figure 2, Supplementary Table 7). We were also able to qualitatively and quantitatively demonstrate
that beak-assisted climbing is not responsible for any convergence in the morphology of cervical
vertebrae amongst parrots with similar incidences of this behavior (Figure 2, Supplementary Figure
4). The neck of parrots also displays significant integration with the head, forelimb and hindlimb
however these integrative relationships often disappear when body mass is accounted for
(Supplementary Table 8).

Evolution of morphological form is often multifactorial in nature due to the interplay of phylogenetic
constraints, allometry, ecology and integration with other anatomical components (26–28,51).
Indeed, both evolutionary allometry (phylogeny and body size) and phenotypic integration have
previously been shown to be some of the factors that influence skull shape variation in parrots (26).
The factors that influence morphological variation are therefore broadly similar across the head and
neck of Psittaciformes as phylogeny, body mass, and integration all significantly influence the shape
variation of the psittaciform neck. A key difference between the skull and neck of parrots is that
dietary preference accounts for more morphological variation in the neck of parrots (up to 15% vs
2.4%, Supplementary Table 4). We had expected to observe the opposite pattern as the skull directly
manipulates and processes food, not the neck (7,26,52,53). Diet is still a minor (~15%) component of
neck morphological variation however, and the differences observed here may be due to
discrepancies in dietary classification schemes between the two studies (26). Allometry accounted
for a smaller proportion of cervical shape variation than diet (between 4.4% and 7.5%), however it
evidently plays an important role in the integrative relationships between the neck, head and
appendicular skeleton as the significance of head-neck, neck-forelimb and neck-hindlimb integration
often disappears when body mass is taken into account (Supplementary Table 8). Across many of
these integrative relationships we observed that larger parrots clustered together with extreme PLS1
and PLS2 scores (Figure 3) and this clustering may indicate that body mass may be a controlling
factor in psittaciform neck integration. Since beak-assisted climbing requires the cooperation of the
psittaciform craniocervical and hindlimb skeleton, a coordinated morphological response may be
required in order for this multi-body system behavior to occur in larger parrots (16,51,54–56).

Across multiple morphological scales we observe that parrots with different diets and neck use
behaviors often have similar neck morphologies (Figures 1 & 2, Supplementary Tables 4-7) and we
infer this is a feature of many-to-one mapping (18,19). Although we do observe a significant
relationship between diet, neck use and vertebral morphology, we find that these relationships are

weak (Supplementary Table 4) and that dietary and neck use groups often heavily overlap in
morphospace (Figures 1 & 2). We also find very few significant differences between the pattern of
morphological variation across the entire neck among dietary and behavioral groups (Supplementary
Table 6). The neck of parrots provides much of the propulsion during beak-assisted climbing. Indeed,
kinematic data suggests it is capable of producing relatively greater contractile forces than the
human neck (6,57). It has also been anecdotally reported that all parrots engage in this behavior
(21). In response to this we hypothesized that beak-assisted climbing would be a highly influential
factor in the morphological variance of the cervical column across Psittaciformes, and that this
behavior was leading to morphological convergence of cervical vertebrae. In light of the present
results, we reject this hypothesis and suggest that form and function are somewhat decoupled in the
psittaciform cervical column. Instead of a tight relationship between form and function, we suggest
that many-to-one mapping of form to function is allowing parrots to occupy a diverse range of
dietary niches and to utilize a wide array of neck use behaviors (18,19), including tripedal locomotion
(6,16) and 'beakiation' (9). By displaying neither significant morphological convergence or clear
morphological adaptations to dietary ecology or neck use we hypothesize that changes to muscle
activation patterns and neuromuscular innovation may be responsible for tripedal locomotion and
array of neck use behaviors in Psittaciformes, as has previously been suggested (6,16). The
neuromuscular pathways associated with neck use behaviors such as beak-assisted climbing must
allow for movements of the craniocervical system to be incorporated into the locomotor cycle (6).
This may not require a radical neuromuscular innovation as this pathway already exists to
accommodate avian head-bobbing (58), and may have been modified by parrots to allow for tripedal
locomotion (6,16). Since the neuromuscular pathways associated with the parrot neck are already
optimizes for a wide variety of behaviors, these pathways may display plasticity in their ability to
adapt to novel functions such as beak-assisted climbing (6,16) and 'beakiation' (9).

The findings presented here for parrots may have implications for the mechanisms behind broader
avian neck evolution, as the ecological signal in cervical morphological variation is similarly low
across Aves (3,20,30). The similarly weak influence of ecology on neck morphology across all birds
suggests that many-to-one mapping may be present across Aves. This extrapolation could explain
why there is an apparent disconnect between avian cervical form and function: the overall
morphological construction of the avian neck is highly conserved and only adapts to behaviors that
require specialized kinematic forces, yet the neck still participates in a disparate array of behaviors
(3,4). Although further work is required to formally test the presence of many-to-one mapping in the
parrot and avian cervical column, it has been shown to be a common feature of organismal design
(18,19) that weakens the effect of convergent evolution (59).

Conclusions

This work represents the first quantification of the presence of tripedal locomotion across parrots
and finds that beak-assisted climbing is not a ubiquitous feature of Psittaciformes. We also find that
tripedal locomotion, alongside a multitude of other factors, governs a small portion of morphological
variability of the parrot neck. Parrots with similar cervical morphologies appear to be able to use
their necks to access a wide variety of food types and to both participate and not participate in beak-
assisted climbing. This suggests that many-to-one mapping of cervical form to function is a feature of
the neck of Psittaciformes, and potentially a feature of neck construction across extant Aves. Without
the presence of clear vertebral adaptations to beak-assisted climbing we suggest changes to the
neuromuscular control of the cervical column have underpinned the evolution of tripedal
locomotion in parrots and this may be a modification of existing neural pathways associated with
avian head-bobbing.

Figures & tables

Figure 1: Morphospaces for each of the psittaciform cervical regions studied, grouped by dietary
preference. A) C2 cervical vertebrae, B) C25% cervical vertebrae, C) C50% vertebrae, D) C75%
vertebrae, E) Last cervical vertebrae. Warped meshes of cervical vertebrae display changes across
PC1 and PC2 in anterior (left along PC1, bottom along PC2) view and left lateral (right along PC1, top
along PC2) view.

Figure 2: Phylomorphospace of PC1 and PC2 for the C2 vertebrae of parrots. Points are coloured by
incidence of beak-assisted climbing. Next to the 4 species of parrot that experience the highest
incidence of BAC are photographs of that species as well as the mesh of that species' C2 vertebrae.
Bottom photo: *Lathamus discolor* (photo credit David Irving, Macaulay Library ID ML613044549).
Bottom-left photo: *Psittacus erithacus* (photo credit Manuel-Fernandez-Bermejo, Macaulay Library
ID ML613685886). Top photo: *Rhychopsitta pachyrhychna* (photo credit Ken Chamberlain, Macaulay
Library ID ML608581146). Right photo: *Oreopsittacus arfaki* (photo credit Robert Tizard, Macaulay
Library ID ML613799167).

Figure 3: 2BPLS plots of vertebral shape versus forelimb proportion (Ai-Av), vertebral shape versus
head mass (Bi-Bv) and vertebral shape versus hindlimb proportion (Ci-Cv). Arrows beside the Y-axes
indicate what portion of the limb is increasing as PLS2 scores increase: purple denote the proximal
portion is lengthening and yellow indicates the distal portion is lengthening. Points within each
2BPLS plot are coloured by body mass. Warped meshes of cervical vertebrae for each plot display
shape changes across PLS1. The forelimb is represented by *Ara ararauna* in A, the skull morphology
displayed in B is that of *Micropsitta finschii*, the hindlimb is represented by *Chalcopsitta atra* in C.

Supplementary Figure 1: Visual representation of the landmark scheme used throughout this study.
Mesh is the C25% vertebrae of *Pionites melanocephalus*.

Supplementary Figure 2: Morphospace of vertebral shape across the neck of all parrots studied.
Colours denote vertebral region. Warped meshes of cervical vertebrae display shape change across
PC1 and PC2 in anterior (left along PC1, bottom along PC2) view and left lateral (right along PC1, top
along PC2).

Supplementary Figure 3: Phenotypic trajectory plot depicting patterns of shape change across the
neck of parrots with different dietary niches. Point and line colours denote dietary preference and
point shape denotes vertebral region.

Supplementary Figure 4: Phylomorphospace plots for the C25% (A), C50% (B), C75% (C) and last
cervical vertebrae (D) of 44 species of parrots. Points are coloured by incidence of beak-assisted
climbing.

Supplementary Table 1: Specimen information and metadata for all studied species. Asterisks
indicate outlier taxa that were removed from the neck use incidence analysis.

Supplementary Table 2: Landmark scheme used as part of the geometric morphometric component
of this analysis.

Supplementary Table 3: Neck-use behaviour data for all species. Asterisks indicate outlier taxa that
were removed from the neck use incidence analysis.

Supplementary Table 4: Results from the pMANOVA (phylogenetic multivariate ANOVA) analyses.

Supplementary Table 5: Results summary of post-hoc tests performed after pMANOVA.

Supplementary Table 6: Results summary of the phenotypic trajectory analysis (PTA). MD =
magnitude of differences between trajectories, TC = trajectory correlations, SD = trajectory shape
differences.

Supplementary Table 7: Behavioural MANOVA results table

Supplementary Table 8: 2BPLS results table

References

1.   Heers AM, Dial KP. From extant to extinct: locomotor ontogeny and the evolution of avian flight.
Trends in Ecology & Evolution. 2012 May 1;27(5):296–305.

2.   Marek RD. The Surrogate Arm: Functional Morphology of the Avian Cervical Column [Internet]
[Ph.D.]. [England]: The University of Liverpool (United Kingdom); 2020 [cited 2023 Aug 31].
Available from:
https://www.proquest.com/docview/2451874026/abstract/9CB75106C01A45B2PQ/1

3.   Marek RD, Falkingham PL, Benson RBJ, Gardiner JD, Maddox TW, Bates KT. Evolutionary
versatility of the avian neck. Proceedings of the Royal Society B: Biological Sciences. 2021 Mar
3;288(1946):20203150.

4.   Marek RD. A surrogate forelimb: Evolution, function and development of the avian cervical
spine. Journal of Morphology. 2023;284(10):e21638.

5.   Bhullar BAS, Hanson M, Fabbri M, Pritchard A, Bever GS, Hoffman E. How to Make a Bird Skull:
Major Transitions in the Evolution of the Avian Cranium, Paedomorphosis, and the Beak as a
Surrogate Hand. Integrative and Comparative Biology. 2016 Sep 1;56(3):389–403.

6.   Young MW, Dickinson E, Flaim ND, Granatosky MC. Overcoming a 'forbidden phenotype': the
parrot's head supports, propels and powers tripedal locomotion. Proceedings of the Royal
Society B: Biological Sciences. 2022 May 18;289(1975):20220245.

7.   Young MW, Wilken AT, Manafzadeh AR, Schurr AF, Bastian A, Dickinson E, et al. The dual function
of prokinesis in the feeding and locomotor systems of parrots. Journal of Experimental Biology.
2023 Dec 8;226(24):jeb246659.

8.   Reader LL, Carrier DR, Goller F, Isaacs MR, Moore Crisp A, Barnes CJ, et al. Climbing parrots
achieve pitch stability using forces and free moments produced by axial–appendicular couples.
Journal of Experimental Biology. 2022 Jan 7;225(1):jeb242305.

9.   Dickinson E, Young MW, Granatosky MC. Beakiation: how a novel parrot gait expands the
locomotor repertoire of living birds. Royal Society Open Science. 2024 Jan 31;11(1):231397.

10. Bertram JE. Concepts in locomotion: levers, struts, pendula and springs. Understanding
mammalian locomotion: concepts and applications. 2016;79–109.

11. Willey JS, Biknevicius AR, Reilly SM, Earls KD. The tale of the tail: limb function and locomotor
mechanics in Alligator mississippiensis. Journal of Experimental Biology. 2004 Jan 22;207(3):553–
63.

12. Pridmore P. Submerged walking in the epaulette shark Hemiscyllium ocellatum (Hemiscyllidae)
and its implications for locomotion in rhipidistian fishes and early tetrapods. Zoology.
1994;98:278–97.

13. Zinner D, Fickenscher G, Roos C, Mittermeier R, Rylands A, Wilson D. Handbook of the mammals
of the world. Primates. 2013;

14. O'Connor SM, Dawson TJ, Kram R, Donelan JM. The kangaroo's tail propels and powers
pentapedal locomotion. Biology letters. 2014;10(7):20140381.

15. Schmitt D, Rose MD, Turnquist JE, Lemelin P. Role of the prehensile tail during ateline
locomotion: experimental and osteological evidence. American Journal of Physical Anthropology:
The Official Publication of the American Association of Physical Anthropologists.
2005;126(4):435–46.

16. Young MW, Webster C, Tanis D, Schurr AF, Hanna CS, Lynch SK, et al. What does climbing mean
exactly? Assessing spatiotemporal gait characteristics of inclined locomotion in parrots. J Comp
Physiol A [Internet]. 2023 May 4 [cited 2023 Oct 24]; Available from:
https://doi.org/10.1007/s00359-023-01630-0

17. Alfaro ME, Bolnick DI, Wainwright PC. Evolutionary Consequences of Many-to-One Mapping of
Jaw Morphology to Mechanics in Labrid Fishes. The American Naturalist. 2005 Jun;165(6):E140–
54.

18. Wainwright PC, Alfaro ME, Bolnick DI, Hulsey CD. Many-to-One Mapping of Form to Function: A
General Principle in Organismal Design?1. Integrative and Comparative Biology. 2005 Apr
1;45(2):256–62.

19. Wainwright PC. Functional Versus Morphological Diversity in Macroevolution. Annual Review of
Ecology, Evolution, and Systematics. 2007;38(1):381–401.

20. Marek RD, Felice RN. The neck as a keystone structure in avian macroevolution and mosaicism.
BMC Biology. 2023 Oct 13;21(1):216.

21. Dilger WC. The comparative ethology of the African parrot genus Agapornis. Zeitschrift für
Tierpsychologie. 1960;

22. Brockway BF. Ethological studies of the budgerigar (Melopsittacus undulatus): Non-reproductive
behavior. Behaviour. 1964;22(3–4):193–222.

23. Webster C, Jusufi A, Liu D. A comparative survey of climbing robots and arboreal animals in
scaling complex environments. In Springer; 2019. p. 31–45.

24. Gutiérrez-Ibáñez C, Amaral-Peçanha C, Iwaniuk AN, Wylie DR, Baron J. Online repositories of
photographs and videos provide insights into the evolution of skilled hindlimb movements in
birds. Commun Biol. 2023 Aug 15;6(1):1–11.

25. Bright JA, Marugán-Lobón J, Cobb SN, Rayfield EJ. The shapes of bird beaks are highly controlled
by nondietary factors. Proceedings of the National Academy of Sciences. 2016 May
10;113(19):5352–7.

26. Bright JA, Marugán-Lobón J, Rayfield EJ, Cobb SN. The multifactorial nature of beak and skull
shape evolution in parrots and cockatoos (Psittaciformes). BMC Evolutionary Biology. 2019 May
17;19(1):104.

27. Goswami A, Smaers JB, Soligo C, Polly PD. The macroevolutionary consequences of phenotypic
integration: from development to deep time. Philos Trans R Soc Lond B Biol Sci. 2014 Aug
19;369(1649):20130254.

28. Felice RN, Randau M, Goswami A. A fly in a tube: Macroevolutionary expectations for integrated
phenotypes. Evolution. 2018 Dec 1;72(12):2580–94.

29. Evans KM, Larouche O, Watson SJ, Farina S, Habegger ML, Friedman M. Integration drives rapid
phenotypic evolution in flatfishes. Proceedings of the National Academy of Sciences. 2021 May
4;118(18):e2101330118.

30. Böhmer C, Plateau O, Cornette R, Abourachid A. Correlated evolution of neck length and leg
length in birds. Royal Society Open Science. 2019 May 8;6(5):181588.

31. Tobias JA, Pigot AL. Integrating behaviour and ecology into global biodiversity conservation
strategies. Philosophical Transactions of the Royal Society B: Biological Sciences. 2019 Jul
29;374(1781):20190012.

32. Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G. Meshlab: an open-source
mesh processing tool. In Salerno, Italy; 2008. p. 129–36.

33. Jones KE, Benitez L, Angielczyk KD, Pierce SE. Adaptation and constraint in the evolution of the
mammalian backbone 06 Biological Sciences. BMC Evolutionary Biology. 2018;18(1).

34. Jones KE, Dickson BV, Angielczyk KD, Pierce SE. Adaptive landscapes challenge the "lateral-to-
sagittal" paradigm for mammalian vertebral evolution. Current Biology. 2021 May 10;31(9):1883-
1892.e7.

35. Böhmer C, Rauhut OWM, Wörheide G. Correlation between Hox code and vertebral morphology
in archosaurs. Proceedings of the Royal Society B: Biological Sciences. 2015 Jul
7;282(1810):20150077.

36. Terray L, Plateau O, Abourachid A, Böhmer C, Delapré A, de la Bernardie X, et al. Modularity of
the Neck in Birds (Aves). Evol Biol. 2020 Jun 1;47(2):97–110.

37. Brassey CA, Gardiner JD. An advanced shape-fitting algorithm applied to quadrupedal mammals:
improving volumetric mass estimates. Royal Society Open Science. 2015 Aug;2(8):150302.

38. Navalón G, Bright JA, Marugán-Lobón J, Rayfield EJ. The evolutionary relationship among beak
shape, mechanical advantage, and feeding ecology in modern birds*. Evolution. 2019 Mar
1;73(3):422–35.

39. Field DJ, Lynner C, Brown C, Darroch SAF. Skeletal Correlates for Body Mass Estimation in
Modern and Fossil Flying Birds. PLOS ONE. 2013 Nov 29;8(11):e82000.

40. Ramírez-Herranz M, Sayol F, Rios RS, Antonelli A, Squeo FA. The origin of excavator nesting
behavior and its impact on the evolution of Neotropical parrots. Oikos. 2024;n/a(n/a):e10155.

41. Romero-Vidal P, Blanco G, Hiraldo F, Díaz-Luque JA, Luna Á, Lera D, et al. Nesting innovations in
neotropical parrots associated to anthropogenic environmental changes. Ecology and Evolution.
2023;13(9):e10462.

42. Brightsmith DJ. Competition, predation and nest niche shifts among tropical cavity nesters:
phylogeny and natural history evolution of parrots (Psittaciformes) and trogons (Trogoniformes).
Journal of Avian Biology. 2005;36(1):64–73.

43. Granatosky MC, Young MW, Herr V, Chai C, Raidah A, Kairo JN, et al. Positional Behavior of
Introduced Monk Parakeets (Myiopsitta monachus) in an Urban Landscape. Animals. 2022
Jan;12(18):2372.

44. Lehner PN. Design and Execution of Animal Behavior Research: An Overview. Journal of Animal
Science. 1987 Nov 1;65(5):1213–9.

45. Altmann J. Observational Study of Behavior: Sampling Methods. Behaviour. 1974 Jan 1;49(3–
4):227–66.

46. Adams DC, Otárola-Castillo E. geomorph: an R package for the collection and analysis of
geometric morphometric shape data. Methods in ecology and evolution. 2013;4(4):393–9.

47. Collyer M, Adams D. Phenotypic trajectory analysis: comparison of shape change patterns in
evolution and ecology. 2013;

48. Stayton CT, Stayton MCT. Package 'convevol.' 2015;

49. Grossnickle DM, Brightly WH, Weaver LN, Stanchak KE, Roston RA, Pevsner SK, et al. Challenges
and advances in methods for measuring phenotypic convergence [Internet]. bioRxiv; 2023 [cited
2023 Oct 2]. p. 2022.10.18.512739. Available from:
https://www.biorxiv.org/content/10.1101/2022.10.18.512739v3

50. Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution. 2012;3(2):217–23.

51. Marroig G, Shirai LT, Porto A, de Oliveira FB, De Conto V. The Evolution of Modularity in the
Mammalian Skull II: Evolutionary Consequences. Evol Biol. 2009 Mar 1;36(1):136–48.

52. Snively E, Russell AP, Powell GL, Theodor JM, Ryan MJ. The role of the neck in the feeding
behaviour of the Tyrannosauridae: inference based on kinematics and muscle function of extant
avians. Journal of Zoology. 2014;292(4):290–303.

53. van der Leeuw AHJ, Bout RG, Zweers GA. Control of the Cranio-Cervical System During Feeding in
Birds1. American Zoologist. 2001 Dec 1;41(6):1352–63.

54. Hildebrand M, Bramble DM, Liem KF, Wake DB. Functional vertebrate morphology. Harvard
University Press; 1985.

55. Cant JG. Positional behavior and body size of arboreal primates: a theoretical framework for field
studies and an illustration of its application. American journal of physical anthropology.
1992;88(3):273–83.

56. Granatosky MC. A review of locomotor diversity in mammals with analyses exploring the
influence of substrate use, body mass and intermembral index in primates. Journal of Zoology.
2018;306(4):207–16.

57. Valkeinen H, Ylinen J, Mälkiä E, Alen M, Häkkinen K. Maximal force, force/time and
activation/coactivation characteristics of the neck muscles in extension and flexion in healthy
men and women at different ages. European journal of applied physiology. 2002;88:247–54.

58. Nyakatura JA, Andrada E. On vision in birds: coordination of head-bobbing and gait stabilises
vertical head position in quail. Frontiers in zoology. 2014;11:1–10.

59. Thompson CJ, Ahmed NI, Veen T, Peichel CL, Hendry AP, Bolnick DI, et al. Many-to-one form-to-
function mapping weakens parallel morphological evolution. Evolution. 2017 Nov
1;71(11):2738–49.

Incidence of beak-assisted climbing (%)