The microbiome interacts with the circadian clock and dietary composition to regulate metabolite cycling in the *Drosophila* gut

Yueliang Zhang [a,b], Sara B. Noya [a], Yongjun Li [a], Jichao Fang[b] and Amita Sehgal [a]*.

[a] HHMI, Chronobiology and Sleep Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia PA 19104

[b] Institute of Plant Protection, Jiangsu Academy of Agricultural Sciences, Key Lab of Food Quality and Safety of Jiangsu Province–State Key Laboratory Breeding Base, Nanjing, China

*Correspondence to: amita@pennmedicine.upenn.edu

The gut microbiome plays a key role in the maintenance of host metabolic homeostasis and health. Most metabolic processes cycle with a 24-hour rhythm, but the extent to which the microbiome influences metabolite cycling under different conditions, such as variations in dietary composition, remains largely unknown. In this study, we utilized high temporal resolution metabolite profiling of the *Drosophila* gut to investigate the role of the microbiome in metabolite cycling. We find that the microbiome increases the number of oscillating metabolites despite the previous finding that it dampens transcript cycling in the gut. Time restricted feeding (TF) also promotes metabolite cycling and does so to a larger extent in germ-free flies, thereby increasing cycling in these flies to levels comparable to those in microbiome containing flies. Enhancement of cycling by the microbiome depends upon a circadian clock, which also maintains phase in the face of changes in the microbiome. Interestingly, a high-protein diet increases microbiome-dependent metabolite cycling, while a high sugar diet suppresses it. Gene Ontology identifies amino acid metabolism as the metabolic pathway most affected by changes in the gut microbiome, the circadian clock and timed feeding, suggesting that it is subject to regulation by multiple inputs. Collectively, our observations highlight a key role of the gut microbiome in host metabolite cycling, and reveal a complex interaction with internal and external factors.

Introduction

To adapt to the 24-hour light and temperature cycle caused by the Earth's rotation, organisms have evolved endogenous circadian clocks comprised of auto-regulating clock genes(Sehgal, 2017). Circadian clocks can be in the brain (i.e. central) or in peripheral tissues, and show differential responses to external entraining cues. In general, central clocks are mainly regulated by light signals, while dietary signals affect the rhythm of several peripheral circadian clocks(Albrecht, 2012; Stephan, 2002). Central and peripheral clocks coordinate with each other to achieve synchronization of physiological and behavioral circadian rhythms. Disrupting circadian rhythms can result in a range of metabolic and neurological disorders (Castanon-Cervantes et al., 2010;

44 Rijo-Ferreira and Takahashi, 2019; Turek et al., 2005), underscoring the importance of
45 a well-synchronized circadian system.
46
The gut microbiome consists of microorganisms, including bacteria, viruses, protists,
archaea and fungi, that coexist symbiotically with hosts. Indeed, compelling evidence
shows that gut microbiomes play a vital role in host nutrition, immunity, and
neurodevelopment (Flint, 2012; Seo et al., 2023; Thaiss et al., 2016c). About 10% to 15%
of the microbe species in the mammalian gut exhibit rhythms of abundance, and this is
considered to be a key determinant of host health and fitness (Liang et al., 2015; Thaiss
et al., 2016b; Thaiss et al., 2014). Time restricted feeding (TF), which restricts food
intake to a specific daily interval, enhances oscillations of the microbiome and improves
health conditions in animals with obesity or metabolic syndrome (Chaix et al., 2019;
Lundell et al., 2020; Villanueva et al., 2019). On the other hand, western diets (high fat,
high protein, and high sugar), which significantly increase the occurrence of metabolic
diseases, can disrupt microbiome rhythms (Hariri and Thibault, 2010; Ko et al., 2020;
Na et al., 2013). In addition, gut microbiomes regulate circadian rhythms of gene
expression in metabolic tissues (Thaiss et al., 2016a; Zhang et al., 2023), however their
impact on host metabolite cycling remains largely unknown.
62
We previously reported that the gut microbiome in *Drosophila* does not cycle, but it
regulates cycling of the host transcriptome. Specifically, it suppresses transcript rhythms
in the gut while TF enhances these rhythms (Zhang et al., 2023). Here we sought to test
whether the microbiome regulates metabolite oscillations and how the host clock and
dietary manipulations interact with those cycles. Thus, we assayed metabolite rhythms
in wild type flies with and without a microbiome in different dietary paradigms and in
clock mutant flies. Overall, we observed that the microbiome increases metabolite
cycling. However, in the absence of a host clock, the microbiome decreases cycling of
metabolites in total while dramatically affecting the phase of all. A TF feeding paradigm
increases metabolite cycling, although less so in animals that contain a microbiome. On
the other hand, a protein-rich diet enhances microbiome-dependent cycling, but this is
not the case in sugar-rich diet. Overall, our results suggest the gut microbiome plays a
critical role in maintaining cycles of metabolic activity in the gut, in a manner that
depends on the host circadian clock, diet composition and feeding behavior.
77
78
79 **Methods**
80
81 **Generation and maintenance of fly lines**
82
The w^{118} *iso31, per^{01}* fly lines present in the study were maintained on standard
cornmeal/molasses medium at 25°C under 12:12 LD conditions. The germ-free flies
used in this study are the same as in the previous report(Feltzin et al., 2019; Zhang et
al., 2023). The Newborn fly embryos within 12 hours were rinsed in 100% ethanol,
dechorionated in 10% bleach for 2 minutes, then immediately rinsed three times in
germ-free PBS. The germ-free embryos were transferred to autoclaved standard germ-
free molasses-cornmeal-yeast medium containing 1 mM kanamycin, 650 μM ampicillin

(61-238-RH, MediaTech), and 650 µM doxycycline (D9891, Sigma-Aldrich). Germ-free
flies were maintained on germ-free mediums with three antibiotics for four generations
and then maintained on the same medium without antibiotics for the next four
generations. Bacterial contamination of flies was monitored by homogenizing larvae in
germ-free PBS. Aspirate supernatant after brief centrifugation and monitoring the
bacterial growth on DeMan, Rogosa and Sharpe (MRS)-agar plates(Lekova, 2017).

Diets Preparation and Sample collection

Diets preparation followed the protocols below. Normal chow diets: 64.67g of corn meal,
27.1g of dry yeast, 8g of agar, and 61.6ml of molasses in 1L ddH$_2$O. For high protein
diets: 64.67g of corn meal, 150g of dry yeast, 8g of agar, and 61.6ml of molasses in 1L
of ddH$_2$O. For high sugar diets: 64.67g of corn meal, 27.1g of dry yeast, 8g of agar,
61.6ml of molasses, and 243g of sucrose in 1L of ddH$_2$O(Bedont et al., 2021; Morris et
al., 2012). After autoclaving the diets at high temperature, temperature was allowed to
drop to 40°C and then 10.17ml of 20% Tegosept was added to each 1L diet and mixed
thoroughly. Germ-free conditions were maintained throughout. For all treatments, 5-7
day old adults laid eggs on their respective diets (is031 and per^{01} flies were fed normal
chow diets, while is031 flies were fed high protein or high sugar diets). The subsequent
generation of newly hatched adults continued to feed on their respective diets for 3 days.
Then, newly hatched female adult flies aged 5-7 days were separated by gender and
subjected to the feeding paradigm in the presence of light-dark cycles beginning on day
5. In the timed feeding experiment, both ad libitum (AF) flies and those subjected to
timed feeding (TF) were transferred to germ-free medium vials at zeitgeber time 0 (ZT0),
which corresponds to the actual time of 9:00 am. They were then relocated into either a
new germ-free medium or a 1.1% agar vial at ZT10 (7pm) for a 14-hour fast. In the ad
lib feeding experiment, per^{01} flies were fed normal chow diets, while iso31 flies received
high protein and high sugar diets. AF flies were maintained in germ-free medium vials at
all times, with germ-free fresh diets replaced according to the timed feeding schedule.
Fly female guts were dissected on the fifth day at ZT0, ZT6, ZT12, and ZT18 within a
12:12 light-dark (LD) cycle. This dissection took place after the flies had been
continuously fed for 4 days under ad lib or timed feeding conditions. For each condition,
4 repeat samples were collected both from microbiome-containing and germ-free
female flies (>180 guts per sample, which corresponds to at least 10 mg for each
sample). All experimental processes were carried out on ice, and samples quickly
transferred to dry ice for storage after each sample was collected.

Metabolite profiling

Primary metabolites: Samples were extracted using the Matyash extraction buffer
which includes MTBE, MeOH, and H$_2$O. The organic (upper) phase was dried down and
submitted for resuspension and injection onto the LC while the aqueous (bottom) phase
was dried down and submitted to derivatization for GC. 10 ul of Methoxamine in pyridine
was added to the aqueous phase, after which it was shaken at 30 C for 1.5 hours. Then
91 uL of MSTFA + FAMEs was added to each sample and shaking was continued at 37
C for 0.5 hours to finish derivatization. Samples were then vialed, capped, and injected

onto the instrument. We use a 7890A GC coupled with a LECO TOF. 0.5 uL of derivatized sample is injected using a splitless method onto a RESTEK RTX-5SIL MS column with an Intergra-Guard at 275C with a helium flow of 1 mL/min. The GC oven is set to hold at 50C for 1 min then ramped at 20C/min to 330C and held for 5 min. The transferline is set to 280C while the EI ion source is set to 250C. The Mass spec parameters collect data from 85m/z to 500m/z at an acquisition rate of 17 spectra/sec.

Lipids: Samples were extracted using the Matyash extraction procedure, which includes MTBE, MeOH, and H_2O. The organic (upper) phase was dried down and submitted for resuspension and injection onto the LC while the aqueous (bottom) phase was dried down and submitted to derivatization for GC. It was resuspended with 110 uL of a solution of 9:1 methanol: toluene and 50 ng/mL CUDA. This was then shaken for 20 seconds, sonicated for 5 minutes at room temperature, and then centrifuged for 2 minutes at 16100 rcf. The samples were then aliquoted into three parts. 33 uL were aliquoted into each of two vials with a 50 uL glass insert for positive and negative mode lipidomics. The last part was aliquoted into an eppendorf tube to be used as a pool. The samples are then loaded up on an Agilent 1290 Infinity LC stack. The positive mode was run on an Agilent 6530 with a scan range of m/z 120-1200 Da with an acquisition speed of 2 spectra/s. Positive mode has between 0.5 and 2 uL injected onto an Acquity Premier BEH C18 1.7 µm, 2.1 x 50 mm Column. The gradient consists of two mobile phases, of which only B is controlled by the software as follows: 0 min 15% (B), 0.75 min 30% (B), 0.98 min 48% (B), 4.00 min 82% (B), 4.13-4.50 min 99% (B), 4.58-5.50 min 15% (B) with a flow rate of 0.8 mL/min. The other sample aliquot was run in negative mode, which was run on Agilent 1290 Infinity LC stack, and injected on the same column, with the same gradient and using an Agilent 6546 QTOF mass spec. The acquisition rate was 2 spectra/s with a scan range of m/z 60-1200 Da. The mass resolution for the Agilent 6530 is 10,000 for ESI (+) and 30,000 for ESI (-) for the Agilent 6546.

Biogenic Amines: Sample extraction for biogenic amines is a Liquid-Liquid extraction using the Matyash extraction containing MTBE, Methanol and Water and creating a biphasic partition. The polar phase is then dried down to completeness and run on a Waters Premier Acquity BEH Amide column. A short 4-minute Liquid Chromatography method is used for separation of polar metabolites from a starting condition of 100% LCMS H_2O with 10 mM ammonium formate and 0.125% formic acid to an end condition of 100% ACN: H_2O 95:5 (v/v) with 10 mM ammonium formate and 0.125% formic acid. A Sciex Triple-ToF scans from 50-1500 m/z with MS/MS collection from 40-1000 selecting from the top 5 ions per cycle.

Excreta quantification (Ex-Q) assay and feeding time course

For 24 h feeding we used the Ex-Q assay. 15 females were pre-fed on the three experimental diets described above for three days. On day 4 flies were transferred to the same food supplemented with 2.5mg/mL of FD&C blue FCF brilliant blue dye for 48 h. Flies were switched back to the specific diet without the dye for an extra 24h. Excrement was suspended into 150uL of MilliQ water. Absorbance was measured at

620 nm. Raw data was divided by the number of flies and further normalized to the minimum value of the entire dataset. For time courses females were pre-fed on the three experimental diets described above for three days. On day 4, groups of three flies were transferred to the same food supplemented with 2.5mg/mL of FD&C blue FCF brilliant blue dye for 4 h (ie. for the ZT2 collection, flies were transferred to the blue food at ZT22 and collected at ZT2). Flies were collected and homogenized using a Tissue Lyser in 200 ul of PBS. After centrifugation to remove debris, absorbance was measured at 620 nm. Data was divided by the number of flies and further normalized to the minimum of each dataset, high protein and high sugar separately.

Data processing and enrichment analysis

After acquisition primary metabolomics data is converted from LECO peg files to NetCDF files for processing. Files are submitted to in-house data processor BinBase for annotation and a report is exported into an excel file. In excel data is curated by blank reduction with a fold change of 3 or lower by using the max peak height from samples/ average blank peak height. After reduction peak heights are normalized by sum normalization. This is done using the un-normalized peak height of the feature divided by the mTIC (sum of the annotated metabolites) of the respective sample multiplied by the average mTIC of the samples. Normalized peak heights are then reported in an excel document. Chromatograms first undergo a quality control check in which internal standards are examined for consistency of peak height and retention time. Raw data files are then processed using an updated version of MS-DIAL software which identifies and aligns peaks and then annotates peaks using both an in-house mzRT library and MS/MS spectral matching with NIST/MoNA libraries. All MS/MS annotations are then manually curated by a lab member to ensure that only high-quality compound identifications are included in the final report. Enrichment analysis was conducted through the online software MetaboAnalyst 5.0.

Statistical analysis

Statistical details of experiments can be found in the figure legends. Circadian statistical analysis was performed in R using JTK_CYCLEv3.1. JTK_P value < 0.05 and JTK BH.Q < 0.2 deem as cycling metabolites(Pang et al., 2022).

Results

The microbiome enhances gut metabolite cycling

We showed previously that the *Drosophila* gut microbiome does not exhibit obvious oscillations in a 12:12 h LD cycle, but it significantly reduces the number and strength of host transcript cycling (Zhang et al., 2023). To investigate the role of the microbiome in gut metabolite cycling, we assayed metabolite cycling at different times of day in microbiome-containing (AM) and germ-free (AS) fed *ad libitum* (Figure. 1A). Metabolic

profiling was performed separately to characterize primary metabolites, lipids and biogenic amines. Principal component analysis (PCA) of the total expressed primary metabolites did not show significant separation between AM and AS flies at any time of day, indicating that this class of molecules is not changed by loss of the microbiome. However, a clear separation between the two groups was observed for lipids and biogenic amines (Figure. S1A-C).

Given our transcriptomic results we were expecting that the gut microbiome would reduce the number of cycling metabolites, however, we observed the opposite. JTK cycling analysis of the three different types of metabolites revealed that the likelihood of oscillations increased in AM flies as compared to AS flies. 26.9% (227 of 843) primary metabolites were rhythmic in AM flies as compared to 20.3% (171 of 843) in AS flies. A further reduction in AS was observed for lipids whose rhythms dropped from 29.5% (1329 of 4510) cycling in AM to 14.4% (648 of 4510) in AS. Similarly, the proportion of biogenic amines cycling decreased from 39.0% (1689 of 4330) to 24.7% (1068 of 4330) (JTK_P value < 0.05) (Figure. 1B, D, F, H, J and L). In addition, only about half of the cycling metabolites in germ-free flies are shared with microbiome containing flies (Figure S2). Consistent with the role of the microbiome in favoring metabolic rhythms, lipids and biogenic amines that cycle in both AM and AS flies have a higher amplitude in AM flies (Figure S3 A-C). Although the microbiome did not significantly change the overall distribution of phases for the cycling primary metabolites and lipids, it had a substantial effect on the phase of biogenic amine oscillations that shifted from peaking in the middle of the day (ZT6) in the AM flies to a peak in the early morning (ZT0) in AS flies (Figure. 1B, D, F, H, J and L). Morning peaks were also noted for lipids under almost all conditions, which may be driven by the morning peak of feeding.

Analysis of the abundance of cycling metabolites in AM, compared to their non-cycling counterparts in AS, also revealed that lipids and biogenic amines had higher expression levels in AM flies compared to AS flies. However, there was no significant difference between AM and AS flies in abundance of primary metabolites (Figure. S4A, E and I). Likewise, the overall abundance, regardless of cycling, of lipids and biogenic amines was higher in AM flies but primary metabolites were equivalent in AS and AM (Figure. S5A-C).

Overall, this indicates that the microbiome increases the cycling of all metabolites in the gut but has more dramatic effects on lipids and biogenic amines.

The microbiome has less effect on metabolite cycling under timed feeding conditions

Timed feeding affects host metabolism and benefits host health, according to several studies (Chaix et al., 2019; Gill et al., 2015; Lundell et al., 2020). In this study, we aimed to determine how microbiome-mediated metabolite cycling interacts with a timed feeding paradigm in which flies are provided with food only from ZT0 to ZT10. Similarly to *ad libitum* conditions, PCA of the total expressed primary metabolites showed only

minor separation between the four groups— AM, AS, microbiome-containing flies subjected to timed feeding (TM) and germ-free flies subjected to timed feeding (TS)—but a clear separation was observed for lipids and biogenic amines just as seen for AM and AS flies (Figure S1A-C). Consistent with the results of our previous study on transcript cycling, we found that timed feeding significantly increased the number of metabolites cycling in germ-free and microbiome-containing flies (Figure 1B-M, Figure S6A-D and Figure S2B,C,E,F,H and I).

Time restricted feeding coordinated the phase of all three-metabolite classes with TM and TS flies displaying very similar phase distributions. Interestingly, primary metabolites peaked around a single time of day whilst lipids and biogenic amines displayed two distinct phases in timed feeding conditions (Figure 1B-M). Moreover, the observation that TM and TS flies display similar metabolite phases differs from what happens in ad lib conditions in which the phase of biogenic amines and lipids changes in the absence of a microbiome. In particular, metabolites peaking at ZT9 increased with TF in all metabolite groups. This could suggest an anticipation of sleep at the end of the feeding window. Interestingly, a smaller proportion of cyclers is specific for the Ad libitum condition when compared to the TF in both sterile and microbiome containing flies (Figure S2).

Analysis of abundance of cycling metabolites under ad lib feeding and timed feeding conditions revealed that, in general, rhythmic metabolites tend to have higher overall expression levels compared to non-rhythmic metabolites, and this phenomenon is more prominent in the case of cycling biogenic amines (Figure. S4A-L). Additionally, we also looked at the amplitude of metabolites that cycle in both microbiome containing and microbiome free flies under the two feeding paradigms. In this case, the three metabolite groups did not always show a larger amplitude in the timed fed groups, which slightly contradicts the potential benefits of this type of intervention (Figure. S3D-I).

To further determine whether timed feeding modulates the effect of the microbiome, we compared the number of metabolites cycling with TF in AS and AM flies and found that loss of the microbiome had a larger effect in ad lib fed flies. This was reflected in the ratios depicting metabolite cycling in microbiome-containing versus germ-free flies, such that the AM/AS ratio was higher than the TM/TS for lipids and biogenic amines, although not for primary metabolites (Figure 1N-P). Moreover, the TS/AS ratio for lipids and biogenic amines was also higher than TM/AM (Figure, S1D-F). These data suggest that loss of the microbiome has less of an effect on timed fed flies than on ad lib fed, and, conversely, timed feeding has less effect on microbiome-containing flies. Thus, time restricted feeding improves cycling and it may compensate for the lack of a microbiome.

Effects of the microbiome on the cycling metabolome are modulated by the circadian clock.

To determine whether circadian clocks play a role in the enhanced metabolite cycling in microbiome-containing flies, we next subjected per^{01} flies with (per^{01}-AM) and without (per^{01}-AS) a microbiome to ad lib feeding conditions and examined the cycling of metabolites in their guts (Figure 2A). The results showed that the PCA for the three metabolite classes differentiated per^{01}-AM and per^{01}-AS flies (Figure S7A-C). Surprisingly, loss of the microbiome increased metabolite cycling in per^{01} flies, not for primary metabolites, but increasing the proportion of lipids cycling from 17.8% to 22.2% and almost duplicating the proportion of biogenic amines cycling from 18% in per^{01}-AM to 30.5% cycling features in per^{01}-AS flies (Figure 2B, D, F). Moreover, all cycling metabolites exhibited completely opposite phases in per^{01}-AM relative to per^{01}-AS flies (Figure 2C, E and G). Phases also changed for cycling metabolites shared between AM and AS (Figure S8A-C). Along these lines, we found an even lower overlap of cycling metabolites between AM and AS in per^{01} flies (Figure S2) indicating that the clock modulates not just the phase but also the identity of cyclers.

Analysis of the abundance of cycling metabolites in per^{01} flies revealed that, in general, cycling lipids and biogenic amines have higher expression in AM compared with their non-cycling equivalents in AS, whereas primary metabolites show minor difference between the two groups (Figure S9A-C). This result is consistent with the comparison of abundance in microbiome-containing and germ-free iso31 flies.

These datasets suggest a complex interaction between the microbiome and the host clock. It was noticeable that lack of a host clock resulted in reduced metabolite rhythms in microbiome containing flies (Figure S6A, E and Figure S2A, D, G) but enhanced cycling in microbiome-free flies (Figure S6B, F and Figure S2J, K, L). And also caused profound changes in phase (Figure 5). Overall, it appears that the host clock stabilizes metabolite rhythms against changes in the microbiome.

The microbiome enhances metabolite cycling in the presence of high protein diets and decreases cycling with high sugar diets

Changes in diet composition can significantly affect gut microbiome rhythms (Flint, 2012; Frazier et al., 2022). Therefore, we also explored the cyclic metabolome of flies with or without a microbiome and fed on two different diets; a high sugar and a high protein diet (Figure 3A and Figure 4A). From the PCA analysis we noticed that high sugar diets trigger metabolic differences between flies that contain a microbiome and microbiome-free flies, however, high protein diets do not prompt such difference (Figure S7D-F and G-I).

On high protein diets, the lack of a microbiome reduced the number of rhythmic metabolites (Figure 3B, D, F). This was consistent with the role of the microbiome in promoting metabolic rhythms. However, the effects of a high-sugar diet diverged substantially from those of a high-protein diet. With the high sugar diet, rhythmic lipids and biogenic amines increased in germ-free flies, similar to what we observed in a clock mutant background (Figure 4B, D, F). Indeed, when the three metabolite groups were

analyzed together, flies on a high sugar diet showed enhanced cycling of the metabolome when the microbiome was absent (Figure S6I and J).

We also looked at phases of cycling metabolites. In flies fed a high protein diet, a predominant peak around ZT15 was observed for all metabolites regardless of the presence or absence of a microbiome (Figure 3C, E, G). This suggests that protein metabolism imposes a specific metabolic rhythm that is not dependent on the gut microbiome. In case of the high sugar diet, we noticed that the absence of microbes changed the phase distribution of both lipid and biogenic amines but not of primary metabolites (Figure 4C, E and G).

We also compared the dataset generated from flies fed a standard diet versus these nutrient specific paradigms. For flies with a normal microbiome, the fraction of cycling metabolites decreased from 33.5% on standard diet to 22.2% and 24.5% with a high protein and high sugar diet respectively (Figure S6A, G and I). Thus, overall metabolite cycling decreases on high protein and high sugar diets. But while high protein potentiates the effect of the microbiome in promoting cycling, high sugar suppresses it (Figure S6G-J and Figure S2A, D, G, M-R).

Overall, the results suggest that nutrient specific diets interact in different ways with the microbiome to regulate the cycling of gut metabolites.

Circadian clock, dietary composition and the microbiome determine the phase of cycling metabolite in the gut

We further explored how the phase of cycling is affected by the endogenous clock, the diet and the microbiome. The phase of cycling in per^{01} flies was different from that of wild type flies. The average phase of primary metabolites in wild type flies falls around ZT6 while per^{01} flies mainly have a peak phase near ZT3. Lipids lose a morning peak in the clock mutant and most dramatically, biogenic amines go from peaking in the middle of the day to the middle of the night (Figure 5A, B, left panels). This effect of the host clock on the phase of cycling metabolites was even stronger in microbiome free flies (Figure 5A, B, right panels). Phases were also different on the different diets in microbiome-containing flies, with just primary metabolites conserving a peak around ZT6 in both the standard and high sugar diet; in high protein these metabolites showed a peak around ZT15. Lipids exhibit a cycling peak primarily around ZT0 and ZT6 in standard diets. However, the phase shifts to around ZT6 and ZT15 for high protein diets, and to around ZT3 and ZT15 for high-sugar diets. Biogenic amines have a cycling phase that falls around ZT15 in both high-protein and high-sugar diets, while in standard diets, the peak occurs around ZT6 (Figure 5A, C, D left panels). Germ-free flies also show significant effects of diet composition on cycling phase (Figure 5).

Feeding rhythms have no influence on metabolite cycling
To determine whether metabolite cycling is affected by altered feeding, we investigated food intake at different times of day in different genotypes and under different conditions and diets. Given that much has already been published on timed feeding, and the fact

that it does not significantly affect food consumption(Gill et al., 2015; Villanueva et al.,
2019), we chose to investigate feeding rhythms of AM and AS iso31 and per^{01} flies on
different diets. The results showed that, compared to normal chow diets, high-protein
diets decreased food consumption, while high-sugar diets increased food consumption
(Figure S10A). Flies fed high-protein diets displayed two feeding peaks, one in the
morning and one in the evening, as we previously demonstrated with standard diets.
The feeding rhythm showed no significant difference between AM and AS flies, in both
iso31 and per^{01} strains (Figure S10B). However, high-sugar diets reduce the amplitude
of feeding rhythms and eliminate the evening feeding peak. Indeed, the feeding rhythm
nearly disappears in AS flies, regardless of genotype, on a high-sugar diet (Figure
S10C).

Amino acid metabolism is modulated by the microbiome, timed feeding and the
circadian clock.

The extensive metabolic profiling performed here indicates that the gut microbiome is a
central piece in generating daily cycles of metabolites and that depending on the host
circadian clock and the composition of the diet, the extent and directionality of that
modulation varies.

To get a further understanding of the biological implications of this cycling metabolic
datasets we performed pathway enrichment analysis. First, we explored the ad libitum
dataset in which we found that absence of a microbiome reduced metabolic rhythms
(Figure 1). To our surprise, despite the reduction in cycling metabolite number in flies
lacking a microbiome, the pathway enrichment identified numerous pathways as
significantly enriched in the AS only cycling group. These included biosynthesis of
branched (Valine, leucine, isoleucine) and aromatic (Phenylalanine, tyrosine and
tryptophan) amino acids as well as the aminoacyl-tRNA biosynthesis pathway that is
relevant for protein synthesis (Figure 6A). Another amino acid pathway, involving
alanine, aspartate and glutamate, was found significant in AS and AM cycling sets, with
different specific metabolites cycling in these datasets. No pathway related to amino
acids or protein metabolism was specific to the AM cycling metabolites suggesting that,
in general, the microbiome abrogates daily oscillations of protein anabolism.

We found that similar pathways were significant in the cycling dataset from microbiome-
containing flies subjected to a daytime restricted feeding paradigm versus ad libitum
feeding (Figure 6B). Additionally, we determined whether pathways affected by timed
feeding were different in flies with and without a microbiome and found that amino acid
pathways cycle significantly in both groups (Figure 6D). Overall, the data suggest that
amino acid and protein biosynthesis cycles are dampened by the gut microbiome but
potentiated by time restricted feeding and the latter has a larger effect in microbiome
containing flies, indicating a microbiome and TF interaction. An interesting point here is
how timed feeding influences the same amino acid pathways in microbiome-containing
and germ-free flies (e.g., valine, leucine, and isoleucine biosynthesis) but sometimes
through distinct cycling metabolites. For instance, threonine, leucine, isoleucine, 4-

Methyl-2-oxovaleric Acid and valine in TM (Figure S11, Tables S1-21). In TS, cycling metabolites in this pathway are isoleucine and valine. Likewise, specific metabolites of a pathway showed differential regulation in AS and AM flies. In AS on a high sugar diet, L-threonine and leucine cycle. Another amino acid, 4-aminobutyric acid, cycles in both AM and AS flies (Figure S11, Tables S1-21). An acetylated derivative of a natural essential amino acid, N-acetyl-l-leucine, uniquely cycles in AS flies but not in AM flies. On the other hand, the TCA cycle appeared as a shared cycling category in all comparisons (AM vs AS, Figure 6A; AM vs TM, Figure 6B, AM vs per01 AM, Figure 6E), suggesting consistency in the rhythmicity of this pathway. Lastly, we aimed to identify pathways whose cycles were exclusive to microbiome containing flies. We found that alpha-linoleic metabolism cycled significantly in AM flies as compared to AS flies (Figure 6A). This pathway cycled significantly in AM and TM flies (Figure 6B), indicating that its cycling is not dramatically influenced by timed feeding and may be largely dependent on the microbiome. However, in the absence of a microbiome, timed feeding seems to drive cycling of this pathway (see comparison of AS and TS in Figure 6C). In addition, cycling of this pathway requires a functional circadian clock, as it is lost in per^{01}-AM flies. Thus cycling of alpha-linoleic metabolism depends upon the microbiome and the clock and can also be driven by timed feeding.

Consistent with the results from flies fed a normal chow diet, we observed that germ-free flies fed a high-sugar diet showed an enrichment of amino acid-related metabolism in the absence of a microbiome (Figure 6F), which indicates that sugar content did not alter the functional category of cycling metabolites by KEGG analysis. Pathway analysis of cycling metabolites under high protein conditions was not durable due to the reduced number of features.

Discussion

Our previous study reported that the microbiome dampens transcript cycling in the gut (Zhang et al., 2023). By contrast, we now show that the microbiome increases the number of cycling metabolites. At first glance, the two results appear contradictory because enhanced cycling gene expression would appear to be a prerequisite for producing large amounts of metabolites in an oscillatory manner. However, gut microbiomes can effectively provide the host with nutrients and promote metabolite absorption and they may do so in a rhythmic fashion (Flint, 2012; Wang et al., 2017). Thus, in the presence of a microbiome, the gut may not need to drive cycling of its own genes to promote metabolic oscillations since that might be facilitated by a healthy microbiome. In contrast, in germ-free flies the host might need rhythmic expression of metabolism-related genes to achieve nutritional homeostasis. We report that the cycling metabolites in germ-free flies are extensively enriched in amino acid metabolism. This may compensate for the loss of metabolic properties that would normally be conferred by the microbiome. Interestingly, this pathway is also a major target of timed feeding.

In this study, we first explored the cycling of metabolites in wild flies (iso31) under ad lib and timed feeding conditions, both in germ-free and microbiome-containing flies (Zhang et al., 2023). To better understand the role of the circadian clock and the effect of

different diet compositions on metabolite cycling, we further investigated the microbiome-dependence of metabolite oscillations in per^{01} flies fed on normal chow diets and iso31 wild type flies fed on high protein and high sugar diets, respectively. Although these two sets of studies revealed similar numbers of named metabolites, the total numbers of metabolites exhibited some differences. For instance, the first study had 193/843, 456/4510 and 621/4330 named/total primary metabolites, lipids, and biogenic amines, respectively; in the second study, these numbers were 144/623, 513/2245, and 530/2791 respectively. We acknowledge that due to this fact, direct comparison of some of the datasets is not possible. As a strategy to eliminate the influence of different run conditions on the results, we also investigated metabolite cycling profiles using the shared named metabolites across the two batches and found that the results were highly consistent (data not shown). Although not ideal, this suggest that an overall comparison of all the samples included here is appropriate.

Some empirical evidence has found that timed feeding has a beneficial influence on metabolic health (Gill et al., 2015; Lundell et al., 2020; Villanueva et al., 2019). As reported here for metabolite cycling, our previous study found that timed feeding not only enhances gene cycling but also changes the phase of cycling (Zhang et al., 2023). Metabolite cycling was further measured under ad lib and timed feeding conditions in microbiome-containing and germ-free flies. In line with the transcript results, timed feeding significantly enhanced metabolite cycling and changed the phase regardless of the presence of a microbiome. However, the microbiome has less effect in timed fed flies, and the reverse was also true, timed feeding had less effect on flies containing a microbiome relative to germ-free flies. It is possible that the microbiome buffers the cyclic metabolome against strong external zeitgebers such as the time of feeding. Indeed, we previously showed that the gut microbiome stabilizes transcript cycling in the gut such that the phase is not rapidly shifted by acute changes in the light: dark cycle (Zhang et al., 2023). Alternatively, given that the presence of a microbiome also has less effect in a timed feeding paradigm, it is possible that TF and the microbiome have overlapping impact on metabolite cycling. On the other hand, oscillations in amino acid metabolism are characteristic of a timed feeding paradigm, as reported here and by others (Lundell et al., 2020), and are enhanced in germ-free flies as opposed to microbiome-containing flies. Perhaps these oscillations help the host cope with the lack of microbial driven cycles. How the microbiome helps host cycling remains an open question. We did not detect rhythms of microbial species in *Drosophila*, but it is possible that rhythms of feeding drive rhythms of microbiome abundance in the gut (Zhang et al., 2023), given that food is the major source of microbes for flies.

Keeping metabolic rhythms synchronized with the external environment is an important way to maintain optimal fitness for the host. The synchronization is mainly achieved through the circadian clock (Castanon-Cervantes et al., 2010; Dunlap, 1999; Roenneberg and Merrow, 2016), and so mutations in core clock genes can lead to profound metabolic abnormalities(Marcheva et al., 2010; Rudic et al., 2004). To investigate whether the microbiome-dependent increase in the number of cycling metabolites depends on the endogenous circadian clock, metabolite cycling was further investigated in microbiome-containing and germ-free per^{01} flies. Interestingly, the

number of cycling metabolites was significantly decreased in per^{01} AM flies compared
with the per^{01} AS flies. Also, unexpectedly, compared with per^{01} AS flies, cycling
metabolites exhibit an opposite cycling phase in per^{01} AM flies. On the other hand, wild
type flies show only a minor change in the phase of cycling primary metabolites with the
loss of a microbiome. The cycling phase of biogenic amines shows a significant
difference between AM and AS wild type flies but the difference is larger between AM
and AS per^{01} flies. We speculate that the host circadian clock prevents microbial
changes from influencing the phase of metabolite cycling. Thus, in clock mutant flies,
loss of the microbiome has a dramatic effect on phase. Cycling of metabolites involved
in amino acid metabolism occurs in the face of perturbations to the normal e.g. in the
absence of a clock (per^{01} AM) or in wild type flies devoid of a microbiome. This likely
reflects compensation for a function normally served by the clock and microbiome (e.g.
perhaps the clock drives cycling of a microbiome component) and helps to maintain
effective amino acid/protein metabolism. Apart from the influence of the endogenous
clock, investigation of specific cycling metabolites in the amino acid biosynthesis
pathway indicates that although both the microbiome and timed feeding affect this
pathway, they modulate different cycling metabolites - and these processes are also
influenced by diet composition. This suggests that the microbiome, timed feeding,
circadian clock, and diet can utilize distinct cycling metabolites to fulfill similar
metabolism needs. At the same time, the diet may affect the microbiome.

Mice fed on western diets which contain high sugar amounts, show a significant
reduction in the cycling of gut microbiomes, and it has been argued that the changed
feeding rhythm may be the reason for this phenomenon (Frazier et al., 2022; Kohsaka
et al., 2007). In the current study, we found that the microbiome suppresses the number
of cycling metabolites and substantially affects the phase when flies are fed high sugar
diet. Conversely, the microbiome increases the number of cycling metabolites but does
not change the cycling phase when flies are fed high protein diets. High sugar diet also
resulted in phenotypic differences in the guts of AM and AS flies. In general, on normal
chow or high protein diets, AM fly guts had good plasticity and were larger in size than
those from AS flies. However, flies fed a high-sugar diet had smaller gut sizes than flies
fed a high-protein or normal chow diets, and the microbiome-mediated gut plasticity
difference between AM and AS conditions disappeared (data not shown). Moreover,
metabolite abundance analysis showed higher expression of lipids and biogenic amines
in AM flies than AS flies when flies were fed on normal chow or high protein diets, but
on a high sugar diet AS flies had higher expression of lipids and biogenic amines,
although the difference in biogenic amines was not significant (Figure S5). All these
findings suggest that high-sugar diets disrupt microbiome function, and this is likely the
cause of the increased number of cycling metabolites in AS flies compared to AM flies.
While we are attributing differences on high sugar and high protein diets to food
composition, we acknowledge that we cannot exclude an effect of altered caloric intake
as this was difficult to control for or evaluate.

Changes in the feeding rhythm in response to different diet compositions could
contribute to the observed variations in metabolite cycling phenotypes, as seen in mice
fed different diets (Kohsaka et al., 2007). To address some of these questions we

measured food consumption across 24 h in flies on different diets, with and without a microbiome. Results indicate that feeding rhythms are not different between AM and AS flies fed on either normal chow or high-protein diets(Zhang et al., 2023) . Given that AM flies show increased metabolite cycling compared to AS flies on normal chow diets, and AS flies show more cycling on high protein diets, these data do not support control of metabolite cycling by feeding rhythms . Furthermore, the feeding rhythm nearly disappears in AS flies fed high-sugar diets, regardless of whether they are iso31 or per^{01} flies, but AS flies increase metabolite cycling compared to AM flies on high-sugar diets. Thus, there seems to be no obvious link between feeding and metabolite rhythms. Caloric content, rather than nutrition (protein/carbohydrate), may be a significant factor affecting metabolite cycling, but this was difficult to control with the diets we used. We note that flies ate more on a high sugar diet and less on a high protein diet, but whether this accounted for loss of the feeding rhythm or was a consequence of it is unclear. Other factors such as locomotor rhythms and sleep are also likely contributors to metabolite cycling and potential subjects of future study. For instance, sleep reduces metabolic rate, which could affect metabolite levels(Stahl et al., 2017).

Based on these extensive metabolic profiling data, we propose that an interplay of the microbiome, the circadian clock and food intake maintain metabolic homeostasis in the gut. With respect to food, the timing of intake as well as the composition are important (Figure 7). Interestingly, the microbiome and time restricted feeding paradigms can compensate for each other to some extent, suggesting that different strategies can be leveraged to serve organismal health.

Acknowledgements: This work was supported by a grant from the Volkwagen Stiftung (Life?). YZ was supported by the Opening Projects of Shanghai Key Laboratory of Chemical Biology and the National Natural Science Foundation of China (31972308). AS is an Investigator of the HHMI.

References

Albrecht, U. (2012). Timing to perfection: the biology of central and peripheral circadian clocks. Neuron *74*, 246-260.

Bedont, J., Kolesnik, A., Malik, D., Weljie, A., and Sehgal, A. (2021). Polyamine elevation and nitrogen stress are toxic hallmarks of chronic sleep loss in Drosophila melanogaster. bioRxiv, 2021.2010. 2001.462746.

Castanon-Cervantes, O., Wu, M., Ehlen, J.C., Paul, K., Gamble, K.L., Johnson, R.L., Besing, R.C., Menaker, M., Gewirtz, A.T., and Davidson, A.J. (2010). Dysregulation of inflammatory responses by chronic circadian disruption. The Journal of Immunology *185*, 5796-5805.

Chaix, A., Lin, T., Le, H.D., Chang, M.W., and Panda, S. (2019). Time-restricted feeding prevents obesity and metabolic syndrome in mice lacking a circadian clock. Cell metabolism *29*, 303-319. e304.

Dunlap, J.C. (1999). Molecular bases for circadian clocks. Cell *96*, 271-290.

Feltzin, V., Wan, K.H., Celniker, S.E., and Bonini, N.M. (2019). Role and impact of the gut microbiota in a Drosophila model for parkinsonism. BioRxiv, 718825.

Flint, H.J. (2012). The impact of nutrition on the human microbiome. Nutrition reviews *70*, S10-S13.

Frazier, K., Kambal, A., Zale, E.A., Pierre, J.F., Hubert, N., Miyoshi, S., Miyoshi, J., Ringus, D.L., Harris, D.,
and Yang, K. (2022). High-fat diet disrupts REG3γ and gut microbial rhythms promoting metabolic
dysfunction. Cell Host & Microbe 30, 809-823. e806.
Gill, S., Le, H.D., Melkani, G.C., and Panda, S. (2015). Time-restricted feeding attenuates age-related
cardiac decline in Drosophila. Science 347, 1265-1269.
Hariri, N., and Thibault, L. (2010). High-fat diet-induced obesity in animal models. Nutrition research
reviews 23, 270-299.
Ko, G.-J., Rhee, C.M., Kalantar-Zadeh, K., and Joshi, S. (2020). The effects of high-protein diets on kidney
health and longevity. Journal of the American Society of Nephrology 31, 1667-1679.
Kohsaka, A., Laposky, A.D., Ramsey, K.M., Estrada, C., Joshu, C., Kobayashi, Y., Turek, F.W., and Bass, J.
(2007). High-fat diet disrupts behavioral and molecular circadian rhythms in mice. Cell metabolism 6,
414-421.
Lekova, V. (2017). Role and Impact of the Gut Microbiota in a Drosophila Model for Parkinsonism
(University of Pennsylvania).
Liang, X., Bushman, F.D., and FitzGerald, G.A. (2015). Rhythmicity of the intestinal microbiota is
regulated by gender and the host circadian clock. Proceedings of the National Academy of Sciences 112,
10479-10484.
Lundell, L.S., Parr, E.B., Devlin, B.L., Ingerslev, L.R., Altıntaş, A., Sato, S., Sassone-Corsi, P., Barrès, R.,
Zierath, J.R., and Hawley, J.A. (2020). Time-restricted feeding alters lipid and amino acid metabolite
rhythmicity without perturbing clock gene expression. Nature communications 11, 4643.
Marcheva, B., Ramsey, K.M., Buhr, E.D., Kobayashi, Y., Su, H., Ko, C.H., Ivanova, G., Omura, C., Mo, S.,
and Vitaterna, M.H. (2010). Disruption of the clock components CLOCK and BMAL1 leads to
hypoinsulinaemia and diabetes. Nature 466, 627-631.
Morris, S.N.S., Coogan, C., Chamseddin, K., Fernandez-Kim, S.O., Kolli, S., Keller, J.N., and Bauer, J.H.
(2012). Development of diet-induced insulin resistance in adult Drosophila melanogaster. Biochimica et
Biophysica Acta (BBA)-Molecular Basis of Disease 1822, 1230-1237.
Na, J., Musselman, L.P., Pendse, J., Baranski, T.J., Bodmer, R., Ocorr, K., and Cagan, R. (2013). A
Drosophila model of high sugar diet-induced cardiomyopathy. PLoS genetics 9, e1003175.
Pang, Z., Zhou, G., Ewald, J., Chang, L., Hacariz, O., Basu, N., and Xia, J. (2022). Using MetaboAnalyst 5.0
for LC–HRMS spectra processing, multi-omics integration and covariate adjustment of global
metabolomics data. Nature protocols 17, 1735-1761.
Rijo-Ferreira, F., and Takahashi, J.S. (2019). Genomics of circadian rhythms in health and disease.
Genome medicine 11, 1-16.
Roenneberg, T., and Merrow, M. (2016). The circadian clock and human health. Current biology 26,
R432-R443.
Rudic, R.D., McNamara, P., Curtis, A.-M., Boston, R.C., Panda, S., Hogenesch, J.B., and FitzGerald, G.A.
(2004). BMAL1 and CLOCK, two essential components of the circadian clock, are involved in glucose
homeostasis. PLoS biology 2, e377.
Sehgal, A. (2017). Physiology flies with time. Cell 171, 1232-1235.
Seo, D.-o., O'Donnell, D., Jain, N., Ulrich, J.D., Herz, J., Li, Y., Lemieux, M., Cheng, J., Hu, H., and Serrano,
J.R. (2023). ApoE isoform–and microbiota-dependent progression of neurodegeneration in a mouse
model of tauopathy. Science 379, eadd1236.
Stahl, B.A., Slocumb, M.E., Chaitin, H., DiAngelo, J.R., and Keene, A.C. (2017). Sleep-dependent
modulation of metabolic rate in Drosophila. Sleep 40, zsx084.
Stephan, F.K. (2002). The "other" circadian system: food as a Zeitgeber. Journal of biological rhythms 17,
284-292.

Thaiss, C.A., Levy, M., Korem, T., Dohnalová, L., Shapiro, H., Jaitin, D.A., David, E., Winter, D.R., Gury-
BenAri, M., and Tatirovsky, E. (2016a). Microbiota diurnal rhythmicity programs host transcriptome
oscillations. Cell *167*, 1495-1510.
Thaiss, C.A., Levy, M., Korem, T., Dohnalová, L., Shapiro, H., Jaitin, D.A., David, E., Winter, D.R., Gury-
BenAri, M., and Tatirovsky, E. (2016b). Microbiota diurnal rhythmicity programs host transcriptome
oscillations. Cell *167*, 1495-1510. e1412.
Thaiss, C.A., Zeevi, D., Levy, M., Zilberman-Schapira, G., Suez, J., Tengeler, A.C., Abramson, L., Katz, M.N.,
Korem, T., and Zmora, N. (2014). Transkingdom control of microbiota diurnal oscillations promotes
metabolic homeostasis. Cell *159*, 514-529.
Thaiss, C.A., Zmora, N., Levy, M., and Elinav, E. (2016c). The microbiome and innate immunity. Nature
535, 65-74.
Turek, F.W., Joshu, C., Kohsaka, A., Lin, E., Ivanova, G., McDearmon, E., Laposky, A., Losee-Olson, S.,
Easton, A., and Jensen, D.R. (2005). Obesity and metabolic syndrome in circadian Clock mutant mice.
Science *308*, 1043-1045.
Villanueva, J.E., Livelo, C., Trujillo, A.S., Chandran, S., Woodworth, B., Andrade, L., Le, H.D., Manor, U.,
Panda, S., and Melkani, G.C. (2019). Time-restricted feeding restores muscle function in Drosophila
models of obesity and circadian-rhythm disruption. Nature communications *10*, 1-17.
Wang, Y., Kuang, Z., Yu, X., Ruhn, K.A., Kubo, M., and Hooper, L.V. (2017). The intestinal microbiota
regulates body composition through NFIL3 and the circadian clock. Science *357*, 912-916.
Zhang, Y., Li, Y., Barber, A.F., Noya, S.B., Williams, J.A., Li, F., Daniel, S.G., Bittinger, K., Fang, J., and
Sehgal, A. (2023). The microbiome stabilizes circadian rhythms in the gut. Proceedings of the National
Academy of Sciences *120*, e2217532120.

698
699
700 **Figures and legends**
701
702

Figure 1. The microbiome reduces the effect of timed feeding on the cycling of lipids and biogenic amines.

(A) Schematic showing the experimental protocol for 1) analysis of gut metabolite
cycling in microbiome-containing (AM) and sterile (AS) iso31 flies fed on normal chow
diets under ad lib and 2) timed feeding (TF) conditions, respectively.
(B-E) Pie chart depicting the fraction of cycling primary metabolites and polar histogram
plots of the peak phase for cycling primary metabolites in microbiome-containing and
sterile flies under ad lib and timed feeding conditions (AM, TM, AS and TS), respectively,
using a JTK_cycle cutoff of $P < 0.05$.
(F-I) Pie chart depicting the fraction of cycling lipids and polar histogram plots of the
peak phase for cycling lipids in AM, TM, AS and TS flies, respectively, using a
JTK_cycle cutoff of $P < 0.05$.
(J-M) Pie chart depicting the fraction of cycling biogenic amines and polar histogram
plots of the peak phase for cycling biogenic amines in AM, TM, AS and TS flies,
respectively, using a JTK_cycle cutoff of $P < 0.05$.
(N-P) The ratio of primary metabolites, lipids and biogenic amines that cycle in
microbiome-containing versus sterile flies under ad lib feeding (AM/AS) and timed
feeding (TM/TS) conditions, respectively.
722
723

Figure 2. The effect of the microbiome on metabolite cycling depends on a functional circadian clock.

(A) Schematic showing the experimental protocol for analysis of gut metabolite cycling in microbiome-containing (AM) and sterile (AS) *per01* flies fed on normal chow diets.

(B) Pie chart depicting the fraction of cycling primary metabolites, and (C) polar histogram plots of the peak phase for cycling primary metabolites in AM and AS *per01* flies fed on normal chow diets, using a JTK_cycle cutoff of P < 0.05.

(D) Pie chart depicting the fraction of cycling lipids and (E) polar histogram plots of the peak phase for cycling lipids in AM and AS *per01* flies fed on normal chow diets, using a JTK_cycle cutoff of P < 0.05.

(F) Pie chart depicting the fraction of cycling biogenic amines and (G) polar histogram plots of the peak phase for cycling biogenic amines in AM and AS *per01* flies fed on normal chow diets, using a JTK_cycle cutoff of P < 0.05.

Figure 3. The microbiome increases the number of cycling of metabolites in flies fed on high protein diets.

(A) Schematic showing the experimental protocol for analysis of gut metabolite cycling in microbiome-containing (AM) and sterile (AS) Iso31 flies fed on high protein diets.
(B) Pie chart depicting the fraction of cycling primary metabolites, and (C) polar histogram plots of the peak phase for oscillating primary metabolites in AM and AS iso31 flies fed on high protein diets, using a JTK_cycle cutoff of P < 0.05.
(D) Pie chart depicting the fraction of cycling lipids, and (E) polar histogram plots of the peak phase for oscillating lipids in AM and AS iso31 flies fed on high protein diets, using a JTK_cycle cutoff of P < 0.05.
(F) Pie chart depicting the fraction of cycling biogenic amines, and (G) polar histogram plots of the peak phase for oscillating biogenic amines in AM and AS iso31 flies fed on high protein diets, using a JTK_cycle cutoff of P < 0.05.

756
757

758
759
Figure 4. The microbiome decreases the number of cycling metabolites in iso31
flies fed on high sugar diets.

(A) Schematic showing the experimental protocol for analysis of gut metabolite cycling
in microbiome-containing (AM) and sterile (AS) Iso31 flies fed on high sugar diets.
(B) Pie chart depicting the fraction of cycling primary metabolites, and (C) polar
histogram plots of the peak phase for cycling primary metabolites in AM and AS iso31
flies fed on high sugar diets, using a JTK_cycle cutoff of $P < 0.05$.
(D) Pie chart depicting the fraction of cycling lipids, and (E) polar histogram plots of the
peak phase for cycling lipids in AM and AS iso31 flies fed on high sugar diets, using a
JTK_cycle cutoff of $P < 0.05$.
(F) Pie chart depicting the fraction of cycling biogenic amines and (G) polar histogram
plots of the peak phase for cycling biogenic amines in AM and AS iso31 flies fed on high
sugar diets, using a JTK_cycle cutoff of $P < 0.05$.
773
774

Figure 5. Effects of the circadian clock and dietary composition on the phase of metabolite cycling in microbiome-containing and sterile flies (phases are for cycling metabolites reported in Figure1-4).

(A-D) Polar histogram plots of the peak phase for oscillating primary metabolites, lipids, and biogenic amines, respectively, in microbiome-containing and sterile flies, using a JTK_cycle value of $P < 0.05$. A: iso31 flies fed on normal chow diets; B: per^{01} flies fed on normal chow diets; C: iso31 flies fed on high protein diets; D: iso31 flies fed on high sugar diets.

784
785
786

Figure 6. Amino acid metabolism is modulated by the microbiome, timed feeding
and the circadian clock.

(A) KEGG enrichment analysis of oscillating metabolites unique to AM flies, shared in
AM and AS flies and unique to AS flies. Data only show the P < 0.05 results.
(B) KEGG enrichment analysis of oscillating metabolites unique to AM flies, shared in
AM and TM flies and unique to TM flies. Data only show the P < 0.05 results.
(C) KEGG enrichment analysis of oscillating metabolites unique to AS flies, shared in
AS and TS flies and unique to TS flies. Data only show the P < 0.05 results.
(D) KEGG enrichment analysis of oscillating metabolites unique to TM flies, shared in
TM and TS flies and unique to TS flies. Data only show the P < 0.05 results.

(E) KEGG enrichment analysis of oscillating metabolites unique to *per*^{01} AM flies,
shared in AM and AS of *per*^{01} flies and unique in *per*^{01} AS flies. Data only show the P <
0.05 results.
(F) KEGG enrichment analysis of oscillating metabolites unique to high sugar fed AM
flies, shared in AM and AS of *iso31* flies fed on high sugar diets and unique in high
sugar fed AS flies. Data only show the P < 0.05 results.
804
805

806
807   **Figure 7. Model showing effects of the microbiome, timed feeding, the circadian**
808   **clock and diet composition on metabolite rhythms in the gut.**
809
810   The microbiome broadly increases metabolite rhythms on a normal chow diet and its
811   effects are enhanced by high protein diet and the circadian clock. Timed feeding
812   promotes cycling itself, but attenuates the effect of the microbiome on cycling, as does a
813   high sugar diet. Effects of each of these manipulations on phase are also indicated.
814

Supplement Figures

Figure S1. Timed feeding has less effect on cycling metabolites in microbiome-
containing flies.

(A-C) A: Principal component analysis (PCA) of primary metabolites, lipids and biogenic
amine abundance, respectively, at each time-point in a 12:12 LD cycle for AM, TM, AS,
and TS flies.
(D-F) The ratio of primary metabolites, lipids and biogenic amines that cycle in
microbiome-containing versus sterile flies under ad lib feeding (TS/AS) and timed
feeding (TM/AM) conditions, respectively.

Figure S2. The percentage of unique and shared cycling metabolites in different treatment groups.

(A-C) Venn diagram illustrates the percentage of unique and overlapping oscillating primary metabolites between AM and AS, AM and TM, AS and TS iso31 flies fed on normal chow diets; (D-F) Venn diagram illustrates the percentage of unique and overlapping oscillating lipids between AM and AS, AM and TM, AS and TS iso31 flies fed on normal chow diets; (G-I) Venn diagram illustrates the percentage of unique and overlapping oscillating biogenic amines between AM and AS, AM and TM, AS and TS iso31 flies fed on normal chow diets; (J-L) Venn diagram illustrates the percentage of unique and overlapping oscillating primary metabolites, lipids and biogenic amines between AM and AS per^{01} flies fed on normal chow diets; (M-O) Venn diagram illustrates the percentage of unique and overlapping oscillating primary metabolites, lipids and biogenic amines between AM and AS iso31 flies fed on high protein diets; (P-R) Venn diagram illustrates the percentage of unique and overlapping oscillating primary metabolites, lipids and biogenic amines between AM and AS iso31 flies fed on high sugar diets.

Figure S3. The amplitude of metabolite cycling is sometimes, but not always, increased by the presence of a microbiome or by timed feeding.

(A-C) The comparison of amplitudes for cycling primary metabolites, lipids and biogenic
amines, respectively, that overlap in AM and AS flies. Data are mean ± SEM, ****p <
0.0001 shown by Student's t test.
(D-F) The comparison of amplitudes for cycling primary metabolites, lipids and biogenic
amines, respectively, that overlap in AM and TM flies. Data are mean ± SEM, **p < 0.01
and ****p < 0.0001 shown by Student's t test.
(G-I) The comparison of amplitudes for cycling primary metabolites, lipids and biogenic
amines, respectively, that overlap in AS and TS flies. Data are mean ± SEM, ***p <
0.001 shown by Student's t test.
867

Primary metabolites

Lipids

Biogenic amines

Figure S4. Lipids and biogenic amines show higher abundance in microbiome-containing flies.

A-D: Average expression levels of gut primary metabolites that cycle in either microbiome-containing or sterile flies or both conditions (lost, shared and gained). The number indicates the average normalized counts in each condition for primary metabolites that cycle with $p < 0.05$, JTK_cutoff. Data are mean ± SEM, comparisons were with Student's t test. A: AS versus AM; B: TM versus AM; C: TS versus AS; D**:** TS versus TM.

E-H: Average expression levels of gut lipids that cycle in either microbiome-containing or sterile flies or both conditions (lost, shared and gained). The number indicates the average normalized counts in each condition for lipids that cycle with $p < 0.05$, JTK_cutoff. Data are mean ± SEM, comparisons were with Student's t test. E: AS versus AM; F: TM versus AM; G: TS versus AS; H**:** TS versus TM.

I-L: Average expression levels of gut biogenic amines that cycle in either microbiome-containing or sterile flies or both conditions (lost, shared and gained). The number indicates the average normalized counts in each condition for biogenic amines that cycle with $p < 0.05$, JTK_cutoff. Data are mean ± SEM, comparisons were with Student's t test. I: AS versus AM; J: TM versus AM; K: TS versus AS; L**:** TS versus TM.

Iso31 flies fed
normal chow diets

Per[01] flies fed
normal chow diets

Iso31 flies fed
high protein diets

Iso31 flies fed
high sugar diets

Figure S5. Abundance analysis of all metabolites in microbiome containing and sterile flies.

(A-C) Average expression levels of gut metabolites in microbiome-containing and sterile
flies. iso31 flies are fed on normal chow diets. A: AS versus AM for primary metabolites;
B: AS versus AM for lipids; C: AS versus AM for biogenic amines.
(D-F) Average expression levels of gut metabolites between microbiome-containing and
sterile flies. per^{01} flies are fed on normal chow diets. D: AS versus AM for primary
metabolites; E: AS versus AM for lipids; F: AS versus AM for biogenic amines.
(G-I) Average expression levels of gut metabolites between microbiome-containing and
sterile flies. iso31 flies are fed on high protein diets. G: AS versus AM for primary
metabolites; H: AS versus AM for lipids; I: AS versus AM for biogenic amines.
(J-L) Average expression levels of gut metabolites between microbiome-containing and
sterile flies iso31 flies are fed on high sugar diets. J: AS versus AM for primary
metabolites; K: AS versus AM for lipids; L: AS versus AM for biogenic amines.
904
905

906
907
908
Figure S6. Loss of the circadian clock or high protein or high sugar diet
decreases metabolite cycling in microbiome-containing flies.

(A,B) Pie charts depicting the fraction of metabolites cycling in AM and AS iso31 flies
fed on normal chow diets; (C,D) Pie chart depicting the fraction of metabolites cycling in
TM and TS iso31 flies fed on normal chow diets; (E,F) Pie charts depicting the fraction
of metabolites cycling in AM and AS per^{01} flies fed on normal chow diets; (G,H) Pie chart
depicting the fraction of metabolites cycling in AM and AS iso31 flies fed on high protein

diets; (I,J) Pie chart depicting the fraction of metabolites cycling in AM and AS iso31 flies
fed on high sugar diets, using a JTK_cycle cutoff of P < 0.05.

**Figure S7. Principal component analysis (PCA) of total expressed metabolites in
per01 flies fed on normal chow diets, iso31 flies fed on high protein and high sugar
diets.**

(A-C) Principal component analysis (PCA) of metabolite abundance at each time-point
in a 12:12 LD cycle for *per01* AM and AS flies fed on normal chow diets.
(D-F) Principal component analysis (PCA) of metabolite abundance at each time-point
in a 12:12 LD cycle for iso31 AM and AS flies fed on high protein diets
(G-I) Principal component analysis (PCA) of metabolite abundance at each time-point in
a 12:12 LD cycle iso31 AM and AS flies fed on high sugar diets.

934
935
936

Figure S8. Phase changes for cycling metabolites shared between *per^{01}*-AM and *per^{01}*-AS.

(A-C) Polar histogram plots of the peak phase for shared oscillating primary, lipids, and biogenic amines, respectively, in *per^{01}* microbiome-containing and *per^{01}* sterile flies, using a JTK_cycle value of P < 0.05.

944
945

Figure S9. Effect of the microbiome on expression levels of cycling metabolites in clock mutants and under different dietary conditions.

A-C: Average expression levels of gut metabolites that cycle in microbiome-containing versus sterile *per01* flies fed on normal chow diets. The middle group in each panel shows metabolites that cycle under both conditions, while those on either side indicate metabolites that cycle in only one condition. The number indicates the average normalized counts in each condition for metabolites that cycle with p < 0.05, JTK_cycle. Data are mean ± SEM, comparisons were with Student's t test. A: AS versus AM for primary metabolites; B: AS versus AM for lipids; C: AS versus AM for biogenic amines.

D-F: Average expression levels of gut metabolites that cycle in microbiome-containing versus sterile iso31 flies fed on high protein diets. The middle group in each panel shows metabolites that cycle under both conditions, while those on either side indicate metabolites that cycle in only one condition. The number indicates the average normalized counts in each condition for metabolites that cycle with p < 0.05, JTK_cycle. Data are mean ± SEM, comparisons were with student's t test. D: AS versus AM for primary metabolites; E: AS versus AM for lipids; F: AS versus AM for biogenic amines.

G-I: Average expression levels of gut metabolites that cycle in microbiome-containing versus sterile flies fed on high sugar diets. The middle group in each panel shows metabolites that cycle under both conditions, while those on either side indicate metabolites that cycle in only one condition. The number indicates the average normalized counts in each condition for metabolites that cycle with p < 0.05, JTK_cycle.

968 Data are mean ± SEM, comparisons were with Student's t test. G: AS versus AM for
969 primary metabolites; H: AS versus AM for lipids; I: AS versus AM for biogenic amines.
970

**Figure S10. Feeding rhythms do not show an obvious link with metabolite cycling
in different genotypes and under different conditions.**

(A). 24h feeding in iso31 and per^{01} flies with or without a microbiome on three different
diets. Each data point represents feeding of groups of 15 flies. (B and C). Time course
of food intake in iso31 and per^{01} flies with or without a microbiome in the high protein (B)
and high sugar (C) condition. Each data point represents a 4h feeding window ending at
the time of day specified in the x axis. 3 flies per data point. $P < 0.05$ by JTK_cycle was
used as the cutoff for cycling.

is031 flies fed standard diets

is031 flies fed standard diets

is031 flies fed standard diets

is031 flies fed standard diets

per^{01} flies fed standard diets

is031 flies fed high protein diets

is031 flies fed high sugar diets

Figure S11. Examples of cycling metabolites in iso31 and *per^{01}* flies maintained
under different conditions

Conditions include iso31 or *per^{01}* flies maintained with a microbiome or in germfree
conditions on standard diet as well as iso31 flies on high protein or high sugar.
Metabolites that cycle under two different conditions (common) or are unique to one
condition (unique) are indicated. A: iso31 fed standard diets in AM and AS flies, B: iso31
fed standard diets in AM and TM flies, C: iso31 fed standard diets in AS and TS flies, D:
iso31 fed standard diets in TM and TS flies, E: *per^{01}* fed standard diets in AM and AS
flies, F: iso31 fed high protein diets in AM and AS flies, G: iso31 fed high sugar diets in
AM and AS flies. (P: JTK p-value A: JTK amplitude). P < 0.05 by JTK_cycle was used
as the cutoff for cycling.

994