The Early Dodder Gets the Host: Decoding the Coiling Patterns of *Cuscuta campestris* with

Automated Image Processing

Max Bentelspacher[1, a], Erik J. Amézquita[1,2, a, b], Supral Adhikari[1], Jaime Barros[1], So-Yon Park[1, b]

[1]Division of Plant Science and Technology, University of Missouri, Columbia, MO, 65211, USA

[2]Department of Mathematics, University of Missouri, Columbia, MO, 65211, USA

[a] These authors contributed equally to this work.

[b] Corresponding authors: So-Yon Park (parksoy@missoui.edu) and Erik J. Amézquita

(eah4d@missouri.edu)

Abstract

Cuscuta spp., (dodder) is a rootless and leafless parasitic plant posing significant agricultural

challenges. In this study, we aimed to elucidate the dynamics of the coiling patterns in *Cuscuta*

campestris and examine the role of circadian rhythms in its host-seeking ability. Using time-lapse

photography, we recorded the circumnutation and coiling movements of *C. campestris* at different

inoculation times on non-living hosts. Subsequent image analyses were facilitated through an in-

house Python-based image analysis pipeline. We observed that the coiling efficacy of *C.*

campestris varied with the inoculation time of day, showing higher success and faster initiation in

morning than in evening. These observations suggest that *Cuscuta*, despite lacking leaves and a

developed chloroplast, can discern photoperiod changes, which significantly determine its

parasitic efficiency. The automated image analysis confirmed the reliability of our Python pipeline,

aligning closely with manual annotations. This study provides significant insights into the parasitic

strategies of *C. campestris* and demonstrates the potential of integrating computational image

analysis in plant biology for exploring complex plant behaviors. Furthermore, this method

provides an efficient tool for investigating plant movement dynamics, laying the foundation for

future studies on mitigating the economic impacts of parasitic plants.

KEYWORDS: Parasitic plants, *Cuscuta campestris*, Circumnutation, Circadian rhythm,

Timelapse photography, Image analysis.

INTRODUCTION

Cuscuta species, commonly referred to as dodders, is an obligate stem parasitic plant of the Convolvulaceae family, widely recognized as morning glories (1). *Cuscuta* infestations can have profound economic implications. It can infect and considerably reduce the yield of more than 25 different crop species and has been found in at least 55 countries across all continents (2). *Cuscuta* infestations are difficult to control, and they can reduce the yield of tomato fields (3), cranberry vines (4), carrots (5), and alfalfa (6) by more than 75%. *Cuscuta* is resistant to most of the commercially available herbicides (7), and a single plant can produce up to 15,000 seeds that can remain viable for at least 10 years (8). As environmental patterns are altered due to climate change, species distribution models predict that the field and host ranges of *Cuscuta* can expand even further (9, 10). Therefore, new control strategies focus on determining the molecular and environmental conditions that affect *Cuscuta*'s ability to detect and attach itself successfully to new hosts during its seedling stage (11).

Cuscuta propagates through either seeds or stem cuttings (12, 13). Following germination from seed or attachment to a host via stem cuttings, *Cuscuta* initiates a counterclockwise rotational movement, referred to as circumnutation. Since this movement is crucial for *Cuscuta* to identify and secure a suitable host plant, circumnutation is the most essential process in early parasitism for the species. After recognition and successful attachment, *Cuscuta* starts producing feeding sites known as haustoria, which penetrate the vascular system of the host plant to obtain water and nutrients for its own survival (14-16). Without attachment to a host, *Cuscuta* cannot survive independently due to its lack of leaves and roots. Unraveling these circumnutating, coiling, and attaching dynamics are key to understand how *Cuscuta* survives and interact with its environment.

Circumnutation, first described by Charles Darwin (17), is a phenomenon characterized by the combination of circular movement and axial growth for the independent and autonomous movements of young and growing organs such as the flowers, stems, shoots, tips, petioles, tendrils, or roots of plants, including *Cuscuta* (18-22). Notably, these movements can be affected by diverse factors, including organ morphology, light, biotic and abiotic stresses, and temperature (18, 22).

Interestingly, circumnutation movement is influenced by the circadian clock (18, 23). The circadian clock helps plants adapt to varying light conditions by regulating physiological and behavioral processes through an endogenous 24-hour rhythm (24-26). In response to changing light conditions, the circadian clock uses external cues, such as light and temperature, to

synchronize its internal rhythm with the external environment. Notably, plant circadian rhythms
are self-sustaining (free running) and have an intrinsic, endogenous component that allows it to
maintain a 24-hour cycle even in the absence of external cues. For example, when sunflowers are
transferred from light/dark (LD) transition to constant light (LL), their circumnutation amplitude
and period remain consistent as the LD transition for a few days (18).

The conventional method for phenotyping and characterizing circumnutation and other
developmental traits typically entails direct visual examination and annotation. This process is
both time-consuming and prone to mistakes, which limits the number of phenotypes observed, and
plant samples analyzed. In a digital data-dominated era, open-sourced high-throughput plant
phenotyping (HTPP) pipelines are crucial to gain better and more profound insights from image-
based data (27, 28). For example, 2D image-processing HTTP pipelines can automatically track
size changes in *Arabidopsis* leaves (29), detect plant-pathogen interactions in excavated maize
roots (30), characterize leaf venation patterns for grasses (31), and even trace circumnutation
movements made by *Arabidopsis* stems (32). However, adapting existing HTPP pipelines for *C.*
campestris phenotyping presents challenges due to several unique features of the organism. These
include its absence of leaves and root system, as well its highly variable movement and twinning
patterns.

In this study, we introduce a Python-based image processing pipeline designed to phenotype
key traits associated with *C. campestris* circumnutation and coiling. This pipeline allows the
automated detection of coiling location, angle, initiation, and completion times, along with the
duration of coiling stages in between. Subsequently, we applied the pipeline to gain a better
understanding of circumnutation and coiling of *C. campestris* and investigated how circadian
rhythm influences successful inoculation. The findings obtained from this automated image
analysis tool aligned with those derived by manual annotation, confirming the reliability of the
Python pipeline. The pipeline is openly accessible as a series of commented Jupyter notebooks to
encourage appropriate modifications for different image data or experimental setups. This
approach not only enhances the precision in tracking plant movements but also paves the way for
future research on the mechanisms underlying parasitic plant interactions and potential sustainable
weed control strategies.

MATERIALS AND METHODS

Plant growth and *C. campestris* inoculation

The experimental workflow of our study is illustrated in Figure 1. *C. campestris* seedlings were inoculated on one-month-old beet plants (*Beta vulgaris)* in a greenhouse with a temperature of 25-30°C and day/night cycle of 16/8 hours. Three weeks after the initial inoculation, mature *C. campestris* stem segments were collected from the greenhouse. To evaluate the potential for inducing coiling under our experimental conditions, preliminary tests were conducted by attaching these *C. campestris* stems to 3-week-old *Arabidopsis* Col-0 and to bamboo skewers (30 cm in length and 4 mm in diameter), enabling a comparison of coiling patterns on living versus non-living materials (Fig. 2a-b, Suppl. Video 1). The *C. campestris* stem segments used in the inoculation experiments measured between 6-8 cm, with a shoot tip ranging from 4-6 cm in length and a single offshoot branch. The *C. campestris* stems were attached to their host or skewer with pieces of 3M Magic tape of approximately 3-3.5 cm in length and 4 mm in width. The tape was applied on the shoot tip at half its length. To ensure optimal inoculation, the shoot tips were aligned facing left at an angle of 30-60 degrees, considering that *C. campestris* shoot tips exhibit an anti-clockwise movement pattern. The attached shoot tips were kept under 24-hour constant lightening using far-red enriched lights with an intensity of 100 µmol/m^2 (GE R30 Soft White Medium Base, model #30711) as previously described in (33) (Supp Fig. 1). For the time lapse experiment, every sampling setup consisted of five *C. campestris* stems attached to five skewers roughly equidistant from each other. This setup was repeated seven times for three different inoculation times: 9 AM, 12 PM, and 4 PM, resulting in 35 individual stems for every inoculation time.

Camera setting, time lapse, and videos

Photos were captured on a Panasonic DC-FZ80 digital camera. The aspect ratio was set to 16:9, picture quality to 4896 x 2752 pixels, aperture value of 4.0, shutter speed of 100, ISO 1600, and an interval time of 96 seconds. The white balance was set manually based on the light conditions in the room before the timelapse. The focus was set using automatic settings, then switched to manual prior to capture. Flash, image stabilizing, and automatic white balance settings were turned off. From the ruler attached next to the skewers, 14 pixels correspond to 1 mm length. Shotcut software (https://shotcut.org) was used to make and render timelapse videos with the image

124 sequence option. The videos were rendered at 30fps using a libx265 HEVC codec and exported as

125 an mp4 file (Suppl. Videos 2-4).

127 Manual observation

According to *C. campestris* movement and coiling patterns on the bamboo skewer, we

manually observed success rate of coiling, coiling initiation, and completion times while *C.*

campestris stems coiled on bamboo skewers. Successful inoculations were quantified by

evaluating whether the *C. campestris* stems exhibited complete coiling around the bamboo skewers

showing a stable physical interaction. Only *C. campestris* stems that successfully coiled on

bamboo skewers were considered for analysis. Coiling initiation was defined as the time from the

point of contact until the *C. campestris* stems began its characteristic anti-clockwise twining

motion. Completion of inoculation was determined when the main shoots ceased to move

significantly around the surrogate host (Fig. 2c). Time intervals for initiation and completion of

coiling were recorded and averaged in a spreadsheet for each inoculation event within the

experiment.

140 Automatic detection and image analysis

An in-house SciPy-based Python image processing script was developed to automatically

extract movement and position information from the individual standstill photos described above.

(Suppl. Videos 5-9). This way we divided the *C. campestris* coiling process into nine partially

overlapping steps: 1) inoculation, 2) Coil 1 initiation, 3) Coil 1 positioning, 4) Coil 1 tightening,

5) Coil 1 completion, 6) Coil 2 initiation, 7) Coil 2 positioning, 8) Coil 2 tightening, and 9)

completion, as explained below. The script is based on a combination of color, hue, and saturation

thresholding and elementary image morphology operations. All the image processing and detailed

analyses steps, limitations, and considerations can be read as Jupyter notebooks available at the

GitHub repository listed at the end. The exact hyperparameters were first manually tuned for a

single photo and these values were later used for the rest of the images; this was possible since all

the photos have comparable illumination and color values.

First, the background was thresholded (black in Fig. 3b); the centers of the skewers were

approximated with ordinary least-squares best fit lines (red in Fig. 3b); and the top edge of the tape

pieces was located (orange in Fig. 3b). Next, the image was split into 5 overlapping sub-images,

each centered on a different skewer; left and right limits determined by neighboring skewers; bottom margin determined by tape (Fig. 3c-g). The *C. campestris* in each section was identified as the union of large, connected components near the central skewer (Fig. 3h-l). The resulting segmented *C. campestris* was skeletonized (34) (gray in Fig. 3m-q). Subsequent phenotyping was limited to instances where the skeleton crossed in front the center of the skewer (white dot in Fig. 3r-v). These crossing instances were modeled by weighted linear RANSACs (35), where pixels closer to the crossing site had higher weights (light blue and mustard in Fig. 3s-v). The crossing angle was determined as the complement of the angle between the skewer central line and the RANSAC line (pink in Fig. 3s-v). It is important to note that an angle of 0 degrees indicates that the *C. campestris* crosses perpendicularly to the skewer. Repeating these steps for all snapshots resulted in location and angle time series for Coil 1 and Coil 2 for every individual *C. campestris* (Fig. 4a). Missing time points were interpolated using linear splines and outliers were removed following a Savitzky-Golay filter (36). A *C. campestris* stem was determined as successfully coiled if it presented relevant angle and position information for both Coil 1 and Coil 2.

As illustrated in Figure 4, Coil 1 is initiated at 7.7 hours after inoculation (HAI) when the *C. campestris* crosses in front of the skewer for the first time. Immediately after, Coil 1 *positions* itself for the next 4.3 hours—both its angle and position vary more than 5 degrees and 1 mm respectively with respect to their final values at 24 HAI. In this case, Coil 1 stops positioning at 12.0 HAI, as its position remains stable for the remainder of the recording. Coil 1 then *tightens* for the next 4.8 hours —its angle remains variable while its position is stable. Finally, at 16.8 HAI, Coil 1 is completed as its angle also remains stable for the remainder of the recording. The coil stabilization time is simply the difference between completion and initiation times, which is 9.1 hours in this case. Note that this is the same as the sum of positioning and tightening periods. The times and periods for Coil 2 are analogous. In this case, Coil 2 initiates at 15.5 HAI, it exhibits no positioning period, and it tightens for 3.8 hours to be completed at 19.3 HAI so that it is stabilized after 3.8 hours. Additionally, the time for *C. campestris* to turn 360° was determined as the difference between Coil 1 and Coil 2 initiation times (7.8 hours). Finally, the whole coiling process time for a *C. campestris* was determined as the time difference between the initiation of Coil 1 and the completion of Coil 2 (11.6 hours). An animated version of this example can be seen in Supplementary Video 5.

Statistical analysis

A Mann-Whitney U-test (37) was performed to compare if the reported position, angle, and time values had statistically different average values for different pairs of initial inoculation times or different coils. This test was chosen since it does not assume any special conditions on the underlying distributions. This test was used whenever comparing manual or automatic observations.

Spearman's rank correlation coefficients were computed between the time to turn 360° and the different time periods related to the coiling —positioning, tightening, stabilization times— as well as between positioning and tightening times. Additionally, p-values associated to these coefficients were computed following a t-distribution. These analyses were carried out separately for different inoculation times and for Coil 1 and Coil 2. (Supp Fig 4). The Spearman coefficient was chosen because it does not assume linearity.

RESULTS

C. campestris **coiled on both living and non-living hosts in 24 hours**

Within 24 hours following *C. campestris* inoculation, the shoot tips successfully coiled on both bamboo skewers and *Arabidopsis* plants (Fig. 2a-b and Suppl. Video 1). This indicates that *C. campestris* can coil around a host within a day, and that bamboo skewers are suitable surrogate hosts for studying *C. campestris* coiling movements. These findings are consistent with previous research showing *C. campestris*'s ability to coil around non-living materials and form haustoria when exposed to an appropriate blue/far-red light ratio (12, 33). Consequently, skewers were used as surrogate hosts for subsequent analysis. Furthermore, we conducted manual observations to assess the success rate of coiling, as well as to track the initiation and completion times of coiling, while closely monitoring *C. campestris* stems as they coiled around bamboo skewers (Fig 2c).

Successful rate of *C. campestris* **coiling under different inoculation times**

Manual observations indicated a 100% success rate for *C. campestris* coiling at 9 AM (35 stems), whereas the success rates were approximately 83% at both 12 PM and 4 PM, with 29 stems each (Fig 5a, left panel; Supplementary Table 1, and Suppl. Video 2-4). Automated observations showed a similar trend with lower rates for each time point: 97.1% at 9 AM (34 stems), 71.4% at

12 PM (25 stems), and 80% at 4 PM (28 stems) (Fig 5b, right panel; Supp table 2). This discrepancy is primarily attributed to difficulties in distinguishing the color between the *C. campestris* stem and the skewers in some replicates and in extracting data from skewers that did not remain completely stationary.

Initiation and completion of *C. campestris* coiling under different inoculation times

Manual observations revealed average coiling initiation times of 5.3, 5.9, and 8.5 hours after inoculation (HAI) for *C. campestris* inoculated at 9 AM, 12 PM, and 4 PM, respectively (Fig 5b, left panel; Supp table 3, and Suppl. Video 2-4). Notably, significant differences were observed in initiation times between *C. campestris* inoculated at 9 AM and 4 PM (p-value $< 10^{-5}$), as well as between 12 PM and 4 PM (p-value $< 10^{-3}$). Interestingly, stems inoculated at 4 PM exhibited delayed coiling initiation compared to those at 9 AM, potentially due to longer resting periods or circumnutation before recognizing the bamboo skewer. However, no significant differences were observed between inoculation at 9 AM and 12 PM. A similar trend was observed in the completion times, with an average time of 14.4, 16.5, and 18.0 HAI for 9 AM, 12 PM, and 4 PM, respectively (Fig 5c, left panel; Supp table 3, and Suppl. Video 2-4). In this case, the completion time was significantly different between *C. campestris* inoculated at 9 AM vs 4 PM (p-value $< 10^{-3}$).

Similar conclusions can be drawn from the automatic observations. Coil 1 was initiated on average at 4.8, 5.3, and 7.9 HAI for 9 AM, 12 PM, and 4 PM, respectively (Fig 5b, middle panel; Supp table 4). Coil 2 was initiated at 7.8, 8.7, and 11.9 HAI for 9 AM, 12 PM, and 4 PM, respectively (Fig 5b, right panel; Supp table 5). These times are significantly different between *C. campestris* inoculated at 9 AM and 4 PM (p-values $< 10^{-5}$), and between 12 PM and 4 PM (p-values $< 10^{-4}$) for both Coil 1 and Coil 2. There were no significant differences between 9 AM and 12 PM. Do notice that the automated observations are based on when the *C. campestris* crossed in front of the skewer without regard the context of the movement, which most likely explains why the automatically reported times are earlier than the manual ones.

The automatic observations show that Coil 1 was completed on average at 14.8, 15.9, and 16.7 HAI for 9 AM, 12 PM, and 4 PM, respectively (Fig 5c, middle panel; Supp table 4). Coil 2 was completed at 13.3, 15.2, and 17.4 HAI for 9 AM, 12 PM, and 4 PM, respectively (Fig 5c, right panel; Supp table 5). We observed that Coil 2 tends to reach a stable state slightly earlier than Coil

1, but these differences were not significant. There are no significant differences in completion time for Coil 1 with respect to inoculation time. However, there is a significant difference in Coil 2 completion times between 9 AM and 4 PM (p-value $< 10^{-3}$). Since the automatic annotation focuses solely on the behavior of the *C. campestris* in front of the skewer, it ignores the possibility of the *C. campestris*'s upper tip still moving while the coil in front of the skewer is stationary. This could explain why the automatically reported completion times of Coil 2 happen earlier than the manual ones. Nonetheless, despite different annotating criteria, both manual and automatic observations highlight significant differences in coiling initiation and completion times among *C. campestris* inoculated at different times, with those inoculated at 9 AM consistently showing earlier coiling compared to those inoculated at 4 PM.

The automated pipeline tracks the coiling stabilization, twisting times, angles, positions, and gaps of the *C. campestris* coils.

The automatic observations revealed consistent findings across different inoculation times. Firstly, there were no significant variations in the gap size between coils or in the time required to complete a full 360° coil around the skewer (Supp Fig 2a-b). Additionally, examination of Coil 1 and Coil 2 angles at 24 HAI showed no statistical differences in their final angle and position across all inoculation times. The average angle of Coil 1 was 23.5, 21.8, and 30.4 degrees for *C. campestris* inoculated at 9 AM, 12 PM, and 4 PM, respectively (Supp Fig 2c, left panel), while for Coil 2, the angles were 8.1, 9.0, and 13.9 degrees, respectively (Supp Fig 2c, right panel). Notably, there were no statistical differences in final angles and positions observed across all inoculation times, whether examining Coil 1 or Coil 2 (Supp Fig 2c-d). However, a marked contrast was observed between the final angle of Coil 1 and Coil 2 for every inoculation time (p-values $< 10^{-4}$) (Supp Fig 2c). Specifically, Coil 1 tended to be more angled or slanted, while Coil 2 generally aligned parallel to the ground. These results indicate that the coiling dynamics of *C. campestris*, including the speed of wrapping and the final positioning of the coils, are not influenced by the time of day at which inoculation occurs, but they might be affected by the mechanical differences between the first and second coils.

Similar trends were observed when examining the time periods of different coiling stages. There were no significant differences in the positioning, tightening, and stabilization period durations for either Coil 1 or Coil 2 across all inoculation times (Supp Fig 3a-c). However, there

was a striking statistical difference between the positioning duration of Coil 1 and Coil 2 across all inoculation times (p-values < 10^{-4}). Coil 1 tended to position for about 6.0, 5.6, and 5.6 hours, while Coil 2 only positioned for 1.4, 2.6, and 2.0 hours, for *C. campestris* inoculated at 9 AM, 12 PM, and 4 PM, respectively. These time differences between coils are not observed in tightening time duration, but they reemerge when considering stabilization times (p-values < 10^{-3}). In this case, Coil 1 tends to stabilize within 10.0, 10.6, and 9.0 hours compared to 5.6, 7.0, and 5.4 hours for Coil 2 for inoculations at 9 AM, 12 PM, and 4 PM, respectively. In other words, Coil 1 tends to move up and down the skewer for a while before finding a stationary position, while Coil 2 remains in place soon after it has twined with just slight adjustments in angle, suggesting mechanical differences between these two coils (Supp Fig 3a-c).

There were no significant differences when comparing the whole duration of the coiling process across all inoculation times. This conclusion is supported whenever using either the manual or automated annotation criteria (Supp Fig 3d). Such result suggests that while *C. campestris* inoculated at 9 AM tends to initiate coiling considerably earlier than *C. campestris* inoculated at 4 PM, once they start, the mechanistic coiling processes and stages have similar durations. This could indicate that *C. campestris* inoculated at 9 AM reaches a stationary state considerably earlier than *C. campestris* inoculated at 4 PM mainly because the former has an earlier head start (Fig 6). This might also indicate that the mechanisms involved in coiling initiation are independent from those involved in turning and twining. This final observation is also supported by a lack of correlation between the time it takes *C. campestris* to complete a 360° turn and the time it spends positioning, tightening, or stabilizing, regardless of the inoculation time or the coil observed (Supp Fig 4).

DISCUSSION

Throughout evolution, plants have developed a plethora of survival strategies. Among these strategies, parasitic plants have evolved unique adaptations to exploit other plant species for their resources. These unique members of the plant kingdom exhibit either facultative or obligatory dependence on other plant species for survival. They are also classified either as stem or root parasites, depending on the point of attachment to a host (38). *Cuscuta* has evolved specialized haustoria structures and exhibits remarkable mobility, while undergoing atrophy in its root and leaf organs (14, 39). Despite the critical role of mobility, such as circumnutation and coiling, in

the successful parasitism of *Cuscuta*, studies investigating this aspect have been limited and primarily focused on manual observations, often in the context of responses to volatiles or light wavelengths (40-43). Addressing this gap, our study aimed to provide a combined analysis of *Cuscuta* coiling dynamics, using both manual observations and automated detection methods.

We found that *C. campestris* inoculations at 9 AM resulted in the highest success rate of inoculation and showed the fastest coiling initiation around the host (Fig. 5a-b). In contrast, inoculations at 4PM showed a lower success rate and an approximately 1.5-fold slower coiling initiation compared to those at 9 AM and 12 PM, possibly due to an extended resting stage or prolonged circumnutation movement (Fig 5b and 6). These observations suggest a difference in circumnutation and coiling behavior of *C. campestris* depending on the time of day. These observations give evidence that *C. campestris* perform circumnutation best in morning, moderately in afternoon, and lower in the evening, indicating that *C. campestris* has preferable times for active circumnutation and subsequent coiling. Interestingly, similar diurnal patterns have been observed in sunflowers, where young sunflowers exhibited a wide amplitude of circumnutation from midnight to noon, gradually decreasing in the afternoon and reaching a resting stage from evening (around 6PM) until midnight (18). This indicates that like sunflowers, rootless and leafless *C. campestris* retains the light/dark period even though its stem is cut. Moreover, it suggests that *C. campestris* has its own internal oscillator capable of detecting the circadian clock, regulating circumnutation for successful parasitism independent of host-derived hormones or signals. It remains unclear if the purpose of circumnutation and coiling movements in *C. campestris*; are influenced by its circadian clock and if these movements serve as a strategic tactic for successful parasitism. Further experiments are required to fully understand this biological aspect of *Cuscuta*.

Previous studies have shown that *Cuscuta* has loose or tight coils under different wavelengths, critically affecting its haustoria development (41, 43, 44). When *Cuscuta* presents loose coils under red light or a high red/far red (R/FR) ratio, there was no haustoria development, while low R/FR ratio and blue lights induced tight coils and successful haustoria development. An interesting discovery from previous studies, although not directly focused on this aspect, is that the final coil at the upper section of loose coils consistently exhibited higher angles compared to tight coils, even though *Cuscuta* wraps around the host multiple times, with all coils varying in angles. Notably, haustoria organs were produced from the flattened upper coil of tight coils, indicating that the angle of the top coil is important for the haustoria development (41). In this study, we

separately analyzed Coil 1, Coil 2, and stabilization of coiling (positioning and tightening) mathematically. As a result, there were significant angle and time differences when comparing Coil 1 and Coil 2. On one hand, Coil 1 tended to position for 5.7 hours and stabilize after 9.9 hours, coming to a final angle of 25° with respect to the base. On the other hand, Coil 2 tended to position just for 3.0 hours and stabilize after 6.0 hours, coming to a final angle of 10° with respect to the base. Therefore, Coil 1 tended to exhibit much more mobility after contacting the skewer to then stay at a slanted angle, whereas Coil 2 tended to be more limited in movement and settled at a flat angle. Thus, we suggest that our approach has capability to distinguish mechanistic differences between loose or tight coils, as previously described (41), and predict the development of the haustoria organ prior to its emergence from the inoculated *Cuscuta* stem.

Moreover, the automated pipeline of this study can determine more nuanced time points and periods related to *Cuscuta*'s coiling process. For instance, through these automatic analyses, it was observed that the time it takes *C. campestris* to twist around the skewer is independent of the time it takes to stabilize its coiling, regardless of the inoculation time. This suggests that twisting and stabilization follow two different mechanisms. This agrees with prior reports where twining and haustoria development could be independently induced from each other using different light stimuli (43). Another observation from automated annotations is that once *Cuscuta* initiates coiling, the period duration for each of its coiling stages —positioning, tightening, and stabilization— is independent of the time of inoculation, even though *Cuscuta* inoculated during the morning initiates considerably quicker than the one inoculated during afternoon. It was also observed that all phenotypes related to final coil positions and angles were also independent of inoculation time.

At the moment, the proposed pipeline can only extract phenotypes associated to *C. campestris* 's stem angle and position with respect to the skewer by contrasting the skewers' and parasites' color. A more careful choice of background and skewer color can provide clearer images from which the whole *C. campestris* stem could be digitally isolated. It would be thus possible to automatically extract more movement and circumnutation characteristics, and even refine the criteria to determine coil initiation and completion so that they coincide with those used in a manual setting.

Circumnutation and coiling represent crucial behaviors for stem parasitic plants, facilitating their effective parasitic lifestyle. This research endeavors to advance computational image analysis

tools capable of monitoring *Cuscuta* movement under controlled laboratory conditions within a
short period of time. Through these invested tools, we aim to delve into the biological,
morphological, biochemical, and physiological aspects of *Cuscuta* mobility. Such investigations
promise fundamental insights into the intricate interactions between plants, particularly between
hosts and parasitic species, thereby laying the groundwork for parasitic plant management
strategies.

In summary, our study introduces a Python-based image processing pipeline designed to
phenotype key traits associated with *Cuscuta* circumnutation and coiling. This tool automates the
detection of plant movement parameters such as coiling location, angle, initiation, and completion
times, as well as the durations of various coiling stages. We utilized this pipeline to better
understand the coiling behavior and circadian rhythms of *Cuscuta*. The data generated by this
automated tool agreed with manual annotations, underscoring its accuracy and reliability.
Additionally, we have made the pipeline publicly available in the form of Jupyter notebooks on
GitHub. This facilitates adaptations to diverse imaging datasets and experimental conditions,
broadening its applicability. Our approach not only refines the accuracy of tracking plant
movements but also lays the groundwork for further exploration into the mechanisms of parasitic
plant interactions, potentially leading to sustainable methods for managing parasitic weeds.

AUTHOR CONTRIBUTIONS

SP, EA, and JB conceived the ideas and designed the methodology. MB and SA conducted lab work and analyzed the data. EA performed image analysis and script development. MB and EA led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGMENTS

This research received support from the USDA-AFRI grant number 2023-67013-39896 awarded to SP, as well as funding from the Research Council, the College of Agriculture, Food, and Natural Resources (CAFNR), and the Interdisciplinary Plant Group (IPG) at the University of Missouri. Grammarly generated responses to the following AI prompts: Prompts created by Grammarly- "Improve it".

CONFLICTING INTEREST STATEMENT

All authors declare that they have no conflicts of interest.

DATA AVAILABILITY STATEMENT

The developed Python-based pipeline is available as a collection of Jupyter notebooks at https://github.com/ejamezquita/cuscuta/ . The datasets used and/or analyzed during the current study are available here:

https://youtube.com/playlist?list=PLZkYcVyQr2u4tT0yoZAkrMqzQxRIDvxru&feature=shared

ORCID

Max Bentelspacher https://orcid.org/0009-0004-7357-917X

Erik J. Amézquita https://orcid.org/0000-0002-9837-0397

Supral Adhikari https://orcid.org/0000-0002-9914-2986

Jaime Barros https://orcid.org/0000-0002-9545-312X

So-Yon Park https://orcid.org/0000-0002-2126-3382

REFERENCES

1.   S. Riviere, C. Clayson, K. Dockstader, M. A. R. Wright, M. Costea, To attract or to repel? Diversity, evolution and role of the "most peculiar organ" in the Cuscuta flower (dodder, Convolvulaceae)—the infrastaminal scales. *Plant Syst. Evol.* **299**, 529-552 (2013).

2.   M. J. Kogan, W. T. Lanini, Biology and management of Cuscuta in crops. *Ciencia E Investigacion Agraria* **32**, 165-180 (2005).

3.   Y. Goldwasser, W. T. Lanini, M. R. M. Sazo, Control of Field Dodder (Cuscuta campestris) Parasitizing Tomato with ALS-Inhibiting Herbicides. *Weed Technol.* **26**, 740-746 (2012).

4.   K. M. Ghantous, H. A. Sandler (2017) Evaluations of pre- and postemergence herbicides for dodder management in cranberry. (International Society for Horticultural Science (ISHS), Leuven, Belgium), pp 379-386.

5.   J. B. Colquhoun, C. M. Konieczka, R. A. Rittmeyer, Swamp Dodder (Cuscuta gronovii) Management in Carrot Production. *Weed Technol.* **23**, 408-411 (2009).

6.   J. S. Mishra, Biology and management of Cuscuta species. *Indian Journal of Weed science* **41**, 1-11 (2009).

7.   T. Nadler-Hassar, B. Rubin, Natural tolerance of Cuscuta campestris to herbicides inhibiting amino acid biosynthesis. *Weed Res.* **43**, 341-347 (2003).

8.   M. Saric-Krsmanovic, S. Vrbnicanin, Field dodder - how to control it? *Pestic. Fitomed.* **30**, 137-145 (2015).

9.   J. Masanga *et al.*, Physiological and ecological warnings that dodders pose an exigent threat to farmlands in Eastern Africa. *Plant Physiol.* **185**, 1457-1467 (2021).

10.   C. Cai *et al.*, Implications of climate change for environmental niche overlap between five Cuscuta pest species and their two main Leguminosae host crop species. *Weed Sci.* **70**, 543-552 (2022).

11.   M. Hartenstein, M. Albert, K. Krause, The plant vampire diaries: a historic perspective on Cuscuta research. *J. Exp. Bot.* **74**, 2944-2955 (2023).

12.   V. Hegenauer, M. Welz, M. Körner, M. Albert, Growth Assay for the Stem Parasitic Plants of the Genus Cuscuta. *Bio Protoc* **7**, e2243 (2017).

13.   F. M. Ashton, J. M. Hutchison, Germination of Field Dodder (Cuscuta campestris). *Weed Sci.* **28**, 330-333 (1980).

14.   M.-Y. Jhu, N. R. Sinha, Cuscuta species: Model organisms for haustorium development in stem holoparasitic plants. *Frontiers in plant science* **13**, 1086384 (2022).

15.   G. Kim, J. H. Westwood, Macromolecule exchange in *Cuscuta*–host plant interactions. *Curr. Opin. Plant Biol.* **26**, 20-25 (2015).

16.   K. Shimizu, K. Aoki, Development of Parasitic Organs of a Stem Holoparasitic Plant in Genus Cuscuta. *Frontiers in Plant Science* **10** (2019).

17.   C. Darwin, F. Darwin, *The power of movement in plants* (John Murray, 1880).

18.   M. Stolarz, Circumnutation as a visible plant action and reaction: physiological, cellular and molecular basis for circumnutations. *Plant Signal Behav* **4**, 380-387 (2009).

19.   D. E. Moulton, H. Oliveri, A. Goriely, Multiscale integration of environmental stimuli in plant tropism produces complex behaviors. *Proceedings of the National Academy of Sciences* **117**, 32226-32237 (2020).

20. A. H. Brown, Circumnutations: from Darwin to space flights. *Plant Physiol.* **101**, 345 (1993).

21. D. Agostinelli, A. DeSimone, G. Noselli, Nutations in plant shoots: endogenous and exogenous factors in the presence of mechanical deformations. *Frontiers in Plant Science* **12**, 608005 (2021).

22. Y. Wu, J. Xie, L. Wang, H. Zheng, Circumnutation and Growth of Inflorescence Stems of Arabidopsis thaliana in Response to Microgravity under Different Photoperiod Conditions. *Life* **10**, 26 (2020).

23. K. Niinuma, N. Someya, M. Kimura, I. Yamaguchi, H. Hamamoto, Circadian Rhythm of Circumnutation in Inflorescence Stems of Arabidopsis. *Plant and Cell Physiology* **46**, 1423-1427 (2005).

24. C. R. McClung, Plant circadian rhythms. *Plant Cell* **18**, 792-803 (2006).

25. K. Greenham, C. R. McClung, Integrating circadian dynamics with physiological processes in plants. *Nature Reviews Genetics* **16**, 598-610 (2015).

26. N. Creux, S. Harmer, Circadian Rhythms in Plants. *Cold Spring Harb Perspect Biol* **11** (2019).

27. N. Fahlgren, M. Gehan, I. Baxter, Lights, camera, action: high-throughput plant phenotyping is ready for a close-up. *Curr. Opin. Plant Biol.* **24**, 93-99 (2015).

28. J. L. Araus, J. E. Cairns, Field high-throughput phenotyping: the new crop breeding frontier. *Trends Plant Sci.* **19**, 52-61 (2014).

29. L. G. Swartz *et al.*, OPEN leaf: an open-source cloud-based phenotyping system for tracking dynamic changes at leaf-specific resolution in Arabidopsis. *The Plant Journal* **116**, 1600-1616 (2023).

30. L. D. Pierz, D. R. Heslinga, C. R. Buell, M. J. Haus, An image-based technique for automated root disease severity assessment using PlantCV. *Applications in Plant Sciences* **11**, e11507 (2023).

31. J. M. Robil *et al.*, grasviq: an image analysis framework for automatically quantifying vein number and morphology in grass leaves. *The Plant Journal* **107**, 629-648 (2021).

32. Y. Mao, H. Liu, Y. Wang, E. D. Brenner, A deep learning approach to track Arabidopsis seedlings' circumnutation from time-lapse videos. *Plant Methods* **19**, 18 (2023).

33. V. Bernal-Galeano, K. Beard, J. H. Westwood, An artificial host system enables the obligate parasite Cuscuta campestris to grow and reproduce in vitro. *Plant Physiol.* **189**, 687-702 (2022).

34. T. Y. Zhang, C. Y. Suen, A fast parallel algorithm for thinning digital patterns. *Communications of the ACM* **27**, 236-239 (1984).

35. M. A. Fischler, R. C. Bolles, Random sample consensus: a paradigm for model fitting with applications to image analysis and automated cartography. *Commun. ACM* **24**, 381–395 (1981).

36. A. Savitzky, M. J. Golay, Smoothing and differentiation of data by simplified least squares procedures. *Analytical chemistry* **36**, 1627-1639 (1964).

37. H. B. Mann, D. R. Whitney, On a test of whether one of two random variables is stochastically larger than the other. *The annals of mathematical statistics*, 50-60 (1947).

38. J. Těšitel, Functional biology of parasitic plants: a review. *Plant Ecology and Evolution* **149**, 5-20 (2016).

39. G. Sun *et al.*, Large-scale gene losses underlie the genome evolution of parasitic plant Cuscuta australis. *Nature Communications* **9**, 2683 (2018).

508   40.   J. B. Runyon, M. C. Mescher, C. M. De Moraes, Volatile chemical cues guide host
509   location and host selection by parasitic plants. *Science* **313**, 1964-1967 (2006).
510   41.   T. Yokoyama, A. Watanabe, M. Asaoka, K. Nishitani, Germinating seedlings and mature
511   shoots of Cuscuta campestris respond differently to light stimuli during parasitism but not
512   during circumnutation. *Plant, Cell Environ.* **46**, 1774-1784 (2023).
513   42.   T. Furuhashi, K. Furuhashi, W. Weckwerth, The parasitic mechanism of the
514   holostemparasitic plant Cuscuta. *J. Plant Interactions* **6**, 207-219 (2011).
515   43.   K. Furuhashi, K. Iwase, T. Furuhashi, Role of Light and Plant Hormones in Stem
516   Parasitic Plant (Cuscuta and Cassytha) Twining and Haustoria Induction. *Photochem.*
517   *Photobiol.* **97**, 1054-1062 (2021).
518   44.   H. Pan, Y. Li, L. Chen, J. Li, Molecular Processes of Dodder Haustorium Formation on
519   Host Plant under Low Red/Far Red (R/FR) Irradiation. *International Journal of*
520   *Molecular Sciences* **23**, 7528 (2022).

Fig 1. Experimental workflow for monitoring the coiling movement of *Cuscuta*. After acquiring time-lapse images of *Cuscuta* stems subjected to varying inoculation times, each measurement was analyzed through manual observation or automatic detection.

Fig 2. *Cuscuta* stems were wrapped around *Arabidopsis* stems and bamboo skewers. **(a)** *Cuscuta* stems were attached on *Arabidopsis* Col-0 and bamboo skewers and incubated for 24 hours. **(b)** Zoomed-in images of lined boxes in **(a)**. **(c)** Timelapse images were captured to observe the coiling movement of *Cuscuta* stems under different inoculation times. Scale bar: 1 cm. HAI: hours after inoculation. Black arrows indicate *Cuscuta* stems.

Fig. 3. Image processing and analysis of *Cuscuta* images. **(a)** Standstill image of the *Cuscuta* experiment setup as initial input. **(b)** The background was thresholded out (black), the center of each skewer was approximated with a best-fit line (red line), and the tape upper border was identified (orange line). Further steps only considered the skewer section above such tape (cyan). **(c–g)** The image was then split into 5 sub-images, each centered at a different skewer, from which **(h–l)** the relevant *Cuscuta* (white) was segmented and **(m–q)** skeletonized (gray). Further analysis focused solely on instances where the skeleton crosses the center of the skewer. **(r)** Example where no information is extracted as the *Cuscuta* has not yet passed in front of the skewer. **(s–v)** The position where *Cuscuta*'s skeleton crosses the skewer is recorded (white star) as well as the complementary angle (pink pie) made by the first coil (cyan line) and **(v)** second coil (mustard line). The skewer is shown in purple and its midline approximation is shown in red. The yellow scalebar corresponds to 56 pixels which approximates to 2 mm for all images.

Fig. 4. Automated detection of essential time points of *Cuscuta* coiling process. **(a)** For a single

Cuscuta, the angle and position information of both coils is extracted from all the standstill images

(Fig. 3). The scatter plot represents the actual observations. The solid lines represent the time series

obtained after removing outliers and smoothening the signal. **(b)** Sample standstill sub-images

represented by the time series. Raw (R) and its corresponding processed (P) sub-image are

presented side by side for visual comparison. The yellow scalebar corresponds to 2 mm for all sub-

images. In this analysis, the time series contains no information prior to 7.7 HAI since *Cuscuta*

has not crossed in front of the skewer yet. Similarly, no information on Coil 2 was recorded prior

to 15.5 HAI. Refer to Supplementary Video 5 for an animated example. **(c)** From the time series,

for each coil we can immediately determine the time points when it initiated, when its position

stopped changing, and when its angle stopped changing. By identifying these timepoints (in blue),

we can deduce the duration of coiling stages (in gray).

Fig 5. Examining the coiling behavior of *Cuscuta* through manual and automatic detection

methods. The successful rate of *Cuscuta* coiling (n=35) **(a)**, coiling initiation time **(b)**, and coiling

completion time **(c)** were measured by manual and automatic detection methods. ns: Not

significant, p-value of ≥ 0.05; *: p-value of 0.05 to 0.005, **: p-value of 0.005 to 0.0005, and ***:

p-value of < 0.0005.

Fig 6. A summary of *Cuscuta* coiling movement using automatic detection methods with three

different inoculation times (9 AM, 12 PM and 4 PM). Blue bars and yellow bars indicate average

time of rest and coiling period. Dark and light gray shading indicates actual light/dark cycle and

subjective darkness, respectively.

Fig 1.

(a)

0.0 HAI   18.0 HAI

(b)

0.0 HAI   18.0 HAI

(c)

Inoculation
(0.0 HAI)

Coiling initiation
(4.1 HAI)

Completion
(18.0 HAI)

Fig 2.

Fig 3.

Fig 4.

Fig 5.

Fig 6.