ATG6 interacting with NPR1 increases *Arabidopsis thaliana* resistance to *Pst* DC3000/*avrRps4* by increasing its nuclear accumulation and stability

Baihong Zhang[1], Shuqin Huang[1], Shuyu Guo[2], Yixuan Meng[1], Yuzhen Tian[1], Yue Zhou[1], Hang Chen[1], Xue Li[1], Jun Zhou[1, *], Wenli Chen[1, *].

[1] MOE Key Laboratory of Laser Life Science & Institute of Laser Life Science, College of Biophotonics, South China Normal University, Guangzhou 510631, China; Guangdong Provincial Key Laboratory of Laser Life Science, College of Biophotonics, South China Normal University, Guangzhou 510631, China;

[2] State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, College of Life Sciences, Inner Mongolia University, Hohhot, 010070, PR China；Key Laboratory of Herbage and Endemic Crop Biotechnology, and College of Life Sciences, Inner Mongolia University, Hohhot, 010070, China;

*Corresponding Author: Wenli Chen, e-mail address: chenwl@scnu.edu.cn; Jun Zhou, e-mail address: zhoujun@scnu.edu.cn.

E-mail address: zhangbyhome@126.com (Baihong Zhang); huangshuqin@m.scnu.edu.cn (Shuqin Huang); guoshuyu1220@163.com (Shuyu Guo); 2215341147@qq.com (Yixuan Meng); tianyzh@126.com (Yuzhen Tian); 307758744@qq.com (Yue Zhou); 2674876291@qq.com (Hang Chen); 1374834169@qq.com (Xue Li).

Word counts: 4190 (excluding materials and methods)

Figures: 8

Supplemental Figures: 15

Supplemental movies: 2

Supplemental Data: 2

Running title: ATG6 interacting with NPR1 increases *Arabidopsis thaliana* resistance to *Pst* DC3000/*avrRps4*

 Highlight

We unveil a novel relationship in which ATG6 positively regulates NPR1 in

plant immunity.

Abstract

Autophagy-related gene 6 (ATG6) plays a crucial role in plant immunity.

Nonexpressor of pathogenesis-related genes1 (NPR1) acts as a signaling hub

of plant immunity. However, the relationship between ATG6 and NPR1 is

unclear. Here, we find that ATG6 directly interacts with NPR1. *ATG6*

overexpression significantly increased nuclear accumulation of NPR1.

Furthermore, we demonstrate that *ATG6* increases NPR1 protein levels and

improves its stability. Interestingly, ATG6 promotes the formation of SINCs

(SA-induced NPR1 condensates)-like condensates. Additionally, ATG6 and

NPR1 synergistically promote the expression of *pathogenesis-related* genes.

Further results showed that silencing *ATG6* in *NPR1-GFP* exacerbates *Pst*

DC3000/*avrRps4* invasion, while double overexpression of *ATG6* and *NPR1*

synergistically inhibits *Pst* DC3000/*avrRps4* invasion. In summary, our

findings unveil an interplay of NPR1 with ATG6 and elucidate important

molecular mechanisms for enhancing plant immunity.

Keywords: ATG6, NPR1, Nucleus, Cytoplasm, Stability, *Pst* DC3000/*avrRps4*.

Abbreviations：

artificial miRNA^{ATG6} (amiRNA^{ATG6}); Autophagy-related genes (ATGs);

Cycloheximide (CHX); Gold nanoparticles (AuNPs); Nonexpressor of

pathogenesis-related genes 1 (NPR1); Nuclear localization sequence (NLS);

P. syringae pv. tomato (*Pst*); SA-induced NPR1 condensates (SINCs);

Salicylic acid (SA); *UBQ10::ATG6-mCherry × 35S::NPR1-GFP* (*ATG6-*

mCherry × NPR1-GFP).

Introduction

Plants are constantly challenged by pathogens in nature. In order to survive and reproduce, plants have evolved complex mechanisms to cope with attack by pathogens (Jones and Dangl, 2006). Nonexpressor of pathogenesis-related genes 1 (NPR1) is a key regulator of plant immunity (Chen *et al.*, 2021b). It contains the BTB/POZ (Broad Compex, Tramtrack, and BricaBrac/Pox virus and Zinc finger) domain in the N-terminal region, the ANK (Ankyrin repeats) domain in the middle region, and the nuclear localization sequence (NLS) in the C-terminal region (Cao *et al.*, 1997; Rochon *et al.*, 2006). NPR1 is a receptor of SA (salicylic acid) mainly localized as an oligomer in the cytoplasm and sensitive to the surrounding redox state (Tada *et al.*, 2008; Wu *et al.*, 2012). SA mediates the dynamic oligomer to monomer response of NPR1 (Tada *et al.*, 2008) and promotes translocation of NPR1 into the nucleus, which increases plant resistance to pathogens by activating the expression of immune-related genes (Kinkema *et al.*, 2000; Chen *et al.*, 2021b).

NPR1 is mainly degraded by the ubiquitin proteasome system (UPS) (Spoel *et al.*, 2009; Saleh *et al.*, 2015; Skelly *et al.*, 2019). An increasing researches have shown that autophagy and the UPS pathway play overlapping roles in regulating intracellular protein homeostasis (Zhou *et al.*, 2014; Marshall *et al.*, 2015; Kikuchi *et al.*, 2020). Our previous study showed that ATGs (autophagy-related genes) are involved in NPR1 turnover (Gong *et al.*, 2020). Autophagy negatively regulates *Pst* DC3000/*avrRpm1*-induced programmed cell death (PCD) via the SA receptor NPR1 (Yoshimoto *et al.*, 2009). These results imply that ATGs might be involved in plant immunity through the regulation of NPR1 homeostasis. However, the detailed mechanism has not yet been elucidated.

ATG6 is the homologues of yeast Vps30/Atg6 and mammalian BECN1/Beclin1 (Xu *et al.*, 2017). It is a common and required subunit of the

class III phosphatidylinositol 3-kinase (PtdIns3K) lipid kinase complexes, which regulates autophagosome nucleation in *Arabidopsis thaliana* (*Arabidopsis*) (Qi *et al.*, 2017; Wang *et al.*, 2020). The homozygous *atg6* mutant is lethal, suggesting that ATG6 is essential for plant growth and development (Fujiki *et al.*, 2007; Qin *et al.*, 2007; Harrison-Lowe and Olsen, 2008; Patel and Dinesh-Kumar, 2008). In *Arabidopsis*, *N. benthamiana* and wheat, ATG6 or its homologues was reported to act as a positive regulator to prevent the invasion of *P. syringae pv. tomato* (*Pst*) DC3000 and *Pst* DC3000/*avrRpm1* bacteria (Patel and Dinesh-Kumar, 2008), *N. benthamiana* mosaic virus (TMV) (Liu *et al.*, 2005), turnip mosaic virus (TuMV) (Li *et al.*, 2018), pepper mild mottle virus (PMMoV) (Jiao *et al.*, 2020), and *Blumeria graminis f. sp. tritici (Bgt)* fungus (Yue *et al.*, 2015). Several research teams have also elucidated that ATG6 interacted with Bax Inhibitor-1 (NbBI-1) (Xu *et al.*, 2017) and RNA-dependent RNA polymerase (RdRp) (Li *et al.*, 2018) to suppress pathogen invasion. However, the mechanism by which ATG6 suppresses pathogen invasion by regulating NPR1 has not yet been reported.

Here, we show that ATG6 and NPR1 synergistically enhance *Arabidopsis* resistance to *Pst* DC3000/*avrRps4* infiltration. We discover that ATG6 increases NPR1 protein levels and nuclear accumulation of NPR1. Moreover, ATG6 can stabilize NPR1 and promote the formation of SINCs (SA-induced NPR1 condensates)-like condensates. Our study revealed a unique mechanism in which NPR1 cooperatively increases plant immunity with ATG6.

Results

NPR1 physically interacts with ATG6 in vitro and in vivo

To examine the relationship between ATGs and NPRs, we predicted that some ATGs might interact with NPRs. In a yeast two-hybrid (Y2H) screen, we identified that NPR1, NPR3 and NPR4 could interact with ATG6 and several ATG8 isoforms (**Fig. S1** and **Results S1** in the Supplemental Data 1). In this study, we mainly investigated the relationship between ATG6 and NPR1

during the process of plant immune response. Firstly, the NPR1 truncations NPR1-N (1-328AA, containing the BTB/POZ domain) and NPR1-C (328-594AA, containing the ANK domain and NLS) were used to identify the interaction domains between NPR1 and ATG6. The results showed that NPR1-C interacted with full-length ATG6 in yeast (**Fig. 1a**, line 3), indicating that the C terminus of NPR1 was necessary for the interaction between NPR1 and ATG6. The interaction between NPR1 and SnRK2.8 was used as a positive control (Lee *et al.*, 2015). Secondly, pull-down assays were performed *in vitro*. NPR1-His bound GST-ATG6, but not GST (**Fig. 1b**). Furthermore, co-immunoprecipitation (Co-IP) assays were performed in *N. benthamiana*, as shown in **Fig. 1c**, ATG6-mCherry was co-immunoprecipitated with NPR1-GFP. In **Fig. 1d**, fluorescence signals of NPR1-GFP and ATG6-mCherry were co-localized in the nucleus under normal and 1 mM SA treatment conditions. The interaction between ATG6 and NPR1 was also verified by a bimolecular fluorescence complementation (BiFC) assay (**Fig. S2**). These results demonstrate that ATG6 interacts with NPR1 both *in vitro* and *in vivo*.

ATG6 co-localized with NPR1 in the nucleus

Remarkably, under normal and SA treatment conditions, we found that ATG6 is localized in the cytoplasm and nucleus, and it co-localized with NPR1 in the nucleus (**Fig. 1d**). Nuclear localization of ATG6 was also observed in *N. benthamiana* transiently transformed with ATG6-mCherry and ATG6-GFP under normal and SA treatment conditions (**Fig. 2a** and **b**). ATG6-GFP co-localizes with the nuclear localization marker nls-mCherry (**Fig. 2b**). The nuclear localization signal of ATG6 was also observed in *UBQ10::ATG6-GFP* overexpressing *Arabidopsis* (**Fig. S3a**). To exclude the possibility that the observed localization of ATG6-GFP is due to free GFP. ATG6-GFP and free GFP levels in *UBQ10::ATG6-GFP Arabidopsis* and *N.benthamiana* were detected before and after SA treatment. Notably, no free GFP was detected and this means that the fluorescence signal observed by confocal microscopy

is ATG6-GFP, not free GFP (**Fig. S3d** and **e**).

Moreover, the nuclear and cytoplasmic fractions were separated. Under

SA treatment, ATG6-mCherry and ATG6-GFP were detected in the

cytoplasmic and nuclear fractions in *N. benthamiana* (**Fig. 2c and d**). Similarly,

ATG6 was also detected in the nuclear fraction of *UBQ10::ATG6-GFP* and

UBQ10::ATG6-mCherry overexpressing plants, and SA promoted both

cytoplasm and nuclear accumulation of ATG6 (**Fig. 2e and f**). Additionally, we

obtained *ATG6* and *NPR1* double overexpression of *Arabidopsis*

UBQ10::ATG6-mCherry × 35S::NPR1-GFP (*ATG6-mCherry ×NPR1-GFP*) by

crossing and screening (**Fig. S4a**). In *ATG6-mCherry × NPR1-GFP*, we

observed co-localization of ATG6-mCherry with NPR1-GFP in the nucleus

(**Fig. S3b**). These results are consistent with the prediction of the subcellular

location of ATG6 in the *Arabidopsis* subcellular database (https://suba.live/)

(**Fig. S3c**). Additionally, we have conducted an investigation into the

localization of endogenous ATG6 in Col. Our results demonstrate that

endogenous ATG6 is present in both the nucleus and cytoplasm, and we have

observed that SA treatment promotes the accumulation of ATG6 in the

nucleus (**Fig. S5**). Together, these findings suggest that ATG6 is localized to

both cytoplasm and nucleus, and co-localized with NPR1 in the nucleus.

ATG6 overexpression increased nuclear accumulation of NPR1

Previous studies have shown that the nuclear localization of NPR1 is

essential for improving plant immunity (Kinkema *et al.*, 2000; Chen *et al.*,

2021b). We observed that a stronger nuclear localization signal of NPR1-GFP

in *ATG6-mCherry × NPR1-GFP* leaves than that in *NPR1-GFP* under normal

condition and 0.5 mM SA treatment for 3 h (**Fig. 3a and b**). These findings

indicate that ATG6 might increase nuclear accumulation of NPR1. To exclude

the possibility that the observed localization of NPR1-GFP is due to free GFP,

we detected the levels of NPR1-GFP and free GFP in *ATG6-mCherry* x

NPR1-GFP plants before and after SA treatment. Only ~ 10 % of free GFP

was detected in *ATG6-mCherry* x *NPR1-GFP* plants before and after SA

treatment, confirming that the observed localization of NPR1-GFP is not due to free GFP (**Fig. S4b**). Furthermore, the nuclear and cytoplasmic fractions of *ATG6-mCherry × NPR1-GFP* and *NPR1-GFP* were separated. Under normal conditions, the nuclear fractions NPR1-GFP in *ATG6-mCherry × NPR1-GFP* and *NPR1-GFP* were relatively weaker (**Fig. 3c**), which differs from the above observation (**Fig. 3a**). We speculate that this phenomenon might be attributed to the rapid turnover of NPR1 in the nucleus (Spoel *et al.*, 2009; Saleh *et al.*, 2015). Consistent with the fluorescence distribution results, the nuclear fractions of NPR1-GFP in *ATG6-mCherry × NPR1-GFP* were significantly higher than those in *NPR1-GFP* under 0.5 mM SA treatment for 3 and 6 h (**Fig. 3c**). Furthermore, *Agrobacterium* harboring ATG6-mCherry and NPR1-GFP were transiently transformed to *N. benthamiana* leaves. After 1 d, the leaves were treated with 1 mM SA for 8 and 20 h. Subsequently nucleoplasmic separation experiments were performed. Similar to *Arabidopsis*, increased nuclear accumulation of NPR1 was found when *ATG6* was overexpressed (**Fig. 3e**). Notably, we found that the ratio (nucleus NPR1/total NPR1) in *ATG6-mCherry × NPR1-GFP* was not significantly different from that in *NPR1-GFP* after SA treatment, and a similar phenomenon was observed in *N. benthamiana* (**Fig. 3d and f**). These results suggested that the increased nuclear accumulation of NPR1 in *ATG6-mCherry* x *NPR1-GFP* plants might attributed to higher levels and more stable NPR1 rather than the enhanced nuclear translocation of NPR1 facilitated by ATG6. Furthermore, we validated the functionality of the ATG6-GFP and ATG6-mCherry fusion proteins utilized in this study by examining the phenotypes of *ATG6-GFP* and *ATG6-mCherry Arabidopsis* plants under carbon starvation conditions (**Fig. S6** and **Results S2** in the Supplemental Data 1).

ATG6 increases endogenous SA levels and promotes the expression of NPR1 downstream target genes

NPR1 localized in the nucleus is essential for activation of immune gene expression (Kinkema *et al.*, 2000; Chen *et al.*, 2021b). In our study, we

observed that *ATG6* overexpression increased nuclear accumulation of NPR1
(**Fig. 3**) and demonstrated an interaction between ATG6 and NPR1 in the
nucleus (**Fig. 1d**). Therefore, we speculate that ATG6 might regulate NPR1
transcriptional activity. Notably, the expression level of *ICS1* in *ATG6-mCherry*
× *NPR1-GFP* seedlings was significantly higher than that in *NPR1-GFP* under
normal and SA treatment conditions (**Fig. S7**). Free SA levels in *ATG6-*
mCherry × *NPR1-GFP* were also significantly higher compared to *NPR1-GFP*
under *Pst* DC3000/*avrRps4* treatment. While there was no significant
difference was observed under normal condition (**Fig. 4a**), this may be related
to free SA consumption, as it can be converted to bound SA (Ding and Ding,
2020). In addition, the expression of *PR1* (*pathogenesis-related* gene) and
PR5 in *ATG6-mCherry* × *NPR1-GFP* was significantly higher than that of
NPR1-GFP under normal and SA treatment conditions (**Fig. 4b and c**). The
expression of *PR1* and *PR5* in *ATG6-mCherry* was significantly higher than
that of Col under *Pst* DC3000/*avrRps4* treatment (**Fig. S8**). These results
support the role of ATG6 in facilitating the expression of NPR1 downstream
PR1 and *PR5* genes.

ATG6 increases NPR1 protein levels and the formation of SINCs-like
condensates

Interestingly, similar to previous reports (Zavaliev *et al.*, 2020), SA promoted
the translocation of NPR1 into the nucleus, but still a significant amount of
NPR1 was present in the cytoplasm (**Fig. 3c and e**). Previous studies have
shown that SA increased NPR1 protein levels and facilitated the formation of
SINCs in the cytoplasm, which are known to promote cell survival (Zavaliev *et*
al., 2020). In our experiments, we observed that under SA treatment, the
protein levels of NPR1 in *ATG6-mCherry* × *NPR1-GFP* was significantly
higher than that in *NPR1-GFP* (**Fig. 5a**). To further support our conclusions,
we proceeded to silence *ATG6* in *NPR1-GFP* (*NPR1-GFP*/silencing *ATG6*)
and subsequently assessed the protein level of NPR1-GFP before and after
SA treatment. Our findings revealed that the protein level of NPR1-GFP in

NPR1-GFP/silencing *ATG6* under SA treatment was notably lower than that in

the *NPR1-GFP*/Negative control (**Fig. S9**). Under SA treatment, the protein

levels of NPR1-GFP in *N. benthamiana* co-transformed with ATG6-mCherry +

NPR1-GFP was also significantly higher than that of mCherry + NPR1-GFP

(**Fig. 5b**). We also detected the expression of *NPR1* was detected. It is worth

noting that NPR1 up-regulation was more obvious in Col after 3 h treatment

with *Pst* DC3000/*avrRps4*. After 6 h treatment with *Pst* DC3000/*avrRps4*,

there was no significant difference in the expression of NPR1 between Col

and *ATG6-mCherry* (**Fig. S10**). These results suggest that ATG6 increases

NPR1 protein levels. After SA treatment, more SINCs-like condensates

fluorescence were observed in *N. benthamiana* co-transformed with ATG6-

mCherry + NPR1-GFP compared to mCherry + NPR1-GFP (**Fig. 5c-d and**

Supplemental movie 1-2). Taken together, these results suggest that ATG6

increases the protein levels of NPR1 and promotes the formation of SINCs-

like condensates, possibly caused by ATG6 increasing SA levels *in vivo*.

ATG6 maintains the protein stability of NPR1

Maintaining the stability of NPR1 is critical for enhancing plant immunity

(Skelly *et al.*, 2019). To further verify whether ATG6 regulates NPR1 stability,

we co-transfected NPR1-GFP with ATG6-mCherry or mCherry in *N.*

benthamiana and performed cell-free degradation assays. Our results showed

that NPR1-GFP degradation was significantly delayed when *ATG6* was

overexpressed (**Fig. S11**). A similar trend was observed in *Arabidopsis*, where

the NPR1-GFP protein in *ATG6-mCherry* × *NPR1-GFP* showed a slower

degradation rate compared to *NPR1-GFP* during 0~180 min time period in a

cell-free degradation assay (**Fig. 6a and b**). Moreover, when *Arabidopsis*

seedlings were treated with cycloheximide (CHX) to block protein synthesis,

we found that NPR1-GFP in *NPR1-GFP* was degraded after CHX treatment

for 3~9 h and the half-life of NPR1-GFP is ~3 h, while the half-life of NPR1-

GFP in *ATG6-mCherry* × *NPR1-GFP* is ~9 h (**Fig. 6c and d**). In addition, we

also analyzed the degradation of NPR1-GFP in *NPR1-GFP* and *NPR1-*
GFP/*atg5* following 100 µM cycloheximide (CHX) treatment. The results show
that the degradation rate of NPR1-GFP in *NPR1-GFP*/*atg5* plants was
similarly to that in *NPR1-GFP* plants (**Fig. 6e and f**). These results indicate
that ATG6 plays a role in maintaining the stability of NPR1, which may also be
related to the fact that ATG6 promotes an increase in free SA *in vivo*, since SA
has the function of increasing NPR1 stability (Ding *et al.*, 2016; Skelly *et al.*,
2019).

ATG6 and NPR1 jointly inhibit invasion of Pst DC3000/avrRps4

The mRNA expression levels of *ATG6* in Col were significantly increased after
6 h, 12 h and 24 h under *Pst* DC3000/*avrRps4* treatment (**Fig. 7a**). Similarly,
both the *ATG6* gene and protein were significantly up-regulated under 0.5 mM
SA treatment (**Fig. 7b and c**). These results suggest that the expression of
ATG6 could be induced by *Pst* DC3000/*avrRps4* and 0.5 mM SA treatment.

Considering that ATG6 increases NPR1 protein levels (**Fig. 5a-b**) and
promotes its nuclear accumulation (**Fig. 3**), as well as maintains NPR1
stability (**Fig. 6**), then we studied the role of ATG6-NPR1 interactions in plant
immunity. However, studying the function of ATG6 is challenging due to the
lethality of homozygous *atg6* mutant (Qin *et al.*, 2007; Harrison-Lowe and
Olsen, 2008; Patel and Dinesh-Kumar, 2008). According to our previous
report (Lei *et al.*, 2020; Zhang *et al.*, 2023), *ATG6* was silenced using artificial
miRNA^{ATG6} (amiRNA^{ATG6}) delivered by the gold nanoparticles (AuNPs). First,
the effect of *ATG6* silencing in Col on the plant immune response was
investigated. Similar to *atg5*, Col/silencing *ATG6* exhibited more active growth
of *Pst* DC3000/*avrRps4* than Col/negative control (NC) after *Pst*
DC3000/*avrRps4* infiltration for 3 days (**Fig. 7d**). Furthermore, according to
the previously reported methods (Ohira *et al.*, 2017; Gomez *et al.*, 2022), we
generated two amiRNA^{ATG6} lines (amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2)
designed against *ATG6* and placed under the control of a β-estradiol inducible

promoter. There were no significant phenotypic differences in amiRNA^{ATG6} # 1

compared to the Col, while amiRNA^{ATG6} # 2 exhibited a slight leaf

developmental defect (**Fig. 7e** and **f**). Subsequently, we investigated the

expression of *ATG6* following treatment with 100 µM β-estradiol. Our results

showed that, after 100 µM β-estradiol treatment for 1~3 d, the expression of

ATG6 in both amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2 lines was significantly

lower than that in Col. Specifically, the expression of *ATG6* in the amiRNA^{ATG6}

#1 and amiRNA^{ATG6} *#2* lines decreased by 50~70% compared with Col (**Fig.**

7g). Furthermore, to assess the function of ATG6 in plant immune, we

performed infiltrations of *Pst* DC3000/*avrRps4* after 100 uM β-estradiol

treatment for 24 h. We compared the growth of *Pst* DC3000/*avrRps4* in the

amiRNA^{ATG6} lines and Col. The results clearly demonstrate that the growth of

Pst DC3000/*avrRps4* in amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2 was significantly

more compared to Col (**Fig. 7h**). Moreover, we silenced *ATG6* in *NPR1-GFP*

(*NPR1-GFP*/silencing *ATG6*), and *NPR1-GFP*/*atg5* (crossed *NPR1-GFP* with

atg5 to obtain *NPR1-GFP*/*atg5*) was used as an autophagy-deficient control.

There was more *Pst* DC3000/*avrRps4* growth in *NPR1-GFP*/silencing *ATG6*

and *NPR1-GFP*/*atg5* compared to *NPR1-GFP*/NC after *Pst* DC3000/*avrRps4*

infiltration (**Fig. 7i**). In contrast, the growth of *Pst* DC3000/*avrRps4* in *NPR1-*

GFP, *ATG6-mCherry*, *ATG6-mCherry* × *NPR1-GFP* was significantly lower

than that in Col and *npr1* (**Fig. 7j**) and was the lowest in *ATG6-mCherry* ×

NPR1-GFP (**Fig. 7j**).

These results confirm that ATG6 and NPR1 jointly enhance *Arabidopsis*

resistance to inhibit *Pst* DC3000/*avrRps4* invasion. Together, these results

suggest that ATG6 improves plant resistance to pathogens by regulating

NPR1.

Discussion

318   Although SA signaling and autophagy are related to the plant immune

319   system (Yoshimoto *et al.*, 2009; Munch *et al.*, 2014; Wang *et al.*, 2016), the

connection of these two processes in plant immune processes and their interaction is rarely reported. Previous studies have shown that unrestricted pathogen-induced PCD requires SA signalling in autophagy-deficient mutants. SA and its analogue benzo (1,2,3) thiadiazole-7-carbothioic acid (BTH) induce autophagosome production (Yoshimoto *et al.*, 2009). Moreover, autophagy has been shown to negatively regulates *Pst* DC3000/*avrRpm1*-induced PCD via the SA receptor NPR1 (Yoshimoto *et al.*, 2009), implying that autophagy regulates SA signaling through a negative feedback loop to limit immune-related PCD. Here, we demonstrated that ATG6 increases NPR1 protein levels and nuclear accumulation (**Fig. 3** and **5**). Additionally, ATG6 also maintains the stability of NPR1 and promotes the formation of SINCs-like condensates (**Fig. 5** and **6**). These findings introduce a novel perspective on the positive regulation of NPR1 by ATG6, highlighting their synergistic role in enhancing plant resistance.

Our results confirmed that *ATG6* overexpression significantly increased nuclear accumulation of NPR1 (**Fig. 3**). ATG6 also increases NPR1 protein levels and improves NPR1 stability (**Fig. 5 and 6**). Therefore, we consider that the increased nuclear accumulation of NPR1 in *ATG6-mCherry* x *NPR1-GFP* plants might result from higher levels and more stable NPR1 rather than the enhanced nuclear translocation of NPR1 facilitated by ATG6. To verify this possibility, we determined the ratio of NPR1-GFP in the nuclear localization versus total NPR1-GFP. Notably, the ratio (nucleus NPR1/total NPR1) in *ATG6-mCherry* × *NPR1-GFP* was not significantly different from that in *NPR1-GFP*, and there is a similar phenomenon in *N. benthamiana* (**Fig. 3c-f**). Further we analyzed whether ATG6 affects NPR1 protein levels and protein stability. Our results show that ATG6 increases NPR1 protein levels under SA treatment and ATG6 maintains the protein stability of NPR1 (**Fig. 5 and 6**). These results suggested that the increased nuclear accumulation of NPR1 by ATG6 result from higher levels and more stable NPR1.

NPR1 is an important signaling hub of the plant immune response.

Nuclear localization of NPR1 is essential to enhance plant resistance (Kinkema *et al.*, 2000; Chen *et al.*, 2021b), it interacts with transcription factors such as TGAs in the nucleus to activate expression of downstream target genes (Chen *et al.*, 2019; Chen *et al.*, 2021a). A recent study showed that nuclear-located ATG8h recognizes C1, a geminivirus nuclear protein, and promotes C1 degradation through autophagy to limit viral infiltration in solanaceous plants (Li *et al.*, 2020). Here, we confirmed that ATG6 is also distributed in the nucleus and ATG6 is co-localized with NPR1 (**Fig. 1d** and **2**), suggesting that ATG6 interact with NPR1 in the nucleus. ATG6 synergistically inhibits the invasion of *Pst* DC3000/*avrRps4* with NPR1. Chen et al. found that in the nucleus, NPR1 can recruit enhanced disease susceptibility 1 (EDS1), a transcriptional coactivator, to synergistically activate expression of downstream target genes (Chen *et al.*, 2021a). Previous studies have shown that acidic activation domains (AADs) in transcriptional activators (such as Gal4, Gcn4 and VP16) play important roles in activating downstream target genes. Acidic amino acids and hydrophobic residues are the key structural elements of AAD (Pennica *et al.*, 1984; Cress and Triezenberg, 1991; Van Hoy *et al.*, 1993). Chen et al. found that EDS1 contains two ADD domains and confirmed that EDS1 is a transcriptional activator with AAD (Chen *et al.*, 2021a). Here, we also have similar results that *ATG6* overexpression significantly enhanced the expression of *PR1* and *PR5* (**Fig. 4b-c and S6**), and that the ADD domain containing acidic and hydrophobic amino acids is also found in ATG6 (148-295 AA) (**Fig. S12**). We speculate that ATG6 might act as a transcriptional coactivator to activate *PRs* expression synergistically with NPR1.

A recent study showed that SA not only enhances plant resistance by increasing NPR1 nuclear import and transcriptional activity, but also promotes cell survival by coordinating the distribution of NPR1 in the nucleus and cytoplasm (Zavaliev *et al.*, 2020). Notably, NPR1 accumulated in the cytoplasm recruits other immunomodulators (such as EDS1, PAD4 etc) to

form SINCs to promote cell survival (Zavaliev *et al.*, 2020). Similarly, we also

found that NPR1 accumulated abundantly in the cytoplasm after SA treatment

and that ATG6 significantly increased NPR1 protein levels (**Fig. 3c, 3e and**

5a-b). Obviously, the accumulation of NPR1 in the cytoplasm may be related

to ATG6 synergizing with NPR1 to enhance plant resistance. Interestingly,

ATG6 overexpression significantly increased the formation of SINCs-like

condensates (**Fig. 5c-d** and **Supplemental movie 1-2**), which should also be

a way for ATG6 and NPR1 to synergistically resist invasion of pathogens. We

consider that ATG6 promotes the formation of SINCs-like condensates

through the dual action of endogenous and exogenous SA. Considering that

ATG6 promotes SINCs-like condensates formation, we further examined

changes in cell death in Col, amiRNA^{ATG6} # 1, amiRNA^{ATG6} # 2, *npr1*, *NPR1-*

GFP, *ATG6-mCherry* and *ATG6-mCherry × NPR1-GFP* plants. The results of

Taipan blue staining showed that *Pst* DC3000/*avrRps4*-induced cell death in

npr1, amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2 was significantly higher compared

to Col (**Fig. S13**). Conversely, *Pst* DC3000/*avrRps4*-induced cell death in

ATG6-mCherry, *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* was

significantly lower compared to Col. Notably, *Pst* DC3000/*avrRps4*-induced

cell death in *ATG6-mCherry × NPR1-GFP* was significantly lower compared

ATG6-mCherry and *NPR1-GFP* (**Fig. S13**). These results suggest that ATG6

and NPR1 jointly inhibit *Pst* DC3000/*avrRps4*-induced cell dead.

ATG6 is a common and required subunit of PtdIns3K lipid kinase

complexes, which regulates autophagosome nucleation in *Arabidopsis* (Qi *et*

al., 2017; Bozhkov, 2018). In this study, we also found that ATG6 can

maintain the stability of NPR1. Thus, to confirm whether the regulation of

NPR1 protein stability by ATG6 is autophagy-dependent, we used autophagy

inhibitors (Concanamycin A, ConA and Wortmannin, WM) to detect the

degradation of NPR1-GFP. Cell-free degradation assays showed that 100 µM

MG115 treatment significantly inhibited the degradation of NPR1-GFP.

However, 5 µM concanamycin A and 30 µM Wortmannin treatment did not

significantly delay NPR1 degradation (**Fig. S14**). In addition, we also analyzed the degradation of NPR1-GFP in *NPR1-GFP* and *NPR1-GFP*/*atg5* following 100 μM cycloheximide (CHX) treatment. The results show that the degradation rate of NPR1-GFP in *NPR1-GFP*/*atg5* plants was similarly to that in *NPR1-GFP* plants (**Fig. 6e and f**). These results suggest that autophagy defects do not affect the protein stability of NPR1, and ATG6 promoting NPR1 protein stability maybe its unique autophagy-independent role.

An increasing number of studies have shown that ATGs differentially affect plant immunity. Deletion of ATGs (ATG5, ATG7, ATG10 etc.) leads to reduced resistance of plants to necrotrophic pathogens (Lai *et al.*, 2011; Lenz *et al.*, 2011; Minina *et al.*, 2018). ATGs can directly interact with other proteins to positively regulate plant immunity. In *N. benthamiana*, ATG8f interacts the effector protein βC1 of the cotton *leaf curl multan virus* and promotes its degradation to limit pathogen invasion (Haxim *et al.*, 2017). Notably, ATG18a can interact with WRKY33 transcription factor to synergistically against *Botrytis* invasion (Lai *et al.*, 2011). Our evidence shows that ATG6 interacts with NPR1 and works together to counteract pathogen invasion by positively regulating NPR1 and SA levels *in vivo*. In conclusion, we unveil a novel relationship in which ATG6 positively regulates NPR1 in plant immunity (**Fig. 8**). ATG6 interacts with NPR1 to synergistically enhance plant resistance by regulating NPR1 protein levels, stability, nuclear accumulation, and formation of SINCs-like condensates.

Material and Methods

Plasmid construction

Details of plasmid construction methods are listed in Methods S1 of Supplemental Data 1, primer used are listed in Table S1 and S2 of Supplemental Data 1. The mapping of vectors is listed in Supplemental Data 2.

Plant material

Arabidopsis

35S::NPR1-GFP (in *npr1-2* background) and *npr1-1* were kindly provided by

Dr. Xinnian Dong of Duke University; *atg5-1* (SALK_020601).

UBQ10::ATG6-mCherry, *UBQ10::ATG6-GFP and* amiRNA*^{ATG6}* lines were

obtained by *Agrobacterium* transformation (Clough and Bent, 1998). *ATG6*,

NPR1 double overexpression of *Arabidopsis* (*ATG6-mCherry* × *NPR1-GFP*)

and *NPR1-GFP*/*atg5* were obtained by crossing respectively.

Full description of the *Arabidopsis* screening is included Methods S2 of

Supplemental Data 1. Details of plant material are listed in Table S3 of

Supplemental Data 1.

Growth conditions

Arabidopsis thaliana

All *Arabidopsis thaliana* (*Arabidopsis*) seeds were treated in 10% sodium

hypochlorite for 7 min, washed with ddH_2O, and treated in 75% ethanol for 30

s, finally washed three times with ddH_2O. Seeds were sown in 1/2 MS with 2%

sucrose solid medium, vernalized at 4℃ for 2 days.

For 7-day-old *Arabidopsis* seedling cultures, the plates were placed under

the following conditions: daily cycle of 16 h light (~80 $\mu molm^{-2} \cdot s^{-1}$) and 8 h

dark at 23 ± 2℃.

For 3-week-old *Arabidopsis* cultures, after 7 days of growth on the plates,

the seedlings were transferred to soil for further growth for 2 weeks under the

same conditions (Zhang *et al.*, 2018).

N. benthamiana

For 3-week-old *N. benthamiana* cultures, seeds were sown in the soil and

vernalized at 4℃ for 2 days. After 10 days of growth on soil , the seedlings

were transferred to soil for further growth for 2 weeks under the same

conditions (Jiao *et al.*, 2019).

Treatment conditions

Treatment of 7-day-old seedlings

For SA treatment

7-day-old *Arabidopsis* seedlings were transferred to 1/2 MS liquid medium

containing 0.5 mM SA for 0, 3 and 6 h, respectively. The corresponding results

are shown in Fig. 2e, f, 3c, d and 5a.

For cycloheximide (CHX) treatment

473 Seedlings of *Arabidopsis* (7 days) were transferred to 1/2 MS liquid medium

474 containing 100 µM CHX for 0, 3, 6 and 9 h, respectively. The corresponding

475 results are shown in Fig. 6c and e.

Treatment of 3-week-old Arabidopsis

For silencing ATG6 in Col and NPR1-GFP

As previously described (Lei *et al.*, 2020; Zhang *et al.*, 2022; Zhang *et al.*,

2023), 1 mM gold nanoparticles (AuNPs) were synthesized. The artificial

microRNA (amiRNA)^{ATG6} (UCAAUUCUAGGAUAACUGCCC) was designed

based on the Web MicroRNA Designer (http://wmd3.weigelworld.org/)

platform. The complementary sequence of amiRNA^{ATG6} is located on the

eighth exon of the *ATG6* gene. The sequence of

"UUCUCCGAACGUGUCACGUTT" was used as a negative control (NC). NC

is a universal negative control without species specificity (Gao *et al.*, 2018; Lei

et al., 2020). amiRNA^{ATG6} and amiRNA^{NC} synthesized by Suzhou

GenePharma. AuNPs (1 mM) and amiRNA^{ATG6} (20 µM) were incubated at a

9:1 ratio for 30 min at 25℃, 50 rpm. After incubation, a mixture of AuNPs and

amiRNA^{ATG6} was diluted 15-fold with the infiltration buffer (pH 5.7, 10 mM

MES, 10 mM MgCl$_2$) and infiltrated through the abaxial leaf surface into 3-

week-old Col or *NPR1-GFP* for 1-3 days. The third day was chosen as

material for *ATG6* silencing. After the third day of AuNPs-amiRNA^{ATG6} and

AuNPs-amiRNA^{NC} infiltration, *Pst* DC3000/*avrRps4* was infiltrated, and then

growth of *Pst* DC3000/*avrRps4* was detected.

For β-estradiol treatment

496 100 µM β-estradiol was infiltrated to treat 3-week-old *Arabidopsis* leaves. After

497 24 h of treatment with β-estradiol, *Pst* DC3000/*avrRps4* was infiltrated and

498 then growth of *Pst* DC3000/*avrRps4* was detected after 3 d.

For Pst DC3000/avrRps4 infiltration

Infiltration with *Pst* DC3000/*avrRps4* was performed as previously described (Wang *et al.*, 2016; Skelly *et al.*, 2019). Full description of the *Pst* DC3000/*avrRps4* culture is included in Methods S3 of Supplemental Data 1.

For SA treatment

For 3-week-old Col, 0.5 mM SA was infiltrated into the leaves for 0, 2, 4, 6 and 8 h. The corresponding results are shown in Fig. 7b and c.

Yeast two-hybrid assay

Yeast two-hybrid experiments were performed according to the previously described protocol (Fu *et al.*, 2012). Full description of Yeast two-hybrid is included in Methods S4 of Supplemental Data 1.

Pull down assays in vitro

500 µL of GST, GST-ATG6, SnRK2.8-GST were incubated with GST-tag Purification Resin (Beyotime, P2250) for 2 h at 4°C. The mixture was then centrifuged at 1500 g for 1 min at 4°C, and the resin was washed three times with PBS buffer. Next, the GST-tag purification resin was incubated with the NPR1-His for 2 h at 4°C. After washing three times with PBS buffer, 2 × sample buffers were added to the resin and denatured at 100 °C for 10 mins. The resulting samples were then used for western blotting analysis. Full description of prokaryotic proteins expression is included in Methods S5 of Supplemental Data 1.

Co-immunoprecipitation

0.5 g leaves of *N. benthamiana* transiently transformed with ATG6-mCherry + GFP and ATG6-mCherry + NPR1-GFP were fully ground in liquid nitrogen and homogenized in 500 µL of lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 5% Glycerol, 0.2% NP40, 1 mM PMSF, 40 µM MG115, protease inhibitor cocktail 500× and phosphatase inhibitor cocktail 5000×). The samples were then incubated on ice for 30 mins, and centrifuged at 10142 g (TGL16, cence, hunan, China) for 15 mins at 4°C. The supernatant (500 µL) was incubated with 20 µL of GFP-Trap Magnetic Agarose beads (ChromoTek, gtma-20) in a 1.5 mL Eppendorf tube for 2 h by rotating at 4°C.

After incubation, the GFP-Trap magnetic Agarose beads were washed three times with cold wash buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA) and denatured at 75°C for 10 minutes after adding 2 × sample buffer. Western blotting was performed with antibodies to ATG6 and GFP.

Nuclear and cytoplasmic separation

Nuclear and cytoplasmic separation were performed according to the previously described method (Kinkema *et al.*, 2000). Full description of nuclear and cytoplasmic separation is given in Methods S6 of Supplemental Data 1.

Protein degradation in vitro

Protein degradation assays were performed according to a previously described method (Spoel *et al.*, 2009; Saleh *et al.*, 2015). Full description of protein degradation is included in Methods S7 of Supplemental Data 1.

Protein Extraction and Western Blotting Analysis

Protein extraction and western blotting were performed as previously described (Lei *et al.*, 2020; Zhang *et al.*, 2022). Protein was denatured at 100°C for 10 mins. NPR1 protein was denatured at 75°C for 10 mins (Lei *et al.*, 2020). Full description is included in Methods S8 of Supplemental Data 1. Antibody information is presented in Table S4 of Supplemental Data 1.

Confocal microscope observation

For nuclear localization of NPR1-GFP observation

7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* were sprayed with 0.5 mM SA for 0 and 3 h. GFP and mCherry fluorescence signals in leaves were observed under the confocal microscope (Zeiss LSM880). At least 20 image sets were obtained and analyzed.

For the Bimolecular Bluorescence Complementation assay (BiFC)

Agrobacterium was infiltrated into *N. benthamiana* as previously described (Jiao *et al.*, 2019). Fluorescence signals were observed after 3 days. The full description of BiFC is contained in Methods S9 and S10 of Supplemental Data 1.

For the observation of SINCs-like condensates

Agrobacterium was infiltrated into *N. benthamiana*. After 2 days, the leaves were treated in 1 mM SA solution for 24 h, and then fluorescence signals were observed. At least 20 image sets were obtained and analyzed. A full description of SINCs-like condensates observation is included in Methods S11 of Supplemental Data 1.

For growth of Pst DC3000/avrRps4

A low dose (OD_{600} = 0.001) of *Pst* DC3000/*avrRps4* was used for the infiltration experiments. After 3 days, the colony count was counted according to a previous description (Wang *et al.*, 2016; Lei *et al.*, 2020). Full description of the growth of *Pst* DC3000/*avrRps4* is given in Methods S12 of Supplemental Data 1.

Free SA measurement

Free SA was extracted from 3-week-old *Arabidopsis* using a previously described method (Wang *et al.*, 2016; Gong *et al.*, 2020). Free SA was measured by High-performance liquid chromatography (Shimadzu LC-6A, Japan). Detection conditions: 294 nm excitation wavelength, 426 nm emission wavelength.

Real-Time Quantitative PCR (RT-qPCR)

Total RNA was extracted from *Arabidopsis* (100 mg) using Trizol RNA reagent (Invitrogen, 10296-028, Waltham, MA, USA). RT-qPCR assays were performed as previously described (Zhang *et al.*, 2018; Zhang *et al.*, 2022). All primers for RT-qPCR are listed individually in Table S5 of Supplemental Data 1. Full description of RT-qPCR is included in Methods S13 of Supplemental Data 1.

Trypan Blue Staining

The leaves of 3-week-old Col, amiRNA^{ATG6} # 1, amiRNA^{ATG6} # 2, *npr1*, *NPR1-GFP*, *ATG6-mCherry* and *ATG6-mCherry* × *NPR1-GFP* plants, located in the fifth and sixth positions, were infiltrated with *Pst* DC3000/*avrRps4*. After 3 days, the leaves were excised and subjected to a 1 min boiling step in trypan

blue staining buffer (consisting of 10 g phenol, 10 mL glycerol, 10 mL lactic acid, 10 mL ddH$_2$O, and 10 mg trypan blue), followed by destaining 3 times at 37℃ in 2.5 mg/mL chloral hydrate.

Statistical Analysis

All quantitative data in this study were presented as mean ± SD. The experimental data were analyzed by a two-tailed Student's *t*-test. Significance was assigned at *P* values < 0.05 or < 0.01.

Acknowledgments: We thank Dr. Xinnian Dong (Duke University, USA), Dr. ZhengQing Fu (University of South Carolina) and Dr. Sheng Li (South China Normal University) for their help and contribution.

Author contribution: B.-H.Z., J.Z. and W.-L.C. designed and supervised the research; B.-H.Z. and S.-Q.H. performed majority experiments; B.-H.Z., Y.-X.M. and H.C. synthesized gold nanoparticles (AuNPs); B.-H.Z. and S.-Q.H., Y.-Z.T. performed western blotting analysis; B.-H.Z.,Y.Z and X.L performed confocal experiment; B.-H.Z. and S.-Q.H. analyzed the data; B.-H.Z. and S.-Q.H. prepared figures; B.-H.Z. and S.-Q.H. wrote the manuscript; J.Z. and W.-L.C. supervised and completed the writing; S.-Y.G. drew working model and edited language. J.Z. and W.-L.C. provided reagents/materials/analysis tools; J.Z. and W.-L.C. agrees to serve as the corresponding author responsible for contact and ensures communication.

Conflicts of interest: The author declares that there are no conflicts of interest.

Funding: This research was supported by the National Natural Science Foundation of China [Grant Number 31570256].

Data availability: The authors confirm that the data supporting the results of this study are available in the article and its supplementary materials.

Reference

Bozhkov PV. 2018. Plant autophagy: mechanisms and functions. J Exp Bot **69**, 1281-1285.

Cao H, Glazebrook J, Clarke JD, et al. 1997. The Arabidopsis NPR1 gene

that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. Cell **88**, 57-63.

Chen H, Li M, Qi G, et al. 2021a. Two interacting transcriptional coactivators cooperatively control plant immune responses. Sci Adv **7**, eabl7173.

Chen J, Mohan R, Zhang Y, et al. 2019. NPR1 Promotes Its Own and Target Gene Expression in Plant Defense by Recruiting CDK8. Plant Physiol **181**, 289-304.

Chen J, Zhang J, Kong M, et al. 2021b. More stories to tell: NONEXPRESSOR OF PATHOGENESIS-RELATED GENES1, a salicylic acid receptor. Plant Cell Environ.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J **16**, 735-743.

Cress WD, Triezenberg SJ. 1991. Critical structural elements of the VP16 transcriptional activation domain. Science **251**, 87-90.

Ding P, Ding Y. 2020. Stories of Salicylic Acid: A Plant Defense Hormone. Trends Plant Sci **25**, 549-565.

Ding Y, Dommel M, Mou Z. 2016. Abscisic acid promotes proteasome-mediated degradation of the transcription coactivator NPR1 in Arabidopsis thaliana. Plant J **86**, 20-34.

Fu ZQ, Yan S, Saleh A, et al. 2012. NPR3 and NPR4 are receptors for the immune signal salicylic acid in plants. Nature **486**, 228-232.

Fujiki Y, Yoshimoto K, Ohsumi Y. 2007. An Arabidopsis homolog of yeast ATG6/VPS30 is essential for pollen germination. Plant Physiol **143**, 1132-1139.

Gao L, Wu K, Liu Z, et al. 2018. Chromatin Accessibility Landscape in Human Early Embryos and Its Association with Evolution. Cell **173**, 248-259 e215.

Gomez RE, Chambaud C, Lupette J, et al. 2022. Phosphatidylinositol-4-phosphate controls autophagosome formation in Arabidopsis thaliana. Nature Communications **13**.

Gong W, Li B, Zhang B, et al. 2020. ATG4 Mediated Psm ES4326/AvrRpt2-

Induced Autophagy Dependent on Salicylic Acid in Arabidopsis Thaliana. Int J

Mol Sci **21**.

Harrison-Lowe NJ, Olsen LJ. 2008. Autophagy protein 6 (ATG6) is required

for pollen germination in Arabidopsis thaliana. Autophagy **4**, 339-348.

Haxim Y, Ismayil A, Jia Q, et al. 2017. Autophagy functions as an antiviral

mechanism against geminiviruses in plants. Elife **6**.

Jiao Y, Srba M, Wang JC, et al. 2019. Correlation of Autophagosome

Formation with Degradation and Endocytosis Arabidopsis Regulator of G-

Protein Signaling (RGS1) through ATG8a. International Journal of Molecular

Sciences **20**.

Jiao YB, An MN, Li XD, et al. 2020. Transcriptomic and functional analyses

reveal an antiviral role of autophagy during pepper mild mottle virus infection.

Bmc Plant Biology **20**.

Jones JD, Dangl JL. 2006. The plant immune system. Nature **444**, 323-329.

Kikuchi Y, Nakamura S, Woodson JD, et al. 2020. Chloroplast Autophagy

and Ubiquitination Combine to Manage Oxidative Damage and Starvation

Responses. Plant Physiol **183**, 1531-1544.

Kinkema M, Fan W, Dong X. 2000. Nuclear localization of NPR1 is required

for activation of PR gene expression. Plant Cell **12**, 2339-2350.

Lai Z, Wang F, Zheng Z, et al. 2011. A critical role of autophagy in plant

resistance to necrotrophic fungal pathogens. Plant J **66**, 953-968.

Lee HJ, Park YJ, Seo PJ, et al. 2015. Systemic Immunity Requires SnRK2.8-

Mediated Nuclear Import of NPR1 in Arabidopsis. Plant Cell **27**, 3425-3438.

Lei WX, An ZS, Zhang BH, et al. 2020. Construction of gold-siRNA(NPR1)

nanoparticles for effective and quick silencing of NPR1 in Arabidopsis thaliana.

Rsc Advances **10**, 19300-19308.

Lenz HD, Haller E, Melzer E, et al. 2011. Autophagy differentially controls

plant basal immunity to biotrophic and necrotrophic pathogens. Plant J **66**,

818-830.

Li F, Zhang C, Li Y, et al. 2018. Beclin1 restricts RNA virus infection in plants

through suppression and degradation of the viral polymerase. Nat Commun **9**, 681 1268.

Li F, Zhang M, Zhang C, et al. 2020. Nuclear autophagy degrades a 683 geminivirus nuclear protein to restrict viral infection in solanaceous plants. 684 New Phytol **225**, 1746-1761.

Liu Y, Schiff M, Czymmek K, et al. 2005. Autophagy regulates programmed 686 cell death during the plant innate immune response. Cell **121**, 567-577.

Marshall RS, Li FQ, Gemperline DC, et al. 2015. Autophagic Degradation of 688 the 26S Proteasome Is Mediated by the Dual ATG8/Ubiquitin Receptor 689 RPN10 in Arabidopsis. Molecular Cell **58**, 1053-1066.

Minina EA, Moschou PN, Vetukuri RR, et al. 2018. Transcriptional 691 stimulation of rate-limiting components of the autophagic pathway improves 692 plant fitness. J Exp Bot **69**, 1415-1432.

Munch D, Rodriguez E, Bressendorff S, et al. 2014. Autophagy deficiency 694 leads to accumulation of ubiquitinated proteins, ER stress, and cell death in 695 Arabidopsis. Autophagy **10**, 1579-1587.

Ohira MJ, Hendrickson DG, McIsaac RS, et al. 2017. An estradiol-inducible 697 promoter enables fast, graduated control of gene expression in fission yeast. 698 Yeast **34**, 323-334.

Patel S, Dinesh-Kumar SP. 2008. Arabidopsis ATG6 is required to limit the 700 pathogen-associated cell death response. Autophagy **4**, 20-27.

Pennica D, Goeddel DV, Hayflick JS, et al. 1984. The amino acid sequence 702 of murine p53 determined from a c-DNA clone. Virology **134**, 477-482.

Qi H, Xia FN, Xie LJ, et al. 2017. TRAF Family Proteins Regulate Autophagy 704 Dynamics by Modulating AUTOPHAGY PROTEIN6 Stability in Arabidopsis. 705 Plant Cell **29**, 890-911.

Qin G, Ma Z, Zhang L, et al. 2007. Arabidopsis AtBECLIN 1/AtAtg6/AtVps30 707 is essential for pollen germination and plant development. Cell Res **17**, 249-708 263.

Rochon A, Boyle P, Wignes T, et al. 2006. The coactivator function of

Arabidopsis NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal cysteines. Plant Cell **18**, 3670-3685.

Saleh A, Withers J, Mohan R, et al. 2015. Posttranslational Modifications of the Master Transcriptional Regulator NPR1 Enable Dynamic but Tight Control of Plant Immune Responses. Cell Host Microbe **18**, 169-182.

Skelly MJ, Furniss JJ, Grey H, et al. 2019. Dynamic ubiquitination determines transcriptional activity of the plant immune coactivator NPR1. Elife **8**.

Spoel SH, Mou Z, Tada Y, et al. 2009. Proteasome-mediated turnover of the transcription coactivator NPR1 plays dual roles in regulating plant immunity. Cell **137**, 860-872.

Tada Y, Spoel SH, Pajerowska-Mukhtar K, et al. 2008. Plant immunity requires conformational changes [corrected] of NPR1 via S-nitrosylation and thioredoxins. Science **321**, 952-956.

Van Hoy M, Leuther KK, Kodadek T, et al. 1993. The acidic activation domains of the GCN4 and GAL4 proteins are not alpha helical but form beta sheets. Cell **72**, 587-594.

Wang P, Nolan TM, Yin Y, et al. 2020. Identification of transcription factors that regulate ATG8 expression and autophagy in Arabidopsis. Autophagy **16**, 123-139.

Wang XD, Gao YY, Yan QQ, et al. 2016. Salicylic acid promotes autophagy via NPR3 and NPR4 in Arabidopsis senescence and innate immune response. Acta Physiologiae Plantarum **38**.

Wu Y, Zhang D, Chu JY, et al. 2012. The Arabidopsis NPR1 protein is a receptor for the plant defense hormone salicylic acid. Cell Rep **1**, 639-647.

Xu G, Wang S, Han S, et al. 2017. Plant Bax Inhibitor-1 interacts with ATG6 to regulate autophagy and programmed cell death. Autophagy **13**, 1161-1175.

Yoshimoto K, Jikumaru Y, Kamiya Y, et al. 2009. Autophagy negatively regulates cell death by controlling NPR1-dependent salicylic acid signaling during senescence and the innate immune response in Arabidopsis. Plant Cell

21, 2914-2927.

Yue J, Sun H, Zhang W, et al. 2015. Wheat homologs of yeast ATG6 function

in autophagy and are implicated in powdery mildew immunity. BMC Plant Biol

15, 95.

Zavaliev R, Mohan R, Chen T, et al. 2020. Formation of NPR1 Condensates

Promotes Cell Survival during the Plant Immune Response. Cell **182**, 1093-

1108 e1018.

Zhang B, Huang S, Meng Y, et al. 2023. Gold nanoparticles (AuNPs) can

rapidly deliver artificial microRNA (AmiRNA)-ATG6 to silence ATG6

expression in Arabidopsis. Plant Cell Rep **42**, 1191-1201.

Zhang B, Su L, Hu B, et al. 2018. Expression of AhDREB1, an AP2/ERF

Transcription Factor Gene from Peanut, Is Affected by Histone Acetylation and

Increases Abscisic Acid Sensitivity and Tolerance to Osmotic Stress in

Arabidopsis. Int J Mol Sci **19**.

Zhang J, Lei W, Meng Y, et al. 2022. Expression of PEI-coated gold

nanoparticles carrying exogenous gene in periwinkle mesophyll cells and its

practice in huanglongbing research. iScience **25**, 104479.

Zhou J, Zhang Y, Qi J, et al. 2014. E3 ubiquitin ligase CHIP and NBR1-

mediated selective autophagy protect additively against proteotoxicity in plant

stress responses. PLoS Genet **10**, e1004116.

Figure Legends

Figure 1. Physical interaction between NPR1 and ATG6. (**a**). Interaction of

NPR1 with ATG6 in yeast. The CDS of *ATG6*, *NPR1*, *NPR1-N* (1~984 bp),

NPR1-C (984~1782 bp) and *SnRK2.8* were fused to pGADT7 (AD) and

pGBKT7 (BD), respectively. Co-transformation of NPR1-BD + AD, BD +

ATG6-AD, BD + SnRK2.8-AD, NPR1-N-BD + AD, NPR1-C-BD + AD were

used as negative controls. The interaction of NPR1-BD and SnRK2.8-AD was

used as a positive control. Yeast growth on SD/-Trp-Leu-His-Ade media

represents interaction. Numbers represent the dilution fold of yeast. 0, -1 (10

fold dilution), -2 (100 fold dilution), -3 (1000 fold dilution). (**b**). In vitro pull-
down assays of NPR1-His with GST-ATG6 fusion protein. NPR1-His
prokaryotic proteins were incubated with GST-tag Purification Resin
conjugated with GST-ATG6, GST and SnRK2.8-GST. Western blotting
analysis with anti-GST and anti-His. Black asterisk indicate SnRK2.8-GST
bands. Red asterisk indicate GST-ATG6 bands. (**c**). Co-immunoprecipitation
of NPR1 with ATG6 in vivo. Total protein was extracted from *N. benthamiana*
transiently transformed with ATG6-mCherry + GFP and ATG6-mCherry +
NPR1-GFP, followed by IP with GFP-Trap. Western blots analysis with ATG6
and GFP antibodies. (**d**). Co-localization of NPR1-GFP and ATG6-mCherry in
N. benthamiana under normal and SA treatment conditions. NPR1-GFP were
co-expressed with ATG6-mCherry in *N. benthamiana* for 2 d followed by
confocal observation. Scale bar, 100 μm. All experiments were performed with
three biological replicates.

Figure 2. ATG6 is localized in the cytoplasm and nucleus. (**a**). The nuclear
localization of ATG6-mCherry in *N. benthamiana*. Scale bar, 100 μm. (**b**). Co-
localization of ATG6-GFP and nls-mCherry in *N. benthamiana*. Scale bar, 100
μm. (**c**). Subcellular fractionation of ATG6-mCherry in *N. benthamiana* after 1
mM SA treatment. Black asterisk (*) indicate ATG6-mCherry bands. (**d**).
Subcellular fractionation of ATG6-GFP in *N. benthamiana* after 1 mM SA
treatment. Black asterisk (*) indicate ATG6-GFP bands. (**e**). Subcellular
fractionation of ATG6-GFP in *ATG6-GFP Arabidopsis* after 0.5 mM SA
treatment. (**f**). Subcellular fractionation of ATG6-mCherry in *ATG6-mCherry*
Arabidopsis after 0.5 mM SA treatment. In (**c**-**f**), ATG6-mCherry (**c** and **f**) and
ATG6-GFP (**d** and **e**) were detected using ATG6 or GFP antibody. Actin and
H3 were used as cytoplasmic and nucleus internal reference, respectively.
Numerical values indicate quantitative analysis of ATG6-mCherry and ATG6-
GFP using image J. All experiments were performed with three biological
replicates.

Figure 3. ATG6 increases the nuclear accumulation of NPR1 under SA

treatment. (**a**). Confocal images of NPR1-GFP nuclear localization in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* under normal and 0.5 mM SA spray for 3 h. Scale bar, 50 μm. (**b**). Nuclear localization numbers of NPR1-GFP (per section) in (**a**), n > 10. (**c**). Subcellular fractionation of NPR1-GFP in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after 0.5 mM SA treatment for 0, 3 and 6 h. (**d**). The ration of NPR1 in the nucleus/total NPR1 in (**c**). (**e**). Subcellular fractionation of NPR1-GFP in *N. benthamiana* after 1 mM SA treatment for 0, 8 and 20 h. (**f**). The ration of NPR1 in the nucleus/total NPR1 in (**e**). In (**c** and **e**), cytoplasmic and nuclear proteins were extracted from *Arabidopsis* or *N. benthamiana*. NPR1-GFP were detected using GFP antibody. Actin and H3 were used as cytoplasmic and nucleus internal reference, respectively. Numerical values indicate quantitative analysis of NPR1-GFP using image J. All experiments were performed with three biological replicates.

Figure 4. *ATG6 increases endogenous SA levels and promotes the expression of NPR1 downstream target genes*

(**a**). Level of free SA in 3-week-old *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after *Pst* DC3000/*avrRps4* for 12 h. (**b-c**). Expression of *PR1* (**b**), *PR5* (**c**) under normal and SA treatment conditions. * or ** indicates that the significant difference compared to the control is at the level of 0.05 or 0.01 (Student *t* test *p* value, * *p*< 0.05 or ** *p*< 0.01). All experiments were performed with three biological replicates.

Figure 5. ATG6 increases the NPR1 protein levels and the formation of SINCs-like condensates. (**a**). NPR1-GFP protein levels in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after 0.5 mM SA treatment for 0, 3, 6 and 9 h. Numerical values indicate quantitative analysis of NPR1-GFP protein using image J. (**b**). NPR1-GFP protein levels in *N. benthamiana*. ATG6-mCherry + NPR1-GFP, NPR1-GFP + mCherry were co-expressed in *N. benthamiana*. After 2 days, leaves were treated with 1 mM SA for 8, 20, 24 h. Total proteins were extracted and analyzed. Numerical values

indicate quantitative analysis of NPR1-GFP protein using image J. (**c**). ATG6

promotes the formation of SINCs-like condensates. ATG6-mCherry + NPR1-

GFP, NPR1-GFP + mCherry were co-expressed in *N. benthamiana*. After 2

days, leaves were treated with 1 mM SA for 24 h. Confocal images obtained

at excitation with wavelengths of 488 nm, scale bar = 50 µm. (**d**). SINCs-like

condensates numbers of per section in (**c**), n > 10 sections. ** indicates that

the significant difference compared to the control is at the level of 0.01

(Student *t* test *p* value, ** *p*< 0.01). All experiments were performed with three

biological replicates.

Figure 6. ATG6 improves the protein stability of NPR1.

(**a**). NPR1-GFP degradation assay in *Arabidopsis*. Total proteins from 7-day-

old seedlings of *NPR1-GFP* and *ATG6-mCherry* × *NPR1-GFP* were extracted,

using Actin as an internal reference. "M" indicates MG115 treatment. (**b**).

Quantification of NPR1-GFP degradation rates in (**a**) using Image J. In (**a**

and **b**), the extracts were incubated for 0～180 min at room temperature

(25℃), the degradation rate of NPR1-GFP was analyzed. (**c**). NPR1-GFP

protein turnover. 7-day-old *NPR1-GFP* and *ATG6-mCherry* × *NPR1-GFP*

seedlings were treated with 100 µM cycloheximide (CHX) for different times.

Total proteins were analyzed, Actin was used as an internal reference. (**d**).

Quantification of NPR1-GFP protein turnover rates in (**c**) using Image J. (**e**).

NPR1-GFP protein turnover. 7-day-old *NPR1-GFP* and *NPR1-GFP*/*atg5*

seedlings were treated with 100 µM cycloheximide (CHX) for different times.

Total proteins were analyzed, Actin was used as an internal reference. (**f**).

Quantification of protein levels of NPR1-GFP in (**e**) using Image J. All

experiments were performed with three biological replicates.

Figure 7. ATG6 and NPR1 jointly inhibit the growth of *Pst*

DC3000/*avrRps4* .

(**a**). Expression of *ATG6* under *Pst* DC3000/*avrRps4* infiltration in 3-week-old

Col leaves. (**b**). Expression of *ATG6* in the presence of 0.5 mM SA in 3-week-

old Col leaves. (**c**). The protein levels of ATG6 after 0.5 mM SA in 3-week-old

Col leaves. Total leaf proteins from *Arabidopsis* were analyzed, Actin was

used as an internal reference. Numerical values indicate quantitative analysis

of ATG6 protein using image J. (**d**). Growth of *Pst* DC3000/*avrRps4* in

Col/silencing ATG6 and Col/negative control (NC). (**e**). Phenotypes of 16-day-

old amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2. Bar, 1 cm. (**f**). Phenotypes of 23-

day-old amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2. Bar, 3 cm. (**g**). Expression of

ATG6 in Col, amiRNA^{ATG6} # 1 and amiRNA^{ATG6} # 2 under infiltration treatment

of 100 µM β-estradiol. (**h**). Growth of *Pst* DC3000/*avrRps4* in *Arabidopsis*

leaves of amiRNA^{ATG6} # 1, amiRNA^{ATG6} # 2 and Col. (**i**). Growth of *Pst*

DC3000/*avrRps4* in *NPR1-GFP*/silencing *ATG6* and *NPR1-GFP*/NC. (**j**).

Growth of *Pst* DC3000/*avrRps4* in *Arabidopsis* leaves of Col, amiRNA^{ATG6} # 1,

amiRNA^{ATG6} # 2, *npr1*, *NPR1-GFP*, *ATG6-mCherry* and *ATG6-mCherry* ×

NPR1-GFP. In (**d**, **h-j**), a low dose of *Pst* DC3000/*avrRps4* (OD600 = 0.001)

was infiltrated. After 3 days, the growth of *Pst* DC3000/*avrRps4* was counted.

* or ** indicates that the significant difference compared to the control is at the

level of 0.05 or 0.01 (Student *t* test *p* value, * $p< 0.05$ or ** $p< 0.01$). All

experiments were performed with three biological replicates.

Figure 8. Working model for NPR1 regulation by ATG6. ATG6 interacts

directly with NPR1 to increase NPR1 protein level and stability, thereby

promoting the formation of SINCs-like condensates and increasing the

nuclear accumulation of NPR1. ATG6 synergistically activates *PRs* expression

with NPR1 to jointly enhance resistance to inhibit *Pst* DC3000/*avrRps4*

invasion in *Arabidopsis*.

List Supplemental Information

Supplemental Data 1

Supplemental Figure

Fig. S1 Physical interaction between NPRs and ATGs in yeast.

Fig. S2 Bimolecular fluorescence complementation assay of NPR1 with ATG6

in *N. benthamiana*.

Fig. S3 The nuclear localization of ATG6 in *Arabidopsis*.

Fig. S4 Identification of *ATG6-mCherry* × *NPR1-GFP* plants.

Fig. S5 Subcellular fractionation of endogenous ATG6 in Col after 0.5 mM SA

treatment for 0, 3, 6 and 20 h.

Fig. S6 Overexpression of *ATG6* delayed dark-induced leaf senescence.

Fig. S7 Expression of *ICS1* under normal and SA treatment conditions.

Fig. S8 Expression of *PR1* and *PR5* in Col and *ATG6-mCherry* under normal

and *Pst* DC3000/*avrRps4* treatment.

Fig. S9 The protein level of NPR1-GFP in *NPR1-GFP*/silencing *ATG6* and

NPR1-GFP/Negative control.

Fig. S10 Expression of *NPR1* in Col and *ATG6-mCherry* under normal and *Pst*

DC3000/*avrRps4* treatment.

Fig S11. ATG6 improves the protein stability of NPR1 in *N. benthamiana.*

Fig. S12 Structural analysis of acidic activation domains in ATG6.

Fig. S13 ATG6 and NPR1 jointly inhibit *Pst* DC3000/*avrRps4*-induced cell

dead.

Fig. S14 NPR1-GFP degradation assay in *ATG6-mCherry* x *NPR1-GFP*

Arabidopsis.

Fig. S15. Verification of ATG6 antibody specificity.

Supplemental Table

Table S1. Plasmid in this study.

Table S2. Primers for vector construction.

Table S3. Plant Materials.

Table S4. Antibody Information.

Table S5. Primers of RT-qPCR.

Supplemental Results

Results S1. NPR1 and its paralogues NPR3/NPR4 physically interact with

multiple ATGs.

Results S2. Overexpression of *ATG6* delays carbon starvation-induced leaf senescence, and *ATG6-GFP* and *ATG6-mCherry* fusion proteins are functional.

Supplement Methods

Methods S1. Plasmid construction

Methods S2. *Arabidopsis* screening.

Methods S3. For *Pst* DC3000/*avrRps4* culture and infiltration.

Methods S4. Yeast two-hybrid assay.

Methods S5. Prokaryotic protein expression.

Methods S6. Nuclear and cytoplasmic separation of NPR1-GFP.

Methods S7. Protein degradation analysis.

Methods S8. Protein extraction and western blotting.

Methods S9. For the treatment of 3-week-old *N. benthamiana*.

Methods S10. For the bimolecular fluorescence complementation assay.

Methods S11. For SINCs-like condensates observation.

Methods S12. For the growth of *Pst* DC3000/*avrRps4.*

Methods S13. Real-Time Quantitative PCR (RT-qPCR).

Accession numbers

Supplemental Moves

Supplemental Move 1. Localization of NPR1-GFP in *N. benthamiana* co-expressed NPR1-GFP and mCherry.

Supplemental Move 2. Localization of NPR1-GFP in *N. benthamiana* co-expressed NPR1-GFP and ATG6-mCherry.

Supplemental Data 2

Vector mapping used in this study.

Figure 1. Physical interaction between NPR1 and ATG6. (a). Interaction of NPR1 with ATG6 in yeast. The CDS of *ATG6*, *NPR1*, *NPR1-N* (0-328AA) and *NPR1-C* (328-594AA), *SnRK2.8* were fused to pGADT7 (AD) and pGBKT7 (BD), respectively. Co-transformation of NPR1-BD + AD, BD + ATG6-AD, BD + SnRK2.8-AD, NPR1-N-BD + AD, NPR1-C-BD + AD were used as negative controls. The interaction of NPR1-BD and SnRK2.8-AD was used as a positive control. Yeast growth on SD/-Trp-Leu-His-Ade media represents interaction. Numbers represent the dilution fold of yeast. 0, -1 (10 fold dilution), -2 (100 fold dilution) and -3 (1000 fold dilution). **(b).** In vitro pull-down assays of NPR1-His with GST-ATG6 fusion protein. NPR1-His prokaryotic proteins were incubated with GST-tag Purification Resin conjugated with GST-ATG6, GST and GST-SnRK2.8. Western blotting analysis with anti-GST and anti-His. Black asterisk (*) indicate GST-SnRK2.8 bands. Red asterisk (*) indicate GST-ATG6 bands. **(c).** Co-immunoprecipitation of NPR1 with ATG6 in *vivo*. Total protein was extracted from *N. benthamiana* transiently transformed with ATG6-mCherry + GFP and ATG6-mCherry + NPR1-GFP, followed by IP with GFP-Trap. Western blots analysis with ATG6 and GFP antibodies. **(d).** Co-localization of NPR1-GFP and ATG6-mCherry in *N. benthamiana* under normal and SA treatment conditions. NPR1-GFP were co-expressed with ATG6-mCherry in *N. benthamiana* for 2 d followed by confocal observation. Scale bar, 100 μm. All experiments were performed with three biological replicates.

Figure 2. ATG6 is localized in the cytoplasm and nucleus. (a). The nuclear localization of ATG6-mCherry in *N. benthamiana*. Scale bar, 100 μm. **(b).** Co-localization of ATG6-GFP and nls-mCherry in *N. benthamiana*. Scale bar, 100 μm. **(c).** Subcellular fractionation of ATG6-mCherry in *N. benthamiana* after 1 mM SA treatment. Black asterisk (*) indicate ATG6-mCherry bands. **(d).** Subcellular fractionation of ATG6-GFP in *N. benthamiana* after 1 mM SA treatment. Black asterisk (*) indicate ATG6-GFP bands. **(e).** Subcellular fractionation of ATG6-GFP in *ATG6-GFP Arabidopsis* after 0.5 mM SA treatment. **(f).** Subcellular fractionation of ATG6-mCherry in *ATG6-mCherry Arabidopsis* after 0.5 mM SA treatment. In **(c-f)**, ATG6-mCherry (c and f) and ATG6-GFP (d and e) were detected using ATG6 or GFP antibody. Actin and H3 were used as cytoplasmic and nucleus internal reference, respectively. Numerical values indicate quantitative analysis of ATG6-mCherry and ATG6-GFP using image J. All experiments were performed with three biological replicates.

Figure 3. ATG6 increases the nuclear accumulation of NPR1 under SA treatment. (a). Confocal images of NPR1-GFP nuclear localization in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* under normal and 0.5 mM SA spray for 3 h. Scale bar, 50 μm. **(b).** Nuclear localization numbers of NPR1-GFP (per section) in **(a)**, n > 10. **(c).** Subcellular fractionation of NPR1-GFP in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after 0.5 mM SA treatment for 0, 3 and 6 h. **(d).** The ration of NPR1 in the nucleus/total NPR1 in **(c). (e).** Subcellular fractionation of NPR1-GFP in *N. benthamiana* after 1 mM SA treatment for 0, 8 and 20 h. Black asterisk (*) indicate ATG6-mCherry bands **(f)** The ration of NPR1 in the nucleus/total NPR1 in **(e).** In (**c** and **e**), cytoplasmic and nuclear proteins were extracted from *Arabidopsis* or *N. benthamiana*. NPR1-GFP were detected using GFP antibody. Actin and H3 were used as cytoplasmic and nucleus internal reference, respectively. Numerical values indicate quantitative analysis of NPR1-GFP using image J. All experiments were performed with three biological replicates.

Figure 4. ATG6 increases endogenous SA levels and promotes the expression of NPR1 downstream target genes. (a). Level of free SA in 3-week-old *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after *Pst* DC3000/*avrRps4* for 12 h.**(b-c).** Expression of *PR1* **(b)**, *PR5* **(c)** under normal and 0.5 mM SA treatment conditions. * or ** indicates that the significant difference compared to the control is at the level of 0.05 or 0.01 (Student t test p value, * p< 0.05 or ** p< 0.01). All experiments were performed with three biological replicates.

Figure 5. ATG6 increases the NPR1 protein levels and the formation of SINCs. (a). NPR1-GFP protein levels in 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* after 0.5 mM SA treatment for 0, 3, 6 and 9 h. Numerical values indicate quantitative analysis of NPR1-GFP protein using image J. **(b).** NPR1-GFP protein levels in *N. benthamiana.* ATG6-mCherry + NPR1-GFP, NPR1-GFP + mCherry were co-expressed in *N. benthamiana.* After 2 days, leaves were treated with 1 mM SA for 8, 20, 24 h. Total proteins were extracted and analyzed. Numerical values indicate quantitative analysis of NPR1-GFP protein using image J. **(c).** ATG6 promotes the formation of SINCs. ATG6-mCherry + NPR1-GFP, NPR1-GFP + mCherry were co-expressed in *N. benthamiana.* After 2 days, leaves were treated with 1 mM SA for 24 h. Confocal images obtained at excitation with wavelengths of 488 nm, scale bar = 50 μm. **(d).** SINCs numbers of per section in **(c)**, n > 10 sections. ** indicates that the significant difference compared to the control is at the level of 0.01 (Student t test p value, ** p< 0.01). All experiments were performed with three biological replicates.

Figure 6. ATG6 improves the protein stability of NPR1. (a). NPR1-GFP degradation assay in *Arabidopsis*. Total proteins from 7-day-old seedlings of *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* were extracted, using Actin as an internal reference. "M" indicates MG115 treatment. **(b).** Quantification of NPR1-GFP degradation rates in **(a)** using Image J. In **(a and b)**, the extracts were incubated for 0 ~ 180 min at room temperature (25℃), the degradation rate of NPR1-GFP was analyzed. **(c).** NPR1-GFP protein turnover. 7-day-old *NPR1-GFP* and *ATG6-mCherry × NPR1-GFP* seedlings were treated with 100 μM cycloheximide (CHX) for different times. Total proteins were analyzed, Actin was used as an internal reference. **(d).** Quantification of NPR1-GFP protein turnover rates in **(c)** using Image J. **(e).** NPR1-GFP protein turnover.7-day-old *NPR1-GFP* and *NPR1-GFP/atg5* seedlings were treated with 100 μM cycloheximide (CHX) for different times. Total proteins were analyzed, Actin was used as an internal reference. **(f).** Quantification of protein levels of NPR1-GFP in **(e)** using Image J. All experiments were performed with three biological replicates.

Figure 7. ATG6 and NPR1 jointly inhibit the growth of *Pst* DC3000/*avrRps4*. (a). Expression of *ATG6* under *Pst* DC3000/*avrRps4* injection in 3-week-old Col leaves. **(b)**. Expression of *ATG6* in the presence of 0.5 mM SA in 3-week-old Col leaves. **(c)**. The protein levels of ATG6 after 0.5 mM SA in 3-week-old Col leaves. Total leaf proteins from *Arabidopsis* were analyzed, Actin was used as an internal reference. Numerical values indicate quantitative analysis of ATG6 protein using image J. **(d)**. Growth of *Pst* DC3000/*avrRps4* in Col/silencing *ATG6* and Col/negative control (NC). **(e)**. Phenotypes of 16-day-old amiRNA*ATG6* # 1 and amiRNA*ATG6* # 2. Bar, 1 cm. **(f)**. Phenotypes of 23-day-old amiRNA*ATG6* # 1 and amiRNA*ATG6* # 2. Bar, 3 cm. **(g)**. Expression of *ATG6* in Col, amiRNA*ATG6* # 1 and amiRNA*ATG6* # 2 under injection treatment of 100 μM β-estradiol. **(h)**. Growth of *Pst* DC3000/*avrRps4* in *Arabidopsis* leaves of amiRNA*ATG6* # 1, amiRNA*ATG6* # 2 and Col. **(i)**.Growth of *Pst* DC3000/*avrRps4* in *NPR1-GFP*/silencing *ATG6* and *NPR1-GFP*/NC. **(j)**. Growth of *Pst* DC3000/*avrRps4* in *Arabidopsis* leaves of Col, amiRNA*ATG6* # 1, amiRNA*ATG6* # 2, *npr1*, *NPR1-GFP*, *ATG6-mCherry* and *ATG6-mCherry* × *NPR1-GFP*. In (d, h-j), a low dose of *Pst* DC3000/*avrRps*4 (OD_{600} = 0.001) was infiltrated. After 3 days, the growth of *Pst* DC3000/*avrRps4* was counted. * or ** indicates that the significant difference compared to the control is at the level of 0.05 or 0.01 (Student t test p value, * p< 0.05 or ** p< 0.01). All experiments were performed with three biological replicates.

Figure 8. Working model for NPR1 regulation by ATG6. ATG6 interacts directly with NPR1 to increase NPR1 protein level and stability, thereby promoting the formation of SINCs-like condensates and increasing the nuclear accumulation of NPR1. ATG6 synergistically activates *PRs* expression with NPR1 to jointly enhance resistance to inhibit *Pst* DC3000/*avrRps4* invasion in *Arabidopsis*.