The TCRs of pathogenic Th17 cells in arthritogenic mice are shifted

toward a Treg-like repertoire

Mara A. Llamas-Covarrubias[1,2,3**], Atsushi Tanaka[4,5,6], Martin Loza[7], Diego Diez[8], Zichang

Xu[1], Ee Lyn Lim[5], Shunsuke Teraguchi[2,9], Shimon Sakaguchi[4,5], Daron M. Standley[1,2 *]

[1]Laboratory of Systems Immunology, WPI Immunology Frontier Research Center, Osaka

University, Suita, Japan; [2]Department of Genome Informatics, Research Institute for

Microbial Diseases (RIMD), Osaka University, Suita, Japan; [3]Departamento de Biología

Molecular y Genómica, Centro Universitario de Ciencias de la Salud, Universidad de

Guadalajara, Guadalajara, México; [4]Department of Experimental Pathology, Institute for

Frontier Medical Sciences, Kyoto University, Kyoto, Japan; [5]Laboratory of Experimental

Immunology, WPI Immunology Frontier Research Center, Osaka University, Suita, Japan;

[6]Department of Frontier Research in Tumor Immunology, Center of Medical Innovation

and Translational Research, Graduate School of Medicine, Osaka University, Osaka, Japan;

[7]Laboratory of functional analysis in silico, Human Genome Center, The University of

Tokyo, Tokyo, Japan; [8]Quantitative Immunology Research Unit, Immunology Frontier

Research Center, Osaka University, Suita, Japan; [9]Faculty of Data Science, Shiga

University, Hikone, Japan

Correspondence:

*Daron M. Standley. 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan. +81-6-6879-8366.

standley@biken.osaka-u.ac.jp

** Mara A. Llamas-Covarrubias. 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan. +81-6-

6879-8366. mara.llamas@biken.osaka-u.ac.jp

Conflict of interest:

Summary

Mutations in the ZAP-70 gene that cause moderate attenuation of T cell receptor (TCR)

signaling in mice can result in autoimmune manifestations. One explanation for this

pathology is a shift in the regulatory-conventional (Treg-Tconv) T cell repertoire

composition. To test this hypothesis, we characterized the single-cell gene expression

profiles and TCR repertoires of Tconv and Treg CD4+ T cells of arthritic (ZAC), poised

(SKG) ZAP-70 mutant, and wild-type (WT) mice. We identified a group of Th17 cells

which exhibited a pathogenic signature and occurred exclusively in inflamed joints of ZAC

mice. Such pathogenic signature was uniquely detected in CD4+ T cells obtained from

inflamed joints of RA patients. Overall, the Tconv repertoires of ZAP-70 mutant mice were

increasingly similar to the repertoires of WT Tregs, and this effect was most notable in the

subset of pathogenic Th17 cells. Our results support a model where, upon moderate ZAP-

70-mediated signal weakening, T cells that would normally develop into Tregs, instead

develop into self-reactive Tconvs, resulting in a breakdown in self-tolerance and

susceptibility to autoimmune arthritis.

Main text

49
Introduction

Autoimmune diseases encompass a diverse group of disorders characterized by aberrant T

and B cell responses against the body's own tissues. These conditions are a major source of

morbidity and disability worldwide, disproportionately affecting young women[1]. Despite

extensive research, the molecular mechanisms driving the development and activation of

self-reactive immune cells remain poorly understood. Previous studies have implicated a

range of factors in autoimmune disease pathogenesis, including viral infections[2], X

chromosome-linked ribonucleoproteins [3], and the HLA presentation of misfolded proteins

[4], among others. Independent of these models, our group recently demonstrated that a

moderate, systemic reduction in the T cell receptor (TCR) signaling protein ZAP-70 results

in autoimmune arthritis and colitis in mice [5], while more severe impairment has been

shown to lead to immunodeficiency [6].

The link between weakened TCR signaling and autoimmune disease appears to hinge on

TCR repertoire composition rather than signaling intensity in the periphery. During T cell

maturation, the repertoire is shaped by thymic selection, a process that ensures T cells are

both functional (MHC-restricted, through positive selection) and self-tolerant (through

negative selection). High-affinity interactions between TCRs and self-peptide-MHC

complexes typically lead to either clonal deletion or selection into the Treg population [7].

However, a systemic weakening of TCR signaling could disrupt this thymic selection,

allowing T cells with moderately to strongly self-reactive TCRs to evade negative selection

and enter the mature conventional T cell (Tconv) repertoire. This concept, formally known

as the "Repertoire Shift" hypothesis, is illustrated in Fig. 1a.

While the "Repertoire Shift" hypothesis aligns with previous observations in ZAP-70

mutant mice (5, 8, 9), it has not been rigorously tested. Advances in repertoire sequencing

and single-cell resolution technologies have now enabled the first direct test of this

hypothesis. To investigate it, we analyzed the transcriptome and immune repertoires of

CD4+ T cells from wild-type (WT), and two ZAP-70 mutant mouse models: non-arthritic

SKG, and arthritic ZAC, at the single-cell level. Our findings revealed distinct migratory

and pathogenic populations of Th17 cells in arthritic mice. Notably, these pathogenic cells

exhibited high expression of osteoclastogenic genes, were localized specifically to inflamed

joints, and were absent in both WT and SKG mice. Strikingly, a population with a similar

pathogenic signature was uniquely identified in T cells from human RA joints. Moreover,

the TCR repertoire of these pathogenic Th17 cells was unique and convergent. Overall, we

found an increased similarity of ZAP-70 Tconv repertoires to the repertoires of large public

WT Treg datasets, as demonstrated by two independent metrics. In addition, the Treg-like

nature of ZAP-70 mutant Tconv TCRs was most notable in the subset of pathogenic Th17

cells. Additionally, the overall similarity between Treg and Tconv repertoires was

significantly diminished in arthritic mice, suggesting that autoimmunity may result from a

depletion of critical Treg populations and the expansion of self-reactive Tconvs. Our

findings suggest that restoring the balance between Treg and Tconv repertoires could be a

promising therapeutic strategy for treating autoimmune diseases.

Results

Arthritic ZAC mice harbor two groups of Th17 cells with distinct gene programs in their joints

The ZAP-70 mutant mice (ZAC, SKG) used in this study are characterized by the spontaneous development of autoimmune arthritis (Fig. 1b), with disease progression typically accelerated in the SKG model through the stimulation of innate immunity(5, 8, 9). Unlike other widely used mouse models of arthritis, such as collagen-induced arthritis (CIA)(10), the arthritis in ZAP-70 mutant mice is genuinely autoimmune, mediated primarily by T-cells, and closely resembles human RA(5, 8, 9). Therefore, ZAP-70 mutant mouse models are particularly well-suited for investigating self-reactive TCR repertoires. To study the transcriptomes and immune repertoires of self-reactive Tconv ($CD3^+CD4^+Foxp3^-$) cells in-depth, we sorted cells from the joints, draining lymph nodes (dLN), and spleens of arthritic ZAC mice and performed RNA+V(D)J single-cell sequencing. Gene expression analysis revealed a considerable overlap in the transcriptional profiles of spleens and dLNs but a distinct profile for joint cells (Supplementary Fig. 1a). Unsupervised clustering delineated nine gene expression clusters, including two (clusters 1 and 4) expressing canonical Th17 markers (*Rorc, Il17a, Il17f*) but with different gene expression programs and tissue abundances (Supplementary Note 1, Fig. 1c-e, Supplementary Fig. 1b-d). Cluster 1 was abundant in joints, but was also detected in the

spleen and dLNs. Relative to cluster 4, cells in cluster 1 differentially expressed *Klf2*, *S1pr1*

and chemokine receptors, and it was enriched in the regulation of T cell migration (GO:

20000404) and cell chemotaxis (GO: 0060326) gene ontology (GO) Terms (Fig. 1f-g,

Supplementary Table 4-5) suggesting a phenotype of migratory cells. In contrast, cluster 4

was almost exclusively observed in joints, and highly expressed *Tnfsf11* (encoding for

RANKL) along with other osteoclastogenic genes (*Csf1*, *Anxa2*), and overall, the osteoclast

differentiation (GO:0030316) GO term was significantly enriched (Fig. 1d, f-g,

Supplementary Table 5). Moreover, this cluster uniquely expressed transcription factors

associated with a pathogenic Th17 phenotype: *Rbpj*(11)*, Crem*(12)*, Atf3*(13)*, Bhlhe40*(14)*,*

and *Igfbp7*(15). These results indicate that cluster 4 contains pathogenic Th17 cells with an

increased osteoclastogenic potential. Thus, we annotated clusters 1 and 4 as migratory and

pathogenic Th17, respectively. Interestingly, cluster 1 and cluster 4 could be clearly

distinguished by their differential expression of *Il22* or *Tnfsf11* (Fig. 1h). Trajectory

inference of the different Tconv cell states resulted in the identification of the pathogenic

Th17 cluster as one of the endpoint lineage states (Supplementary Fig. 1e), consistent with

their effector phenotype. Of note, we did not detect clear and specific expression of marker

genes of previously reported populations of pathogenic Th17 cells (Th17.1(16),

exFoxP3(17)) in cluster 4, suggesting that it represents a distinct group of pathogenic cells

(Supplementary Fig. 1f-g).

To contextualize our findings in arthritic ZAC mice, we replicated the analyses in WT and

non-arthritic SKG mice. Here, we sampled Tconv from their spleens and integrated the

ZAC spleen data in order to facilitate direct comparison of clusters (Supplementary Note 2,

Supplementary Fig. 2a-g). This approach revealed that the general Th17 cell abundance

increased progressively from WT to SKG to ZAC mice (Supplementary Fig. 2d). To

analyze whether the transcriptional programs of these Th17 cells were similar to those in

the migratory and pathogenic Th17 cells, we calculated gene module scores for their

signature genes (Supplementary Table 6). Expression of the migratory Th17 signature was

clearly detected and gradually increased from WT, to SKG to ZAC splenic Th17 cells;

however, the pathogenic Th17 signature was largely undetected (Supplementary Fig. 3a-b).

Thus, we demonstrated the presence of two distinct groups of Th17 cells in arthritic ZAC

mice: migratory Th17 cells, expressing *Il22*, and pathogenic Th17 cells, which highly

express *Tnfsf11* and uniquely localize within inflamed joints.

The transcriptional signature of pathogenic Th17 cells is present in inflamed joints of

human RA but not in other autoimmune diseases or non-autoimmune arthritis

To investigate the presence of cells resembling pathogenic Th17 cells in human tissues, we

calculated migratory and pathogenic Th17 gene signature module scores in publicly

available single-cell RNAseq datasets. We analyzed data obtained from three sources:

inflamed synovia(18) and PBMCs(19) of RA patients; synovia and infrapatellar fat pad

(IPFP) of osteoarthritis (OA) patients(20); peripheral CD4+ T cells from healthy controls

and patients with other autoimmune diseases(21). Detailed descriptions of the datasets are

provided in Supplementary Table 7.

Our analysis revealed that the migratory Th17 signature was clearly detected in all datasets

(Fig. 1i, Supplementary Fig. 3c-e). In contrast, expression of the pathogenic Th17 signature

was nearly undetectable in all samples (Supplementary Fig. 3c-e) except in CD4$^+$ T cells

derived from inflamed joints of RA patients (Fig. 1i); in addition, its expression was non-

overlapping with the expression of the migratory Th17 signature, mirroring our

observations in ZAC joints. Notably, as in ZAC joints, the expression of Th17.1 or

exFoxP3 markers was not specifically detected in the cluster of cells expressing the

pathogenic Th17 gene signature in human RA joints (Supplementary Fig. 4a-d). Since OA

is a form of non-autoimmune arthritis(22), our results indicate that cells with a pathogenic

Th17 transcriptional signature are likely to be specifically involved in the pathogenesis of

human autoimmune arthritis.

Pathogenic Th17 cells are expanded, convergent, and contain public clones

We next asked whether the TCR repertoires of the two Th17 clusters were also distinct.

First, we examined clonal expansion and found that large and medium clones were

abundant in both Th17 cell groups (Fig. 2a, Supplementary Fig. 5a-b). Next, we evaluated

the repertoire similarity of both Th17 groups to other Tconv populations and between each

other by using the Morisita-Horn index. At the exact paired CDR3 amino acid sequence

level, the repertoire of migratory Th17 cells was largely similar to that of other Tconv

repertoires from ZAC tissues (Fig. 2b). In contrast, the pathogenic Th17 repertoire was less

similar to other Tconv cells, and the similarity between the two Th17 cell repertoires was

rather low (Fig. 2b). These results suggest that pathogenic Th17 cells possess a unique

repertoire, presumably due to the targeting of a specific set of antigens.

We next evaluated intra-repertoire TCR similarity as a measure of repertoire convergence.

For this purpose, we used TCRdist(23), which computes the distance between the CDR1,

CDR2 and CDR3 amino acid sequences of single or paired TCR chains. Using TCRdist, we

independently calculated distances among all receptors in the two Th17 clusters, as well as

the naïve cluster for reference. Because a bias in the distribution of the distances to lower

values indicates higher similarity within the receptors, it can be interpreted as an indication

of repertoire convergence. We found that the TCR distance distribution of the pathogenic

Th17 repertoire trended lower in paired and single chain analysis, especially for the alpha

chain (Fig. 2c, Supplementary Fig. 5c). This result indicates that the repertoire of

pathogenic Th17 cells is more convergent than that of migratory Th17 cells. Finally, we

identified a total of 9 unique clones shared in two or more individuals (public clones) and

these included expanded clones within the Th17 compartments (Supplementary Fig. 5d,

Supplementary Table 8). Together, these results indicate that although the Th17 response in

arthritic ZAC mice is polyclonal, the repertoire of tissue-specific pathogenic Th17 cells is

unique, convergent, and public. Given their specific location in joints, and exclusive

similarity to Th17 T cells in human RA, our results suggest that such pathogenic Th17 cells

may recognize a restricted set of self-antigens in the affected tissues.

TCR repertoires of pathogenic Th17 cells are more Treg-like than Tconv-like

The repertoire shift hypothesis postulates that ZAP-70 hypomorphic mutations would lead

to the appearance of TCR repertoires in Tconvs that would typically be deleted or found in

WT Tregs(5, 9) (Fig. 1a). To test this prediction, we used a machine learning model, TiRP

(TCR-intrinsic regulatory potential), that measures the intrinsic Treg propensity of a given

TCR, based on sequence features(24). It has previously been shown that self-reactive TCRs

typically have higher TiRP scores than non self-reactive TCRs(24). Consistent with these

reports, our experimentally determined Treg datasets exhibited higher TiRP scores than WT

Tconv cells, and their respective Tconv compartments (Supplementary Fig. 6a-b). In

contrast, the ZAP-70 mutant Tconvs exhibited significantly higher TiRP scores than WT

Tconvs or even WT Tregs (Supplementary Fig. 6a-b). Furthermore, we found that

pathogenic Th17 TiRP scores were significantly higher than those of WT Tconvs (Fig. 2d-

e). These finding support a model where TCRs that would normally be deleted or develop

into Tregs instead develop into pathogenic Tconvs upon moderate TCR signal weakening.

We next focused on the Treg and Tconv TCR sequences that have been identified in

previous studies (Supplementary Table 10) (25-28) in order to confirm whether or not we

could observe similarities to the TCRs in our study. To this end, we treated each of our

experimentally-determined (WT, SKG and ZAC Tconv and Treg) TCRs as a "query" that

was used to search reference WT Tconv or WT Treg datasets (Logunova(25), Lu(26),

Ko(27), Wolf(28)). For each query, we computed the number of reference WT Tconv and

WT Treg sequence matches to CDR1, CDR2 and CDR3. The number of matches

(normalized by the total size of each reference dataset) was deemed "Treg-like", "Tconv-

like" or "not significant" using a hypergeometric p-value cutoff of 0.001 (Fig. 2f).

Considering the largest (Logunova(25)) reference dataset, we found that the number of

Treg-like TCRs within the SKG and ZAC Tconv T cells was approximately twice as high

as that within WT Tconvs, and was higher even than that of query WT Tregs (Fig. 2g). In

addition, the fraction of Tconv-like TCRs was notably reduced within ZAP-70 mutant

Tconv queries (Fig. 2h). Similar trends were observed for the other reference datasets

(Supplementary Fig. 6c). Significantly, the ZAC pathogenic Th17 repertoire exhibited the

highest fraction of Treg-like TCRs along with the smallest fraction of Tconv-like TCRs

(Fig. 2h, Supplementary Fig. 6d), indicating that the TCRs of pathogenic Th17 cells were

largely absent from WT Tconv repertoires. Of note, pathogenic Th17 cells from ZAC mice

did not show any significant WT Treg or WT Tconv matches within the Ko et al., 2020(27)

dataset (Supplementary Fig. 6d). These observations support TiRP predictions that ZAP-70

mutant Tconv repertoires are more Treg-like than Tconv-like, and that this trend is most

pronounced in the pathogenic Th17 cells.

Evidence of missing TCR specificities in ZAP-70 mutant WT Treg repertoires

A second consequence of weakening of TCR signaling is that it generates "holes" in the

normal Treg repertoire. It has been reported that missing specificities in the Treg repertoire

have a large impact on the suppressive function against self-reactive T cells(29, 30). This

raises the possibility that an absence of Treg specificities within ZAP-70 mutant mice

repertoires contribute to a loss in tolerance toward specific self-antigens. To explore this

hypothesis, we analyzed the Treg-Tconv repertoire similarity within each mouse and

compared it across groups (WT, SKG, ZAC). The Morisita-Horn index for exact CDR3

overlap showed a notably reduced Treg-Tconv similarity in ZAC mice relative to WT or

SKG, especially at the single chain level (Fig. 3a). The Treg-Tconv overlap was generally

distributed across effector phenotypes (Tconv: intermediate, Th17 and Tfh. Treg:

intermediate and effector) in non-arthritic mice (Supplementary Fig. 7a-h), suggesting a

suppressive potential for shared Tregs towards activated Tconvs; however, such enrichment

was not observed in ZAC mice. To further test these observations, we again employed

TCRdist(23) and calculated repertoire-repertoire correlations. We found that WT Treg and

WT Tconv repertoires were well correlated, whereas SKG and ZAC Tconv repertoires

exhibited lower correlation with their Treg repertoires. Indeed, the ZAC Treg repertoire

was poorly correlated with other groups in general, indicating a reduced similarity to any

other tested repertoires (Fig. 3b). Together, these results are consistent with a scenario

wherein ZAP-70 mutant were unable to select certain Treg TCR specificities normally

present in WT mice and capable of suppressing autoimmune arthritis. This, along with a

more self-reactive Th17 population, contributes to a loss of tolerance and the development

of autoimmune arthritis.

Discussion

In this study, we provide a comprehensive characterization of the Treg and Tconv

phenotypes and repertoires of WT, non-arthritic SKG and arthritic ZAC mice at the single

cell level. We identified two transcriptionally distinct populations of Th17 cells within

tissues of arthritic ZAC mice: migratory and pathogenic Th17 cells. Pathogenic Th17 cells

have an osteoclastogenic transcriptional signature, are almost exclusively detected in joints

and harbor unique and convergent TCRs with a greater similarity to the TCRs of WT Treg

cells, which are generally understood to be biased toward self-reactivity(31-34).

It is possible that the pathogenic Th17 population is only acquired within joints by cells

harboring self-reactive TCRs, and that these cells are critical for the establishment of

disease; whereas migratory Th17 cells may contribute to disease progression and

maintenance and represent a prior differentiation state for the cells that later acquire a

pathogenic Th17 phenotype. This hypothesis is supported by the observation of a degree of

clone sharing between migratory and pathogenic Th17 cells, as well as by trajectory

prediction. Our results are also consistent with previous research showing that in

experimental autoimmune encephalomyelitis (EAE), Th17 cells progress from a self-

renewing cellular, to an effector pre-Th1-like phenotype characterized by the expression of

chemokine receptors, to a Th17/Th1-like effector phenotype that exhibits a pathogenic

signature with high expression of *Tnfsf11*. Notably, in line with our findings, these cells

also display differential enrichment in tissues from lymph nodes to central nervous system

(CNS); however, their TCR repertoires were not explored(35).

In a typical immune response, T cells that infiltrate target tissues include both antigen-

specific cells and those activated in a bystander manner, recruited by inflammatory signals.

In autoimmune diseases, the proportion of tissue-infiltrating T cells that are truly self-

reactive remains uncertain. Identifying these self-reactive T cells is essential for

understanding the mechanisms driving these disorders and developing more targeted

therapeutic strategies. Our study identified a pathogenic subset of joint-infiltrating T cells

with a TCR repertoire that is distinctly skewed towards self-reactivity. The selective

enrichment of these pathogenic cells in inflamed joints, both in mouse models and human

RA synovia, underscores their self-reactive nature and highlights their potential role in

recognizing tissue-specific self-antigens.

While other pathogenic Th17 cell subtypes in autoimmune arthritis have been described in

the literature(16)'(17), the transcriptional profile of the cells identified in our study is

distinct, suggesting a unique subset. Although some degree of self-reactivity has been

reported for Th17.1 cells(36), the specific TCRs involved, and the characteristics of their

repertoire remain largely unexplored. In a separate study, a population of T peripheral

helper (Tph) cells, characterized by their expansion and expression of B cell-helper factors,

was identified in the joints of seropositive RA patients(37). These cells were later shown to

proliferate in autologous mixed lymphocyte reaction (AMLR) assays, indicating self-

reactivity(38). However, seropositive RA represents only a subset of RA patients(39), and

it is unlikely that the same antigens drive both B cell help and antibody production, as well

as local inflammation and tissue destruction. Therefore, our identification of self-reactive

cells involved in osteoclastogenesis and tissue destruction is critical for uncovering tissue

epitopes relevant across different clinical groups of RA. Furthermore, the single-cell

approach in our study enables the retrieval of full and paired-chain TCR sequences,

offering the potential for future determination of antigen specificity in the context of a

complete TCR repertoire.

In line with this, we observed a number of public clones within ZAC Tconv cells,

particularly concentrated within both Th17 phenotypes, and notably expanded. Among the

CDR3 sequences of public clones, we identified only one exact match to a TCR in the

VDJdb database(40) of TCR sequences with known antigen specificities (Supplementary

Table 9). Unexpectedly, the match corresponded to a CD8 TCR, and its target peptide was

presented in a class I MHC. Although the presence of "class-mismatched" T cells has been

reported in the literature(41-43), it remains unknown whether this phenomenon occurs in

this or other self-reactive clones in ZAC mice. Consequently, the identity of any potential

epitopes or amino acid motifs remains an open problem.

In addition to these observations on the pathogenic phenotype of ZAC Th17 cells, we

carried out the first systematic investigation at single cell resolution of the repertoire shift

of conventional T cells upon TCR signal weakening. Our results are in line with the

observation that T cells infiltrating prostatic autoimmune lesions in Aire[-/-] mice express

receptors preferentially found in Tregs from Aire[+/+] mice(44). In addition, exFoxP3 cells

display a higher osteoclastogenic potential as compared to Th17 derived from naïve CD4[+]

T cells, presumably due to their higher affinity to self-antigens(17). Together with our

findings, these reports point to a key role of TCRs normally occurring in the Treg

compartment in the recognition of self-antigens, and suggests that aberrant expression of

such TCRs in Tconv cells may drive autoimmune disease.

Finally, the repertoire shift hypothesis raises the prospect of loss of certain TCR

specificities within the Treg compartment. Considering that antigen recognition by the Treg

TCR appears to be less degenerate than in Tconv cells(29, 45, 46), the loss of certain Treg

specificities could profoundly impact tolerance towards specific self-antigens(30). While it

is generally acknowledged that Treg and Tconv repertoires are distinct, a small proportion

of their repertoires typically overlap(47-49). In this study, we observed a reduction in the

similarity between ZAC Treg and Tconv repertoires, specially within activated Tconv cell

groups, suggesting a divergence in the Treg-Tconv repertoires that could result in the loss

of critical Treg function. Therefore, it is possible to speculate that moderate TCR

weakening by ZAP-70 mutation may not only shift the Tconv repertoire but also deplete the

Treg repertoire of critical TCRs. The antigen specificity of such TCRs remains unknown,

and thus worthy of future study as a starting point for development of TCR-transgenic Treg

immunotherapy for the treatment of autoimmune diseases(50).

Taken together, the impact of weakened TCR signaling on the development of autoimmune

disease is multifaceted. Firstly, it allows T cells with self-reactive TCRs to escape negative

selection or diversion into the Treg lineage, instead differentiating into self-reactive Tconvs.

These cells can subsequently acquire a pathogenic phenotype and accumulate in affected

tissues. Simultaneously, the Treg repertoire may lose some critical TCRs, leading to both a

shift and bifurcation in the repertoire. In addition, there is evidence that Treg effector

activity upon TCR weakening is significantly disrupted(5), rendering Tregs insufficient for

the suppression of self-reactive Tconvs (Fig. 4).

It should be noted that our study has several limitations, such as the limited sample size and

repertoire coverage that can be practically analyzed by current single cell sequencing

approaches. In addition, the pathogenic potential of Th17 cells, and their self-reactivity

have not been further experimentally assessed. Moreover, the antigen specificity of public

clones could only be predicted, which yielded only a single match to a functionally

annotated TCR. Future *in vitro* and *in vivo* experiments will be required to elucidate the

specific role and antigen specificities of Th17 cell populations in disease establishment and

progression. In spite of these limitations, our research expands the cellular and molecular

level understanding of the breakdown in tolerance during autoimmune disease and proposes

a straightforward path to discovering TCRs involved in RA.

Methods

Sex as a biological variable

The experimental design and analyses in this study were conducted without distinguishing

between sexes, and sex was not considered as a biological variable.

Mice

BALB/c and SKG mice were acquired from CLEA Japan. SKG(8), ZAC(5) and FIG

(Foxp3-IRES-GFP knock-in(51)) mice were previously described. In this study, FIG

BALB/c WT, FIG SKG and FIG ZAC mice were used and generated as previously reported

(5). All animals were maintained in Specific pathogen-free (SPF) facilities.

Droplet-based single cell RNA sequencing.

CD4$^+$Foxp3$^-$ or Foxp3$^+$ cells from FIG BALB/c WT, FIG SKG or FIG ZAC mice were

sorted by FACSAria SORP (BD Biosciences). For FIG BALB/c WT and FIG SKG mice,

cells were obtained from spleens, processed with the Chromium Single Cell 5' Library &

Gel bead kit (10x Genomics), loaded onto a Chromium Single Cell G Chip, and

encapsulated in a Chromium single cell controller (10x Genomics) to generate single-cell

gel beads in the emulsion (GEMs) following manufacturer's protocol. After encapsulation,

reverse transcription and cDNA amplification were performed, followed by target

enrichment of V(D)J segments and the independent construction of V(D)J and Gene

expression libraries as recommended by the manufacturer's instructions.

For FIG ZAC mice, cells from inflamed joint, draining popliteal lymph node (dLN) and

spleen from each replicate were barcoded with TotalSeq hashtag antibodies (BioLegend)

and then pooled by tissue and cell type (i.e., Joint Tconv, Joint Treg, etc.) before the

generation of GEMs. Next, RNA and V(D)J libraries were produced as described above

and an additional library for the barcoded antibodies was prepared. HiSeq3000 and

NovaSeq6000 were used for sequencing.

Single-cell RNAseq data processing

Raw reads were aligned to the mouse reference genome (GRCm38 and GRCm38 VDJ

v5.0.0, from 10× Genomics), filtered, and demultiplexed to create gene expression matrices

by Cell Ranger (v.5.0). The resulting data was analyzed in R software (v 4.1.2) with Seurat

v4.0 package(52, 53).

Where necessary, hashtag demultiplexing was performed by the *HTODemux* function from

Seurat and only data from cells classified as singlets was kept for further steps. Next, low-

quality data was removed from all datasets, this means cells with unique feature counts less

than 200 or over 2500, and cells in which mitochondrial reads represented >5% of reads.

Only cells annotated to a single alpha and a single beta chain in the V(D)J result were

retained for further analysis. Normalization was performed by the "LogNormalize" method

from Seurat and the top 2,000 variable genes were identified in each dataset independently.

Batch correction and data integration were performed in combinations of data as follows: 1)

Tconvs from ZAC tissues; 2) Tregs from ZAC tissues; 3) Tconvs from spleen (including

WT, SKG and ZAC samples); 4) Tregs from spleen (including WT, SKG and ZAC

samples). Batch correction was performed with the Canek R package(54) using the

intersecting variable genes for the datasets to be integrated in each run.

After integration and QC, the final datasets for analysis consisted of: 6,509 cells for ZAC

Tconv, 5,306 cells for ZAC Treg, 24,743 cells for Spleen Tconv and 25,227 cells for

Spleen Treg (Supplementary Table 1). Uniform manifold approximation and projection

(UMAP) was used to visualize gene expression profiles and clusters from the principal

components (PCs) representation of the integrated datasets; the number of PCs were

independently selected on each dataset as the elbow suggested in variance plot of PCs.

Public single-cell RNAseq data was obtained as count matrices and pre-processed

accordingly for QC, normalization, batch correction and data integration (when required).

Single cell gene expression analysis

Clustering and cell-type annotation

The identification of cell clusters based on gene expression data was performed in the PCs

representation of integrated datasets with the *FindClusters* function from Seurat using a

resolution parameter of 0.5. Clusters were annotated by the examination of expression of

canonical Treg or Tconv markers together with the inspection of markers defining each

cluster and enriched GO Biological Process Terms (Supplementary Notes 1-4). Additional

exclusion of artifactual populations was performed after clustering, including cells lacking

expression of CD3 (*Cd3d*, *Cd3e*, and *Cd3g)*, invariant NKT (iNKT) cells, and clusters with

high expression of Hemoglobin genes (*Hba-a1, Hba-a2, Hbb-bt, Hbb-bs*).

DEG and gene set enrichment analysis

Identification of cluster marker genes was performed with *FindAllMarkers* function in

Seurat. Significant markers were considered with a fold change (FC) ≥1.2 and adjusted p

value ≤0.05, and were used for gene set enrichment analysis.

When the comparison of two populations was required, DEGs were identified with the

FindMarkers function with the min.pct parameter set at 0.25. Genes with adjusted p value

≤0.05 where considered significantly changed and were classified as upregulated if FC ≥1.5

or downregulated if FC ≤0.67.

Gene set enrichment analysis was performed with enrichR (v3.1) package(55) using the

GO_Biological_Process_2021 database as reference.

For the generation of ZAC migratory and pathogenic Th17 gene signatures for module

score calculation, significant markers were considered if fold change (FC) ≥2 and adjusted

p value ≤0.05.

Trajectory analysis

Trajectory analyses were performed by using the Slingshot Bioconductor package

(v2.10.0)(56) with the PCs representation of integrated datasets and their respective cluster

labels as inputs. We set the parameter *start.clus='naive'* to use the cluster of Naive cells as

the starting point of the inferred trajectories.

Module score calculation

Calculation of expression levels from Th17 ZAC programs in spleen and public human

scRNAseq datasets was performed by the *AddModuleScore* function from Seurat using the

specified gene lists or their corresponding human orthologs (Supplementary Table 6) and

default parameters.

For public scRNAseq datasets, clustering was performed as described above. Where it was

possible to annotate Th17 cells, these were separated in an independent Seurat object and

processed for module score calculation and visualization. If Th17 cells were not clearly

identified, the whole CD4$^+$ T cell clusters were separated in independent Seurat objects for

module score calculation.

Single TCR V(D)J sequencing analysis

Clonal expansion and repertoire overlap

Clonal expansion was assessed by the calculation of clonal proportions, then, clones were

classified by their size as: small (clonal proportion $<1\text{x}10^{-3}$), medium (clonal proportion

from $1\text{x}10^{-3} – 1\text{x}10^{-2}$) and large (clonal proportion $>1\text{x}10^{-2}$). Given that only a small sample

size could be recovered for some replicates, which inflates clonal proportions of lowly

abundant clones, we labeled singletons (clone size=1) separately.

The Morisita-Horn index is a typical metric used in ecology to calculate the similarity of

two ecosystems in terms of species distribution and abundance, and it has been introduced

as a metric for repertoire overlap analysis; its values go from 0 to1, where 1 represents a

perfect overlap(57). The Morisita-Horn index was calculated between repertoires of each

individual replicate, at the paired or single CDR3 amino acid level by the Immunarch

(v0.9.0) R package. The median Morisita-Horn indices were compared by a Wilcoxon test

using a significance level of 0.05. In addition, clonotype tracking across Treg and Tconv

phenotypes was performed by Immunarch.

Clonal publicity

Public clones were defined as TCRs sharing paired gene usage and CDR3 amino acid

sequences and occurring in more than one subject.

TCR distances and repertoire-repertoire correlation

In order to evaluate if the overall features of selected repertoires show convergence, we

calculated receptor distances with TCRdist. TCRdist allows for the calculation of a distance

measure between TCRs by comparing the CDR1, CDR2 and CDR3 amino acid sequences

of single or paired TCR chains(23), then, a neighbor distance metric consisting of the

weighted average of the distance of a given TCR to the nearest 25[th] percentile of the other

TCRs in the repertoire can be calculated (weighted neighbor distance). Then, distance

distributions of selected repertoires were compared and distributions biased toward lower

distances were interpreted as an indication of repertoire convergence. In order to restrict the

convergence analysis to the qualitative composition of the repertoires, we used unique

clone sequences i.e., clonal counts where not considered. Median distances of Naïve,

migratory and pathogenic Th17 cells were compared by a Wilcoxon test with a significance

level of 0.05. Finally, the epitope-epitope (or repertoire-repertoire) neighbor distance score

correlation metric from the TCRdist algorithm was computed for assessment of intra Treg-

Tconv repertoire similarity in pooled WT, SKG and ZAC reperotires. Briefly, a score is

assigned to each TCR in one repertoire which is the average distance to the nearest TCRs in

a reference repertoire. Then, the sets of scores obtained by two different repertoires can be

compared over the entire merged set of TCRs and the Pearson correlation score is

calculated. In principle, two similar repertoires will be well correlated.

TiRP score calculation

Lagattuta et al proposed a Treg-propensity scoring system for the TCR to quantify the

likelihood of Treg cells based on the TCR feature on Treg fate (TiRP score)(24). The

source code was obtained from https://github.com/immunogenomics/TiRP. The TiRP score

of this study was calculated using the default parameters. For each TCR sequence, the v-

gene and amino acid of CDR3 were utilized as input information. Median TiRP scores were

compared across groups by a Wilcoxon test with a significance level of 0.05.

Published sequence preparation

The published mouse Treg and Tconv TCR beta chain sequence data were obtained from

four studies(25-28). The detailed accession ID and references were summarized in

Supplementary Table 14. The annotation of raw sequences was performed by mixcr

(3.0.13)(58) with align, assemble, assembleContigs, and exportClones process. Sequences

with complete frameworks and CDR regions were used in this study.

Sequence searching and comparison

We implemented the TCR sequence searching using Interclone method(59). The source

code was deposited in https://gitlab.com/sysimm/interclone and the searching parameter

using the default setting. In brief, InterClone regards the CDR1,2,3 of one TRB seq as a

pseudo sequence and considers one pair of hitting using the following CDR similarity and

coverage thresholds: CDR1 $\geq$ 90%, CDR2 $\geq$ 90%, CDR3 $\geq$ 80%, and Coverage $\geq$ 90%. We

searched Tconv and Treg TCR templates from the same reference using identical inputs.

Then we compare the corresponding hit numbers with Tconv and Treg templates for each

query and calculate the significant Tconv or Treg-biased query sequence using a

hypergeometric p-value cutoff of 0.001. We then quantified the total number of matches in

terms of the query repertoire size and normalized the results to the matches in the query

WT Tconvs (Fig. 4a).

539   Statistical analysis

For identification of cluster markers, significant markers were considered with a FC $\geq$1.2

and adjusted p value $\leq$0.05. For DEG analysis between two cell populations, genes with

adjusted p value $\leq$0.05 where considered significantly changed and were classified as

upregulated if FC $\geq$1.5 or downregulated if FC $\leq$0.67. The median Morisita-Horn indices,

median distances of Naïve, migratory and pathogenic Th17 repertoires and median TiRP

scores were compared by a Wilcoxon test using a significance level of 0.05. For TCR

sequence searching and comparison, we calculated the significant Tconv or Treg-biased

query sequence using a hypergeometric p-value cutoff of 0.001.

Study approval

Animal experiments were approved by the institutional review board of Osaka University,

and animals were treated in accordance with the institutional guidelines of the Immunology

Frontier Research Center, Osaka University.

Data availability

The datasets generated during the current study have been deposited in the DDBJ Sequence

Read Archive database (accession numbers DRA010311, DRA010223), and NCBI Gene

Expression Omnibus (accession number GSE180432).

Author contributions

D.S. and M.L.C. conceived the study. M.L.C., S.T. and A.T. designed experiments. M.L.C.,

E.L., and A.T performed experiments. M.L.C., M.L.L., D.D., Z.X., performed data analysis.

M.L., A.T. and D.S. interpreted data and wrote the paper. S.T. and Z.X. designed custom

software for TCRs analysis. D.M. and S.S. supervised the project. All the authors read and

approved the final manuscript.

Acknowledgements

This work was funded by the Japan Agency for Medical Research and Development (AMED), grant number JP223fa627002 and Platform Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research) under JP21am0101108; and by JSPS KAKENHI Grant Numbers JA23H034980 (to D.S.), 20K16286 (to M.L.C.), and 26860331, 17K15723, 22H02920 (to A.T.). M.L.C. received a postdoctoral scholarship from the Mexican National Council of Science and Technology (CONACYT) number 2018-000022-01EXTV-00314 to perform this research.

576   **References**

577   1.   Pisetsky DS. Pathogenesis of autoimmune disease. *Nat Rev Nephrol.*

578   2023;19(8):509-24.

579   2.   Robinson WH, Younis S, Love ZZ, Steinman L, and Lanz TV. Epstein-Barr virus

580   as a potentiator of autoimmune diseases. *Nat Rev Rheumatol.* 2024.

581   3.   Dou DR, Zhao Y, Belk JA, Zhao Y, Casey KM, Chen DC, et al. Xist

582   ribonucleoproteins   promote   female   sex-biased   autoimmunity.   *Cell.*

583   2024;187(3):733-49 e16.

584   4.   Mori S, Kohyama M, Yasumizu Y, Tada A, Tanzawa K, Shishido T, et al. Neoself-

585   antigens are the primary target for autoreactive T cells in human lupus. *Cell.*

586   2024;187(21):6071-87 e20.

587   5.   Tanaka A, Maeda S, Nomura T, Llamas-Covarrubias MA, Tanaka S, Jin L, et al.

588   Construction of a T cell receptor signaling range for spontaneous development of

589   autoimmune disease. *J Exp Med.* 2023;220(2).

590   6.   Siggs OM, Miosge LA, Yates AL, Kucharska EM, Sheahan D, Brdicka T, et al.

591   Opposing functions of the T cell receptor kinase ZAP-70 in immunity and tolerance

592   differentially   titrate   in   response   to   nucleotide   substitutions.   *Immunity.*

593   2007;27(6):912-26.

594   7.   Ashby KM, and Hogquist KA. A guide to thymic selection of T cells. *Nat Rev*

595   *Immunol.* 2024;24(2):103-17.

8.   Sakaguchi N, Takahashi T, Hata H, Nomura T, Tagami T, Yamazaki S, et al.
Altered thymic T-cell selection due to a mutation of the ZAP-70 gene causes
autoimmune arthritis in mice. *Nature*. 2003;426(6965):454-60.

9.   Tanaka S, Maeda S, Hashimoto M, Fujimori C, Ito Y, Teradaira S, et al. Graded
attenuation of TCR signaling elicits distinct autoimmune diseases by altering
thymic T cell selection and regulatory T cell function. *J Immunol*.
2010;185(4):2295-305.

10.   Brand DD, Latham KA, and Rosloniec EF. Collagen-induced arthritis. *Nat Protoc*.
2007;2(5):1269-75.

11.   Meyer Zu Horste G, Wu C, Wang C, Cong L, Pawlak M, Lee Y, et al. RBPJ
Controls Development of Pathogenic Th17 Cells by Regulating IL-23 Receptor
Expression. *Cell Rep*. 2016;16(2):392-404.

12.   Ohl K, Nickel H, Moncrieffe H, Klemm P, Scheufen A, Foll D, et al. The
transcription factor CREM drives an inflammatory phenotype of T cells in
oligoarticular juvenile idiopathic arthritis. *Pediatr Rheumatol Online J*.
2018;16(1):39.

13.   Glal D, Sudhakar JN, Lu HH, Liu MC, Chiang HY, Liu YC, et al. ATF3 Sustains
IL-22-Induced STAT3 Phosphorylation to Maintain Mucosal Immunity Through
Inhibiting Phosphatases. *Front Immunol*. 2018;9:2522.

14.   Lin CC, Bradstreet TR, Schwarzkopf EA, Jarjour NN, Chou C, Archambault AS, et
al. IL-1-induced Bhlhe40 identifies pathogenic T helper cells in a model of
autoimmune neuroinflammation. *J Exp Med*. 2016;213(2):251-71.

15. DiToro D, Harbour SN, Bando JK, Benavides G, Witte S, Laufer VA, et al. Insulin-
Like Growth Factors Are Key Regulators of T Helper 17 Regulatory T Cell Balance
in Autoimmunity. *Immunity.* 2020;52(4):650-67 e10.

16. Basdeo SA, Cluxton D, Sulaimani J, Moran B, Canavan M, Orr C, et al. Ex-Th17
(Nonclassical Th1) Cells Are Functionally Distinct from Classical Th1 and Th17
Cells and Are Not Constrained by Regulatory T Cells. *J Immunol.*
2017;198(6):2249-59.

17. Komatsu N, Okamoto K, Sawa S, Nakashima T, Oh-hora M, Kodama T, et al.
Pathogenic conversion of Foxp3+ T cells into TH17 cells in autoimmune arthritis.
Nat Med. 2014;20(1):62-8.

18. Etori K, Tanaka S, Tamura J, Hattori K, Kagami SI, Nakamura J, et al. Fibroblast
growth factor receptor 1 as a potential marker of terminal effector peripheral T
helper cells in rheumatoid arthritis patients. *Rheumatology (Oxford).*
2023;62(11):3763-9.

19. Zhang B, Zhang Y, Xiong L, Li Y, Zhang Y, Zhao J, et al. CD127 imprints
functional heterogeneity to diversify monocyte responses in inflammatory diseases.
J Exp Med. 2022;219(2).

20. Tang S, Yao L, Ruan J, Kang J, Cao Y, Nie X, et al. Single-cell atlas of human
infrapatellar fat pad and synovium implicates APOE signaling in osteoarthritis
pathology. *Sci Transl Med.* 2024;16(731):eadf4590.

21.   Yasumizu Y, Takeuchi D, Morimoto R, Takeshima Y, Okuno T, Kinoshita M, et al.

Single-cell transcriptome landscape of circulating CD4(+) T cell populations in

autoimmune diseases. *Cell Genom.* 2024;4(2):100473.

22.   Tong L, Yu H, Huang X, Shen J, Xiao G, Chen L, et al. Current understanding of

osteoarthritis pathogenesis and relevant new approaches. *Bone Res.* 2022;10(1):60.

23.   Dash P, Fiore-Gartland AJ, Hertz T, Wang GC, Sharma S, Souquette A, et al.

Quantifiable predictive features define epitope-specific T cell receptor repertoires.

Nature. 2017;547(7661):89-93.

24.   Lagattuta KA, Kang JB, Nathan A, Pauken KE, Jonsson AH, Rao DA, et al.

Repertoire analyses reveal T cell antigen receptor sequence features that influence T

cell fate. *Nat Immunol.* 2022;23(3):446-57.

25.   Logunova NN, Kriukova VV, Shelyakin PV, Egorov ES, Pereverzeva A,

Bozhanova NG, et al. MHC-II alleles shape the CDR3 repertoires of conventional

and regulatory naive CD4(+) T cells. *Proc Natl Acad Sci U S A.*

2020;117(24):13659-69.

26.   Lu DR, Wu H, Driver I, Ingersoll S, Sohn S, Wang S, et al. Dynamic changes in the

regulatory T-cell heterogeneity and function by murine IL-2 mutein. *Life Sci*

Alliance. 2020;3(5).

27.   Ko A, Watanabe M, Nguyen T, Shi A, Achour A, Zhang B, et al. TCR Repertoires

of Thymic Conventional and Regulatory T Cells: Identification and

Characterization of Both Unique and Shared TCR Sequences. *J Immunol.*

2020;204(4):858-67.

28.   Wolf KJ, Emerson RO, Pingel J, Buller RM, and DiPaolo RJ. Conventional and

661   Regulatory CD4+ T Cells That Share Identical TCRs Are Derived from Common

662   Clones. *PLoS One.* 2016;11(4):e0153705.

663   29.   Pohar J, Simon Q, and Fillatreau S. Antigen-Specificity in the Thymic Development

664   and Peripheral Activity of CD4(+)FOXP3(+) T Regulatory Cells. *Front Immunol.*

665   2018;9:1701.

666   30.   Ooi JD, Petersen J, Tan YH, Huynh M, Willett ZJ, Ramarathinam SH, et al.

667   Dominant protection from HLA-linked autoimmunity by antigen-specific regulatory

668   T cells. *Nature.* 2017;545(7653):243-7.

669   31.   Jordan MS, Boesteanu A, Reed AJ, Petrone AL, Holenbeck AE, Lerman MA, et al.

670   Thymic selection of CD4+CD25+ regulatory T cells induced by an agonist self-

671   peptide. *Nat Immunol.* 2001;2(4):301-6.

672   32.   Hsieh CS, Zheng Y, Liang Y, Fontenot JD, and Rudensky AY. An intersection

673   between the self-reactive regulatory and nonregulatory T cell receptor repertoires.

674   *Nat Immunol.* 2006;7(4):401-10.

675   33.   Aschenbrenner K, D'Cruz LM, Vollmann EH, Hinterberger M, Emmerich J, Swee

676   LK, et al. Selection of Foxp3+ regulatory T cells specific for self antigen expressed

677   and presented by Aire+ medullary thymic epithelial cells. *Nat Immunol.*

678   2007;8(4):351-8.

679   34.   Lee HM, Bautista JL, Scott-Browne J, Mohan JF, and Hsieh CS. A broad range of

680   self-reactivity drives thymic regulatory T cell selection to limit responses to self.

681   *Immunity.* 2012;37(3):475-86.

35.   Gaublomme JT, Yosef N, Lee Y, Gertner RS, Yang LV, Wu C, et al. Single-Cell

Genomics Unveils Critical Regulators of Th17 Cell Pathogenicity. *Cell.*

2015;163(6):1400-12.

36.   Paroni M, Maltese V, De Simone M, Ranzani V, Larghi P, Fenoglio C, et al.

Recognition of viral and self-antigens by T(H)1 and T(H)1/T(H)17 central memory

cells in patients with multiple sclerosis reveals distinct roles in immune surveillance

and relapses. *J Allergy Clin Immunol.* 2017;140(3):797-808.

37.   Rao DA, Gurish MF, Marshall JL, Slowikowski K, Fonseka CY, Liu Y, et al.

Pathologically expanded peripheral T helper cell subset drives B cells in rheumatoid

arthritis. *Nature.* 2017;542(7639):110-4.

38.   Sakuragi T, Yamada H, Haraguchi A, Kai K, Fukushi JI, Ikemura S, et al.

Autoreactivity of Peripheral Helper T Cells in the Joints of Rheumatoid Arthritis. *J*

Immunol. 2021;206(9):2045-51.

39.   Cunningham KY, Hur B, Gupta VK, Arment CA, Wright KA, Mason TG, et al.

Patients with ACPA-positive and ACPA-negative rheumatoid arthritis show

different serological autoantibody repertoires and autoantibody associations with

disease activity. *Sci Rep.* 2023;13(1):5360.

40.   Shugay M, Bagaev DV, Zvyagin IV, Vroomans RM, Crawford JC, Dolton G, et al.

VDJdb: a curated database of T-cell receptor sequences with known antigen

specificity. *Nucleic Acids Res.* 2018;46(D1):D419-D27.

41.   Boyle LH, Goodall JC, and Gaston JS. Major histocompatibility complex class I-

restricted alloreactive CD4+ T cells. *Immunology.* 2004;112(1):54-63.

42.   Boyle LH, Goodall JC, Opat SS, and Gaston JS. The recognition of HLA-B27 by

705   human CD4(+) T lymphocytes. *J Immunol.* 2001;167(5):2619-24.

706   43.   Singh NK, Alonso JA, Devlin JR, Keller GLJ, Gray GI, Chiranjivi AK, et al. A

707   class-mismatched TCR bypasses MHC restriction via an unorthodox but fully

708   functional binding geometry. *Nat Commun.* 2022;13(1):7189.

709   44.   Malchow S, Leventhal DS, Lee V, Nishi S, Socci ND, and Savage PA. Aire

710   Enforces Immune Tolerance by Directing Autoreactive T Cells into the Regulatory

711   T Cell Lineage. *Immunity.* 2016;44(5):1102-13.

712   45.   Kieback E, Hilgenberg E, Stervbo U, Lampropoulou V, Shen P, Bunse M, et al.

713   Thymus-Derived Regulatory T Cells Are Positively Selected on Natural Self-

714   Antigen through Cognate Interactions of High Functional Avidity. *Immunity.*

715   2016;44(5):1114-26.

716   46.   Leonard JD, Gilmore DC, Dileepan T, Nawrocka WI, Chao JL, Schoenbach MH, et

717   al. Identification of Natural Regulatory T Cell Epitopes Reveals Convergence on a

718   Dominant Autoantigen. *Immunity.* 2017;47(1):107-17 e8.

719   47.   Hsieh CS, Liang Y, Tyznik AJ, Self SG, Liggitt D, and Rudensky AY. Recognition

720   of the peripheral self by naturally arising CD25+ CD4+ T cell receptors. *Immunity.*

721   2004;21(2):267-77.

722   48.   Pacholczyk R, Ignatowicz H, Kraj P, and Ignatowicz L. Origin and T cell receptor

723   diversity of Foxp3+CD4+CD25+ T cells. *Immunity.* 2006;25(2):249-59.

724   49.   Al Khabouri S, Benson RA, Prendergast CT, Gray JI, Otto TD, Brewer JM, et al.

725   TCRbeta Sequencing Reveals Spatial and Temporal Evolution of Clonal CD4 T

Cell Responses in a Breach of Tolerance Model of Inflammatory Arthritis. *Front*

Immunol. 2021;12:669856.

50.   Eggenhuizen PJ, Cheong RMY, Lo C, Chang J, Ng BH, Ting YT, et al. Smith-

specific regulatory T cells halt the progression of lupus nephritis. *Nat Commun.*

2024;15(1):899.

51.   Ohkura N, Hamaguchi M, Morikawa H, Sugimura K, Tanaka A, Ito Y, et al. T cell

receptor stimulation-induced epigenetic changes and Foxp3 expression are

independent and complementary events required for Treg cell development.

Immunity. 2012;37(5):785-99.

52.   Satija R, Farrell JA, Gennert D, Schier AF, and Regev A. Spatial reconstruction of

single-cell gene expression data. *Nat Biotechnol.* 2015;33(5):495-502.

53.   Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Mauck WM, 3rd, et al.

Cell Hashing with barcoded antibodies enables multiplexing and doublet detection

for single cell genomics. *Genome Biol.* 2018;19(1):224.

54.   Loza M, Teraguchi S, Standley DM, and Diez D. Unbiased integration of single cell

transcriptome replicates. *NAR Genom Bioinform.* 2022;4(1):lqac022.

55.   Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr:

interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC*

Bioinformatics. 2013;14:128.

56.   Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell

lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics.*

2018;19(1):477.

57. Chiffelle J, Genolet R, Perez MA, Coukos G, Zoete V, and Harari A. T-cell

repertoire analysis and metrics of diversity and clonality. *Curr Opin Biotechnol.*

2020;65:284-95.

58. Bolotin DA, Poslavsky S, Mitrophanov I, Shugay M, Mamedov IZ, Putintseva EV,

et al. MiXCR: software for comprehensive adaptive immunity profiling. *Nat*

Methods. 2015;12(5):380-1.

59. Wilamowski J, Xu Z, Ismanto HS, Li S, Teraguchi S, Llamas- Covarrubias MA, et

al. InterClone: Store, Search and Cluster Adaptive Immune Receptor Repertoires.

bioRxiv. 2022:2022.07.31.501809.

Figure legends

Figure 1. Two transcriptionally different Th17 cell populations exist in inflamed joints of arthritic ZAC mice. a) a) Normally, during T cell development in thymus, high-intensity interactions between a TCR and self-pMHC will induce either clonal deletion or selection into the Treg repertoire (upper plot); however, a systemic weakening of TCR signaling could disrupt this mechanism by allowing T cells with moderately to strongly self-reactive TCRs to evade negative selection and enter the mature conventional Tconv repertoire (lower plot). These cells would later be activated by self-peptides in the periphery, accumulate in target tissues and orchestrate tissue damage. **b)** Lateral view of a hindpaw of a 23-week old arthritic ZAC mouse. **c)** UMAP plot of Tconv cells obtained from joints, draining lymph node (dLN) and spleen of arthritic ZAC mice colored by cluster. **d)** Proportion of annotated Tconv cell populations in tissues of arthritic ZAC mice. **e)** Relative expression of canonical Tconv markers across gene expression clusters. Circle color and intensity represent the average expression whereas circle size represents the percentage of cells from each cluster expressing the gene. **f)** Volcano plot of the differentially expressed genes (DEG) among cluster 1 and cluster 4. **g)** Top 5 gene Ontology (GO) Terms enriched in DEGs from cluster 1 vs cluster 4. The color intensity of bars represents the odds ratio (OR) of enrichment. **h)** UMAP plot of Tconv cells from tissues of arthritic ZAC mice colored according to their expression of *Il17a*, *Il17f*, *Il22* or *Tnfsf11*. **i)** Module scores for the gene signatures of migratory Th17 and pathogenic Th17 cells in a public scRNAseq dataset of CD4[+] T cells from inflamed synovia of RA patients (n=3)(18). Created with BioRender.com

Figure 2. The TCR repertoire of pathogenic Th17 cells is unique, convergent, and self-

reactive. a) UMAP plot of Tconv cells from tissues of arthritic ZAC mice, colored

according to clonal proportion classification. **b)** Morisita-Horn index of the paired CDR3

amino acid similarity between the migratory and pathogenic Th17 repertoires (gray bar);

and between both Th17 populations vs the other Tconv phenotypes in ZAC tissues (brown

and green bars). **c)** Distribution of neighbor distances to the nearest 25th percentile

(methods) in the alpha chain repertoires of naïve, migratory Th17 and pathogenic Th17

cells. **d)** Distribution of TiRP scores for WT Tconv, and overall Tconv, migratory Th17,

and pathogenicTh17 repertoires of ZAC mice. **e)** Median TiRP score change of overall

Tconv, migratory Th17, and pathogenicTh17 repertoires of ZAC mice relative to WT

Tconv. **f)** Description of the method employed for the assessment of repertoire similarity to

reference WT Treg and WT Tconv repertoires. **(g-h)** Normalized fraction of query WT,

SKG and ZAC Tconv repertoires **(g)** or WT Tconv, WT Treg, ZAC migratory and

pathogenic Th17 repertoires **(h)** matching with reference WT Tconv (magenta section) or

WT Treg repertoires (blue section) obtained from a representative reference dataset(25). *:

$p<0.05$; **:$p<0.01$; Wilcoxon Test.

Figure 3. Intra-Treg-Tconv similarity in WT, SKG and ZAC repertoires. a) Morisita-

Horn index calculated at the exact CDR3 amino acid level between the Tconv and Treg

compartments of WT, SKG and ZAC mice. **b)** Heatmap of weighted neighbor distance

correlations between WT, SKG and ZAC Treg and Tconv repertoires as calculated by

TCRdist (methods), higher correlation scores indicate closer repertoires.

Figure 4. Consequences of weakened TCR signaling in the shaping of Treg and Tconv repertoires and autoimmunity. The SKG (W163C) and ZAC (H165A) mutations in the ZAP-70 gene reduce its affinity for phosphorylated CD3 ζ. As a result, TCR signaling is attenuated. A systemic weakening of TCR signaling would disrupt the T cell selection process by allowing the maturation of self-reactive T cells into the Tconv compartment. Simultaneously, a bifurcation of the Treg-Tconv repertoires is observed, with the consequent loss of certain antigen-specific Treg cells that could effectively suppress autoimmune responses in the periphery. These events together, favor the activation of self-reactive T conv cells, their differentiation into pathogenic phenotypes and their accumulation in target tissues. Created with BioRender.com

Supplementary Figure 1. Single-cell transcriptomic analysis of Tconv cells obtained from tissues of arthritic ZAC mice. a) UMAP plot of integrated Tconv cells obtained from joints, draining lymph node (dLN) and spleen of arthritic ZAC mice. Each panel is colored according to cell source. **b)** Proportion of annotated Tconv cell populations in tissues of each arthritic ZAC mice. **c)** Expression of top 5 gene markers defining each cluster; clusters were subsampled to 100 cells for visualization. **d)** Heatmap of the top 5 enriched GO Terms in each cluster, color scale represents -log10 of adjusted p values. **(e-g)** UMAP plots of Tconv cells from tissues of arthritic ZAC mice colored according to their

expression of **e)** inferred pseudotime values **f)** markers of Th17.1 cells: *Ifng*, *Tbx21* and

Cxcr3; and **g)** markers of exFoxP3 cells: *Sox4*, *Ccr6*, *Ccl20*, *Il23r*, and *Il17re.*

Supplementary Figure 2. Single-cell transcriptomic and repertoire analysis of Tconv

cells obtained from spleens of WT, SKG and ZAC mice. UMAP plots of integrated

Tconv cells obtained from spleens of WT (n=2), SKG (n=2) and arthritic ZAC mice (n=2).

Colored by **(a)** cell source; and **(b)** cluster identity. Unsupervised clustering by the Louvain

algorithm was applied, and clusters were annotated as described in the Supplementary

Notes. **c)** Relative expression of canonical Tconv markers across gene expression clusters.

Circle color and intensity represents average gene expression whereas circle size represents

the percentage of cells from each cluster expressing the gene. **d)** Proportion of annotated

Tconv cell populations in spleens of WT, SKG, and arthritic ZAC mice. **e)** Expression of

top 5 gene markers defining each cluster. Clusters were subsampled to 100 cells for

visualization. Significant enriched GO Terms in each cluster are depicted to the right, if no

GO Terms were significantly enriched, top GO Term is shown and font color is grayed out.

f) Heatmap of overlap between cell annotations in the ZAC Tconv dataset (columns) and

the WT, SKG, ZAC spleen (Spleen Tconv) dataset (rows). Cell colors represent the

proportion of ZAC Tconv annotations within Spleen Tconv annotations (i.e., each column

sums up to 1). **g)** UMAP plot of integrated Th17 cells annotated in spleens of WT (n=2),

SKG (n=2) and arthritic ZAC mice (n=2). Each panel is colored according to cell source. **h)**

Distribution of clonal proportion classes across gene expression clusters in spleens of WT,

SKG and ZAC Tconv cells.

Supplementary Figure 3. The pathogenic Th17 gene signature is uniquely detected in CD4$^+$ T cells from inflamed synovia of rheumatoid arthritis (RA) patients. Gene signatures of migratory Th17 and pathogenic Th17 cells were obtained and used for calculation of module scores in public scRNAseq datasets of **a)** Th17 cells obtained from spleens of WT, SKG and ZAC mice (n=2 each), **c)** CD4+ T cells from peripheral blood of RA patients (n=1)(19), **d)** CD4+ T cells from synovia and infrapatellar fat pad (IPFP) tissue of osteoarthritis (OA) patients (n=3)(20), **e)** Circulating Th17 cells from healthy donors (n=3), and patients with autoimmune diseases (AD) other than RA: myasthenia gravis (MG) (n=3), multiple sclerosis (MS) (n=4), systemic lupus erythematosus (SLE) (n=3)(21); **b**) Violin plot of module scores for the gene signatures of migratory Th17 (left) and pathogenic Th17 (right) cells in Th17 cells from spleens of WT, SKG and arthritic ZAC mice.

Supplementary Figure 4. CD4$^+$ T cells from RA synovia expressing the pathogenic Th17 gene signature are different from Th17.1 or exFoxP3 cells. UMAP plot of CD4$^+$ T cells from inflamed synovia of RA patients (n=3)(18), cells are colored according to the density of expression of individual markers of Th17.1 (*Ifng*, *Tbx21* and *Cxcr3*) **(a)** and exFoxP3 cells (*Sox4*, *Ccr6*, *Ccl20*, *Il23r*, and *Il17re*) **(c)**; or as module scores **(b, d)**. The region corresponding to the expression of the pathogenic Th17 gene signature is enclosed by a dashed line in **b** and **d**.

Supplementary Figure 5. Repertoire analysis of Tconv cells obtained from tissues of

arthritic ZAC mice. Distribution of clonal proportion classes across **(a)** ZAC Tconv gene

expression clusters. **(b)** gene expression cluster separated by mouse replicate. **c)**

Distribution of neighbor distances to the nearest 25th percentile (methods) in the paired

chain (left) and beta chain (right) repertoires of Naïve, migratory Th17 and pathogenic

Th17 cells. **d)** Public clones and their clonal proportion classification and distribution

across Tconv phenotypes in each mouse replicate

Supplementary Figure 6. Similarity of TCR repertoires of ZAP-70 mutant Tconv cells,

to reference WT Treg and WT Tconv repertoires. a) Distribution of TCR-intrinsic

regulatory potential (TiRP) scores for Tconv and Treg repertoires of WT, SKG and ZAC

mice. **b)** Median TiRP score change of Tconv and Treg repertoires of WT, SKG and ZAC

mice relative to the WT Tconv repertoire. c) Normalized fraction of query WT Tconv, WT

Treg, and SKG and ZAC Tconv repertoires matching with reference WT Tconv (magenta

section) or WT Treg repertoires (blue section) obtained from reference datasets(26-28). **d)**

Normalized fraction of query WT Tconv, WT Treg, and ZAC migratory and pathogenic

Th17 repertoires matching with reference WT Tconv or WT Treg repertoires obtained from

reference datasets(26-28). No significant matches to reference WT Treg or WT Tconv in

the Ko et al., 2020 dataset were found in pathogenic Th17 cells.

Supplementary Figure 7. Intra-Treg-Tconv similarity in WT, SKG and ZAC

repertoires. a) Morisita-Horn index calculated at the exact CDR3 amino acid level

between the Tconv and Treg compartments of WT, SKG and ZAC mice, split by Tconv

phenotype (x axis). **(b-h)** Clonotype tracking plots of intra-Treg-Tconv shared clones

across Treg and Tconv gene expression clusters for WT **(b-c)**, SKG **(d-e)**, and ZAC **(f-h)**

mice. No Treg-Tconv sharing was detected in ZAC Replicate 3.

Figure 1

Figure 1. Two transcriptionally different Th17 cell populations exist in inflamed joints of arthritic ZAC mice. a) Normally, during T cell development in thymus, high-intensity interactions between a TCR and self-pMHC will induce either clonal deletion or selection into the Treg repertoire (upper plot); however, a systemic weakening of TCR signaling could disrupt this mechanism by allowing T cells with moderately to strongly self-reactive TCRs to evade negative selection and enter the mature conventional Tconv repertoire (lower plot). These cells would later be activated by self-peptides in the periphery, accumulate in target tissues and orchestrate tissue damage. **b)** Lateral view of a hindpaw of a 23-week old arthritic ZAC mouse. **c)** UMAP plot of Tconv cells obtained from joints, draining lymph node (dLN) and spleen of arthritic ZAC mice colored by cluster. **d)** Proportion of annotated Tconv cell populations in tissues of arthritic ZAC mice. **e)** Relative expression of canonical Tconv markers across gene expression clusters. Circle color and intensity represent the average expression whereas circle size represents the percentage of cells from each cluster expressing the gene. **f)** Volcano plot of the differentially expressed genes (DEG) among cluster 1 and cluster 4. **g)** Top 5 gene Ontology (GO) Terms enriched in DEGs from cluster 1 vs cluster 4. The color intensity of bars represents the odds ratio (OR) of enrichment. **h)** UMAP plot of Tconv cells from tissues of arthritic ZAC mice colored according to their expression of *Il17a*, *Il17f*, *Il22* or *Tnfsf11*. **i)** Module scores for the gene signatures of migratory Th17 and pathogenic Th17 cells in a public scRNAseq dataset of CD4[+] T cells from inflamed synovia of RA patients (n=3[18]. Created with BioRender.com

Figure 2

Figure 2. The TCR repertoire of pathogenic Th17 cells is unique, convergent, and self-reactive. a) UMAP plot of Tconv cells from tissues of arthritic ZAC mice, colored according to clonal proportion classification. **b)** Morisita-Horn index of the paired CDR3 amino acid similarity between the migratory and pathogenic Th17 repertoires (gray bar); and between both Th17 populations vs the other Tconv phenotypes in ZAC tissues (brown and green bars). **c)** Distribution of neighbor distances to the nearest 25th percentile (methods) in the alpha chain repertoires of naïve, migratory Th17 and pathogenic Th17 cells. **d)** Distribution of TiRP scores for WT Tconv, and overall Tconv, migratory Th17, and pathogenicTh17 repertoires of ZAC mice. **e)** Median TiRP score change of overall Tconv, migratory Th17, and pathogenicTh17 repertoires of ZAC mice relative to WT Tconv. **f)** Description of the method employed for the assessment of repertoire similarity to reference WT Treg and WT Tconv repertoires. **(g-h)** Normalized fraction of query WT, SKG and ZAC Tconv repertoires **(g)** or WT Tconv, WT Treg, ZAC migratory and pathogenic Th17 repertoires **(h)** matching with reference WT Tconv (magenta section) or WT Treg repertoires (blue section) obtained from a representative reference dataset[25]. *: p<0.05; **:p<0.01; Wilcoxon Test.

Figure 3

Figure 3. Intra-Treg-Tconv similarity in WT, SKG and ZAC repertoires. a) Morisita-Horn index calculated at the exact CDR3 amino acid level between the Tconv and Treg compartments of WT, SKG and ZAC mice. **b)** Heatmap of weighted neighbor distance correlations between WT, SKG and ZAC Treg and Tconv repertoires as calculated by TCRdist (methods), higher correlation scores indicate closer repertoires.

Figure 4

Figure 4. Consequences of weakened TCR signaling in the shaping of Treg and Tconv repertoires and autoimmunity. The SKG (W163C) and ZAC (H165A) mutations in the ZAP-70 gene cause a reduction in its affinity for phosphorylated CD3ζ and the consequent attenuation of TCR signaling. A systemic weakening of TCR signaling would disrupt the T cell selection process by allowing the maturation of self-reactive T cells into the Tconv compartment. Simultaneously, a bifurcation of the Treg-Tconv repertoires is observed, with the consequent loss of certain antigen-specific Treg cells that could effectively suppress autoimmune responses in the periphery. These events together, favor the activation of self-reactive T conv cells, their differentiation into pathogenic phenotypes and their accumulation in target tissues. Created with BioRender.com