Pathogenic Th17 cells in arthritogenic mice share T cell receptor features with Wild-type Tregs

Mara A. Llamas-Covarrubias[1,2,3*], Atsushi Tanaka[4,5,6], Martin Loza[7], Diego Diez[8], Zichang Xu[1], Ee Lyn Lim[5], Shunsuke Teraguchi[2,9], Shimon Sakaguchi[4,5], Daron M. Standley[1,2 *]

[1]Laboratory of Systems Immunology, WPI Immunology Frontier Research Center, Osaka University, Suita, Japan; [2]Department of Genome Informatics, Research Institute for Microbial Diseases (RIMD), Osaka University, Suita, Japan; [3]Departamento de Biología Molecular y Genómica, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara, México; [4]Department of Experimental Pathology, Institute for Frontier Medical Sciences, Kyoto University, Kyoto, Japan; [5]Laboratory of Experimental Immunology, WPI Immunology Frontier Research Center, Osaka University, Suita, Japan; [6]Department of Frontier Research in Tumor Immunology, Center of Medical Innovation and Translational Research, Graduate School of Medicine, Osaka University, Osaka, Japan; [7]Laboratory of functional analysis in silico, Human Genome Center, The University of Tokyo, Tokyo, Japan; [8]Quantitative Immunology Research Unit, Immunology Frontier Research Center, Osaka University, Suita, Japan; [9]Faculty of Data Science, Shiga University, Hikone, Japan

[*]Corresponding authors:

22   Laboratory of Systems Immunology, WPI Immunology Frontier Research Center, Osaka

23   University, Suita, Japan. Tel: +81-6-6879-8366; Fax: +81-6-6879-8366. Email:

24   standley@biken.osaka-u.ac.jp; mara.llamas@biken.osaka-u.ac.jp

Abstract

Mutations in the ZAP-70 gene that cause moderate attenuation of T cell receptor (TCR) signaling in mice, can result in autoimmune manifestations akin to rheumatoid arthritis (RA). Here, we characterized the single-cell gene expression profiles and TCR repertoires of conventional (Tconv) and regulatory (Treg) CD4$^+$ T cells of arthritic (ZAC), poised (SKG) ZAP-70 mutant, and wild-type (WT) mice. We identified two Th17 cell subtypes in the joints of ZAC mice that were characterized by distinct transcriptional profiles and TCR repertoires, one of which exhibited a pathogenic signature and occurred exclusively in inflamed joints. Such a pathogenic signature was also uniquely detected in CD4$^+$ T cells obtained from inflamed joints of human RA patients. The TCR repertoire of pathogenic Th17 cells showed signs of increased intra-repertoire similarity (convergence) and was skewed toward the WT Treg rather than the WT Tconv repertoire. In addition, the overall similarity between the Treg and Tconv repertoires was severely reduced in arthritic mice. Our results support a model where, upon moderate ZAP-70-mediated signal weakening, T cells that would normally develop into Tregs, instead develop into self-reactive Tconvs, resulting in a breakdown in self-tolerance and susceptibility to autoimmune arthritis.

Introduction

The development of autoimmune diseases is influenced by various factors, with the presence of self-reactive conventional T cells (Tconvs) being a critical element[1]. During T cell

maturation, thymocytes expressing T cell receptors (TCRs) with high affinity to self-peptide-

MHC complexes are typically eliminated from the repertoire via negative selection or

diverted to the T regulatory (Treg) cell lineage[2,3]. Nonetheless, a subset of self-reactive cells

persists in the periphery of healthy individuals, usually kept in check by the suppressive

activity of Tregs[4,5]. Antigen recognition by the TCR is key for both Treg development in the

thymus and for their function in peripheral tissues[6-8]; and, it is generally accepted that the

Treg repertoire is biased to self-recognition[9-12]. Therefore, disruption of the normal balance

between Treg and Tconv repertoires can bias the immune system toward autoimmunity.

It has been previously reported that specific mutations in the ZAP-70 gene that moderately

weaken downstream TCR signaling upon antigen recognition, altering the Treg-Tconv

repertoire balance, predispose mice to an autoimmune disease closely resembling rheumatoid

arthritis (RA)[13-15]. SKG mice, which are characterized by a W163C mutation in the ZAP-70

gene that causes a fifteen-fold reduction in ZAP-70 affinity for phosphorylated CD3ζ[13],

develop disease upon secondary stimulation of innate immune responses (i.e. by mannan

injection). In contrast, ZAC mice, which carry an H165A mutation that causes only an eight-

fold reduction, spontaneously develop chronic autoimmune arthritis, even under specific

pathogen-free (SPF) conditions[15]. The underlying mechanism has been ascribed to disruption

of the normal thymic selection process that allows T cells with moderately to strongly self-

reactive TCRs to escape negative selection[14,15]. As a consequence, in ZAP-70 mutant mice,

some T cells harboring TCRs that would normally be observed in the Treg repertoire instead

develop into self-reactive Tconvs. Here, we refer to this mechanism as the "repertoire shift"

hypothesis.

In order to test this hypothesis, we examined the transcriptome and immune repertoires of

CD4$^+$ T cells in WT, non-arthritic SKG and arthritic ZAC mice at the single-cell level. We

found migratory and pathogenic populations of Th17 cells in arthritic mice. Interestingly,

pathogenic cells highly expressed the osteoclastogenic factor *Tnfsf11* (encoding for RANKL)

and uniquely localize in inflamed joints but were absent in WT and SKG mice. A group of

cells with a similar pathogenic signature was uniquely identified in T cells from human RA

joints. Furthermore, the TCR repertoire of pathogenic Th17 cells was unique, convergent,

and overlapped with the repertoire of WT Tregs. Finally, the overall similarity between the

Treg and Tconv repertoires was severely reduced in arthritic mice, suggesting that a depletion

of critical components of the Treg repertoire accompanies expansion of Tconvs bearing

TCRs that are biased toward self-reactivity. Our data thus supports the repertoire shift

hypothesis and suggests that the autoimmunity observed upon moderate reduction of TCR

signaling strength results from T cells bearing self-reactive TCRs that evade negative

selection. As a consequence, TCRs typical of Tregs are expressed in Tconv cells, and this,

along with a depletion of specific receptors in the Treg compartment, lead to autoimmunity.

Results

Arthritic ZAC mice harbor two groups of Th17 cells with distinct gene programs in their joints

The ZAP-70 mutant mice (ZAC, SKG) used in this study are characterized by the spontaneous development of autoimmune arthritis (Fig. 1a), with disease progression typically accelerated in the SKG model through the stimulation of innate immunity. Unlike other widely used mouse models of arthritis, such as collagen-induced arthritis (CIA)[16], the arthritis in ZAP-70 mutant mice is genuinely autoimmune, mediated primarily by T-cells, and closely resembles human RA [13-15]. Therefore, ZAP-70 mutant mouse models are particularly well-suited for investigating self-reactive TCR repertoires. To study the transcriptomes and immune repertoires of self-reactive Tconv (CD3$^+$CD4$^+$Foxp3$^-$) cells in-depth, we sorted cells from the joints, draining lymph nodes (dLN), and spleens of arthritic ZAC mice and performed RNA+V(D)J single-cell sequencing. Gene expression analysis revealed a considerable overlap in the transcriptional profiles of spleens and dLNs but a distinct profile for joint cells (Supplementary Fig. 1a). Unsupervised clustering delineated nine gene expression clusters with similar distribution in each tissue across replicates (Supplementary Note 1, Fig. 1b-c, Supplementary Fig. 1b). Two clusters (1 and 4) showed canonical Th17 markers (*Rorc, Il17a, Il17f*), but were distinct in their gene expression programs and tissue abundances (Fig. 1b-d, Supplementary Fig. 1b-d). Cluster 1 was abundant in joints, representing ~50% of cells obtained from this tissue, but was also detected in the spleen and dLNs at smaller proportions (~15-17%). Relative to cluster 4, cells in cluster 1 differentially expressed *Klf2*, *S1pr1* and chemokine receptors such as *Ccr2*, *Ccr4* and *Ccr6*,

suggesting a phenotype of migratory cells (Fig. 1e, Supplementary Table 4). Consistently,

pathway analysis revealed an enrichment in the regulation of T cell migration (GO:

20000404) and cell chemotaxis (GO: 0060326) gene ontology (GO) Terms (Fig. 1f,

Supplementary Table 5). In contrast, cluster 4 was almost exclusively observed in joints,

representing approximately 27% of joint cells but less than 2% of dLN cells and less than

1% of Spleen cells (Fig. 1c, Supplementary Fig.1b). The top differentially expressed gene

(DEG) in cluster 4 relative to cluster 1 was *Tnfsf11*, which codes for RANKL, a critical factor

involved in osteoclast differentiation and activation that plays a central role in bone

destruction in RA[17] (Fig. 1e). Two other osteoclastogenic genes (*Csf1*, *Anxa2*) were highly

expressed in this cluster as well. Overall, we observed a significant enrichment in the

osteoclast differentiation (GO:0030316) GO term in cluster 4 as compared to cluster 1

(Supplementary Table 5). In addition, transcripts involved in the response to TNF stimuli

(for example: *Tnfrsf18*, *Tnfrsf9*, *Tnfrsf4*, and *Tnf*; encoding for GITR, 4-1BB, OX40, and

TNF-α respectively), and the cellular response to tumor necrosis factor (GO:0072356) GO

terms were enriched in cluster 4. Moreover, this cluster uniquely expressed transcription

factors associated with a pathogenic Th17 phenotype: *Rbpj*[18], *Crem*[19], *Atf3*[20], *Bhlhe40*[21], and

Igfbp7[22] (Fig. 1e). Finally, the transcription factor *Klf2*, a known inhibitor of NF-κB-

dependent genes that also prevents Tfh differentiation and RORγt expression[23,24] was absent

from cluster 4 (Supplementary Fig. 1e). These results indicate that cluster 4 contains

pathogenic Th17 cells with an increased potential to activate osteoclasts and thus promote

bone damage. Interestingly, cluster 1 and cluster 4 could be clearly distinguished by their

differential expression of *Il22* or *Tnfsf11* (Fig. 1g). Finally, we annotated clusters 1 and 4 as

migratory and pathogenic Th17, respectively. In order to explore the dynamics of the

different Tconv cell states, we performed trajectory inference setting the naïve cluster of cells

as the starting point in pseudotime. This resulted in the identification of the pathogenic Th17

cluster as one of the endpoint lineage states (Fig. 1h), consistent with their effector phenotype.

A population of pathogenic Th17 cells characterized by the expression of Th1 cell markers

(*Ifng*; *Tbx21*, encoding for T-bet; and *Cxcr3;* Th17.1 cells)[25], and a group of osteoclastogenic

Th17 cells derived from CD4$^+$ T cells previously expressing FoxP3 (exFoxP3) and

characterized by the expression of *Sox4*, *Ccr6*, *Ccl20*, *Il23r* and *Tnfsf11*[26] have been

previously reported to be enriched in arthritic joints; however, other than *Tnfsf11* we did not

detect clear and specific expression of these markers in cluster 4, suggesting that it represents

a distinct group of cells (Supplementary Fig. 1f-g).

To contextualize our findings in arthritic ZAC mice tissues, we replicated the analyses in WT

and non-arthritic SKG mice. For this, we sampled Tconv from the spleens of SKG and WT

mice, and integrated the ZAC spleen data in order to facilitate direct comparison of clusters

(Supplementary Note 2, Supplementary Fig. 2a-g). This approach revealed that the general

Th17 cell abundance increased progressively from WT (less than 1%) to SKG (2-9%) to ZAC

(~36%) mice (Supplementary Fig. 2d). To analyze whether the transcriptional programs of

these Th17 cells are similar to the transcriptional programs of the two Th17 cells identified

in ZAC tissues, we calculated gene module scores for the signature genes from migratory

and pathogenic Th17 cells (Supplementary Table 6). Expression of the migratory Th17

signature was clearly detected and gradually increased from WT, to SKG to ZAC splenic

Th17 cells; however, the pathogenic Th17 signature was largely undetected (Fig. 2a-b,

Supplementary Fig. 2h). This result indicates that pathogenic Th17 cells are uniquely present

in the inflamed joints of arthritic mice.

In summary, we demonstrated the presence of two distinct groups of Th17 cells in arthritic

ZAC mice: migratory Th17 cells, expressing *Il22* and the transcription factor *Klf2*; and,

pathogenic Th17 cells, which highly express *Tnfsf11* and uniquely localize within inflamed

joints.

The transcriptional signature of pathogenic Th17 cells is present in inflamed joints of

human RA but not in other autoimmune diseases or non-autoimmune arthritis

To investigate the presence of cells resembling pathogenic Th17 cells in human tissues, we

calculated migratory and pathogenic Th17 gene signature module scores in publicly available

single-cell RNAseq datasets. We analyzed data obtained from inflamed synovia[27] and

PBMCs[28] of RA patients; and, for comparison, we also analyzed data from the synovia and

infrapatellar fat pad (IPFP) of osteoarthritis (OA) patients[29] as well as peripheral CD4+ T

cells from healthy controls and patients with other autoimmune diseases (myasthenia gravis,

multiple sclerosis, and systemic lupus erythematosus)[30]. Detailed descriptions of the datasets

are provided in Supplementary Table 7.

Our analysis revealed that the migratory Th17 signature was clearly detected in CD4$^+$T cells

from inflamed synovia or peripheral blood from RA patients (Fig. 2c-d), as well as in CD4$^+$T

cells obtained from joint tissues of OA patients (Fig. 2e), and in circulating Th17 cells in

healthy controls and patients with other autoimmune diseases (Fig. 2f).

In contrast, expression of the pathogenic Th17 signature was nearly undetectable in all

samples (Fig. 2 d-f) except in CD4$^+$ T cells derived from inflamed joints of RA patients (Fig.

2c) where not only the highest expression of this gene signature was detected, but its

expression was found to be non-overlapping to the expression of the migratory Th17

signature, mirroring our observations in ZAC joints. Notably, as in ZAC joints, the

expression of Th17.1 or exFoxP3 markers was not specifically detected in the cluster of cells

expressing the pathogenic Th17 gene signature in human RA joints (Supplementary Fig. 3a-

d).

OA is the most common form of arthritis and is mainly provoked by the mechanical wear

and tear of joints, and, even though inflammation is involved in its pathogenesis, it is not a

disease of autoimmune origin[31]. Thus, our results indicate that cells with a pathogenic Th17

transcriptional signature are likely to be specifically involved in the pathogenesis of human

autoimmune arthritis.

Pathogenic Th17 cells are expanded, convergent, and contain public clones

Having identified two groups of potentially self-reactive Th17 cells with characteristic

transcriptomic features, we next asked whether their TCR repertoires were also distinct. First,

we examined the degree of clonal expansion by calculating the proportion of a whole

repertoire that each clone represented (clonal proportion) in each replicate, and subsequently

classified clones as: small, medium or large (methods). We found that large and medium

clones were abundant in both Th17 cell groups (Fig. 3a-b, Supplementary Fig. 4a). Next, we

evaluated the repertoire similarity of both Th17 groups to other Tconv populations and

between each other by using the Morisita-Horn index[32]. At the exact paired CDR3 amino

acid sequence level, the repertoire of migratory Th17 cells was largely similar to other Tconv

repertoires from ZAC tissues (Fig. 3c). On the other hand, the pathogenic Th17 repertoire

was less similar to other Tconv cells. Consistently, the similarity between the two Th17 cell

repertoires was rather low (Fig. 3c). These results suggest that pathogenic Th17 cells possess

a unique repertoire, presumably due to the targeting of a specific set of antigens.

Given the unique repertoire of pathogenic Th17 cells, we next evaluated intra-repertoire TCR

similarity as a measure of repertoire convergence. For this purpose, we used TCRdist[33],

which calculates the distance between the CDR1, CDR2 and CDR3 amino acid sequences of

single or paired TCR chains. For this analysis, we independently calculated the TCR

distances among all receptors in the two Th17 clusters, as well as the naïve cluster for

reference. A bias in the distribution of the distances to lower values indicates a higher

similarity of its receptors, and therefore can be interpreted as an indication of repertoire

convergence. We found that the TCR distance distribution of the pathogenic Th17 repertoire

trended lower in paired and single chain analysis and was significantly lower in the alpha

chain (Fig. 3d). This result indicates that the repertoire of pathogenic Th17 cells is more

convergent than that of migratory Th17 cells, especially in the alpha chain. Given their

specific location in joints, and the evidence of similar cells occurring in human RA but not

OA joints, our results suggest that pathogenic Th17 may recognize a particular self-antigen

in the affected tissue and that the alpha chain may play a key role in such recognition.

Finally, we identified a total of 9 unique clones shared in two or more individuals (public

clones) (Fig. 3e, Supplementary Table 8). Interestingly, public clone 1 occurred in 3 out of 4

replicates, suggesting that this clone is likely to be involved in the recognition of relevant

arthritogenic self-antigens (Fig. 3e).

Together, these results indicate that although the Th17 response in arthritic ZAC mice is

polyclonal, the repertoire of tissue-specific pathogenic Th17 cells is unique and shows high

TCR convergence, suggesting the recognition of a limited set of self-antigens. Moreover, a

number of highly expanded public clones was observed within the inflamed tissue.

Effector Treg cells are abundant in the joints of arthritic ZAC mice but potentially

suppressed by cell crosstalk with self-reactive Tconvs

As discussed previously, the failure of Tregs to suppress self-reactive T cells and subsequent

inflammation is a key factor in the development of autoimmune diseases. To investigate the

Treg landscape in ZAP-70 mutant mice, we sorted Treg cells from the joints, dLN, and

spleens of arthritic ZAC mice as well as from spleens of SKG and WT mice and performed

single-cell RNA and V(D)J sequencing. In the tissues of ZAC mice, we identified seven gene

expression clusters, including two groups of effector cells (*Ccr2, Cxcr3, Gzmb*) and one

cluster of cells with a tissue resident memory (Trm) signature (*Il7r, Sell, Cd44, Cd69, Itgae*)

(Supplementary Fig. 5a-c), which were notably expanded and abundant in joints

(Supplementary Note 3, Supplementary Fig. 5a-g, Supplementary Table 10, 11). Conversely,

in WT and SKG Tregs, we observed limited effector and Trm cells, while naïve cells were

abundant (Supplementary Note 4, Supplementary Fig. 6a-f). Remarkably, despite the evident

expansion in the Treg compartment of ZAC mice, only one public clone could be identified

(Supplementary Table 12). These findings suggest that Trm and effector Treg cells undergo

expansion in the tissues of arthritic ZAC mice.

Despite the abundance of effector Treg cells in the sampled tissues, ZAC mice were overly

autoreactive. To elucidate the disparities in cell activity among WT, SKG, and ZAC effector

Tregs, we performed DEG analysis across mouse models. Our analysis revealed that while

the transcriptional programs of WT and SKG Treg cells were largely similar, several genes

involved in TNF signaling were uniquely expressed in ZAC effector Tregs (Supplementary

Fig. 7a-b, Supplementary Table 13). While the exact effect of TNF signaling on Treg effector

function remains unclear, studies have suggested its suppressive effects on Treg function in

RA patients[34].

Tregs can exert their suppressive functions by various mechanisms, including physical

interactions with target cells or secretion of soluble factors[35,36]. Therefore, we sought to

identify potentially significant cell-cell crosstalk events between Tconv Th17 cells and

effector Tregs cells by predicting cell communication events using Cellphone DB[37].

Consistent with the increased TNF signaling activity we observed in ZAC mice, there was a

gradual increase in the number of predicted cell communication events involving members

of the TNF family from WT, SKG, and ZAC Th17 cells toward Treg cells (Supplementary

Fig. 7c). Thus, our findings suggest that despite the abundance of effector Treg cells in ZAC

mice, their function is impaired and one of the mechanisms may involve cell crosstalk with

pathogenic Th17 cells through members of the TNF signaling family.

TCR repertoires of ZAP-70 mutant Tconv, particularly pathogenic Th17, cells are more

WT Treg- than WT Tconv-like

Because T cells harboring self-reactive TCRs would likely be deleted or develop into Tregs

in WT mice, the repertoire shift hypothesis postulates that ZAP-70 hypomorphic mutations

would lead to the appearance of TCR repertoires in Tconvs that would be typically found in

WT Tregs, or normally deleted[14,15]. A natural way of testing this hypothesis would be to

compare the similarity of the repertoires of WT Tregs and SKG or ZAC Tconvs. If ZAP-70

mutant Tconvs were more similar to WT Tregs than to WT Tconvs, it would support the

repertoire shift hypothesis. To address this question, we turned to previously published TCR

reference datasets[38-41] in order to consolidate WT Tconv and WT Treg reference data

(Supplementary Table 14). For the calculation, we treated each of our experimentally

determined (WT Tconv, WT Treg, SKG Tconv, SKG Treg, ZAC Tconv, or ZAC Treg) TCRs

as a "query" that we used to search a reference WT Tconv or WT Treg dataset (Logunova[38],

Lu[39], Ko[40], Wolf[41]). For each of these queries, we computed the number of reference WT

Tconv and WT Treg CDR1, CDR2 and CDR3 sequence matches (methods) and, based on

the number of matches along with the total size of each reference dataset, deemed a given

query as "WT Treg-like", "WT Tconv-like" or "not significant" using a hypergeometric p-

value cutoff of 0.001. We then quantified the total number of matches of each type in terms

of the query repertoire size. We then expressed these values relative to the WT Tconv query

group (Fig. 4a). Considering first the largest (Logunova[38]) reference dataset, we found that

the fraction of WT Treg-like TCRs within SKG and ZAC Tconv queries was approximately

twice as high as the fraction in WT Tconvs, and even higher than the fraction in query WT

Tregs (Fig. 4b). In addition, the fraction of WT Tconv-like TCRs was notably reduced within

ZAP-70 mutant Tconv queries (Fig. 4b). Similar trends were observed in the other reference

datasets (Supplementary Fig. 8a). Furthermore, the ZAC pathogenic Th17 repertoire showed

an increased fraction of WT Treg-like TCRs and consistently displayed the smallest fraction

of WT Tconv-like TCRs (Fig. 4c, Supplementary Fig. 8b), indicating that the TCRs of

pathogenic Th17 cells were largely absent from WT Tconv repertoires, but similar to WT

Treg repertoires. Of note, pathogenic Th17 cells from ZAC mice did not show any significant

WT Treg or WT Tconv matches within the Ko et al., 2020[40] dataset (Supplementary Fig. 8b).

These observations support the repertoire shift hypothesis: ZAP-70 mutant Tconv repertoires

are more similar to WT Treg repertoires than to WT Tconv repertoires, and this trend is most

pronounced in the pathogenic Th17 cells.

Because the query-reference matching procedure described above requires a number of ad

hoc normalization steps, we next sought a simpler procedure that could be used as an

independent check. To this end, we took advantage of a machine learning tool that enables

the quantification of the regulatory potential of a TCR (TCR-intrinsic regulatory potential,

TiRP) based on TCR beta chain features[42]. As expected, our experimentally determined Treg

datasets exhibited higher TiRP scores than WT Tconv cells, and their respective Tconv

compartments (Supplementary Fig. 8c-d). Importantly, the ZAP-70 mutant Tconvs also

exhibited significantly higher TiRP scores than WT Tconvs, and higher than WT Tregs (Fig.

4d-e). Furthermore, when considering the ZAC Th17 phenotypes, we found that pathogenic

Th17 TiRP scores were significantly higher compared to those of WT Tconvs (Fig. 4f-g).

The higher TiRP scores in self-reactive T cells[42] suggest that the TCRs in Tconv cells from

ZAP-70 mutant mice, particularly those in pathogenic Th17 cells, are biased towards self-

reactivity. These results agree with those of the query-reference matching procedure, and,

again, support the repertoire shift hypothesis as an explanation for how ZAP-70 mutation-

mediated TCR signal weakening leads to the emergence of self-reactive Th17 cells that

develop a pathogenic phenotype and accumulate in joints upon activation.

Evidence of missing TCR specificities in ZAP-70 mutant WT Treg repertoires

A second consequence of the repertoire shift hypothesis is that a weakening of TCR signaling

impacts the Treg compartment by generating "holes" in the normal Treg repertoire. It has

been reported that missing specificities in the Treg repertoire have a large impact on the

suppressive function against self-reactive T cells[8,43]. This raises the possibility that an

absence of Treg specificities within ZAP-70 mutant mice repertoires contributes to a loss in

tolerance toward specific self-antigens. To explore this hypothesis, we analyzed the Treg-

Tconv repertoire similarity within each mouse and compared it across groups (WT, SKG,

ZAC). The Morisita-Horn index for exact CDR3 overlap showed a notably reduced Treg-

Tconv similarity in ZAC mice relative to WT or SKG, especially at the single chain level

(Fig. 5a). The Treg-Tconv overlap was generally distributed across effector phenotypes

(Tconv: intermediate, Th17 and Tfh. Treg: intermediate and effector) in non-arthritic mice

(Supplementary Fig. 9a-h), suggesting a suppressive potential for shared Tregs towards

activated Tconvs; however, such enrichment was not observed in ZAC mice. To further test

these observations, we again employed TCRdist[33] and calculated repertoire-repertoire

correlations (methods) for pooled WT, SKG and ZAC repertoires: here, a pair of similar

repertoires exhibit high correlation coefficients. We found that WT Treg and WT Tconv

repertoires were well correlated, whereas SKG and ZAC Tconv repertoires exhibited lower

correlation with their Treg repertoires. Indeed, the ZAC Treg repertoire was poorly correlated

with other groups in general, indicating a reduced similarity to any other tested repertoires

(Fig. 5b). Together, these results are consistent with a scenario wherein ZAP-70 mutant were

unable to select certain Treg TCR specificities, normally present in WT mice and capable of

suppressing autoimmune arthritis; which, along with a more self-reactive Th17 population,

contributes to a loss of tolerance and the development of autoimmune arthritis.

Discussion

In this study, we provide a comprehensive characterization of the Treg and Tconv phenotypes

and repertoires of WT, non-arthritic SKG and arthritic ZAC mice at the single cell level. We

identified two transcriptionally distinct populations of Th17 cells within tissues of arthritic

ZAC mice: migratory Th17 with high expression of *Il22*, and pathogenic Th17 cells, highly

expressing *Tnfsf11*. Pathogenic Th17 cells have a unique transcriptional signature, and are

almost exclusively detected in joints. In addition, pathogenic cells may target a restricted set

of antigens, as they harbor unique and convergent TCRs with a greater similarity to the TCRs

of WT Treg cells, which are generally understood to be biased toward self-reactivity[9-12].

Investigations into the contribution of high-affinity versus lower-affinity T cell specificities

to autoimmune disease establishment and progression, as observed in experimental

autoimmune encephalomyelitis (EAE), have revealed a critical role of high-affinity

autoreactive cells in initiating disease that can be maintained by lower-affinity cells once

established[44]. In our study, it is possible that the pathogenic Th17 population comprises cells

critical for the establishment of disease, given their unique repertoire and tissue location;

while migratory Th17 cells may contribute to disease progression and maintenance and

represent a prior differentiation state for cells that later acquire a pathogenic Th17 phenotype.

Such dynamics would be in line with increased diversity and reduced clonal dominance in

established disease stages[45,46].

Consistently, our trajectory prediction suggests that pathogenic Th17 cells represent a

terminal differentiation state, preceded by the migratory Th17 phenotype. This observation

also aligns with previous research showing that in EAE Th17 cells progress from a self-

renewing cellular state primarily enriched in lymph nodes, to an effector pre-Th1-like

phenotype found in both lymph nodes and the central nervous system (CNS) and

characterized by the expression of chemokine receptors, to a Th17/Th1-like effector

phenotype that is concentrated in the CNS and exhibits a pathogenic signature with high

expression of *Tnfsf11*[47].

In addition to these observations on the pathogenic phenotype of ZAC Th17 cells, we carried

out the first systematic investigation at single cell resolution of the repertoire shift of

conventional T cells upon TCR signal weakening. Our results are consistent with the

observation that T cells infiltrating prostatic autoimmune lesions in Aire$^{-/-}$ mice express

receptors preferentially found in Tregs from Aire$^{+/+}$ mice[48]. In addition, it was shown that

exFoxP3 cells display a higher osteoclastogenic potential as compared to Th17 derived from

naïve CD4$^+$ T cells, presumably due to their higher affinity to self-antigens[26]. Together with

our findings, these reports point to a key role of TCRs normally occurring in the Treg

compartment in the recognition of self-antigens, and suggests that aberrant expression of

such TCRs in Tconv cells may drive autoimmune disease. Finally, the unique identification

of CD4$^+$ T cells expressing the pathogenic Th17 gene signature in inflamed synovia from RA

patients highlights their potential involvement in autoimmune arthritic inflammation,

motivating further identification of their tissue-specific self-antigens.

Remarkably, we observed a number of public clones within ZAC Tconv cells, particularly

concentrated within both Th17 phenotypes, and notably expanded. The significance of public

clones, in the development of autoimmune conditions, has been previously documented[49,50].

Among the CDR3 sequences of public clones, we identified only one exact match to a TCR

in the VDJdb database[51] of TCR sequences with known antigen specificities (Supplementary

Table 9). Unexpectedly, the match corresponded to a CD8 TCR, and its target peptide was

presented in a class I MHC. Although the presence of "class-mismatched" T cells has been

reported in the literature[52-54], it remains unknown whether this phenomenon occurs in this or

other self-reactive clones in ZAC mice. Consequently, the identity of any potential epitopes

or amino acid motifs remains an open problem.

As migratory and pathogenic Th17 cells were predicted to be related both by trajectory

analysis and by the existence of a limited number of shared clones, we speculate that the

pathogenic Th17 phenotype is only acquired within joints and only by cells bearing self-

reactive TCRs. The data supports a model wherein these cells are critical in triggering and

establishing disease in target tissues, whereas migratory Th17 cells play a supporting

inflammatory role, but are not strongly self-reactive.

Finally, the repertoire shift hypothesis raises the prospect of loss of certain TCR specificities

within the Treg compartment. Considering that antigen recognition by the Treg TCR appears

to be less degenerate than in Tconv cells[8,55,56], the loss of certain Treg specificities could

profoundly impact tolerance towards specific self-antigens[43]. While it is generally

acknowledged that Treg and Tconv repertoires are distinct, a small proportion of their

repertoires typically overlap[46,57,58]. In this study, we observed a reduction in the similarity

between ZAC Treg and Tconv repertoires, specially within activated Tconv cell groups,

suggesting a divergence in the Treg-Tconv repertoires that could result in the loss of critical

Treg function. This result is consistent with the observation of a significantly reduced

proportion of potentially self-reactive receptors in the Treg repertoires of autoimmune Aire[-]

[/-] mice[48]. Therefore, it is possible to speculate that moderate TCR weakening by ZAP-70

mutation may not only shift the Tconv repertoire but also deplete the Treg repertoire of

relevant TCRs. The functional identity of such critical TCRs remains unknown, and thus

worthy of future study as a starting point for development of TCR-transgenic Treg

immunotherapy for the treatment of autoimmune diseases[59].

Taken together, the consequences of weakened TCR signaling in the development of

autoimmune disease are multilayered. Firstly, it allows cells with self-reactive TCRs to evade

negative selection or deflection into the Treg repertoire by differentiation into self-reactive

Tconvs. At the same time, Tregs are affected in at least two ways: on the one hand, the Treg

repertoire may become deficient in some normally-occurring and potentially critical TCRs,

resulting in not a shift, but also a bifurcation of the repertoire. On the other hand, Treg effector

activity upon TCR weakening is significantly disrupted[15], rendering Tregs insufficient for

the suppression of self-reactive Tconvs (Fig. 6).

It should be noted that our study has several limitations, such as the limited sample size and

repertoire coverage that can be practically analyzed by current single cell sequencing

approaches. In addition, the pathogenic potential of Th17 cells, their self-reactivity, as well

as their potential to suppress Treg function have not been further experimentally assessed.

Moreover, the antigen specificity of public clones could only be predicted, which yielded

only a single match to a functionally annotated TCR. This TCR was described as being

specific to a viral epitope presented via MHC class I. Future *in vitro* and *in vivo* experiments

will be required to elucidate the specific role and antigen specificities of Th17 cell

populations in disease establishment and progression. In spite of these limitations, our

research expands the cellular and molecular level understanding of the breakdown in

tolerance during autoimmune disease and proposes a straightforward path to discovering

specific TCR-antigen interactions involved in RA.

Methods

Mice

Animal experiments were approved by the institutional review board of Osaka University,

and animals were treated in accordance with the institutional guidelines of the Immunology

Frontier Research Center, Osaka University. BALB/c and SKG mice were acquired from

CLEA Japan. SKG[13], ZAC[15] and FIG (Foxp3-IRES-GFP knock-in[60]) mice were previously

described. In this study, FIG BALB/c WT, FIG SKG and FIG ZAC mice were used and

generated as previously reported [15]. All animals were maintained in Specific pathogen-free

(SPF) facilities.

Droplet-based single cell RNA sequencing.

466 CD4$^+$Foxp3$^-$ or Foxp3$^+$ cells from FIG BALB/c WT, FIG SKG or FIG ZAC mice were sorted

467 by FACSAria SORP (BD Biosciences). For FIG BALB/c WT and FIG SKG mice, cells were

468 obtained from spleens, processed with the Chromium Single Cell 5' Library & Gel bead kit

469 (10x Genomics), loaded onto a Chromium Single Cell G Chip, and encapsulated in a

470 Chromium single cell controller (10x Genomics) to generate single-cell gel beads in the

471 emulsion (GEMs) following manufacturer's protocol. After encapsulation, reverse

472 transcription and cDNA amplification were performed, followed by target enrichment of

473 V(D)J segments and the independent construction of V(D)J and Gene expression libraries as

474 recommended by the manufacturer's instructions.

475 For FIG ZAC mice, cells from inflamed joint, draining popliteal lymph node (dLN) and

476 spleen from each replicate were barcoded with TotalSeq hashtag antibodies (BioLegend) and

477 then pooled by tissue and cell type (i.e., Joint Tconv, Joint Treg, etc.) before the generation

478 of GEMs. Next, RNA and V(D)J libraries were produced as described above and an

479 additional library for the barcoded antibodies was prepared. HiSeq3000 and NovaSeq6000

480 were used for sequencing.

482 **Single-cell RNAseq data processing**

484 Raw reads were aligned to the mouse reference genome (GRCm38 and GRCm38 VDJ v5.0.0,

485 from 10× Genomics), filtered, and demultiplexed to create gene expression matrices by Cell

Ranger (v.5.0). The resulting data was analyzed in R software (v 4.1.2) with Seurat v4.0

package[61,62].

Where necessary, hashtag demultiplexing was performed by the *HTODemux* function from

Seurat and only data from cells classified as singlets was kept for further steps. Next, low-

quality data was removed from all datasets, this means cells with unique feature counts less

than 200 or over 2500, and cells in which mitochondrial reads represented >5% of reads.

Only cells annotated to a single alpha and a single beta chain in the V(D)J result were retained

for further analysis. Normalization was performed by the "LogNormalize" method from

Seurat and the top 2,000 variable genes were identified in each dataset independently.

Batch correction and data integration were performed in combinations of data as follows: 1)

Tconvs from ZAC tissues; 2) Tregs from ZAC tissues; 3) Tconvs from spleen (including WT,

SKG and ZAC samples); 4) Tregs from spleen (including WT, SKG and ZAC samples).

Batch correction was performed with the Canek R package[63] using the intersecting variable

genes for the datasets to be integrated in each run.

After integration and QC, the final datasets for analysis consisted of: 6,509 cells for ZAC

Tconv, 5,306 cells for ZAC Treg, 24,743 cells for Spleen Tconv and 25,227 cells for

Spleen Treg (Supplementary Table 1). Uniform manifold approximation and projection

(UMAP) was used to visualize gene expression profiles and clusters from the principal

components (PCs) representation of the integrated datasets; the number of PCs were

independently selected on each dataset as the elbow suggested in variance plot of PCs.

Public single-cell RNAseq data was obtained as count matrices and pre-processed

accordingly for QC, normalization, batch correction and data integration (when required).

Single cell gene expression analysis

Clustering and cell-type annotation

The identification of cell clusters based on gene expression data was performed in the PCs representation of integrated datasets with the *FindClusters* function from Seurat using a resolution parameter of 0.5. Clusters were annotated by the examination of expression of canonical Treg or Tconv markers together with the inspection of markers defining each cluster and enriched GO Biological Process Terms (Supplementary Notes 1-4). Additional exclusion of artifactual populations was performed after clustering, including cells lacking expression of CD3 (*Cd3d*, *Cd3e*, and *Cd3g)*, invariant NKT (iNKT) cells, and clusters with high expression of Hemoglobin genes (*Hba-a1, Hba-a2, Hbb-bt, Hbb-bs*).

DEG and gene set enrichment analysis

Identification of cluster marker genes was performed with *FindAllMarkers* function in Seurat. Significant markers were considered with a fold change (FC) ≥1.2 and adjusted p value ≤0.05, and were used for gene set enrichment analysis.

When the comparison of two populations was required, DEGs were identified with the *FindMarkers* function with the min.pct parameter set at 0.25. Genes with adjusted p value ≤0.05 where considered significantly changed and were classified as upregulated if FC ≥1.5 or downregulated if FC ≤0.67.

Gene set enrichment analysis was performed with enrichR (v3.1) package[64] using the

GO_Biological_Process_2021 database as reference.

For the generation of ZAC migratory and pathogenic Th17 gene signatures for module score

calculation, significant markers were considered if fold change (FC) $\geq$2 and adjusted p value

$\leq$0.05.

Trajectory analysis

Trajectory analyses were performed by using the Slingshot Bioconductor package (v2.10.0)[65]

with the PCs representation of integrated datasets and their respective cluster labels as inputs.

We set the parameter *start.clus='naive'* to use the cluster of Naive cells as the starting point

of the inferred trajectories.

Module score calculation

Calculation of expression levels from Th17 ZAC programs in spleen and public human

scRNAseq datasets was performed by the *AddModuleScore* function from Seurat using the

specified gene lists or their corresponding human orthologs (Supplementary Table 6) and

default parameters.

For public scRNAseq datasets, clustering was performed as described above. Where it was

possible to annotate Th17 cells, these were separated in an independent Seurat object and

processed for module score calculation and visualization. If Th17 cells were not clearly

identified, the whole CD4$^+$ T cell clusters were separated in independent Seurat objects for

module score calculation.

Cell-cell crosstalk prediction

We used Cellphone DB[37] to predict cell-cell communication. After converting mouse gene

IDs to human orthologues, we ran CellPhoneDB v4.0.0 in statistical analysis mode, for the

subsets of effector Treg together with Intermediate or Th17 Tconvs. Results were visualized

by the ktplots R package. We manually searched for all significant interactions in each mouse

model. Only statistically significant interactions with a reasonable biological meaning are

shown in detail in the results.

Single TCR V(D)J sequencing analysis

Clonal expansion and repertoire overlap

Clonal expansion was assessed by the calculation of clonal proportions, then, clones were

classified by their size as: small (clonal proportion $<1 \times 10^{-3}$), medium (clonal proportion from

$1 \times 10^{-3} - 1 \times 10^{-2}$) and large (clonal proportion $>1 \times 10^{-2}$). Given that only a small sample size

could be recovered for some replicates, which inflates clonal proportions of lowly abundant

clones, we labeled singletons (clone size=1) separately.

The Morisita-Horn index is a typical metric used in ecology to calculate the similarity of two

ecosystems in terms of species distribution and abundance, and it has been introduced as a

metric for repertoire overlap analysis; its values go from 0 to1, where 1 represents a perfect

overlap[32]. The Morisita-Horn index was calculated between repertoires of each individual

replicate, at the paired or single CDR3 amino acid level by the Immunarch (v0.9.0) R package.

The median Morisita-Horn indices were compared by a Wilcoxon test using a significance

level of 0.05. In addition, clonotype tracking across Treg and Tconv phenotypes was

performed by Immunarch.

Clonal publicity

Public clones were defined as TCRs sharing paired gene usage and CDR3 amino acid

sequences and occurring in more than one subject.

TCR distances and repertoire-repertoire correlation

In order to evaluate if the overall features of selected repertoires show convergence, we

calculated receptor distances with TCRdist. TCRdist allows for the calculation of a distance

measure between TCRs by comparing the CDR1, CDR2 and CDR3 amino acid sequences of

single or paired TCR chains[33], then, a neighbor distance metric consisting of the weighted

average of the distance of a given TCR to the nearest 25th percentile of the other TCRs in the

repertoire can be calculated (weighted neighbor distance). Then, distance distributions of

selected repertoires were compared and distributions biased toward lower distances were

interpreted as an indication of repertoire convergence. In order to restrict the convergence

analysis to the qualitative composition of the repertoires, we used unique clone sequences

i.e., clonal counts where not considered. Median distances of Naïve, migratory and

pathogenic Th17 cells were compared by a Wilcoxon test with a significance level of 0.05.

Finally, the epitope-epitope (or repertoire-repertoire) neighbor distance score correlation

metric from the TCRdist algorithm was computed for assessment of intra Treg-Tconv

repertoire similitude. Briefly, a score is assigned to each TCR in one repertoire which is the

average distance to the nearest TCRs in a reference repertoire. Then, the sets of scores

obtained by two different repertoires can be compared over the entire merged set of TCRs

and the Pearson correlation score is calculated. In principle, two similar repertoires will be

well correlated.

TiRP score calculation

Lagattuta et al proposed a Treg-propensity scoring system for the TCR to quantify the

likelihood of Treg cells based on the TCR feature on Treg fate (TiRP score)[42]. The source

code was obtained from https://github.com/immunogenomics/TiRP. The TiRP score of this

study was calculated using the default parameter. For each TCR sequence, the v-gene and

amino acid of CDR3 were utilized as input information. Median TiRP scores were compared

across groups by a Wilcoxon test with a significance level of 0.05.

Published sequence preparation

The published mouse Treg and Tconv TCR beta chain sequence data were obtained from

four studies[38-41]. The detailed accession ID and references were summarized in

Supplementary Table 14. The annotation of raw sequences was performed by mixcr

(3.0.13)[66] with align, assemble, assembleContigs, and exportClones process. The sequences

that have complete sequences of frameworks and CDRs regions were utilized in this study.

Sequence searching and comparison

We implemented the TCR sequence searching using Interclone method[67]. The source code

was deposited in https://gitlab.com/sysimm/interclone and the searching parameter using the

default setting. In brief, InterClone regards the CDR1,2,3 of one TRB seq as a pseudo

sequence and considers one pair of hitting using the following CDR similarity and coverage

thresholds: $CDR1 \geq 90\%$, $CDR2 \geq 90\%$, $CDR3 \geq 80\%$, and $Coverage \geq 90\%$. We first

searched Tconv and Treg TCR templates from the same reference using identical query

inputs. Then we compare the corresponding hitting number with Tconv and Treg templates

for each query and calculate the significant Tconv or Treg-biased query sequence using a

hypergeometric p-value cutoff of 0.001. We then quantified the total number of matches in

terms of the query repertoire size and normalized the results to the matches in the query WT

Tconvs (Fig. 4a).

Statistical analysis

For identification of cluster markers, significant markers were considered with a $FC \geq 1.2$ and

adjusted p value ≤ 0.05. For DEG analysis between two cell populations, genes with adjusted

p value ≤ 0.05 where considered significantly changed and were classified as upregulated if

$FC \geq 1.5$ or downregulated if $FC \leq 0.67$. The median Morisita-Horn indices, median distances

of Naïve, migratory and pathogenic Th17 repertoires and median TiRP scores were compared

by a Wilcoxon test using a significance level of 0.05. For TCR sequence searching and

comparison, we calculated the significant Tconv or Treg-biased query sequence using a

hypergeometric p-value cutoff of 0.001.

Data availability

The datasets generated during the current study have been deposited in the DDBJ Sequence

Read Archive database (accession numbers DRA010311, DRA010223), and NCBI Gene

Expression Omnibus (accession number GSE180432).

References

Richards, D. M., Kyewski, B. & Feuerer, M. Re-examining the Nature and Function of Self-Reactive T cells. *Trends Immunol* **37**, 114-125 (2016). https://doi.org:10.1016/j.it.2015.12.005

Goodnow, C. C., Sprent, J., Fazekas de St Groth, B. & Vinuesa, C. G. Cellular and genetic mechanisms of self tolerance and autoimmunity. *Nature* **435**, 590-597 (2005). https://doi.org:10.1038/nature03724

Kronenberg, M. & Rudensky, A. Regulation of immunity by self-reactive T cells. *Nature* **435**, 598-604 (2005). https://doi.org:10.1038/nature03725

Moon, J. J. *et al.* Quantitative impact of thymic selection on Foxp3+ and Foxp3- subsets of self-peptide/MHC class II-specific CD4+ T cells. *Proc Natl Acad Sci U S A* **108**, 14602-14607 (2011). https://doi.org:10.1073/pnas.1109806108

Cebula, A. *et al.* Dormant pathogenic CD4(+) T cells are prevalent in the peripheral repertoire of healthy mice. *Nat Commun* **10**, 4882 (2019). https://doi.org:10.1038/s41467-019-12820-3

Levine, A. G., Arvey, A., Jin, W. & Rudensky, A. Y. Continuous requirement for the TCR in regulatory T cell function. *Nat Immunol* **15**, 1070-1078 (2014). https://doi.org:10.1038/ni.3004

Li, M. O. & Rudensky, A. Y. T cell receptor signalling in the control of regulatory T cell differentiation and function. *Nat Rev Immunol* **16**, 220-233 (2016). https://doi.org:10.1038/nri.2016.26

Pohar, J., Simon, Q. & Fillatreau, S. Antigen-Specificity in the Thymic Development and Peripheral Activity of CD4(+)FOXP3(+) T Regulatory Cells. *Front Immunol* **9**, 1701 (2018). https://doi.org:10.3389/fimmu.2018.01701

Jordan, M. S. *et al.* Thymic selection of CD4+CD25+ regulatory T cells induced by an agonist self-peptide. *Nat Immunol* **2**, 301-306 (2001). https://doi.org:10.1038/86302

Hsieh, C. S., Zheng, Y., Liang, Y., Fontenot, J. D. & Rudensky, A. Y. An intersection between the self-reactive regulatory and nonregulatory T cell receptor repertoires. *Nat Immunol* **7**, 401-410 (2006). https://doi.org:10.1038/ni1318

11   Aschenbrenner, K. *et al.* Selection of Foxp3+ regulatory T cells specific for self
antigen expressed and presented by Aire+ medullary thymic epithelial cells. *Nat*
Immunol **8**, 351-358 (2007). https://doi.org:10.1038/ni1444

12   Lee, H. M., Bautista, J. L., Scott-Browne, J., Mohan, J. F. & Hsieh, C. S. A broad
range of self-reactivity drives thymic regulatory T cell selection to limit responses to
self. *Immunity* **37**, 475-486 (2012). https://doi.org:10.1016/j.immuni.2012.07.009

13   Sakaguchi, N. *et al.* Altered thymic T-cell selection due to a mutation of the ZAP-70
gene causes autoimmune arthritis in mice. *Nature* **426**, 454-460 (2003).
https://doi.org:10.1038/nature02119

14   Tanaka, S. *et al.* Graded attenuation of TCR signaling elicits distinct autoimmune
diseases by altering thymic T cell selection and regulatory T cell function. *J Immunol*
185, 2295-2305 (2010). https://doi.org:10.4049/jimmunol.1000848

15   Tanaka, A. *et al.* Construction of a T cell receptor signaling range for spontaneous
development of autoimmune disease. *J Exp Med* **220** (2023).
https://doi.org:10.1084/jem.20220386

16   Brand, D. D., Latham, K. A. & Rosloniec, E. F. Collagen-induced arthritis. *Nat*
Protoc **2**, 1269-1275 (2007). https://doi.org:10.1038/nprot.2007.173

17   Komatsu, N. & Takayanagi, H. Mechanisms of joint destruction in rheumatoid
arthritis - immune cell-fibroblast-bone interactions. *Nat Rev Rheumatol* **18**, 415-429
(2022). https://doi.org:10.1038/s41584-022-00793-5

18   Meyer Zu Horste, G. *et al.* RBPJ Controls Development of Pathogenic Th17 Cells by
Regulating IL-23 Receptor Expression. *Cell Rep* **16**, 392-404 (2016).
https://doi.org:10.1016/j.celrep.2016.05.088

19   Ohl, K. *et al.* The transcription factor CREM drives an inflammatory phenotype of T
cells in oligoarticular juvenile idiopathic arthritis. *Pediatr Rheumatol Online J* **16**, 39
(2018). https://doi.org:10.1186/s12969-018-0253-x

20   Glal, D. *et al.* ATF3 Sustains IL-22-Induced STAT3 Phosphorylation to Maintain
Mucosal Immunity Through Inhibiting Phosphatases. *Front Immunol* **9**, 2522 (2018).
https://doi.org:10.3389/fimmu.2018.02522

21   Lin, C. C. *et al.* IL-1-induced Bhlhe40 identifies pathogenic T helper cells in a model
of autoimmune neuroinflammation. *J Exp Med* **213**, 251-271 (2016).
https://doi.org:10.1084/jem.20150568

22   DiToro, D. *et al.* Insulin-Like Growth Factors Are Key Regulators of T Helper 17
Regulatory T Cell Balance in Autoimmunity. *Immunity* **52**, 650-667 e610 (2020).
https://doi.org:10.1016/j.immuni.2020.03.013

23   Jha, P. & Das, H. KLF2 in Regulation of NF-kappaB-Mediated Immune Cell
Function and Inflammation. *Int J Mol Sci* **18** (2017).
https://doi.org:10.3390/ijms18112383

24   Lee, J. Y. *et al.* The transcription factor KLF2 restrains CD4(+) T follicular helper
cell differentiation. *Immunity* **42**, 252-264 (2015).
https://doi.org:10.1016/j.immuni.2015.01.013

25   Basdeo, S. A. *et al.* Ex-Th17 (Nonclassical Th1) Cells Are Functionally Distinct from
Classical Th1 and Th17 Cells and Are Not Constrained by Regulatory T Cells. *J*
Immunol **198**, 2249-2259 (2017). https://doi.org:10.4049/jimmunol.1600737

26 Komatsu, N. *et al.* Pathogenic conversion of Foxp3+ T cells into TH17 cells in
autoimmune arthritis. *Nat Med* **20**, 62-68 (2014). https://doi.org:10.1038/nm.3432

27 Etori, K. *et al.* Fibroblast growth factor receptor 1 as a potential marker of terminal
effector peripheral T helper cells in rheumatoid arthritis patients. *Rheumatology*
(Oxford) **62**, 3763-3769 (2023). https://doi.org:10.1093/rheumatology/kead220

28 Zhang, B. *et al.* CD127 imprints functional heterogeneity to diversify monocyte
responses in inflammatory diseases. *J Exp Med* **219** (2022).
https://doi.org:10.1084/jem.20211191

29 Tang, S. *et al.* Single-cell atlas of human infrapatellar fat pad and synovium
implicates APOE signaling in osteoarthritis pathology. *Sci Transl Med* **16**, eadf4590
(2024). https://doi.org:10.1126/scitranslmed.adf4590

30 Yasumizu, Y. *et al.* Single-cell transcriptome landscape of circulating CD4(+) T cell
populations in autoimmune diseases. *Cell Genom* **4**, 100473 (2024).
https://doi.org:10.1016/j.xgen.2023.100473

31 Tong, L. *et al.* Current understanding of osteoarthritis pathogenesis and relevant new
approaches. *Bone Res* **10**, 60 (2022). https://doi.org:10.1038/s41413-022-00226-9

32 Chiffelle, J. *et al.* T-cell repertoire analysis and metrics of diversity and clonality.
Curr Opin Biotechnol **65**, 284-295 (2020).
https://doi.org:10.1016/j.copbio.2020.07.010

33 Dash, P. *et al.* Quantifiable predictive features define epitope-specific T cell receptor
repertoires. *Nature* **547**, 89-93 (2017). https://doi.org:10.1038/nature22383

34 Valencia, X. *et al.* TNF downmodulates the function of human CD4+CD25hi T-
regulatory cells. *Blood* **108**, 253-261 (2006). https://doi.org:10.1182/blood-2005-11-
4567

35 Dikiy, S. & Rudensky, A. Y. Principles of regulatory T cell function. *Immunity* **56**,
240-255 (2023). https://doi.org:10.1016/j.immuni.2023.01.004

36 Liu, Z. *et al.* Immune homeostasis enforced by co-localized effector and regulatory T
cells. *Nature* **528**, 225-230 (2015). https://doi.org:10.1038/nature16169

37 Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal-fetal interface
in humans. *Nature* **563**, 347-353 (2018). https://doi.org:10.1038/s41586-018-0698-6

38 Logunova, N. N. *et al.* MHC-II alleles shape the CDR3 repertoires of conventional
and regulatory naive CD4(+) T cells. *Proc Natl Acad Sci U S A* **117**, 13659-13669
(2020). https://doi.org:10.1073/pnas.2003170117

39 Lu, D. R. *et al.* Dynamic changes in the regulatory T-cell heterogeneity and function
by murine IL-2 mutein. *Life Sci Alliance* **3** (2020).
https://doi.org:10.26508/lsa.201900520

40 Ko, A. *et al.* TCR Repertoires of Thymic Conventional and Regulatory T Cells:
Identification and Characterization of Both Unique and Shared TCR Sequences. *J*
Immunol **204**, 858-867 (2020). https://doi.org:10.4049/jimmunol.1901006

41 Wolf, K. J., Emerson, R. O., Pingel, J., Buller, R. M. & DiPaolo, R. J. Conventional
and Regulatory CD4+ T Cells That Share Identical TCRs Are Derived from Common
Clones. *PLoS One* **11**, e0153705 (2016).
https://doi.org:10.1371/journal.pone.0153705

42   Lagattuta, K. A. *et al.* Repertoire analyses reveal T cell antigen receptor sequence
features that influence T cell fate. *Nat Immunol* **23**, 446-457 (2022).
https://doi.org:10.1038/s41590-022-01129-x
43   Ooi, J. D. *et al.* Dominant protection from HLA-linked autoimmunity by antigen-
specific   regulatory   T   cells.   *Nature*   **545**,   243-247   (2017).
https://doi.org:10.1038/nature22329
44   Yi, J. *et al.* Antigen-specific depletion of CD4(+) T cells by CAR T cells reveals
distinct roles of higher- and lower-affinity TCRs during autoimmunity. *Sci Immunol*
7, eabo0777 (2022). https://doi.org:10.1126/sciimmunol.abo0777
45   Klarenbeek, P. L. *et al.* Inflamed target tissue provides a specific niche for highly
expanded T-cell clones in early human autoimmune disease. *Ann Rheum Dis* **71**,
1088-1093 (2012). https://doi.org:10.1136/annrheumdis-2011-200612
46   Al Khabouri, S. *et al.* TCRbeta Sequencing Reveals Spatial and Temporal Evolution
of Clonal CD4 T Cell Responses in a Breach of Tolerance Model of Inflammatory
Arthritis.   *Front   Immunol*   **12**,   669856   (2021).
https://doi.org:10.3389/fimmu.2021.669856
47   Gaublomme, J. T. *et al.* Single-Cell Genomics Unveils Critical Regulators of Th17
Cell   Pathogenicity.   *Cell*   **163**,   1400-1412   (2015).
https://doi.org:10.1016/j.cell.2015.11.009
48   Malchow, S. *et al.* Aire Enforces Immune Tolerance by Directing Autoreactive T
Cells into the Regulatory T Cell Lineage. *Immunity* **44**, 1102-1113 (2016).
https://doi.org:10.1016/j.immuni.2016.02.009
49   Menezes, J. S. *et al.* A public T cell clonotype within a heterogeneous autoreactive
repertoire is dominant in driving EAE. *The Journal of Clinical Investigation* **117**,
2176-2185 (2007). https://doi.org:10.1172/JCI28277
50   Zhao, Y. *et al.* Autoimmune susceptibility imposed by public TCRbeta chains. *Sci
Rep* **6**, 37543 (2016). https://doi.org:10.1038/srep37543
51   Shugay, M. *et al.* VDJdb: a curated database of T-cell receptor sequences with known
antigen   specificity.   *Nucleic   Acids   Res*   **46**,   D419-D427   (2018).
https://doi.org:10.1093/nar/gkx760
52   Boyle, L. H., Goodall, J. C. & Gaston, J. S. Major histocompatibility complex class
I-restricted   alloreactive   CD4+   T   cells.   *Immunology*   **112**,   54-63   (2004).
https://doi.org:10.1111/j.1365-2567.2004.01857.x
53   Boyle, L. H., Goodall, J. C., Opat, S. S. & Gaston, J. S. The recognition of HLA-B27
by   human   CD4(+)   T   lymphocytes.   *J   Immunol*   **167**,   2619-2624   (2001).
https://doi.org:10.4049/jimmunol.167.5.2619
54   Singh, N. K. *et al.* A class-mismatched TCR bypasses MHC restriction via an
unorthodox but fully functional binding geometry. *Nat Commun* **13**, 7189 (2022).
https://doi.org:10.1038/s41467-022-34896-0
55   Kieback, E. *et al.* Thymus-Derived Regulatory T Cells Are Positively Selected on
Natural Self-Antigen through Cognate Interactions of High Functional Avidity.
Immunity **44**, 1114-1126 (2016). https://doi.org:10.1016/j.immuni.2016.04.018

56 Leonard, J. D. *et al.* Identification of Natural Regulatory T Cell Epitopes Reveals
Convergence on a Dominant Autoantigen. *Immunity* **47**, 107-117 e108 (2017).
https://doi.org/10.1016/j.immuni.2017.06.015
57 Hsieh, C. S. *et al.* Recognition of the peripheral self by naturally arising CD25+ CD4+
T   cell   receptors.   *Immunity*   **21**,   267-277   (2004).
https://doi.org/10.1016/j.immuni.2004.07.009
58 Pacholczyk, R., Ignatowicz, H., Kraj, P. & Ignatowicz, L. Origin and T cell receptor
diversity of Foxp3+CD4+CD25+ T cells. *Immunity* **25**, 249-259 (2006).
https://doi.org/10.1016/j.immuni.2006.05.016
59 Eggenhuizen, P. J. *et al.* Smith-specific regulatory T cells halt the progression of
lupus nephritis. *Nat Commun* **15**, 899 (2024). https://doi.org/10.1038/s41467-024-
45056-x
60 Ohkura, N. *et al.* T cell receptor stimulation-induced epigenetic changes and Foxp3
expression are independent and complementary events required for Treg cell
development.   *Immunity*   **37**,   785-799   (2012).
https://doi.org/10.1016/j.immuni.2012.09.010
61 Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction
of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).
https://doi.org/10.1038/nbt.3192
62 Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and
doublet detection for single cell genomics. *Genome Biol* **19**, 224 (2018).
https://doi.org/10.1186/s13059-018-1603-1
63 Loza, M., Teraguchi, S., Standley, D. M. & Diez, D. Unbiased integration of single
cell transcriptome replicates. *NAR Genom Bioinform* **4**, lqac022 (2022).
https://doi.org/10.1093/nargab/lqac022
64 Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. *BMC Bioinformatics* **14**, 128 (2013). https://doi.org/10.1186/1471-
2105-14-128
65 Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018). https://doi.org/10.1186/s12864-018-
4772-0
66 Bolotin, D. A. *et al.* MiXCR: software for comprehensive adaptive immunity
profiling. *Nat Methods* **12**, 380-381 (2015). https://doi.org/10.1038/nmeth.3364
67 Wilamowski, J. *et al.* InterClone: Store, Search and Cluster Adaptive Immune
Receptor   Repertoires.   *bioRxiv*,   2022.2007.2031.501809   (2022).
https://doi.org/10.1101/2022.07.31.501809
840

Acknowledgements

This work was funded by the Japan Agency for Medical Research and Development (AMED),

grant number JP223fa627002 and Platform Project for Supporting Drug Discovery and Life

Science Research (Basis for Supporting Innovative Drug Discovery and Life Science

Research) under JP21am0101108; and by JSPS KAKENHI Grant Numbers JA23H034980

(to D.S.), 20K16286 (to M.L.C.), and 26860331, 17K15723, 22H02920 (to A.T.).

848
Author contributions

D.S. and M.L.C. conceived the study. M.L.C., S.T. and A.T. designed experiments. M.L.C.,

E.L., and A.T performed experiments. M.L.C., M.L.L., D.D., Z.X., performed data analysis.

M.L., A.T. and D.S. interpreted data and wrote the paper. S.T. and Z.X. designed custom

software for TCRs analysis. D.M. and S.S. supervised the project. All the authors read and

approved the final manuscript.

Competing interests

A.T. reported grants from Shionogi & Co., Ltd. outside the submitted work. No other

disclosures were reported.

859
860   **Correspondence** and requests for materials should be addressed to D.M. or M.L.C

Figure 1

Figure 1. Two transcriptionally different Th17 cell populations exist in inflamed joints of arthritic ZAC mice. a) Lateral view of a hindpaw of a 23-week old arthritic ZAC mouse. **b)** UMAP plot of Tconv cells obtained from joints, draining lymph node (dLN) and spleen of arthritic ZAC mice colored by cluster. Unsupervised clustering was applied by the Louvain algorithm and clusters were annotated as described in the text. **c)** Proportion of annotated Tconv cell populations in joints, dLN and spleen of arthritic ZAC mice. **d)** Relative expression of canonical Tconv markers across gene expression clusters. Circle color and intensity represent the average expression whereas circle size represents the percentage of cells from each cluster expressing the gene. **e)** Volcano plot of the differentially expressed genes (DEG) among cluster 1 (migratory Th17) and cluster 4 (pathogenic Th17). **f)** Top 5 gene Ontology (GO) Terms enriched in DEGs from cluster 1 (migratory Th17) vs cluster 4 (pathogenic Th17). The color intensity of bars represents the odds ratio (OR) of enrichment. **g)** UMAP plot of Tconv cells from tissues of arthritic ZAC mice, cells are colored according to their expression of *Il17a*, *Il17f*, *Il22* or *Tnfsf11*. **h)** UMAP plot of Tconv cells from tissues of arthritic ZAC mice, with cells colored according to their inferred pseudotime values; cluster centers are connected according to trajectory prediction by Slingshot's pipeline.

Figure 2

Figure 2. The pathogenic Th17 gene signature is uniquely detected in CD4+ T cells from inflamed synovia of rheumatoid arthritis (RA) patients. Gene signatures of migratory Th17 and pathogenic Th17 cells were obtained and used for calculation of module scores in public scRNAseq datasets of **a)** Th17 cells obtained from spleens of WT, SKG and ZAC mice (n=2 each), **c)** CD4+ T cells from inflamed synovia of RA patients (n=3)[27], **d)** CD4+ T cells from peripheral blood of RA patients (n=1)[28], **e)** CD4+ T cells from synovia and infrapatellar fat pad (IPFP) tissue of osteoarthritis (OA) patients (n=3)[29], **f)** Circulating Th17 cells from healthy donors (n=3), and patients with autoimmune diseases (AD) other than RA: myasthenia gravis (MG) (n=3), multiple sclerosis (MS) (n=4), systemic lupus erythematosus (SLE) (n=3)[30]; **b)** Violin plot of module scores for the gene signatures of migratory Th17 (left) and pathogenic Th17 (right) cells in Th17 cells from spleens of WT, SKG and arthritic ZAC mice.

Figure 3

Figure 3. The TCR repertoire of pathogenic Th17 cells is unique and convergent, and contain public clones. a) UMAP plot of Tconv cells from tissues of arthritic ZAC mice, cells are colored according to clonal proportion classification. **b)** Distribution of clonal proportion classes across ZAC Tconv gene expression clusters. **c)** Morisita-Horn index of the paired CDR3 amino acid similitude between the migratory and pathogenic Th17 repertoires (gray bar); and between both Th17 populations vs the other Tconv phenotypes in ZAC tissues (brown and green bars). **d)** Distribution of neighbor distances to the nearest 25th percentile (methods) in the repertoires of Naïve, migratory Th17 and pathogenic Th17 cells. Paired (left) and single alpha (middle) and beta (right) TCR chain distributions are shown. **e)** Public clones and their clonal proportion classification and distribution across Tconv phenotypes in each mouse replicate. *: p<0.05, **p<0.01; Wilcoxon Test.

Figure 4

Figure 4. TCR repertoires of ZAP-70 mutant Tconv cells, especially of pathogenic Th17 cells, are more WT Treg-like than WT Tconv-like. a) Description of the method employed for the assessment of repertoire similitude to reference WT Treg and WT Tconv repertoires. Each of our experimentally determined query TCRs were used to search CDR matches (methods) in a reference WT Tconv or WT Treg dataset[38-41]. We next computed the number of reference WT Tconv and reference WT Treg matches and deemed the query TCR "WT Treg-like", "WT Tconv-like" or "not significant" using a hypergeometric p-value cutoff of 0.001. The total number of matches was calculated in terms of the query repertoire size and results were normalized to the matches in the query WT Tconvs. **(b-c)** Normalized fraction of query WT Tconv and WT Treg as compared to SKG and ZAC Tconv repertoires **(b)** or ZAC migratory and pathogenic Th17 repertoires **(c)** matching with reference WT Tconv (magenta section, negative values) or WT Treg repertoires (blue section, positive values) obtained from a representative reference dataset[38]. **d)** Distribution of TCR-intrinsic regulatory potential (TiRP) scores for Tconv repertoires of WT, SKG and ZAC mice and the WT Treg repertoire. **e)** Median TiRP score change of WT Treg, and SKG and ZAC Tconv repertoires relative to the WT Tconv repertoire. **f)** Distribution of TiRP scores for WT Tconv, and overall Tconv, migratory Th17, and pathogenicTh17 repertoires of ZAC mice. **g)** Median TiRP score change of overall Tconv, migratory Th17, and pathogenicTh17 repertoires of ZAC mice relative to WT Tconv. **p<0.01, ****: p<0.0001; Wilcoxon Test

Figure 5

Figure 5. Intra-Treg-Tconv similitude in WT, SKG and ZAC repertoires. a) Morisita-Horn index calculated at the exact CDR3 amino acid level between the Tconv and Treg compartments of WT, SKG and ZAC mice. Left panel shows the analysis for single alpha chains, middle panel shows single beta chains and right panel depicts paired chains. **b)** Heatmap of weighted neighbor distance correlations between WT, SKG and ZAC Treg and Tconv repertoires as calculated by TCRdist (methods), higher correlation scores indicate closer repertoires.

Figure 6

Figure 6. Consequences of weakened TCR signaling in the shaping of Treg and Tconv repertoires and autoimmunity. The SKG (W163C) and ZAC (H165A) mutations in the ZAP-70 gene cause a reduction in its affinity for phosphorylated CD3z within a range that results in the development of autoimmune disease. The consequences of a weakened TCR signaling in the development of autoimmune disease are multilayered. On one hand, cells with self-reactive TCRs are aberrantly selected into the Tconv compartment instead of the Treg compartment (repertoire shift). This defective selection also deprives the Treg repertoires of specific receptors potentially critical for the proper suppression of autoimmune responses and this is evidenced by a reduced similarity between Treg and Tconv repertoires within an individual (repertoire bifurcation). In addition, Treg effector activity upon TCR weakening is significantly disrupted, rendering Tregs insufficient for the suppression of self-reactive Tconvs. These events together, favor the activation of self-reactive T cells, their differentiation into pathogenic phenotypes and their accumulation in target tissues. Created with BioRender.com