A Targeted Genome-scale Overexpression Platform for Proteobacteria

Authors

Amy B. Banta[1,2], Kevin S. Myers[2,3], Ryan D. Ward[1,2,4], Rodrigo A. Cuellar[5], Michael Place[2],

Claire C. Freeh[1], Emily E. Bacon[1,5], and Jason M. Peters[1,2,6,7,8]✉

Affiliations

[1]Pharmaceutical Sciences Division, School of Pharmacy, University of Wisconsin-Madison,

Madison, WI, USA. [2]DOE Great Lakes Bioenergy Research Center, University of Wisconsin-

Madison, Madison, WI, USA. [3]Wisconsin Energy Institute, University of Wisconsin-Madison,

Madison, WI, USA. [4]Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI,

USA. [5]Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI,

USA. [6]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA.

[7]Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,

Madison, WI, USA. [8]Center for Genomic Science Innovation, University of Wisconsin-Madison,

Madison, WI, USA. ✉e-mail: jason.peters@wisc.edu

Abstract

Targeted, genome-scale gene perturbation screens using Clustered Regularly Interspaced Short
Palindromic Repeats interference (CRISPRi) and activation (CRISPRa) have revolutionized
eukaryotic genetics, advancing medical, industrial, and basic research. Although CRISPRi
knockdowns have been broadly applied in bacteria, options for genome-scale overexpression
face key limitations. Here, we develop a facile approach for genome-scale gene overexpression
in bacteria we call, "CRISPRtOE" (CRISPR transposition and OverExpression). We create a
platform for comprehensive gene targeting using CRISPR-associated transposition (CAST) and
show that transposition occurs at a higher frequency in non-transcribed DNA. We then
demonstrate that CRISPRtOE can upregulate gene expression in Proteobacteria with medical
and industrial relevance by integrating synthetic promoters of varying strength upstream of
target genes. Finally, we employ CRISPRtOE screening at the genome-scale in Escherichia
coli, recovering known antibiotic targets and genes with unexplored roles in antibiotic function.
We envision that CRISPRtOE will be a valuable overexpression tool for antibiotic mode of
action, industrial strain optimization, and gene function discovery in bacteria.

Introduction

Overexpression (OE) studies have been invaluable for gene phenotyping in medical[1–3], industrial[4], and basic contexts[3,5]. When used in functional genomics, OE enables functional interrogation of genes that are normally cryptic under screening conditions[3], mimics natural resistance mechanisms[3], and complements loss of function approaches by identifying genes with oppositely signed phenotypes when knocked down versus overexpressed[1,6]. Clustered Regularly Interspaced Short Palindromic Repeats activation (CRISPRa) has allowed routine, genome-scale overexpression screening in eukaryotes for drug synergy/antagonism[6] or sensitivity/resistance to toxic plant compounds that reduce bioproduct yields[7]. Such screens have been highly effective at identifying direct targets and pathways that mitigate or potentiate drug function. For example, a recent screen in mammalian cells for targets of the anti-cancer drug, rigosertib, leveraged genome-scale CRISPR interference (CRISPRi) knockdowns with CRISPRa to identify target pathways that showed oppositely signed phenotypes[1]. OE has also enhanced strain optimization for industrial purposes, such as a genome-wide CRISPRa screen in *Saccharomyces cerevisiae* for genes involved in resistance to the biofuel production stressor, furfural[7].

Existing genome-scale OE approaches in bacteria face key limitations. Classic approaches, such as random integration of promoter containing transposons to overexpress downstream genes[8], are untargeted and often require hyper-saturated transposon (Tn) libraries to resolve ambiguities related to the effects of Tn insertion versus overexpression. Open Reading Frame (ORF) libraries cloned into plasmids[9] are technically targeted approaches, but are tedious and expensive to construct, limiting their utility in non-model bacteria. Further, the use of plasmids presents additional complications, such as the constant requirement for antibiotic selection for maintenance (complicating antibiotic function screens), and copy number variation that increases experimental noise[10]. Contemporary approaches have sought to mitigate these issues and make genome-scale overexpression more accessible. For instance, trackable multiplex recombineering (TRMR) is a targeted approach that uses recombinase-aided DNA insertion (recombineering) to deliver barcoded promoters or ribosome binding site variants to systematically alter gene expression[11,12]; however, this method is restricted to bacteria with efficient recombineering systems. An innovative new OE technique, "Dub-seq," uses random genomic fragments cloned into multi-copy plasmids that can be quantified by barcode sequencing[13,14], reducing upfront barriers to library design, but complicating downstream analysis and sharing some downsides with other plasmid-based approaches. Moreover, the length of cloned genomic fragments dictates Dub-seq library screening results;

smaller fragments may miss genes that only have phenotypes when co-expressed in operons

(e.g., protein complex subunits) while longer fragments make it more difficult to pinpoint

individual genes responsible for the selected phenotype.

CRISPRa in eukaryotes uses a catalytically inactive Cas9 protein (dCas9) and single

guide RNAs (sgRNAs) to deliver activator proteins to promoters; these activators open

chromatin at the promoter, enhancing transcription activation[15]. In contrast, CRISPRa-based OE

in bacteria requires direct interactions between activator proteins and RNA polymerase (RNAP)

that dictate stringent activator-RNAP spacing and orientation on DNA[16–19]. Recent

improvements to bacterial CRISPRa using dCas9 variants with reduced PAM stringency and

engineered activator proteins partially mitigated the spacing requirements and increased overall

activation[18,19]; however, even in these optimized systems, less than half of targeted promoters

could be upregulated[19]. Accordingly, to our knowledge, no genome-scale bacterial CRISPRa

screens have been reported.

Naturally occurring CRISPR-associated transposon (CAST) systems[20] have been

engineered to generate targeted insertions in bacterial genomes with high efficiency[21–23].

Although several CAST systems have been described[24–26], the Tn*6677* system from *Vibrio*

cholerae (VcCAST) is among the best characterized and most precise[27–30]. In VcCAST, a

nuclease-deficient Type I-F CRISPR system is paired with a Tn*7*-like transposase to achieve

guide RNA-directed transposition. Binding of a VcCAST-gRNA complex spacer to a matching

protospacer results in transposition of DNA flanked by Tn*6677* ends ~49 bp downstream of the

protospacer. Transposon DNA can be inserted in two orientations with respect to the Tn*6677*

ends (R-L or L-R) but is highly biased toward the R-L orientation[21,31]. VcCAST is highly precise,

with nearly all tested guides showing >99% on-target insertion. The utility of CASTs has been

demonstrated for high efficiency editing in Proteobacteria[21], systematically disrupting

transcription factor genes in *Pseudomonas aeruginosa*[23], and even targeting individual strains in

a complex microbiome[22], but existing systems remain sub-optimal for genome-scale functional

genomics. Moreover, the use of Tn*6677* to deliver regulatory elements—such as promoters—

has only recently been explored in a model bacterium[32].

Here we modify VcCAST for genome-scale functional genomics, creating a targeted

overexpression system for bacteria we call, "CRISPRtOE" (CRISPR transposition and

OverExpression). CRISPRtOE delivers a transposon with an outward facing promoter upstream

of genes to facilitate overexpression (Fig. 1a). We demonstrate the utility of CRISPRtOE in

medically and industrially-relevant Proteobacterial species and at the genome-scale in

Escherichia coli to characterize antibiotic action and resistance.

Results

A VcCAST Platform for Genome-scale Functional Genomics

We first generated "CRISPRt," a VcCAST system designed for genome-scale functional genomics. Like previous VcCAST systems[21,22], CRISPRt enables precise insertion of transposons into genomic targets using programmable gRNAs (Fig. 1b). Superimposed on this are additional properties necessary for facile use of CRISPRt screening at the genome-scale: 1) Tn insertion is selectable, resulting in all transconjugants containing an integrated Tn; 2) *cas* helper and Tn plasmids are non-replicative in recipient bacteria, obviating the need for plasmid curing[23] prior to phenotyping; 3) *cas* and Tn components are on separate plasmids, preventing insertion of *cas* genes into the genome in rare events when VcCAST co-integrates the plasmid backbone alongside the Tn (occurring at ~0.01-0.001% frequency) and reducing Tn plasmid size which facilitates conjugation and cargo modification by standard cloning procedures (Fig. 1a, 1b, 1c, and Supplementary Fig. S1a, S1b, and S1c). CRISPRt can be transferred from donor *Escherichia coli* K-12 cells to recipient Proteobacteria (and possibly other phyla) *via* conjugation, and transconjugants containing Tn insertions are recovered at efficiencies consistent with genome-scale library creation.

To test the precision of CRISPRt, we targeted both endogenous (*lacZ*) and heterologous (*gfp*) reporter genes for disruption in *E. coli* K-12 (Supplementary Fig. S1d). Consistent with previous data on VcCAST systems, we found that the vast majority of Tn insertions were on-target. We first assessed targeting of *gfp* and *lacZ* using phenotypic assays (fluorescence measurements for *gfp* and blue-white screening for *lacZ*), finding that 97.6% and 100% of transconjugants showed the expected disruption phenotype, respectively (Supplementary Fig. S2a and S2b). Next, we measured targeting on the genome scale by sequencing genomic DNA at the Tn*6677* transposon junction (Tn-seq) using Illumina-based Next Generation Sequencing (NGS). Tn-seq showed >99.5% on target efficacy for both *gfp* and *lacZ* guide RNAs (Fig. 1d). As expected, Tn insertions largely occurred in a tight window centered at ~49 bp downstream of the protospacer that favored an R-L orientation (89.2% and 74.1% R-L for *gfp* and *lacZ*, respectively, (Fig. 1d and 1e)). Moreover, CRISPRt transconjugants were recovered at an efficiency compatible with genome-scale screening (~10^{-2} to 10^{-3} in *E. coli*, (Supplementary Fig. S2c)). We conclude that CRISPRt has excellent properties for functional genomics while retaining the on-target efficacy of VcCAST.

CRISPRt Efficacy is Higher on Non-transcribed DNA

Although Protospacer Adjacent Motif (PAM) requirements for VcCAST have been well characterized (a "CN" PAM is sufficient[25]), other genomic features that impact guide efficacy are poorly understood. To address this issue, we created a pooled CRISPRt library targeting both strands of an *mScarlet-I* reporter gene that was integrated into the genome of *E. coli* K-12 (46 gRNAs total (Fig. 2a)). Surprisingly, we found that guide efficacy depended on which strand was targeted. To measure relative guide efficacy, we compared the abundance of gRNA spacers in our library before and after transposition into *mScarlet-I* (i.e., in mating donor strains versus recipient transconjugants (Fig. 2b)). Consistent with previous data from smaller sets of spacers[33], we found considerable variability, with ~27-fold difference in efficacy between the most and least active guides. We noticed a clear, but not absolute bias toward higher efficacy in template strand targeting guides (i.e., gRNAs that bind the transcribed strand) versus those that targeted the non-template strand. Other variables, such as PAM identity or distance from the gene start did not show an obvious trend (Supplementary Fig. S3a and S3b).

The bias in strand specificity suggested that either replication or transcription could be impacting CRISPRt efficacy. To test for a replication effect, we flipped the orientation of *mScarlet-I* relative to the origin or replication (Supplementary Fig. S3c). However, we continued to observe strand specificity in the flipped target, seemingly ruling out a role for replication direction in CRISPRt efficacy (Supplementary Fig. S3d and S4a). To test for a transcription effect, we targeted a promoterless version of *mScarlet-I* with our CRISPRt library (Supplementary Fig. S3c). In contrast to the transcribed target, the promoterless *mScarlet-I* gene showed reduced variation in guide efficacy, consistent with transcription affecting CRISPRt function (Supplementary Fig. S3c and S4a). Because our pooled approach did not allow for an absolute measure of CRISPRt activity, we cloned the three most active guides from our *mScarlet-I* library (T03, T09, and T21) and individually tested their activities (Fig. 2c and Supplementary Fig. S4b). We found that transcription reduced recovery of CRISPRt transconjugants (transposition efficiency) by 6.9 to 19.6-fold, depending on the guide tested. Although Tn disruption of transcribed genes is effective, we conclude that CRISPRt efficacy increases on non-transcribed targets.

Targeted and Tunable Overexpression in Proteobacteria

We considered that CRISPRt could be modified into an overexpression system by inserting outward-facing promoters into the Tn*6677* transposon that would cause increased transcription

of downstream genes (CRISPRtOE (Fig. 3a)). We inserted synthetic, constitutive promoters of

varying strength that had been shown to function in various Proteobacteria[34,35] into Tn*6677* with

the goal of creating an OE gradient (Supplementary Fig. S5a and S5b). To quantify the OE

activity of CRISPRtOE, we generated a "test" strain with a promoterless *mScarlet-I* reporter

gene integrated into the *E. coli* K-12 genome. This cassette contained a well-characterized

protospacer upstream of *mScarlet-I* to act as a "landing pad" (LP) for CRISPRtOE transposons

(Fig. 3a and Supplementary Fig. S5c). The test strain showed negligible mScarlet-I expression

in the absence of an upstream CRISPRtOE insertion (Fig. 3b).

CRISPRtOE achieved a gradient of mScarlet-I overexpression across our series of

tested promoters (Fig. 3b). CRISPRtOE containing the strong P_H promoter showed a ~177-fold

increase in mScarlet-I expression, while weaker promoters showed intermediate increases.

CRISPRtOE lacking a promoter (no P) failed to stimulate transcription above background,

although we note that insertions in the reverse orientation (L-R) were excluded from this

analysis due to cryptic promoter activity (Supplementary Fig. S6a and Table S8). We also

observed small variations in overexpression depending on the precise location of insertion,

possibly due to changes in mRNA folding (Supplementary Table S8).

We next sought to demonstrate CRISPRtOE activity in Proteobacteria with medical and

industrial relevance. The Alphaproteobacterium, *Zymomonas mobilis*, is a promising biofuel

producer and emerging model for bacteria with naturally reduced genomes[36]. To assay

CRISPRtOE function in *Z. mobilis*, we again generated an *mScarlet-I* test strain and measured

expression following insertion of CRISPRtOE transposons (Supplementary Fig. S6b and Table

S8). We found that CRISPRtOE could produce a gradient of overexpression in *Z. mobilis* up to

~50-fold. Differences in the fold effects of CRISPRtOE across species likely reflect variation in

promoter activity among our constitutive promoter set.

We then extended this approach to a panel of eight total species including the Gram-

negative ESKAPE pathogens[37] *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *P. aeruginosa*,

and *Enterobacter cloacae*, as well as the industrially and environmentally relevant strains

Pseudomonas putida[38] and *Shewanella oneidensis*[39], focusing our strongest promoter (P_H) to

reduce experimental complexity (Fig. 3c and Supplementary Table S8). We found that

CRISPRtOE could upregulate *mScarlet-I* expression in all tested species with an

overexpression range of 232-fold (*E. cloacae*) to 23-fold (*P. aeruginosa*), likely due to variation

in promoter activity across species. The CRISPRtOE insertion position in most species was

centered at 33 bp upstream of *mScarlet-I*, although *A. baumannii* showed a wider range of

positions for unknown reasons (Supplementary Fig. S6c). We note that the modular design of

CRISPRtOE enables facile swapping of promoter sequences with organism-specific or inducible
promoters to customize expression levels depending on the application. We conclude that
CRISPRtOE is an effective overexpression strategy for Proteobacteria.
204

Pooled CRISPRtOE Recapitulates Known Antibiotic Targets

There is an urgent need to develop new therapeutics to stem the rising tide of antibiotic
resistance[40]. One major bottleneck in the process of translating new antimicrobials to the clinic
is determining the mode of action (MOA), including identifying the direct target. We previously
showed that CRISPRi could be used to determine the direct targets of uncharacterized
antibiotics by observing synergy between knockdown strains and sub-lethal concentrations of
antibiotic[41,42]. However, CRISPRtOE could provide an even more straightforward avenue to
antibiotic target discovery because strains with overexpressed target proteins would outgrow
competitor strains in a pooled context. For instance, it has been shown that OE of *folA*,
encoding the trimethoprim (TMP) target, Dihydrofolate reductase (DHFR), can substantially
increase resistance to TMP[43]. Likewise, *murA*, encoding the fosfomycin (FOS) target, UDP-NAG
enolpyruvyl transferase (MurA), can provide FOS resistance when overexpressed[44].
217   To validate CRISPRtOE as a tool for functional interrogation of endogenous genes and
antibiotic target discovery, we overexpressed and phenotyped the *folA* and *murA* genes in *E.*
coli K-12. As a hedge against variation in integration efficiency and overexpression efficacy, we
tiled the region upstream of *folA* and *murA* with 12 unique spacers per gene and created a
negative control library that inserted into the *lacZ* coding region (Fig. 4a). To test for
overexpression phenotypes, we mixed our *folA* or *murA* library 50:50 with the *lacZ* control library
and grew cells in the presence/absence of TMP or FOS, respectively. Strikingly, we found that
all CRISPRtOE gRNA spacers targeting *folA* or *murA* were effective in increasing competitive
fitness in the context of TMP or FOS, respectively (Fig. 4b and Supplementary Fig. S7a). This
competitive advantage led to a 22-fold increase in the abundance of *folA* OE strains and a 52-
fold increase in *murA* OE strains, relative to *lacZ* controls. To further characterize the impact of
CRISPRtOE on TMP and FOS resistance, we performed Minimal Inhibitory Concentration (MIC)
strip assays on individual insertion isolates (Supplementary Fig. S7b, S7c, and S7d).
CRISPRtOE strains showed dramatically increased resistance, in some cases with an MIC
change that approached the maximum value of the scale. Because FOS showed a high
frequency of spontaneous resistant suppressors in MIC strip assays, we focused on TMP in
subsequent broth microdilution MIC measurements including the wild-type parent strain, a *lacZ*
negative control, and *folA* CRISPRtOE isolates with insertions at various positions upstream of

folA (due to different spacer usage (Fig. 4c)). We observed substantial shifts in MIC (~15 to 30-

fold) between the parent strains and *folA* CRISPRtOE isolates, demonstrating that CRISPRtOE

phenotypes were robust to insertion distance upstream of the target gene. We conclude that

CRISPRtOE reliably upregulates endogenous target genes and can recapitulate antibiotic

targets.

Genome-scale CRISPRtOE Reveals Antibiotic Targets and Resistance Pathways

Buoyed by our success with *folA* and *murA*, we expanded the scope of our pooled CRISPRtOE

screening to the genome-scale. We targeted all protein coding genes in the *E. coli* K-12 genome

with ~10 gRNAs designed to integrate CRISPRtOE transposons within a small window

upstream of genes without disrupting downstream coding sequences (45,767 guides total, (Fig.

5)). Critically, we generated two pooled libraries, with and without promoter H, that allowed us to

distinguish between effects caused by promoter activity (intended overexpression) versus

upstream gene disruption (unintended loss of function). We found that CRISPRtOE effectively

targeted regions upstream of genes across the genome, without obvious hot spots or depleted

regions (Fig. 5a, see "CRISPRtOE No Promoter"). To characterize the insertion of CRISPRtOE

transposons relative to genes, we calculated the distance between transposon ends and gene

starts, finding a tight distribution of insertions largely within 100 bp of the 5′ ends of genes and

centered at ~20 bp upstream (Fig. 5b). This tight CRISPRtOE distribution showed a stark

contrast with the nearly flat distribution of Tn*5* insertions, which are known to occur pseudo-

randomly[45]. For instance, the distinction between CRISPRtOE targeted and Tn*5* pseudo-random

insertion was clearly visible at the *lac* locus (Fig. 5c).

To test the ability of CRISPRtOE to identify antibiotic-relevant gene phenotypes at the

genome-scale, we grew our libraries in the presence or absence of a sub-lethal concentration of

TMP and measured relative strain abundance. Biological replicates showed excellent

agreement at the gene level (R^2 values ranged from 0.95 to 0.97, Supplementary Fig. S8),

underscoring the high reproducibility of genome-scale CRISPRtOE. We found *folA*-associated

spacers as clear positive outliers (>1000-fold enrichment), demonstrating the exquisite

sensitivity and specificity of our CRISPRtOE screen in recapitulating antibiotic targets (Fig. 5a,

5d, and 5e). Substantial *folA* enrichment was only seen in the P_H-containing library not in the

promoterless library, indicating the CRISPRtOE promoter activity, rather than insertion per se,

was responsible for the phenotype (Fig. 5e). Promoter-dependent phenotypes were distinct

across the entire genome, as relative strain abundances of the P_H and no promoter libraries

after TMP treatment were not correlated (R^2 = 0.037, Supplementary Fig. S8). Importantly, we

were able to improve data quality by computationally eliminating aberrantly inserted

transposons based on distance between the intended target (defined by the spacer sequence)

and the insertion position/orientation (defined by Tn-seq). This mostly impacted our "no

promoter" data as cryptic transcription from transposons that inserted in the L-R orientation also

resulted in overexpression phenotypes (Supplementary Fig. S6). Our libraries contained many

more guides than was necessary to uncover phenotypes, as gene-level phenotypes were robust

when the number of spacers per gene was reduced from ~10 to just 3 (R^2 = 0.93,

Supplementary Fig. S9) and considering only guides that targeted the first gene of operons

returned many of the same top hits (Supplementary Fig. S10); these findings suggest that

CRISPRtOE library complexity can be reduced considerably with similar results. Thus, we

anticipate that future CRISPRtOE libraries will achieve genome-scale overexpression with far

fewer guides per gene.

282   Other significant outliers revealed potential TMP resistance or susceptibility mechanisms

(Fig. 5e and Supplementary Fig. S10). One of the most resistant outlier genes was *soxS*,

encoding an AraC-type transcription activator known to regulate genes involved in antibiotic

resistance[46] (Fig. 5e). Indeed, gene set enrichment analysis showed that several members of

the AraC and MarR families[47] were resistant outliers in our screen, including *marR*, *rob*, *soxR*,

and *marA* (FDR = 0.0055, afc test) (Fig. 5e and Supplementary Fig. S10), which notably

includes genes encoding the AcrA and AcrB components of the AcrAB-TolC multidrug efflux

pump[48]. Consistent with this, CRISPRtOE of the *acrA* gene also increased TMP resistance

(Supplementary Fig. S10). TMP treatment reduces cellular methionine levels due to the

involvement of tetrahydrofolate (the product of DHFR) in methionine biosynthesis[49].

CRISPRtOE of the *metR* gene, which encodes a positive regulator of methionine biosynthesis

genes[50], increased TMP resistance, possibly by increasing flux through the pathway. In contrast,

metF overexpression was highly deleterious in the presence of TMP (Supplementary Fig. S10),

likely due to depletion of 5,10-methylenetetrahydrofolate pools normally used by the ThyA-

DHFR pathway to generate tetrahydrofolate[51]. Neither *metR* nor *metF* were recovered in

previous TMP loss-of-function screens[52,53], demonstrating the utility of CRISPRtOE to identify

new players in antibiotic function. Interestingly, CRISPRtOE of *folM*, which is thought to encode

a protein with DHFR activity[54], was highly toxic to TMP-treated cells (Supplementary Fig. S10).

Although unintuitive, this result is consistent with other genome-scale phenotyping studies that

found that disruption of *folM* increases resistance to TMP[52,53]. Finally, genes involved in

biosynthesis of the Enteric Common Antigen (ECA), a key component of the enterobacterial

outer membrane, were functionally enriched among sensitive outliers (FDR = 5.58e-05, afc test)

(Supplementary Fig. S10). Previous work has shown that disruption of the ECA pathway also

increases TMP sensitivity, possibly by sequestering thymidine pools and leading to "hyper-

acting" version of thymine-less death[55]. Our work suggests that disregulation by overexpression,

in addition to disruption, can cause ECA-dependent toxicity in TMP-treated cells. Although ECA

genes often exist in complex operons that may be disrupted by insertion, phenotypes for some

ECA genes (e.g., *rffM*) were strongly dependent on the presence of P_H (Supplementary Fig.

S10). Future studies will be needed to define the precise mechanisms of TMP resistance and

sensitivity observed here. Taken together, genome-scale CRISPRtOE is a facile approach for

comprehensive genetic analysis of antibiotic function.

Discussion

Functional genomics approaches that are robustly scalable and readily applicable to diverse

bacteria are required to bridge the yawning gap between genome sequencing and gene function

assessment. Our work provides a targeted, systematic, and practical approach for genome-

scale gene overexpression in Proteobacteria and possibly beyond. By modifying VcCAST for

use in functional genomics (Fig. 1) and defining non-transcribed regions of DNA as high-

efficiency insertion sites (Fig. 2), we demonstrated targeted and tunable OE in Proteobacteria

with medical, industrial, and basic research relevance (Fig. 3). Our genome-scale OE screen for

targets and modulators of TMP efficacy highlighted the exquisite specificity of CRISPRtOE in

recovering genes and pathways that underpin antibiotic function (Fig. 4 and 5). Given the ease

of implementing CRISPRtOE in the organisms explored here, we anticipate that our approach

will be readily expandable to other microbes and screening conditions. While our manuscript

was under review, another manuscript was published that showed that VcCAST containing

promoters could be used to upregulate genes in *E. coli*[32]. Our work dramatically expands this

approach across multiple Proteobacteria and to the genome scale. We note that the replicative

CAST vector used in that study is suboptimal for use at the genome-scale due to a lack of Tn

selection, the need to cure the plasmid post-editing, and an inability to replicate in important

genera such as *Acinetobacter*.

CRISPRtOE is a valuable genetic tool for both basic and applied biology. The targeted

aspect of CRISPRtOE can be used to generate focused libraries that limit size (number of

gRNAs) and scope (number of targeted genes). Moreover, screen hits can be readily pursued

for downstream mechanistic analysis by individually cloning gRNAs used in the screen to easily

recreate strains outside of the pooled context, a clear benefit over random transposition

approaches such as Tn5. Additionally, generating pooled sub-libraries from screen hits enables

follow up validation at scale. Importantly, the ability to insert transposons with or without

promoters at the same genomic locus will allow researchers to disentangle the effects of Tn

insertion versus overexpression, simplifying hit interpretation. Further, combining CRISPRtOE

with existing gene perturbation approaches such as CRISPRi could enable screening for

overexpression suppressors of essential functions at the genome-scale.

Our proof of principle work demonstrating that genome-scale CRISPRtOE can identify

resistance genes in *E. coli* K-12 holds promise for extending this approach to clinically relevant

pathogens with the goal of improving diagnosis and treatment. Importantly, CRISPRtOE action

is similar to clinically relevant antibiotic resistance mechanisms, such as IS element

transposition and overexpression of downstream resistance genes[56]. CRISPRtOE may also be

valuable for optimization of strains for industrial use. Industrial strain optimization often occurs

through loss of function[57] or directed evolution experiments[58], followed up by plasmid-based

overexpression of mutated pathways. CRISPRtOE streamlines this process in non-model

bacteria, particularly by enabling recreation of optimized strain features in different genetic

backgrounds by targeted transposition. Finally, CRISPRtOE-optimized strains may be

immediately useful in production, as CRISPRtOE as leaves no heterologously expressed *cas*

genes in the recipient strain. Such optimization experiments are currently underway.

CRISPRtOE is not without caveats and there may be scenarios in which it is

outperformed by other overexpression or gene manipulation strategies. CRISPRtOE and other

engineering approaches that insert promoters will alter expression of downstream genes in

operons, a phenomenon known as "polarity." Polar effects could complicate, but not necessarily

preclude identification of gene-level CRISPRtOE phenotypes. Our view is that expression of

downstream genes is often advantageous due to fact that protein complexes are often found in

operons (e.g., efflux pump subunits). We expect that "No promoter" CRISPRtOE libraries will

largely allow for disambiguation of insertion versus overexpression phenotypes, but there may

be rare instances in which the combination of insertion and overexpression causes an

unexpected genetic interaction because operons contain functionally related genes. In this

scenario, Dub-seq or promoter recombineering strategies would be advantageous, as Dub-seq

expresses genes *in trans* and recombineering can be used to introduce more subtle *cis*

mutations that could avoid perturbing upstream genes. Dub-seq can also introduce

heterologous DNA from other species for phenotyping[14], something CRISPRtOE is not designed

to accomplish.

Methods

Strains and growth conditions. Strains are listed in Table S1. Growth and antibiotic selection conditions are also summarized in Table S6. *Escherichia coli*, *Acinetobacter baumannii*, *Enterobacter cloacae*, *Klebsiella pneumoniae,* and *Pseudomonas aeruginosa* were grown in LB broth, Lennox (BD240230) at 37°C (or 30°C for CRISPR-guided transposition) in a flask with shaking at 250 rpm, in a culture tube on a roller drum, in a 96 well deepwell plate with shaking at 900 rpm, or in a Tecan Sunrise plate reader shaking with OD600 measurements every 15 min. *Pseudomonas putida* and *Shewanella oneidensis* were grown in LB in a culture tube on a roller drum or in a 96 well deepwell plate with shaking at 900 rpm at 30°C. *Zymomonas mobilis* was grown in RMG medium (10g yeast extract and 2g KH_2PO_4 monobasic/liter with 2% glucose added after autoclaving) at 30°C statically in a culture tube or deepwell plate. *E. coli* was grown in Mueller Hinton Broth (MHB, BD 275730) for antibiotic minimal inhibitory concentration (MIC) assays. Media was solidified with 1.5-2% agar for growth on plates. Antibiotics were added when necessary: *E. coli* (100 µg/ml carbenicillin (carb), 30 µg/ml kanamycin (kan), or 50 µg/ml spectinomycin (spt)), *A. baumannii* (60 µg/ml kan or 100 µg/ml apramycin (apr)), *E. cloacae* (30 µg/ml kan or 50 µg/ml spt), *K. pneumoniae* (60 µg/ml kan or 50 µg/ml spt), *P. aeruginosa* (30 µg/ml gentamicin (gen) or 1 mg/ml kan), *P. putida* (150 µg/ml apr or 60 µg/ml kan), *S. oneidensis* (60 µg/ml kan or 30 µg/ml gen), and *Z. mobilis* (200 µg/ml gen or 100 µg/ml spt). Diaminopimelic acid (DAP) was added at 300 µM to support growth of *dap- E. coli* strains. Strains were preserved in 15% glycerol at -80°C.

General molecular biology techniques. Plasmids are listed in Table S2. Oligonucleotides are listed in Table S3. *pir*-dependent plasmids were propagated in *E. coli* strain BW25141 (sJMP146) or its derivative sJMP3053. Plasmids were purified using the GeneJet Plasmid Miniprep kit (Thermo K0503), the QIAprep Spin Miniprep Kit (Qiagen 27106), or the Purelink HiPure Plasmid Midiprep kit (Invitrogen K210005). Plasmids were digested with restriction enzymes from New England Biolabs (NEB, Ipswich, MA). Ligations used T4 DNA ligase (NEB M0202) and fragment assembly used NEBuilder Hifi (NEB E2621). Genomic DNA was extracted with the DNeasy Blood & Tissue Kit (Qiagen 69504) or the GeneJet genomic DNA purification kit (Thermo K0721). DNA fragments were amplified by PCR using Q5 DNA polymerase (NEB M0491, for cloning) or One*Taq* DNA Polymerase (NEB M0480, for analysis). PCR products were spin-purified using the Monarch PCR & DNA Cleanup Kit (NEB T1030) or the DNA Clean

& Concentrator-5 kit (Zymo Research, Irvine, CA, D4013). Reactions were purified with 1.8X

Mag-Bind TotalPure NGS magnetic beads (Omega) on a magnetic rack (Alpaqua). DNA was

quantified spectrophometrically using a NanoDrop Lite or fluorometrically using a Qubit with the

HS or BR DNA kit (Thermo). Plasmids were transformed into electrocompetent *E. coli* cells

using a 0.1 cm cuvette (Fisher FB101) and a BioRad Gene Pulser Xcell on the Bacterial 1 *E.*

coli preset protocol (25 µF, 200 ohm, 1800 V) as described in detail previously[59].

Oligonucleotides were synthesized by Integrated DNA Technologies (Coralville, IA) or Agilent

(SurePrint Oligonucleotide library) (Santa Clara, CA). Sequencing was performed by Functional

Biosciences (Madison, WI), Plasmidsaurus (Eugene, OR), Azenta (South Plainfield, NJ), or the

University of Wisconsin-Madison Biotechnology Center Next Generation Sequencing Core

(UWBC NGS core, Madison, WI).

E. coli Δ*lac* strain construction. An *E. coli* MG1655 strain with a *lac* operon deletion (Δ*lac*,

Lac⁻) was constructed by λ-Red-mediated recombination as previously described[60]. Briefly, *E.*

coli MG1655 (sJMP163) harboring the plasmid pSIM6 (pJMP170) encoding the λ-Red proteins

was transformed with a linear DNA fragment encoding an FRT-*kan*-FRT cassette which was

amplified by PCR from plasmid pJMP3099 using primers oJMP912 and oJMP913 which also

have 40 nt homology to the chromosomal insertion site) resulting in strain sJMP3267. The allele

was transferred from sJMP3267 to *E. coli* MG1655 (sJMP163) by P1–*vir*-mediated transduction,

as previously described[61], resulting in strain sJMP3269. The *kan* gene was removed by

transformation with plasmid pCP20 (pJMP3008), encoding a constitutively expressed FLP

recombinase, as previously described[62], resulting in strain sJMP3272. Full deletion of the *lac*

operon in sJMP3272 was confirmed by PCR with flanking primers (oJMP914 and oJMP915)

followed by Sanger sequencing.

Construction of fluorescent reporters and insertion into the *att*$_{Tn7}$ site on the bacterial

chromosome. Fluorescent reporter cassettes are located in a Tn*7* transposon located on an

R6K ori (*pir* dependent) plasmid. The transposons were transferred to the *att*$_{Tn7}$ site ~50 bp

downstream of *glmS* of a recipient bacterial chromosome by co-conjugation of the Tn*7*

transposon donor strain and a second donor strain harboring a plasmid expressing Tn*7*

transposase[63]. Donor strains encode conjugation machinery, encode *pir*, and are DAP

dependent. Briefly, a donor strain with the transposon plasmid, a second donor with the

transposase expression plasmid, and the recipient strain were resuspended from plates to an

OD$_{600}$=9 and mixed in equal proportions (100 µl each). Cells were harvested by centrifugation at

5000 rcf for 2 min (except *A. baumannii* at 9000 rcf) and placed on a nitrocellulose filter

(Millipore HAWP01300) on an LB plate (except *Z. mobilis* on RMG) and incubated at 30°C (*Z.*

mobilis and *P. putida*) or 37°C (all other organisms) for 2-18h. Cells from the filter were

resuspended in 300 µl media, serially diluted, and plated with antibiotic selection for the

transposon (see growth and selection conditions above) and without DAP to select against the

donor strains, and incubated at 30 or 37°C until colonies formed.

A transposon containing a GFP expression cassette was derived from a Mobile-CRISPRi

plasmid[59] by assembling the AscI-EcoRI fragment of pJMP2824 and annealed oligonucleotides

oJMP1420 and oJMP1421 to create pJMP6957. The GFP encoding gene of pJMP6957 was

replaced by an mScarlet-I encoding gene derived from Addgene plasmid 85069 (pJMP7001) to

create pJMP10180 which has an *mScarlet-I* cassette (BbsI sites removed and *mScarlet*

converted to *mScarlet-I*) with the T7A1 constitutive promoter in the Tn*7* transposon. A

promoterless *mScarlet-I* reporter with the CRISPRt *LZ1* protospacer was derived from

pJMP6957, pJMP10180, and synthetic DNA oJMP1599 to create pJMP10206 (see Table S2 for

details).

Construction of CRISPRt/CRISPRtOE system. The CRISPRt/CRISPRtOE donor plasmids

were derived from pSpinR (pSL1765)[21], Mobile-CRISPRi plasmids[63], and synthetic DNA.

Cloning details are located in Supplementary Table S2 for the individual plasmids. In summary,

(1) a functional antibiotic resistance cassette was added to the transposon (pJMP747) and the

BsaI cloning site in the gRNA cassette was altered (pJMP10009 and pJMP10011). A two

plasmid R6K ori (*pir*-dependent) system for selectable CRISPR-guided transposition was

created by combining (1) the TnsABC-TniQ-Cas876 expression cassette from pSpinR with the

amp^R, R6K ori, mobilizable backbone from Mobile-CRISPRi (pJMP2782) to create pCRISPRt-H

(pJMP10233) and (2) the transposon and guide RNA cassette from pJMP10011 with the same

backbone (from pJMP2782) to create pCt-T (pJMP10237 and derivatives) which was further

minimized, optimized, and altered by adding transcription terminators and moving the guide

RNA cassette inside the transposon to produce pCRISPRt-T-gent (pJMP10395) and pCRISPRt-

T-kan (pJMP10397). CRISPRtOE plasmids with either no promoter or various strength

promoters (pCtOE-noP, pCtOE-P$_A$, pCtOE-P$_H$, pCtOE-P$_V$) were derived from these plasmids by

inserting a transcription terminator and promoter cassette derived from synthetic DNA

(oJMP2000 plus oJMP2001-2004) into the above plasmids to produce plasmids oJMP10471-

10479. Apramycin resistant derivatives of some of these plasmids were created by digesting

pJMP10397 and pJMP10478 with XhoI and assembling with a piece of synthetic DNA

(oJMP1946) encoding an apramycin resistance gene.

CRISPRtOE Guide Design. Guide RNAs were designed using custom Python scripts, available

with usage notes at GitHub repositories: https://github.com/ryandward/CRISPRt_gRNA and

https://github.com/ryandward/seq_hunter. Initially, the *Escherichia coli* K-12 RefSeq genome

assembly, in GenBank file format, was downloaded from the NCBI database (Accession

Number: GCF_000005845.2), using the seq_hunter.py script. The CRISPRt_gRNA.py script

was employed to identify all genomic locations with a suitable "CN" PAM sequence.

Subsequently, potential spacers of 32 nucleotides in length were evaluated using awk, and ten

were selected based on the following criteria: 1) Unique occurrence in the genome, 2)

occurrence on the same strand as the downstream gene, 3) at least 95 nucleotides upstream

from the gene annotation**.**

CRISPRt/CRISPRtOE individual guide and guide library construction. Guide sequences

were ligated into the BsaI-cloning site of the pCRISPRt-T/pCtOE plasmids. For cloning

individual spacer sequences, two 36-nucleotide (nt) oligonucleotides (oligos) were annealed so

that they encode the desired sequence flanked by sticky ends compatible to the vector BsaI site

whereas for cloning pooled libraries, DNA fragments encoding spacers were amplified from a

pool of oligos and digested with BsaI prior to ligation.

To prepare the pCRISPRt-T/pCtOE vectors for cloning, plasmid DNA was extracted from a 100

ml culture using a midiprep kit, 2-10 µg plasmid DNA was digested with BsaI-HF-v2 (NEB

R3733) in a 100 µl reaction for 4 h at 37°C and then spin-purified.

For cloning individual guides, pairs of oligonucleotides (2 µM each) were mixed in a 50 µl total

volume of 1X NEB CutSmart buffer, heated at 95°C, 5 min and then cooled to room temperature

~20 min to anneal, and then diluted 1:40 in dH$_2$O prior to ligation. Ligation reactions (10 µl)

contained 1X T4 DNA ligase buffer (NEB), additional DTT (10 mM final) and ATP (0.1 mM final),

and 0.5 µl T4 DNA ligase in addition to 50 ng BsaI-digested/spin-purified vector and 2 µl of 1:40

diluted annealed oligos. Ligations were incubated at 25°C for 2 h and ligase was heat

inactivated for 20 min at 65°C. 1-2 µl ligation was used for electroporation into 50 µl *E. coli*

strain BW25141 (sJMP146). Transformations were serially diluted and plated with selection on

carb to obtain isolated colonies. After confirmation of sequence, individual plasmids were

transformed into *E. coli* mating strain WM6026 (sJMP424 or sJMP3257) with selection on carb

and DAP.

For cloning pooled libraries, fragments were amplified from 90 nt pooled oligos (IDT oPools or

Agilent SurePrint pools) using the following conditions per 100 µl reaction (reaction size was

adjusted ~100-600 µl depending on size of library): 20 µl Q5 buffer, 3 µl GC enhancer, 2 µl

10mM each dNTPs, 5 µl each 10 µM forward and reverse primers, 2 µl 10 nM pooled oligo

library, 1 µl Q5 DNA polymerase, and 186 µl H_2O with the following thermocycling parameters:

98°C, 30s; 15-19 cycles of: 98°C, 15s; 56°C, 15s; 72°C, 15s; 72°C, 10 min; 10°C, hold (cycles

were adjusted to obtain sufficient PCR product without overamplifying/biasing the library). PCR

products (90 bp) were spin-purified and 300 ng was digested with BsaI-HF-v2 (NEB R3733) in a

100 µl reaction for 2 h at 37°C (no heat kill, reaction size was adjusted proportionally depending

on the size of the library). Size and digestion of PCR products were confirmed on a 4% agarose

E-Gel (Thermo). The BsaI-digested PCR product without further purification (3ng/µl) was ligated

into BsaI-digested, spin-purified plasmid as described above except 10 µl of the digested PCR

product was ligated with 500 ng cut plasmid in a 100 µl reaction and ligations were incubated at

16°C for 14 h. Library ligations were purified by spot dialysis on a nitrocellulose filter (Millipore

VSWP02500) against 0.1 mM Tris, pH 8 buffer for 20 min prior to transformation by

electroporation into *E. coli* mating strain WM6026 (sJMP424 or sJMP3257). To obtain sufficient

transformants for large libraries, electrocompetent cells were made 5X more concentrated in the

final step of the preparation[59] and multiple transformations were carried out depending on the

size of the library. Transformations were plated with selection on LB with carb and DAP at a

density of ~30,000-50,000 colonies/150 mm petri plate. Cells (>30X more colonies the number

of guides, e.g. >1.35 million CFU for the 45,000 guide libraries) were scraped from plates and

resuspended in LB + 15% glycerol, density was adjusted to OD_{600}=9, and aliquots were frozen

at -80°C.

Transfer of the CRISPRt/CRISPRtOE system to the bacterial chromosome. CRISPRt and

CRISPRtOE strains were constructed by triparental mating of two *E. coli* donor strains—one

with the pCRISPRt-H plasmid encoding Cas678-TniQ-TnsABC and another with the pCRISPRt-

T or pCtOE plasmid containing the transposon with the guide RNA and antibiotic resistance—

and a recipient strain. See Supplementary Table S6 for a summary of growth media and

conditions and antibiotic concentrations. All conjugations and selection post-conjugation were

carried out at 30°C regardless of the normal incubation temperature for the organism. Isolated

donor and recipient strains were struck out on plates with the appropriate media and antibiotic (if

relevant) concentrations. Colonies were resuspended from plates into the growth medium used

for the recipient to a density of OD_{600}=9. Equal amounts of donors and recipient were mixed

together (along with the appropriate no recipient, no CRISPRt-H donor, or no CRISPRt-T donor

controls), cells were pelleted by centrifugation at 5000 x g (except *A. baumannii* at 9000 x g),

placed on a nitrocellulose filter on an LB plate (except an RMG plate for *Z. mobilis*), and

incubated 12-18 h at 30°C. Cells from the filter were resuspended in 300 μl media, serially

diluted, and plated with antibiotic selection for the transposon (see growth and selection

conditions above) and without DAP to select against the donor strains, and incubated at 30°C

until colonies formed (1-3 days). A typical mating was 100 μl each strain and plating 100 μl

undiluted, and 10^{-1} through 10^{-4} dilutions to obtain isolated colonies. For library construction,

conjugations were scaled up and plated with selection at a density of ~30,000-50,000

colonies/150 mm petri plate. Cells (>30X more colonies the number of guides, e.g. >1.35 million

CFU for the 45,000 guide libraries) were scraped from plates and resuspended in LB + 15%

glycerol, density was adjusted to OD_{600}=10-15, and aliquots were frozen at -80°C.

Analysis of phenotype of *E. coli lacZ* and *gfp*-targeting CRISPRt.

For *gfp* targeted CRISPRt, fluorescence level (GFP) of 80 individual colonies was determined.

Cultures were grown in 300 μl media in 96 well deepwell plates from a single colony to

saturation. Cells were centrifuged at 4,000 x g for 10 min and cell pellets were resuspended in

300 μl 1X PBS. 150 μl was transferred to a 96-well black, clear bottom microplate (Corning

3631) and cell density (OD_{600}) and fluorescence (excitation 475 nm, emission 510 nm) were

measured in a fluorescence microplate reader (Tecan Infinite Mplex). Fluorescence values were

normalized to cell density.

For *lacZ* targeted CRISPRt, the color of 80 individual colonies was determined by patching on

LB agar plates containing X-gal (20 μg/ml). Blue indicates the presence of an intact *lacZ* due to

metabolism of the X-gal by LacZ (betagalactosidase) and white indicates disruption of *lacZ*.

Analysis of individual CRISPRtOE isolates from various Proteobacteria.

<u>Phenotype determination</u>. Fluorescence levels (mScarlet-I) of 12 CRISPRtOE isolates of all 8

species were determined. Cultures of isolates were grown in 300 μl media in 96 well deepwell

plates from a single colony to saturation. Cultures were serially diluted to 1:10,000 (1:100 twice)

into fresh media and grown again to saturation (~13 doublings). Cells were centrifuged at 4,000

x g for 10 min. Cell pellets were resuspended in 300μl 1X PBS and 150 μl was transferred to a

96-well black, clear bottom microplate (Corning 3631). Cell density (OD600) and fluorescence

(excitation 584 nm, emission 607 nms) were measured in a fluorescence microplate reader

(Tecan Infinite Mplex). Fluorescence values were normalized to cell density. The assay was

repeated three times for *E. coli* and *Z. mobilis* and twice for all other strains.

Genotype determination. Insertion positions of 12 CRISPRtOE isolates of all 8 species were

determined. Cultures of isolates were grown in 300 µl media in 96 well deepwell plates, serially

diluted 1:100 in dH$_2$O, heated to 90°C for 3 min prior to use as a template for PCR. Fragments

(~2000 bp) were amplified by PCR with a forward primer in the transposon (oJMP490) and a set

of indexed (4 nt barcode) reverse primers in the *mScarlet-I* coding sequence (oJMP2118-2129)

in a 25 µl reaction containing: 5 µl OneTaq buffer, 0.5 µl 10mM each dNTPs, 0.5 µl each 10 µM

forward and reverse primers, 2 µl diluted culture, 0.25 µl OneTaq DNA polymerase, and 16.25 µl

H$_2$O with the following touchdown PCR program: 94°C, 3 min then 10 cycles of 94°C, 30s,

65°C, 30s (-1°C/cycle), 68°C, 2.5 min, then 25 cycles of 94°C, 30s, 55°C, 30s, 68°C, 2.5 min,

then 68°C, 5 min in a BioRad T100 thermalcycler. Sets of indexed PCRs for each organism

were pooled, spin purified and sequenced by Oxford nanopore long read amplicon sequencing

by Plasmidsaurus. Sequencing data was demultiplexed using a custom Python script

(SplitSamplesSeq.py) and insertion position relative to the *mScarlet-I* ATG start codon was

determined.

CRISPRtOE library growth experiments.

Competition experiment growth. The *E. coli* CRISPRtOE *murA* and *folA* libraries were mixed in

equal volume with the LP-*lacZ* library: 50 µl frozen stock (OD$_{600}$ = 10) of each library (*murA* +

LP-*lacZ* and *folA* + LP-*lacZ*) into 100 ml LB (starting OD$_{600}$ = 0.01) in a 500 ml flask and

incubated shaking at 37°C until OD$_{600}$ = 0.2 (~2.5 h) (timepoint = T0) to revive the cells. These

cultures were diluted to OD$_{600}$ = 0.02 into 4 ml LB plus antibiotic fosfomycin (0.4 µg/ml, for the

murA + LP-*lacZ* libraries) or trimethoprim (0.1 µg/ml, for the *folA* + LP-*lacZ* libraries) or no

antibiotic control in 14 ml snap cap culture tubes (Corning 352059) in duplicate and incubated

with shaking for 18 h at 37°C (T1). These cultures were serially diluted back to OD$_{600}$ = 0.01 into

fresh tubes containing the same media and incubated again with shaking for 18 h at 37°C (T2)

for a total of ~10-15 doublings. Cells were pelleted from 15 ml of culture in duplicate at each

time point T0 and 1ml of culture at timepoints T1 and T2 and stored at -20°C.

Whole genome experiment growth. The *E. coli* CRISPRtOE whole genome libraries with no

promoter (sJMP10704) or promoter H (sJMP10705) were revived by dilution of 100 µl frozen

stock (OD$_{600}$ = 15) into 100 ml LB (starting OD$_{600}$ = 0.015) and incubation in 500 ml flasks

shaking at 37°C until OD_{600} = 0.2 (~2.5 h) (timepoint = T0). These cultures were diluted to OD_{600}

= 0.02 into 4 ml LB plus antibiotic trimethoprim (0.1 μg/ml) or no antibiotic control in 14 ml snap

cap culture tubes (Corning 352059) in duplicate and incubated with shaking for 18 h at 37°C

(T1). These cultures were serially diluted back to OD_{600} = 0.01 into fresh tubes containing the

same media and incubated again with shaking for 18 h at 37°C (T2) for a total of ~10-15

doublings. Cells were pelleted from 15 ml of culture in duplicate at time point T0 and 1ml of

culture at timepoints T1 and T2 and stored at -20°C.

Analysis of individual CRISPRtOE isolates from library competition experiment.

Insertion positions of 8 CRISPRtOE isolates from the *folA* and *murA* libraries were determined.

Cultures of isolates were grown in 3 ml media from an isolated colony, serially diluted 1:100 in

dH_2O, heated to 90°C for 3 min prior to use as a template for PCR. Fragments (~600 bp) were

amplified by PCR with a forward primer in the transposon (oJMP61) and a reverse primer in the

folA or *murA* coding sequence (oJMP2349 and oJMP2348, respectively) in a 50 μl reaction

containing: 10 μl OneTaq buffer, 1.0 μl 10mM each dNTPs, 1.0 μl each 10 μM forward and

reverse primers, 2 μl diluted culture, 0.5 μl OneTaq DNA polymerase, and 16.25 μl H_2O with the

following touchdown PCR program: 94°C, 3 min then 10 cycles of 94°C, 30s, 65°C, 30s (-

1°C/cycle), 68°C, 1 min, then 25 cycles of 94°C, 30s, 55°C, 30s, 68°C, 1 min, then 68°C, 5 min

in a BioRad T100 thermalcycler. PCR products were spin-purified and Sanger sequenced to

determine the transposon insertion position relative to the *mScarlet-I* ATG start codon.

Tnseq analysis of CRISPRt/CRISPRtOE constructs. TnSeq protocol was adapted from

Klompe et al[31]. Genomic DNA (gDNA) was extracted from the equivalent of 1 ml of cells at

OD_{600}=3 (~2x10^9 cells) either resuspended from a plate or from liquid culture and was further

purified by spot dialysis on a nitrocellulose filter (Millipore VSWP02500) against 0.1 mM Tris, pH

8 buffer for 20 min. 1 μg gDNA was digested with 4U MmeI (NEB) in a 50 μl reaction at 37°C, 12

hrs and heat inactivated at 65°C, 20 min. The digest was purified using 1.8X magnetic beads

(Omega) following the manufacturer's protocol, eluting in 20 μl 10 mM Tris, pH 8.0. Adapter

oligos oJMP1995 and P-oJMP1996 (phosphorylated) were annealed by mixing equal volume of

100μM oligos in 10mM Tris, pH8 (50 μM each), heating to 95°C, 5 min followed by cooling to RT

~15 min, and diluted 1:10 (5 μM each) for use. Annealed oligos were ligated onto MmeI-

digested DNA in a 20 μl reaction with 2 μl 10X T4 ligase buffer, 2 μl 100 mM DTT, 2 μl 1 mM

ATP, 12 μl purified MmeI-digested gDNA (50 ng/μl, 600 ng total), 1 μl 5 μM annealed adapter

oligos (250 nM final) and 1 μl T4 DNA ligase. Ligations were incubated 14 h at 16°C and the

enzyme was heat-inactivated for 20 min at 65°C. Ligations were purified using 1.8X magnetic

beads, eluting in 22 µl 10 mM Tris, pH 8. Fragments were amplified by PCR with primers

containing partial adapters for index PCR with Illumina TruSeq adapters in a 100µl reaction

containing: 20 µl 5X Q5 buffer, 3 µl GC enhancer, 2 µl 10 mM each dNTPs, 5 µl each forward

primer (oJMP1997) and reverse primer (oJMP1998 or indexed primers oJMP2022-2033), 10 µl

purified ligation, 1 µl Q5 DNA polymerase, and 54 µl dH$_2$O with the following program: 98°C, 30s

then 18-20 cycles of 98°C, 15s, 66°C, 15s, 72°C, 15s, then 72°C, 10 min in a BioRad C1000

thermal cycler. PCR products were spin-purifed (eluted in 15 µl) and quantified fluorometrically.

Samples were sequenced by the UWBC NGS Core facility or Azenta Amplicon-EZ service.

Briefly, PCR products were amplified with nested primers containing i5 and i7 indexes and

Illumina TruSeq adapters followed by bead cleanup, quantification, pooling and running on a

NovaSeq X Plus (150 bp paired end reads) or MiSeq (250 bp paired end reads). Sequencing

analysis of the initial CRISPRt or CRISPRtOE plasmid libraries from which the strain libraries

were prepared was by amplification with oJMP2011 and oJMP1998 (or barcoded version of

oJMP1998: oJMP2022-2033), followed by spin purification and Illumina sequencing.

Analysis of *E. coli mScarlet-I*-targeting CRISPRt libraries.

Two plasmid libraries (pJMP10505 and pJMP10506) were constructed (gen^R and kan^R) and

used to create CRISPRtOE libraries (sJMP10519, 10520, 10594, 10595, and 10637-10640) in

each of four *E. coli mScarlet-I* reporter strains (sJMP10205, 10269, 10630, 10633). Plasmid

DNA from pJMP10505 and pJMP10506 was amplified with oJMP2011 and oJMP1998. gDNA

from (sJMP10519, 10520, 10594, 10595, and 10637-10640) was processed by TnSeq as

detailed above with oJMP1997 and oJMP2022-2033. Sequencing data was demultiplexed using

SplitSamplesSeq.py and guides were quantified using seal.sh from bbmap.

MIC assays. The minimal inhibitory concentration (MIC) of *E. coli folA* and *murA* CRISPRtOE

isolates was determined by either growth in a microtiter plate or by disc diffusion assays,

respectively. For the broth microdilution assay, CRISPRtOE isolates and WT controls were

grown to saturation from an isolated colony and serially diluted to OD600 = 0.003) and the

CRISPRtOE library was diluted from a glycerol stock to the same density. Trimethoprim was

serially diluted in DMSO at 1000X concentration and then diluted to 2X in MHB. *E. coli folA*

CRISPRtOE isolates, the *folA* CRISPRtOE library, a *lacZ* CRISPRtOE control, and WT cultures

and the media containing antibiotic were mixed in equal proportions and incubated 20 h shaking

at 37°C prior to cell density (OD$_{600}$) determination in a microplate reader. For the MIC strip

assays, *E. coli folA* or *murA* CRISPRtOE isolates, a *lacZ* CRISPRtOE control, and WT cultures

were diluted to OD_{600} = 0.3 and 300 µl was spread on a 100 mm MH agar plate and dried in a

laminar flow cabinet. When dry, MIC strips (Liofilchem, Italy; 920371 (TMP 0.002-32 mg/L) and

920791(FOS 0.064-1024 mg/L)) were applied to the plate and 10 µl serially diluted antibiotics

were applied to the discs. Plates were incubated at 37°C for 18 h. Broth microdilution assays

were repeated 3 times and MIC test strip assays twice.

CRISPRtOE data processing and analysis. All commands and scripts used for processing

and analysis of CRISPRtOE data can be found on GitHub

(https://github.com/GLBRC/CRISPRtOE_Analysis). FASTQ files were separated for each read

(R1 and R2), concatenated, and trimmed to remove transposon-specific sequences using

Cutadapt (v3.4)[64] using default parameters and sequences flanking the desired genome

sequence. The trimmed FASTQ files were aligned to the *E. coli* genome (NCBI RefSeq

Assembly ID GCF_000005845.2) using Bowtie2 (v2.4.4)[65] using default parameters for paired-

end alignment. The alignment SAM file for each sample was filtered using Samtools (v1.13)[66] to

remove unmapped reads (-F 8) and to remove reads that mapped to opposite strands (-F 0x2).

This resulted in paired end reads that aligned to the same DNA strand. BBMap (v38.32)[67] was

used to filter the SAM files to remove read pairs wherein at least one read is less than 2

nucleotides or greater than 50 nucleotides in order to identify specific transposon insertion sites

downstream of spacer sequences. A custom Python script (parsing_sam_for_hisogram.py) and

a custom Rscript (Histogram_CRISPRtOE.R) were used to parse the filtered SAM files and

construct histograms representing the distance between the spacer sequence and the

transposon insertion site.

The SAM files for each sample were further filtered to remove reads there were more than 100

nucleotides apart using the Picard tools (v2.20)[68] "FilterSamReads" function with default

parameters. This resulted in an aligned file where in the spacer sequence was located on the

same strand as the transposon insertion site and in the same location (within 100 nucleotides).

These filtered files were used for all further analyses. Samtools (v1.13)[66] was used to sort and

index these files and DeepTools (v3.5.1)[69] was used to construct BigWig files (normalized using

CPM) and Bed files for visualization. To identify the distance between the transposon insertion

site and the nearest gene downstream, the alignment file was filtered to include only reads < 25

nucleotides (those reads corresponding to the site of transposon insertion) using Samtools

(v1.13)[66] and a custom Python script (site_of_insertion.py) was used to generate a resulting

table of the location of transposon insertion sites. The table was converted to a Bed file using a

custom Python script (table_to_bed.py) and the BEDOPS (v2.4.41)[70] command "closest-

features" was used to identify the distance between transposon insertion site and the nearest

gene downstream. CRISPRtOE distance results from all samples were combined and a custom

Python script (make_table_for_histogram.py) was used to construct the input for R. To compare

CRISPRtOE insertion sites relative to genes to that of traditional Tn-seq, Tn-seq data from

Goodall *et al.*[71] (TL1 samples) were aligned to the same *E. coli* genome (NCBI RefSeq

Assembly ID GCF_000005845.2) using Bowtie2 (v2.4.4)[65] using default parameters and Bed

files were created using DeepTools (v3.5.1)[69] as described above. The Tn-seq data was

processed to identify the distance between transposon insertion and the gene downstream and

compared to CRISPRtOE distance as above. Data were visualized on the IGV Genome

Browser[72] and Circos plots were created using Proksee[73].

A custom Python script (counting_spacers.py) was used to determine the number of times each

spacer appeared in the filtered alignment file. Spacers were retained for analysis if they had a

count ≥5. Pairwise comparison of spacer counts in DMSO and TMP treatment samples was

performed using edgeR[74] and a custom Rscript (comparison_of_spacer_counts_edgeR.R). The

results were plotted using a custom Rscripts (plotting_scatter_plot.R, barplot_with_points.R,

scatter_plot_commands.R). For gene analysis, spacers for each gene were averaged and

genes were retained if they contained >2 spacers for further analysis.

Data availability

727   Sequencing data will be deposited at the Sequence Read Archive. All other data are available

728   upon request.

Code availability

731   Guide design script is available at https://github.com/ryandward/CRISPRt_gRNA. Analysis and

732   demultiplexing scripts are available at https://github.com/GLBRC/CRISPRtOE_Analysis.

References

1. Jost, M. *et al.* Combined CRISPRi/a-Based Chemical Genetic Screens Reveal that Rigosertib

Is a Microtubule-Destabilizing Agent. *Molecular Cell* **68**, 210-223.e6 (2017).

2. Kampmann, M. Elucidating drug targets and mechanisms of action by genetic screens in

mammalian cells. *Chem Commun (Camb)* **53**, 7162–7167 (2017).

3. Kampmann, M. CRISPRi and CRISPRa Screens in Mammalian Cells for Precision Biology
and Medicine. *ACS Chem Biol* **13**, 406–416 (2018).

4. Cámara, E., Lenitz, I. & Nygård, Y. A CRISPR activation and interference toolkit for industrial
Saccharomyces cerevisiae strain KE6-12. *Sci Rep* **10**, 14605 (2020).

5. Dominguez, A. A., Lim, W. A. & Qi, L. S. Beyond editing: repurposing CRISPR-Cas9 for
precision genome regulation and interrogation. *Nat Rev Mol Cell Biol* **17**, 5–15 (2016).

6. Jost, M. & Weissman, J. S. CRISPR Approaches to Small Molecule Target Identification. *ACS*
Chem Biol **13**, 366–375 (2018).

7. Lian, J., Schultz, C., Cao, M., HamediRad, M. & Zhao, H. Multi-functional genome-wide
CRISPR system for high throughput genotype-phenotype mapping. *Nat Commun* **10**, 5794
(2019).

8. Santiago, M. *et al.* A new platform for ultra-high density Staphylococcus aureus transposon
libraries. *BMC Genomics* **16**, 252 (2015).

9. Kitagawa, M. *et al.* Complete set of ORF clones of Escherichia coli ASKA library (a complete
set of E. coli K-12 ORF archive): unique resources for biological research. *DNA Res.* **12**,
291–299 (2005).

10.   Tal, S. & Paulsson, J. Evaluating quantitative methods for measuring plasmid copy
numbers in single cells. *Plasmid* **67**, 167–173 (2012).

11.   Freed, E. F. *et al.* Genome-Wide Tuning of Protein Expression Levels to Rapidly
Engineer Microbial Traits. *ACS Synth Biol* **4**, 1244–1253 (2015).

12.   Warner, J. R., Reeder, P. J., Karimpour-Fard, A., Woodruff, L. B. A. & Gill, R. T. Rapid
profiling of a microbial genome using mixtures of barcoded oligonucleotides. *Nat Biotechnol*
28, 856–862 (2010).

13.   Mutalik, V. K. *et al.* Dual-barcoded shotgun expression library sequencing for high-
throughput characterization of functional traits in bacteria. *Nat Commun* **10**, 308 (2019).

14.   Huang, Y. Y. *et al.* Barcoded overexpression screens in gut Bacteroidales identify genes
with new roles in carbon utilization and stress resistance. 2022.10.10.511384 Preprint at
https://doi.org/10.1101/2022.10.10.511384 (2023).

15.   Mali, P. *et al.* CAS9 transcriptional activators for target specificity screening and paired
nickases for cooperative genome engineering. *Nat Biotechnol* **31**, 833–838 (2013).

16.   Lee, D. J., Minchin, S. D. & Busby, S. J. W. Activating transcription in bacteria. *Annu Rev*
Microbiol **66**, 125–152 (2012).

17. Bikard, D. *et al.* Programmable repression and activation of bacterial gene expression

using an engineered CRISPR-Cas system. *Nucl. Acids Res.* gkt520 (2013)

doi:10.1093/nar/gkt520.

18. Kiattisewee, C. *et al.* Expanding the Scope of Bacterial CRISPR Activation with PAM-

Flexible dCas9 Variants. *ACS Synth Biol* **11**, 4103–4112 (2022).

19. Ho, H.-I., Fang, J. R., Cheung, J. & Wang, H. H. Programmable CRISPR-Cas

transcriptional activation in bacteria. *Mol Syst Biol* **16**, e9427 (2020).

20. Peters, J. E., Makarova, K. S., Shmakov, S. & Koonin, E. V. Recruitment of CRISPR-Cas

systems by Tn7-like transposons. *PNAS* **114**, E7358–E7366 (2017).

21. Vo, P. L. H. *et al.* CRISPR RNA-guided integrases for high-efficiency, multiplexed

bacterial genome engineering. *Nat Biotechnol* **39**, 480–489 (2021).

22. Rubin, B. E. *et al.* Species- and site-specific genome editing in complex bacterial

communities. *Nat Microbiol* **7**, 34–47 (2022).

23. Chen, W. *et al.* Targeted genetic screening in bacteria with a Cas12k-guided

transposase. *Cell Rep* **36**, 109635 (2021).

24. Trujillo Rodríguez, L., Ellington, A. J., Reisch, C. R. & Chevrette, M. G. CRISPR-

Associated Transposase for Targeted Mutagenesis in Diverse Proteobacteria. *ACS Synth Biol*

12, 1989–2003 (2023).

25. Roberts, A., Nethery, M. A. & Barrangou, R. Functional characterization of diverse type I-

F CRISPR-associated transposons. *Nucleic Acids Res* **50**, 11670–11681 (2022).

26. Hsieh, S.-C. & Peters, J. E. Discovery and characterization of novel type I-D CRISPR-

guided transposons identified among diverse Tn7-like elements in cyanobacteria. *Nucleic*

Acids Res **51**, 765–782 (2023).

27. Walker, M. W. G., Klompe, S. E., Zhang, D. J. & Sternberg, S. H. Novel molecular

requirements for CRISPR RNA-guided transposition. *Nucleic Acids Res* **51**, 4519–4535

(2023).

28. Gelsinger, D. R. *et al.* Bacterial genome engineering using CRISPR RNA-guided

transposases. *bioRxiv* 2023.03.18.533263 (2023) doi:10.1101/2023.03.18.533263.

29. Hoffmann, F. T. *et al.* Selective TnsC recruitment enhances the fidelity of RNA-guided

transposition. *Nature* **609**, 384–393 (2022).

30. Halpin-Healy, T. S., Klompe, S. E., Sternberg, S. H. & Fernández, I. S. Structural basis of

DNA targeting by a transposon-encoded CRISPR-Cas system. *Nature* **577**, 271–274 (2020).

31. Klompe, S. E., Vo, P. L. H., Halpin-Healy, T. S. & Sternberg, S. H. Transposon-encoded

CRISPR–Cas systems direct RNA-guided DNA integration. *Nature* **571**, 219 (2019).

32.   Garza Elizondo, A. M. & Chappell, J. Targeted Transcriptional Activation Using a
CRISPR-Associated Transposon System. *ACS Synth Biol* **13**, 328–336 (2024).

33.   Petassi, M. T., Hsieh, S.-C. & Peters, J. E. Guide RNA Categorization Enables Target
Site Choice in Tn7-CRISPR-Cas Transposons. *Cell* **183**, 1757-1771.e18 (2020).

34.   Banta, A. B., Enright, A. L., Siletti, C. & Peters, J. M. A High-efficacy CRISPRi System for
Gene Function Discovery in Zymomonas mobilis. *Appl. Environ. Microbiol.* (2020)
doi:10.1128/AEM.01621-20.

35.   Hall, A. N. *et al.* Tools for Genetic Engineering and Gene Expression Control in
Novosphingobium aromaticivorans and Rhodobacter sphaeroides. *bioRxiv*
2023.08.25.554875 (2023) doi:10.1101/2023.08.25.554875.

36.   Enright, A. L. *et al.* The genetics of aerotolerant growth in an alphaproteobacterium with
a naturally reduced genome. *mBio* e0148723 (2023) doi:10.1128/mbio.01487-23.

37.   Rice, L. B. Federal funding for the study of antimicrobial resistance in nosocomial
pathogens: no ESKAPE. *J Infect Dis* **197**, 1079–1081 (2008).

38.   Fenster, J. A. *et al.* Dynamic and single cell characterization of a CRISPR-interference
toolset in Pseudomonas putida KT2440 for β-ketoadipate production from p-coumarate.
Metab Eng Commun **15**, e00204 (2022).

39.   Hau, H. H. & Gralnick, J. A. Ecology and biotechnology of the genus Shewanella. *Annu*
Rev Microbiol **61**, 237–258 (2007).

40.   Prasad, N. K., Seiple, I. B., Cirz, R. T. & Rosenberg, O. S. Leaks in the Pipeline: a
Failure Analysis of Gram-Negative Antibiotic Development from 2010 to 2020. *Antimicrobial*
Agents and Chemotherapy **66**, e00054-22 (2022).

41.   Peters, J. M. *et al.* A Comprehensive, CRISPR-based Functional Analysis of Essential
Genes in Bacteria. *Cell* **165**, 1493–1506 (2016).

42.   Johnson, R. A. *et al.* Inhibition of Isoleucyl-tRNA Synthetase by the Hybrid Antibiotic
Thiomarinol. *J Am Chem Soc* **143**, 12003–12013 (2021).

43.   Palmer, A. C. & Kishony, R. Opposing effects of target overexpression reveal drug
mechanisms. *Nat Commun* **5**, 4296 (2014).

44.   Couce, A. *et al.* Genomewide overexpression screen for fosfomycin resistance in
Escherichia coli: MurA confers clinical resistance at low fitness cost. *Antimicrob Agents*
Chemother **56**, 2767–2769 (2012).

45.   Ason, B. & Reznikoff, W. S. DNA sequence bias during Tn5 transposition. *J Mol Biol* **335**,
1213–1225 (2004).

46.   Lee, J.-H., Lee, K.-L., Yeo, W.-S., Park, S.-J. & Roe, J.-H. SoxRS-mediated
lipopolysaccharide modification enhances resistance against multiple drugs in Escherichia
coli. *J Bacteriol* **191**, 4441–4450 (2009).

47.   Griffith, K. L., Becker, S. M. & Wolf, R. E. Characterization of TetD as a transcriptional
activator of a subset of genes of the Escherichia coli SoxS/MarA/Rob regulon. *Mol Microbiol*
56, 1103–1117 (2005).

48.   Seo, S. W., Kim, D., Szubin, R. & Palsson, B. O. Genome-wide Reconstruction of OxyR
and SoxRS Transcriptional Regulatory Networks under Oxidative Stress in Escherichia coli
K-12 MG1655. *Cell Rep* **12**, 1289–1299 (2015).

49.   Jones, K. M., Guest, J. R. & Woods, D. D. Folic acid and the synthesis of methionine by
extracts of Escherichia coli. *Biochem J* **79**, 566–574 (1961).

50.   Weissbach, H. & Brot, N. Regulation of methionine synthesis in Escherichia coli. *Mol*
Microbiol **5**, 1593–1597 (1991).

51.   Saint-Girons, I. *et al.* Nucleotide sequence of metF, the E. coli structural gene for 5-10
methylene tetrahydrofolate reductase and of its control region. *Nucleic Acids Res* **11**, 6723–
6732 (1983).

52.   Girgis, H. S., Hottes, A. K. & Tavazoie, S. Genetic architecture of intrinsic antibiotic
susceptibility. *PLoS One* **4**, e5629 (2009).

53.   Nichols, R. J. *et al.* Phenotypic landscape of a bacterial cell. *Cell* **144**, 143–156 (2011).

54.   Giladi, M., Altman-Price, N., Levin, I., Levy, L. & Mevarech, M. FolM, a new
chromosomally encoded dihydrofolate reductase in Escherichia coli. *J Bacteriol* **185**, 7015–
7018 (2003).

55.   Rao, T. V. P. & Kuzminov, A. Exopolysaccharide defects cause hyper-thymineless death
in Escherichia coli via massive loss of chromosomal DNA and cell lysis. *Proc Natl Acad Sci U*
S A **117**, 33549–33560 (2020).

56.   Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile Genetic Elements
Associated with Antimicrobial Resistance. *Clin Microbiol Rev* **31**, e00088-17 (2018).

57.   Martínez-García, E., Aparicio, T., de Lorenzo, V. & Nikel, P. I. New transposon tools
tailored for metabolic engineering of gram-negative microbial cell factories. *Front Bioeng*
Biotechnol **2**, 46 (2014).

58.   Yang, Q. *et al.* Development and characterization of acidic-pH-tolerant mutants of
Zymomonas mobilis through adaptation and next-generation sequencing-based genome
resequencing and RNA-Seq. *Biotechnol Biofuels* **13**, 144 (2020).

59.   Banta, A. B., Ward, R. D., Tran, J. S., Bacon, E. E. & Peters, J. M. Programmable Gene
Knockdown in Diverse Bacteria Using Mobile-CRISPRi. *Curr Protoc Microbiol* **59**, e130
(2020).

60.   Sharan, S. K., Thomason, L. C., Kuznetsov, S. G. & Court, D. L. Recombineering: a
homologous recombination-based method of genetic engineering. *Nat Protoc* **4**, 206–223
(2009).

61.   Thomason, L. C., Costantino, N. & Court, D. L. E. coli genome manipulation by P1
transduction. *Curr Protoc Mol Biol* **Chapter 1**, 1.17.1-1.17.8 (2007).

62.   Cherepanov, P. P. & Wackernagel, W. Gene disruption in Escherichia coli: TcR and KmR
cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant.
Gene **158**, 9–14 (1995).

63.   Peters, J. M. *et al.* Enabling genetic analysis of diverse bacteria with Mobile-CRISPRi.
Nature Microbiology **4**, 244–250 (2019).

64.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**, 10–12 (2011).

65.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357–359 (2012).

66.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

67.   Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. (2014).

68.   Picard Tools - By Broad Institute. https://broadinstitute.github.io/picard/.

69.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

70.   Neph, S. *et al.* BEDOPS: high-performance genomic feature operations. *Bioinformatics*
28, 1919–1920 (2012).

71.   Goodall, E. C. A. *et al.* The Essential Genome of Escherichia coli K-12. *mBio* **9**, e02096-
17 (2018).

72.   Robinson, J. T., Thorvaldsdottir, H., Turner, D. & Mesirov, J. P. igv.js: an embeddable
JavaScript implementation of the Integrative Genomics Viewer (IGV). *Bioinformatics* **39**,
btac830 (2023).

73.   Grant, J. R. *et al.* Proksee: in-depth characterization and visualization of bacterial
genomes. *Nucleic Acids Res* **51**, W484–W492 (2023).

74.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).

Acknowledgements

We thank members of the Peters lab and Michael Thomas for helpful discussions and Joshua
Hyman and the University of Wisconsin-Madison Biotechnology Center for technical assistance
with Illumina sequencing and Samuel Sternberg for plasmids. This work was supported by the
U.S. Department of Energy, Office of Science and Office of Biological and Environmental
Research, Great Lakes Bioenergy Research Center, under Award Number DE-SC0018409, and
by the National Institutes of Health under award number 1R35GM150487-01 to J.M.P.

Author contributions

916   A.B.B. and J.M.P. designed, performed, and interpreted the experiments and wrote the
917   manuscript with input from all authors. K.S.M. developed the data analysis pipeline and
918   performed data analysis. R.D.W. and M.P. developed code. R.A.C., C.C.F., and E.E.B
919   performed experiments.

Competing interests

922   The authors declare no competing interests.

Figure legends

Fig. 1 A mobilizable, selectable, dual-plasmid system for CRISPR-guided transposition. a
Schematic of a targeted transposition gene knockout (CRISPRt) and gene overexpression
(CRISPRtOE) system optimized for functional genomics. **b** Schematic of mechanism of *Vibrio*
cholera Type I-F CRISPR-associated transposon (VcCAST) system[20,21]. **c** Schematic of strain
construction using the CRISPRt targeted transposition system. Plasmids have pir-dependent
origins of replication preventing replication in the bacterial recipient. One plasmid encodes the
minimal Type I-F CRISPR-Cas (Cas678) and Tn*7*-like transposase (TnsABC and TniQ)
machinery, and a second plasmid contains a transposon carrying the guide RNA and antibiotic
resistance expression cassettes (see Fig S1 for details). These plasmids are transferred by co-
conjugation to a recipient bacterium by *E. coli* donor strains with a chromosomal copy of the
RP4 transfer machinery. Inside the recipient cell, the transposon (flanked in yellow) is inserted

onto the chromosome at a site determined by the sequence of the guide RNA. Selection on antibiotic plates lacking diaminopimelic acid (DAP) selects for transconjugants and against the *E. coli* donor strains. **d** Specificity of CRISPRt disruption of *gfp* and *lacZ* in *E. coli* measured by Tnseq. Mapping of location of transposon insertion sites to the *E. coli* genome after CRISPRt targeted transposition with *gfp1* guide (226,671 reads) or *LZ1* guide (216,801 reads). **e** Insertion position (bp downstream of protospacer) frequency of CRISPRt disruption of *gfp* and *lacZ* in *E. coli* (RL orientation). Percentage of on-target, RL orientation insertions for all positions with > 0.01%.

Fig. 2 Transcription impacts efficacy of CRISPRt guides. a Schematic of guides in pooled *mScarlet-I* disruption experiment. Twenty-three guides matching protospacers located on top (T1-24) or bottom (B1-24) strand of the mScarlet-I encoding gene with a range of CN guides (see Table S4 for details). **b** Relative efficacy of CRISPRt *mScarlet-I* guides in a pooled screen targeting *mScarlet-I* expressed from a transposon in the att_{Tn7} site on the *E. coli* chromosome. Relative efficacy is expressed as the log_2-fold change of the frequency in the CRISPRt libraries (sJMP10519 and sJMP10520) vs. the frequency in the original plasmid construct libraries (sJMP10505 and sJMP10506). Average and standard deviation of two separate CRISPRt libraries (gentamicin^R and kanamycin^R) containing all 46 guides. **c** Absolute transposition efficiency of 3 individual CRISPRt guides (T03, T09, T21) targeting *mScarlet-I* cassettes in the *E. coli* att_{Tn7} site either with a promoter (+ transcription) or without a promoter (- transcription) measured by plating efficiency with and without selection (+/- kanamycin) in triplicate. Error represents the average of n=3 assays, * indicates p<0.05, two-tailed t-test.

Fig. 3 Tunable overexpression of chromosomally located genes using CRISPRtOE. a Schematic of CRISPRtOE targeted overexpression system. The chromosomal target is an *mScarlet-I* gene preceded by a 'Landing Pad' (LP) with no promoter, a PAM, and the *LZ1* protospacer (see Fig S5a). The CRISPRtOE construct has a transposon carrying guide RNA and antibiotic resistance expression cassettes and an outward facing promoter (see Supplementary Fig. S5b). Co-conjugation of strains carrying the CRISPRtOE construct and the CRISPRt-H plasmid (harboring the VcCAST machinery) with a promoterless *mScarlet-I* reporter recipient strain results in a CRISPRtOE strain. **b** mScarlet-I fluorescence analysis of *E. coli* CRISPRtOE isolates (no promoter or synthetic promoters A, V, and H) compared to the parent (promoterless *mScarlet-I*) strain. **c** Fold effect of CRISPRtOE mScarlet-I overexpression using Promoter H in eight Alpha- and Gammaprotebacteria (*E. coli* (*Eco*), *Enterobacter cloacae* (*Ecl*),

Klebsiella pneumoniae (*Kpn*), *Acinetobacter baumannii* (*Aba*), *Pseudomonas aeruginosa* (*Pae*),

Pseudomonas putida (*Ppu*), *Zymomonas mobilis* (*Zmo*), and *Shewanella oneidensis* (*Son*)).

CRISPRtOE transposon insertion position and fluorescence measurements were determined for

twelve isolates. Fluorescence measurements were normalized to cell density (OD_{600}). Values

are shown for isolates with on-target, RL orientation CRISPRtOE insertions and fold changes for

on-target and RL orientation CRISPRtOE isolates compared to a strain with promoterless

mScarlet-I (see Table S8 for details). Error is expressed for the median value of 8-12 isolates in

n=3 (*Eco* and *Zmo*) or n=2 (*Ecl*, *Kpn*, *Aba*, *Pae*, *Ppu*, and *Son*) assays.

Fig. 4 CRISPRtOE informs antibiotic mode of action. a Schematic of CRISPRtOE

competitive growth assay. A pooled CRISPRtOE strain library is constructed by amplification of

spacer sequences from a pooled oligo library and transfer onto the chromosome of a recipient

strain. Insertion of the CRISPRtOE transposon is either upstream of genes of interest or within a

control gene (e.g. *lacZ*) in *E. coli*. A CRISPRtOE pooled library screen is conducted by culturing

a CRISPRtOE pooled library and control library for multiple generations in the presence or

absence of a chemical or condition of interest. The change in the composition of the pooled

library before and after treatment is measured by amplification and NGS sequencing of the

spacer sequences. **b** CRISPRtOE pooled library screen. Fitness of pooled strains

overexpressing *folA* compared to the *lacZ* control strains in the presence of trimethoprim (TMP).

c MIC broth microdilution assay of individual CRISPRtOE *folA* strains compared to WT, a

CRISPRtOE *lacZ* isolate, and the CRISPRtOE *folA* pooled library in the presence of TMP (0.04-

20 µg/ml). Error is expressed for 3 separate assays.

Fig. 5 Genome-scale CRISPRtOE. a Genome plot showing positions of CRISPRtOE insertions

in TMP-treated cells. Data are scaled to the maximum number of counts. **b** Positions of

CRISPRtOE insertions upstream of all genes in the genome. Insertion site distance d was

calculated based on the distance in bp from the 3′ end of the Tn to the 5′ end of the gene. **c**

CRISPRtOE insertions into the *lac* locus. The scale is capped at 20 reads for visual clarity. **d**

Volcano plot of gRNA spacer counts in TMP vs DMSO control. **e** Top positive gene hits in TMP

treatment. Genes must have at least a 4-fold change in the promoter H data and show a 4-fold

difference in a comparison between promoter H and no promoter to be considered. Error bars

represent standard deviation (SD).

1002

Fig. S1 Details of CRISPRt plasmids and *E. coli* GFP reporter strain. a Plasmid maps of

CRISPRt-H helper and CRISPRt-T transposon plasmids. Both plasmids in the dual plasmid

system have backbones with an R6K pir-dependent replication origin, an ampicillan resistance

cassette, and conjugative transfer origin. The CRISPRt-H helper plasmid has expression

cassettes for the minimal Vch CRISPR (Cas678) and Transposase (TnsABC, TniQ) machinery

derived from pSpin-R[21]. The CRISPRt-T transposon plasmid encodes a modified VchINT

transposon derived from pSpin-R[21] which includes a guide RNA expression cassette with a

modified BsaI cloning site and an antibiotic expression cassette and transcription terminators

derived from CRISPRi plasmids[63]. The antibiotic resistance cassette is flanked by XhoI sites to

facilitate modification, three variants are shown. **b** Sequence of CRISPRt targeted transposition

system guide RNA expression cassette including the BsaI cloning site for new guide RNAs. **c**

Position of the P$_{veg}$-*sfgfp* expression cassette inserted between *yjaA* and *yjaB*. on the *E. coli*

chromosome.

1016

Fig. S2 CRISPRt disruption efficiency in *E. coli*. a Efficiency of CRISPRt disruption of *sfgfp* in

E. coli using a fluorescence phenotypic assay. The *E. coli yjaA-sfgfp-yjaB* strain (sJMP2688) is

green fluorescent whereas the *E. coli* WT strain (sJMP163) is not. Isolates from CRISPRt

targeted transposition of *E. coli yjaA-sfgfp-yjaB* with the *gfp1* guide (selection for

transconjugants on LB-kan plates) were assayed for fluorescence; 82/84 (97.6%) were white

indicating disruption of *sfgfp*. **b** Efficiency of CRISPRt disruption of *lacZ* in *E. coli* using an X-gal

phenotypic assay. The *E. coli* WT strain (sJMP163) can metabolize X-gal resulting in a blue

colony color whereas the *E. coli lacZYA* deletion strain (sJMP3272) cannot metabolize X-gal

resulting in a white colony color. Isolates from CRISPRt targeted transposition of *E. coli* WT with

the *LZ1* guide (selection for transconjugants on LB-kan plates) were assayed for colony color;

90/90 (100%) were white indicating disruption of *lacZ*. **c** Efficiency of obtaining CRISPRt

transconjugants in *E. coli* using the *LZ1* and *gfp1* guides. Serial dilutions of three technical

replicates of *LZ1* and *gfp1* CRISPRt assays were plated with and without selection (LB-kan or

LB, respectively) and CFUs were compared; Efficiency of CRISPRt targeted transposition in *E.*

coli with the *LZ1* guide was ~1 in 100 and with the *gfp1* guide was ~1 in 1000.

1032

Fig. S3 Transcription impacts efficacy of CRISPRt guides. a. No correlation between PAM

sequence (CA, CC, CG, CT) and efficacy (frequency of insertion) for 46 *mScarlet-I* targeting

guides. **b**. No correlation between CRISPRt guide position and efficacy for 46 *mScarlet-I*

targeting guides. **c.** Schematic of *mScarlet-I* cassettes +/- promoter in *E. coli* chromosome. The

mScarlet-I expression cassette (red) with and without a promoter (orange) along with a

spectinomycin resistance cassette (SmR, brown) are located on the *E. coli* chromosome just

downstream of *glmS* and upstream of *pstS*. The *mScarlet-I* expression cassette is inserted in

one of two orientations so that the direction of transcription is either in the same or the opposite

direction of the leading strand of chromosomal DNA replication. **d** Relative efficacy of CRISPRt

mScarlet-I guides in a pooled screen targeting *mScarlet-I* expressed on the *E. coli* chromosome

in different orientations with and without transcription. Relative efficacy is expressed as the log2-

fold change of the frequency in the CRISPRt libraries (sJMP10519-10520, sJMP10594-10595,

sJMP10637-10640) vs. the frequency in the original plasmid construct libraries (sJMP10505 and

sJMP10506) shown on scatter plots.

1047

Fig. S4 Transcription impacts efficacy of CRISPRt guides. a Relative efficacy of CRISPRt

gRNAs in a pooled screen targeting *mScarlet-I* as in **Fig S3d** with average and standard

deviation of two separate CRISPRt libraries (gentamicin^R and kanamycin^R) containing all 46

guides shown. **b** Fluorescent phenotypes of 80 isolates from each of the *E. coli* CRISPRt

mScarlet-I individual guide libraries (sJMP10641-10646) compared to *E. coli* strains with no

mScarlet-I gene ((-) mScar, sJMP6999) or expressing *mScarlet-I* ((+) mScar, sJMP10205). Only

4/480 (0.83%) retained *mScarlet-I* levels equal to the parent strain indicating an off-target rate of

<1% overall.

1056

Fig. S5 CRISPRtOE construct details. a Detailed schematic of the chromosomally-encoded

promoterless *mScarlet-I* reporter contained on a Tn*7* transposon inserted into the att_{Tn7} site of *E.*

coli. **b** Detailed schematics of CRISPRtOE Tn*7*-like transposon constructs with four different

outward facing promoter sequences (no promoter, and constitutive promoters A, H, and V). **c**

Sequence alignment of CRISPRtOE promoters A, H, and V compared to a consensus *E. coli* σ^{70}

promoter. Promoter A is based on the *lacUV5* promoter. Promoter H and V are synthetic

promoters. Promoter H has an UP element highlighted in yellow. **d** Schematic of CRISPRtOE

transposon insertion into Landing Pad (LP) showing CRISPRtOE Promoter and transposon L-

end relative to LP RBS and *mScarlet-I* coding sequence.

1066

1067   **Fig. S6 Tunable overexpression of chromosomally located genes using CRISPRtOE.**

1068   **a** mScarlet-I fluorescence analysis of *E. coli* CRISPRtOE isolates (no promoter or synthetic

1069   promoters A, V, and H) compared to the parent (promoterless *mScarlet-I*) strain. As in Fig 3b

1070   except values are shown for all CRISPRtOE isolates. **b** mScarlet-I fluorescence analysis of

CRISPRtOE isolates of the Alphaproteobacterium *Zymomonas mobilis* (no promoter or synthetic promoters A, V, and H) compared to the parent (promoterless *mScarlet-I*) strain. Values are shown for isolates with on-target, RL orientation CRISPRtOE insertions. **c** Schematic showing insertion of CRISPRtOE transposon into the 'Landing Pad' upstream of *mScarlet-I*. Insertion position is variable (~45-55bp downstream of protospacer) resulting in a variable amount of the Landing Pad (containing the RBS) remaining upstream of the *mScarlet-I* coding sequence. Insertion positions and orientation of CRISPRtOE isolates in the eight species shown in Fig3 c. Insertion position is expressed relative to start of *mScarlet-I* coding sequence (equal to the remaining LP sequence).

Fig. S7 CRISPRtOE *murA* pooled library screen and *folA* and *murA* MIC assays. **a** Fitness of pooled strains overexpressing *murA* compared to the *lacZ* control strains in the presence of fosfomycin. **b** Schematic of CRISPRtOE insertion position upstream of genes. **c** MIC test strip assays (trimethoprim, 0.002-32 μg/ml) comparing WT and CRISPRtOE *folA* isolates. **d** MIC test strip assays (fosfomycin, 0.064-1024 μg/ml) comparing WT and CRISPRtOE *murA* isolates.

Fig. S8 CRISPRtOE reproducibility. a Biological replicates of CRISPRtOE data at the gene level. **b** Replicate comparison of TMP treatment versus DMSO control. FC = fold change. **c** Comparison of TMP versus DMSO for the promoter H versus no promoter libraries.

Fig. S9 CRISPRtOE robustness with fewer analyzed guides. Scatterplots comparing all spacers measured in our study to subsets of 5, 4, 3, or 2 randomly selected spacers per gene.

Fig. S10 CRISPRtOE TMP screen positive and negative outlier genes. a Top positive gene hits in TMP treatment only considering the first gene in the operon. Genes must have at least a 4-fold change in the promoter H data and show a 4-fold difference in a comparison between promoter H and no promoter to be considered. **b** Genes from the enriched STRING group "DNA binding HTH domain, AraC-type, and Transcription regulators MarR/SlyA-like" (CL:3238) as well as *acrA*. **c** Genes from the enriched GO group "Enterobacterial common antigen biosynthetic process" (GO:0009246) as well as *metF* and *folM*. Error bars represent standard deviation (SD).

Fig 1

a

VcCAST
for genome-scale
functional genomics

Product

gene

Targeted Transposition
and selection

Tn   Tn   🔴 Promoter

Product
Product
Product

CRISPRt
Gene Knockout

ge Tn ne

CRISPRtOE
Gene Overexpression

Tn   gene

b

VcCAST transposon insertion

Tn*7*-like Transposase/Transposon

Minimal Type I-F CRISPR-Cas

Tn*6677*
Transposon

Cas678   gRNA

spacer

TniQ TnsABC

PAM   target DNA

protospacer

~49 bp

Integrated Transposon

c

DAP⁻
E. coli
Donors

H

T

+

+

DAP⁺
Bacterial
Recipient

tri-parental
mating

Abᴿ
DAP⁺
selection

VcCAST for functional genomics

cas6 cas7 cas8 tniQ tnsA tnsB tnsC

pir-dependent CRISPRt Helper plasmid

R r s r Abᴿ L

pir-dependent CRISPRt Transposon plasmid
with gRNA inside Tn

lacZ   gfp

Bacterial genome containing reporter gene

g R r s r Abᴿ L fp

Selectable CRISPRt Knockout

d

CRISPRt insertions into *gfp* (*gfp1* spacer)

Orientation
R 89.2% L
L 10.7% R

99.68% on target

E. coli genome position

CRISPRt insertions into *lacZ* (*LZ1* spacer)

99.94% on target

Orientation
R 74.1% L
L 25.9% R

E. coli genome position

e

gfp1 insertion position (on target, RL orientation)

45	46	47	48	49	50	51	52	53
0.0	0.0	0.0	24.8	67.3	7.3	0.6	0.0	0.0

Insertion Position Downstream of Protospacer

_LZ1_ insertion position (on target, RL orientation)

45	46	47	48	49	50	51	52	53
0.3	0.0	0.1	0.0	96.9	1.9	0.0	0.7	0.0

Insertion Position Downstream of Protospacer

Fig 2

a

CRISPRt *mScarlet-I* spacer library

spacer/protospacer locations

gRNAs bind to template strand

gRNAs bind to non-template strand

b

CRISPRt *mScarlet-I* guide relative efficacy

c

CRISPRt transposition efficiency +/- transcription

Fig 3

a

cas6 | cas7 | cas8 | tniQ | tnsA | tnsB | tnsC

Helper plasmid

+

R | r | s | r | Ab^R | L

Transposon with outward facing promoter

+

LP | mScarlet-I

LZ1

Promoterless *mScarlet-I* in chromosome

CRISPR-guided transposition

Tn | mScarlet-I

CRISPRt Targeted OverExpression

b

E. coli **K-12 CRISPRtOE**

mScarlet-I expression (AU/OD600)

12614

2146

71 | 71 | 73

~177X

~30X

~2X

Parent | LP | no P | P$_V$ | P$_A$ | P$_H$

CRISPRtOE

c

CRISPRtOE in diverse bacteria (Promoter H)

Fold change mScarlet-I expression

| *Eco* | *Ecl* | *Kpn* | *Aba* | *Pae* | *Ppu* | *Zmo* | *Son* |
| 177X | 232X | 158X | 73X | 23X | 62X | 50X | 30X |

Pathogen | Industrial

Fig 4

a

CRISPRtOE pooled gRNA library

gRNAs targeting upstream of gene

CRISPRtOE pooled strain library

drug targets

negative control

CRISPRtOE pooled screen

Growth +/- drug

T0 → T2

Amplify gRNA spacers from genome

Illumina sequencing

Count spacers to identify changes in pool composition with treatment

b

CRISPRtOE Screen (*folA*-trimethoprim)

Relative fitness score (log2-fold change T0 vs. T2)

~22X

lacZ | *folA* — No Drug

lacZ | *folA* — Trimethoprim

c

CRISPRtOE *folA*-trimethoprim MIC

Relative Growth vs. [Trimethoprim] (µg/ml)

- WT
- *lacZ* control
- CtOE library
- CtOE *folA* isolate (-20)
- CtOE *folA* isolate (-29)
- CtOE *folA* isolate (-45)
- CtOE *folA* isolate (-80)

Fig 5

a Genome-scale CRISPRtOE in *E. coli* K-12

b CRISPRtOE insertions upstream of genes

Fig S1

Fig S2

a

Strain	Parent Strain genotype	CRISPRt guide	Fluorescence phenotype
sJMP163	E. coli MG1655 WT	na	white
sJMP2688	E. coli MG1655 gfp	na	green
isolates	E. coli MG1655 gfp	gfp1	82/84 white (97.6% gfp disruption)

b

Strain	Parent Strain genotype	CRISPRt guide	X-gal phenotype
sJMP163	E. coli MG1655 WT	na	blue
sJMP3272	E. coli MG1655 DlacZYA	na	white
isolates	E. coli MG1655 WT	LZ1	90/90 white (100% lacZ disruption)

c

Fig S3

a

Frequency of insertion vs. PAM

Legend: CA (green), CC (blue), CG (yellow), CT (red)

Y-axis: Frequency (L2FC)

X-axis categories: mScarlet-I (+promoter), mScarlet-I (+promoter, flipped), mScarlet-I (-promoter), mScarlet-I (-promoter, flipped)

b

CRISPRt guide efficacy by position

Y-axis: Log2-fold change

X-axis (top): CRISPRt guide (template strand targeting) — T01, T02, T03, T05, T06, T07, T08, T09, T10, T11, T12, T13, T14, T15, T16, T17, T18, T19, T20, T21, T22, T23, T24

X-axis (bottom): CRISPRt guide (non-template strand targeting) — B01, B02, B03, B04, B06, B07, B08, B09, B10, B11, B12, B13, B14, B15, B16, B17, B18, B19, B20, B21, B22, B23, B24

c

CRISPRt transposon insertion

replication

DNA strand	Promoter
leading	+
lagging	+
leading	−
lagging	−

d

Log2-fold change mScarlet-I (+promoter) rep1 vs rep2 — R² = 0.844

Log2-fold change mScarlet-I (+promoter, flipped) rep1 vs rep2 — R² = 0.8192

Log2-fold change mScarlet-I (-promoter) rep1 vs rep2 — R² = 0.7194

Log2-fold change mScarlet-I (-promoter, flipped) rep1 vs rep2 — R² = 0.5483

Fig S4

Fig S5

a

b

c

CRISPRtOE Promoter Sequences

```
A   CTCACTCATTAGGCACCCCAGGCTTTACACTTTATGCTTCCGGCTCGTATAATGTCTAG
H   GGAAAATTTTTTTTTCAAAAGTACTTGAAAATTAATCA•TCCGGCTCGTATAATTCTAGT
V   CACCAAGCCAGTTACCTCGGTTCTTTAATCTTTATGC•TCCGGCTCGTATAGTGCTAGC
    nnnnnnnnnnnnnnnnnnnnnnnnTTGACAnnnnnnnn•nnnnnnnnnTATAATnnnnn
                            -35        spacer        -10
```

Fig S6

a

b

c

position relative to mScarlet-I start

Position of CRISPRtOE transposon in Landing Pad relative to mScarlet-I start (LZ1 guide)														
	# insertions analyzed	RL vs LR orientation		Position relative to ATG start (% of total RL insertions)										
	total RL and LR	%RL	%LR	-35	-34	-33	-32	-31	-30	-13	-2	+19	+108
Eco	48	87.5	12.5	21.4	2.4	71.4	0.0	0.0	4.8	0.0	0.0	0.0	0.0
Aba	12	100.0	0.0	0.0	0.0	66.7	0.0	0.0	0.0	8.3	8.3	8.3	8.3
Ecl	8	87.5	12.5	28.6	0.0	71.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Kpn	10	90.0	10.0	22.2	0.0	77.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pae	12	91.7	8.3	36.4	0.0	63.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ppu	12	83.3	16.7	10.0	10.0	80.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Son	11	100.0	0.0	36.4	9.1	45.5	0.0	0.0	9.1	0.0	0.0	0.0	0.0
Zmo	48	91.7	8.3	0.0	2.3	95.5	2.3	0.0	0.0	0.0	0.0	0.0	0.0

Fig S7

a

CRISPRtOE Screen (*murA*-fosfomycin)

Relative fitness (log2-fold change T0 vs. T2)

~52X enrichment

| murA | lacZ | murA | lacZ |
| No Drug | | Fosfomycin | |

b

~155 bp from promoter to L end of transposon

N bp insertion upstream of ATG

155 + N bp from promoter to ATG

c   Trimethoprim MIC (0.002-32 µg/ml)

CRISPRtOE *folA* isolates

WT (sJMP163)   (-20bp) (sJMP10712)   (-29bp) (sJMP10709)   (-45bp) (sJMP10716)   (-80bp) (sJMP10714)

A   B   A   B   A   B   A   B   A   B

d   Fosfomycin MIC (0.064-1024 µg/ml)

CRISPRtOE *murA* isolates

WT (sJMP163)   (-29bp) (sJMP10718)   (-33bp) (sJMP10717)   (-46bp) (sJMP10720)   (-52bp) (sJMP10719)

A   B   A   B   A   B   A   B   A   B

Fig S8

a

Comparing Replicates With or Without Promoter H − Gene Medians

b

Comparing TMP vs DMSO With or Without Promoter H − Gene Medians

c

FC TMP vs DMSO Comparing Gene Medians for CRISPRtOE With and Without Promoters

Fig S9

Fig S10

a

b

c