1

Intermittent protein restriction elevates food intake and plasma ghrelin in male mice

Volcko, K.L[a], Taghipourbibalan, H[a]., & McCutcheon, J.E.[a]

[a] Department of Psychology, UiT the Arctic University of Norway, Tromsø, Norway

Correspondence

James E. McCutcheon, Department of Psychology, UiT The Arctic University of Norway, Tromsø, Norway

Tel: +47 7764 4593, Email: j.mccutcheon@uit.no

Declaration of Interest

None

Funding

This work was supported by a Tromsø Research Foundation Starting Grant to JEM (19-SG-

JMcC).

Keywords

Protein restriction, body weight, body composition, food intake, ghrelin, sex differences

Abstract

Low-protein diets affect body weight, body composition, food intake, and food preferences in mice.

Furthermore, single periods of protein restriction can have lasting effects on these parameters.

We sought to examine the effect of multiple, short, bouts of protein restriction, relative to long-term

maintenance on either a control (NR) or protein-restricted (PR) diet. We found that male mice

experiencing intermittent protein restriction (IPR) were indistinguishable from NR mice in terms of

body weight and composition, but had food intake and plasma ghrelin as high as mice on PR diet,

even when they were returned to control diet. This was not found in female mice. The results of

this experiment highlight the importance of diet history on food intake and ghrelin levels in male

mice, and the difference in how PR diet affects male and female mice.

1. Introduction

Low-protein diets have wide-ranging effects on physiology and ingestive behavior. In male mice, low-protein diets often lead to a reduction in body weight, relative to mice eating diets that are sufficient in protein (Fontana et al., 2016; Green et al., 2022; Hill et al., 2019, 2022; Laeger et al., 2014; Larson et al., 2017; Trautman et al., 2023; Wu et al., 2021), despite increasing food intake (Fontana et al., 2016; Green et al., 2022; Hill et al., 2019; Laeger et al., 2014; Trautman et al., 2023; Volcko & McCutcheon, 2022). This mismatch appears due to increased energy expenditure observed in mice on low-protein diets (Hill et al., 2019, 2022; Wu et al., 2021). Low-protein diets also affect body composition. Some studies report a reduction in both fat and lean mass (Fontana et al., 2016; Hill et al., 2022; Larson et al., 2017; Trautman et al., 2023; Wu et al., 2021), some in only lean mass (Green et al., 2022), while others have found increases in fat mass (Simpson et al., 2013; Sørensen et al., 2008). Female mice have been studied far less frequently, but low-protein diets do not seem to have as large of an effect, if any, on body weight or body composition (Blais et al., 2018; Green et al., 2022; Larson et al., 2017; Volcko & McCutcheon, 2022); some studies do not find an effect of low-protein diet on food intake in female mice (Green et al., 2022; Larson et al., 2017) while others find increased intake (Blais et al., 2018; Volcko & McCutcheon, 2022).

Food choice is different in mice eating diets sufficient in protein compared to those eating a low-protein diet. Specifically, low-protein diets cause rodents to prefer sources of protein, such as the milk protein casein, over sources of carbohydrate, such as maltodextrin (Chiacchierini et al., 2021; Hill et al., 2019; Murphy et al., 2018; Volcko & McCutcheon, 2022). Despite differences between males and females in metabolic responses to low-protein diets, we have recently shown that this heightened protein preference is nearly identical in male and female mice (Volcko & McCutcheon, 2022).

The effects of low-protein diets can persist long after the animals are eating a normal diet again. For example, offspring of dams fed a low-protein diet during pregnancy and lactation have higher food intake even at a year of age (Qasem et al., 2016), and rats fed a low-protein diet during adolescence or early adulthood and a normal-protein diet thereafter, show elevated food intake and higher fat mass in adulthood than rats who did not experience a period of protein restriction (de Oliveira et al., 2013; Malta et al., 2014). The same group, however, did not find an effect when the diet manipulation occurred later in adulthood (Malta et al., 2016), indicating that the age of the rats was critical in determining the long-term effects of a period of protein restriction.

If a single period of protein restriction, in some developmental windows, can affect later behavior and body composition, one may also wonder if repeated bouts of protein restriction have

an effect. This was explored in a recent study by Torres et al. (2022), in which individual male

mice alternated between control and low-protein diets, with varying times on each diet (from 3 - 9

weeks). This study found that mice regulated body weight and food intake depending on which

diet they were eating (i.e. eating more but gaining less weight when on the low-protein diet

compared to the control diet) (Torres et al., 2022). Because this was a single-case experimental

design, however, it did not examine if repeated periods of protein restriction would have a long-

term effect on body weight and composition compared to if no restriction was experienced. We

were curious to know what would happen to food intake, body weight and composition, and ghrelin

levels if mice alternated between control and protein-restricted diets every seven days, and how

this would compare to mice with continuous access to either NR or PR diet.

2. Methods

2.1 Animals

The experiment was conducted on two cohorts of mice. The first cohort consisted of forty-

two adult male C57BL/6NRj mice, and the second cohort of thirty-six female C57BL/6NRj mice (6

– 8 weeks old). All mice were purchased from Janvier (France) and housed in a temperature- and

humidity-controlled room, maintained on a 12:12 light:dark cycle (lights-on at 07:00). Two mice

shared each cage, separated by a perforated divider that allowed visual, olfactory, and auditory

communication but prevented major physical contact, and allowed individual measurements of

food intake. Water and food were available *ad libitum*. A third of the mice were fed a control diet

(20% casein; D11051801; Research Diets), a third of the mice an isocaloric protein-restricted diet

(5% casein; D15100602; Research Diets), and a third of the mice alternated between the two

diets. The three groups, each with 12-14 mice, were therefore the non-restricted controls (NR),

the chronically protein-restricted (PR), and the intermittently protein-restricted (IPR). All mice were

fed the control diet for a week at the start of the experiment (Fig 1A). Both mice from each cage

experienced the same diet condition. All animal care and experimentation followed the EU

directive 2010/63/EU for animal experiments, and was approved by the National Animal Research

Authority in Norway.

2.2 Body weight and food intake

Body weight was measured three days per week. For male mice, food intake was

measured two days per week: one day immediately before the IPR group changed diets, and one

day immediately after the IPR diets were switched. These two measurements were averaged for

an average intake per week. For female mice, food intake was measured weekly.

2.3 Body composition

Body composition was assessed at baseline (when all mice were on control diet) and after 8 weeks (at the end of the experiment). Mice were weighed early in the light phase, and body composition examined 0-6 hours later in an echoMRI-700 Body Composition Analyzer (echoMRI, Houston, TX, USA), with an equal number of mice from each diet condition measured earlier and later in the day.

2.4 Plasma levels of ghrelin

At the end of the experiment, plasma was collected, with half the mice in each diet group killed at each of two time points. In the "fasted" condition, plasma was collected at the beginning of the dark phase / end of the light phase with food removed 4-6 hours beforehand. In the "fed" condition, plasma was sampled at the beginning of the light phase / end of the dark phase. At the time of plasma collection, mice in the IPR group had switched back to NR diet 7 days earlier. Mice were killed by decapitation under isoflurane anesthesia and trunk blood was collected in K_2 EDTA-coated tubes (16.444.100, Sarstedt) with 1 mg/ml AEBSF (Merck SBR00015) and placed on ice. Blood was then centrifuged, within an hour of collection, for 10 minutes at 3000 rpm and 4°C (Mikro 220R centrifuge). Plasma was separated, acidified with hydrochloric acid to a concentration of 0.05N, and frozen at -70°C until analysis. Active ghrelin was assessed by ELISA (Millipore EZRGRA-90K) on a ClarioStar microplate reader (BMG Labtech).

2.5 Data analysis

Statistical analyses were conducted using JASP (https://jasp-stats.org/). Post-hoc tests were all Tukey corrected.

Body weight per week for each mouse was an average of three measurements. A mixed model ANOVA was conducted, with Group (NR, PR, or IPR) as between-groups factor and Time as within-groups factor. Body composition measures were mixed-design ANOVA with Group and Time (baseline vs end of experiment) as factors.

Food intake per week for each mouse was an average of two measurements for males, and a single weekly measurement for females. Values over 5 g per day, or under 1 g per day, were excluded as they were almost certainly erroneous. There were three occasions in which an estimated value for weekly food intake was used, so that we did not lose data for the entire mouse. In the female cohort, food intake for a single PR mouse for week 4 was estimated by averaging the weekly intake of this mouse on all other weeks. This was similar to the approach taken in the male cohort for a single NR mouse during week 2. In the male cohort a PR mouse had its baseline

intake estimated by using the average intake of all the other mice. The average food intake for

each mouse on weeks when the IPR group was on NR, and when the IPR group was on PR, were

compared between groups. A mixed model ANOVA was conducted, with Group as between-

groups factor and Type of Week (IPR on NR diet, or IPR on PR diet) as within-groups factor.

Plasma ghrelin was analyzed as a factorial ANOVA, with Group and Condition as factors.

Due to the number of samples able to be run on a single ELISA plate, 3 female mice (1 NR, 1 PR,

1 IPR, all from the "fed" condition) were not included. One mouse (male NR fasted) was two

standard deviations above the group mean and was therefore considered an outlier and removed

from the analysis.

3. Results

3.1 Body weight

In males, body weight, as expected, increased over time as evidenced by a main effect of

Time (Fig 1B; $F_{8, 312} = 357.159$, $p < 0.001$). There was also a main effect of Group ($F_{2, 39} = 4.425$,

$p = 0.019$), as well as a Time x Group interaction ($F_{16, 312} = 22.177$, $p < 0.001$). Post-hoc tests on

the interaction revealed that by week 6, IPR mice were significantly heavier than PR mice. By

week 7, NR mice were also heavier than PR mice, and both groups were heavier than PR mice

also at week 8. At no point did IPR mice and NR mice differ in body weight.

In females, body weight also increased over time (Fig. 1C; $F_{8, 264} = 113.22$, $p < 0.001$).

There were no differences between the groups ($F_{2, 33} = 0.047$, $p = 0.954$), nor a Group x Time

interaction.

3.2 Body composition

In males, fat mass increased between baseline and after eight weeks on experimental

diets (Fig 2A; main effect of Time, $F_{1, 39} = 305.706$, $p < 0.001$). There was also a significant main

effect of Group ($F_{2, 39} = 22.91$, $p < 0.001$), and a Group x Time interaction ($F_{2, 39} = 9.813$, $p < 0.001$).

Post-hoc tests on this interaction indicated that although each of the groups had a higher fat mass

at the end of the experiment than at the beginning, PR mice had significantly less fat mass at the

end than either NR mice or IPR mice; NR and PR mice did not differ from one another (Fig 2E).

Analysis of lean mass also showed a significant main effect of Time (Fig 2C; $F_{1, 39} = 50.109$, $p <$

0.001), but no main effect of Group ($F_{2, 39} = 2.178$, $p = 0.2$). The Time x Group interaction was

significant ($F_{2, 39} = 8.5$, $p < 0.001$), and post-hoc tests showed that only the NR mice and IPR mice

gained lean mass over the eight weeks of the experiment (Fig 2G). At the end of the experiment

IPR mice had a higher lean mass than did PR mice, but PR mice and NR mice did not differ in

lean mass.

In females, fat mass increased over time (Fig 2B; $F_{1, 33}$ = 79.768, p < 0.001) but there was

no main effect of Group ($F_{2, 33}$ = 0.644, p = 0.532) nor a Group x Time interaction (Fig 2F; $_{2, 33}$ =

1.058, p = 0.359). A similar pattern of results was found for lean mass, such that lean mass was

higher at the end of the experiment than at baseline (Fig 2D; $F_{1, 33}$ = 329.289 p < 0.001), but there

were no group differences ($F_{2, 33}$ = 0.529, p = 0.722) nor Group x Time interaction (Fig 2H; $F_{2, 33}$ =

1.359, p = 0.271).

Figure 1: Schematic of experimental groups, and body weight. After a week of baseline measurements while on a normal protein diet (NR), some mice continued eating only this diet, some were given a protein-restricted diet (PR), and others alternated between the two diets (IPR) (A). Over the course of the experiment, body weight increased in male (B) and female (C) mice. Male mice on PR diet weighed less than IPR mice from week 6 and onwards, and less than those NR diet on weeks 7 and 8. Groups did not differ in body weight in female mice. Data are mean +/- SEM. Error bars too small to be visible. * PR differs from IPR, # PR differs from NR.

Figure 2: Body composition. Fat mass in male (A) and female (B) mice increased over time. Lean mass was only increased in NR and IPR males (C), but in all groups in female mice (D). Change in fat mass between start and end of the experiment was higher in NR and IPR males than PR males (E), with no differences between groups in females (F). Change in lean mass was also higher in NR and IPR males than PR males (G), with again no differences in females (H). Bars are mean and circles show individual data points. * p < 0.05 vs PR mice.

3.3 Food intake

In males, average food intake consumed each week was different across time (Fig 3A; $F_{8, 312}$ =

7.565, p < 0.001) and between groups ($F_{2, 39}$ = 27.66, p < 0.001), and there was a significant Time

x Group interaction ($F_{16, 312}$ = 2.643, p < 0.001). Post-hoc testing revealed that NR mice ate less

food than did PR mice on weeks 3, 4, 6, and 8, and less than IPR mice did on weeks 6, 7, and 8;

IPR mice were eating NR diet on weeks 6 and 8. Furthermore, we compared the average intake

of mice during weeks in which the IPR group was on NR diet or PR diets (Fig 3C). Here, we found

that intake did not differ by type of week (IPR mice on NR diet vs. PR diet) ($F_{1, 39}$ = 0.004, p =

0.947), but there was a significant main effect of Group ($F_{2, 39}$ = 2.272, p < 0.001) and a Type of

Week x Group interaction ($F_{2, 39}$ = 4.507, p = 0.017). As expected, neither NR mice nor PR mice

ate different amounts of their respective diets depending on whether IPR mice were eating NR or

PR diet, but, interestingly, on average food intake also did not differ in IPR mice depending on

Figure 3: Food intake. Male PR mice ate significantly more than NR mice on weeks 3,4,6 and 8; IPR male mice are more than NR mice on the last three weeks of the experiment (A). In female mice, PR mice ate more than NR on weeks 1 and 6, while IPR ate more than PR on week 8 (B). Average intake on weeks in which IPR was on NR diet or PR diet did not differ in any male groups, but PR and IPR males ate more than NR males regardless of week type (C). In female mice, IPR mice ate more when on PR diet than NR diet; their intake was well-matched to that of the mice on the chronic diets (D). In A and B, data are mean +/- SEM. Error bars are too small to be visible. In C and D, bars are mean and circles show individual data points. ¤ NR differs from PR, * PR differs from IPR, + IPR differs from NR, lower-case letters indicate p < 0.05 (e.g., a is different from b, b* p = 0.05 difference vs NR mice when IPR mice are on PR diet).

which diet they were currently consuming. Regardless of which diet they were eating, IPR mice ate the same amount as PR mice and more than NR mice.

In females, food intake by week had significant main effects of Time (Fig 3B; $F_{8, 264}$ = 15.532, p < 0.001) and Group ($F_{2, 33}$ = 8.46, p < 0.001), as well as a significant Time x Group interaction ($F_{16, 264}$ = 3.05, p < 0.001). Post-hoc tests on the interaction revealed that NR mice and PR mice differed in intakes on weeks 1 and 6, while PR mice and IPR mice differed in intakes on week 8. Average intake on weeks when IPR mice were on PR diet or NR diet similarly had main effects of Type of Week (Fig 3D; $F_{1, 33}$ = 66.354, p < 0.001) and Group ($F_{2, 33}$ = 9.506, p < 0.001), and a Type of Week x Group interaction ($F_{2, 33}$ = 18.108, p < 0.001). Post-hoc tests revealed that intakes of NR mice and PR mice remained stable (as expected), with NR mice eating less than PR mice. IPR mice, on the other hand, ate significantly more when on PR diet than on NR diet. When IPR mice were on NR diet their intake was the same as NR mice and lower than that of PR mice; when IPR mice were on PR diet their intake was the same as PR mice and there was a trend (p=0.053) towards their intake being higher than NR mice.

3.4 Plasma ghrelin

Figure 4: Plasma ghrelin in male mice. Male on PR diet, and IPR mice currently eating NR diet, both had higher plasma ghrelin than did NR mice. Bars are mean and circles show individual data points, lower-case letters indicate p < 0.05 (e.g., a is different from b).

Plasma ghrelin levels in a fed and fasted state were analyzed with repeated measures ANOVA. We found a main effect of Group ($F_{2, 35}$ = 8.041, p = 0.001) and no effect of Condition ($F_{1, 35}$ = 0.529, p = 0.472), nor an interaction ($F_{2, 35}$ = 0.286, p = 0.753). Post-hoc tests revealed that both PR mice and IPR mice had higher plasma ghrelin than did NR mice.

In the female mice, many samples (9 out of 34) were out of range for the ELISA. Of these, all but one were from the fed state. We therefore ran an ELISA only on the fasted state, but found no effect of Group ($F_{2, 14}$ = 2.167, p = 0.151).

4. Discussion

Mice experiencing intermittent protein restriction did not show an intermediate phenotype between that of NR and PR mice. In male IPR mice, IPR mice were indistinguishable from NR control mice in terms of body weight and body composition. One reason could be that the average level of protein restriction was less in the IPR group. They spent half the time eating a 5% protein diet, and half the time eating a 20% protein diet, so on average were eating 12.5% protein. This may be sufficient for maintaining growth, although it is below the 13.6% that has been found to support growth and reproductive health in mice (Goettsch, 1960). This is important to consider because low-protein diets (and especially those low in certain amino acids) have been suggested as ways of improving metabolic health (for example, (Fontana et al., 2016; Rang et al., 2021). The present experiment highlights that at least some of these benefits may be contingent on long-term use of the diet or avoiding frequently switching between a low and normal protein diet. Female mice, regardless of group, did not differ in body weight or composition.

Interestingly, one measure in which intermittent protein restriction differed from normal diet was in food intake. Protein restriction leads to higher caloric intake per day; this has been demonstrated many times in male mice (Fontana et al., 2016; Green et al., 2022; Hill et al., 2019; Laeger et al., 2014; Trautman et al., 2023; Volcko & McCutcheon, 2022) and is sometimes observed in female mice (Blais et al., 2018; Volcko & McCutcheon, 2022). In males, we found that the increase in food intake persisted even when the IPR male mice were consuming the control diet. Presumably, this persistent hyperphagia allowed the mice to compensate for a period when a reduced amount of protein was available. Understanding precise parameters underlying this compensatory behavior – e.g., how persistent is the hyperphagia and, if a choice is available, is it directed towards certain foods? – will be important for understanding any consequences for human health and eating behavior. As such, future experiments should address these questions as well as consider which parts of the brain are altered by protein restriction to promote hyperphagia of both low- and normal-protein food sources. Importantly, a recent paper has reviewed the persistence of both behavioral responses and physiological parameters such as a change in neural activity (Soto & Morrison, 2024). The authors argue that behavior and physiology are dictated not solely by current nutritional requirements but also by historical nutritional conditions.

Female mice on intermittent protein restriction did not show persistent hyperphagia, but were able to adapt their intake to whichever diet they were eating at the time. As others have pointed out, sex is an important factor in the effects of protein restriction (Green et al., 2022). It is worth noting that food intake was measured differently in male and female mice: in males, two 24-

hour measurement periods (one immediately before and one immediately after IPR changed diets)

were averaged to form a mean daily intake per week, whereas in female mice total food consumed

in a week was divided by 7 for a weekly average intake. In female mice we did measure 24-hour

intake on a few occasions, and correlated the intakes using both methods. The R^2 value (0.72 and

0.78, on two separate weeks, p < 0.001 on both occasions) demonstrates that using two methods

may add some variability, but the relationship is strong enough that we feel confident in the major

interpretation of the data: male mice on IPR show persistent high levels of intake regardless of

their diet, whereas female mice on IPR do not.

271   Ghrelin levels, surprisingly, did not differ by sampling time point – i.e., "fed" vs. "fasted" –

in our experiment. In the female cohort many samples from the "fed" state were below the

detection threshold, indicating that condition (fed vs. fasted) did affect ghrelin levels. In the male

cohort this was not the case; although this could be a sex difference, it is also likely that the minor

difference in timing was important. In females, blood was sampled at the end of the dark phase

and at the end of the light phase; in males, blood was sampled at the beginning of the light phase

and at the beginning of the dark phase. This is a small difference in clock time but perhaps a very

significant one in ghrelin secretion. We chose the time points with the assumption that ghrelin

would be high after a short fast and at the beginning of the dark phase when eating normally

occurs, and would be lower at the beginning of the light phase after the mice had experienced ad

libitum overnight access to their respective diets. We did not find this, but the results of this

experiment are nonetheless intriguing. In males there was a main effect of Diet that showed higher

plasma ghrelin in PR mice than in NR mice, similar to what is seen in rats (Chaumontet et al.,

2018). IPR mice, interestingly, had as high plasma ghrelin as PR, despite blood sampling

occurring a week after they had returned to NR diet. Whether or not this potential difference in

ghrelin might underlie some of the increased food consumption seen in PR and IPR mice warrants

further study.

288   Female mice on protein-restricted diet were remarkably similar to their controls eating a

diet with sufficient protein, in all measures but food intake. This is striking when viewed in light of

their protein-seeking behavior. Although we did not measure protein preference in this experiment,

we have previously shown that female mice have as robust a shift in macronutrient preference as

do male mice experiencing protein restriction (Volcko & McCutcheon, 2022). The lack of a change

in body weight or composition in female mice on this diet indicates that they are able to obtain

enough protein by increasing their intake of protein-restricted diet, while males are not able to do

so. One might speculate that this would lead to a greater drive to consume protein in males than

females; in a sense, a male mouse can perhaps be seen as experiencing a higher degree of

restriction on the same diet, because its protein requirements are higher. And yet, we do not see
a commensurate increase in protein appetite. This is intriguing and suggests that the relationship
between motivation to consume protein, and protein need, is not entirely linear.

In summary, here we add support to earlier studies showing that male and female mice
respond differently when challenged with a low-protein diet with respect to body weight, body
composition, and food intake. Moreover, male mice completely compensate for repeated periods
of low-protein diet exposure by increasing food intake, both while on the PR diet and for at least a
week after return to a diet with a normal protein level. A higher level of plasma ghrelin may be at
least partially involved in this hyperphagia, although a causal link was not studied here.
306

307   **Acknowledgements**
308   The authors wish to thank the excellent animal care staff at UiT; Neoma Boardman for use of a
309   centrifuge; and Mette Kongstorp for helping with an ELISA.
310

311   **Author contributions**
312   **Volcko, K.L**: Conceptualization, Investigation, Writing – Original Draft. **Taghipourbibalan, H**:
313   Investigation. **McCutcheon, J.E**: Conceptualization, Supervision, Writing – Review and Editing.
314   All authors have approved this manuscript.
315
316

Figure legends

Figure 1: Schematic of experimental groups, and body weight. After a week of baseline measurements while on a normal protein diet (NR), some mice continued eating only this diet, some were given a protein-restricted diet (PR), and others alternated between the two diets (IPR) (A). Over the course of the experiment, body weight increased in male (B) and female (C) mice. Male mice on PR diet weighed less than IPR mice from week 6 and onwards, and less than those NR diet on weeks 7 and 8. Groups did not differ in body weight in female mice. Data are mean +/- SEM. Error bars too small to be visible. * PR differs from IPR, # PR differs from NR.

Figure 2: Body composition. Fat mass in male (A) and female (B) mice increased over time. Lean mass was only increased in NR and IPR males (C), but in all groups in female mice (D). Change in fat mass between start and end of the experiment was higher in NR and IPR males than PR males (E), with no differences between groups in females (F). Change in lean mass was also higher in NR and IPR males than PR males (G), with again no differences in females (H). Bars are mean and circles show individual data points. * $p < 0.05$ vs PR mice.

Figure 3: Food intake. Male PR mice ate significantly more than NR mice on weeks 3,4,6 and 8; IPR male mice are more than NR mice on the last three weeks of the experiment (A). In female mice, PR mice ate more than NR on weeks 1 and 6, while IPR ate more than PR on week 8 (B). Average intake on weeks in which IPR was on NR diet or PR diet did not differ in any male groups, but PR and IPR males ate more than NR males regardless of week type (C). In female mice, IPR mice ate more when on PR diet than NR diet; their intake was well-matched to that of the mice on the chronic diets (D). In A and B, data are mean +/- SEM. Error bars are too small to be visible. In C and D, bars are mean and circles show individual data points. ¤ NR differs from PR, * PR differs from IPR, + IPR differs from NR, lower-case letters indicate $p < 0.05$ (e.g., a is different from b, b* $p = 0.05$ difference vs NR mice when IPR mice are on PR diet).

Figure 4: Plasma ghrelin in male mice. Male on PR diet, and IPR mice currently eating NR diet, both had higher plasma ghrelin than did NR mice. Bars are mean and circles show individual data points, lower-case letters indicate $p < 0.05$ (e.g., a is different from b).

References

Blais, A., Chaumontet, C., Azzout-Marniche, D., Piedcoq, J., Fromentin, G., Gaudichon, C.,

Tomé, D., & Even, P. C. (2018). Low-protein diet-induced hyperphagia and adiposity are

modulated through interactions involving thermoregulation, motor activity, and protein

quality in mice. *American Journal of Physiology - Endocrinology and Metabolism*, *314*(2),

E139–E151. https://doi.org/10.1152/ajpendo.00318.2017

Chaumontet, C., Recio, I., Fromentin, G., Benoit, S., Piedcoq, J., Darcel, N., & Tomé, D. (2018).

The protein status of rats affects the rewarding value of meals due to their protein content.

Journal of Nutrition, *148*(6), 989–998. https://doi.org/10.1093/jn/nxy060

Chiacchierini, G., Naneix, F., Peters, K. Z., Apergis-Schoute, J., Snoeren, E. M. S., &

McCutcheon, J. E. (2021). Protein appetite drives macronutrient-related differences in

ventral tegmental area neural activity. *Journal of Neuroscience*, *41*(23), 5080–5092.

https://doi.org/10.1523/JNEUROSCI.3082-20.2021

de Oliveira, J. C., Lisboa, P. C., de Moura, E. G., Barella, L. F., Miranda, R. A., Malta, A., da

Silva Franco, C. C., da Silva Ribeiro, T. A., Torrezan, R., Gravena, C., & de Freitas

Mathias, P. C. (2013). Poor pubertal protein nutrition disturbs glucose-induced insulin

secretion process in pancreatic islets and programs rats in adulthood to increase fat

accumulation. *Journal of Endocrinology*, *216*(2), 195–206. https://doi.org/10.1530/JOE-12-

0408

Fontana, L., Cummings, N. E., Arriola Apelo, S. I., Neuman, J. C., Kasza, I., Schmidt, B. A.,

Cava, E., Spelta, F., Tosti, V., Syed, F. A., Baar, E. L., Veronese, N., Cottrell, S. E.,

Fenske, R. J., Bertozzi, B., Brar, H. K., Pietka, T., Bullock, A. D., Figenshau, R. S., …

Lamming, D. W. (2016). Decreased Consumption of Branched-Chain Amino Acids

Improves Metabolic Health. *Cell Reports*, *16*(2), 520–530.

https://doi.org/10.1016/j.celrep.2016.05.092

Goettsch, M. (1960). Comparative Protein Requirement of the Rat and Mouse for Growth,

Reproduction and Lactation Using Casein Diets. *The Journal of Nutrition*, *70*(3), 307–312.

https://doi.org/10.1093/jn/70.3.307

Green, C. L., Pak, H. H., Richardson, N. E., Flores, V., Yu, D., Tomasiewicz, J. L., Dumas, S. N.,

Kredell, K., Fan, J. W., Kirsh, C., Chaiyakul, K., Murphy, M. E., Babygirija, R., Barrett-Wilt,

G. A., Rabinowitz, J., Ong, I. M., Jang, C., Simcox, J., & Lamming, D. W. (2022). Sex and

genetic background define the metabolic, physiologic, and molecular response to protein

restriction. *Cell Metabolism*, *34*(2), 209-226.e5. https://doi.org/10.1016/j.cmet.2021.12.018

Hill, C. M., Albarado, D. C., Coco, L. G., Spann, R. A., Khan, M. S., Qualls-Creekmore, E., Burk,

D. H., Burke, S. J., Collier, J. J., Yu, S., McDougal, D. H., Berthoud, H. R., Münzberg, H., Bartke, A., & Morrison, C. D. (2022). FGF21 is required for protein restriction to extend lifespan and improve metabolic health in male mice. *Nature Communications*, *13*(1), 1–14. https://doi.org/10.1038/s41467-022-29499-8

Hill, C. M., Laeger, T., Dehner, M., Albarado, D. C., Clarke, B., Wanders, D., Burke, S. J., Collier, J. J., Qualls-Creekmore, E., Solon-Biet, S. M., Simpson, S. J., Berthoud, H. R., Münzberg, H., & Morrison, C. D. (2019). FGF21 Signals Protein Status to the Brain and Adaptively Regulates Food Choice and Metabolism. *Cell Reports*, *27*(10), 2934-2947.e3. https://doi.org/10.1016/j.celrep.2019.05.022

Laeger, T., Henagan, T. M., Albarado, D. C., Redman, L. M., Bray, G. A., Noland, R. C., Münzberg, H., Hutson, S. M., Gettys, T. W., Schwartz, M. W., & Morrison, C. D. (2014). FGF21 is an endocrine signal of protein restriction. *Journal of Clinical Investigation*, *124*(9), 3913–3922. https://doi.org/10.1172/JCI74915

Larson, K. R., Russo, K. A., Fang, Y., Mohajerani, N., Goodson, M. L., & Ryan, K. K. (2017). Sex differences in the hormonal and metabolic response to dietary protein dilution. *Endocrinology*, *158*(10), 3477–3487. https://doi.org/10.1210/en.2017-00331

Malta, A., de Moura, E. G., Ribeiro, T. A., Tófolo, L. P., Abdennebi-Najar, L., Vieau, D., Barella, L. F., de Freitas Mathias, P. C., Lisboa, P. C., & de Oliveira, J. C. (2016). Protein-energy malnutrition at mid-adulthood does not imprint long-term metabolic consequences in male rats. *European Journal of Nutrition*, *55*(4), 1423–1433. https://doi.org/10.1007/s00394-015-0960-8

Malta, A., de Oliveira, J. C., da Silva Ribeiro, T. A., Tófolo, L. P., Barella, L. F., Prates, K. V., Miranda, R. A., Elmhiri, G., da Silva Franco, C. C., Agostinho, A. R., Trombini, A. B., Pavanello, A., Gravena, C., Abdennebi-Najar, L., & Mathias, P. C. de F. (2014). Low-protein diet in adult male rats has long-term effects on metabolism. *Journal of Endocrinology*, *221*(2), 285–295. https://doi.org/10.1530/JOE-13-0473

Murphy, M., Peters, K. Z., Denton, B. S., Lee, K. A., Chadchankar, H., & McCutcheon, J. E. (2018). Restriction of dietary protein leads to conditioned protein preference and elevated palatability of protein-containing food in rats. *Physiology and Behavior*, *184*(December 2017), 235–241. https://doi.org/10.1016/j.physbeh.2017.12.011

Qasem, R. J., Li, J., Tang, H. M., Pontiggia, L., & D'mello, A. P. (2016). Maternal protein restriction during pregnancy and lactation alters central leptin signalling, increases food intake, and decreases bone mass in 1 year old rat offspring. *Clinical and Experimental Pharmacology and Physiology*, *43*(4), 494–502. https://doi.org/10.1111/1440-1681.12545

Rang, Y., Ma, S., Yang, J., Liu, H., Suzuki, K., & Liu, C. (2021). A low-protein high-fat diet leads
to loss of body weight and white adipose tissue weight via enhancing energy expenditure in
mice. *Metabolites*, *11*(5). https://doi.org/10.3390/metabo11050301
Simpson, S. J., Huang, X., Hancock, D. P., Gosby, A. K., McMahon, A. C., Solon, S. M. C., Le
Couteur, D. G., Conigrave, A. D., & Raubenheimer, D. (2013). Effects of dietary protein to
carbohydrate balance on energy intake, fat storage, and heat production in mice. *Obesity*,
21(1), 85–92. https://doi.org/10.1002/oby.20007
Sørensen, A., Mayntz, D., Raubenheimer, D., & Simpson, S. J. (2008). Protein-leverage in mice:
The Geometry of macronutrient balancing and consequences for fat deposition. *Obesity*,
16(3), 566–571. https://doi.org/10.1038/oby.2007.58
Soto, P. L., & Morrison, C. D. (2024). Persistent Behavioral and Physiological Effects of Dietary
Protein Restriction. *BioRxiv*.
Torres, F., Khan, S., Fernandez-Kim, S. O., Spann, R., Albarado, D., Wagner, T. J., Morrison, C.
D., & Soto, P. L. (2022). Dynamic effects of dietary protein restriction on body weights, food
consumption, and protein preference in C57BL/6J and Fgf21-KO mice. *Journal of the*
Experimental Analysis of Behavior, *117*(3), 346–362. https://doi.org/10.1002/jeab.745
Trautman, M. E., Braucher, L. N., Elliehausen, C., Zhu, W. G., & Zelenovskiy, E. (2023).
Resistance exercise protects mice from protein-induced fat accretion. 1–30.
Volcko, K. L., & McCutcheon, J. E. (2022). Protein preference and elevated plasma FGF21
induced by dietary protein restriction is similar in both male and female mice. *Physiology*
and Behavior, *257*(October), 113994. https://doi.org/10.1016/j.physbeh.2022.113994
Wu, Y., Li, B., Li, L., Zheng, Y., Douglas, A., Speakman, J. R., Wu, Y., Li, B., Li, L., Mitchell, S.
E., Green, C. L., Agostino, G. D., & Wang, G. (2021). *Article Very-low-protein diets lead to*
reduced food intake and weight loss , linked to inhibition of hypothalamic mTOR signaling ,
in mice ll Article Very-low-protein diets lead to reduced food intake and weight loss , linked
to inhibition of hypothalamic. 888–904. https://doi.org/10.1016/j.cmet.2021.01.017
443
444