Tick-borne pathogens identified in *Hyalomma marginatum* (Acari: Ixodidae) collected from various vertebrate hosts in the South of France.

Bernard Célia[1,2,3,*], Pollet Thomas[1,2], Galon Clémence[4], Joly Kukla Charlotte[1,2,4], Cicculli Vincent[5], Falchi Alessandra[5], Grech-Angelini Sebastien[6], Poli Paul-Eric[7], Bastien Matthieu[3], Combes Benoit[3], Moutailler Sara[4], Holzmuller Philippe[1,2,§,] Vial Laurence[1,2,§]

CIRAD, UMR ASTRE, F-34398 Montpellier, France

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France

French Establishment for Fighting Zoonoses (ELIZ), 54220 Malzéville, France

ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale, Maisons-Alfort, F-94700, France

Bioscope Corse Méditerranée unit UR7310, Faculté de Sciences, Campus Grimaldi, Université de Corse, 20250 Corte, France

Groupement Technique Vétérinaire Corse

INRAE, LRDE, Corte, France

*Corresponding author
Correspondence: celia.bernard@cirad.fr ; celia.b2@orange.fr
§ Joint last authors

ABSTRACT

Hyalomma marginatum is a common ectoparasitic tick of ungulates, lagomorphs, insectivores, ground-foraging birds, observed in Corsica for decades, but whose permanent establishment in mainland France is very recent. This species is known to be one of the main vectors of the Crimean-Congo Hemorrhagic Fever virus, but also of various parasitic, bacterial or viral pathogens. In this study, we investigated the molecular infection rates of numerous tick-borne pathogens in *H. marginatum* ticks mainly sampled on horses, and occasionally on other animal species, from the French Mediterranean rim and Corsica between 2016 and 2020. In total, 1, 195 DNA and RNA purified from individual ticks or pools of ticks were screened for 26 microbial genera or species (viruses, bacteria and parasites), using a high-throughput microfluidic real-time PCR system (BioMark™ dynamic array system, Standard Biotools). For individual ticks and pooled ones, respectively, the most prevalent tick-borne microorganisms were *Francisella*-like endosymbionts at 97.0% and 96.8%, followed by *Rickettsia aeschlimannii* (76.4% and 96.4%), *Theileria* spp. and *Theileria equi* (3,5% and 0%; 1,9% and 5,8%), *Anaplasma phagocytophilum* (3.7% and 6.7%), and West Nile virus (0.1% and 0.4%). *Babesia occultans* (0.9%), *Ehrlichia minasensis* (0.3%), and *Coxiella*-like endosymbionts (0.1%) were only detected in individual ticks. Our study provides an overview of the diversity of microorganisms and tick-borne pathogens detected in the invasive tick *H. marginatum* in

Corsica and the continental departments of the Mediterranean rim. Our study opens up new research perspectives on the epidemiology of tick-borne pathogens carried by *H. marginatum* and on the associated public and veterinary health risks.

Keywords: Tick; *Hyalomma marginatum*; tick-borne pathogens; endosymbionts, microfluidic PCR

Introduction

Worldwide, ticks are considered to be the first vectors of pathogens responsible for animal diseases and second behind mosquitoes for those affecting humans (Jongejan et Uilenberg 2004; Colwell, Dantas-Torres, et Otranto 2011). In France, about forty species of ticks are identified on the territory, most of them transmitting pathogens responsible for important public and veterinary health problems (Boulanger et McCoy 2017; PÉREZ-EID 2007). This specifically includes *Ixodes ricinus*, the primary vector for the agents of Lyme borreliosis, *Borrelia burgdorferi* sensu lato, as well as ticks from the genera *Dermacentor* or *Rhipicephalus* responsible for the transmission of equine piroplasmosis (Gray 1991; Nadal, Bonnet, et Marsot 2022). Recently, Vial et al. (2016) provided strong evidence for the presence of reproducing populations of the exotic tick *Hyalomma marginatum* in parts of southern France. While this tick species has been present in Corsica (a Mediterranean island of southern France) for several decades, it seems to be now established on the Mediterranean coastline of mainland France in many areas such as Pyrénées-Orientales, Aude, Hérault, Gard, Ardèche, Drôme and Var (Bah et al. 2022). This tick specieshas been also detected since many years in Spain (Palomar et al. 2016) and Italy (Satta et al. 2011), showing a possible continuum for the sanitary risks associated to *H. marginatum* in southwestern Europe. As a "generalist" tick, *H. marginatum* presents a large panel of vertebrate hosts, ranging from lagomorphs and birds for the immature stages to large domestic and wild ungulates for the adult ones (Bernard et al., 2022). It is a two-host tick, meaning that larvae and nymphs feed on the same individual, while the nymph detaches and moults into an adult that seeks out a new host for blood feeding and mating. This diversity of hosts expands the pathways for tick introduction via natural or human-driven animal movements and is believed to increase opportunities for pathogen transmission and possible spillover. Ticks of the genus *Hyalomma* are involved in the transmission of many human and animal pathogens in Eurasia and Africa (Bonnet et al. 2023). They can in particular carry parasites and bacteria such as *Theileria* spp., *Babesia* spp. and *Anaplasma* spp., responsible for equine piroplasmosis or bovine anaplasmosis, with often non-specific clinical signs (e.g. hyperthermia, asthenia, abortion...) in cattle or sheep (Wise et al. 2013, Rothschild et al. 2013, Ristic et al., 1981). Other infectious bacteria such as *Coxiella burnetii*, and *Rickettsia* spp. including *R aeschlimannii* (Bakheit et al. 2012; Kilicoglu et al. 2020; Orkun 2022), or viruses such as Bhanja (Genus Bandavirus) and West Nile (Genus Flavivirus) (Formosinho et Santos-Silva 2006; Kolodziejek et al. 2014), can be carried by *H. marginatum*. Additionally, Orthonairoviruses such as the emerging and highly contagious Crimean Congo Hemorrhagic Fever virus (CCHFV) (Bakheit et al. 2012) can be transmitted by this tick spcies. In the Mediterranean basin, two species of *Hyalomma* (*H. marginatum* and *H. anatolicum*) are used to be considered as the main vectors of CCHFV (Perveen et Khan 2022). However, viral RNA has been recently detected in *H. lusitanicum* collected from deer and wild boar in Spain, in higher proportion than in *H. marginatum*, suggesting major role of this additional tick in CCHFV transmission in this country (Moraga-Fernández et al. 2021). In France, based on our knowledge on local tick communities, we considered *H. marginatum* as the main candidate tick species likely to transmit CCHFV (Bernard et al. 2022). In 2023, this was confirmed by the first detection of CCHFV in H. marginatum ticks from Pyrénées-Orientales (Bernard, 2024). Considering the large geographic distribution of *H. marginatum*, from Spain to North Africa and India (Fernández-Ruiz et Estrada-Peña 2021), this questions about local adaptations in the abilities developed by this tick species to maintain and transmit all, or part of the pre-cited pathogens.

The objective of this study was thus to describe the panel of pathogens and/or endosymbiotic microorganisms in *H. marginatum* ticks locally collected in the South of France, mainland and Corsica island. This work was part of a national surveillance program developed a few years ago to assess the risk of emergence of exotic diseases including tick-borne diseases in France, under the aegis of the French Ministry of Agriculture. In Corsica, first investigations were recently conducted on tick species collected from different hosts, especially cattle, and reported in *H. marginatum* high prevalence of *Francisella*-like endosymbionts, *R. aeschlimannii* and in a less extend *Candidatus Ri. Barberiae*, as well as the presence of *Anaplasma marginale* and *A. phagocytophylum* (V. Cicculli et al. 2019; Grech-Angelini et al. 2020). These studies also allowed the detection of new viruses in *H. marginatum*, for example a Pseudocowpox virus of which the zoonotic potential remains unknown (V. Cicculli et al. 2019). On mainland Francewhere *H. marginatum* was not yet established only 15 years ago, such investigation has never been done. Keeping in mind that genomic detection of pathogens does not imply a mandatory vector competence of the tick for the considered microorganisms (Bonnet et al. 2023), this study aimed to assess potential vector

competence by combining pathogen detection with an understanding of *H. marginatum*'s host preferences
and the susceptibility of these hosts to the identified pathogens. The aim was to know whether *H.*
marginatum ticks arrived with their native associated pathogens usually described in endemic
Mediterranean areas including Corsica, and may result in the introduction of new pathogens in France.
Another alternative would be that *H. marginatum* can become a new vector for local French tick-borne
pathogens already present on the territory, with the risk of changing the local transmission dynamics of
such pathogens due to its distinct seasonal activity from other local tick species.
111

Methods

Study area, period and hosts for tick collection

113
114
Among all the *H. marginatum* ticks analyzed in this study, 1,711 came from a large-scale tick sampling
that was conducted on horses (considered as the likely host of *H. marginatum* adult stages), during spring
(the activity period of *H. marginatum* adult stages), in southern France (mainland and Corsica), between
2016 and 2020. This was carried out to determine and explain the distribution of the newly established tick
H. marginatum on the French territory (Bah et al. 2022). Eight French departments forming a link from
Spain to Italy, which are two countries where *H. marginatum* is known to be already present, plus the two
departments of Corsica were sampled. *H. marginatum* ticks were detected in 56 and 38 equestrian
structures in mainland France and in Corsica, respectively. On the mainland, a reference site in Pompignan,
where *H. marginatum* is abundant and monitored for its dynamics of infestation on horses, was also
sampled each year and 173 ticks of the 1,711 ticks came from this site. In Corsica, each year, the whole
island was visited whereas on the mainland, most of *H. marginatum* were obtained in 2017, except for
northern Gard and Ardèche that were investigated in 2019 (Figure 1). Among the 1,711 ticks, 900 collected
in Corsica in 2019 and 2020 were grouped into 222 pools of 1 to 10 individuals according to different
parameters of sex or hosts (Cicculli et al., 2022) and 811 were treated individually.
In Corsica, 118 additional ticks were collected from cattle examined in three farms located in the
northern part of the island, in 2019.
Additionally, 7 and 37 *H. marginatum* ticks were obtained by punctual samplings in Hérault and Gard
on wild boars in 2016 and birds in 2017, respectively. Ticks from birds were all immature stages, and were
mostly associated to the resident avian fauna of Pompignan (Vial et al. 2016). Thus, taking all host species
together, we worked on 973 ticks treated individually and 222 tick pools (1,195 samples).
135

Figure 1: Study area overview : geographic distribution and host diversity. On this map, sites are represented by the host species from which the ticks were collected (horses, cattle, wildboars and birds).

During any tick samplings, animals were examined on the whole body and all tick species were collected. Back from the field, they were characterized under binocular magnifiers using morphological identification keys (PÉREZ-EID 2007), and sorted according to species, locality, host, gender (male/female), developmental stage (larva/nymph/adult), and engorgement status. They were then frozen at -80°C for later molecular analysis. As this study focused on sanitary risks associated to the newly established *H. marginatum*, only specimens from this species were screened for microorganisms.

DNA and RNA extraction

As we looked for DNA and RNA of microorganisms in all tick samples, their treatment for molecular analysis was the same.

After a first washing in bleach for 30 sec followed by three times in distilled water for 1 min, ticks were crushed individually with three steel beads (2.8mm, OZYME, France) in 300μL DMEM (Eurobio Scientific, France) supplemented with 10% fetal calf serum (Dutscher, France), using the homogenizer Precellys® 24 Dual (Bertin, France) at 5500 rpm for 40s. The processing was different for samples of horses coming from Corsica as, prior to be crushed, ticks were pooled by up to 10 individuals per pool, according to locality and developmental stage (Vincent Cicculli et al. 2022).

DNA was extracted using 100μl of the homogenate of crushed ticks according to the Genomic DNA from tissue kit (Macherey-Nagel, Germany) following the manufacturer's instructions. Total DNA per sample was eluted in 50 μl of rehydration solution and stored at −20∘C until further use (Michelet et al, 2014).

Total RNA was extracted from 100μl of the homogenate of crushed ticks using the NucleoSpin RNA kit (Macherey-Nagel, Germany) according to the manufacturer's instructions. Total RNA per sample was eluted in 50 μl of RNA free water and stored at −80∘C until further use (Gondard et al., 2018).

Detection of tick-borne microorganisms' DNA or RNA

The Reverse Transcription Master Mix kit (Standard Biotools, USA) was used on all samples to obtain cDNAs from RNA viruses. The reaction was performed in 5µL of final volume, including 1µL of Reverse Transcription Master Mix, 3µL of RNase-free ultrapure water, and 1µL of extracted RNA. A thermal cycler (Eppendorf, Germany) performed the RT cycles following the program: 25°C for 5 min, 42°C for 30 min and a final step at 85°C for 5 min for a total duration of 40 min.

A pre-amplification step was performed to pre-amplify the DNA/cDNA to increase the signal of the genetic material of the microorganisms compared to those of the tick. The PreAmp Master Mix kit (Standard Biotools, USA) was used. A 0.2X pool containing the primer pairs of the targeted pathogens was made upstream, to obtain a final primer concentration of 200 nM. For one sample, the mix was composed of 1 µL of PreAmp Master mix, 1.25 µl of the 0.2X pool, and 1.5 µl of ultra-pure water. Then 1.25µL of sample (DNA/cDNA mix) was added. A negative control (ultra-pure water) was added to each plate. PCR was then performed using a thermal cycler with the following program: 95°C for 2min, 14 cycles of two steps of 95°C for 15s and a final step of 4min at 60°C.

Then, for the real time microfluidic PCR, twenty-four sets of primers and probes were used to detect predefined microorganisms specifically targeted for this study (Table 1) (Michelet et al. 2014; Gondard et al. 2018). The BioMark™ real-time PCR system (Standard Biotools, USA) was used for high throughput microfluidic real-time PCR amplification using 48.48 dynamic arrays (Standard Biotools). These chips dispense 48 PCR mixes and 48 samples into individual wells, after which on-chip microfluidics assemble PCR reactions in individual chambers prior to thermal cycling resulting in 2,304 individual reactions (Michelet et al. 2014). In a single experiment, 47 individual ticks or tick pools samples (and one negative control) can be tested.

In order to develop the a priori approach to detect pathogens in ticks using real time microfluidic PCR, we first generated a list of microorganisms (TBPs and endosymbionts) (Table 1) with the highest probability of being transmitted or carried by *H. marginatum*, according to data mining of the available literature. From two studies *(Bakheit et al. 2012; Bonnet et al. 2022)*, we identified several potentially *H. marginatum*-borne pathogens: protozoa such as *B. caballi* or *T. annulata*, bacteria such as *R. aechlimannii* and finally viruses such as Thogoto, Dhori and Bhanja viruses (RNA for these viruses has already been detected in ticks), in addition to CCHFV. We then identified microorganisms from scientific literature that mentioned the detection of pathogens in *H. marginatum*, which could be transmitted by this tick or just reflect the infectious status of the animal host on which the tick was collected, as shown in Corsica or other regions of the world: *A. phagocytophilum, A. marginale (Grech-Angelini et al. 2020), E. minasensis (Vincent Cicculli et al. 2019), T. equi (Rocafort-Ferrer et al. 2022)*, the West Nile Virus *(Formosinho et Santos-Silva 2006; Kolodziejek et al. 2014)*, the symbiotic form of *Coxiella burnetii* and *Francisella tularensis* : *Coxiella*-like endosymbiont CLE and *Francisella*-like endosymbiont FLE *(Kilicoglu et al. 2020; Selmi et al. 2019; Buysse et al. 2021; Demir et al. 2020)*. Then, some pathogens known to be transmitted by ticks of the genus *Hyalomma* but not necessarily *H. marginatum*, or even by other tick genera but with a likelihood to replicate in *Hyalomma* ticks, were also added to the chip: Alkhurma virus transmitted by *H. rufipes (Horton et al. 2016; Hoffman et al. 2018)*, as well as Nairobi sheep disease and Hazara viruses that are genetically close to CCHFV (Garrison *et al.*, 2020), *Borrelia miyamotoi* and *Bartonella henselae* that circulate in France in other tick species or even other arthropoda such as fleas *(Cosson et al. 2014; Grech-Angelini et al. 2020)*.

213 **Table 1:** Tick-borne pathogens and endosymbionts targeted by the chip of the high-throughput
214 microfluidic real-time PCR system developed specifically for the study or for specific RT-qPCR (CCHFV).

Genus / Family	Species
Bacteria	
Anaplasma	*A. marginale, A. phagocytophilum, Anaplasma* spp.
Bartonella	*B. henselae, Bartonella* spp.
Borrelia	*B. miyamotoi, Borrelia* spp.
Coxiella	*Coxiella burnetii and Coxiella-like endosymbionts*
Ehrlichia	*Ehrlichia* spp.
Francisella	*F. tularensis, Francisella-like endosymbionts*
Rickettsia	*R. aeschlimannii, Rickettsia* spp.
Protozoan	
Babesia	*B. caballi*
Theileria	*T. equi, T. annulata, Theileria* spp.
Viruses	
Bunyaviridae	Nairobi sheep disease virus
Flaviviridae	West Nile Virus, Alkhurma virus
Nairoviridae	Hazara virus, CCHFV
Orthomyxoviridae	Thogoto virus, Dhori virus
Phenuiviridae	Bhanja virus

215
216
To confirm our results of positive detection of pathogens by the microfluidic chip in field samples,
specific PCRs were performed individually on some samples with the following primers (Table 2).
Conventional PCRs using primers targeting genes or regions other than those of the BioMark™ system were
used. Amplicons were sequenced by Eurofins MWG Operon (Germany), and then assembled using BioEdit
software (Ibis Biosciences, Carlsbad). An online BLAST (National Center for Biotechnology Information) was
used to compare results with published sequences listed in GenBank sequence databases.

Table 2: Primers used to confirm each pathogen detected by the microfluidic chip, through
conventional and/or nested PCR.

Pathogen	Name	PCR type		Amorce name	Sequence
Anaplasma / *Ehrlichia* sp.	EHR	Nested	PCR1	EHR1	GAACGAACGCTGGCGGCAAGC
				EHR2	AGT AYC GRA CCA GAT AGC CGC
			PCR2	EHR3	TGCATAGGAATCTACCTAGTAG
				EHR2	AGT AYC GRA CCA GAT AGC CGC
	msp4	Conventionnal		MSP4AP5	ATG AAT TAC AGA GAA TTG CTT GTA GG
				MSP4AP3	TTA ATT GAA AGC AAA TCT TGC TCC TAT
Babesia / Theileria sp.	BTH	Nested	PCR1	BTH 18S 1st F	GTGAAACTGCGAATGGCTCATTAC
				BTH 18S 1st R	AAGTGATAAGGTTCACAAAACTTCCC
			PCR2	BTH 18S 2nd F	GGCTCATTACAACAGTTATAGTTTATTTG
				BTH 18S 2nd R	CGGTCCGAATAATTCACCGGAT
	babG	Conventionnal		BABGF2	GYY TTG TAA TTG GAA TGA TGG
				BABGR2	CCA AAG ACT TTG ATT TCT CTC
Borrelia sp.	FlaB	Nested	PCR1	FlaB280F	GCAGTTCARTCAGGTAACGG
				FlaRL	GCAATCATAGCCATTGCAGATTGT
			PCR2	flaB	GCATCAACTGTRGTTGTAACATTAACAGG
				FlaLL	ACATATTCAGATGCAGACAGAGGT
Rickettsia sp.	rsfg	Conventionnal		Rsfg877	GGGGGCCTGCTCACGGCGG
				Rsfg1258	ATTGCAAAAAGTACAGTGAACA
	rc.rompB	Nested	PCR1	Rc.rompB.4362p	GTCAGCGTTACTTCTTCGATGC
				Rc.rompB.4,836n	CCGTACTCCATCTTAGCATCAG
			PCR2	Rc.rompB.4,496p	CCAATGGCAGGACTTAGCTACT
				Rc.rompB.4,762n	AGGCTGGCTGATACACGGAGTAA

226
Finally, a specific detection of CCHFV was performed on tick samples. All RNA extracts were screened for CCHFV
by another RT-qPCR targeting a 3' non-coding region of the CCHFV genome using specific primers and probes
(Sas et al., 2018). rRT-PCR Taqman assays were performed in a final volume of 20 µL using the LightCycler 480
RNA Master Hydrolysis Probes Master Mix (Roche Applied Science, Germany) at 1 X final concentration, with 0.5
µM specific primers and 0.25 µM probes and 2 µL RNA. Negative (ultra-pure water) controls were included in
each run. rRT-PCR thermal cycling conditions were as follows: 63°C for 3 min, 95°C for 30 s, 45 cycles at 95°C for
10 s then 60°C for 30 s, followed by cooling at 40°C for 10 s.

234
235 **Statistical analysis**
236
The statistical analysis was conducted using R software (version 4.2.1, R Core Team).
In order to investigate the potential difference in the presence of *Rickettsia* between male and female
ticks, a chi-square test was performed, as a relevant method commonly used to assess associations
between categorical variables. The variables of interest were the presence of *Rickettsia* (1,0) and the
gender (male or female), and the test was conducted to determine if there was a statistically significant
association between gender and the presence of *Rickettsia*.

243
244

245 **Results**

246
247 ***1. Identification of microorganisms detected in H. marginatum***
248
249

DNA and RNA of ten of the targeted tick-borne microorganisms were identified in the 1,195 tick
samples (973 individual ticks and 222 pools of 1 to 10 ticks). Using the microfluidic method on individual
ticks, we detected 2 ticks positive for *Borrelia* spp, 3 for *A. marginale*, 33 for *A. phagocytophilum*, 201 for
Anaplasma spp, 3 for *Ehrlichia* spp, 878 for *R. aeschlimannii*, 957 for *Rickettsia* spp, 8 for *B. henselae*, 4 for
Bartonella spp., 68 for *Theileria* spp, 22 for *T. equi*, 13 for West Nile Virus, 1 for CLE and 1,160 for FLE. In
terms of tick pools, the same method identified 19 positive pools for *Anaplasma* spp., 215 pools for *R.*
aeschlimannii, 214 for *Rickettsia*. spp., 10 for *Theileria*. spp., 5 for *T. equi*, 216 for FLE and 3 for WNV. Except

for the well-known CLE and FLE (and because the pathogenic forms *C. burnettii* and *F. tularensis* were never
detected in this study), any other amplified microorganisms suspected to be pathogenic were sequenced
from some positive samples to be confirmed.

Figure 2: Proportions of *Rickettsia aeschlimannii* infections (A) and other microorganisms (B) according
to geographical sampling sites in *H. marginatum* ticks collected on horses (on the left) and cattle (only in
Corsica) (on the right). The size of the circles is proportional to the number of ticks collected per sampling
site, the color of the department (in A and B) corresponds to the samplig year.

For *Rickettsia* spp*.,* most of samples amplified specifically for *R. aeschlimannii* were also amplified in
Rickettsia spp., using the microfluigdim method. 18 amplicons (5 combining the two targets, 3 amplifying
only *R. aeschlimannii* and 10 amplifying only *Rickettsia* spp.) were sequenced. Of the 10 readable
sequences, all sequences confirmed the presence of *R. aeschlimannii*. One of these sequences has been
deposited in GenBank (AN: PP379722), showing 98.77% homology with a strain isolated from a *H.
marginatum* found in England in 2019 (AN: MT365092) and 97.85% homology with another strain detected
in Corsica in 2019 (AN: MK732478) in a *Rhipicephalus bursa* tick.
274
275 For *Anaplasma* spp., all samples amplifying *A. marginale* were also positive for *Anaplasma* spp., which
276 was not necessarily the case for *A. phagocytophilum*, for which some samples were positive for *A.*

phagocytophilum but not *Anaplasma* spp*.*, and many samples were only amplified using primers targeting *Anaplasma* spp. Nineteen amplicons representative of the four cases (*A. marginale* + *Anaplasma* spp. / *A. phagocytophilum* only / *A. phagocytophilum* + *Anaplasma* spp. / *Anaplasma* spp. only / *A. phagocytophilum* only) were sequenced, and only confirmed samples that combined the amplification of both specific and generic targets. The samples positive only for *Anaplasma* spp. were finally identified as *Devosia* spp. by sequencing confirmation PCR products, which is an ubiquitous soil bacteria that is considered as a PCR cross contaminant. At the end, two *Anaplasma* species were confirmed: *A. phagocytophilum* and *A. marginale*. The sequence of *A. phagocytophilum* (AN: PP227262) showed 99% homology with the reference sequence from an *A. phagocytophilum* strain isolated in a *Rhipicephalus microplus* tick collected on a cow in Taiwan in 2021 (AN: OL690562.1) and 99.12% with a sequence from a strain isolated in an *Ixodes ricinus* tick, in Hérault, France in 2010 (AN : OR426543). For *A. marginale*, the sequence (AN : PP218689) showed 96% homology with a reference sequence from an *A. marginale* strain isolated in a *H. schulzei* tick collected in Iran in 2019 (AN: MK310488) and 95,27 % from a strain isolated in cattle in 2017 in Spain (AN: MT664983).

For *Ehrlichia* spp*.,* two amplicons were sequenced and were confirmed as *E. minasensis*. The resulting sequence deposited in GenBank (AN: PP227261) showed 99% homology with a reference sequence (AN: NR_*148800.1)* of a strain that was isolated from the hemolymph of a *Rhipicephalus (Boophilus) microplus* tick from Brazil in 2019. Moreover, the sequence detected showed 98% homology with a strain (AN : JX629805) detected in a *Rhipicephalus microplus* tick from Brazil in 2016. The detected sequence in Corsica was 100% identical with the brazilian strain (Vincent Cicculli et al. 2019).

For *Theileria* spp., most of samples that were amplified specifically for *T. equi* were also amplified for *Theileria* spp*.*, except two samples positive only for *T.equi*, and other samples were only amplified for *Theileria* spp. No sample was amplified for *T. annulata*. Sixty six amplicons combing the three cases (*T. equi* only / *T. equi* + *Theileria* spp*. / Theileria* spp*.* only) were sequenced and confirmed the presence of *T. equi* for samples amplified for both targets as well as those amplified only for *T. equi*. All amplicons for *Theileria* spp. only produced sequences that could not allow species-specific identification despite repeated sequencing (unreadable sequences). For *T. equi,* the same sequence (AN: PP227191) was obtained for all samples confirmed and showed 99% homology with a reference sequence from a *T. equi* strain isolated in Chile in 2020 (AN: MT463613.1) and 99,81% homology with a sequence from a *Rhipicephalus bursa* tick in Corsica in 2019 (AN: MK732476.1).
Nine ticks detected positive for *Theileria* spp. only were finally confirmed by sequencing as infected by *Babesia occultans*. The sequence deposited in GenBank (AN: PP227192) showed 99% homology with a sequence of *B. occultans* isolated from a *H. excavatum* tick collected on cattle in Turkey in 2022 (AN : OM066180.1).

One individual tick and one pool of ticks from Corsica were positive for the West Nile Virus. The amplification product was sequenced (AN: PP379723) and showed 100% homology with a reference sequence from the complete genome of West Nile Virus detected from a snowy owl in Germany in 2019 (AN : LR743458.1).

3. Proportions of microorganisms in H. marginatum, according to host origin and spatial distribution

Results of individual and pooling detections are presented in Table 3.

Table 3: Microorganisms detected and confirmed in *H. marginatum* ticks, tested individually or by pools. The currently known or assumed pathogens are in black and the endosymbionts in grey.

	Individual ticks (N=973)			Pools of ticks (N=222)		
	Nb. detected	Proportion (%)	Host	Nb. detected	Proportion (%)	Host
Rickettsia aeschlimannii	743	76,4	B, C, H, W	214	95,1	H
Francisella-like endosymbiots	944	97,0	B, C, H, W	216	97,3	H
Anaplasma phagocytophilum	33	3,4	B, C, H	15	6,7	H
Anaplasma marginale	3	0,3	C	0	0	H
Ehrlichia minasensis	3	0,3	C, H	0	0	H
Coxiella-like endosymbiots	1	0,1	H	0	0	H
Theileria spp.	34	3,5	H	0	0	H
Theileria equi	17	1,8	H	13	5,8	H
Babesia occultans	9	0,9	C	0	0	H
West Nile Virus	1	0,1	C	1	0,4	H

B: birds, C: cattle, H: horses, W: wild boars

FLE was the most prevalent microorganism, as detected in more than 95% of individual ticks and pools of ticks of *H. marginatum*. *R. aechlimannii* was detected in more than 75% of the analyzed individual ticks and 95% of pools tick of *H. marginatum*. Conversely, we only identified a CLE in one tick collected on a horse in Var. Then, proportions of infection due to all the other targeted tick-borne pathogens ranged from 0.1% to 3.5% (up to 6.7% in pools of ticks). If we remove the associations between *R. aeschlimannii* that was highly prevalent and the other pathogens, the likelihood of coinfections in *H. marginatum* ticks was very low. Only 3 coinfections were observed, two ticks carrying both *B. occultans* and *A. marginale* and one tick carrying both *E. minasensis* and *A. marginale*.

Although equine piroplasmosis is commonly reported in horses from the South of France (Rocafort-Ferrer et al. 2022) and our ticks were mainly collected on horses, *T. equi* was detected in a very small number of ticks, 17 individual ticks from mainland France and 13 pools of ticks from Corsica, all collected on horses. No *B. caballi* was detected in the analyzed ticks. *B. occultans* was detected in 9 individual ticks collected only on cattle in Corsica. For bacteria, *A. marginale* was only detected in ticks collected on cattle in Corsica, while *A. phagocytophilum* was mainly found in ticks collected from horses but also on cattle and to a lesser extent in ticks collected on birds (n = 2/37). Birds were mainly infected by *R. aeschlimannii* (n = 15/37). *E. minasensis* was detected in 3 ticks, two from Corsican cattle and one from a horse on the mainland. Only one virus, the West Nile virus, was detected in one tick collected from a cow and one pool of ticks collected from a horse, both in Corsica. Conversely, *H. marginatum* ticks were all negative for CCHFV. No specific pathogens, except *R. Aeschlimannii* (n = 6/7), were found in ticks collected on wild boar. Aside from minor differences between ticks collected on horses versus cattle, and considering the significant underrepresentation of hosts other than horses, which limited statistical comparisons, the panel of microorganisms appeared to be quite similar across the different hosts.

Finally, we did not observe significant differences in the presence of most of the detected microorganisms between male and female ticks (*A. phagocytophilum* : χ^2 = 0.46 df = 1, *P* = 0.49 ; *A. marginale* : χ^2 = 1.37, df = 1, *P* = 0.24 ; *E. minasensis* : χ^2 = 1.37, df = 1, *P* = 0.24 ; *Theileria* spp. : χ^2 = 1.37, df = 1, *P* = 0.24 ; *T. equi* : χ^2= 0.001, df = 1, *P*= 0.97). However, *R. aeschlimannii* was significantly much more present in males than females (n(M) = 388 ; n(F) = 340 ; χ^2 = 10.28 , df = 1, *P* = 0.001) and *B. occultans* in females than males (n(M) = 1 ; n(F) = 8 ; χ^2 = 4.14, df = 1, *P* = 0.04).

Discussion

In the present study, several tick-borne microorganisms, including both pathogens and endosymbionts,
were identified in *H. marginatum* ticks primarily collected on horses, with some samples taken from other
animal species, on the island of Corsica as well as on mainland France, where this tick species has recently
settled (Vial et al., 2016). The microfluidic method allowed the detection of eight known or suspected
genera and species of pathogens, specifically *R. aeschlimannii*, *A. phagocytophilum*, *A. marginale*, *E.
minasensis*, *Theileria* spp., *T. equi*, *B. occultans* and West Nile Virus, along with two endosymbionts species
(FLE and CLE), all of which being already demonstrated to circulate in France. The most prevalent
microorganisms detected were FLE and *R. aechlimannii* with a large proportion of ticks infected, while the
infection rates for the other microorganisms were relatively low, ranging from 0.1 to 6.7%, depending on
whether individual or pooled ticks were analysed.
The microfluidic method demonstrated significant potential for high-throughput, targeted detection of
multiple parasites, bacteria and viruses, simultaneously, using only a minimal amount of material enabling
detection at the individual level. However, this approach may overlook unexpected or new microorganisms
present in the sample but that were not included in the pre-defined list of targeted candidates. This
limitation raises concerns when investigating the panel of microorganisms that could be introduced by an
invasive exotic tick species such as *H. marginatum*. Consequently, it was necessary to develop and adapt
the microfluidic array specifically for our study to comprehensively identify the diversity of microorganisms
that could be transmitted by, or at least detected in, such invasive exotic tick species within both its native
and expension geographical zones. Furthermore, our strategy to target not only specific microorganism
species but also broader genera, and to confirm the targets by sequencing, enhanced our detection and
identification capacities. For instance, *B. occultans* was subsequently confirmed in samples that were
initially positive for *Theileria* spp. However, for efficiency reasons, confirmations using sequencing were
performed only on a limited subset of detected samples, resulting in unavoidable extrapolations for
samples exhibiting similar detection patterns. This is illustrated by *Anaplasma* spp. and *A.
phagocytophilum*, for which out of 19 samples sent for sequencing, two sequences confirmed the presence
of *A. phagocytophilum*, while the remaining sequences corresponded to *Devosia* spp., an ubiquitous soil
bacterium contaminating numerous samples and co-amplified by our PCR primers designed for *Anaplasma*
spp. From this observation, the samples that tested positive for *A. phagocytophilum* on the microfluidic
chip and exhibited the same profile as the two samples with confirmed sequences, were all classified as *A.
phagocytophylum*. Conversely, the samples amplifying *Anaplasma* spp. only were ultimately deemed
negative for *Anaplasma*, as we assumed the false positive results were due to contamination by *Devosia*.
This may be a limitation of the microfluigdim method, which avoids to investigate intraspecific diversity
within detected pathogens, but this was clearly not the primary aim of our study. Instead, this high-
throughput approach allowed us, for the first time, to provide a comprehensive panel of microorganisms
encountered in 973 individual *H. marginatum* ticks and 900 ticks grouped in 222 spools collected across 9
French departments, from Spanish to Italian borders and including Corsica island. Despite potential biases
related to differing sampling efforts in Corsica island and mainland France, this method also enabled the
first prevalence estimates of microorganisms (including TBPs) carried by *H. marginatum,* based on their
geographic location and animal hosts. By integrating ecological and epidemiological data on tick-host-
pathogen associations, our study generated first hypotheses regarding the potential role of *H. marginatum*
as vector for the detected pathogens. However, since *H. marginatum* is a hunting tick, all samples in our
study were collected from animals during their blood feeding. This raises questions about the origin of
microorganisms' DNA or RNA detected in ticks; specifically, whether the ticks were truly infected or merely
carried microorganisms from their infected animal hosts via the remaining blood meal in the tick midgut.
While experimental infection studies are scarce, they are crucial for confirmainng a tick species' vector
competence, namely its ability of to acquire a pathogen from an infected host, replicate and maintain it,
and transmit it to a new, naïve host.
Initially, the tick sampling was included in a monitoring program for emergence risk of CCHF, under the
auspices of Ministère de l'Agriculture, de la Souveraineté alimentaire et de la Forêt. CCHFV was not
detected in any of *H. marginatum* tested in our study, despite serological evidence of antibodies against

the virus in cattle from the same areas where the sampled horses were located, which were infested with
H. marginatum (Grech-Angelini et al. 2020). Confirmation of these serological results using the
pseudoplaque neutralization test (PPNT) on CCHFV cultures in BSL-4 conditions suggests that the virus, or
an antigenically similar variant (CCHFV-like), circulates in the natural enzootic cycle between ticks and non-
human vertebrate hosts (Grech-Angelini et al. 2020). In Europe, CCHFV infection prevalence in ticks ranges
from 0.5% to 3.70% (Cicculli et al., 2022), so our sampling effort may have been insufficient to detect at
least one infected tick. Most of the ticks collected between 2016 and 2020 were collected from horses, and
based on studies of *H. marginatum* hosts and their abundance in France, as well as their varying ability to
replicate CCHFV and infect new naïve ticks during blood feeding, most French *H. marginatum* are likely to
be in contact with "poor" hosts, such as birds or horses, which have a low or negligible probability of
transmitting the virus (Bernard et al. 2022). Another hypothesis could be the existence of highly localized
CCHFV transmission hotspots, which may not have been sampled due to difficult access during the optimal
virus' circulation period, an issue that requires further investigation. This hypothesis was supported by the
first detection of CCHFV in *H. marginatum* ticks in France in 2023, where targeted sampling was conducted
(Bernard et al., 2024), unlike the more general sampling in this study. Additionally, the lack of detection of
CCHFV in our study could be attributed to a too low virus load in the ticks to be detected or to a fast
degradation of the genetic material (RNA virus), despite freezing of ticks at -80°C after morphological
identification(Cicculli et al. 2022).
Actually, the only virus detected in the tick samples was WNV, in one individual tick, which was
collected from a Corsican cow, and in one pool of ticks from a Corsican horse. Since cattle are not involved
in the natural cycle of WNV, it is likely that the tick became infected during a previous blood meal, at
immature stage on a bird. Birds play a key role in the WNV cycle and are considered as reservoirs of the
virus, as they are able to reinfect vectors, typically mosquitoes, and possibly ticks that feed on them
(Vaughan, Newman, et Turell 2022). Regarding the tick pool of 10 males collected from a horse, it is
important to note that horses are epidemiological dead-ends for WNV, as they do not produce a sufficient
viremia to infect vectors, including ticks despite larger volumes of blood ingested (David et Abraham 2016).
Additionally, male ticks engorge much less than females even not at all, making it unlikely that the virus
was transmitted from the horse to the ticks. This raises the question of *H. marginatum* being able at least
of transstadial transmission of WNV, although this alone does not confirm full vector competence.
However, horizontal transmission of WNV by *H. marginatum* to an uninfected vertebrate host, in both
immature and adult stages, has been experimentally demonstrated in Portugal (Formosinho et Santos-Silva
2006). Given that WNV can cause neurological disease and death in humans, our findings highlight the
need to intensify monitoring of WNV in *H. marginatum* and further innvestigate the potential of this tick
species to transmit the virus.
Concerning *Francisella* and *Coxiella* detected in our study, they were considered as *Francisella*-like and
Coxiella-like non-pathogenic mutualistic and/or commensal endosymbionts, as previously described
(Buysse et al. 2021; Hussain et al. 2022). In our study, about 97% of *H. marginatum* were infected by FLE,
and only one tick was positive for CLE, which is coherent with previous observations that FLE outcompetes
CLE in ixodid ticks (Kumar et al. 2022). High prevalence of FLE in *H. marginatum* ticks was previously
observed, as it seems to be an ancestral primary endosymbiotic organism (mandatory to tick survival),
being involved in nutritional symbiotic interactions including B vitamins biosynthesis pathways critical for
tick metabolism (vitamin B9 (folate), B2 (riboflavin) and B8 (biotin)) (Buysse et al. 2021)**.** B vitamins are
present at low concentrations in the blood of ticks' hosts compared to the amount required for arthropods
development (Zhong, Jasinskas, et Barbour 2007; Duron et al. 2018; Serrato-Salas et Gendrin 2023),
resulting in the need of production by bacterial endosymbionts. Deprivation of their nutritional symbionts
induced in ticks a complete stop of growth and moulting together with lethal physical abnormalities, which
can be fully restored by exogenous supplementation with B vitamins(Buysse et al. 2021; Cibichakravarthy
et al. 2024).
The second most abundant bacterium identified in *H. marginatum* ticks was *R. aeschlimannii,* detected
in 75% of individual ticks and 95% of tick pools, and thus in much larger proportions than any other bacteria
detected in our study. *R. aeschlimannii* was first isolated from *H. marginatum* ticks in Morocco in 1997
(Beati et al. 1997), and is now regularly reported in ticks across the Mediterranean region (Parola et al.
2014). In Corsica, Grech-Angelini et al (2020) detected *R. aeschlimannii* in 100% of the *H. marginatum* tick
pools they tested. Pooling ticks can inflate infection prevalence estimates , as a pool tests positive even if

only one tick within it is infected. This was evident in our study, where pooled tick samples showed more
frequent positive results compared to individual ticks, with only two negative pools, both containing a
single tick. Given the high prevalence of *R. aeschlimannii* in ticks from both mainland France and Corsica,
and considering its potential pathogenicity in humans, the risk of human exposure to this bacterium likely
exists, even though humans are not common hosts for *H. marginatum*. In Corsica, it has been hypothesized
that the high presence of this bacterium may be correlated to human cases of tick-borne spotted fever
reported on the island (Grech-Angelini et al. 2020). It was previously reported humans cases of spotted
fever in North Africa and South Africa caused by *R. aeschlimannii* infections, and in 2010 a first case
occurred in Southern Europe, in a Greek patient (Germanakis et al. 2013), followed by two addditional
cases in Italy in 2016 and in Germany in 2019 (Tosoni et al. 2016; « Cas suspect de fièvre tachetée à
Rickettsia aeschlimannii en Allemagne », s. d.). Notably, the strain detected in these cases shares 100%
homology with the strain identified in the Greek patient (GeneBank Accession Number: JF803904.1).
However, no human cases of tick-borne spotted fever due to *R. aeschlimannii* have been reported so far
on the French mainland, raising questions about the pathogenic role of *R. aeschlimanii* found in *H.*
marginatum ticks. *R. aeschlimannii* can be transmitted transovarially from *H. marginatum* females to their
offspring, supporting the idea that *H. marginatum* could act as a reservoir for this bacterium (Matsumoto
et al. 2004; Azagi et al. 2017). Given the significantly higher prevalence of *R. aeschlimannii* compared to
other pathogenic bacteria in *H. marginatum*, an alternative hypothesis suggests that it may function as a
secondary symbiont within these ticks (Joly-Kukla et al., 2024). Similar to *Rickettsia peacockii* in
Dermacentor andersoni, secondary symbionts can regulate the development of other infectious *Rickettsia*
(Baldridge et al. 2004). While primary symbionts are virtually fixed in the tick population because vital,
secondary symbionts, though non-essential, may provide ticks with a selective advantage. At this point, it
is unclear whether *R. aeschlimannii* acts as a pathogen, a symbiont, or both. Further investigations are
needed, including its specific detection in tick salivary glands to confirm its ability for horizontal transmition
through tick bite. Additionally, experimental trials involving the removal of *Rickttsia* from ticks would help
clarify its role in tick metabolism and development.

Apart from *R. aeschlimannii*, bacterial DNA detected in *H. marginatum* was mostly represented by the
genus *Anaplasma*, namely *A. phagocytophilum* and *A. marginale, remaining* both at low levels (3,7% in
individual ticks, 6,7 % in tick pools for *A. phagocytophilum* and less than 1% in individual ticks (not detected
in tick pools) for *A. marginale*). *A. phagocytophilum* is a zoonotic obligate intracellular bacterium causing
Tick-Borne Fever in cattle and sheep, Human Granulocytic Anaplasmosis (HGA) in humans, and Equine
Ehrlichiosis (Rikihisa 2011), and *A. marginale* is responsible for the erythrocytic bovine anaplasmosis that
can affect various species of domestic and wild ruminants, including cattle, and that is largely distributed
in France (Aubry et Geale 2011). A recent study showed that in the French Pays Basque, 22.4% of collected
Rhipicephalus bursa ticks contained *A. phagocytophilum* DNA (Dahmani et al. 2017), indicating other tick
species more adapted to carry *A. phagocytophilum*. This TBP has been reported in several European
countries (Matsumoto et al. 2006). HGA cases have been documented in mainland France, with a first case
identified in 2003 and others reported sporadically since then (Edouard et al. 2012). Human cases have
also been recorded in continental Italy (Ruscio et Cinco 2003). This is not the first time *A. phagocytophilum*
has been detected in *H. marginatum*, as one infected tick was found in Israel in 2007 (Keysary et al. 2007),
and three in Tunisia in 2012 (M'ghirbi et al. 2012). As observed in our study, the prevalence of *A.*
phagocytophilum infection in ticks has consistently been low, despite its high presence in cattle
(Woldehiwet 2006) and horses, with 67% of horses testing positive (M'ghirbi et al. 2012). These animals
likely serve as hosts for the adult stages of *H. marginatum*. In France, *A. phagocytophilum* is present in
about 22% of rodent species across different regions (Chastagner et al., 2016), 0.6% of cattle, 0.4% of birds
and 3.3% of common blackbirds (Rouxel et al., 2024), but 0% of horses studied (Dugat et al., 2017). Under
these conditions, the prevalence of infection in ticks carrying feeding on infected hosts would not increase
significantly. Additionally, our detection of the bacterium in non-engorged adult ticks raises questions
about their potential role as reservoirs and vectors for *A. phagocytophilum*, especially given that the hosts
of the tick's immature and adult stages (birds, cattle, horses) are susceptible to this bacterium and that
transovarial transmission occurs in other tick species, what deserves further studies on the vector
competence of *H. marginatum* for *Anaplasma* bacteria. Additionally to *A. phagocytophilum*, *A. marginale*
DNA was found in 3 *H. marginatum* ticks collected from cattle in Corsica, which is consistent with previous
reports in ticks from Corsica (Dahmani et al. 2017; V. Cicculli et al. 2019), including *H. marginatum* (Grech-

Angelini et al. 2020). The primary vectors of this bacterium are *Rhipicephalus* spp. and *Dermacentor* spp. found in tropical and subtropical regions but also in Europe. The activity periods of these ticks may overlap with those of *H. marginatu*m, suggesting the tick could become infected while feeding on an infected animal. The bacterium tipically circulates in late spring and early summer, which coincide with *H. marginatum*'s period of activity, as well as in autumn. Although not considered as a primary vector of this bacterium (Atif 2015), our findings indicate that the presence of this pathogen in our ticks likely reflects their host's infection, as we only detected infected ticks from cattle.

Another bacterium identified in *H. marginatum* was *E. minasensis*, a recent member of the genus *Ehrlichia* and responsible for bovine Ehrliciosis*.* In 2014, it was shown to naturally infect cattle in Brazil and cause clinical Ehrlichiosis in an experimentally infected calf (Aguiar et al. 2014). The distribution of *E. minasensis* extends beyond the Americas (Gajadhar et al. 2010; Cruz et al. 2012; Aguiar et al. 2014), with reports of the bacterium in Pakistan (Rehman et al. 2019), Ethiopia (Hailemariam et al. 2017), South Africa (Iweriebor et al. 2017), Israel (Thomson et al. 2018), and it was recently detected in Corsica island in 3.7% of *H. marginatum* collected from cattle (Vincent Cicculli et al. 2019). Although the vector competence of ticks for this bacterium has yet to be assessed, *E. minasensis* has already been identified in several tick species including *R. microplus* (Cruz et al. 2012; Carvalho et al. 2016), *R. appendiculatus* (Iweriebor et al. 2017) and *H. anatolicum* (Rehman et al. 2019). In addition to infecting bovines (Gajadhar et al. 2010; Aguiar et al. 2014; Hailemariam et al. 2017), *E. minasensis* infects also cervids (Lobanov et al. 2012), dogs (Thomson et al. 2018), and horses (Muraro et al. 2021). In our study, *E. minasensis* was detected in three ticks, two from cattle in Corsica and one from a horse in mainland France. All three ticks were fully engorged and still attached to their host when collected, suggesting this detection may simply be due to their feeding on infected animals. However, we cannot rule out the potetial vector competence of *H. marginatum* for this newly described pathogen, which requires further investigation.

Concerning protozoans, DNA detected in *H. marginatum* was mostly represented by the genus *Theileria*. This was *T. equi*, one of the causative agents of Equine Piroplasmosis (EP) in France, which can also be due to *B. caballi* (Wise et al. 2013) that was not detected in *H. marginatum* in our study raising questions on whether this pathogen is not or no longer circulating in horses in our study area in South of France, which has been previously observed (Rocafort-Ferrer et al. 2022), or that the primers used for PCR amplification also failed to detect *B. caballi*, which desserves futher experiments on the methods to detect the diversity of piroplasms circulating in France. *T. equi* infects wild and domestic equines worldwide, and are enzootic in many countries including several European countries bordering the Mediterranean sea such as France (Tirosh-Levy et al. 2020). This parasite is transmitted by hard tick species belonging to several genera, including *Dermacentor*, *Rhipicephalus*, *Amblyomma*, *Hyalomma*, and *Haemaphysalis* (Scoles and Ueti 2015; Tirosh-Levy et al. 2020). This is not the first instance of this parasite beeing identified in *H. marginatum* in France; a previous study reported that 43% of *H. marginatum* ticks collected from horses in Camargue tested positive for the parasite in 2022 (Rocafort-Ferrer et al. 2022). Notably, our infection rate was considerably lower than that observed in this study. This significant discrepancy may be attributed to the specific design of the tick collection, which focused on stables where cases of piroplasmosis were frequently reported for Rocafort-Ferrer et al. versus stables of all kinds in our study. EP caused by *T. equi* is known to be widespread in France, particularly in the southern regions, with many asymptomatic carrier horses (Rocafort-Ferrer et al. 2022). Although the horses in this study were asymptomatic, a high proportion of horses in the same study area had antibodies against *T. equi* (Guidi et al. 2014), indicating that the parasite circulates in the area. But this is opposite to the low proportion of ticks positive for *T. equi* we found. This is all the more surprising since, once infected with *T. equi*, horses can remain infected for the rest of their lives, becoming asymptomatic carriers of the parasite, and thus constituting a source of infection for uninfected ticks (Friedhoff et al. 1990). Our approach has highlighted the likely underestimation of the infection rate, as we used primers that did not allow the detection of all genotypes potentially circulating in France. In fact, Joly-Kukla et al. (2024) used the same method and later re-evaluated *T. equi* infection rates by targetting a different gene (18S rRNA) through qPCR, which can detect multiple *T. equi* genotypes. Furthermore, our findings may be influenced by the possibility that the areas sampled in our study do not have high levels of parasite circulation, contrary to Camargue (Rocafort-Ferrer et al. 2022). Certain tick species, such as *Rhipicephalus turanicus*, which are present in Camargue but less common in other parts of southern France, are known vectors of the parasites responsible for EP. This could account for a higher local circulation of the parasite and, consequently, a greater prevalence of

infection in ticks. However, this is not an isolated instance; previous studies have also shown cases where
the prevalence of infection in horses was high although the infection rate in ticks remained low. For
example, Nadal (2019) reported that where 50% of horses were infected, yet only 3% of the ticks tested
positive, questionning whether an insufficient parasite load to be infectious through blood meal or a
moderate role of reservoir played by ticks. This could also be linked to our observation that about 5% of
our collected ticks tested positive for *Theileria* only at genus level, suggesting a low circulation in horses
and therefore a very low transmission to ticks, avoiding precise identification of the parasite species
because of insufficient molecular material. While concerns were raised about the risk of introducing
pathogens belonging to the genus *Theileria* causing Tropical Theileriosis with the arrival of *H. marginatum*
in France*,* our study found no*T. annulata* DNA in the collected ticks, neither did we detect *T. buffeli*, a
benign cattle pathogen previously detected in Corsica (ICTTD, 2000). Interstingly, in about two-thirds of
the cases where only *Theileria* spp. was identified, we were not able to determine the specific species of
Theileria because of unsuccessful sequencing. Several hypotheses can be considered for these cases. For
ticks collected from cattle, we can wonder if it is *T. buffeli* that is detected, given its presence in cattle, even
though no species other than *T. equi* was characterized. Alternatively, it may be *T. equi*, as our primers may
not have been suitable for detecting all genotypes circulating in France, which requires the definition of
new primers to improve the detection of all *Theileria* species potentially transmitted by ticks..

We have finally detected *B. occultans* by sequencing in nine cattle ticks from Corsica amplifying first
Theileria spp. with the micro array. Some studies conducted in Turkey and Tunisia, have previously
detected *B. occultans* in *H. rufipes*, *H. marginatum*, *R. turanicus*, and *H. excavatum* ticks (Aktas, Vatansever,
et Ozubek 2014; Ros-García et al. 2011; Orkun 2019; Orkun et al. 2020). So far, only *H. rufipes* has been
confirmed as a vector of *B. occultans* (Ozubek et al. 2020) and *H. marginatum* is not described as a potential
vector of this pathogen. In our study, we can assume that ticks engorged on cows carrying this parasite and
got infected during their blood feeding. Nevertheless, it cannot be ruled out that an evolutive adaptation
could lead to *H. marginatum*-borne transmission of *B. occultans* in southern Europe. This deserves further
investigations in all tick species collected from cattle populations, as recently done in Morocco (Elhachimi
et al. 2021).

Except for known endosymbionts, and the suspected endosymbiotic status of *R. aeschlimanii*, our study
revealed that the infection prevalence of endemic pathogens remains very low in the tick *H. marginatum*,
settled in the south of France since 2016. The vector competence of this tick species for *T.equi* and *A.*
phagocytophilum is still uncertain, but it has to be considered as it is almost demonstrated for CCHFV
(Kiwan 2024, personal communication), confirming the risk of emergence of *Hyalomma marginatum*-borne
pathogens. Additionally, we did not identify any new exotic pathogens that could have have been
introduced by *H. marginatum* from its original distribution area. Based on this information, and considering
that the epidemiological situation is dynamic and may evolve, we propose that *H. marginatum* serves as
valuable sentinel arthropod for anticipating significant health risks for humans or animals in France.
Therefore, it warrants ongoing monitoring due to its potential to spread on long distances through bird
migrations and terrestrial animal trade.

Acknowledgements

We would like to thank all the people who were involved in the field tick collections. We would like to thank Maxime Duhayon and Karine Huber for technical support in managing tick processing and biological material extraction. We would like to thank all cattle and horse owners for collaborating with us on this study.

Funding

The authors thank the funders who made this work possible: French Ministry of Agriculture—General Directorate for Food (DGAl, grant agreement: SPA17 number 0079-E), European Funds for Regional Development (FEDER, Grand-Est), French Establishment for Fighting Zoonoses (ELIZ) and the Association Nationale Recherche Technologie (ANRT, grant agreement number: 2019-1145), Défi clé RIVOC (Occitanie Region): Holis-tiques project. This research was funded by grant from the French Agency for Food, Environmental and Occupational Health & Safety (ANSES).

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. Thomas Pollet, Sara Moutailler and Laurence Vial are recommenders for PCI infections.

Author contributions

Conceptualization: CB, MB, BC, LV, PH; Data curation: CB, CG, CJK, VC, SGA, PEP, SM, PH, LV ; Funding acquisition: MB, BC, SM, PH, LV; ; Methodology: CB, CG, SGA, SM, PH, LV ; Project administration: CB, TP, MB, BC, SM, PH, LV; Supervision: CB, TP, MB, BC, SM, PH, LV; Validation: all authors; Writing – original draft: CB, PH, LV; Writing – review and editing : all authors.

Data, scripts and codes availability

Data are available online: https://gitlab.cirad.fr/astre/data_tickpathogens_cbernard2025

References

Aguiar, Daniel M., Thayza F. Ziliani, Xiaofeng Zhang, Andreia L. T. Melo, Ísis A. Braga, Rute Witter, Leodil C. Freitas, et al. 2014. « A Novel Ehrlichia Genotype Strain Distinguished by the TRP36 Gene Naturally Infects Cattle in Brazil and Causes Clinical Manifestations Associated with Ehrlichiosis ». *Ticks and Tick-Borne Diseases* 5 (5): 537-44. https://doi.org/10.1016/j.ttbdis.2014.03.010.

Aktas, Munir, Zati Vatansever, et Sezayi Ozubek. 2014. « Molecular Evidence for Trans-Stadial and Transovarial Transmission of Babesia Occultans in Hyalomma Marginatum and Rhipicephalus Turanicus in Turkey ». *Veterinary Parasitology* 204 (3-4): 369-71. https://doi.org/10.1016/j.vetpar.2014.05.037.

Atif, Farhan Ahmad. 2015. « Anaplasma Marginale and Anaplasma Phagocytophilum: Rickettsiales Pathogens of Veterinary and Public Health Significance ». *Parasitology Research* 114 (11): 3941-57. https://doi.org/10.1007/s00436-015-4698-2.

Aubry, P., et D. W. Geale. 2011. « A Review of Bovine Anaplasmosis ». *Transboundary and Emerging*
Diseases 58 (1): 1-30. https://doi.org/10.1111/j.1865-1682.2010.01173.x.

Azagi, Tal, Eyal Klement, Gidon Perlman, Yaniv Lustig, Kosta Y. Mumcuoglu, Dmitry A.
Apanaskevich, et Yuval Gottlieb. 2017. « Francisella-Like Endosymbionts and Rickettsia
Species in Local and Imported Hyalomma Ticks ». *Applied and Environmental Microbiology* 83
(18): e01302-17. https://doi.org/10.1128/AEM.01302-17.

Bah, Madiou Thierno, Vladimir Grosbois, Frédéric Stachurski, Facundo Muñoz, Maxime Duhayon,
Ignace Rakotoarivony, Anaïs Appelgren, et al. 2022. « The Crimean-Congo haemorrhagic fever
tick vector Hyalomma marginatum in the south of France: Modelling its distribution and
determination of factors influencing its establishment in a newly invaded area ». *Transboundary*
and Emerging Diseases 69 (5): e2351-65. https://doi.org/10.1111/tbed.14578.

Bakheit, M, A Latif, Z Vatansever, U Seitzer, et J Ahmed. 2012. « The Huge Risks Due to Hyalomma
Ticks ». https://doi.org/10.1007/978-3-642-28842-5_8.

Baldridge, Gerald D., Nicole Y. Burkhardt, Jason A. Simser, Timothy J. Kurtti, et Ulrike G. Munderloh.
2004. « Sequence and Expression Analysis of the ompA Gene of Rickettsia Peacockii, an
Endosymbiont of the Rocky Mountain Wood Tick, Dermacentor Andersoni ». *Applied and*
Environmental Microbiology 70 (11): 6628-36. https://doi.org/10.1128/AEM.70.11.6628-6636.2004.

Beati, L., M. Meskini, B. Thiers, et D. Raoult. 1997. « Rickettsia Aeschlimannii Sp. Nov., a New Spotted
Fever Group Rickettsia Associated with Hyalomma Marginatum Ticks ». *International Journal*
of Systematic Bacteriology 47 (2): 548-54. https://doi.org/10.1099/00207713-47-2-548.

Bernard, Célia, Philippe Holzmuller, Madiou Thierno Bah, Matthieu Bastien, Benoit Combes, Ferran
Jori, Vladimir Grosbois, et Laurence Vial. 2022. « Systematic Review on Crimean–Congo
Hemorrhagic Fever Enzootic Cycle and Factors Favoring Virus Transmission: Special Focus on
France, an Apparently Free-Disease Area in Europe ». *Frontiers in Veterinary Science* 9.
https://www.frontiersin.org/articles/10.3389/fvets.2022.932304.

Bonnet, Sarah I., Stéphane Bertagnoli, Alessandra Falchi, Julie Figoni, Johanna Fite, Thierry Hoch, Elsa
Quillery, et al. 2023. « An Update of Evidence for Pathogen Transmission by Ticks of the Genus
Hyalomma ». *Pathogens* 12 (4): 513. https://doi.org/10.3390/pathogens12040513.

Bonnet, Sarah I., Gwenaël Vourc'h, Alice Raffetin, Alessandra Falchi, Julie Figoni, Johanna Fite, Thierry
Hoch, Sara Moutailler, et Elsa Quillery. 2022. « The Control of Hyalomma Ticks, Vectors of the
Crimean-Congo Hemorrhagic Fever Virus: Where Are We Now and Where Are We Going? »
PLoS Neglected Tropical Diseases 16 (11): e0010846. https://doi.org/10.1371/journal.pntd.0010846.

Boulanger, Nathalie, et Karen D. McCoy, éd. 2017. *Tiques et maladies à tiques : Biologie, écologie évolutive,*
épidémiologie. *Tiques et maladies à tiques : Biologie, écologie évolutive, épidémiologie*. Didactiques.
Marseille: IRD Éditions. https://doi.org/10.4000/books.irdeditions.9001.

Buysse, Marie, Anna Maria Floriano, Yuval Gottlieb, Tiago Nardi, Francesco Comandatore, Emanuela
Olivieri, Alessia Giannetto, et al. 2021. « A dual endosymbiosis supports nutritional adaptation
to hematophagy in the invasive tick Hyalomma marginatum ». Édité par Bruno Lemaître,
Patricia J Wittkopp, Bruno Lemaître, Sukanya Narasimhan, et Alejandro Manzano-Marin. *eLife*
10 (décembre):e72747. https://doi.org/10.7554/eLife.72747.

Carvalho, Izabelle T. S., Andréia L. T. Melo, Leodil C. Freitas, Rodolfo V. Verçoza, Alvair S. Alves,
Jackeliny S. Costa, Cristiane S. Chitarra, et al. 2016. « Minimum Infection Rate of Ehrlichia
Minasensis in Rhipicephalus Microplus and Amblyomma Sculptum Ticks in Brazil ». *Ticks and*
Tick-Borne Diseases 7 (5): 849-52. https://doi.org/10.1016/j.ttbdis.2016.04.004.

« Cas suspect de fièvre tachetée à Rickettsia aeschlimannii en Allemagne ». s. d. Consulté le 28 janvier
2025. https://www.vidal.fr/actualites/23772-cas-suspect-de-fievre-tachetee-a-rickettsia-
aeschlimannii-en-allemagne.html.

Cibichakravarthy, Balasubramanian, Neta Shaked, Einat Kapri, et Yuval Gottlieb. 2024.
« Endosymbiont-derived metabolites are essential for tick host reproductive fitness ». *mSphere*
9 (7): e00693-23. https://doi.org/10.1128/msphere.00693-23.

Cicculli, V., L. Capai, Y. Quilichini, S. Masse, A. Fernández-Alvarez, L. Minodier, P. Bompard, R. Charrel, et A. Falchi. 2019. « Molecular Investigation of Tick-Borne Pathogens in Ixodid Ticks Infesting Domestic Animals (Cattle and Sheep) and Small Rodents (Black Rats) of Corsica, France ». *Ticks and Tick-Borne Diseases* 10 (3): 606-13. https://doi.org/10.1016/j.ttbdis.2019.02.007.

Cicculli, Vincent, Apolline Maitre, Nazli Ayhan, Stevan Mondoloni, Jean-Christophe Paoli, Laurence Vial, Xavier N. de Lamballerie, Remi Charrel, et Alessandra Falchi. 2022. « Lack of Evidence for Crimean-Congo Hemorrhagic Fever Virus in Ticks Collected from Animals, Corsica, France ». *Emerging Infectious Diseases* 28 (5): 1035-38. https://doi.org/10.3201/eid2805.211996.

Cicculli, Vincent, Shirley Masse, Lisandru Capai, Xavier de Lamballerie, Remi Charrel, et Alessandra Falchi. 2019. « First detection of Ehrlichia minasensis in Hyalomma marginatum ticks collected from cattle in Corsica, France ». *Veterinary Medicine and Science* 5 (2): 243-48. https://doi.org/10.1002/vms3.140.

Colwell, Douglas D., Filipe Dantas-Torres, et Domenico Otranto. 2011. « Vector-Borne Parasitic Zoonoses: Emerging Scenarios and New Perspectives ». *Veterinary Parasitology* 182 (1): 14-21. https://doi.org/10.1016/j.vetpar.2011.07.012.

Cosson, Jean-François, Lorraine Michelet, Julien Chotte, Evelyne Le Naour, Martine Cote, Elodie Devillers, Marie-Lazarine Poulle, et al. 2014. « Genetic Characterization of the Human Relapsing Fever Spirochete Borrelia Miyamotoi in Vectors and Animal Reservoirs of Lyme Disease Spirochetes in France ». *Parasites & Vectors* 7 (1): 233. https://doi.org/10.1186/1756-3305-7-233.

Cruz, Alejandro Cabezas, Erich Zweygarth, Mucio Flavio Barbosa Ribeiro, Julia Angelica Gonçalves da Silveira, Jose de la Fuente, Libor Grubhoffer, James J. Valdés, et Lygia Maria Friche Passos. 2012. « New species of Ehrlichia isolated from Rhipicephalus (Boophilus) microplus shows an ortholog of the E. canis major immunogenic glycoprotein gp36 with a new sequence of tandem repeats ». *Parasites & Vectors* 5 (1): 291. https://doi.org/10.1186/1756-3305-5-291.

Dahmani, Mustapha, Bernard Davoust, Frédéric Rousseau, Didier Raoult, Florence Fenollar, et Oleg Mediannikov. 2017. « Natural Anaplasmataceae Infection in Rhipicephalus Bursa Ticks Collected from Sheep in the French Basque Country ». *Ticks and Tick-Borne Diseases* 8 (1): 18-24. https://doi.org/10.1016/j.ttbdis.2016.09.009.

David, Shoba, et Asha Mary Abraham. 2016. « Epidemiological and Clinical Aspects on West Nile Virus, a Globally Emerging Pathogen ». *Infectious Diseases (London, England)* 48 (8): 571-86. https://doi.org/10.3109/23744235.2016.1164890.

Demir, Samiye, Sedef Erkunt Alak, Ahmet Efe Köseoğlu, Cemal Ün, Muhammed Nalçacı, et Hüseyin Can. 2020. « Molecular Investigation of Rickettsia Spp. and Francisella Tularensis in Ticks from Three Provinces of Turkey ». *Experimental & Applied Acarology* 81 (2): 239-53. https://doi.org/10.1007/s10493-020-00498-y.

Duron, Olivier, Olivier Morel, Valérie Noël, Marie Buysse, Florian Binetruy, Renaud Lancelot, Etienne Loire, et al. 2018. « Tick-Bacteria Mutualism Depends on B Vitamin Synthesis Pathways ». *Current Biology* 28 (12): 1896-1902.e5. https://doi.org/10.1016/j.cub.2018.04.038.

Edouard, Sophie, Christelle Koebel, François Goehringer, Cristina Socolovschi, Benoit Jaulhac, Didier Raoult, et Philippe Brouqui. 2012. « Emergence of Human Granulocytic Anaplasmosis in France ». *Ticks and Tick-Borne Diseases* 3 (5): 403-5. https://doi.org/10.1016/j.ttbdis.2012.10.002.

Elhachimi, Latifa, Carolien Rogiers, Stijn Casaert, Siham Fellahi, Thomas Van Leeuwen, Wannes Dermauw, Félix Valcárcel, et al. 2021. « Ticks and Tick-Borne Pathogens Abound in the Cattle Population of the Rabat-Sale Kenitra Region, Morocco ». *Pathogens (Basel, Switzerland)* 10 (12): 1594. https://doi.org/10.3390/pathogens10121594.

Fernández-Ruiz, Natalia, et Agustín Estrada-Peña. 2021. « Towards New Horizons: Climate Trends in Europe Increase the Environmental Suitability for Permanent Populations of Hyalomma Marginatum (Ixodidae) ». *Pathogens (Basel, Switzerland)* 10 (2): 95. https://doi.org/10.3390/pathogens10020095.

Formosinho, P., et M. M. Santos-Silva. 2006. « Experimental Infection of Hyalomma Marginatum Ticks
with West Nile Virus ». *Acta Virologica* 50 (3): 175-80.
Gajadhar, Alvin A., Vladislav Lobanov, W. Brad Scandrett, John Campbell, et Batol Al-Adhami. 2010.
« A Novel Ehrlichia Genotype Detected in Naturally Infected Cattle in North America ».
Veterinary Parasitology 173 (3): 324-29. https://doi.org/10.1016/j.vetpar.2010.06.034.
Germanakis, Antonis, Dimosthenis Chochlakis, Emmanouil Angelakis, Yannis Tselentis, et Anna
Psaroulaki. 2013. « Rickettsia Aeschlimannii Infection in a Man, Greece ». *Emerging Infectious*
Diseases 19 (7): 1176-77. https://doi.org/10.3201/eid1907.130232.
Gondard, Mathilde, Lorraine Michelet, Athinna Nisavanh, Elodie Devillers, Sabine Delannoy, Patrick
Fach, Anna Aspan, et al. 2018. « Prevalence of tick-borne viruses in Ixodes ricinus assessed by
high-throughput   real-time   PCR ».   *Pathogens   and   Disease*   76   (8):   fty083.
https://doi.org/10.1093/femspd/fty083.
Gray, J. S. 1991. « The Development and Seasonal Activity of the Tick Ixodes Ricinus: A Vector of Lyme
Borreliosis. » *Review of Medical and Veterinary Entomology* 79 (6): 323-33.
Grech-Angelini, Sébastien, Frédéric Stachurski, Muriel Vayssier-Taussat, Elodie Devillers, François
Casabianca, Renaud Lancelot, Gerrit Uilenberg, et Sara Moutailler. 2020. « Tick-Borne
Pathogens in Ticks (Acari: Ixodidae) Collected from Various Domestic and Wild Hosts in
Corsica (France), a Mediterranean Island Environment ». *Transboundary and Emerging Diseases*.
https://doi.org/10.1111/tbed.13393.
Hailemariam, Zerihun, Jürgen Krücken, Maximilian Baumann, Jabbar S. Ahmed, Peter-Henning
Clausen, et Ard M. Nijhof. 2017. « Molecular Detection of Tick-Borne Pathogens in Cattle from
Southwestern   Ethiopia ».   *PLOS   ONE*   12   (11):   e0188248.
https://doi.org/10.1371/journal.pone.0188248.
Hoffman, Tove, Mats Lindeborg, Christos Barboutis, Kiraz Erciyas-Yavuz, Magnus Evander, Thord
Fransson, Jordi Figuerola, et al. 2018. « Alkhurma Hemorrhagic Fever Virus RNA in
Hyalomma rufipes Ticks Infesting Migratory Birds, Europe and Asia Minor ». *Emerging*
Infectious Diseases 24 (5): 879-82. https://doi.org/10.3201/eid2405.171369.
Horton, Katherine C., Nermeen T. Fahmy, Noha Watany, Alia Zayed, Abro Mohamed, Ammar Abdo
Ahmed, Pierre E. Rollin, et Erica L. Dueger. 2016. « Crimean Congo Hemorrhagic Fever Virus
and Alkhurma (Alkhumra) Virus in Ticks in Djibouti ». *Vector-Borne and Zoonotic Diseases* 16
(10): 680-82. https://doi.org/10.1089/vbz.2016.1951.
Hussain, Sabir, Nighat Perveen, Abrar Hussain, Baolin Song, Muhammad Umair Aziz, Jehan Zeb, Jun
Li, David George, Alejandro Cabezas-Cruz, et Olivier Sparagano. 2022. « The Symbiotic
Continuum Within Ticks: Opportunities for Disease Control ». *Frontiers in Microbiology* 13.
https://www.frontiersin.org/articles/10.3389/fmicb.2022.854803.
Iweriebor, Benson C., Elia J. Mmbaga, Abiodun Adegborioye, Aboi Igwaran, Larry C. Obi, et Anthony
I. Okoh. 2017. « Genetic profiling for Anaplasma and Ehrlichia species in ticks collected in the
Eastern   Cape   Province   of   South   Africa ».   *BMC   Microbiology*   17   (1):   45.
https://doi.org/10.1186/s12866-017-0955-0.
Jongejan, F., et G. Uilenberg. 2004. « The Global Importance of Ticks ». *Parasitology* 129 Suppl:S3-14.
https://doi.org/10.1017/s0031182004005967.
Keysary, Avi, Robert F. Massung, Moshe Inbar, Arian D. Wallach, Uri Shanas, Kosta Y. Mumcuoglu, et
Trevor Waner. 2007. « Molecular Evidence for Anaplasma phagocytophilum in Israel ».
Emerging Infectious Diseases 13 (9): 1411-12. https://doi.org/10.3201/eid1309.070455.
Kilicoglu, Yunus, Abdurrahman Anil Cagirgan, Gulnur Serdar, Selma Kaya, Yuksel Durmaz, et Yunus
Gur. 2020. « Molecular Investigation, Isolation and Phylogenetic Analsysis of Coxiella Burnetii
from Aborted Fetus and Ticks ». *Comparative Immunology, Microbiology and Infectious Diseases* 73
(décembre):101571. https://doi.org/10.1016/j.cimid.2020.101571.
Kolodziejek, Jolanta, Mihai Marinov, Botond J. Kiss, Vasile Alexe, et Norbert Nowotny. 2014. « The
Complete Sequence of a West Nile Virus Lineage 2 Strain Detected in a Hyalomma
Marginatum Marginatum Tick Collected from a Song Thrush (Turdus Philomelos) in Eastern

Romania in 2013 Revealed Closest Genetic Relationship to Strain Volgograd 2007 ». *PLOS ONE*
9 (10): e109905. https://doi.org/10.1371/journal.pone.0109905.

Kumar, Deepak, Surendra Raj Sharma, Abdulsalam Adegoke, Ashley Kennedy, Holly C. Tuten,
Andrew Y. Li, et Shahid Karim. 2022. « Recently Evolved Francisella-Like Endosymbiont
Outcompetes an Ancient and Evolutionarily Associated Coxiella-Like Endosymbiont in the
Lone Star Tick (Amblyomma Americanum) Linked to the Alpha-Gal Syndrome ». *Frontiers in*
Cellular and Infection Microbiology 12:787209. https://doi.org/10.3389/fcimb.2022.787209.

Lobanov, V. A., A. A. Gajadhar, B. Al-Adhami, et H. M. Schwantje. 2012. « Molecular Study of Free-
Ranging Mule Deer and White-Tailed Deer from British Columbia, Canada, for Evidence of
Anaplasma Spp. and Ehrlichia Spp. » *Transboundary and Emerging Diseases* 59 (3): 233-43.
https://doi.org/10.1111/j.1865-1682.2011.01259.x.

Matsumoto, K., G. Joncour, B. Davoust, Pierre-Hugues Pitel, A. Chauzy, E. Collin, H. Morvan, N.
Vassallo, et P. Brouqui. 2006. « Anaplasma Phagocytophilum Infection in Cattle in France ».
Annals of the New York Academy of Sciences 1078 (1): 491. https://doi.org/10.1196/annals.1374.092.

Matsumoto, K., P. Parola, P. Brouqui, et D. Raoult. 2004. « Rickettsia Aeschlimannii in Hyalomma Ticks
from Corsica ». *European Journal of Clinical Microbiology and Infectious Diseases* 23 (9): 732-34.
https://doi.org/10.1007/s10096-004-1190-9.

M'ghirbi, Youmna, Hèla Yaïch, Abderazek Ghorbel, et Ali Bouattour. 2012. « Anaplasma
phagocytophilum in horses and ticks in Tunisia ». *Parasites & Vectors* 5 (août):180.
https://doi.org/10.1186/1756-3305-5-180.

Michelet, Lorraine, Sabine Delannoy, Elodie Devillers, Gérald Umhang, Anna Aspan, Mikael
Juremalm, Jan Chirico, et al. 2014. « High-throughput screening of tick-borne pathogens in
Europe ». *Frontiers in Cellular and Infection Microbiology* 4.
https://www.frontiersin.org/articles/10.3389/fcimb.2014.00103.

Moraga-Fernández, Alberto, Francisco Ruiz-Fons, Miguel A. Habela, Lara Royo-Hernández, Rafael
Calero-Bernal, Christian Gortazar, José de la Fuente, et Isabel G. Fernández de Mera. 2021.
« Detection of New Crimean–Congo Haemorrhagic Fever Virus Genotypes in Ticks Feeding
on Deer and Wild Boar, Spain ». *Transboundary and Emerging Diseases* 68 (3): 993-1000.
https://doi.org/10.1111/tbed.13756.

Muraro, Lívia S., Aneliza de O. Souza, Tamyres N. S. Leite, Stefhano L. Cândido, Andréia L. T. Melo,
Hugo S. Toma, Mariana B. Carvalho, et al. 2021. « First Evidence of Ehrlichia Minasensis
Infection in Horses from Brazil ». *Pathogens* 10 (3): 265.
https://doi.org/10.3390/pathogens10030265.

Nadal, Clémence, Sarah I. Bonnet, et Maud Marsot. 2022. « Eco-Epidemiology of Equine Piroplasmosis
and Its Associated Tick Vectors in Europe: A Systematic Literature Review and a Meta-
Analysis of Prevalence ». *Transboundary and Emerging Diseases* 69 (5): 2474-98.
https://doi.org/10.1111/tbed.14261.

Orkun, Ömer. 2019. « Molecular Investigation of the Natural Transovarial Transmission of Tick-Borne
Pathogens in Turkey ». *Veterinary Parasitology* 273 (septembre):97-104.
https://doi.org/10.1016/j.vetpar.2019.08.013.

———. 2022. « Comprehensive Screening of Tick-Borne Microorganisms Indicates That a Great Variety
of Pathogens Are Circulating between Hard Ticks (Ixodoidea: Ixodidae) and Domestic
Ruminants in Natural Foci of Anatolia ». *Ticks and Tick-Borne Diseases* 13 (6): 102027.
https://doi.org/10.1016/j.ttbdis.2022.102027.

Orkun, Ömer, Ayşe Çakmak, Serpil Nalbantoğlu, et Zafer Karaer. 2020. « Turkey Tick News: A
Molecular Investigation into the Presence of Tick-Borne Pathogens in Host-Seeking Ticks in
Anatolia; Initial Evidence of Putative Vectors and Pathogens, and Footsteps of a Secretly Rising
Vector Tick, Haemaphysalis Parva ». *Ticks and Tick-Borne Diseases* 11 (3): 101373.
https://doi.org/10.1016/j.ttbdis.2020.101373.

Ozubek, Sezayi, Reginaldo G. Bastos, Heba F. Alzan, Abdullah Inci, Munir Aktas, et Carlos E. Suarez. 2020. « Bovine Babesiosis in Turkey: Impact, Current Gaps, and Opportunities for Intervention ». *Pathogens* 9 (12): 1041. https://doi.org/10.3390/pathogens9121041.

Palomar, Ana M., Aránzazu Portillo, David Mazuelas, Lidia Roncero, Juan Arizaga, Ariñe Crespo, Óscar Gutiérrez, et al. 2016. « Molecular Analysis of Crimean-Congo Hemorrhagic Fever Virus and Rickettsia in Hyalomma Marginatum Ticks Removed from Patients (Spain) and Birds (Spain and Morocco), 2009–2015 ». *Ticks and Tick-Borne Diseases* 7 (5): 983-87. https://doi.org/10.1016/j.ttbdis.2016.05.004.

Parola, Philippe, Christopher D. Paddock, Cristina Socolovschi, Marcelo B. Labruna, Oleg Mediannikov, Tahar Kernif, Mohammad Yazid Abdad, et al. 2014. « Update on Tick-Borne Rickettsioses around the World: a Geographic Approach ». *Clinical Microbiology Reviews* 27 (1): 166. https://doi.org/10.1128/CMR.00104-13.

PÉREZ-EID, Claudine. 2007. *Les tiques : identification, biologie, importance médicale et vétérinaire*. Lavoisier.

Perveen, Nighat, et Gulfaraz Khan. 2022. « Crimean–Congo hemorrhagic fever in the Arab world: A systematic review ». *Frontiers in Veterinary Science* 9 (septembre):938601. https://doi.org/10.3389/fvets.2022.938601.

Rehman, Abdul, Franz J. Conraths, Carola Sauter-Louis, Jürgen Krücken, et Ard M. Nijhof. 2019. « Epidemiology of Tick-Borne Pathogens in the Semi-Arid and the Arid Agro-Ecological Zones of Punjab Province, Pakistan ». *Transboundary and Emerging Diseases* 66 (1): 526-36. https://doi.org/10.1111/tbed.13059.

Rikihisa, Yasuko. 2011. « Mechanisms of Obligatory Intracellular Infection with Anaplasma phagocytophilum ». *Clinical Microbiology Reviews* 24 (3): 469-89. https://doi.org/10.1128/CMR.00064-10.

Ristic, Miodrag. 1981. « Anaplasmosis ». In *Diseases of Cattle in the Tropics: Economic and Zoonotic Relevance*, édité par Miodrag Ristic et Ian McIntyre, 327-44. Current Topics in Veterinary Medicine and Animal Science. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-015-6895-1_27.

Rocafort-Ferrer, Gloria, Agnès Leblond, Aurélien Joulié, Magalie René-Martellet, Alain Sandoz, Valérie Poux, Sophie Pradier, Séverine Barry, Laurence Vial, et Loïc Legrand. 2022. « Molecular Assessment of Theileria Equi and Babesia Caballi Prevalence in Horses and Ticks on Horses in Southeastern France ». *Parasitology Research* 121 (3): 999-1008. https://doi.org/10.1007/s00436-022-07441-7.

Ros-García, A., Y. M'ghirbi, A. Bouattour, et A. Hurtado. 2011. « First Detection of Babesia Occultans in Hyalomma Ticks from Tunisia ». *Parasitology* 138 (5): 578-82. https://doi.org/10.1017/S0031182011000060.

Rothschild, Chantal M. 2013. « Equine Piroplasmosis ». *Journal of Equine Veterinary Science* 33 (7): 497-508. https://doi.org/10.1016/j.jevs.2013.03.189.

Ruscio, M., et M. Cinco. 2003. « Human granulocytic ehrlichiosis in Italy: first report on two confirmed cases. » *Annals of the New York Academy of Sciences*. https://www.semanticscholar.org/paper/Human-granulocytic-ehrlichiosis-in-Italy%3A-first-on-Ruscio-Cinco/b5b6d8ff69ae6c5dff525b35b7a3f4f7048af442.

Satta, Giuseppe, Valentina Chisu, Pierangela Cabras, Francesco Fois, et Giovanna Masala. 2011. « Pathogens and symbionts in ticks: a survey on tick species distribution and presence of tick-transmitted micro-organisms in Sardinia, Italy ». *Journal of Medical Microbiology* 60 (1): 63-68. https://doi.org/10.1099/jmm.0.021543-0.

Selmi, Rachid, Mourad Ben Said, Aymen Mamlouk, Houcine Ben Yahia, et Lilia Messadi. 2019. « Molecular Detection and Genetic Characterization of the Potentially Pathogenic Coxiella Burnetii and the Endosymbiotic Candidatus Midichloria Mitochondrii in Ticks Infesting Camels (Camelus Dromedarius) from Tunisia ». *Microbial Pathogenesis* 136 (novembre):103655. https://doi.org/10.1016/j.micpath.2019.103655.

Serrato-Salas, Javier, et Mathilde Gendrin. 2023. « Involvement of Microbiota in Insect Physiology:
Focus on B Vitamins ». Édité par Brian Weiss et Jacob Yount. *mBio* 14 (1): e02225-22.
https://doi.org/10.1128/mbio.02225-22.
Thomson, Kirsty, Tal Yaaran, Alex Belshaw, Lucia Curson, Laurence Tisi, Sarah Maurice, et Guy
Kiddle. 2018. « A New TaqMan Method for the Reliable Diagnosis of Ehrlichia Spp. in Canine
Whole Blood ». *Parasites & Vectors* 11 (1): 350. https://doi.org/10.1186/s13071-018-2914-5.
Tirosh-Levy, Sharon, Yuval Gottlieb, Monica L. Mazuz, Igor Savitsky, et Amir Steinman. 2020.
« Infection Dynamics of Theileria Equi in Carrier Horses Is Associated with Management and
Tick   Exposure ».   *Ticks   and   Tick-Borne   Diseases*   11   (6):   101508.
https://doi.org/10.1016/j.ttbdis.2020.101508.
Tosoni, A., A. Mirijello, A. Ciervo, F. Mancini, G. Rezza, F. Damiano, R. Cauda, A. Gasbarrini, G.
Addolorato, et Internal Medicine Sepsis Study Group. 2016. « Human Rickettsia Aeschlimannii
Infection: First Case with Acute Hepatitis and Review of the Literature ». *European Review for*
Medical and Pharmacological Sciences 20 (12): 2630-33.
Vaughan, Jefferson A., Robert A. Newman, et Michael J. Turell. 2022. « Bird Species Define the
Relationship between West Nile Viremia and Infectiousness to Culex Pipiens Mosquitoes ».
PLOS Neglected Tropical Diseases 16 (10): e0010835. https://doi.org/10.1371/journal.pntd.0010835.
Vial, Laurence, Frédéric Stachurski, Agnès Leblond, Karine Huber, Gwenaël Vourc'h, Magalie René-
Martellet, Isabelle Desjardins, et al. 2016. « Strong Evidence for the Presence of the Tick
Hyalomma Marginatum Koch, 1844 in Southern Continental France ». *Ticks and Tick-Borne*
Diseases 7 (6): 1162-67. https://doi.org/10.1016/j.ttbdis.2016.08.002.
Wise, L.n., L.s. Kappmeyer, R.h. Mealey, et D.p. Knowles. 2013. « Review of Equine Piroplasmosis ».
Journal of Veterinary Internal Medicine 27 (6): 1334-46. https://doi.org/10.1111/jvim.12168.
Zhong, Jianmin, Algimantas Jasinskas, et Alan G. Barbour. 2007. « Antibiotic Treatment of the Tick
Vector Amblyomma Americanum Reduced Reproductive Fitness ». *PloS One* 2 (5): e405.
https://doi.org/10.1371/journal.pone.0000405.