Coxiella burnetii-containing vacuoles interact with host recycling-endosomal proteins Rab11a and Rab35 for vacuolar expansion

Brooke A. Hall[1], Kristen E. Senior[1], Nicolle T. Ocampo[2], and Dhritiman Samanta[1]

[1]Department of Microbiology and Immunology, College of Graduate Studies, Midwestern University, Glendale, AZ, USA

[2]Arizona College of Osteopathic Medicine, Midwestern University, Glendale, AZ, USA

* Correspondence:
Dhritiman Samanta
Email: dsaman@midwestern.edu

Keywords: *Coxiella burnetii*, recycling endosomes, Rab11a, Rab35, Intracellular Bacterial Pathogen, confocal microscopy

Abstract

Coxiella burnetii is a gram-negative obligate intracellular bacterium and a zoonotic pathogen that causes human Q fever. Although acute Q fever manifests as atypical pneumonia, chronic infection may lead to life-threatening endocarditis. The lack of effective antibiotics and a licensed vaccine for *Coxiella* in the U.S. warrants further research into *Coxiella* pathogenesis. Within the host cells, *Coxiella* replicates in an acidic phagolysosome-like vacuole termed *Coxiella*-containing vacuole (CCV). Previously, we have shown that the CCV pH is critical for *Coxiella* survival and that the *Coxiella* Type 4B secretion system regulates CCV pH by inhibiting the host endosomal maturation pathway. However, the trafficking pattern of the 'immature' endosomes in *Coxiella*-infected cells remained unclear. Our recent CCV localization screen with host Rab proteins revealed that recycling endosome-associated proteins Rab11a and Rab35 localize to the CCV during infection, suggesting that CCV interacts with host recycling endosomes during maturation. Interestingly, only a subset of CCVs were Rab11a or Rab35-positive at any given time point. A quantitation of Rab11a/Rab35-positive CCVs at 3- and 6-days post-infection (dpi) revealed that at 3 dpi, ~52% of CCVs were positive for Rab11a, whereas only ~39% CCVs were positive for Rab35. This pattern reversed at 6-dpi, when only ~22% of CCVs were positive for Rab11a and ~64% of CCVs were positive for Rab35. These data suggest that the CCV preferentially interacts with Rab11a and Rab35-positive recycling endosomes depending on the stage of maturation. Furthermore, we observed a significant increase in Rab11a and Rab35 fluorescent intensity in *Coxiella*-infected cells compared to mock cells suggesting that *Coxiella* increases the recycling endosome content in infected cells. Finally, a siRNA-mediated knockdown of both Rab11a and Rab35 resulted in significantly smaller CCVs, suggesting that recycling endosomal Rab proteins are essential for CCV expansion. Overall, our data, for the

39 first time, show that the CCV dynamically interacts with host recycling endosomes for vacuolar
40 expansion and potentially uncovers novel host cell factors essential for *Coxiella* pathogenesis.

1   Introduction

Coxiella burnetii is a highly infectious, zoonotic, bacterial pathogen and the causative agent of
human Q (query) fever. In acute cases, Q fever manifests as a flu-like illness, whereas in about 5% of
patients, the disease progresses to a chronic stage that often leads to life-threatening endocarditis [1].
The treatment plan for chronic Q fever involves an 18-month-long combined antibiotic therapy and
currently, there is no Q fever vaccine licensed in the U.S., presenting challenges in both treating and
preventing the disease. Q fever has emerged as a significant public health concern worldwide [2-10].
In the course of a natural infection, *Coxiella* is transmitted through the respiratory route,
where alveolar macrophages within the lung parenchyma endocytose the bacteria [11, 12]. Following
entry into host cells, *Coxiella* resides in a tight-fitting phagosome that expands and matures over the
next 48-72 hours by interacting with the host endocytic pathway [13]. The phagosome maturation
pathway involves sequential fusion with the host early and late endosomes, incorporating their
respective membrane markers Rab5 and Rab7 [13, 14] and subsequent fusion with lysosomes,
acquiring lysosomal markers such as LAMP1 (lysosome-associated membrane glycoprotein-1) and
vacuolar ATPase [15-19]. The phagosome-lysosome fusion is a unique event in the *Coxiella* life cycle
that generates an acidic phagolysosome-like vacuole with a pH of ~5.2 [20, 21], within which the
bacteria multiply. The mature phagosome, termed the *Coxiella*-containing vacuole (CCV), is crucial
to *Coxiella* metabolism, replication, and pathogenesis [14, 22, 23].
Previously, using a ratiometric fluorescence-based pH measurement assay, we demonstrated
that although the CCV is acidic, further escalation in CCV acidity is bactericidal to *Coxiella* [20] and
therefore, *Coxiella* actively regulates CCV acidity in a type 4B secretion system (T4BSS)-dependent
manner [21]. Specifically, we showed that *Coxiella* T4BSS inhibits the host endosomal maturation
pathway, markedly reducing the number of acidic, mature endosomes and lysosomes available for
fusion with the CCV [21]. Since the CCV attains its luminal acidity by fusing with mature
endosomes and lysosomes, inhibiting endosomal maturation indirectly aids *Coxiella* in regulating the
CCV pH [21]. One notable outcome of *Coxiella* inhibition of endosomal maturation is the increase in
mean endosomal pH (pH ~5.2) in the infected cells. Since mature endosomes have a significantly
lower pH (~ 4.7), we define the endosomes with an elevated pH as 'immature' endosomes. Because
endosomal maturation is crucial to cargo transport and degradation [24], the 'immature' endosomes
in *Coxiella*-infected cells may be impaired in these functions and therefore detrimental to host cells.
However, the fate and the trafficking pattern of the 'immature' endosomes in *Coxiella*-infected cells
remained unclear. In this study, we aim to characterize the endosomal trafficking in *Coxiella*-infected
cells by using HeLa cells expressing endosome-specific fluorescent markers analyzed by quantitative
confocal microscopy.
The Ras-associated binding (Rab) proteins are the largest family in the Ras superfamily
GTPases and are the master regulators of all vesicular trafficking in mammalian cells [25]. Rab
proteins localize on the cytosolic surface of endosomes and perform functions like cargo selection
and sorting, vesicle transport along microtubules, and vesicle tethering and fusion with target
organelles [25]. The human genome encodes more than 60 Rab proteins that are associated with
specific endosome populations to regulate their trafficking, sorting, and recycling events. Rab
proteins cycle between an active, GTP-bound, membrane-recruited form and an inactive, GDP-
bound, cytosolic form to associate and dissociate with specific endosomes [24, 25]. Therefore, the
localization of a specific Rab protein provides key information about the trafficking pattern of a
population of endosomes in mammalian cells.

To understand the trafficking of the *Coxiella*-generated 'immature' endosomes, we ectopically
expressed EGFP-tagged Rab proteins in the infected host cells and analyzed their localization by
confocal microscopy. To our surprise, we observed that the CCV, in addition to acquiring Rab5 and
Rab7, acquired the recycling endosomal markers Rab11a and Rab35, suggesting that the CCV also
interacts with the host recycling endosomes during maturation. Furthermore, we observed an increase
in Rab11a and Rab35-positive endosomes in the infected cells, indicating that *Coxiella* enhances the
recycling endosome content in the infected cells. Finally, silencing Rab11a and Rab35 using siRNA
resulted in significantly smaller CCVs, suggesting that recycling endosomes facilitate CCV
expansion. Our study is the first report suggesting a CCV-host recycling endosome interaction
indicating the importance of recycling endosomes in the *Coxiella* intracellular life cycle.

2   Materials and methods

2.1   Bacteria and mammalian cells

Coxiella burnetii Nine Mile Phase II (NMII) (Clone 4, RSA 439) and mCherry-expressing *C. burnetii*
NMII [26] were grown and stored as previously described [21]. Human cervical epithelial cells
(HeLa, ATCC CCL-2) were maintained in Roswell Park Memorial Institute (RPMI) 1640 medium
with glutamine (Cat. 10-040-CV, Corning, New York, NY) and 10% fetal bovine serum (FBS; Cat.
S11150, R&D Systems, Inc., Minneapolis, MN). The genome equivalents (GE) for the *Coxiella*
stocks were determined by qPCR using primers for the *Coxiella burnetii dotA* gene [27]. The
multiplicities of infection (MOI) of the bacteria stocks were optimized for HeLa cells and each type
of culture vessels to ~1 internalized bacterium per cell at 37$°$C and 5% CO_2.

2.2   Preparation of EGFP-tagged Rab plasmids

The plasmids encoding EGFP-tagged Rab proteins were generous gifts from Dr. Mary Weber,
University of Iowa [28]. The plasmids were first transformed into competent *E. coli* DH5α (Cat.
C2988J, New England Biolabs, Ipswich, MA) and stored at -80$°$C. A single colony of the
transformed *E. coli* was grown in 5 mL lysogeny broth (LB) containing 50 µg/mL kanamycin or 100
µg/mL carbenicillin overnight, which was then used to inoculate a 150 mL LB culture and incubated
overnight. The GeneJET Endo-free Plasmid Maxiprep Kit (Cat. K0861, Thermo Fisher Scientific
Baltics UAB, Lithuania) was used to isolate and purify plasmid DNA from bacterial cultures
according to the manufacturer's protocol. Plasmids were then concentrated using the MilliporeSigma
Amicon Ultra-0.5 Centrifugal Filter Units (Cat. UFC503096, Fisher Scientific, Waltham, MA) and
stored at -20$°$C until used for cell transfection.

2.3   Transfection and immunofluorescent assay

HeLa cells, grown in RPMI with 10% FBS (hereafter referred to as 10% RPMI), were plated in a 24-
well plate ($2x10^4$ cells per well) and simultaneously transfected with 0.4 µg of respective EGFP-Rab
plasmids using FuGENE 6 transfection reagent (Cat. E269A, Promega, Madison, WI) per the
manufacturer's protocol. At 24 hours post-transfection, the cells were infected with mCherry-
Coxiella in 250 uL 10% RPMI for 1 h, washed extensively with sterile phosphate-buffered saline
(PBS; HyClone, Cat. SH30256.01, Cytiva, Marlborough, MA), and incubated in 10% RPMI at 37$°$C.
On the day before the indicated time points, cells were trypsinized, counted, and diluted to $1x10^5$
cells/mL. Diluted cells were then plated onto coverslips placed in a separate 24-well plate ($5x10^4$
cells per coverslip) and allowed to adhere overnight. The next day, cells were fixed in 2.5%
paraformaldehyde (PFA; Cat. 15710, Electron Microscopy Sciences, Hatfield, PA) for 15 min,
washed in PBS, and blocked/permeabilized for 20 min in 1% bovine serum albumin (BSA; Cat.

BP9700-100, Fisher BioReagents, Pittsburgh, PA) and 0.1% saponin (Cat. S0019, TCI America,
Portland, OR) in PBS. Cells were then incubated in rabbit anti-LAMP1 (1:1000; Cat. PA1-654A,
Thermo Fisher Scientific, Waltham, MA) primary antibody for 1h followed by Alexa Fluor secondary
antibody (1:1000; Life Technologies) for 1 h to label the CCV. Following washing with PBS,
coverslips were mounted using ProLong Gold Antifade Mountant with 4', 6'-diamidino-2
phenylindole (DAPI; Cat. P36941, Invitrogen, Carlsbad, CA) and visualized on a Leica Stellaris
confocal microscope using a 63x oil immersion objective. Images were processed and analyzed in
ImageJ FIJI [29].

2.4 Enumeration of Rab11a/Rab35-positive CCVs

HeLa cells were plated in a 24-well plate and transfected with plasmids encoding EGFP-tagged
Rab11a or Rab35 using FuGENE 6 transfection reagent according to the manufacturer's protocol. At
24 hours post-transfection, the cells were infected with mCherry-*Coxiella* in 250 μL 10% RPMI for 1
h, washed in PBS, and incubated in 10% RPMI. On the day before the indicated time points, cells
were trypsinized, diluted to 1×10^5 cells/mL, and plated onto coverslips in a 24-well plate (5×10^4 cells
per coverslip). The next day, the cells were subjected to immunofluorescent assay (IFA) where they
were fixed with 2.5% PFA for 15 min and blocked/permeabilized with 1% BSA in PBS with 0.1%
saponin for 20 min. Cells were then incubated in rabbit anti-LAMP1 (1:1000; Cat. PA1-654A,
Thermo Fisher) primary antibody for 1 h followed by Alexa Fluor 647 goat anti-rabbit secondary
antibody (1:1000; Cat. A21245, Invitrogen) for 1 h to label the CCVs. Following washing with PBS,
the coverslips were mounted with ProLong Gold Antifade Mountant with DAPI. The cells were
visualized with a Leica Stellaris confocal microscope using a 63x oil immersion objective and images
were imported and analyzed in ImageJ FIJI. A Rab-LAMP1 colocalization at the CCV membrane
was considered positive for Rab trafficking to the CCV. Approximately 100 CCVs were scored for
Rab11a and Rab35 localization and the percentage of Rab11a- and Rab35-positive CCVs were
determined. The statistical difference in the percentage of Rab11a and Rab35-positive CCVs at each
time-point was determined by Welch's t-test from three independent experiments (N=3).

2.5 Quantification of Rab11a and Rab35 content in *Coxiella*-infected cells

HeLa cells were either mock-infected or infected with mCherry-*Coxiella* in 24-well plates (2.5×10^4
cells per well; two wells per condition) for 1 h, washed extensively with PBS, and incubated in 10%
RPMI. For 3-day experiments, the cells were trypsinized at 2 dpi, diluted to 1×10^5 cells/mL, and
replated onto coverslips in a new 24-well plate (5×10^4 cells per coverslip). At 3 dpi, cells were fixed
with 2.5% PFA for 15 min and washed thoroughly with PBS. For 6-day experiments, cells were
trypsinized, diluted, and replated first at 3 dpi and again at 5 dpi (5×10^4 cells per coverslip). Next day,
cells were fixed with 2.5% PFA and washed thoroughly with PBS. For both 3-day and 6-day
experiments, following fixation, the cells were blocked/permeabilized with 1% BSA in PBS with
0.1% saponin for 20 min. Cells were then incubated overnight at 4°C in rabbit anti-Rab35 (1:100;
Cat. 11329-2-AP, Proteintech, Rosemont, IL) or for 1 h at room temperature (RT) in rabbit anti-
Rab11a (1:500; Cat. 20229-1-AP, Proteintech) primary antibody. The coverslips were washed with
PBS and then incubated in Alexa Fluor 488 goat anti-rabbit (1:1000; Cat. A11034, Invitrogen)
secondary antibody for 1 h at RT. Upon washing with PBS, the coverslips were mounted using
ProLong Gold with DAPI. Cells were visualized with a Leica Stellaris confocal microscope using a
63x oil immersion objective. To quantify Rab content from mock and WT-infected cells, identically
captured and processed images were imported into ImageJ FIJI and the total fluorescent intensity of
Rab11a or Rab35 was measured and normalized to cell area. At least 25 cells were analyzed per

condition in each of three independent experiments (N=3). The statistical difference in Rab11a and
Rab35 intensities between mock and *Coxiella*-infected cells was measured by Welch's t-test.

2.6   RNA interference and immunoblotting

HeLa cells (10^5 cells per well, 6-well plate) were reverse transfected with 50 nM small-interfering
RNA (siRNA) SMARTpools specific for human Rab11a (ON-TARGETplus, Cat. L-004726-00-0005,
Dharmacon Inc., Horizon Discovery Ltd., Lafayette, CO), Rab35 (ON-TARGETplus, Cat. L-009781-
00-0005, Dharmacon Inc.), or a non-targeting (NT) control (Cat. D-001810-10-05, Dharmacon Inc.)
using DharmaFECT-1 transfection reagent (Cat. T-2001-01, Dharmacon Inc.) in RPMI with 5% FBS
according to the manufacturer's protocol. At 48 h post-transfection, cells were infected with
mCherry-*Coxiella* for 1 h at 37°C. Following washing with PBS, cells were harvested by
trypsinization and subjected to a second round of siRNA transfection in a 24-well plate ($2.5x10^4$ cells
per well). On the day before the indicated time points, cells were trypsinized, resuspended to $1x10^5$
cells/mL, and either replated onto coverslips ($2.5x10^4$ cells per coverslip) or into another 24-well
plate without coverslips. The next day, the cells on coverslips were fixed with 2.5% PFA and
subjected to immunofluorescent assay with rabbit anti-LAMP1 primary and Alexa Fluor 488
secondary antibodies to label the CCVs. Concurrently, mock-infected cells from the second 24-well
plate were lysed with RIPA lysis buffer (Cat. 89900, Thermo Fisher) supplemented with HALT
protease inhibitor (Cat. 87786, Thermo Fisher) on ice. The soluble fraction from the cell lysates were
collected and protein concentrations were determined using the Pierce Rapid Gold BCA kit (Cat.
A53226, Thermo Fisher). To confirm Rab11a and Rab35 silencing, 10 µg of total protein for both NT
and Rab siRNA-transfected samples were resolved in 4-20% gradient SDS-PAGE (Mini-PROTEAN
Gel, Cat. 4561094, Bio-Rad Laboratories, Hercules, CA) and subsequently transferred to a PVDF
membrane for immunoblotting (Immun-Blot PVDF Membrane, Cat. 1620174, Bio-Rad). After
blocking in Bio-Rad EveryBlot buffer (Cat. 12010020, Bio-Rad) for 5 min, the membranes were then
probed separately using rabbit anti-Rab11a (Cat. 20229-1-AP, Proteintech), rabbit anti-Rab35 (Cat.
11329-2-AP, Proteintech), and anti-β-tubulin (BT7R) (HRP-conjugated loading control; Cat. MA5-
16308-HRP, Invitrogen) antibodies in 1% BSA in PBS. After washing, the membranes were
incubated with horseradish peroxidase-conjugated goat anti-rabbit (1:1000; Cat. 31460, Thermo
Fisher) secondary antibody in EveryBlot blocking buffer, and developed using enhanced
chemiluminescence (ECL) reagent (Cat. 170-5060, Bio-Rad).

2.7   Measurement of CCV area

HeLa cells were reverse-transfected with NT control, Rab11a, or Rab35-specific siRNAs as
described above, and at 48 h post-transfection, were infected with mCherry-*Coxiella* for 1 h. Upon
washing with PBS, cells were subjected to a second round of siRNA transfection as described above.
On the day before the indicated time points, cells were trypsinized, resuspended to $1x10^5$ cells/mL,
and replated onto coverslips ($2.5x10^4$ cells/coverslip). The next day, the coverslips were fixed with
2.5% PFA and cells were immunostained with rabbit anti-LAMP1 primary antibody and Alexa Fluor
secondary antibody to label the CCVs. Mounted coverslips were visualized using a Keyence BZ-
X710 fluorescent microscope. Images of individual CCVs were captured and processed identically
between conditions, and the CCV areas from exported images were measured in ImageJ FIJI. A 20
µm scale bar placed on a representative image for each sample was used for setting the scale in
ImageJ. CCV areas were expressed as pixels and plotted using GraphPad Prism. At least 25 CCVs
were measured per condition for each of the three independent experiments (N=3). The statistical
differences in CCV areas between non-targeting and Rab11a/Rab35-specific siRNA-transfected cells
were measured by Welch's t-test.

2.8   Data analyses

Fluorescent and confocal images were processed and analyzed in ImageJ FIJI software [29]. All statistical analyses were performed using unpaired t-test with Welch's correction (Welch's t-test) using GraphPad Prism (GraphPad, La Jolla, CA). A p-value < 0.05 was considered significant. Data are plotted as mean ± standard error of mean (SEM).

3   Results

3.1   Recycling endosomal markers Rab11a and Rab35 localize to the CCV during infection

We previously observed that *Coxiella* actively regulates the CCV acidity by inhibiting host endosomal maturation in a T4BSS-dependent manner [21]. However, by impeding endosomal maturation, *Coxiella* gives rise to population of 'immature' endosomes with a higher mean pH compared to mature endosomes. As the immature endosomes may be impaired in cargo trafficking and degradation, they may be detrimental to the host cells. However, whether *Coxiella* manipulates the trafficking of the 'immature' endosomes in the infected cells has not been clear. Since Rab proteins are the master regulators of endocytic trafficking, we hypothesized that *Coxiella* manipulates one or more Rab proteins to modulate endosomal trafficking in the infected cells. To test this, we screened 13 major Rab proteins in *Coxiella*-infected cells for their interaction with the CCV. Plasmids encoding EGFP-tagged Rab proteins (Table 1; a gift from Dr. Mary Weber, University of Iowa) were maxiprepped from *E. coli* culture and stored in endotoxin free elution buffer. HeLa cells were transfected with the EGFP-Rab plasmids and infected with mCherry-*Coxiella*. At 3 dpi, the cells were immunostained with LAMP1 primary antibody to label the CCVs. The cells were then visualized with a Leica Stellaris confocal microscope where 488 and 647 nm lasers were used for imaging the Rab proteins and LAMP1, respectively. Images were processed and analyzed in ImageJ FIJI and a Rab-LAMP1 colocalization at the CCV was considered positive for Rab trafficking to the CCV. EGFP tagged-Rab5 and Rab7 were included as positive controls in the screen as Rab5 and Rab7 are known to localize to the CCV during infection [30, 31]. As expected, all CCVs were positive for Rab5, Rab7, and LAMP1. In addition to that, we interestingly observed that some CCVs were also positive for Rab11a and Rab35 as evidenced by their colocalization with LAMP1 at the CCV membrane (Figure 1). Both Rab11a and Rab35 are associated with host recycling endosomes, where they regulate the 'slow' and 'fast' endosomal recycling pathways, respectively [32-34]. Therefore, these data suggest that, in addition to interacting with early endosomes, late endosomes, and lysosomes, at least a subset of CCVs also interacts with the host recycling endosomes in both the 'slow' and the 'fast' recycling pathways. A CCV area comparison between the Rab11a/Rab35-positive and the negative CCVs showed no significant difference in areas between the positive and the negative CCVs (data not shown). Moreover, *Coxiella* colonies appeared unaffected by the presence of Rab11a and Rab35 at the CCV. This study is the first report suggesting that the CCV interacts with host recycling endosomes and acquires recycling endosomal markers during maturation.

Table 1. EGFP-tagged Rab proteins used in this study

Rab proteins	Cellular localization	Membrane traffic function	References	Traffics to CCV
Rab1a	ER, Golgi	ER to Golgi	[35]	No
Rab4a	Early endosome	Protein recycling, transport to PM	[36]	No
Rab5	Early endosome, PM	Early endosome fusion	[37]	Yes

Rab6a	Golgi	Endosome to Golgi, intra-Golgi transport	[38]	No
Rab7	Late endosomes, lysosomes	Late endosome to lysosome	[37]	Yes
Rab8	Cell membrane, primary cilia	Exocytosis, TGN to PM	[39]	No
Rab9	Late endosomes, Golgi	Endosome to TGN	[40]	No
Rab11a	Recycling endosomes, Golgi	TGN/RE to PM	[41]	
Rab14	Golgi, early endosomes	TGN/RE to PM	[42]	No
Rab18	Golgi, lipid droplets	Lipid droplet formation	[43]	No
Rab33	Golgi	Autophagosome formation	[44]	No
Rab34	Golgi	Intra-Golgi transport	[45]	No
Rab35	Recycling endosomes, PM	RE to PM	[46]	Yes

255 ER – Endoplasmic reticulum; PM – plasma membrane; TGN – Trans-Golgi Network; RE –
256 Recycling endosomes

3.2 CCV-Rab11a and CCV-Rab35 interactions are dynamic and temporally regulated

In our Rab localization screen, we interestingly observed that only a subset of the CCVs were
positive for Rab11a or Rab35 at a given time-point. This led us to hypothesize that the interaction
between the CCV and Rab11a/Rab35 are dynamic in nature. To test this, we quantified the
percentage of Rab11a- and Rab35-positive CCVs in a population of *Coxiella*-infected HeLa cells at 3
and 6 dpi. Rab11a- or Rab35-positive and negative CCVs were identified in ImageJ FIJI by a
colocalization of Rab and LAMP1 at the CCV membrane, as described above (Figure 2A). Our
quantitation data revealed that at 3 dpi, ~52% of CCVs were positive for Rab11a, whereas only ~39%
of CCVs were positive for Rab35 (Fig. 2B-C). Interestingly this pattern reversed at 6 dpi, when only
~22% of CCVs were positive for Rab11 and ~64% of CCVs were positive for Rab35 (Fig. 2B-C).
Together, these data suggest that, while CCVs preferentially interact with the Rab11a-positive
endosomes at 3pi, at 6 dpi, the preferential interaction shifts towards the Rab35-positive endosomes.
Therefore, our data indicate that Rab11a and Rab35 interaction with the CCV are dynamic and varies
depending on the stage of infection.

3.3 *Coxiella* infection leads to increased recycling endosome content in infected cells

Our observation that *Coxiella* interacts with two different populations of recycling endosomes in the
host cells led us to hypothesize that *Coxiella* manipulates host recycling endosomes during infection.
To test this hypothesis, we quantitated the total cellular Rab11a and Rab35 content in mock and
Coxiella-infected HeLa cells by immunofluorescent assay with anti-Rab11a or anti-Rab35 antibody.
EGFP-Rab11 or EGFP-Rab35 transfected HeLa cells were not used in this experiment because of
potential differences in ectopic expression level of Rab proteins between individual cells. Mock and
Coxiella-infected HeLa cells were immunostained using anti-Rab11a or anti-Rab35 antibodies at 3
and 6 dpi and identically captured images were analyzed in ImageJ FIJI. The total fluorescent
intensity of Rab11 or Rab35 was measured from mock and *Coxiella*-infected cells normalized to cell
area and were considered indicative of total recycling endosome content in each cell. Our data
revealed a ~19% increase in Rab11a fluorescent intensity ($p < 0.0001$; Figures 3A, B) and a ~27%
increase in Rab35 fluorescent intensity in infected cells compared to mock at 3 dpi ($p < 0.0001$;
Figures 4A, B). However, to our surprise, we did not observe any difference in Rab11a or Rab35

fluorescent intensities between mock and *Coxiella*-infected cells at 6 dpi (Figures 3C, 4C). These
data suggest that *Coxiella* increases the recycling endosome content in infected host cells at 3 dpi, but
not at 6 dpi.

3.4   Rab11a and Rab35 silencing inhibits CCV expansion

Based on our observation that the CCV dynamically interacts with recycling endosome-associated
Rab11a and Rab35, and that *Coxiella* increases the recycling endosome content of infected cells, we
hypothesized that Rab11a and Rab35 positively affects CCV maturation. To test this, we examined
CCV expansion in Rab11a or Rab35 knockdown HeLa cells compared to a non-targeting (NT)
control using our dual-hit siRNA knockdown protocol described previously (Figure 5A) [21]. Since
CCV interaction with Rab11a is higher at 3 dpi, and that with Rab35 is higher at 6 dpi, we measured
the CCV areas in Rab11a and Rab35 knockdown cells at 3 dpi and 6 dpi, respectively, to detect
maximum effect of the respective knockdowns. Immunoblotting with cell lysates from NT, siRab11a,
and siRab35-transfected cells revealed significant depletion of both Rab11a (Figure 5B) and Rab35
(Figure 5C) at 3 and 6 dpi, respectively. The CCV area measurement data revealed a ~45% reduction
in CCV area in Rab11a knockdown cells compared to NT ($p<0.0001$) at 3 dpi (Figure 5D), and a
~28% reduction in CCV area in Rab35 knockdown cells ($p<0.0001$) at 6 dpi (Figure 5E). These data
suggest that both Rab11a and Rab35 positively affect CCV expansion at 3 and 6 dpi, respectively,
indicating that host recycling endosomes are essential for CCV maturation.

4   Discussion

We previously established that *Coxiella* T4BSS inhibits host endosomal maturation for bacterial
survival and pathogenesis. However, in the process, *Coxiella* generates a population of 'immature'
endosomes with an elevated mean pH relative to mature endosomes. To generate a comprehensive
understanding of endosomal trafficking in the *Coxiella*-infected cells, we screened 13 well-studied
Rab proteins to assess their localization in *Coxiella*-infected cells. We discovered that Rab11a and
Rab35, the molecular markers of recycling endosomes, localize to the CCV, although only a subset of
CCVs were found positive for Rab11a or Rab35 at a given time point. Our data also revealed that
Coxiella increases the recycling endosome content of the infected cells at 3 dpi, and Rab11a and
Rab35 knockdown resulted in significantly smaller CCVs at 3 and 6 dpi, respectively, suggesting that
host recycling endosomes are essential for CCV expansion.
314   Mammalian endosomal trafficking can be broadly classified into degradative and recycling
pathways [47]. While the degradative pathway traffics the cargo through early and late endosomes
and finally delivers it to lysosomes for degradation, the recycling pathway sorts and re-exports
essential membrane components including receptors, carrier molecules, and cell adhesion molecules
that have been internalized [33]. For the first time, we present findings suggesting an interaction
between the CCV and the host recycling endosomes. This intriguing observation implies that the
CCV not only engages with the host endocytic degradative endosomes but also with recycling
endosomes, potentially incorporating their contents and membranes during maturation. Mammalian
cells exhibit two simultaneous pathways for cargo recycling namely the 'fast' and the 'slow'
recycling pathways [48]. In the 'fast' pathway, the internalized cargo is directly transported back to
the plasma membrane from early endosomes. However, the 'slow' pathway first traffics the cargo to a
perinuclear vesicle known as the endocytic recycling compartment (ERC), within which, the cargo is
sorted and then redirected to the plasma membrane via recycling endosomes [33, 49]. Rab35
predominantly regulates the 'fast' recycling pathway [32, 34], while Rab11a serves as a major
regulator of the 'slow' pathway [33, 34, 50]. Therefore, our data suggest that the CCV interacts with
both the 'fast' and the 'slow' recycling pathways during maturation. Moreover, our observation that

Rab11a localizes to the CCVs significantly more at 3 dpi whereas Rab35 is more prevalent at CCVs
at 6 dpi suggest a preference for interaction with the 'slow' recycling pathway during the early stages
of maturation and a shift towards a preferential interaction with the 'fast' recycling pathway as the
CCV matures further.
While the significance of the interactions between CCV and the host recycling endosomes
remains uncertain, multiple studies propose that similar interactions with intracellular bacterial
vacuoles play a crucial role in providing nutrients to the vacuole or in ensuring vacuolar stability. For
example, *Chlamydophila pneumoniae*-containing inclusions recruit the mammalian Rab11/Rab14
adapter protein Fip2, facilitating the early-stage recruitment of Rab11/14 to the inclusion membrane
during infection [51, 52]. Rab11/14 positive recycling endosomes trafficked nutrients, such as
transferrin to the proximity of *C. pneumoniae* inclusion, although whether transferrin is delivered
into the inclusion lumen remained unclear [53-55]. Indeed, a recent study with fluorescent transferrin
(Tf488) revealed that in *Coxiella*-infected HeLa cells, Tf488-containing vesicles also trafficked to the
proximity of CCVs [56]. Moreover, Rab11-positive recycling endosomes trafficked transferrin in
K562 cells [50]. Therefore, it is possible that recycling endosomes traffic transferrin to the CCV
aiding in its maturation. Further experimentation is needed to examine this intriguing hypothesis.
Another possible benefit of intracellular pathogens subverting recycling endosomes is to augment
vacuole stability. For example, *Chlamydia trachomatis* inclusion protein CT229 interacts with Rab35
to maintain the inclusion integrity [28], and *Legionella pneumophila*-containing vacuoles (LCV)
acquire Rab11 and Rab35 for stability [57]. It is possible that Rab11 and Rab35 plays a critical role in
maintaining CCV stability. However, we did not observe a disintegration of CCV in Rab11a and
Rab35 knockdown HeLa cells, raising a possibility that *Coxiella* employes redundant mechanisms to
maintain CCV stability.
As mentioned before, we observed that only a subset of CCVs are positive for Rab11a or
Rab35 at a given time point, and the proportion of Rab11a- and Rab35-positive CCVs in a population
of infected cells vary over time. previous studies including ours have only observed stable fusion
events between the CCV and host endosomes and lysosomes [14, 20, 21, 58]. The dynamic nature of
CCV-Rab11a/Rab35 interactions could be a result of two possibilities: first, only a subset of CCVs
preferentially interacts with the recycling endosomes at a given time point and the number of
preferred CCVs interacting with a particular Rab protein varies depending on the stage of infection,
and second, it is possible that the recycling endosomes make transient contact with the CCVs in a
'kiss-and-run' manner [59], to release their content into the CCV and/or to receive recyclable cargo
from the CCV and then dissociate. However, further experimentation is required to test these
hypotheses and could be an intriguing area of future investigation.
Our observation that *Coxiella* infection led to an increased recycling endosome content at 3
dpi builds upon the previous observation that *Coxiella* expands the endosomal compartment in
infected cells [56]. It is possible that increased number of recycling endosomes accumulate more
Tf488 in *Coxiella*-infected cells observed in that study [56]. Interestingly, in our study, we did not
observe an increase in the recycling endosome content at 6 dpi. This may be a result of an already
low number of endosomes in the *Coxiella*-infected cells at 6 dpi [21]. Nonetheless, these data, taken
together with our previous data indicate that *Coxiella* has a dual effect on host endosomes: while it
reduces the number of mature endosomes in infected cells [21], *Coxiella* simultaneously elevates the
number of recycling endosomes during infection. While the full characterization of these regulations
requires further testing, we propose a model that *Coxiella* reroutes the 'immature' endosomes from
the degradative pathway into the recycling pathway, potentially protecting the host cell from the
detriments of the 'immature' endosomes.
Our siRNA-mediated knockdown experiments revealed that Rab11a and Rab35 are essential
for CCV expansion at 3 and 6 dpi, respectively. A recent study with Rab11a knockout HeLa cells
revealed that Rab11a deficiency increases the number of late endosomes and lysosomes within the

cells [60]. It is possible that in our experiments, Rab11a knockdown led to an increased number of
lysosomes that is detrimental to CCV acidity and *Coxiella* growth. Future research analyzing the
lysosomal pH and enzyme activity in *Coxiella*-infected, Rab11a/Rab35 knockdown cells will help
further characterize the CCV-recycling endosome interactions.
Overall, our data revealed novel host-*Coxiella* interactions that appear to be beneficial for
CCV maturation. Future research into characterizing the role of Rab11a and Rab35 in *Coxiella*
pathogenesis, and the molecular mechanisms of *Coxiella* enhancement of recycling endosome
content will potentially identify novel therapeutic targets for *Coxiella*.

5   Acknowledgments

We thank Stacey Gilk (University of Nebraska Medical Center) for providing the parental HeLa cells
and mCherry-*Coxiella burnetii*. We are also thankful to Mary Weber (University of Iowa) for
providing the EGFP-tagged Rab plasmids. We thank Manan Damani (Midwestern University) for
assistance with confocal and fluorescent microscopy. We also thanks Vijitha Kantety and Ruthvik
Gundala for helpful suggestions.

393
394
Figure 1. Host recycling endosome-associated proteins Rab11a and Rab35 localize to the CCV.
Representative images of *Coxiella*-infected HeLa cells, expressing EGFP-tagged Rab proteins. HeLa
cells were transfected with plasmids encoding EGFP-Rab proteins and at 24 h post-transfection, were
infected with mCherry-*Coxiella* for 1 h. At 3 dpi, cells plated on coverslips were fixed and subjected
to immunofluorescent assay with anti-LAMP1 antibody to label the CCVs. Cells were visualized
with a Leica Stellaris confocal microscope and images were analyzed in ImageJ FIJI. CCVs, where
LAMP1 and Rab proteins colocalized at the membrane (shown in insets) were considered positive for
a specific Rab protein. As expected, CCVs were positive for both Rab5 and Rab7 at 3 dpi.
Additionally, we observed that Rab11a and Rab35 also localized to CCVs. Notably, only a subset of
CCVs were positive for Rab11a or Rab35.

405
406
Figure 2. CCV-Rab11a and CCV-Rab35 interactions are dynamic and temporally regulated.
(A) Representative images showing Rab11a and Rab35-positive and negative CCVs. HeLa cells were
transfected with plasmids encoding EGFP-Rab11a or EGFP-Rab35 and at 24 h post-transfection,
were infected with mCherry-*Coxiella* for 1 h. At 3 dpi, cells plated on coverslips were fixed and
subjected to immunofluorescent assay with anti-LAMP1 antibody to label the CCVs. Cells were
visualized with a Leica Stellaris confocal microscope and images were analyzed in ImageJ FIJI.
CCVs, where LAMP1 and Rab proteins colocalized at the membrane (shown in insets) were
considered positive for a specific Rab protein. CCVs were scored for Rab11a or Rab35 localization
and percentage of positive CCVs were calculated. **(B-C)** Quantification of Rab11a- and Rab35-
positive and negative CCVs at 3 and 6 dpi revealed that Rab11a localized to significantly more CCVs
at 3 dpi compared to 6 dpi, whereas the pattern reversed for Rab35, which localized to more CCVs at
6 dpi. Data shown as mean±SEM of at least 100 infected cells per condition in each of three
independent experiments (N=3) as analyzed by Welch's t-test; *, P<0.05.

Figure 3. *Coxiella* increases Rab11a content in infected host cells. (A) Representative images showing Rab11a-positive recycling endosome content of mock and *Coxiella*-infected HeLa cells at 3 and 6 dpi. HeLa cells were mock infected or infected with mCherry-*Coxiella* for 1 h. On the days before the indicated time points, the cells were plated on coverslips. The next day, cells were fixed and subjected to immunofluorescent assay with anti-Rab11a antibody to uniformly stain the Rab11a-positive recycling endosomes. Cells were visualized with a Leica Stellaris confocal microscope and the images were analyzed in ImageJ FIJI. Identically captured images were imported into ImageJ FIJI and the total fluorescent intensity of Rab11a was measured and normalized to cell area. **(B-C)** Quantification of Rab11a fluorescent intensity revealed an increase in Rab11a content in *Coxiella*-infected cells at 3 dpi but not at 6 dpi. Data shown as mean±SEM of at least 25 cells per condition in each of three independent experiments (N=3) as analyzed by Welch's t-test; ****, P<0.0001; ns, non-significant.

434
Figure 4. *Coxiella* increases Rab35 content in infected host cells. (A) Representative images
showing Rab35-positive recycling endosome content of mock and *Coxiella*-infected HeLa cells at 3
and 6 dpi. HeLa cells were mock infected or infected with mCherry-*Coxiella* for 1 h. On the days
before the indicated time points, the cells were plated on coverslips. The next day, cells were fixed
and subjected to an immunofluorescent assay with anti-Rab35 antibody to uniformly stain the Rab35-
positive recycling endosomes. Cells were visualized with a Leica Stellaris confocal microscope and
the images were analyzed in ImageJ FIJI. Identically captured images were imported into ImageJ FIJI
and the total fluorescent intensity of Rab35 was measured and normalized to cell area. **(B-C)**
Quantification of Rab35 fluorescent intensity revealed an increase in Rab35 content in *Coxiella*-
infected cells at 3 dpi but not at 6 dpi. Data shown as mean±SEM of at least 25 cells per condition in
each of three independent experiments (N=3) as analyzed by Welch's t-test; ****, P<0.0001; ns, non-
significant.

447
448
Figure 5. Rab11a and Rab35 are essential for CCV expansion. (A) Schematic diagram of dual
siRNA transfection protocol in HeLa cells to silence Rab11a and Rab35. Cells were transfected with
Rab11a or Rab35-specific siRNA (siRab11a or siRab35, respectively) or a control non-targeting
siRNA (NT) and incubated. At 48 h post-transfection cells were infected with mCherry-*Coxiella* and
subjected to a second round of siRNA transfection. At the indicated time points, cells were fixed and
subjected to immunofluorescent assay with anti-LAMP1 antibody to label the CCVs. **(B-C)**
Immunoblots revealed depletion of both Rab11a and Rab35 in siRNA-transfected cells at 3 and 6 dpi,
respectively. **(D-E)** measurement of the CCV area revealed significantly smaller CCVs in siRab11a-
transfected cells at 3 dpi and siRab35-transfected cells at 6 dpi. Data shown as mean±SEM of at least
25 cells per condition in each of three independent experiments (N=3) as analyzed by Welch's t-test;
****, P<0.0001; ns, non-significant.

REFERENCES

460
461
1.   Maurin, M. and D. Raoult, *Q fever.* Clin Microbiol Rev, 1999. **12**(4): p. 518-53.

2.   Eldin, C., et al., *Q fever in French Guiana.* Am J Trop Med Hyg, 2014. **91**(4): p. 771-6.

3.   Angelakis, E., et al., *Coxiella burnetii-positive PCR in febrile patients in rural and urban*
Africa. Int J Infect Dis, 2014. **28**: p. 107-10.

4.   White, B., T. Brooks, and R.A. Seaton, *Q fever in military and paramilitary personnel in*
conflict zones: case report and review. Travel Med Infect Dis, 2013. **11**(2): p. 134-7.

5.   Amitai, Z., et al., *A large Q fever outbreak in an urban school in central Israel.* Clin Infect
Dis, 2010. **50**(11): p. 1433-8.

6.   Porter, S.R., et al., *Q fever in Japan: an update review.* Vet Microbiol, 2011. **149**(3-4): p. 298-
306.

7.   Lyytikainen, O., et al., *An outbreak of sheep-associated Q fever in a rural community in*
Germany. Eur J Epidemiol, 1998. **14**(2): p. 193-9.

8.   Alonso, E., et al., *Q Fever Outbreak among Workers at a Waste-Sorting Plant.* PLoS One,
2015. **10**(9): p. e0138817.

9.   Bjork, A., et al., *First reported multistate human Q fever outbreak in the United States, 2011.*
Vector Borne Zoonotic Dis, 2014. **14**(2): p. 111-7.

10.   Nett, R.J., S.D. Helgerson, and A.D. Anderson, *Clinician assessment for Coxiella burnetii*
infection in hospitalized patients with potentially compatible illnesses during Q fever
outbreaks and following a health alert, Montana, 2011. Vector Borne Zoonotic Dis, 2013.
13(2): p. 128-30.

11.   Graham, J.G., et al., *Development of an Ex Vivo Tissue Platform To Study the Human Lung*
Response to Coxiella burnetii. Infect Immun, 2016. **84**(5): p. 1438-1445.

12.   Dragan, A.L., R.C. Kurten, and D.E. Voth, *Characterization of Early Stages of Human*
Alveolar Infection by the Q Fever Agent Coxiella burnetii. Infect Immun, 2019. **87**(5).

13.   Howe, D. and L.P. Mallavia, *Coxiella burnetii exhibits morphological change and delays*
phagolysosomal fusion after internalization by J774A.1 cells. Infect Immun, 2000. **68**(7): p.
3815-21.

14.   Voth, D.E. and R.A. Heinzen, *Lounging in a lysosome: the intracellular lifestyle of Coxiella*
burnetii. Cell Microbiol, 2007. **9**(4): p. 829-40.

15.   Magunda, F., et al., *Anaplasma marginale actively modulates vacuolar maturation during*
intracellular infection of its tick vector, Dermacentor andersoni. Appl Environ Microbiol,
2016. **82**(15): p. 4715-4731.

16. Huang, B., et al., *The Anaplasma phagocytophilum-occupied vacuole selectively recruits Rab-*
GTPases that are predominantly associated with recycling endosomes. Cell Microbiol, 2010.
12(9): p. 1292-307.

17. Clemens, D.L. and M.A. Horwitz, *Characterization of the Mycobacterium tuberculosis*
phagosome and evidence that phagosomal maturation is inhibited. J Exp Med, 1995. **181**(1):
p. 257-70.

18. Horwitz, M.A., *The Legionnaires' disease bacterium (Legionella pneumophila) inhibits*
phagosome-lysosome fusion in human monocytes. J Exp Med, 1983. **158**(6): p. 2108-26.

19. Connor, M.G., et al., *Yersinia pestis Requires Host Rab1b for Survival in Macrophages.* PLoS
Pathog, 2015. **11**(10): p. e1005241.

20. Mulye, M., et al., *Elevated Cholesterol in the Coxiella burnetii Intracellular Niche Is*
Bacteriolytic. mBio, 2017. **8**(1).

21. Samanta, D., et al., *Coxiella burnetii Type 4B Secretion System-dependent manipulation of*
endolysosomal maturation is required for bacterial growth. PLoS Pathog, 2019. **15**(12): p.
e1007855.

22. Newton, H.J., J.A. McDonough, and C.R. Roy, *Effector protein translocation by the Coxiella*
burnetii Dot/Icm type IV secretion system requires endocytic maturation of the pathogen-
occupied vacuole. PLoS One, 2013. **8**(1): p. e54566.

23. Ghigo, E., M.I. Colombo, and R.A. Heinzen, *The Coxiella burnetii parasitophorous vacuole.*
Adv Exp Med Biol, 2012. **984**: p. 141-69.

24. Huotari, J. and A. Helenius, *Endosome maturation.* EMBO J, 2011. **30**(17): p. 3481-500.

25. Wandinger-Ness, A. and M. Zerial, *Rab proteins and the compartmentalization of the*
endosomal system. Cold Spring Harb Perspect Biol, 2014. **6**(11): p. a022616.

26. Beare, P.A., et al., *Characterization of a Coxiella burnetii ftsZ mutant generated by Himar1*
transposon mutagenesis. J Bacteriol, 2009. **191**(5): p. 1369-81.

27. Coleman, S.A., et al., *Temporal analysis of Coxiella burnetii morphological differentiation.* J
Bacteriol, 2004. **186**(21): p. 7344-52.

28. Faris, R., et al., *Chlamydia trachomatis CT229 Subverts Rab GTPase-Dependent CCV*
Trafficking Pathways to Promote Chlamydial Infection. Cell Rep, 2019. **26**(12): p. 3380-3390
e5.

29. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat
Methods, 2012. **9**(7): p. 676-82.

30. Beron, W., et al., *Coxiella burnetii localizes in a Rab7-labeled compartment with autophagic*
characteristics. Infect Immun, 2002. **70**(10): p. 5816-21.

31. Romano, P.S., et al., *The autophagic pathway is actively modulated by phase II Coxiella*
burnetii to efficiently replicate in the host cell. Cell Microbiol, 2007. **9**(4): p. 891-909.

32.   Chaineau, M., M.S. Ioannou, and P.S. McPherson, *Rab35: GEFs, GAPs and effectors.* Traffic,
2013. **14**(11): p. 1109-17.

33.   Grant, B.D. and J.G. Donaldson, *Pathways and mechanisms of endocytic recycling.* Nat Rev
Mol Cell Biol, 2009. **10**(9): p. 597-608.

34.   Ullrich, O., et al., *Rab11 regulates recycling through the pericentriolar recycling endosome.* J
Cell Biol, 1996. **135**(4): p. 913-24.

35.   Plutner, H., et al., *Rab1b regulates vesicular transport between the endoplasmic reticulum
and successive Golgi compartments.* J Cell Biol, 1991. **115**(1): p. 31-43.

36.   van der Sluijs, P., et al., *The small GTP-binding protein rab4 controls an early sorting event
on the endocytic pathway.* Cell, 1992. **70**(5): p. 729-40.

37.   Chavrier, P., et al., *Localization of low molecular weight GTP binding proteins to exocytic
and endocytic compartments.* Cell, 1990. **62**(2): p. 317-29.

38.   Goud, B., et al., *Small GTP-binding protein associated with Golgi cisternae.* Nature, 1990.
345(6275): p. 553-6.

39.   Huber, L.A., et al., *Rab8, a small GTPase involved in vesicular traffic between the TGN and
the basolateral plasma membrane.* J Cell Biol, 1993. **123**(1): p. 35-45.

40.   Yoshimura, S., et al., *Functional dissection of Rab GTPases involved in primary cilium
formation.* J Cell Biol, 2007. **178**(3): p. 363-9.

41.   Ren, M., et al., *Hydrolysis of GTP on rab11 is required for the direct delivery of transferrin
from the pericentriolar recycling compartment to the cell surface but not from sorting
endosomes.* Proc Natl Acad Sci U S A, 1998. **95**(11): p. 6187-92.

42.   Junutula, J.R., et al., *Rab14 is involved in membrane trafficking between the Golgi complex
and endosomes.* Mol Biol Cell, 2004. **15**(5): p. 2218-29.

43.   Dejgaard, S.Y., et al., *Rab18 and Rab43 have key roles in ER-Golgi trafficking.* J Cell Sci,
2008. **121**(Pt 16): p. 2768-81.

44.   Zheng, J.Y., et al., *A novel Rab GTPase, Rab33B, is ubiquitously expressed and localized to
the medial Golgi cisternae.* J Cell Sci, 1998. **111 (Pt 8)**: p. 1061-9.

45.   Wang, T. and W. Hong, *Interorganellar regulation of lysosome positioning by the Golgi
apparatus through Rab34 interaction with Rab-interacting lysosomal protein.* Mol Biol Cell,
2002. **13**(12): p. 4317-32.

46.   Kouranti, I., et al., *Rab35 regulates an endocytic recycling pathway essential for the terminal
steps of cytokinesis.* Curr Biol, 2006. **16**(17): p. 1719-25.

47.   O'Sullivan, M.J. and A.J. Lindsay, *The Endosomal Recycling Pathway-At the Crossroads of
the Cell.* Int J Mol Sci, 2020. **21**(17).

48. Hopkins, C.R. and I.S. Trowbridge, *Internalization and processing of transferrin and the*
transferrin receptor in human carcinoma A431 cells. J Cell Biol, 1983. **97**(2): p. 508-21.

49. Allgood, S.C. and M.R. Neunuebel, *The recycling endosome and bacterial pathogens.* Cell
Microbiol, 2018. **20**(7): p. e12857.

50. Green, E.G., et al., *Rab11 is associated with transferrin-containing recycling compartments in*
K562 cells. Biochem Biophys Res Commun, 1997. **239**(2): p. 612-6.

51. Molleken, K. and J.H. Hegemann, *Acquisition of Rab11 and Rab11-Fip2-A novel strategy for*
Chlamydia pneumoniae early survival. PLoS Pathog, 2017. **13**(8): p. e1006556.

52. Baetz, N.W. and J.R. Goldenring, *Rab11-family interacting proteins define spatially and*
temporally distinct regions within the dynamic Rab11a-dependent recycling system. Mol Biol
Cell, 2013. **24**(5): p. 643-58.

53. Leiva, N., A. Capmany, and M.T. Damiani, *Rab11-family of interacting protein 2 associates*
with chlamydial inclusions through its Rab-binding domain and promotes bacterial
multiplication. Cell Microbiol, 2013. **15**(1): p. 114-29.

54. Rejman Lipinski, A., et al., *Rab6 and Rab11 regulate Chlamydia trachomatis development*
and golgin-84-dependent Golgi fragmentation. PLoS Pathog, 2009. **5**(10): p. e1000615.

55. Ouellette, S.P. and R.A. Carabeo, *A Functional Slow Recycling Pathway of Transferrin is*
Required for Growth of Chlamydia. Front Microbiol, 2010. **1**: p. 112.

56. Larson, C.L. and R.A. Heinzen, *High-Content Imaging Reveals Expansion of the Endosomal*
Compartment during Coxiella burnetii Parasitophorous Vacuole Maturation. Front Cell
Infect Microbiol, 2017. **7**: p. 48.

57. Allgood, S.C., et al., *Legionella Effector AnkX Disrupts Host Cell Endocytic Recycling in a*
Phosphocholination-Dependent Manner. Front Cell Infect Microbiol, 2017. **7**: p. 397.

58. Howe, D., et al., *Maturation of the Coxiella burnetii parasitophorous vacuole requires*
bacterial protein synthesis but not replication. Cell Microbiol, 2003. **5**(7): p. 469-80.

59. Ryan, T.A., *Kiss-and-run, fuse-pinch-and-linger, fuse-and-collapse: the life and times of a*
neurosecretory granule. Proc Natl Acad Sci U S A, 2003. **100**(5): p. 2171-3.

60. Zulkefli, K.L., et al., *A role for Rab11 in the homeostasis of the endosome-lysosomal pathway.*
Exp Cell Res, 2019. **380**(1): p. 55-68.