Times birds need to evolve mating cues under allopatry and parapatry

Richard M. Sibly[1*] and Robert N. Curnow[2]

[1] School of Biological Sciences, University of Reading, UK. r.m.sibly@reading.ac.uk

[2] Department of Mathematics and Statistics, University of Reading, UK. r.n.curnow@reading.ac.uk

*Corresponding author. Mailing address School of Biological Sciences, University of Reading, Reading
RG66EX UK. Email r.m.sibly@reading.ac.uk

Running head: Times birds need to evolve mating cues

Keywords: Sexual imprinting; phenotype matching; pseudomagic traits; fixation time; fixation
probability; nascent speciation

Abstract

The time needed for the evolution of mating cues that distinguish species, such as species-specific
songs or plumage coloration, has received little attention. Here we show that the expected time
needed for a neutral substitution at a mating-cue locus under allopatry is approximately equal to the
reciprocal of mutation rate, but the time needed under parapatry can be much shorter. The time
needed is reduced by a factor of 4 x (population size) x (selection coefficient on the mating cue). In
the simplest case of parapatric speciation the mating cue is an adaptive trait which is acted on by
selection directly, but this does not in general result in species-specific songs or plumage coloration.
Alternatively, (Sibly and Curnow, 2022) showed that, with phenotype matching, local adaptation
induces selection on a mating cue even if the locus coding for the mating cue is not linked to those
producing local adaptation. Detailed analysis of two cases shows how mating cue selection then
depends on the migration rates between the locally-adapted populations and the selective
conditions within them. Comparison with data from molecular phylogenies suggests if mating cues
are selectively neutral, only parapatric, not allopatric, speciation can account for the times recorded
for the evolution of species-specific songs or plumage coloration.

Introduction

Speciation has been most extensively studied in birds. Bird species are generally distinguished by the
colour plumage and/or song used in mate choice, and these mating cues are generally learnt from
their parents or close kin in a process known as sexual imprinting. Sexual imprinting seems to be a
general feature of birds, shown to exist in over 100 species belonging to 15 different orders (ten Cate
and Vos, 1999; Verzijden et al., 2012). Novel mating cues that distinguish species evolve during the
process of speciation but the time needed for the evolution of novel mating cues, the subject of the
present paper, has received little attention. Analysis of DNA sequences shows that in some cases new
mating cues have evolved rapidly. Noteworthy examples occur in the tanagers (family Thraupidae)
including southern capuchinos seedeaters (*Sporophila*) and Darwin's finches. The Neotropical
southern capuchinos radiated within the past million years to form 10 predominantly sympatric
species that differ primarily in male plumage coloration and song (Campagna et al., 2017; Campagna
et al., 2012; Turbek et al., 2021). Eight of these species emerged in less than 50,000 generations
(Hejase et al., 2020). In Darwin's finches, radiations of ground and tree finches began around
100,000–300,000 years ago (Lamichhaney et al., 2015). The purpose of this paper is to estimate the
time to fixation of a neutral substitution for a novel mating cue under models of allopatric and

parapatric speciation described below. The estimated times to fixation will then be compared with
the above data.

Speciation in birds – and many other species – is currently considered generally to occur after a
period of allopatry, during which a population becomes split into geographically isolated
subpopulations which remain separate and diverge over thousands of generations, and eventually
can no longer interbreed (Tobias et al., 2020). Allopatric speciation is not the only possible speciation
mechanism, however, and in recent decades attention has been given to the role of divergent
natural selection, which leads to local adaptation and, potentially, speciation in the presence of gene
flow (see reviews in, e.g., (Janicke et al., 2019; Kopp et al., 2018)). Here in addition to allopatric
speciation we consider in particular the case of parapatric speciation. The difference between
allopatric and parapatric speciation is that the former occurs in a single environment whereas the
latter has migration between regions that experience different environmental selection pressures.
Parapatric speciation is thought to be more likely under phenotype matching than under alternative
methods of mate choice (Kopp et al., 2018). Phenotype matching means that individuals with a
particular trait tend to mate with others with the same trait, and is a likely result of sexual
imprinting. In this paper we assume that phenotype matching is complete, so individuals only mate
with others with the same trait. With phenotype matching, speciation in populations that initially
exchange genes (parapatric speciation) occurs readily in locally adapted populations.

We consider here two types of parapatric speciation. In the first, which we designate Type 1
parapatric speciation, adaptive traits are used as mating cues, i.e., are 'magic traits'. The term magic
trait refers to an adaptive trait that is used as a mating cue (see, e.g., (Kopp et al., 2018)). An
example is choosing mates on the basis of having a beak size that is advantageous in the niche in
which the population lives. The combination of phenotype matching and a magic trait is known to be
particularly favourable for speciation (Smadja and Butlin, 2011). Treating a related case, (Servedio
and Burger, 2020) used a deterministic haploid model to analyse what they term 'pseudomagic
traits', in which a mating cue locus and an ecological trait locus are separate but linked. (Servedio
and Burger, 2020) show that evolutionary outcomes are similar to Type 1 if the ecological trait locus
is tightly linked to the mating cue locus. (Sibly and Curnow, 2022) used a deterministic diploid model
to show that local adaptation induces selection on the mating cues even though the loci controlling
the mating cues are not linked to the loci controlling local adaptation. We refer to this as Type 2
parapatric speciation. Speciation occurs in (Sibly and Curnow, 2022)'s model because individuals in
each niche avoid mating with incomers who predominantly carry disadvantageous alleles. The model
incorporates a (Felsenstein, 1981) "two-allele mechanism" sensu (Butlin et al., 2021), with the key
difference that Felsenstein's two-allele models assume linkage between a locus controlling
assortative mating and local adaptation loci, whereas (Sibly and Curnow, 2022)'s does not. In (Sibly
and Curnow, 2022)'s model the loci are not close to each other in the genome—a situation allowing
much greater evolutionary potential. Whether the population will evolve towards speciation
depends in (Sibly and Curnow, 2022)'s model as in Felsenstein's on the balance between migration
and selection: some selection at the ecological locus is necessary. (Sibly and Curnow, 2022)'s results
were based on simulations that assumed complete phenotype matching, an infinite population
divided equally between two niches, migration rates the same in both directions, and local
adaptation the result of two alleles P and Q at a single diploid locus. The mating cue is controlled by
a single locus with only D alleles prior to a new mating cue produced by a C allele arising by
mutation. C is assumed dominant to D. We retain this notation below.

Our aim here is to use standard methods of population genetics to compare the expected times for
new mating cues to evolve under allopatry and parapatry. Bird speciation is used as a key example
though the results have wider application. Although avian coloration genetics has shown that many

genes may be involved in plumage coloration (Price-Waldman and Stoddard, 2021), the number of
possible mutations that could result in phenotypic change is not known. Our initial analysis is
therefore based on calculating times at a single given locus, but times are reduced if there are several
loci that can result in phenotypic change, as described in the Discussion. The paper is in two sections.
In the first we show that the time needed for a new neutral mating cue to evolve under allopatry is
very long. The time needed under parapatry is generally much shorter. In the second section we
calculate for selected cases the time needed under Type 2 parapatry, in which local adaptation
induces selection on mating cues even though the loci controlling the mating cues are not close to
the loci controlling local adaptation. This second section offers some insight into how the time
needed under Type 2 parapatry depends on the selection coefficients in the niches containing the
parapatric population.

Section 1: Time for a new mating cue to spread to fixation

In this section we compare times for a new mating cue to arise and spread to fixation under
allopatry and parapatry.

Methods

We assume mating cues are controlled by a single locus which is homozygous prior to a new mating
cue arising by mutation. The alleles at this locus are labelled D alleles. Any mutation of D can
produce a new mating cue. We label C the first such mutation that spreads to fixation, and assume C
dominant to D. In Type 1 parapatric speciation, the mating cue is an adaptive trait and so directly
subject to natural selection. In allopatry or Type 2 parapatry, mating cues are not subject to
selection directly, i.e. they are selectively neutral, but under Type 2 parapatry they may be acted on
by selection indirectly as a result of phenotype matching (Sibly and Curnow, 2022). Our aim in this
section is to use standard methods of population genetics to calculate the times for new mating cues
to arise and spread to fixation as a result of the mating cue selection coefficient acting directly or
indirectly on C. This is assumed to be zero under allopatry, but positive under parapatry.

The time of arising of the first mutated allele C that goes to fixation, here labelled T_a, and the time
from this mutation arising to fixation, T_f, are random variables, so the expected time to the fixation
of C, E[T], can be calculated from:

$E[T] = E[T_a] + E[T_f].$ (1)

The time of arising of the first mutated allele that goes to fixation depends on the mutation rate. We
let μ be the rate at which new neutral mutations arise per allele per generation. We begin by
calculating the expected time needed to acquire a new C mating cue under allopatry.

Timings under allopatry

Under allopatry we assume that mating cues are selectively neutral. To calculate the expected time
of arising of the first C mutation that goes to fixation, $E[T_a]$, we begin by calculating the probability of
a neutral mutation arising and going to fixation. This is the product of the probability of a mutation
arising, μ, and the probability it goes to fixation. Assuming large N and small μ, the per generation
probability of a mutation arising is approximately $2N\mu$, where N is the number of individuals in the
population. This is because there are 2N alleles at the focal locus and the chance of each mutating is
μ. The probability the arising mutation goes to fixation is 1/(2N) if the mutation is neutral ((Kimura
and Ohta, 1971), see also (Otto and Whitlock, 2013)). So the probability of a neutral mutation arising

and going to fixation is $2N\mu/(2N) = \mu$, and the expected time of arising of mutations that go to
fixation is the reciprocal of this, $1/\mu$.

Based on diffusion approximations with large N and continuous time, the expected time from the
arising of a neutral mutation until it reaches fixation is approximately 4N generations, conditional on
the allele fixing ((Kimura and Ohta, 1971), see also (Otto and Whitlock, 2013)). The expected time to
the fixation of C is now given by eqn (1) as:

Expected time to the fixation of C, E[T] $\approx \dfrac{1}{\mu} + 4N$ (2)

Timings under parapatry

We consider here two types of parapatric speciation, both of which assume the existence of
phenotype matching. In Type 1 parapatric speciation, adaptive traits are used as mating cues, i.e.,
are 'magic traits'. In this case mate choice may facilitate the spread of the adaptive gene(s) and
completion of local adaptation and speciation coincide. In Type 2 parapatric speciation, local
adaptation induces selection on a mating cue that is not linked to the adaptive trait(s). The strength
of selection on the mating cue then depends on how selection operates on the adaptive trait. The
overall selection coefficient on the mating cue depends on the selection coefficients operating on
the adaptive trait – positive in one niche and negative in the other. The relationship between these
selection coefficients is described in detail for Type 2 parapatry in Section 2, below. In considering
the timing of parapatric speciation, what is important is the overall selection pressure on the trait
under consideration, which could be either the adaptive trait or a mating cue distinct from the
adaptive trait. The overall selection pressure on the trait can be quantified by an *overall selection*
coefficient, here designated s. In this section s is taken to be constant. This is approximately true in
Type 2 parapatry except for the first few generations of the spread of the mating cue, when it is
higher (example in Fig. 2b, below).

We now use equation 1 to calculate the expected time to the fixation of C, E[T]. The probability of a
mutation arising is approximately $2N\mu$, as before. The probability the arising mutation goes to
fixation is approximately 2s if s is small ((Kimura and Ohta, 1971), see also (Otto and Whitlock,
2013)). So the probability of a selected mutation arising and going to fixation is $4N\mu s$, and the
expected time of arising of selected mutations that go to fixation, T_a, is the reciprocal of this,
$1/(4N\mu s)$.

The time from C arising to fixation can be obtained from equations given by (Charlesworth, 2020).
These equations apply to cases where the fixation index, F, is constant and known. Under parapatry,
however, as C spreads, mating is increasingly non-random and F increases. This situation has not
been modelled, and we therefore consider two limiting cases: F = 0 (random mating), and F positive
and constant. It turns out that the time from C arising to fixation is longer for random than for non-
random (F > 0) mating, so random-mating times represent a limiting case – actual times will be
shorter than those for random mating.

With random mating, for a dominant allele at an autosomal locus with weak selection and a large
population, (Charlesworth, 2020) showed that the time from C arising to fixation in units of 2N ,
conditional on the allele fixing, is approximately:

$\qquad (2\gamma)^{-1} + \gamma^{-1}[p_2{}^{-1} + \ln(p_1 q_2 q_1{}^{-1} p_2{}^{-1}) - p_1{}^{-1}] + 1.7724\gamma^{-0.5}$ (3)

where $\gamma = 2Ns$, assumed >>1, $p_1 = 1 - (2\gamma)^{-1}$, $p_2 = 0.8862\gamma^{-0.5}$, q_1=1-p_1, q_2=1-p_2, and N_e is taken
to be N.

With assortative mating and fixation index F, (Charlesworth, 2020) showed that the time is
approximately:

$$(2\gamma)^{-1} + \gamma^{-1}[-\ln(q_1 F) + F^{-1}\ln(Fp_2{}^{-1})] + 0.6321(F\gamma)^{-1} \qquad (4)$$

In deriving eqn 4 it is assumed that $F > O(s)$, and $p_2{}^{-1}$ is large. Numerical evaluations of eqns 3 and
4 reported in Supplementary Material show that times from C arising to fixation are always lower
without random mating than with it; the ratio declines as F or Ns increase, getting below 5% when
Ns>100,000 and F>0.1. The expected time to the fixation of a selected mutation for the limiting case
of random mating can now be approximated from equation 1 as:

E[T]= E[T_a]+ E[T_f] $\approx \dfrac{1}{4N\mu s} + 2N[(2\gamma)^{-1} + \gamma^{-1}[p_2{}^{-1} + \ln(p_1 q_2 q_1{}^{-1} p_2{}^{-1}) - p_1{}^{-1}] + 1.7724\gamma^{-0.5}]$

$\qquad (5)$

Equations 3 – 5 hold provided that s < 0.02 and Ns is large (Charlesworth, 2020).

Results

The expected times to fixation of a mating cue for allopatry and parapatry are shown in relation to
the selection coefficient *s* in Fig. 1, for the limiting case of random mating. Both T and T_a are plotted
but in Fig. 1a these overlay if N < 100,000 indicating that time from C arising to fixation is then
negligible in comparison with the time of arising of the first mutated allele that goes to fixation, T_a.
Fig. 1a shows that mating cues generally evolve substantially faster under parapatry than under
allopatry. With μ=10^{-8}, *s* = 0.02 and a population size of 10,000, cues evolve in 100,000 generations
under parapatry, compared with 100 million generations needed under allopatry. In a population of
100,000, cues evolve in a few thousand generations under parapatry. Even for a population of 100
individuals, cues evolve faster under parapatry than allopatry provided *s* > 0.03.

Fig. 1a shows the situation when the mutation rate μ=10^{-8}. If the mutation rate is higher, time to
fixation is reduced, and the contribution of the time from C arising to fixation is more pronounced.
These effects are illustrated in Fig. 1b, where μ=10^{-5}. The colour coding is the same in Figs. 1a and b.
All times are reduced in Fig. 1b in comparison with Fig. 1a. Times to fixation under parapatry are
closer together than in Fig. 1a because of increases in the contribution of the time from C arising to
fixation: these increase with N. Over the ranges of s and N shown in Fig. 1b, times to fixation range
from under a thousand if s=0.2 and N=1,000, to a little over 100,000 if s= 0.001 and N=1,000,000.
These are much less than the 5 million generations needed under allopatry.

Fig. 1. Expected times to fixation of a mating cue under allopatry and parapatry for the limiting case
of random mating. Times to fixation are plotted against the overall selection coefficient *s* on log-log
scales for two mutation rates: a) μ=10^{-8}; b) μ=10^{-5}. The lines for allopatry are shown for comparison

with those for parapatry but the selection coefficient under allopatry is assumed to be zero. The
allopatry lines satisfy equation 2. The other lines are for parapatric populations, for which the
numbers in the right-hand columns indicate population size in equation 5. The lines for both E[T] and
E[T_a] are plotted and these generally overlay. Where they do not overlay the E[T_a] line has the suffix
T_a, the E[T] line has no suffix. The cases in which the lines overlay are in a) where N < 100,000 and
allopatry; and in b) where N < 1,000. Dashed lines indicate regions where Ns <5, i.e., failure of the
assumption Ns large that is used to derive eqns 3 – 5.

The results shown in Fig. 1 show the effects of selection on the time needed for a trait to evolve. The
case labelled 'Allopatry' corresponds to the trait being neutral. This is the limiting case as the
selection coefficient *s* tends to zero. The cases when *s* > 0 are considered as cases of parapatry.
Although these results apply to any evolving traits, our focus here is on discussing them in relation to
allopatry and parapatry. Fig. 1 suggests that the time needed for a new mating cue to evolve can be
much shorter under parapatry than under allopatry. If the mutation rate μ=10⁻⁸, the value commonly
reported in reptiles, birds, fish and mammals (Bergeron et al., 2023), then in populations under
100,000 the time needed for a new mating cue to evolve depends almost entirely on the time of
arising of the first mutated allele C that goes to fixation, as shown in Fig. 1a. The times needed for a
new mating cue to evolve under allopatry and parapatry then simplify to $\frac{1}{\mu}$ and $\frac{1}{4N\mu s}$ respectively.
Thus, the time needed under parapatry is lower than that needed under allopatry by a factor of $4Ns$.
On the log-log scales of Fig. 1, the relationship for allopatry is $\log(time\ needed) = -\log \mu$. The
relationship for parapatry is $\log(time\ needed) = -\log \mu - \log 4 - \log N - \log s$. So, *time needed*
can be decreased by an order of magnitude by increasing population size *N* by an order of
magnitude, or increasing the overall selection coefficient *s* by an order of magnitude.

If the mutation rate is higher, time to fixation is reduced, and the contribution of the time from C
arising to fixation is more pronounced (example in Fig. 1b). The net effect is that the time needed for
a new mating cue to evolve is still much shorter under parapatry than under allopatry. Allowing for
changes in the y-axis scale, results are similar to those shown in Fig. 1 as mutation rate is varied from
μ=10⁻⁸ to μ=10⁻² (results not shown).

In this section we have shown that the time needed for a new mating cue to evolve can be much
shorter under parapatry than under allopatry, as illustrated in Fig. 1. This is true of all three types of
parapatric speciation.

Section 2: Dependence of the mating cue selection coefficient on the selective regimes in two
niches in (Sibly and Curnow, 2022)'s Type 2 parapatric populations

One objective of this paper is to calculate the time needed for a new mating cue to evolve when local
adaptation induces selection on mating cues even though the loci controlling the mating cues are not
linked to the loci controlling local adaptation – i.e., Type 2 parapatry. The mating cue is assumed to
be coded by a C allele arising by mutation at a locus not linked to those producing local adaptation.
Fig. 1 illustrates how the advantages of parapatry depend on the value of the overall selection
coefficient acting on the C allele. In this section we consider how the overall selection coefficient on
the mating cue relates to the selection coefficients operating on the adaptive trait – positive in one
niche and negative in at least one other. An indication of the answer to this question can be obtained
using (Sibly and Curnow, 2022)'s two-niche model of Type 2 parapatric speciation.

Methods

In (Sibly and Curnow, 2022)'s model generations are discrete and individuals die after mating. At the
start of each generation individuals in each niche mate with others of the same mating-cue
phenotype, and all mating individuals obtain the same number of offspring. The number of offspring
of each genotype that survive in each niche is the product of its initial frequency and its fitness.
Population regulation then returns population numbers to their initial values, after which some
individuals migrate between niches. Local adaptation is the result of two alleles P and Q at a single
locus. PP has fitness 1 in both niches. QQ has fitness $1+s_1$ in niche 1 and $1+s_2$ in niche 2. s_2 is assumed
positive and s_1 negative. Whether or not local adaptation occurs depends not only on the values of s_1
and s_2 and the level of dominance of Q, but also on the migration rates between the two niches. For
the case that population sizes in the two niches are equal and migration rates in both directions are
the same, (Sibly and Curnow, 2022) used deterministic simulation of the evolutionary process to map
the set of migration and selection rates for which local adaptation occurs, and showed that in all
analysed cases, the evolutionary outcome is speciation if a C allele coding for a new neutral mating
cue arises by mutation. C allele was originally introduced into the simulation at low frequency
(0.05%). As the C allele spread the CD and PQ heterozygotes were eliminated, so that eventually the
population consisted only of the CCQQ and DDPP genotypes. The C allele increased in frequency
approximately exponentially until close to its final value (example in Fig. 2). During the exponential
phase the mating cue selection coefficient on C alleles was obtained by simulations carried out as in
(Sibly and Curnow, 2022). By repeating this procedure over a range of values of s_1, s_2 and migration
rate m, a picture was built up of the dependence of s on s_1, s_2 and m.

Fig. 2. Example of the exponential increase of the C allele in the (Sibly and Curnow, 2022) model. a)
The frequency of the CCQQ and CDQQ genotypes in the two niches. The C allele was originally
introduced at low frequency (1%) into niche 2 as a CDQQ genotype. The evolutionary outcome is
that only two genotypes persist in the two niches, CCQQ and DDPP (DDPP is not shown here). b) The
resulting increase of the C allele is approximately linear on a log scale, indicating that the overall
selection coefficient s is constant except for the first few generations, when it is higher. Parameter
values were m=0.067, s_2 = 2.0, s_1 = -0.4. Simulation carried out as in (Sibly and Curnow, 2022).

Results

Fig. 3 shows how the mating cue selection coefficient on the C allele, *s*, is affected by migration rate
m and the selection coefficient s_2 in niche 2, for two values of s_1. s_2 and s_1 represent the strength of
selection for and against the Q allele in niches 2 and 1 respectively. Note first that selection for C, i.e.
s > 0, only occurs within a restricted region in the m s2 plane. This is the region in which local
adaptation is possible. Outside this region C is not selected or is selected against, so s ≤ 0. The
highest values of the selection coefficient are 0.013 in Fig. 3a and 0.12 in Fig. 3b. They occur on

horizontal ridges where the opposing selection conditions in the two niches are close to being equal
and opposite. Thus, the ridge occurs at $s_2 \sim 0.05$ in Fig. 3a and at $s_2 \sim 0.40$ in Fig. 3b. High values of s
also occur on vertical ridges at $m \sim 0.05$ in Fig. 3a and $m \sim 0.15$ in Fig. 3b. Either side of the ridges, s
declines down to zero at the edges of the region in which local adaptation is possible. To the left s
declines to zero when $m = 0$. $m = 0$ corresponds to allopatry since there is no migration between
niches. Towards the bottom of Fig. 3, s declines steeply and becomes negative as s_2 declines to zero.
When s_2 is zero, s is negative because there is no selection for CCQQ in niche 2 and selection against
CCQQ in niche 1. s is zero at the upper right of the figure, because local adaptation is not possible
there: the region consists entirely of Q alleles, and there is then no selection for C alleles.

In sum, as in (Sibly and Curnow, 2022) there is no selection for the mating cue outside the region is
which local adaptation occurs. Within the region of local adaptation, selection on the mating cue
seemingly increases with migration rate from zero up to some optimum, and then declines back
down to zero at the edge of the adaptive region. Mating cue selection is also relatively high if the
opposing selection conditions in the two niches are approximately equal and opposite, i.e., $s_2 \sim - s_1$.

The cases analysed assume complete phenotype matching, populations divided equally between two
niches, migration rates the same in both directions, and local adaptation the result of two alleles P
and Q at a single locus. The mating cue selection coefficient s is at most 0.12 in the cases analysed in
Fig. 3, within the range of validity of the assumptions used to derive eqns 3 - 5. Complete phenotypic
matching is assumed in Fig. 3, but if some individuals mate with others of a different phenotype then
the mating cue selection coefficient would be reduced. The evolution of choosiness could be
explored using the methods and results of (Aubier et al., 2023) as a starting point, and theoretical
equations with which to investigate this and other parameter variants are available in (Sibly and
Curnow, 2022), though further work is needed to see what happens when key genes are not
dominant.

Fig. 3. 3D contour plots showing how the mating cue selection coefficient, *s*, is affected by migration
rate *m* and s_2, for two values of s_1. Mating cue selection coefficients *s* were calculated over grids of
points in (Sibly and Curnow, 2022)'s Fig. 2 as follows: (a) corresponds to (Sibly and Curnow, 2022)'s
bottom left panel; (b) to their bottom right panel. In these simulations Q was dominant to P.
Simulations carried out as in (Sibly and Curnow, 2022). *s* was calculated in the phase of exponential
increase of C, generation 50 in (a) and 20 in (b). The contour plots were obtained by interpolation
between grid points using Minitab 21.

Discussion

Our main results, illustrated in Fig. 1, show quantitatively how the time needed for a mating cue to evolve depends on the strength of selection for the mating cue. These results hold for any newly evolving allele but our focus in this paper is on mating cues. The results demonstrate quantitatively how much shorter is the time needed for a new mating cue to evolve under parapatry compared with under allopatry. The time needed can be reduced by several orders of magnitude, depending on population sizes and the strength of selection, as shown in Fig. 1. How the strength of selection for the mating cue depends on the strength of selection for an ecological trait in (Sibly and Curnow, 2022)'s Type 2 parapatry is addressed in Fig. 3. To see the relationship between Figs. 1 and 3, consider a fictitious example: suppose an F_{ST} value of 0.31 is recorded for an ecological trait that varies between two niches. Using the results of (Sibly and Curnow, 2023), the selection coefficients on the ecological trait would be 0.4 in one niche and -0.4 in the other if their absolute values were equal. Fig. 3 shows that the mating cue selection coefficient would then be around 0.1 if the migration rate was 0.15, in which case from Fig. 1 the expected speciation time would be about 26,000 generations if the population size was 100,000. For smaller populations or lower migration rates the expected speciation time would be longer. The take-home message is that from measured F_{ST} values for an ecological trait, if we have some indication of migration rates we can get an idea of expected speciation times.

These results assume that phenotypic change can only occur as a result of genetic change at a single locus. If there are several loci that potentially could code for mating cues, the number being given the symbol x, then μ should be replaced by μx throughout the analysis. This is because there are now $2Nx$ alleles at the x focal loci and the chance of each mutating is μ, so the per generation probability of a mutation arising is approximately $2N\mu x$. The rest of the analyses follow as before with μ replaced by μx. The effect is identical to that of increasing μ, so that for example in comparison with Fig. 1a, Fig. 1b could result either from increasing μ by three orders of magnitude or from an equivalent increase in x. So speciation times are reduced the more loci there are that potentially could code for mating cues.

The results in Fig. 1 hold for all types of parapatric speciation. If the mating cue evolves by Type 1 parapatric speciation, the mating cue is the same as the adaptive trait – a prime example is beak size. Evolution of species-specific songs or plumage coloration – if these are not themselves adaptive – would then have to await a relevant neutral mutation arising and going to fixation. The timings would be long, the same as under allopatry. By contrast to the above, species-specific songs or plumage colorations evolve faster under Type 2 parapatric speciation on the assumption that local adaptation has already occurred. The case of (Servedio and Burger, 2020)'s pseudomagic traits would be intermediate between Type 1 and Type 2. The mating cue trait would then need to evolve close to the ecological trait in locations comprising some proportion y, say, of the genome. Using the argument of the preceding paragraph, the chance of a mutation arising in these locations would be a fraction y of the chance of a mutation arising anywhere in the genome, so μx would be replaced by $\mu x y$ throughout the analyses, and relative to Type 2, speciation times would increase by a factor of $\frac{1}{y}$. If for example y was 1/1000 of the genome then speciation times would increase by 1000. In sum, species-specific songs or plumage colorations evolve fastest under Type 2 parapatric speciation on the assumption that local adaptation has already occurred.

The assumptions and approximations used in deriving equations 2 – 5 merit discussion. It has been assumed that mating cues under allopatry are selectively neutral, but this might not be true if sexual selection operated on the mating cues during allopatry. However to get differences between the geographic regions it would be necessary for sexual selection to operate differently in the different regions. Sexual selection depends on mate-choosers – often females – have preferences for selected

traits, so these preferences also would need to differ between the geographic regions. So while
possible in some cases this seems unlikely as a general explanation, and we contend that our
assumption of neutrality of mating cues under allopatry is an appropriate theoretical starting point. It
is also assumed that assortative mating has no costs, but being choosy may reduce the chance of
mating, reducing the mating cue selection coefficient in the parapatric models (see, e.g., (Kopp and
Hermisson, 2008; Schneider and Burger, 2006)). The probability of a mutation arising is $1 -$
$(1 - \mu)^{2N}$ and this is approximated as $2N\mu$. This is a good approximation even when N=1,000,000:
with μ =10^{-8} the approximation then gives 0.0200 compared to the exact value of 0.0198. In deriving
equations $3 - 5$, s is taken to be constant, though in reality it is higher in (Sibly and Curnow, 2022)'s
Type 2 parapatry in the first few generations of the spread of the mating cue, as shown in Fig. 2b. The
effect of this will be to shorten the time to fixation of a mating cue under Type 2 parapatry. Fig. 1
only shows s values up to 0.2 and Ns large, the regions in which (Charlesworth, 2020)'s eqns $3 - 5$ are
valid. In sum, since the values of *s* may often be less than 0.1 (Fig. 3), our conclusions are unlikely to
be affected by the approximations used in deriving equations $2 - 5$.

We are now in a position to compare with data the expected times to fixation of a mating cue under
allopatry and parapatry shown in Fig. 1. As described in the Introduction, analysis of molecular
phylogenies suggest that eight species of southern capuchinos seedeaters (*Sporophila*) evolved
distinct male plumage coloration and in some cases song in less than 50,000 generations, and in
Darwin's finches radiations of ground and tree finches began around 100,000–300,000 years ago and
at least in some cases involved species-specific songs (Grant and Grant, 2018). These timings of
around 100,000 generations are three orders of magnitude less that the time of 10^8 generations that
Fig. 1 suggests would be required under allopatry or Type 1 parapatry, but are similar to those
resulting from Type 2 parapatry. Although there are several simplifications in our argument, it seems
likely that only Type 2 parapatric speciation can account for the times recorded for the evolution of
species-specific songs or plumage coloration.

Our results encourage further research on species that are in the process of speciating or have very
recently speciated. We expect to find that the species uses sexual imprinting (or some other method
of phenotypic matching) as has been shown to exist in Darwin's finches (Grant and Grant, 2018).
Comparing the genomes of very recently diverged species, under Type 2 parapatry we expect to find
divergence at just one plumage/song locus, since the evolution of more loci, while they may clarify
the signal, are likely to add substantially to the time taken. In addition to the divergence at the
plumage/song locus we expect to find divergence at one or more loci conferring local adaptation(s),
which we expect to have diverged earlier, on the premise that, under Type 2 parapatry, local
adaptation is in general a necessary precursor to the evolution of a new mating cue.

Acknowledgements. We are grateful to Brian Charlesworth for comments on the manuscript.

Conflict of Interest: None.

Supplementary Material: Numerical evaluation of times to fixation given by equations 3 and 4.

Aubier, T. G., Buerger, R., Servedio, M. R., 2023. The effectiveness of pseudomagic traits in promoting
premating isolation. Proceedings of the Royal Society B-Biological Sciences 290,
doi:10.1098/rspb.2022.2108.
Bergeron, L. A., Besenbacher, S., Zheng, J., Li, P., Bertelsen, M. F., Quintard, B., Hoffman, J. I., Li, Z., St.
Leger, J., Shao, C., Stiller, J., Gilbert, M. T. P., Schierup, M. H., Zhang, G., 2023. Evolution of the
germline mutation rate across vertebrates. Nature 615, 285-291, doi:10.1038/s41586-023-
05752-y.
Butlin, R. K., Servedio, M. R., Smadja, C. M., Bank, C., Barton, N. H., Flaxman, S. M., Giraud, T.,
Hopkins, R., Larson, E. L., Maan, M. E., Meier, J., Merrill, R., Noor, M. A. F., Ortiz-Barrientos,

D., Qvarnstrom, A., 2021. Homage to Felsenstein 1981, or why are there so few/many
species? Evolution 75, 978-988, doi:10.1111/evo.14235.
Campagna, L., Repenning, M., Silveira, L. F., Fontana, C. S., Tubaro, P. L., Lovette, I. J., 2017. Repeated
divergent selection on pigmentation genes in a rapid finch radiation. Science Advances 3,
doi:10.1126/sciadv.1602404.
Campagna, L., Benites, P., Lougheed, S. C., Lijtmaer, D. A., Di Giacomo, A. S., Eaton, M. D., Tubaro, P.
L., 2012. Rapid phenotypic evolution during incipient speciation in a continental avian
radiation. Proceedings of the Royal Society B-Biological Sciences 279, 1847-1856,
doi:10.1098/rspb.2011.2170.
Charlesworth, B., 2020. How Long Does It Take to Fix a Favorable Mutation, and Why Should We
Care? American Naturalist 195, 753-771, doi:10.1086/708187.
Felsenstein, J., 1981. Skepticism towards Santa Rosalia, or why are there so few kinds of animals.
Evolution 35, 124-138, doi:10.2307/2407946.
Grant, P. R., Grant, B. R., 2018. Role of sexual imprinting in assortative mating and premating isolation
in Darwin's finches. Proceedings of the National Academy of Sciences of the United States of
America 115, E10879-E10887, doi:10.1073/pnas.1813662115.
Hejase, H. A., Salman-Minkov, A., Campagna, L., Hubisz, M. J., Lovette, I. J., Gronau, I., Siepel, A.,
2020. Genomic islands of differentiation in a rapid avian radiation have been driven by recent
selective sweeps. Proceedings of the National Academy of Sciences of the United States of
America 117, 30554-30565, doi:10.1073/pnas.2015987117.
Janicke, T., Marie-Orleach, L., Aubier, T. G., Perrier, C., Morrow, E. H., 2019. Assortative Mating in
Animals and Its Role for Speciation. American Naturalist 194, 865-875, doi:10.1086/705825.
Kimura, M., Ohta, T., 1971. Theoretical Aspects of Population Genetics. Princeton University Press,
Princeton, N.J.
Kopp, M., Hermisson, J., 2008. Competitive speciation and costs of choosiness. Journal of
Evolutionary Biology 21, 1005-1023, doi:10.1111/j.1420-9101.2008.01547.x.
Kopp, M., Servedio, M. R., Mendelson, T. C., Safran, R. J., Rodriguez, R. L., Hauber, M. E., Scordato, E.
C., Symes, L. B., Balakrishnan, C. N., Zonana, D. M., van Doorn, G. S., 2018. Mechanisms of
Assortative Mating in Speciation with Gene Flow: Connecting Theory and Empirical Research.
American Naturalist 191, 1-20, doi:10.1086/694889.
Lamichhaney, S., Berglund, J., Almen, M. S., Maqbool, K., Grabherr, M., Martinez-Barrio, A.,
Promerova, M., Rubin, C. J., Wang, C., Zamani, N., Grant, B. R., Grant, P. R., Webster, M. T.,
Andersson, L., 2015. Evolution of Darwin's finches and their beaks revealed by genome
sequencing. Nature 518, 371-375, doi:10.1038/nature14181.
Otto, S. P., Whitlock, M. C., 2013. Fixation Probabilities and Times. eLS,
doi:10.1002/9780470015902.a0005464.pub3.
Price-Waldman, R., Stoddard, M. C., 2021. Avian Coloration Genetics: Recent Advances and Emerging
Questions. Journal of Heredity 112, 395-416, doi:10.1093/jhered/esab015.
Schneider, K. A., Burger, R., 2006. Does competitive divergence occur if assortative mating is costly?
Journal of Evolutionary Biology 19, 570-588, doi:10.1111/j.1420-9101.2005.01001.x.
Servedio, M. R., Burger, R., 2020. The effectiveness of pseudomagic traits in promoting divergence
and enhancing local adaptation. Evolution 74, 2438-2450, doi:10.1111/evo.14056.
Sibly, R. M., Curnow, R. N., 2022. Sexual imprinting leads to speciation in locally adapted populations.
Ecology and Evolution 12,
, doi:h t t p s : //d o i .org/10.1002/ece3.9479.
Sibly, R. M., Curnow, R. N., 2023. Allele frequencies and selection coefficients in locally adapted
populations. Journal of Theoretical Biology 565, doi:10.1016/j.jtbi.2023.111463.
Smadja, C. M., Butlin, R. K., 2011. A framework for comparing processes of speciation in the presence
of gene flow. Molecular Ecology 20, 5123-5140, doi:10.1111/j.1365-294X.2011.05350.x.
ten Cate, C., Vos, D. R., 1999. Sexual imprinting and evolutionary processes in birds: A reassessment.
In: Slater, P. J. B., et al., Eds.), Advances in the Study of Behavior, Vol. 28, Vol. 28, pp. 1-31.

Tobias, J. A., Ottenburghs, J., Pigot, A. L., 2020. Avian diversity: speciation, macroevolution, and
ecological function. Annual Review of Ecology, Evolution and Systematics 51, 533-560.
Turbek, S. P., Semenov, G. A., Enbody, E. D., Campagna, L., Taylor, S. A., 2021. Variable Signatures of
Selection Despite Conserved Recombination Landscapes Early in Speciation. Journal of
Heredity 112, 485-496, doi:10.1093/jhered/esab054.
Verzijden, M. N., ten Cate, C., Servedio, M. R., Kozak, G. M., Boughman, J. W., Svensson, E. I., 2012.
The impact of learning on sexual selection and speciation. Trends in Ecology & Evolution 27,
511-519, doi:10.1016/j.tree.2012.05.007.
474