1

Bacteria in honeybee crops are decoupled from those in floral nectar and bee

mouths

Magdalena L. Warren[1]*, Kaoru Tsuji[1,2,3]*, Leslie E. Decker[1], Manabu Kishi[4,5], Jihoon

Yang[6], Adina C. Howe[6], Tadashi Fukami[1,7]

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA

[2]Center for Ecological Research, Kyoto University, 2-Hirano, Otsu, Shiga, 520-2113,

Japan

[3]Department of Biology, Graduate School of Science, Kobe University, Kobe, 657-8501,

Japan

[4]Japanese Apricot Laboratory, Wakayama Fruit Tree Experiment Station, Minabe,

Higashi-Honjo 1416-7, Wakayama, 645-0021, Japan

[5]Hidaka Promotion Bureau, Wakayama Prefecture, Gobo, Yukawa 651, Wakayama,

644-0011, Japan

[6]Department of Agricultural and Biosystems Engineering, Iowa State University, Ames,

Iowa, USA

[7]Department of Earth System Science, Stanford University, Stanford, CA 94305, USA

* These authors contributed equally to this work

Competing Interests statement: This work was supported by the Office of International

Affairs and the FSI Japan Fund of Stanford University.

Abstract

Bacteria in the honeybee gut have garnered much attention as a factor affecting bee

health. However, unlike the hindgut, which has been the primary focus of this research,

the crop, or the honey stomach, is often assumed to be dominated by environmentally

acquired transient taxa that matter little to the bees. To evaluate this assumption, we

examined bacterial taxa in the crop and the mouth of foraging adults of *Apis mellifera*

and *A. cerana japonica* and the floral nectar of the Japanese apricot, *Prunus mume*,

visited by the bees in the Minabe-Tanabe region of Japan. We found that the crop was

distinct from both the mouth and the nectar in bacterial composition even though all

samples were collected simultaneously at the same locations, whereas mouth and

nectar samples were indistinguishable from each other. Furthermore, the crop remained

similar in bacterial composition and alpha and beta diversity from summer to winter,

whereas the mouth showed a sharp drop in alpha diversity and a large increase in beta

diversity from summer to winter. These results refute the conventional assumption,

suggesting instead that the crop contains a conserved bacterial community that is

largely distinct from the environmental taxa that the bees encounter as they forage in

the environment. We also found, however, that strains of a representative crop-

associated species, *Apilactobacillus kunkeei*, could be season- and host species-

specific. Together, these findings suggest that further studies of crop-associated

bacterial communities are needed to better understand the relationship between

honeybees and their gut bacteria.

Introduction

Microbes in the honeybee gut are receiving increasing attention not only as a factor affecting the health of the agriculturally vital insects, but also as a model system for gaining basic understanding of gut-associated microbiota (1-3). One component of the honeybee gut, the crop, contains species such as *Apilactobacillus kunkeei* and *Bombella apis* that may affect the survival and pathogen resistance of the foraging adults that host them and the larvae that the adults feed (2, 4-6). However, microbes in the crop are not as well studied as those in the hindgut, which are generally considered more important to honeybee health (7-10).

Given its function for temporary storage and transport of floral nectar (11), it seems reasonable to assume that the crop is occupied mainly by the transient microbes acquired from the environment via foraging of nectar (12). One way to investigate this assumption of environment–crop matching is to compare microbes in the crop of foraging bees and those in their mouth as well as in the floral nectar they forage for. If the assumption is true, the crop should be similar to both the mouth and the floral nectar in bacterial composition (4, 13). To our knowledge, no study has directly compared bacteria in crop, mouth, and nectar for this purpose, leaving untested the assumption that the crop microbiota is dominated by environmentally derived transients.

Here we test the hypothesis that crop-associated bacteria show species composition that mirror those observed in mouth- and nectar-associated bacteria. To this end, we compare crop-, mouth-, and nectar-associated bacteria in samples that were collected

simultaneously at the same locations. Our crop and mouth samples come from foraging

adults of the introduced *Apis mellifera* and the native *A. cerana japonica* (hereafter *A.*

cerana) collected in the Minabe-Tanabe region of Wakayama Prefecture in Japan. In

this region, farmers use both species of honeybees to pollinate the winter-blooming

Japanese apricot, *Prunus mume* (14). Our nectar samples come from *P. mume* flowers

near the hives of the *A. mellifera* and *A. cerana* bees that we caught for crop and mouth

sampling.

Methods

Study sites

Japanese apricot orchards in the Minabe-Tanabe region are embedded within a

mountainous countryside landscape that has recently become internationally

recognized. In 2015, the FAO designated the region as a Globally Important Agricultural

Heritage System for the agricultural practice in the region that embraces both human

needs and biodiversity conservation. Many of the apricot orchards are located adjacent

to natural and managed forests, which are thought to provide stable habitats for the

native *Apis cerana*. The two *Apis* species are the main pollinators of the apricot, which

blooms from mid or late February to early or mid-March, during which few other flower-

visiting insects are available (14). All our bee and nectar collection sites were located

within this landscape, but at least 1.0 km away from their respective nearest

neighboring collection site in each season (Fig. 1, Table S1).

Bee collection

A total of 12 *P. mume* orchards in the study region were visited in the summer of 2018

and the winter of 2019. Seventy-two and 149 honeybees were collected in the summer

of 2018 and the winter of 2019, respectively, from colonies in or near these apricot

orchards (Table S1). Honeybees were collected as they flew near their hives located

within or near the orchards. At orchards where beekeepers maintained *A. ceran*a hives,

A. cerana bees were collected near their hives whose locations varied from tens of

meters to about a hundred meters from the orchard and were often situated at the edge

of the local forest. The collected bees were individually placed in a sterile, sealed plastic

vial immediately after the collection. The vials were kept for up to two hours in a cool

box with ice packs in it for transport to the Japanese Apricot Laboratory located within

100 m from one of the collection sites (Higashi-Honjo) (Fig. 1).

After we brought the collected bees into the laboratory in the Japanese Apricot

Laboratory facility, the bees were kept cool in the box for up to two additional hours so

that the bees would stay sluggish for ease of handling. The captured bees were then

taken out of the vials, and their wings were fixated with adhesive tape to facilitate

handling. The taped bees were made to drink approximately half of 20 µl of sterile 20%

sucrose solution placed on the lid of a sterile 1.5-ml microcentrifuge tube. After the

feeding, the remaining solution was placed in the microcentrifuge tube with the lid now

closed. These solutions were later used to study the bacteria associated with the mouth.

We refer to these solutions as mouth samples. Since we had the bees extend their

proboscis into the sterile sucrose solution, our method is likely to have captured mostly

the bacteria from the surface of the proboscis rather than those present within the

mandibles.

Immediately after the sampling of mouth-associated bacteria, we dissected the bees by

gently pulling the abdomen using tweezers to expose the crop, making sure the crop did

not touch other parts of the bee body. We then removed the crop from the bee body

using another set of tweezers. Both tweezers were sterilized before each use by dipping

in 70% ethanol and then flaming. Each crop sample was separately placed in 40 µl of

sterile water within sterile microcentrifuge tubes and homogenized using a sterile plastic

pestle. All mouth and crop samples were immediately stored at -80°C until further

processing.

We collected crop and mouth samples not just in winter during the *P. mume* flowering

season, but also in summer, to determine whether crop and mouth differed in the way

bacterial composition changed from summer to winter. Hives of *A. cerana*, many of

which were made of carved tree trunks, were maintained by local beekeepers

throughout the year in the region. On the other hand, most *A. mellifera* hives in the

region were transported elsewhere in the country while *P. mume* was not in bloom.

However, some were kept locally throughout the year, allowing us to collect bees in both

seasons. One site (Nishi-Honjo) was available for bee collection in both summer and

winter, allowing bacterial taxa from the same site and hives to be compared between

seasons. In addition, at two of our sites (Higashi-Honjo and Kamihaya), we were able to

collect both *A. cerana* and *A. mellifera*, enabling comparison of the two species

collected at the same locations (Table S1).

Nectar collection

In addition to the mouth and crop samples, we also collected samples of *P. mume* floral

nectar on the same days the bees were collected in the winter of 2019. At each of the

nine orchards where we sampled bees that winter (Table S1), we collected 16 nectar

samples (except at one site, Kamihaya, where we instead collected 24 nectar samples),

for a total of 152 nectar samples. Each sample was collected by probing 10 flowers on a

randomly selected branch of an individual *P. mume* tree with a sterile 0.5μl

microcapillary tube and dispensing the collected nectar into a PCR tube containing 40 μl

of PCR-quality sterile deionized water. The collected nectar samples were immediately

placed in a box with ice packs during the transport to the laboratory. Within six hours,

the nectar samples were brought back to the Japanese Apricot Laboratory. The samples

were kept at -80°C until further processing for DNA extraction, except during a one-day

transport to the laboratory at Kyoto University in Shiga, Japan, and another one-day

transport from there to Stanford University in California, USA, during which the samples

were kept at about -5 °C.

Microbial DNA extraction

In our laboratory at Stanford University, the Qiagen DNeasy Blood and Tissue kit and

Qiagen Genomic DNA Buffer set (Qiagen, Germantown, MD, USA) were used to extract

DNA from all the samples. The extraction followed the gram-positive bacteria protocol

along with the following modifications. For each of the crop, mouth, and nectar samples

collected, 10 µL of the thawed sample was combined with 2 µL of zymolyase enzyme

(MP Biomedicals, Burlingame, CA, USA) within a microcentrifuge tube. These mixtures

were incubated at 30°C for 30 minutes. Once the incubation period was complete, 10 µL

of a 500:1 G2 (from the Qiagen Genomic DNA Buffer set) buffer to RNAse A (Qiagen)

mixture was added to each DNA and zymolyase mixture. 20 µL of ProteinaseK (Qiagen)

and 180 µL of Buffer ATL (Qiagen) were added to each sample and vortexed to mix. The

samples were then incubated at 56°C overnight. The next morning the samples were

mixed and centrifuged to collect any condensation on the caps, and 410 µL of AL-

Ethanol (Qiagen) mixture were added to each sample and shaken vigorously to mix.

The rest of the protocol was followed as stated in the DNeasy Blood & Tissue Handbook

(2018) with the only exception being that the final elution of DNA was done with 50 µL of

Buffer AE (Qiagen) rather than 200 µL to increase the final DNA concentration in the

eluate. The final extracted DNA from each sample was stored in sterile microcentrifuge

tubes at 4°C until library preparation.

Bacterial amplicon sequencing

The highly variable (V4) region of the 16S rRNA gene was amplified using primers 515F

(5´-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGTGYCAGCMGCCGCGGTAA-

3´) and 806R

(5´GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGGACTACNVGGGTWTCTAAT

-3´) (15). Both primers included an Illumina (San Diego, CA, USA) adapter on the 5´

ends used to attach unique Nextera XT indexes for sample identification. Each 10 µL

first step PCR reaction consisted of 3.2 μL of PCR-grade water, 5 μL of MyTaq HS Red

Mix (Bioline, Taunton, MA, USA), 0.4 μL each of the forward and reverse primers, and 1

μL of the extracted DNA sample. PCR conditions were 95ºC for 2 minutes, followed by

35 cycles of 95ºC for 20 seconds, 50ºC for 20 seconds, 72ºC for 50 seconds, and a final

extension at 72ºC for 10 minutes with storage at 4ºC. Amplification was confirmed by

running samples on an agarose gel, and the resulting amplicons were purified using

Sera-Mag SpeedBeads (Sigma-Aldrich, St. Louis, MO, USA). The second step 10 μL

PCR reaction consisted of 3.2 μL of PCR-grade water, 5 μL of MyTaq HS Red Mix

(Bioline), 0.4 μL each of the combined Nextera XT index primers 1 and 2, and 1 μL of

the first step PCR product. PCR conditions for this second step were 95ºC for 2

minutes, followed by 8 cycles of 95ºC for 20 seconds, 50ºC for 20 seconds, 72ºC for 50

seconds, and a final extension at 72ºC for 10 minutes with storage at 4ºC. Amplification

was confirmed by running samples on an agarose gel, the barcoded amplicons were

purified once again using Sera-mag SpeedBeads (Sigma-Aldrich), and the resulting

clean product was pooled evenly across samples. The pooled samples were combined

to a concentration of 4 nM using a Qubit 4 Fluorometer (Thermo Fisher, Waltham, MA,

USA), and then the entire amplicon libraries were pooled together. The final DNA

concentration and quality of the pooled amplicon libraries were quantified with a

Fragment Analyzer (Agilent, Santa Clara, CA, USA), and then sequenced on an Illumina

MiSeq using a 2 X 300 cycle sequencing kit with a 15% PhiX spike-in (16) at the

Stanford Genomic Sequencing Service Center.

Quantitative real-time PCR (qPCR) of 16S rRNA gene

A subset of the crop and mouth samples was selected for 16S rRNA gene

quantification. These 112 samples were made up of mouth and crop samples from 28

Apis mellifera and 28 *A. cerana* bees, 14 each from the summer and winter sites. To

ensure we captured differences in bacterial load in Nishi-Honjo, the one site where we

did the crop and mouth sampling in both seasons, we included all samples from this site

in this analysis. The remaining *A. cerana* and *A. mellifera* samples were randomly

selected from the other sites. The total number of bacterial 16S rRNA genes was

quantified in the extracted DNA samples. Targeted genes were amplified using the

CFX96^{TM} Real-Time System (C1000TM Thermal Cycler, Bio-Rad, Hercules, CA, USA).

All reactions were performed using SsoFast™ EvaGreen® Supermix (Bio-Rad)

according to the manufacturer's instructions and in triplicate using 10 µL of SsoFast

EvaGreen Supermix, 0.6 µL of each primer, 2 µL DNA samples, and nuclease-free

water to a final volume of 20 µL. 16S rRNA genes were amplified using the primer sets

1055YF(ATGGYTGTCGTCAGCT) and 1392R (ACGGGCGGTGTGTAC), with the PCR

conditions as follows: 2 minutes at 50°C and 10 minutes at 95°C, followed by 40 cycles

of 15 seconds at 95°C and 1 minute at 58°C (17). The detection limit (1.6 x 10^2 gene

copies per reaction) for quantitative real-time PCR was determined using 55 standard

curves for 16S rRNA genes that were constructed using the serial 10-fold dilutions from

10^{-1} to 10^{-7} of known concentrations of synthesized oligonucleotides (Integrated DNA

Technologies, Inc., Coralville, IA, USA). The gene copy number was determined from

CT values equal to or higher than the negative controls were considered as below the

detection limit.

16S rRNA gene amplicon-sequencing data processing

The raw amplicon sequences were trimmed using the Cutadapt software (18). The

Divisive Amplicon Denoising Algorithm 2 (DADA2) pipeline (19) was used to merge

paired-end sequences, quality filter these sequences, remove chimeric reads, and

cluster sequences into amplicon sequence variants (ASVs). Taxonomy assignments for

each ASV were assigned using the SILVA database (20). ASVs present in the DNA

extraction and first and second step PCR negative controls as well as those identified

as chloroplast or mitochondria were removed from the dataset. Any phyla less than 1%

abundant among all samples were trimmed (21). This preprocessing resulted in several

samples containing no sequence reads. Consequently, these samples were excluded

from further analysis, resulting in a total of 218 crop samples, 161 mouth samples, and

82 nectar samples used for downstream analysis. Note that the taxon previously called

Parasaccharibacter apium (5) has been re-named *Bombella apis* (22). ASVs were

clustered into operational taxonomic units (OTUs) with a 97% similarity threshold using

the DECIPHER and speedyseq R packages (23, 24).

Statistical analysis

All analyses were conducted using R version 4.0.4 (25). The OTU dataset was analyzed

with the phyloseq and vegan packages (21, 26). Venn diagrams of sample types and

seasons at the genus level were created using the MicrobiotaProcess package (27, 28).

Differential abundances of bacteria agglomerated to the genus level were explored with

the DESeq2 package version 1.30.1 (ref. 29). The taxa identified as significantly

differentially abundant in the different contrasts (i.e., winter mouth vs. winter crop, winter

nectar vs. winter crop, winter mouth vs. winter nectar, and summer crop vs. summer

mouth; Fig. S1) were aggregated. Of these taxa, those that were present in at least 50%

of the sample type groups (Table S2) were further analyzed with a heatmap. The

heatmap was created with the pheatmap package (30). Differential abundance taxa

phylogeny was made with a GTR model using the phangorn package (31). Sample

cladogram was based on a weighted UniFrac distance (32). Alpha diversity (Shannon

diversity index) in each sample was calculated and compared by fitting a linear mixed

effects model of Shannon diversity index as the response and site as a random effect

with 2000 parametric bootstraps (33). Shannon-Weaver evenness was calculated using

the evenness function from the microbiome package (34). Permutation multivariate

analysis of variance using a weighted UniFrac distance matrix (32) was used to test the

effects of season, sample type, site, and host species on bacterial OTU composition.

The beta diversity of samples was visualized with a Principal Coordinate Analysis, and

individual sample distances from the centroid and the homogeneity of dispersion of the

samples were calculated. We also did Multinomial Species Classification Method

(CLAM) tests, using the clamtest package (35), to identify OTUs found more frequently

in either winter or summer, those found frequently in both seasons, and those too rare

to categorize as summer- or winter-specialists or generalists.

Apilactobacillus kunkeei *strain-level phylogenetic analysis*

Fifty µL of each thawed honeybee crop sample were aliquoted onto De Man, Rogosa

and Sharpe (MRS) agar (Sigma-Aldrich), spread thoroughly using a sterile loop, and

incubated aerobically at 37°C for approximately 48 hours. Half of each single white

colony was picked and placed in 10 μL of MilliQ (EMD Millipore, Burlington, MA, USA)

water, and the other half was streaked onto a new MRS agar plate with a sterile loop.

The 16S rRNA gene of these colonies was amplified using the universal primers 27F

and 1492R in a 25 μL PCR reaction that included 12.5 μL of MyTaq Red Mix (Bioline),

8.75 μL of PCR-grade water, 1.25 μL each of the forward and reverse primers, 0.25 μL

of dimethyl sulfoxide (DMSO), and 1 μL of each diluted colony. The resulting product

was sequenced with Sanger sequencing, and colonies identified as *Apilactobacillus*

kunkeei were once again isolated and placed in 10 μL of MilliQ (EMD Millipore) water.

The variable segments of three housekeeping genes, *lepA*, *recG*, and *rpoB* were

amplified using primers described in Tamarit et al. 2015 and Ogier et al. 2019 (ref. 4,

36). The 25 μL PCR reaction was made up of 11 μL of PCR-grade water, 12 μL of

MyTaq Red Mix (Bioline), 0.5 μL of each forward and reverse primer, and 1 μL of the

diluted colony. The annealing temperature for the PCR reaction was set to 47°C for *lepA*

and recG and 60°C for *rpoB* for 30 seconds (4, 36). The amplified PCR product was

sequenced with Sanger sequencing. The sequences were aligned with MAFFT (37) and

trimmed with TrimAI (38) with a penalty for more than 50% gaps. Individual gene trees

and the partitioned analysis for the multi-gene alignment were inferred using IQ-TREE

(39, 40). Statistical support for the phylogenies was calculated using 1000 replicates of

both a Shimodaira-Hasegawa-like approximation likelihood test and ultrafast

bootstrapping along with a Bayesian-like transformation of aLRT (41-43). The partition

model was selected by the IQ-TREE ModelFinder (Table S3) (44). *Apilactobacillus*

apinorum Fhon13 (ref. 45; Table S4) served as the outgroup for this phylogenetic tree.

Outlier long branches were trimmed with TreeShrink (46).

Results

Crop-mouth-nectar comparisons

Among winter samples, we found more than seven times as many bacterial OTUs in the

crop (a total of 657) as in the mouth and the nectar (a total of 77 and 81, respectively)

(Fig. 2). Mean OTU diversity per sample was about six times higher in the crop (1.78)

than in the mouth (0.37) and the nectar (0.36) (crop-mouth: Welch's t = 17.1, p < 0.001;

crop-nectar: Welch's t = 15.8, p < 0.001). Similarly, mean evenness was about two

times higher in the crop (0.74) than in the mouth (0.42) and the nectar (0.38) (crop-

mouth: Welch's t = 6.8, p < 0.001; crop-nectar: Welch's t = 7, p < 0.001) (Fig. 3).

Of sample type (crop, moth, or nectar), bee species (*A. mellifera* or *A. cerana*), and

sampling site (12 sites shown in Fig. 1), only sample type was a significant predictor of

bacterial OTU composition (PERMANOVA R^2 = 0.10 p = 0.001). Specifically, mouth and

nectar samples were more similar to each other than either mouth and crop samples

were, or nectar and crop samples were to each other (Fig. 4; PERMANOVA mouth-

nectar R^2 = 0.03, p = 0.005; mouth-crop R^2 = 0.11, p = 0.001; nectar-crop R^2 = 0.08, p =

0.001).

In fact, most OTUs in the crop were distinct from those in the mouth and the nectar, with

only 9% (4-7% in *A. cerana* and 6-9% in *A. mellifera*; Fig. S2) of the crop OTUs also

present in the mouth or the nectar (Fig. 2). Taxa frequently found in the crop, such as

Bombella, Apibacter, Gilliamella, Lactobacillus, and *Snodgrassella*, were mostly absent

in the nectar and only infrequently present in the mouth (Figs. 5 and S3). Furthermore,

some OTUs belonging to *Acinetobacter,* Comamonadaceae, *Flavobacterium*, and

Pseudomonas were consistently found in the crop at all local sites examined, whereas

there were no taxa that were present in mouth or nectar samples at all sites.

Summer-winter difference in crop vs. in mouth

Although OTU richness was higher in summer than in winter in both the crop and the

mouth, the magnitude of this difference was eight times greater in the mouth than in the

crop (Figs. 3 and S4). In terms of OTU composition, most of the crop samples were

clustered together regardless of season (winter vs. summer, PERMANOVA R^2 = 0.03, p

= 0.001), whereas the mouth samples formed two distinct clusters, corresponding to

summer and winter (PERMANOVA R^2 = 0.20, p = 0.001; Figs. 4 and 5, Fig. S3). These

two mouth clusters differed in the amount of variation among samples, with winter

showing higher variation than summer (Figs. 4 and S5) despite no significant difference

between them in total bacterial load (Fig. S6).

The CLAM test (35) identified 27% of the OTUs in the crop as generalists, meaning that

they were similarly frequent in winter and summer (Fig. 6). In contrast, the CLAM test

classified only 2% of the OTUs in the mouth as generalists, with the majority of others

identified as either summer- or winter-specialists (76 and 8%, respectively; Fig. 6).

Apilactobacillus kunkeei *strains in the crop*

The multi-gene tree of *Apilactobacillus kunkeei* strains suggested potential season- and

Apis species-specificity. Specifically, the tree had one clade consisting of only isolates

from summer *A. cerana* samples and two other clades consisting of isolates from

summer and winter *A. mellifera* samples (Figs. 7 and S7). Those associated with *A.*

cerana originated from five bees, three from Nishi-Honjo and two from Ichiigawa. Those

associated with *A. mellifera* originated from nine bees: six summer bees from Oshine 1

and three winter bees, one each from Shimanose, Yamauchi, and Kamihaya.

Discussion

Based on the conventional assumption that bacterial assemblage in the crop primarily

consists of transient microbes acquired via nectar foraging (3, 45, 47, 48), we expected

a high degree of overlap among nectar, mouth, and crop samples. We did find high

overlap between the nectar and the mouth, but to our surprise, most of the crop taxa

were clearly distinct from those found in the nectar and the mouth (Figs. 2 and 5).

Several OTUs of *Acinetobacter* and *Pseudomonas* that were relatively prevalent in both

crop and mouth samples in our study have been commonly found in nectar as well (49-

53). However, most of the taxa that were more prevalent in the crop were rare or

entirely absent in nectar and mouth samples (Figs. 5, S1 and S3).

In addition, our results indicate that the crop maintained a high degree of taxonomic

consistency between winter and summer, a pattern that is in sharp contrast to the large

compositional changes seen in the mouth (Figs. 3, 4 and 6). Taxa common in the crop

in both seasons included Comamonadaceae, *Acinetobacter*, Saprospiraceae,

Flavobacterium, *Gilliamella*, *Bombella*, *Apilactobacillus*, *Pseudomonas*, *Fluviicola*,

Polynucleobacter, *Snodgrassella*, *Microbacteriaceae*, and *Rhodococcus* species (Fig.

5). *Apilactobacillus* and *Bombella*, which are among the most commonly reported taxa

in the crop of foraging adults, have also been found frequently in food stores, larvae,

and queens (3-5, 45, 48, 54-56). Although the data we present in this study are

correlational, preventing us from establishing causal relationships, we speculate that the

crop may serve as a consistent reservoir of these taxa that can influence the health of

the adult bees and the larvae that they tend (3, 45).

We also found evidence for potential strain-level specificity among *Apilactobacillus*

kunkeei isolates from collected foragers. One clade was made up of summer *A. cerana*

isolates from two different sites, and two others contained only *A. mellifera* isolates from

both seasons (Figs. 7 and S7). Although a larger sample size than we had in this study

as well as full genome sequencing are needed to ascertain the putative strain-level

differences, previous studies have shown differences in specific house-keeping genes

to suggest strain-level host specificity between *Apis* and *Bombus* and even among *Apis*

species (57-59). As in these previous studies, some of our isolates clustered more

frequently by host species than by site (60, 61), potentially suggesting host specificity

and possibly specific host-symbiont relationships between *Apis* species and *A. kunkeei*

strains, even though this possibility remains speculative at this stage.

A few caveats should be considered in interpreting our findings. First, our analysis is

mostly based on relative abundance (60), even though we used qPCR to attempt to

estimate at least total absolute abundance in crop and mouth samples (Figs. S5 and

S6). Absolute abundance of each taxon, which may be more relevant to bacterial effects

on the host (13, 60), can differ even when relative abundance is the same. Second, we

had only one site, Nishi-Honjo, where bees were sampled in both summer and winter. A

greater and more even number of bees at each site in multi-year seasonal sampling

would have made our results most robust. Nevertheless, our analysis of several

hundred bee and nectar samples all sampled within the same landscape clearly refutes

the hypothesis that crop communities simply reflect what is available in the

environment.

In conclusion, our results suggest that the composition of crop microbes may be more

deterministic than generally assumed, questioning the notion that crop microbes are

environmentally acquired transients that are unimportant to the bees. Although less well

studied than midgut and hindgut microbes, a small but increasing number of recent

studies indicate that some crop microbes do alter the health of individuals and hives (6,

62-64). Our data reinforce the idea that it would help to study how microbial

communities are assembled and maintained not just in the midgut and the hindgut, but

also the crop, to better understand the relationship between honeybees and their gut

microbes.

Acknowledgments

We thank Ryota Nakahaya of Minabe town, Kenichi Hirohata of Tanabe city, and the

staff members at the Japanese Apricot Laboratory of the Wakayama Fruit Tree

Experiment Station for logistical help; Michiko Futaba, Masaru Horiguchi, Tsutomu

Matsuba, Motoki Morikawa, Toshio Nishikawa, Atsuo Okada, Toshiaki Sawa, Yoshinori

Taira, Tamotsu Touroku, Ikunori Yamamoto, Toshihiro Yamanouchi, and Shinichi

Yamazaki for allowing us to collect bees from their hives and orchards; Junji

Takabayashi for laboratory support; Callie Chappell for graphical abstract help; and Taro

Maeda, Junichi Takahashi, and the members of the community ecology group at

Stanford University for discussion. This work was supported by the Office of

International Affairs and the FSI Japan Fund of Stanford University.

Figure legends

Figure 1. Map of collection sites. Foraging adults of *Apis cerana* and *A. mellifera* were

collected from a total of 12 sites (Table S1) in or near Japanese apricot orchards in the

summer of 2018 and the winter of 2019 in the Minabe-Tanabe region, the location of

which is indicated by the red square on the map of Japan. Nectar samples were also

collected from apricot flowers at each of the winter collection sites. Sites are labeled

with circles colored by collection season (summer = white, winter = black, both seasons

= grey). The location of the Japanese Apricot Laboratory is indicated by the red outline

of the Higashi-Honjo site.

Figure 2. The crop contains more common and constant OTUs than the mouth and the

nectar. Only 9% of winter crop OTUs were shared with the mouth or nectar samples

while the mouth and nectar shared about 50% of their OTUs. Venn diagrams are

colored by sample type with OTU counts in bold.

Figure 3. The crop presents more consistent species richness. Species richness

(Shannon diversity) in the crop varied less from summer to winter than the mouth, and

winter mouth and nectar samples had similar species richness across sites. Nishi-

Honjo_W and Nishi-Honjo_S are the samples collected at the Nishi-Honjo site (see

Table S1) in winter and summer, respectively. Each data point represents a single

sample collected from the labeled site, colored by sample type and shaped by host

species. Mean diversity is depicted by the large center point and the error bars depict

the 95% confidence interval for samples from each site.

Figure 4. Principal Coordinate Analysis of all samples presents similar composition across season in the crop but not the mouth. Winter mouth and nectar samples were highly similar (PERMANOVA sample type: $R^2 = 0.07$, p = 0.001, season: $R^2 = 0.05$ p = 0.001, host species: $R^2 = 0.003$, p = 0.154). Each dot represents a sample from an individual bee, colored by sample type. *Apis mellifera* samples are depicted with triangles and the solid ellipse while *Apis cerana* samples are depicted with circles and the dashed ellipse. Ellipses mark the 95% confidence interval, and axes are labeled with corresponding percent variation explained.

Figure 5. There is a greater prevalence of differentially abundant taxa among crop groups than mouth and nectar groups. Samples were grouped by season, sampling site, host *Apis* species, and sample type as illustrated by colored bars above the heatmap and described in Table S2. For ease of visualization, heatmap columns are differentially abundant taxa that were present in at least 50% of one of the sample types e.g., at least 50% of all the mouth groups. The full heatmap with individual samples and all differentially abundant taxa can be found in Fig. S3. The OTUs were agglomerated to genus, colored by prevalence within groups, arranged according to their phylogeny. The y-axis dendrogram depicts the weighted UniFrac distance between samples.

Figure 6. There are bacteria that have high affinity for the crop. Generalist bacteria, i.e., those with no special affinity to one environment or, in this case, season, are greater between the summer and winter crop samples (about 27% of the OTUs) than the

corresponding mouth samples. The mouth presents only about 2% generalists, with

most OTUs classified as summer (76%) specialists. Scatterplots depict the log

abundance of each OTU, with each point representing a single OTU color coded by its

category. OTUs from both *A. mellifera* and *A. cerana* are included. The identity line is

plotted in a dotted grey pattern. The proportion of OTUs classified in each category is

depicted in the corresponding bar plots below the scatterplots, with categories colored

as in the scatterplot.

Figure 7. Multi-gene maximum likelihood tree of *Apilactobacillus kunkeei* isolates shows

Apis species specific clades. The phylogeny is the result of a multi-gene alignment

partitioned analysis of 49 *A. kunkeei* isolates and *A. apinorum* Fhon13 (see Tables S3

and S4). In the depicted tree, substitution frequencies are represented by branch length.

Isolates are labeled starting with an "S" for summer or "W" for winter, followed by a "C"

for *Apis cerana* and an "M" for *Apis mellifera* host species, the host ID number, and

lastly the isolate number. Isolates that occurred in both species are colored in grey while

those that are phylogenetically similar with high statistical support and associated with a

single host species are colored by host species. Statistical support was estimated from

1000 Shimodaira-Hasegawa-approximate likelihood ratio test (SH-aLRT) pseudo

replicates and ultrafast bootstraps (UFboot). Nodes with an SH-aLRT score greater than

0 are labeled with respective SH-aLRT/UFboot values. *Apilactobacillus apinorum* root is

not depicted for ease of visualization. Supplementary Figure 7 shows the full tree with

statistical support on all nodes and *A. apinorum* branch.

References

1.   Moran NA, Hansen AK, Powell JE, Sabree ZL. Distinctive Gut Microbiota of Honey Bees Assessed Using Deep Sampling from Individual Worker Bees. *PLoS One*. 2012;7(4):e36393.

2.   Kwong WK, Moran NA. Gut microbial communities of social bees. *Nat Rev Microbiol*. 2016;14(6):374-84.

3.   Anderson KE, Ricigliano VA, Copeland DC, Mott BM, Maes P. Social Interaction is Unnecessary for Hindgut Microbiome Transmission in Honey Bees: The Effect of Diet and Social Exposure on Tissue-Specific Microbiome Assembly. *Microb Ecol*. 2023;85:1498-513.

4.   Tamarit D, Ellegaard KM, Wikander J, Olofsson T, Vásquez A, Andersson SGE. Functionally Structured Genomes in *Lactobacillus kunkeei* Colonizing the Honey Crop and Food Products of Honeybees and Stingless Bees. *Genome Biol Evol*. 2015;7(6):1455-73.

5.   Corby-Harris V, Snyder LA, Schwan MR, Maes P, McFrederick QS, Anderson KE. Origin and Effect of Alpha 2.2 Acetobacteraceae in Honey Bee Larvae and Description of *Parasaccharibacter apium* gen. nov., sp. nov. *Appl Environ Microbiol*. 2014;80(24):7460-72.

6.   Arredondo D, Castelli L, Porrini MP, Garrido PM, Eguaras MJ, Zunino P, Antúnez K. *Lactobacillus kunkeei* strains decreased the infection by honey bee pathogens *Paenibacillus larvae* and *Nosema ceranae*. *Benef Microbes*. 2018;9(2):279-90.

7.   Kwong WK, Mancenido AL, Moran NA. Immune system stimulation by the native gut microbiota of honey bees. *Roy Soc Open Sci*. 2017;4(2):170003.

8.   Zheng H, Powell JE, Steele MI, Dietrich C, Moran NA. Honeybee gut microbiota

promotes host weight gain via bacterial metabolism and hormonal signaling. *Proc Natl*

Acad Sci USA. 2017;114(18):4775-80.

9.   Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. Disentangling

metabolic functions of bacteria in the honey bee gut. *PLoS Biol*. 2017;15(12):e2003467.

10.   Engel P, Martinson VG, Moran NA. Functional diversity within the simple gut

microbiota of the honey bee. *Proc Natl Acad Sci USA*. 2012;109(27):11002-7.

11.   Hart AG, Ratnieks FLW. Why do honey-bee (*Apis millifera*) foragers transfer

nectar to several receivers? Information improvement through multiple sampling in a

biological system. *Behav Ecol Sociobiol*. 2001;49(4):244-50.

12.   Decker LE, San Juan PA, Warren ML, Duckworth CE, Gao C, Fukami T. Higher

Variability in Fungi Compared to Bacteria in the Foraging Honey Bee Gut. *Microb Ecol*.

2022;8:1-5.

13.   Kešnerová L, Emery O, Troilo M, Liberti J, Erkosar B, Engel P. Gut microbiota

structure differs between honeybees in winter and summer. *ISME J*. 2020;14(3):801-14.

14.   Maeda T, Hiraiwa MK, Shimomura Y, Oe T. Weather conditions affect pollinator

activity, fruit set rate, and yield in Japanese apricot. *Sci Hortic*. 2023;307:111522.

15.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al.

Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq

platforms. *ISME J*. 2012;6(8):1621-4.

16.   Toju H, Vannette RL, Gauthier M-PL, Dhami MK, Fukami T. Priority effects can

persist across floral generations in nectar microbial metacommunities. *Oikos*.

2018;127(3):345-52.

17.   Ritalahti KM, Amos BK, Sung Y, Wu Q, Koenigsberg SS, Löffler FE. Quantitative

PCR targeting 16S rRNA and reductive dehalogenase genes simultaneously monitors

multiple Dehalococcoides strains. *Appl Environ Microbiol*. 2006;72(4):2765-74.

18.   Martin M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet J*. 2011;17:10-2.

19.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.

DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods*.

2016;13(7):581-3.

20.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, et al. The SILVA

and "all-species living tree project (LTP)" taxonomic frameworks. *Nucleic Acids Res*.

2014;42(D1):D643-D8.

21.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One*. 2013;8(4):e61217.

22.   Smith EA, Martin-Eberhardt SA, Miller DL, Parish AJ, Newton ILG. Draft Genome

Sequence of a *Bombella apis* Strain Isolated from Honey Bees. *Microbiol Resour*

Announc. 2019;8(47):e01329-19.

23.   Wright ES. Using DECIPHER v2. 0 to analyze big biological sequence data in R.

R J. 2016;8(1).

24.   McLaren M. speedyseq: Faster implementations of common phyloseq functions.

2020.

25.   Team RC. R: A Language and Environment for Statistical Computing. In:

Computing RffS, editor. Vienna, Austria2023.

26.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al.

vegan: Community Ecology Package. R package version 2.5-6. 2019.

27.   Xu S, Zhan L, Tang W, Wang Q, Dai Z, Zhou L, et al. MicrobiotaProcess: A

comprehensive R package for deep mining microbiome. *Innovation*. 2023;4(2).

28.   Chen H. _VennDiagram: Generate High-Resolution Venn and Euler Plots_. R

package version 1.7.3. 2022.

29.   Love MI, Huber W, Anders S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550.

30.   Kolde R. _pheatmap: Pretty Heatmaps_. R package version 1.0.12. 2019.

31.   Schliep K. phangorn: phylogenetic analysis in R. *Bioinformatics*. 2011;27:592e3.

32.   Hamady M, Lozupone C, Knight R. Fast UniFrac: facilitating high-throughput

phylogenetic analyses of microbial communities including analysis of pyrosequencing

and PhyloChip data. *ISME J*. 2010;4(1):17-27.

33.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models

using lme4. *J Stat Softw*. 2015;67(1):1-48.

34.   Lahti L, Shetty S. microbiome R package. 2017.

35.   Chazdon RL, Chao A, Colwell RK, Lin S-Y, Norden N, Letcher SG, et al. A novel

statistical method for classifying habitat generalists and specialists. *Ecology*.

2011;92(6):1332-43.

36.   Ogier J-C, Pagès S, Galan M, Barret M, Gaudriault S. rpoB, a promising marker

for analyzing the diversity of bacterial communities by amplicon sequencing. *BMC*

Microbiol. 2019;19(1):171.

37.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol*. 2013;30(4):772-80.

38.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*.

2009;25(15):1972-3.

39.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von

Haeseler A, Lanfear R. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Mol Biol Evol*. 2020;37(5):1530-4.

40.   Chernomor O, von Haeseler A, Minh BQ. Terrace Aware Data Structure for

Phylogenomic Inference from Supermatrices. *Syst Biol*. 2016;65(6):997-1008.

41.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New

Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the

Performance of PhyML 3.0. *Syst Biol*. 2010;59(3):307-21.

42.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2:

Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol*. 2017;35(2):518-22.

43.   Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. Survey of Branch

Support Methods Demonstrates Accuracy, Power, and Robustness of Fast Likelihood-

based Approximation Schemes. *Syst Biol*. 2011;60(5):685-99.

44.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*.

2017;14(6):587-9.

45. Dyrhage K, Garcia-Montaner A, Tamarit D, Seeger C, Näslund K, Olofsson TC, et

al. Genome evolution of a symbiont population for pathogen defense in honeybees.

Genome Biol Evol. 2022;14(11):evac153.

46. Mai U, Mirarab S. TreeShrink: fast and accurate detection of outlier long

branches in collections of phylogenetic trees. *BMC Genomics*. 2018;19(5):23-40.

47. Anderson KE, Maes P. Social microbiota and social gland gene expression of

worker honey bees by age and climate. *Sci Rep*. 2022;12(1):10690.

48. Anderson KE, Sheehan TH, Mott BM, Maes P, Snyder L, Schwan MR, et al.

Microbial ecology of the hive and pollination landscape: bacterial associates from floral

nectar, the alimentary tract and stored food of honey bees (Apis mellifera). *PLoS One*.

2013;8(12):e83125.

49. Alvarez-Perez S, Baker LJ, Morris MM, Tsuji K, Sanchez VA, Fukami T, et al.

Acinetobacter pollinis sp. nov., Acinetobacter baretiae sp. nov. and Acinetobacter

rathckeae sp. nov., isolated from floral nectar and honey bees. *Int J Syst Evol Microbiol*.

2021;71(5):004783.

50. Álvarez-Pérez S, Herrera CM, Vega C. Zooming-in on floral nectar: a first

exploration of nectar-associated bacteria in wild plant communities. *FEMS Microbiol*

Ecol. 2012;80(3):591-602.

51. Ben Belgacem Z, Bijttebier S, Verreth C, Voorspoels S, Van de Voorde I, Aerts G,

et al. Biosurfactant production by *Pseudomonas* strains isolated from floral nectar. *J*

Appl Microbiol. 2015;118(6):1370-84.

52. Fridman S, Izhaki I, Gerchman Y, Halpern M. Bacterial communities in floral

nectar. *Environ Microbiol Rep*. 2012;4(1):97-104.

53.   Morris MM, Frixione NJ, Burkert AC, Dinsdale EA, Vannette RL. Microbial

abundance, composition, and function in nectar are shaped by flower visitor identity.

FEMS Microbiol Ecol. 2020;96(3):fiaa003.

54.   Vásquez A, Forsgren E, Fries I, Paxton RJ, Flaberg E, Szekely L, Olofsson TC.

Symbionts as Major Modulators of Insect Health: Lactic Acid Bacteria and Honeybees.

PLoS One. 2012;7(3):e33188.

55.   Corby-Harris V, Maes P, Anderson KE. The bacterial communities associated

with honey bee (Apis mellifera) foragers. *PLoS One*. 2014;9(4):e95056.

56.   Rangberg A, Mathiesen G, Amdam GV, Diep DB. The paratransgenic potential of

Lactobacillus kunkeei in the honey bee *Apis mellifera*. *Benef Microbes*. 2015;6(4):513-

23.

57.   Koch H, Abrol DP, Li J, Schmid-Hempel P. Diversity and evolutionary patterns of

bacterial gut associates of corbiculate bees. *Mol Ecol*. 2013;22(7):2028-44.

58.   Kwong WK, Engel P, Koch H, Moran NA. Genomics and host specialization of

honey bee and bumble bee gut symbionts. *Proc Natl Acad Sci USA*.

2014;111(31):11509-14.

59.   Kwong WK, Medina LA, Koch H, Sing K-W, Soh EJY, Ascher JS, et al. Dynamic

microbiome evolution in social bees. *Sci Adv*. 2017;3(3):e1600513.

60.   Almeida EL, Ribiere C, Frei W, Kenny D, Coffey MF, O'Toole PW. Geographical

and Seasonal Analysis of the Honeybee Microbiome. *Microb Ecol*. 2023;85:765-78.

61.   Ellegaard KM, Suenami S, Miyazaki R, Engel P. Vast Differences in Strain-Level

Diversity in the Gut Microbiota of Two Closely Related Honey Bee Species. *Curr Biol*.

2020;30(13):2520-31.e7.

62.   Chege M, Kinyua J, Paredes JC. Lactobacillus kunkeei impacts the health of

honey bees, Apis mellifera scutellata, and protects the bees against the opportunistic

pathogen Serratia marcescens. *Int J Trop Insect Sci*. 2023;43:1947-55.

63.   Corby-Harris V, Snyder L, Meador CAD, Naldo R, Mott B, Anderson KE.

Parasaccharibacter apium , gen. nov., sp. nov., Improves Honey Bee (Hymenoptera:

Apidae) Resistance to *Nosema*. *J Econ Entomol*. 2016;109(2):537-43.

64.   Miller DL, Smith EA, Newton ILG. A Bacterial Symbiont Protects Honey Bees

from Fungal Disease. *mBio*. 2021;12(3):e00503-21.

Figure 1

Figure 2

Figure 3

Sample type crop mouth nectar **Bee species** ⊙ *Apis cerana* △ *Apis mellifera*

Figure 4

Figure 5

Figure 6

Figure 7

0.01