Bacteria in honeybee crops are decoupled from those in floral nectar and bee mouths

Magdalena L. Warren[1*†], Kaoru Tsuji[1,2,3*], Leslie E. Decker[1], Manabu Kishi[4,5], Jihoon Yang[6], Adina C. Howe[6], Tadashi Fukami[1,7†]

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA

[2]Center for Ecological Research, Kyoto University, 2-Hirano, Otsu, Shiga, 520-2113, Japan

[3]Department of Biology, Graduate School of Science, Kobe University, Kobe, 657-8501, Japan

[4]Japanese Apricot Laboratory, Wakayama Fruit Tree Experiment Station, Minabe, Higashi-Honjo 1416-7, Wakayama, 645-0021, Japan

[5]Hidaka Promotion Bureau, Wakayama Prefecture, Gobo, Yukawa 651, Wakayama, 644-0011, Japan

[6]Department of Agricultural and Biosystems Engineering, Iowa State University, Ames, Iowa, USA

[7]Department of Earth System Science, Stanford University, Stanford, CA 94305, USA

* These authors contributed equally to this work

[†]Co-corresponding authors:

mlwarren@stanford.edu; ORCID: 0000-0003-0115-5066

fukamit@stanford.edu; ORCID: 0000-0001-5654-478

Acknowledgments

We thank Ryota Nakahaya of Minabe town, Kenichi Hirohata of Tanabe city, and the

staff members at the Japanese Apricot Laboratory of the Wakayama Fruit Tree

Experiment Station for logistical help; Michiko Futaba, Masaru Horiguchi, Tsutomu

Matsuba, Motoki Morikawa, Toshio Nishikawa, Atsuo Okada, Toshiaki Sawa, Yoshinori

Taira, Tamotsu Touroku, Ikunori Yamamoto, Toshihiro Yamanouchi, and Shinichi

Yamazaki for allowing us to collect bees from their hives and orchards; Junji

Takabayashi for laboratory support; Callie Chappell for graphical abstract help; and Taro

Maeda, Junichi Takahashi, and the members of the community ecology group at

Stanford University for discussion. We also thank Josh Neufeld, Irene Newton, Julie

Olson, and five anonymous reviewers for comments on early versions of this

manuscript. This work was supported by the Office of International Affairs and the

Freeman Spogli Institute for International Studies Japan Fund of Stanford University.

Abstract

Bacteria in the honeybee gut are a well-recognized factor affecting bee health. However, the primary focus of this research has been the hindgut, while the crop, or honey stomach, is assumed to be dominated by environmentally acquired transient taxa that matter little to the bees. To evaluate this assumption, we examined bacterial taxa in the crop and mouth of *Apis mellifera* and *A. cerana japonica* foragers and in the nectar of *Prunus mume* flowers visited by the bees in the Minabe-Tanabe region of Japan. We found that in bacterial composition, the crop was distinct from both the mouth and the nectar, whereas mouth and nectar samples were indistinguishable. Furthermore, the crop remained similar in bacterial composition and diversity, while the mouth showed a sharp drop in alpha diversity and a large increase in beta diversity, from summer to winter. These results refute the conventional assumption, suggesting instead that the crop contains a conserved bacterial community largely distinct from environmental taxa. We also found that strains of a crop-associated species, *Apilactobacillus kunkeei*, could be season- and host species-specific. Together, these findings suggest that crop-associated bacterial communities should be studied further to better understand the relationship between honeybees and their gut bacteria.

Keywords: microbial ecology; honeybee microbiome; nectar microbiome; plant-pollinator-microbe interactions

Introduction

Microbes in the honeybee gut are receiving increasing attention not only as a factor affecting the health of the agriculturally vital insects, but also as a model system for gaining basic understanding of gut-associated microbiota [1, 2]. One component of the honeybee gut is used for the transport of liquids. This organ, which is an expandable portion of the esophagus hereby termed the crop, but also known as the honey stomach or honey sac [3], contains species such as *Apilactobacillus kunkeei* and *Bombella apis* that may affect the survival and pathogen resistance of the foraging adults that host them and the larvae that the adults feed [1, 4-6]. However, microbes in the crop are not as well studied as those in the hindgut, which are generally considered more important to honeybee health [7, 8].

Given its function for temporary storage and transport of floral nectar [9], it seems reasonable to assume that the crop is occupied mainly by the transient microbes acquired from the environment via foraging of nectar [10, 11]. One way to investigate this assumption of environment–crop matching is to compare microbes in the crop of foraging bees and those in their mouth as well as in the floral nectar they forage for. If the assumption is true, the crop should be similar to both the mouth and the floral nectar in bacterial composition [4, 12]. To our knowledge, no study has directly compared bacteria in crop, mouth, and nectar for this purpose, leaving untested the assumption that the crop microbiota is dominated by environmentally derived transients.

Here we test the hypothesis that crop-associated bacteria show species composition

that mirror those observed in mouth- and nectar-associated bacteria. To this end, we

compare crop-, mouth-, and nectar-associated bacteria in samples that were collected

simultaneously at the same locations. Our crop and mouth samples come from actively

foraging adults of the introduced *Apis mellifera* and the native *A. cerana japonica*

(hereafter *A. cerana*) collected in the Minabe-Tanabe region of Wakayama Prefecture in

Japan. In this region, farmers use both species of honeybees to pollinate the winter-

blooming Japanese apricot, *Prunus mume* [13]. Our nectar samples come from *P.*

mume flowers collected in the winter near the hives of the *A. mellifera* and *A. cerana*

bees that we caught for crop and mouth sampling.

Materials and Methods

Study sites

Japanese apricot orchards in the Minabe-Tanabe region are embedded within a

mountainous countryside landscape that has recently become internationally

recognized. In 2015, the Food and Agricultural Organization designated the region as a

Globally Important Agricultural Heritage System for the agricultural practice in the region

that embraces both human needs and biodiversity conservation. Many of the apricot

orchards are located adjacent to natural and managed forests, which are thought to

provide stable habitats for the native *A. cerana*. The two *Apis* species are the main

pollinators of the apricot, which blooms from mid or late February to early or mid-March,

during which few other flower-visiting insects are available [13]. In the winter, *A.*

mellifera colonies are supplemented with a sugar solution, while *A. cerana* colonies are

not. Although there may have been several different blooming species during our

collections in winter and summer, our interest in nectar focused on the economically

important *P. mume.* All our bee and nectar collection sites were located within this

landscape, but at least 1.0 km away from their respective nearest neighboring collection

site in each season (Fig. 1a, Table S1).

Bee collection

A total of 12 *P. mume* orchards in the study region were visited in the summer of 2018

and the winter of 2019. Seventy-two and 149 honeybees were collected in the summer

of 2018 and the winter of 2019, respectively, from hives in or near these apricot

orchards (Table S1). At each collection site, there were about four to 10 *A. cerana* hives

or 10 to 40 *A. mellifera* hives. Honeybees were collected as they flew near their hives

located within or near the orchards. At orchards where the beekeepers maintained *A.*

*ceran*a hives, *A. cerana* bees were collected near their hives whose locations varied

from tens of meters to about a hundred meters from the orchard and were often situated

at the edge of the local forest. The collected bees were individually placed in a sterile,

sealed plastic vial immediately after the collection. The vials were kept for up to two

hours in a cool box with ice packs in it for transport to the Japanese Apricot Laboratory

located within 100 m from one of the collection sites (Higashi-Honjo) (Fig. 1a).

Fig. 1 Collection sites and crop dissection. (**a**) Foraging adults of *Apis cerana* and *A. mellifera* were collected from a total of 12 sites (see Table S1) in or near Japanese apricot orchards in the summer of 2018 and the winter of 2019 in the Minabe-Tanabe region, the location of which is indicated by the red square on the map of Japan. Nectar samples were also collected from apricot flowers at each of the winter collection sites. Sites are labeled with circles colored by collection season (summer = white, winter = black, both seasons = grey). The location of the Japanese Apricot Laboratory is indicated by the red outline of the Higashi-Honjo site. (**b**) The crop was distinctly separated from the rest of the alimentary tract. (**c**) The crop of each honeybee was sterilely dissected and placed in an aliquot of sterile water

After we brought the collected bees into the laboratory in the Japanese Apricot

Laboratory facility, the bees were kept cool in the box for up to two additional hours so

that the bees would stay sluggish for ease of handling. The captured bees were then

taken out of the vials, and their wings were fixated with adhesive tape to facilitate

handling. The taped bees were made to drink approximately half of 20 μL of sterile 20%

sucrose solution placed on the lid of a sterile 1.5-mL microcentrifuge tube. After the

feeding, the remaining solution was placed in the microcentrifuge tube with the lid now

closed. These solutions were later used to study the bacteria associated with the mouth.

We refer to these solutions as mouth samples. As we had the bees extend their

proboscis into the sterile sucrose solution, our method is likely to have captured mostly

the bacteria from the surface of the proboscis rather than those present within the

mandibles.

Immediately after the sampling of mouth-associated bacteria, we dissected the bees by

gently pulling the abdomen using tweezers to expose the crop, making sure the crop did

not touch other parts of the bee body. We then removed the crop from the bee body

using another set of tweezers (Fig. 1b). Both tweezers were sterilized before each use

by dipping in 70% ethanol and then flaming. Each crop sample was separately placed in

40 μL of sterile water within sterile microcentrifuge tubes (Fig. 1c) and homogenized

using a sterile plastic pestle. All mouth and crop samples were immediately stored at -

80°C until further processing.

We collected crop and mouth samples not just in winter during the *P. mume* flowering

season, but also in summer, to determine whether crop and mouth differed in the way

bacterial composition changed from summer to winter. Hives of *A. cerana*, many of

which were made of carved tree trunks, were maintained by local beekeepers

throughout the year in the region. In contrast, most *A. mellifera* hives in the region were

transported elsewhere in the country while *P. mume* was not in bloom. However, some

were kept locally throughout the year, allowing us to collect bees in both seasons. One

site (Nishi-Honjo) was available for bee collection in both summer and winter, allowing

bacterial taxa from the same site and hives to be compared between seasons. In

addition, at two of our sites (Higashi-Honjo and Kamihaya), we were able to collect both

A. cerana and *A. mellifera*, enabling comparison of the two species collected at the

same locations (Table S1).

Nectar collection

In addition to the mouth and crop samples, we also collected samples of *P. mume* floral

nectar on the same days the bees were collected in the winter of 2019. At each of the

nine orchards where we sampled bees that winter (Table S1), we collected 16 nectar

samples (except at one site, Kamihaya, where we instead collected 24 nectar samples),

for a total of 152 nectar samples. Each sample was collected by probing 10 flowers on a

randomly selected branch of an individual *P. mume* tree with a sterile 0.5μl

microcapillary tube and dispensing the collected nectar into a PCR tube containing 40

μL of PCR-quality sterile deionized water. The collected nectar samples were

immediately placed in a box with ice packs during the transport to the laboratory. Within

six hours, the nectar samples were brought back to the Japanese Apricot Laboratory.

The samples were kept at -80°C until further processing for DNA extraction, except

during a one-day transport to the laboratory at Kyoto University in Shiga, Japan, and

another one-day transport from there to Stanford University in California, USA, during

which the samples were kept at about -5 °C.

Microbial DNA analysis

179 Using the nectar and bee samples obtained as above, we conducted microbial DNA

180 extraction, bacterial amplicon sequencing, quantitative real-time PCR (qPCR) of 16S

181 rRNA gene, and processing of the amplicon sequencing data (for details, see Text S1).

Statistical analysis

All analyses were conducted using R version 4.0.4 [14]. The ASV_{97} dataset was

analyzed with the phyloseq and vegan packages [15, 16]. Venn diagrams of sample

types and seasons at the Genus level were created using the MicrobiotaProcess

package [17, 18]. Differential abundances of bacteria agglomerated to the Genus level

were explored with the DESeq2 package version 1.30.1 [19]. The taxa identified as

significantly differentially abundant in the different contrasts (i.e. winter mouth vs. winter

crop, winter nectar vs. winter crop, winter mouth vs. winter nectar, and summer crop vs.

summer mouth; Fig S1) were aggregated. Of these taxa, those that were present in at

least 50% of the sample type groups (Table S2) were further analyzed with a heatmap.

The heatmap was created with the ComplexHeatmap package [20]. Differential

abundance taxa phylogeny was made with a GTR model using the phangorn package

[21]. Sample cladogram was based on a weighted UniFrac distance [22]. Alpha diversity

(Shannon index) in each sample was calculated and compared by fitting a linear mixed

effects model of Shannon index as the response and site as a random effect with 2000

parametric bootstraps [23]. Shannon-Weaver evenness was calculated using the

evenness function from the microbiome package [24]. Permutation multivariate analysis

of variance using a weighted UniFrac distance matrix [22] was used to test the effects of

season, sample type, site, and host species on bacterial ASV_{97} composition. The beta

diversity of relative abundance and absolute abundance samples was visualized with a

Principal Coordinate Analysis, and individual sample distances from the centroid and

the homogeneity of dispersion of the samples were calculated. We also did Multinomial

Species Classification Method (CLAM) tests, using the clamtest package [25], to identify

$ASVs_{97}$ found more frequently in either winter or summer, those found frequently in both

seasons, and those too rare to categorize as summer- or winter-associated, or

generalists.

Apilactobacillus kunkeei *strain-level phylogenetic analysis*

Fifty µL of each thawed honeybee crop sample were aliquoted onto De Man, Rogosa

and Sharpe (MRS) agar (Sigma-Aldrich), spread thoroughly using a sterile loop, and

incubated aerobically at 37°C for approximately 48 hours. Half of each single white

colony was picked and placed in 10 µL of MilliQ (EMD Millipore, Burlington, MA, USA)

water, and the other half was streaked onto a new MRS agar plate with a sterile loop.

The 16S rRNA gene of these colonies was amplified using the universal primers 27F

and 1492R in a 25 µL PCR amplification that included 12.5 µL of MyTaq Red Mix

(Bioline), 8.75 µL of PCR-grade water, 1.25 µL each of the forward and reverse primers,

0.25 µL of dimethyl sulfoxide (DMSO), and 1 µL of each diluted colony. The resulting

product was sequenced with Sanger sequencing, and colonies identified as

Apilactobacillus kunkeei were once again isolated and placed in 10 µL of MilliQ (EMD

Millipore) water. The variable segments of three housekeeping genes, *lepA, recG,* and

rpoB were amplified using previously described primers [4, 26]. The 25 μL PCR

amplification was made up of 11 μL of PCR-grade water, 12 μL of MyTaq Red Mix

(Bioline), 0.5 μL of each forward and reverse primer, and 1 μL of the diluted colony. The

annealing temperature for the PCR amplification was set to 47°C for *lepA* and *recG* and

60°C for *rpoB* for 30 seconds [4, 26]. The amplified PCR product was sequenced with

Sanger sequencing. As previously described [27], the sequences were aligned with

MAFFT [28] and trimmed with TrimAl [29] with a penalty for more than 50% gaps.

Individual gene trees and the partitioned analysis for the multi-gene alignment were

inferred using IQ-TREE [27, 30]. Statistical support for the phylogenies was calculated

using 1000 replicates of both a Shimodaira-Hasegawa-like approximation likelihood test

and ultrafast bootstrapping along with a Bayesian-like transformation of aLRT [27, 31,

32]. The partition model was selected by the IQ-TREE ModelFinder (Table S3) [27, 33].

Apilactobacillus apinorum Fhon13 (Table S4) [27] served as the outgroup for this

phylogenetic tree. Outlier long branches were trimmed with TreeShrink [34].

Results

Crop–mouth–nectar comparisons

Among winter samples, we found more than seven times as many bacterial $ASVs_{97}$ in

the crop (a total of 655) as in the mouth and the nectar (a total of 77 and 81,

respectively) (Fig. 2). Mean ASV_{97} diversity per sample was about six times higher in the

crop (1.78) than in the mouth (0.37) and the nectar (0.36) (crop–mouth: Welch's t =

17.1, *p-value* < 0.001; crop–nectar: Welch's t = 15.8, *p-value* < 0.001). Similarly, mean

evenness was about two times higher in the crop (0.74) than in the mouth (0.42) and

the nectar (0.38) (crop–mouth: Welch's t = 6.8, *p-value* < 0.001; crop–nectar: Welch's t

= 7, *p-value* < 0.001) (Fig. 3).

Fig. 2 The crop contains more common and constant $ASVs_{97}$ than the mouth and the nectar. Only 9% of 655 winter crop $ASVs_{97}$ were shared with the mouth or nectar samples whereas the mouth and nectar shared about 50% of their 77 and 81 $ASVs_{97,}$ respectively. Venn diagrams are colored by sample type with ASV_{97} counts in bold

Fig. 3 The crop presents more consistent species richness. Species richness (Shannon index) in the crop varied less from summer to winter than the mouth, and winter mouth and nectar samples had similar species richness across sites. Nishi-Honjo_W and Nishi-Honjo_S are the samples collected at the Nishi-Honjo site (see Table S1) in winter and summer, respectively. Each data point represents a single sample collected from the labeled site, colored by sample type, and shaped by host species. Mean diversity is depicted by the large center point and the error bars depict the bootstrapped 95% confidence interval for samples from each site.

Of sample type (crop, mouth, or nectar), bee species (*A. mellifera* or *A. cerana*), and

sampling site (Fig. 1a), only sample type was a significant predictor of bacterial ASV_{97}

composition (PERMANOVA R^2 = 0.10 *p-value* = 0.001). Specifically, mouth and nectar

samples were more similar to each other in bacterial composition than either mouth and

crop samples or nectar and crop samples were to each other (Fig. 4; PERMANOVA

mouth–nectar R^2 = 0.03, *p-value* = 0.002; mouth–crop R^2 = 0.11, *p-value* = 0.001;

nectar–crop R^2 = 0.08, *p-value* = 0.001).

Fig. 4 Principal coordinate analysis of all samples presents similar composition across season in the crop but not the mouth. Winter mouth and nectar samples were highly similar (PERMANOVA sample type: R^2 = 0.07, *p-value* = 0.001, season: R^2 = 0.05 *p-value* = 0.001, host species: R^2 = 0.003, *p-value* = 0.154). Each dot represents a sample from an individual bee, colored by sample type. *Apis mellifera* samples are depicted with triangles and the solid ellipse whereas *Apis cerana* samples are depicted with circles and the dashed ellipse. Ellipses mark the 95% confidence interval, and axes are labeled with corresponding percent variation explained

In fact, most ASVs$_{97}$ in the crop were distinct from those in the mouth and the nectar,

with only 9% (4-8% in *A. cerana* and 6-9% in *A. mellifera*; Fig. S2) of the crop ASVs$_{97}$

also present in the mouth or the nectar (Fig. 2). Taxa frequently found in the crop, such

as *Bombella, Gilliamella, Lactobacillus*, and *Snodgrassella*, were mostly absent in the

nectar and only infrequently present in the mouth (Figs. 5 and S3). Furthermore, some

ASVs$_{97}$ belonging to *Acinetobacter,* Comamonadaceae, and *Flavobacterium* were

consistently found in the crop at all local sites examined, whereas *Acinetobacter* was

present in only seven of the nine nectar sites and 11 of the 15 mouth sites, and there

were no other taxa that were present in mouth or nectar samples at all sites.

Fig. 5 There is a greater prevalence of differentially abundant taxa among crop groups than mouth and nectar groups. Samples were grouped by season, sampling site, host *Apis* species, and sample type as illustrated by colored bars to the left of the heatmap and described in Table S2. For ease of visualization, heatmap columns are differentially abundant taxa that were present in at least 50% of one of the sample groups e.g., at least 50% of all the mouth groups. Burkholderia* denotes the genus Burkholderia-Caballeronia-Paraburkholderia. The full heatmap with individual samples and all differentially abundant taxa can be found in Figure S3. The ASVs$_{97}$ were agglomerated to genus, colored by prevalence within groups, and arranged according to their phylogeny. The y-axis dendrogram depicts the weighted UniFrac distance between samples

These results held true even when we excluded those taxa that are known as the core

species of the hindgut bacterial community (*Gilliamella*, *Lactobacillus*, and

Snodgrassella) from the dataset used for the analysis (Fig. S4).

Summer–winter difference in crop vs. in mouth

Although ASV_{97} richness was higher in summer than in winter in both the crop and the

mouth, the magnitude of this difference was eight times greater in the mouth than in the

crop (Figs. 3 and S5). In terms of ASV_{97} composition, most of the crop samples were

clustered together regardless of season in both the relative and absolute abundance

datasets (relative abundance: winter vs. summer, PERMANOVA R^2 = 0.03, *p-value* =

0.001; absolute abundance: winter vs. summer, PERMANOVA R^2 = 0.04, *p-value* =

0.021), whereas the mouth samples formed two distinct clusters, corresponding to

summer and winter (relative abundance: PERMANOVA R^2 = 0.20, *p-value* = 0.001;

absolute abundance: PERMANOVA R^2 = 0.31, *p-value* = 0.001; Figs. 4-6a). These two

mouth clusters differed in the amount of bacterial composition variation among samples,

with winter mouth samples showing higher variation than summer mouth samples (Figs.

4 and 6a) despite no significant difference between them in total bacterial load (Fig. 6b).

In contrast to the mouth, the bacterial load in the crop was significantly higher in the

summer than in the winter in the *A. mellifera* crop (but not in *A. cerana* crop; Fig. 6b).

Fig. 6 Consideration of absolute abundance estimates in analysis of bacterial composition. (**a**) As in Figure 4, principal coordinate analysis of samples that takes into account absolute total bacterial abundance indicates greater compositional similarity between crop samples than mouth samples across season. Specifically, mouth samples cluster more closely in the summer and are more dispersed in the winter than the crop samples (betadisper distance to median: summer-mouth = 0.21, winter-mouth = 0.40, summer-crop = 0.26, winter-crop = 0.32; betadisper Tukey HSD summer-winter mouth: *p-value* < 0.001, summer-winter crop: *p-value* = 0.008). Overall, sample type and season but not host species are significant predictors of composition (PERMANOVA sample type: R^2 = 0.03, *p-value* = 0.009, season: R^2 = 0.12, *p-value* = 0.001, host species: R^2 = 0.01, *p-value* = 0.214). Each point represents a single sample, is color coded by sample type, and shaped by bee species. Ellipses represent the 95% confidence interval for each bee species. Percent variation explained by each axis is labeled. (**b**) Mouth samples tended to have a more even, lower bacterial load across season than did crop samples. The *Apis mellifera* but not the *Apis cerana* crop samples had a higher bacterial load in the summer than the winter. The mouth maintained the same bacterial load across seasons. Total 16S rRNA gene copies per 10 μl are presented on the log scale, with each point depicting a sample from an individual bee colored by season (summer = white, winter = black). Wilcoxon test significance is labeled with *** = *p-value* < 0.001 and ns = not significant

296   The CLAM test [25] identified 28% of the $ASVs_{97}$ in the crop as generalists, meaning

297   that they were similarly frequent in winter and summer. In contrast, the CLAM test

298   classified only 2% of the $ASVs_{97}$ in the mouth as generalists, with the majority of others

299   identified as either summer- or winter-associated (76 and 8%, respectively; Fig. 7).

Fig. 7 There are bacteria that have high affinity for the crop. Generalist bacteria, i.e., those with no special affinity to one environment or, in this case, season, are greater between the summer and winter crop samples (about 27% of the ASVs$_{97}$) than the corresponding mouth samples. The mouth presents only about 2% generalists, with most ASVs$_{97}$ classified as summer-associated (76%). Scatterplots depict the log abundance of each ASV$_{97}$, with each point representing a single ASV$_{97}$ color coded by its category. ASVs$_{97}$ from both *A. mellifera* and *A. cerana* are included. The identity line is plotted in a dotted grey pattern. The proportion of ASVs$_{97}$ classified in each category is depicted in the corresponding bar plots below the scatterplots, with categories colored as in the scatterplot

Apilactobacillus kunkeei *strains in the crop*

The multi-gene tree of *Apilactobacillus kunkeei* strains suggested potential season- and

Apis species-specificity. Specifically, the tree had one clade consisting of only isolates

from summer *A. cerana* samples and two other clades consisting of isolates from

summer and winter *A. mellifera* samples (Figs. 8 and S6). Those associated with *A.*

cerana originated from five bees, three from Nishi-Honjo and two from Ichiigawa. Those

associated with *A. mellifera* originated from eight bees: five summer bees from Oshine 1

and three winter bees, one each from Shimanose, Yamauchi, and Kamihaya.

Fig. 8 Multi-gene maximum likelihood tree of *Apilactobacillus kunkeei* isolates contains *Apis* species specific clades. The phylogeny is the result of a multi-gene alignment partitioned analysis of 49 *A. kunkeei* isolates and *A. apinorum* Fhon13 (see Tables S3 and S4). In the depicted tree, substitution frequencies are represented by branch length. Isolates are labeled starting with an "S" for summer or "W" for winter, followed by a "C" for *Apis cerana* and an "M" for *Apis mellifera* host species, the host ID number, and lastly the isolate number. Isolates that occurred in both species are colored in grey whereas those that are phylogenetically similar with high statistical support and associated with a single host species are colored by host species. Statistical support was estimated from 1000 Shimodaira-Hasegawa-approximate likelihood ratio test (SH-aLRT) pseudo replicates and ultrafast bootstraps (UFboot). Nodes with an SH-aLRT score greater than 0 are labeled with respective SH-aLRT/UFboot values. *Apilactobacillus apinorum* root is not depicted for ease of visualization. Figure S6 presents the full tree with statistical support on all nodes and *A. apinorum* branch

Discussion

Based on the conventional assumption that bacterial assemblage in the crop primarily

consists of transient microbes acquired via nectar foraging [2, 27, 35, 36], we expected

a high degree of bacterial overlap among nectar, mouth, and crop samples. We did find

high overlap between the nectar and the mouth in the winter, but to our surprise, most of

the crop taxa were clearly distinct from those found in the nectar and the mouth (Figs. 2

and 5) [11]. Several $ASVs_{97}$ of *Acinetobacter* and *Pseudomonas* that were relatively

prevalent in both crop and mouth samples in our study have been commonly found in

nectar as well [37-39]. However, most of the taxa that were more prevalent in the crop

were rare or entirely absent in nectar and mouth samples (Figs. 5, S2 and S3) [11].

In addition, our results indicate that the crop maintained a high degree of taxonomic

consistency between winter and summer despite differences in bacterial load in *A.*

mellifera crop samples, a pattern that is in sharp contrast to the large compositional

changes seen in the mouth between seasons and between bee species (Figs. 3, 4, 6a

and 7). Taxa common in the crop in both seasons included Comamonadaceae,

Acinetobacter, Saprospiraceae, *Flavobacterium*, *Gilliamella*, *Bombella*, *Apilactobacillus*,

Pseudomonas, *Fluviicola*, *Polynucleobacter*, *Snodgrassella*, and *Microbacteriaceae*

species (Fig. 5). *Apilactobacillus* and *Bombella*, previously called *Parasaccharibacter*

apium [5, 40], which are among the most commonly reported taxa in the crop of

foraging adults, have also been found frequently in food stores, larvae, and queens [2,

4, 5, 27, 36, 41-43]. Although the data we present in this study are correlational,

preventing us from establishing causal relationships, we speculate that the crop may

serve as a consistent reservoir of these taxa that can influence the health of the adult

bees and the larvae that they tend [2, 27].

We also found evidence for potential strain-level specificity among *Apilactobacillus*

kunkeei isolates from collected foragers. One clade was made up of summer *A. cerana*

isolates from two different sites, and two others contained only *A. mellifera* isolates from

both seasons (Figs. 8 and S6). Although a larger sample size than we had in this study

as well as full genome sequencing are needed to ascertain the putative strain-level

differences, previous studies have shown differences in specific house-keeping genes

to suggest strain-level host specificity between *Apis* and *Bombus*, and even among *Apis*

species [44, 45]. As in these previous studies, some of our isolates clustered more

frequently by host species than by site [46, 47], potentially suggesting host specificity

and possibly specific host-symbiont relationships between *Apis* species and *A. kunkeei*

strains, even though this possibility remains speculative at this stage.

A few caveats should be considered in interpreting our findings. First, our analysis is

mostly based on relative abundance [46], even though we used qPCR to attempt to

estimate at least total absolute abundance in crop and mouth samples (Fig. 6b).

Absolute abundance of each taxon, which may be more relevant to bacterial effects on

the host [12, 46], can differ even when relative abundance is the same. However, we did

see similar bacterial composition patterns in our analysis of both relative and absolute

abundance (Figs. 4 and 6a). Second, we found that the crop did not consistently have a

higher bacterial load than the mouth (Fig. 6b). Although resident populations can be

persistent regardless of size, there may be more to the crop community than is

suggested by the compositional data. Third, we had only one site, Nishi-Honjo, where

bees were sampled in both summer and winter. A greater and more even number of

bees at each site in multi-year seasonal sampling would have made our results most

robust. Nevertheless, our analysis of several hundred bee and nectar samples all

sampled within the same landscape clearly refutes the hypothesis that crop

communities simply reflect what is available in the environment.

In conclusion, our results suggest that the composition of crop bacteria may be more

deterministic than generally assumed, questioning the notion that crop bacteria are

environmentally acquired transients that are unimportant to the bees. Although less well

studied than midgut and hindgut microbes, a small but increasing number of recent

studies indicate that some crop microbes do alter the health of individuals and hives [6,

48-50]. Our data reinforce the idea that it would help to study how microbial

communities are assembled and maintained not just in the midgut and the hindgut, but

also the crop, to better understand the relationship between honeybees and their gut

microbes.

References

1.   Kwong WK, Moran NA (2016) Gut microbial communities of social bees. Nat Rev

Microbiol 14: 374-384. doi: 10.1038/nrmicro.2016.43

2.   Anderson KE, Ricigliano VA, Copeland DC, Mott BM, Maes P (2023) Social

interaction is unnecessary for hindgut microbiome transmission in honey bees:

the effect of diet and social exposure on tissue-specific microbiome assembly.

Microb Ecol 85: 1498-1513. doi: 10.1007/s00248-022-02025-5

3.   Winston ML (1987) The biology of the honey bee. harvard university press

4.   Tamarit D, Ellegaard KM, Wikander J, Olofsson T, Vásquez A, Andersson SGE

(2015) Functionally structured genomes in *Lactobacillus kunkeei* colonizing the

honey crop and food products of honeybees and stingless bees. Genome Biol

Evol 7: 1455-1473. doi: 10.1093/gbe/evv079

5.   Corby-Harris V, Snyder LA, Schwan MR, Maes P, McFrederick QS, Anderson KE

(2014) Origin and effect of Alpha 2.2 Acetobacteraceae in honey bee larvae and

description of *Parasaccharibacter apium* gen. nov., sp. nov. Appl Environ

Microbiol 80: 7460-7472. doi: 10.1128/AEM.02043-14

6.   Arredondo D, Castelli L, Porrini MP, Garrido PM, Eguaras MJ, Zunino P, Antúnez

K (2018) *Lactobacillus kunkeei* strains decreased the infection by honey bee

pathogens *Paenibacillus larvae* and *Nosema ceranae*. Benef Microbes 9: 279-

290. doi: 10.3920/BM2017.0075

7.   Kwong WK, Mancenido AL, Moran NA (2017) Immune system stimulation by the

native gut microbiota of honey bees. Roy Soc Open Sci 4: 170003.

8.   Zheng H, Powell JE, Steele MI, Dietrich C, Moran NA (2017) Honeybee gut

microbiota promotes host weight gain via bacterial metabolism and hormonal

signaling. Proc Natl Acad Sci USA 114: 4775-4780. doi:

10.1073/pnas.1701819114

9.   Hart AG, Ratnieks FLW (2001) Why do honey-bee (*Apis millifera*) foragers

transfer nectar to several receivers? Information improvement through multiple

sampling in a biological system. Behav Ecol Sociobiol 49: 244-250. doi:

10.1007/s002650000306

10.   Decker LE, San Juan PA, Warren ML, Duckworth CE, Gao C, Fukami T (2022)

Higher variability in fungi compared to bacteria in the foraging honey bee gut.

Microb Ecol 85: 330-334. doi: 10.1007/s00248-021-01922-5

11.   Tiusanen M, Becker-Scarpitta A, Wirta H (2024) Distinct Communities and

Differing Dispersal Routes in Bacteria and Fungi of Honey Bees, Honey, and

Flowers. Microb Ecol 87: 100. doi: 10.1007/s00248-024-02413-z

12.   Kešnerová L, Emery O, Troilo M, Liberti J, Erkosar B, Engel P (2020) Gut

microbiota structure differs between honeybees in winter and summer. ISME J

14: 801-814. doi: 10.1038/s41396-019-0568-8

13.   Maeda T, Hiraiwa MK, Shimomura Y, Oe T (2023) Weather conditions affect

pollinator activity, fruit set rate, and yield in Japanese apricot. Sci Hortic 307:

111522. doi: 10.1016/j.scienta.2022.111522

14.   R Core Team (2023) R: a language and environment for statistical computing. In:

Computing, RffS (ed.), Vienna, Austria.

15.   McMurdie PJ, Holmes S (2013) phyloseq: An R package for reproducible

422   interactive analysis and graphics of microbiome census data. PLoS One 8:

423   e61217. doi: 10.1371/journal.pone.0061217

424   16.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin

425   PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H

426   (2019) vegan: Community ecology package. R package version 2.5-6.

427   17.   Xu S, Zhan L, Tang W, Wang Q, Dai Z, Zhou L, Feng T, Chen M, Wu T, Hu E

428   (2023) MicrobiotaProcess: A comprehensive R package for deep mining

429   microbiome. Innovation 4.

430   18.   Chen H (2022) VennDiagram: generate high-resolution Venn and Euler plots. R

431   package version 1.7.3.

432   19.   Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and

433   dispersion for RNA-seq data with DESeq2. Genome Biol 15: 550. doi:

434   10.1186/s13059-014-0550-8

435   20.   Gu Z, Eils R, Schlesner M (2016) Complex heatmaps reveal patterns and

436   correlations in multidimensional genomic data. Bioinformatics 32: 2847-2849. doi:

437   10.1093/bioinformatics/btw313

438   21.   Schliep K (2011) phangorn: phylogenetic analysis in R. Bioinformatics 27:

439   592e593.

440   22.   Hamady M, Lozupone C, Knight R (2010) Fast UniFrac: facilitating high-

441   throughput phylogenetic analyses of microbial communities including analysis of

442   pyrosequencing and PhyloChip data. ISME J 4: 17-27.

23.   Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects

models using lme4. J Stat Softw 67: 1-48. doi:

https://doi.org/10.18637/jss.v067.i01

24.   Lahti L, Shetty S (2017) microbiome R package.

25.   Chazdon RL, Chao A, Colwell RK, Lin S-Y, Norden N, Letcher SG, Clark DB,

Finegan B, Arroyo JP (2011) A novel statistical method for classifying habitat

generalists and specialists. Ecology 92: 1332-1343. doi:

https://doi.org/10.1890/10-1345.1

26.   Ogier J-C, Pagès S, Galan M, Barret M, Gaudriault S (2019) *rpoB*, a promising

marker for analyzing the diversity of bacterial communities by amplicon

sequencing. BMC Microbiol 19: 171. doi: 10.1186/s12866-019-1546-z

27.   Dyrhage K, Garcia-Montaner A, Tamarit D, Seeger C, Näslund K, Olofsson TC,

Vasquez A, Webster MT, Andersson SG (2022) Genome evolution of a symbiont

population for pathogen defense in honeybees. Genome Biol Evol 14: evac153.

28.   Katoh K, Standley DM (2013) MAFFT Multiple sequence alignment software

version 7: improvements in performance and usability. Mol Biol Evol 30: 772-780.

doi: 10.1093/molbev/mst010

29.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses.

Bioinformatics 25: 1972-1973.

30.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von

Haeseler A, Lanfear R (2020) IQ-TREE 2: new models and efficient methods for

phylogenetic inference in the genomic era. Mol Biol Evol 37: 1530-1534. doi:

10.1093/molbev/msaa015

31.   Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O (2011) Survey of

branch support methods demonstrates accuracy, power, and robustness of fast

likelihood-based approximation schemes. Syst Biol 60: 685-699. doi:

10.1093/sysbio/syr041

32.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2017) UFBoot2:

improving the ultrafast bootstrap approximation. Mol Biol Evol 35: 518-522. doi:

10.1093/molbev/msx281

33.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017)

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat

Methods 14: 587-589. doi: 10.1038/nmeth.4285

34.   Mai U, Mirarab S (2018) TreeShrink: fast and accurate detection of outlier long

branches in collections of phylogenetic trees. BMC Genomics 19: 23-40.

35.   Anderson KE, Maes P (2022) Social microbiota and social gland gene

expression of worker honey bees by age and climate. Sci Rep 12: 10690. doi:

10.1038/s41598-022-14442-0

36.   Anderson KE, Sheehan TH, Mott BM, Maes P, Snyder L, Schwan MR, Walton A,

Jones BM, Corby-Harris V (2013) Microbial ecology of the hive and pollination

landscape: bacterial associates from floral nectar, the alimentary tract and stored

food of honey bees (*Apis mellifera*). PLoS One 8: e83125.

37.   Alvarez-Perez S, Baker LJ, Morris MM, Tsuji K, Sanchez VA, Fukami T, Vannette

RL, Lievens B, Hendry TA (2021) *Acinetobacter pollinis* sp. nov., *Acinetobacter*

488   *baretiae* sp. nov. and *Acinetobacter rathckeae* sp. nov., isolated from floral nectar

489   and honey bees. Int J Syst Evol Microbiol 71: 004783.

490 38. Fridman S, Izhaki I, Gerchman Y, Halpern M (2012) Bacterial communities in

491   floral nectar. Environ Microbiol Rep 4: 97-104. doi: https://doi.org/10.1111/j.1758-

492   2229.2011.00309.x

493 39. Morris MM, Frixione NJ, Burkert AC, Dinsdale EA, Vannette RL (2020) Microbial

494   abundance, composition, and function in nectar are shaped by flower visitor

495   identity. FEMS Microbiol Ecol 96: fiaa003. doi: 10.1093/femsec/fiaa003

496 40. Smith EA, Anderson KE, Corby-Harris V, McFrederick QS, Parish AJ, Rice DW,

497   Newton ILG (2021) Reclassification of seven honey bee symbiont strains as

498   *Bombella apis*. Int J Syst Evol Microbiol 71: 004950. doi: 10.1099/ijsem.0.004950

499 41. Vásquez A, Forsgren E, Fries I, Paxton RJ, Flaberg E, Szekely L, Olofsson TC

500   (2012) Symbionts as major modulators of insect health: lactic acid bacteria and

501   honeybees. PLoS One 7: e33188. doi: 10.1371/journal.pone.0033188

502 42. Corby-Harris V, Maes P, Anderson KE (2014) The bacterial communities

503   associated with honey bee (*Apis mellifera*) foragers. PLoS One 9: e95056.

504 43. Rangberg A, Mathiesen G, Amdam GV, Diep DB (2015) The paratransgenic

505   potential of *Lactobacillus kunkeei* in the honey bee *Apis mellifera*. Benef

506   Microbes 6: 513-523. doi: 10.3920/BM2014.0115

507 44. Kwong WK, Engel P, Koch H, Moran NA (2014) Genomics and host

508   specialization of honey bee and bumble bee gut symbionts. Proc Natl Acad Sci

509   USA 111: 11509-11514. doi: 10.1073/pnas.1405838111

45. Kwong WK, Medina LA, Koch H, Sing K-W, Soh EJY, Ascher JS, Jaffé R, Moran

NA (2017) Dynamic microbiome evolution in social bees. Sci Adv 3: e1600513.

doi: doi:10.1126/sciadv.1600513

46. Almeida EL, Ribiere C, Frei W, Kenny D, Coffey MF, O'Toole PW (2023)

Geographical and seasonal analysis of the honeybee microbiome. Microb Ecol

85: 765-778. doi: 10.1007/s00248-022-01986-x

47. Ellegaard KM, Suenami S, Miyazaki R, Engel P (2020) Vast differences in strain-

level diversity in the gut microbiota of two closely related honey bee species.

Curr Biol 30: 2520-2531.e2527. doi: 10.1016/j.cub.2020.04.070

48. Chege M, Kinyua J, Paredes JC (2023) *Lactobacillus kunkeei* impacts the health

of honey bees, *Apis mellifera* scutellata, and protects the bees against the

opportunistic pathogen *Serratia marcescens*. Int J Trop Insect Sci 43: 1947-1955.

doi: 10.1007/s42690-023-01103-6

49. Corby-Harris V, Snyder L, Meador CAD, Naldo R, Mott B, Anderson KE (2016)

Parasaccharibacter apium, gen. nov., sp. nov., improves honey bee

(Hymenoptera: Apidae) resistance to *Nosema*. J Econ Entomol 109: 537-543.

doi: 10.1093/jee/tow012

50. Miller DL, Smith EA, Newton ILG (2021) A bacterial symbiont protects honey

bees from fungal disease. mBio 12: e00503-00521. doi: 10.1128/mBio.00503-21

Funding

This work was supported by the Office of International Affairs and the Freeman Spogli

Institute for International Studies Japan Fund of Stanford University.

Competing Interests

The authors declare no competing interests.

Author Contributions

KT, MK, and TF designed the study and collected the samples. MLW did the DNA

extraction, sequencing preparation, and bioinformatic analyses. JY and ACH did the

quantitative PCR processing and analyses. MLW and LED did the statistical analyses.

MLW wrote the first draft of the manuscript. All authors contributed to and approved the

manuscript.

Data Availability

The datasets generated during and/or analyzed during the current study are available in

GenBank, https://www.ncbi.nlm.nih.gov (BioProject ID no. PRJNA1076572 and

accession no. PP412473-PP412519, PP341993-PP342038, PP382363-PP382409,

PP382266-PP382314, PP382315-PP382362), or are included in this published article

and its supplementary information files. All custom scripts are available on GitHub at

https://github.com/mlwarren20/ume_bee_analysis.