Synthesis and Biological Assessment of Chalcone and Pyrazoline Derivatives as Novel Inhibitor for ELF3-MED23 Interaction

Soo-Yeon Hwang,[1,#] Kyung-Hwa Jeon,[1,#] Hwa-Jong Lee,[1] Inhye Moon,[1] Sehyun Jung,[1] Seul-Ah Kim,[1] Hyunji Jo,[1] Seojeong Park,[1] Misun Ahn,[1] Soo-Yeon Kwak,[2] Younghwa Na[2,*] and Youngjoo Kwon,[1,*]

[1]*College of Pharmacy & Graduate School of Pharmaceutical Sciences, Ewha Womans University, Seoul, 03760, Korea*

[2]*College of Pharmacy, CHA University, Pocheon, 487-010, Korea*

*To whom correspondence should be addressed. Y. Na: Phone: 82-31-8017-9896. Fax: 82-31-8017-9420. E-mail: yna7315@cha.ac.kr, Y. Kwon: Phone: 82-2-3277-4653. Fax: 82-2-3277-3051. E-mail: ykwon@ewha.ac.kr.

#These authors are equal contributors

Abstract

HER2 overexpression significantly contributes to the aggressive nature and recurrent patterns observed in various solid tumors, notably gastric cancers. Trastuzumab, HER2-targeting monoclonal antibody drug, has shown considerable clinical success, however, readily emerging drug resistance emphasizes the pressing need for improved interventions in HER2-overexpressing cancers. To address this, we proposed targeting the protein-protein interaction (PPI) between ELF3 and MED23 as an alternative therapeutic approach to trastuzumab. In this study, we synthesized a total of 26 compounds consisting of 10 chalcones, 7 pyrazoline acetyl, and 9 pyrazoline propionyl derivatives, and evaluated their biological activity as potential ELF3-MED23 PPI inhibitors. Upon systematic analysis, candidate compound **10** was selected due to its potency in downregulating SEAP activity and its effect on HER2 protein and mRNA levels. Compound **10** effectively disrupted the binding interface between the ELF3 TAD domain and the 391-582 amino acid region of MED23, resulting in successful inhibition of the ELF3-MED23 PPI. This intervention led to a substantial reduction in HER2 levels and its downstream signals in the HER2-positive gastric cancer cell line. Subsequently, compound **10** induced significant apoptosis and anti-proliferative effects, demonstrating superior *in vitro* and *in vivo* anticancer activity overall. We found that the anticancer activity of compound **10** was not only restricted to trastuzumab-sensitive cases, but was also valid for trastuzumab-refractory clones. This suggests its potential as a viable therapeutic option for trastuzumab-resistant gastric cancers. In summary, compound **10** could be a novel alternative therapeutic strategy for HER2-overexpressing cancers, overcoming the limitations of trastuzumab.

Keywords HER2-overexpressing cancers; Gastric cancer; Trastuzumab-resistance; Protein-protein interaction inhibitor; 4-Methoxy-3-((3-methylbut-2-en-1-yl)oxy)benzaldehyde

1. Introduction

Human epidermal growth factor receptor 2 (HER2) belongs to the HER family, encompassing EGFR (HER1, erbB1), HER2 (HER2/neu, erbB2), HER3 (erbB3), and HER4 (erbB4). Known as a proto-oncogene, HER2 plays a pivotal role in various cancer-related processes, including cell migration, proliferation, differentiation, and adhesion [1-3]. Its activation occurs through homo- or hetero-dimerization with other members of the HER family, triggering downstream cellular signaling pathways like Ras-Raf-MAPK and PI3K-AKT-mTOR [2, 4]. Distinct from other HER family proteins, HER2 does not necessitate ligand-binding for dimer formation or downstream signal activation. Moreover, the structure of the HER2 extracellular domain adopts an open conformation, making it more prone to undergo homo- or heterodimerization. This predisposition of HER2 as a dimerization partner, coupled with its tendency for overexpression, contributes to uncontrolled cell proliferation and the progression of tumors.

Numerous studies have consistently identified HER2 overexpression across diverse cancer types including breast, colorectal, ovarian, pancreatic, prostate, bladder, lung, gastric, and gastroesophageal cancers [5]. The frequency of HER2 overexpression varies: approximately 20% in breast cancers [6], 5–30% in ovarian cancers [7], 7–8% in colorectal cancers [8], 0–82% in pancreatic cancers [9], and 7.3–20.2% in gastric and gastroesophageal cancers [10]. In these cases, HER2 status serves as an important prognostic marker, correlating positively with tumor size, invasiveness, and lymph node metastasis [2, 6, 8, 10, 11]. The significance of HER2 is evident in the development of separate treatment guidelines tailored specifically for HER2-positive cases in several cancer subtypes such as breast, colorectal, and gastric cancers. This often involves employing trastuzumab, a humanized monoclonal antibody drug targeting the extracellular domain of HER2 protein [12-14].

Trastuzumab has significantly improved the clinical outcomes of numerous patients with

HER2-positive cancers, but its therapeutic efficacy is considered limited due to readily

developing resistance [15]. The emergence of this persistent refractoriness to trastuzumab has

highlighted the need for the establishment of better therapeutic approaches that maintain

effectiveness regardless of the development of resistance. While HER2 gene amplification is

the primary mechanism responsible for HER2 overexpression, high transcription rates of

HER2 per gene copy have also been observed to contribute [16]. In breast cancer, it has been

reported that radiation or endocrine therapy upregulates HER2 gene transcription in HER2-

low or HER2-negative subtypes, leading to treatment resistance [16-18]. Rather than

targeting already overexpressed HER2 protein, our focus has shifted to reducing HER2

expression at the genetic level. This involves interrupting the interaction between E74-Like

Factor 3 (ELF3) and mediator of RNA polymerase II transcription subunit 23 (MED23) – a

transcription factor (TF) and a coactivator for HER2, respectively. As an essential TF for

HER2 gene expression, ELF3 directly binds to the ETS transcriptional response element

located in the HER2 promoter and interacts with MED23 to drive HER2 overexpression [19].

Natural and synthetic chalcones have been reported to show a variety of biological efficiency,

including anti-proliferative, anticancer, antioxidant, anti-inflammatory, or anti-infective

activities [20-22]. Chalcone derivatives have shown potential as lead compounds in the field

of novel drug discovery due to their promising biological activities and have also been

sourced as precursors of flavonoids in higher plants. Licochalcone derivatives, initially

identified in *licorice* root, possess alkyl-substituted forms of the typical chalcone structure

[23], exhibiting various biological activities such as anticancer, antioxidant, anti-

inflammatory activities [24-26]. Pyrazole, an organic heterocyclic compound, is characterized

by a 5-membered ring comprising three carbon atoms and two adjacent nitrogen atoms [27].

Pyrazoline, an unsaturated pyrazole derivative, contains a single endocyclic double bond on

the N-2 position, distinguishing it from pyrazole. Numerous synthetic pyrazoline analogues

have demonstrated notable anticancer activities [28].

Chalcone   **Pyrazoline**   **Licochalcone E**   **YK-1**

In our previous study, pyrazoline analogue YK-1 was developed as an effective inhibitor

targeting the interaction between ELF3 and MED23. YK-1 exhibited remarkable efficacy,

demonstrating significant *in vitro* and *in vivo* anticancer activity specifically against HER2-

overexpressing breast and gastric cancers [29]. As part of our ongoing efforts to discover a

potent inhibitor of the ELF3-MED23 PPI for potential anticancer therapies, we designed and

synthesized 26 novel compounds, including chalcones, pyrazoline acetyl, and pyrazoline

propionyl analogs **(Figure 1)**. This exploration led us to elicit a highly potent compound **10**

that exhibited excellent *in vitro* and *in vivo* anticancer activity against HER2-positive gastric

cancer cells.

Chalcone   Pyrazoline   Licochalcone E

YK-1   Proposed compounds

109   **Figure 1.** Structure of proposed compounds.

2. Results

2.1. Identification of compound 10 as potent ELF3-MED23 PPI inhibitor

We designed and synthesized 26 novel compounds, including chalcones, pyrazoline acetyl, and pyrazoline propionyl analogs based on the YK1 structure (**Figure 2**). To screen out compounds exhibiting superior inhibitory activity against ELF3-MED23 PPI, we first performed secreted alkaline phosphatase (SEAP) assay [29, 30] using 26 synthesized compounds. As in our previous studies, gefitinib and CI-1033 (Canertinib) were used as positive controls for the experiment [29, 30] (**Figure 3A**). Cell viability tests were conducted in parallel with the SEAP assay under the same experimental condition, in order to confirm that the decrease in fluorescence observed in the SEAP assay was specifically due to ELF3-MED23 disruption, and not related to non-specific cytotoxicity of the compounds (**Figure 3B**). Overall, a total of 12 compounds displayed nearly 100% inhibition at 10 μM, showing the most excellent significant SEAP inhibitory activities among the entire series. Most of these have a chalcone moiety (compounds **1-10** in **group 1**), while compounds **16** and **17** have a pyrazoline acetyl moiety. Among the remaining pyrazoline acetyl derivatives in **group 2**, compounds **11**, **13**, **14**, and **15** also showed inhibitory effects, but their extent was relatively low (compounds **11, 14,** and **15**; below 30% inhibition, compound **13**; 68% inhibition). In case of the molecules of pyrazoline propionyl skeleton in **group 3**, only compound **26** displayed a moderate inhibitory rate of 63.8%, while majority of them did not show noticeable inhibitory effects (**Figure 3A**). To select the compounds that can actually induce downregulation of HER2 in cancer cells, we applied the top 12 compounds with the highest SEAP inhibitory activity to NCI-N87, a well-known HER2-positive gastric cancer cell line, and evaluated changes in the HER2 protein expression level (**Figure 3C**). Of the tested molecules, only compounds **3**, **5**, and **10** were found to significantly reduce the HER2

levels by more than 50%. However, unlike compound **3** and **10**, the downregulation of HER2

induced by compound **5** did not occur at the mRNA level. This indicates that only compounds

3 and **10** were capable of transcriptionally downregulating HER2 by disrupting the ELF3-

MED23 interaction (**Figure 3D**). Since compound **10** remarkably decreased the HER2 level

of NCI-N87 by more effectively disrupting the ELF3-MED23 PPI compared to compound **3**

(**Figure 3E**), we finally selected compound **10** as the candidate molecule and performed

further experiments.

Group 1

1 $R_1 = CH_3$, $R_2 = H$
2 $R_1 = CH_3$, $R_2 = $ Propenyl
3 $R_1 = CH_3$, $R_2 = $ 3-Butenyl
4 $R_1 = CH_3$, $R_2 = $ 2-Butenyl
5 $R_1 = CH_3$, $R_2 = $ 3-Methyl-but-2-enyl
6 $R_1 = H$, $R_2 = CH_3$
7 $R_1 = $ Propenyl , $R_2 = CH_3$
8 $R_1 = $ 3-Butenyl, $R_2 = CH_3$
9 $R_1 = $ 2-Butenyl, $R_2 = CH_3$
10 $R_1 = $ 3-Methyl-but-2-enyl, $R_2 = CH_3$

Group 2

11 $R_1 = CH_3$, $R_2 = H$
12 $R_1 = CH_3$, $R_2 = $ Propenyl
13 $R_1 = CH_3$, $R_2 = $ 3-Butenyl
14 $R_1 = CH_3$, $R_2 = $ 2-Butenyl
15 $R_1 = H$, $R_2 = CH_3$
16 $R_1 = $ Propenyl , $R_2 = CH_3$
17 $R_1 = $ 2-Butenyl, $R_2 = CH_3$

Group 3

18 $R_1 = CH_3$, $R_2 = H$
19 $R_1 = CH_3$, $R_2 = $ Propenyl
20 $R_1 = CH_3$, $R_2 = $ 3-Butenyl
21 $R_1 = CH_3$, $R_2 = $ 2-Butenyl
22 $R_1 = CH_3$, $R_2 = $ 3-Methyl-but-2-enyl
23 $R_1 = H$, $R_2 = CH_3$
24 $R_1 = $ Propenyl , $R_2 = CH_3$
25 $R_1 = $ 2-Butenyl, $R_2 = CH_3$
26 $R_1 = $ 3-Methyl-but-2-enyl, $R_2 = CH_3$

144   **Figure 2. Structures of the prepared compounds.**

Figure 3. Identification of compound 10 as potent ELF3-MED23 PPI inhibitor. **(A)** Synthesized compounds were screened to evaluate their inhibitory activity against ELF3–MED23 PPI. All compounds were used at 10 μM for 12 h ($n = 3$, mean ± S.D.). **(B)** Cell viability was also measured in parallel using the same conditions as for the reporter gene assay in (A) ($n = 3$, mean ± S.D.). **(C)** Changes in the HER2 levels were evaluated by treating NCI-N87 cells with the compounds that exhibited high SEAP inhibitory activity in (A). **(D)** Effect of compound **3**, **5**, and **10** on mRNA level of HER2 in NCI-N87 was assessed (16 h treatment at 10 μM, $n = 3$, mean ± S.D., ANOVA, ***$p < 0.001$ *vs*. CON). **(E)** PPI inhibitory activities of compounds **3** and **10** were evaluated against endogenous ELF3 and MED23.Compound **25** was used as negative control.

2.2. Qualitative structure-activity relationship (SAR) of prepared compounds

Field-based approaches primarily focus on characterizing the molecular properties of

compounds based on their interaction fields. By providing comprehensive 3D information

regarding charge distribution, non-polar interactions, and shape properties of diverse

molecules, this method allows detailed insights into how distinct electrostatic, hydrophobic,

and steric fields of ligands contribute to their biological activity or inactivity [31, 32]. Based

on this concept, we utilized FieldTemplater™ software to 3-dimensionally align all 26

compounds according to their field points. We subsequently analyzed their configurations

through Activity Atlas™ software [33]. We finally visualized the results in 3D activity cliff

summary plots of positive/negative electrostatics and favorable/unfavorable hydrophobics

(**Figure 4**). According to the electrostatic and hydrophobic plots, the positive/negative and

hydrophobic field points of the compounds demonstrating relatively high HER2 inhibitory

activity (as exemplified by compounds **3** and **5**) were found to be well-distributed within the

favored areas. Especially, the negative field formed around the ketone moiety within the

chalcone skeleton, and the hydrophobic field generated by 3-methyl-but-2-enyl side chain of

compound **10** are assumed to be important for the HER2 inhibitory effect induced by

interrupting ELF3-MED23 PPI. This assumption arises from the observation that the

overall activity slightly decreased when both the negative and hydrophobic field point were

less distributed in the mentioned regions (**Figure 4A and 4B**; compounds **3** and **5**). For the

compounds displaying low potency (represented by compounds **12**, **15**, **20** and **24**), their

positive/negative electrostatic and hydrophobic potentials shared unfavorable distribution

patterns. This suggests that conformation of these compounds is less likely to

complementarily fit into the binding interface of ELF3-MED23 PPI (**Figure 4A and 4B**;

compounds **12**, **15**, **20**, and **24**).

Figure 4. Qualitative SAR of prepared compounds. Activity cliff summary for the prepared compound series were plotted regarding the contributions of electrostatic **(A)** and hydrophobic **(B)** fields. The field point pattern was depicted as spheres (Red, positive electrostatics; Cyan, negative electrostatics; Yellow brown, hydrophobic fields).

2.3. Compound 10 as a transcriptional regulator of HER2 by inhibiting ELF3-MED23 PPI

Previously, we have verified that the TAD domain of ELF3 (129–145 residues; 17 amino acids) interacts with the 391–582 residues of MED23, thereby regulating HER2 at the transcriptional level. Specifically, within ELF3, residues S137 to E144 were identified as essential residues for this interaction [29]. Based upon the results, we performed *in vitro* fluorescence polarization (FP) assay by utilizing the $(His)_6$-$MED23_{391-582}$ protein and fluorescein isothiocyanate (FITC)-labeled $ELF3_{129-145}$ peptide (**Figure 4A**). Similar to the positive control, unlabeled $ELF3_{137-144}$ peptide, compound **10** significantly reduced the FP signal induced by the binding of FITC-$ELF3_{129-145}$ and $(His)_6$-$MED23_{391-582}$, demonstrating that compound **10** directly inhibits the ELF3-MED23 PPI. While the K_i value of unlabeled $ELF3_{137-144}$ peptide calculated from the K_d of FITC-$ELF3_{129-145}$ and $(His)_6$-$MED23_{391-582}$ interaction was determined to be 4.47±0.03 μM, compound **10** showed a 6.6-fold lower value of 0.68±0.08 μM, falling within the nanomolar range (**Figure S1 and Figure 5A**). Through GST-pull down assay, we then confirmed that ELF3-MED23 PPI inhibitory effect of compound **10** is also valid at the cell level (**Figure 5B**). To investigate whether intracellular compound **10** also acts on the binding interface of ELF3 and the 391-582 amino acid region of MED23 within the cell system, we performed split luciferase complementation assay using ELF3- and $MED23_{391-582}$-ligated with N-terminal (Nluc) and C-terminal (Cluc) fragments of split luciferase, respectively (**Figure 5C**). Without compound **10**, ELF3 and $MED23_{391-582}$ successfully interacted within the cell system, subsequently inducing the complementation of Nluc and Cluc to generate a significant amount of bioluminescence (32.7-fold change vs.

control). However, compound **10** markedly reduced this luciferase activity in a dose-

dependent manner. This result verified that compound **10** is also capable of disrupting the

ELF3-MED23 PPI at the cellular level. In fact, the successful inhibition of the ELF3-MED23

PPI by compound **10** ultimately resulted in the attenuation of HER2 promoter activity

(**Figure 5D**). This led to a significant downregulation of HER2 expression and its

downstream signaling molecules such as pAKT and pMAPK in a concentration- and time-

dependent manner (**Figure 5E**).

Figure 5. Compound 10 as a transcriptional regulator of HER2 by inhibiting ELF3-MED23 PPI.
(A) Effect of compound **10** on the FP (mP) induced by the binding of FITC-ELF3$_{129-145}$ peptide to (His)$_6$-MED23$_{391-582}$ protein was evaluated in cell-free system. Unlabeled ELF3$_{137-144}$ peptide was used as positive control. IC$_{50}$ and K$_i$ values were calculated from the FP assay results ($n = 3$, mean ± S.D.). **(B)** Intracellular PPI inhibitory effect of compound **10** (5 μM, 12 h treatment) against ELF3-MED23 was evaluated through GST-pull down assay using GST-ELF3$_{WT}$ and 3xFLAG-MED23. **(C)** Impact of compound **10** on the relative luciferase activity generated by Nluc-ELF3$_{WT}$ and Cluc-MED23$_{391-582}$ interaction was evaluated (20 h treatment at indicated concentrations, $n = 3$, mean ± S.D., ANOVA, ***$p < 0.001$). **(D)** Effect of compound **10** on the overall HER2 promoter activity was assessed (20 h treatment at indicated concentrations, $n = 3$, mean ± S.D., ANOVA, *$p < 0.05$, **$p < 0.01$, and ***$p <$

226   0.001). **(E)** Changes in the HER2 and its downstream signaling pathway were evaluated by treating
227   compound **10** in dose- and time-dependent manner.

2.4. Compound 10 as a potent anticancer agent for HER2-positive gastric cancer cells

Based upon the obtained results so far, we proceeded to evaluate whether compound **10**-

mediated inhibition of ELF3-MED23 PPI could lead to significant anticancer activity against

HER2-positive gastric cancer cells in both *in vitro* and *in vivo* settings. First, by treating NCI-

N87 cells with varying concentrations of compound **10**, we found that apoptosis was

significantly induced in a concentration-dependent manner (**Figure 6A**). This effect was

further validated by the dose- and time-dependent increases in pro-apoptotic markers, cleaved

PARP (c-PARP) and cleaved caspase 3 (c-Caspase 3) (**Figure 6B**). Accordingly, compound

10 significantly decreased the cell proliferation of NCI-N87 in a concentration-dependent

manner (**Figure 6C**), indicating its potent anticancer activity in the *in vitro* system. To

evaluate its potency in the *in vivo* setting, we then established an NCI-N87 xenograft mouse

model and administered compound **10** intravenously. Notably, the administration of 4 mg/kg

of compound **10** resulted in significant retardation of tumor growth (**Figure 6D**), leading to a

marked reduction in the final tumor volume, approximately 3.5-fold decrease compared to the

control group (**Figure 6E and 6F**). Through Immunohistochemistry (IHC) analysis, we

confirmed that the proliferation marker, Ki67, was remarkably downregulated (a 2.3-fold

decrease compared to the control) in the tumor tissues administered with compound **10**

(**Figure 6G, right panel**). This reduction in Ki67 expression was accompanied by a

significant decrease in HER2 levels in the tumor tissue (a 1.7-fold decrease compared to the

control), suggesting that the overall anti-tumor activity of compound **10** is primarily due to its

capacity to inhibit HER2 expression (**Figure 6G, left panel**).

250
251

Figure 6. Compound 10 as a potent anticancer agent for HER2-positive gastric cancer cells. (A) Apoptosis induced by compound **10** in parental NCI-N87 was assessed by treatment with the compound in a dose-dependent manner (24 h treatment at indicated concentrations, $n = 3$, mean ± S.D., ANOVA, ns = non-significant, ***$p < 0.001$ vs. CON). **(B)** Changes in the pro-apoptotic markers were evaluated by treating compound **10** in dose- and time-dependent manner. **(C)** Anti-proliferative effect of compound **10** (10 µM) was evaluated against NCI-N87 cell line (10 days of

treatment at indicated concentrations, $n = 3$, mean $\pm$ S.D., ANOVA, ***$p < 0.001$ vs. CON). **(D)**
Tumor growth inhibitory effect of compound **10** was evaluated using NCI-N87 xenograft mouse
model ($n = 5$ per group; intravenous (IV) injection at 4 mg/kg every 3 days). Tumor volumes were
evaluated at the indicated time points by measuring the length and width of the tumor with calipers
using the equation (length x width^2)/2. Data was indicated as mean $\pm$ S.E.M. **(E)** Photograph of the
tumors collected from the vehicle- and compound **10**-treated mice. **(F)** Final volume changes were
assessed for the tumors excised from each experimental group ($n = 5$, mean $\pm$ S.E.M., Student's t-test,
***$p < 0.001$ vs. CON) **(G)** IHC analysis was conducted against HER2 and Ki67 in the tumors. Score
quantification was performed using Image J software (10 independent fields per sample were
evaluated, mean $\pm$ S.D., Student's t test, ***$p < 0.001$ vs. CON).

2.5. Compound 10 as a novel strategy to overcome trastuzumab resistance

As previously mentioned, trastuzumab has been a primary first-line therapy for various

HER2-overexpressing cancer subtypes, including gastric cancer [34, 35]. However, one of the

significant drawback of this drug is the development of resistance, which readily occurs

within 1 year of medication [36]. In our previous study, we demonstrated that inhibiting the

ELF3-MED23 PPI to downregulate HER2 at the transcriptional level could serve as a

potentially promising alternative therapeutic option to trastuzumab [29]. Consequently, we

sought to determine whether compound **10** holds the capability to overcome trastuzumab

resistance. For this assessment, we utilized the trastuzumab-refractory NCI-N87 (NCI-N87

TR) cell line, which exhibited a 5.3-fold lower response rate to 10 μg/mL trastuzumab

compared to the parental NCI-N87 cells (Parent *vs.* TR; 50.1% *vs.* 9.5% growth inhibition)

(**Figure 7A**). At a concentration of 10 μg/mL, trastuzumab exhibited no discernible impact on

the level of HER2 or its downstream signaling molecules such as p-AKT and p-MAPK in

NCI-N87 TR cells. In contrast, compound **10** effectively attenuated these pathways by

inhibiting HER2 expression (**Figure 7B**). Accordingly, compound **10** markedly inhibited the

long-term proliferation of NCI-N87 TR, unlike trastuzumab (**Figure 7C**). Moreover, the

dose-dependent treatment of compound **10** significantly induced apoptosis in NCI-N87 TR

cells (**Figure 7D**). These findings collectively suggest that compound **10** demonstrates

anticancer activity not only against parental HER2-positive cancer cells but also against their

trastuzumab-refractory clones.

Figure 7. Compound 10 as a novel strategy to overcome trastuzumab resistance. (A)
Trastuzumab (10 μg/ml, 24 h)-induced growth inhibitory effects were assessed against parental and
TR NCI-N87 cell lines ($n = 3$, mean ± S.D., Student's t-test, ***$p < 0.001$ *vs.* parent). **(B)**
Trastuzumab- and compound **10**-mediated changes of HER2 and its downstream signaling pathway
were evaluated (16 h treatment at indicated concentrations). **(C)** Anti-proliferative effects of
trastuzumab (10 μg/mL) and compound **10** (10 μM) were evaluated against NCI-N87 TR cell line.
Each compound was treated for 10 days ($n = 3$, mean ± S.D., ANOVA, ns = non-significant, *$p < 0.05$
vs. CON). **(D)** Apoptosis induced by compound **10** in NCI-N87 TR was assessed by FACS analysis
with treatment of the compound in a dose-dependent manner (24 h treatment at indicated
concentrations, $n = 3$, mean ± S.D., ANOVA, ns = non-significant, ***$p < 0.001$ *vs.* CON).

3. Discussion

As HER2-positivity is generally responsible for more aggressive behavior of diverse cancer subtypes, HER2 has long been regarded as a crucial biomarker and a primary therapeutic target. Consequently, numerous researchers have consistently sought to develop new HER2-targeting agents beyond trastuzumab, and indeed, several drugs such as Enhertu® (fam-trastuzumab deruxtecan-nxki) and Tukysa® (tucatinib) have been identified and approved by the FDA in recent years [37, 38].

To date, numerous HER2-targeting drugs are clinically available. Among them, trastuzumab, the first FDA-approved HER2-targeting drug, still remains the most commonly used drug of molecular targeted therapy for HER2-positive breast cancer, improving the survival of patients with this molecular subtype [39, 40]. However, despite initial responses to trastuzumab or trastuzumab conjugate regimens, approximately 70% of HER2-positivie breast cancer patients develop resistance within the first year of treatment [39-41]. This high incidence of trastuzumab resistance highlights the persistent challenges of relapse and drug resistance in the treatment of HER2-positive breast cancer.

As HER2 targeting primarily involves the already overexpressed HER2 and its downstream signaling pathways, typically driven by the gene amplification [42], alternative approaches have been explored to downregulate HER2 signaling from the gene expression level. Specifically, strategies have been investigated to inhibit the interaction between transcription factor (TF), ELF3, and its coactivator, MED23 [29, 30, 43-45]. ELF3, a critical regulator of HER2 gene expression, acts on the ETS transcriptional response element of HER2 promoter and interacts with MED23 to promote HER2 overexpression [19, 46]. Our previous work has provided structural insights into the regulation of the ELF3-MED23 PPI [29]. Expanding on the findings that interrupting the ELF3-MED23 PPI can effectively serve as a viable

alternative strategy for inhibiting HER2 compared to trastuzumab, we designed a series of potential ELF3-MED23 inhibitors incorporating chalcone and pyrazoline moieties.

This study primarily aims to identify novel small molecules that can inhibit HER2 with outstanding anticancer activity, which can be broadly applicable to diverse HER2-positive cancers, irrespective of their resistance to trastuzumab. Following a comprehensive biological evaluation, we successfully identified compound **10,** demonstrating a highly potent ELF3-MED23 inhibitory efficacy ($K_i = 0.68\pm0.08$ μM). Based on these findings, we confirmed that this compound exhibits excellent anticancer activity in both *in vitro* and *in vivo* settings against HER2-overexpressing gastric cancer cells and those exhibiting resistance to trastuzumab.

This study highlights that the chalcone moiety can efficiently bind to the ELF3-MED23 interface, suggesting that this structure can potentially serve as a useful scaffold in the development of PPI inhibitors, specifically for downregulating HER2 and its associated signaling cascade. The results also clearly demonstrates that inhibition of ELF3-MED23 PPI through the chalcone derivative can serve as a potential therapeutic strategy targeting HER2 in anticancer treatments. By combining our findings on the efficacy of chalcone derivatives with our understanding of the complexity of the ELF3-MED23 interaction and its role in HER2 regulation, this study makes a valuable contribution, paving the way for designing targeted therapies. Consequently, we anticipate that these insights will provide valuable contributions towards development of new agents, the progression of treatments, and the enhancement of the therapeutic landscape for HER2-positive cancers.

4. Materials and methods

4.1 Chemistry

The synthesized compounds were classified into three groups: **group 1** encompassing

chalcone compounds, **group 2** comprising N-1-acetyl-4,5-dihydropyrazoline compounds, and

group 3 consisting of *N*-1-propionyl-4,5-dihydropyrazoline compounds (**Scheme 1** and

Figure 2). **Group 1** chalcone compounds were synthesized through a modified *Claisen-*

Schmidt condensation reaction using 3,4,5-trimethoxyacetohenone and the appropriate

aromatic aldehyde under 50% NaOH base condition, yielding between 47.8% to 88.4%.

Analysis through [1]H-NMR spectroscopy confirmed the trans conformation of chalcone

compounds, evidenced by two doublet peaks with coupling constants around 15.6 Hz,

representing α and β hydrogens in the conjugated ketone structure. **Group 2** acetyl-

pyrazoline compounds were obtained by reacting **group 1** chalcone compounds with

hydrazine and acetic acid, yielding between 50.8% to 82.4%. [1]H-NMR spectra of **group 2**

compounds displayed characteristic peaks of three double of doublets at 3.1, 3.7, and 5.5 ppm,

indicating each hydrogen at *C*-4 and *C*-5 positions in the pyrazoline ring structure.

Compounds **11** - **17** also showed a methyl peak as a singlet at about 2.4 ppm, corresponding

to the acetyl group of *N*-1 position. Interestingly, during this pyrazoline ring construction

process, chalcone compounds **5** and **10** yielded compounds **11** and **16** with the *O*-substituted

3-methyl-but-2-enyl moiety deleted instead of the desired acetyl compounds. **Group 3**

propionyl-pyrazoline compounds were also prepared in the same manner as **group 2**

compounds with yields of 4.9% to 79.7%, except for propionic acid conditions. [1]H-NMR

spectra confirmed the presence of a propionyl substituent at the *N*-1 position of the pyrazoline

ring, displaying a set of triplet and double of quartet peaks at about 1.2 and 2.7 ppm,

respectively, corresponding to the ethyl peak of the propionyl group. All synthesized

compounds are consistent with the spectral data.

Scheme 1. General synthesis methods for the target compounds.

Chemicals and reagents used were obtained from Aldrich Chemical Co. and others were from company like TCI. Melting points were measured without correction in open capillaries with Barnstead Electrothermal melting point apparatus, Manual MELTEMP (Model No: 1202D). Chromatographic separations were monitored by thin-layer chromatography using a commercially available pre-coated Merck Kieselgel 60 F254 plate (0.25 mm) and detected by visualizing under UV at 254 and 365 nm. Silicagel column chromatography was carried out with Merck Kieselgel 60 (0.040-0.063 mm). All solvents used for chromatography were directly used without distillation. The purity was assessed by HPLC (Shimadzu LC-20AD) analysis under the following conditions; column, SunFire C18 (4.6 mm × 150 mm, 5 mm); mobile phase, condition : A (water) and B (acetonitrile) using a linear gradient of 50-70% B in 0-15 min, 70% B in 15-20 min,100% B in 20-25 min and 50% B in 25-30 min, flow rate; 1.0 mL/min; detection, diode array detector (Shimadzu Spd-M20A). The purity of compound is described as percent (%) and retention time was given in minutes. NMR spectra were recorded on Varian AS 400 (^{1}H NMR at 400 MHz and ^{13}C NMR at 100 MHz) with

tetramethylsilane as an internal standard. Chemical shift (d) values are expressed in ppm and

coupling constant (J) values in hertz (Hz). The melting points were measured on Gallenkamp

Melting Point Apparatus without correction. Detailed methods for synthesis of each

compound were indicated in supplementary information.

4.2. Secreted Alkaline Phosphatase (SEAP) Assay

SEAP assay was performed as described previously [29]. The mammalian expression vector,

pBJ-GAL4-ELF3 was co-transfected with the reporter gene, pG5IL2SX to 293T human

kidney cells. Compounds were treated for 16 h, and each of the cultured medium was

analyzed to measure the changes in SEAP activity.

4.3. Western blot analysis

Western blot analysis was performed as described previously [29]. For the experiment,

indicated cells were seeded in 6-well plates and incubated until they reached ~70%

confluence. Stock solutions were prepared in DMSO and PBS for compound **10** and

trastuzumab, respectively. The stock solutions were individually treated with varying

volumes to meet the final concentration as indicated. Media were exchanged to serum-free

media right before the compound treatment. All of the antibodies utilized in this study are

summarized in Table S1.

4.4. RNA extraction and quantitative real-time PCR

Cells were seeded in 60 mm cell culture dishes at a density of 5×10^5 cells per dish and

incubated overnight. The cells were washed with PBS and treated with serum-free medium

containing compounds at indicated concentrations. Total RNA from the compound treated

cells was extracted using the Tri-RNA Reagent (FAVORGEN Biotech Corp., Taiwan) and

cDNA was synthesized from the extracted RNA using PrimeScriptTM RT Reagent Kit

(Takara Bio Inc., Japan) according to the manufacturer's instruction. Quantitative analysis of

the *ERBB2* was performed using a SensiFAST™ SYBR Hi-ROX kit (Bioline, Canada). PCR

amplification condition was applied as described previously [29]. Relative quantity of mRNA

was calculated using the $\Delta\Delta$Ct method and normalized by *GAPDH*. The primer sequences

used in this study are summarized in Table S2.

4.5. Co-immunoprecipitation (Co-IP) assay

NCI-N87 cells were seeded in 100 mm cell culture dish. When they reached to 70~80%

confluency, 10 μM of each candidate compounds were treated for 16 hours. Cell lysates were

prepared under the same procedure utilized in western blot analysis. 400 μg of each protein

aliquots were incubated with MED23 antibody or rabbit-IgG antibody for 6 hours under

rotary agitation at 4 °C. Rabbit-IgG treated samples were compared as control. 15 μl of the

protein-A/G PLUS-agarose beads (sc-2003, Santa Cruz, CA) were then added to each the

samples and again rotated at 4 °C for 8 hours. The incubated samples were centrifuged for

20 min under 13000 rpm. After discarding the supernatant, the beads were washed with 200

μl lysis buffer for 3 times. 15 μl of 2X loading buffer (1M Tris-HCl (pH6.8), glycerol, 10%

SDS, H_2O, bromophenol blue, 2-mercaptoethanol) was added right after removing the

supernatant from last washing step, and boiled at 100 °C for 5 min to denature the protein and

detach from the beads. The samples were loaded on SDS-PAGE after centrifugation at 1300

rpm for 10 min. The following western blot protocol was same as above.

4.6. Fluorescence Poliarization (FP) assay

FP assay was conducted following a previously described method [29]. Different

concentrations of (His)$_6$-MED23$_{391-582}$ and 10 nM of FITC-labeled ELF3$_{129-145}$ peptide were

combinded in assay buffer to determine the K_d value in advance. The calculations were

performed using Prism 6.0 (GraphPad Software, USA) with the least-squares non-linear fit

method. The FP signals were measured in millipolarization (mP) units using the Infinite F200

PRO microplate reader (Tecan Group Ltd., Switzerland) at excitation/emission wavelengths

of 485/535 nm. For the displacement assays, varying concentrations of compound **10** and

unlabeled ELF3$_{137-144}$ peptide were separately added to the mixture containing 80 nM of

(His)$_6$-MED23$_{391-582}$ and 10 nM of FITC-ELF3$_{129-145}$. Activity of compound **10** was assessed

by determining the IC$_{50}$ of each compound, using the four-parameter logistic equation with

Table Curve 2D program (SPSS Inc.). Finally, the K_i values were calculated based on the

Cheng-Prusoff equation: $K_i = IC_{50}/1+([Ligand]/K_d)$.

4.7. Glutathione S-Transferase (GST) pull-down assay

Indicated plasmids were transduced to HEK293 cell using JetPRIME® (Polyplus transfection,

France). 5 μM of compound **10** was co-applied to the system. Cell lysis was prepared under

the same procedure as for the western blot analyses [29]. 1000 μg of cell lysates were

incubated overnight with Glutathione Sepharose™ beads (GE Healthcare, UK) at 4 °C on a

rotator. Beads were washed 3 times with ice-cold 1x PBS, and eluted with elution buffer (20

mM Glutathione, 100 mM Tris-HCl (pH 8.0), 120 mM NaCl, 10% glycerol). The precipitated

proteins were analyzed via western blot.

4.8. Split-luciferase complementation assay

All split luciferase biosensors were made by In-Fusion® HD Cloning kit using firefly

luciferase from the pGL3-basic vector (Promega, USA) template. Luciferase was split into

two fragments (Nluc and Cluc). Full-length cDNA for ELF3 was fused with the N-terminal

fragments of the split luciferase (Nluc), and the MED23$_{391-582}$ fragment was fused with the C-

terminal fragments of the split luciferase (Cluc). Both fragments were cloned into the

p3Xflag-myc-CMV26 vector. The HEK293 cells were co-transfected with Nluc-ELF3 and

Cluc-MED23$_{391-582}$. After incubation for 6 hours, the transfected HEK293 cells were washed

with PBS and treated with compound in a serum-free medium for 20 h. HT59 was treated at

different concentrations.

4.9. Luciferase promoter assay

HEK293 cells were plated in 60 mm culture dishes and transfected with 1 μg of pGL2-HER2

alone or in combination with 0.5 μg of pcDNA3.1-flag-ELF3 and p3Xflag-myc-CMV26-

MED23. All transfections were conducted using jetPRIME® Transfection Reagent (Polyplus-

transfection, FRANCE). After incubation for 6 hours, the culture medium was replaced with

serum-free medium containing different concentrations of HT59 for 20 h. After 20 h,

luciferase activities were measured with the Tecan Infinite® 200 PRO microplate reader

(Tecan, Tecan Group Ltd., Switzerland) using the Luciferase Assay system (Promega, USA),

according to the manufacturer's protocols.

4.10. Annexin V/PI Double Staining Apoptosis Assay

NCI-N87 cells were seeded in 60 mm dishes at a density of 5×10^5 cells per dish. When cells

reached 80% confluence, the cells were treated with 5, 10, and 20 μM of compound **10** for 16

hours. Cells were washed with PBS and harvested using Trypsin-EDTA and centrifugation at

3200 rpm for 3 min. FITC-Annexin V apoptosis detection kit 1 (BD Pharmingen[TM]) was used

to evaluate compound-induced apoptosis. Pellets were washed with PBS and incubated with

100 μL of 1×Annexin V binding buffer containing propidium iodide and FITC-Annexin V for

20 min in the dark at room temperature. The samples were diluted by adding 400 μL of

1×Annexin V binding buffer and then analyzed using Fluorescence-Activated Cell Sorting

(FACS) (BD Biosciences, USA). At least 10,000 cells were measured for each sample.

4.11. Clonogenic assay

Cells were seeded at a density of 1,000 cells per well in 6-well plates and incubated overnight.

After the cells were treated with serum-free medium containing compounds at the indicated

concentrations for 24 hours, the medium was replaced to growth medium without compounds.

The media was changed every 3 days. The plates were incubated for 9 days to allow colony

formation. The colonies were washed with PBS, fixed with 100% methanol for 3 min at room

temperature, and then stained with crystal violet (0.5% in 100% Methanol) for 10 min. The

images were obtained using ChemiDoc (bio-image analyzer, BR179-8280) and the number of

colony was counted using the image J program.

4.12. Tumor xenografts

NCI-N87 cells (5×10^6 cells) were implanted into the flank of 5-week-old female athymic

nude mice (Envigo, USA) using 100 μL of 1x PBS. The mice were then assigned randomly to

different groups once the average tumor volume reached 90-100 mm^3. Subsequently, the drug

was administered intravenously to NCI-N87 xenografts, with six repetitions at 3-day intervals.

Compound **10** was given at the concentration of 4 mg/kg, prepared in

DMAC/Tween80/saline (5:10:85) mixture along with saline. Tumor size changes were

monitored for additional days after the last drug injection until the average tumor size of the

control group reached 2000-2500 mm^3. Mice were sacrificed 25 days after the first drug

injection, and tumors were immediately excised from each mouse. The relative tumor sizes

were determined by measuring the tumor length (L) and width (W) with calipers and the

formula $(L \times W^2) / 2$ was used for calculations.

4.13. IHC Assay for Xenograft Mouse Model

Tumors excised from the xenograft mouse model were processed into paraffin-embedded

block sections for IHC. The experiments were carried out following standard protocols. The

specified HER2 and Ki67 primary antibodies were incubated at 4 °C overnight and then

washed 3 times with 1x PBS. Subsequently, they were exposed to secondary antibodies,

developed using a Vectastain ABC kit (Vector Laboratories, USA), and stained with DAB

solution (Dako, Carpinteria, USA), all in accordance with the manufacturers' protocols. The

IHC staining was further counterstained with hematoxylin (USA), and was evaluated under

light microscopy at 200× magnification. A semi-quantitative IHC score was adopted for

evaluation, where the final scores were calculated by multiplying the intensity and fraction

score (percentage of samples counted at each scale), leading to a range from 0 to 300. All

imaging and evaluation were conducted using an Axiophot 2 apparatus (Carl Zeiss

MicroImaging Inc., Thornwood, NY, USA), at the Drug Development Research Core Center.

The details of the applied antibodies and dilution ratios can be found in Table S1.

4.14. In silico qualitative SAR analysis

For the qualitative SAR analysis Cresset's Flare v7.1 software was utilized. All the prepared

28 compounds were primarily aligned in FieldTemplater™ to generate conformations and

molecular fields for each compound. Using compound 10 as the reference molecule,

qualitative Activity Atlas model was subsequently constructed under the default settings [47]

to visualize the activity cliff summary of electrostatics and hydrophobics in 3D plots.

4.15. Development of trastuzumab resistant NCI-N87

To develop trastuzumab-resistance NCI-N87 cells, NCI-N87 cells were continuously treated

with gradually increased amount of trastuzumab. Cells were cultured in medium containing 3

to 10 μg/ml of trastuzumab until 30th passage.

4.16. Statistical analysis

All experiments were performed at least three times and all results were expressed as mean ±

standard deviation. Statistics were analyzed via one-way analysis of variance (ANOVA) or

Student's t-test with Prism 6.0 (GraphPad Software, USA). Differences between two values

were considered statistically significant when the *p*-values (demonstrated as single, double or

triple asterisks) were <0.05, <0.01, and <0.001.

Acknowledgements

This work was supported by grants from the National Research Foundation of Korea (NRF),

funded by the Korean government (MSIT) (NRF-2018R1A5A2025286, 2021M3E5E7024855,

2022R1A2C2092053), and by a grant from the Korea Basic Science Institute (National

Research Facilities and Equipment Center), funded by the Ministry of Education

(2021R1A6C101A442).

References

[1] Iqbal N., Iqbal N., Human Epidermal Growth Factor Receptor 2 (HER2) in Cancers:
Overexpression and Therapeutic Implications, Molecular biology international. 2014, (2014) 852748.
10.1155/2014/852748.

[2] Gravalos C., Jimeno A., HER2 in gastric cancer: a new prognostic factor and a novel therapeutic
target, Annals of oncology : official journal of the European Society for Medical Oncology. 19,
(2008) 1523-9. 10.1093/annonc/mdn169.

[3] Kelly C.M., Janjigian Y.Y., The genomics and therapeutics of HER2-positive gastric cancer-from
trastuzumab and beyond, Journal of gastrointestinal oncology. 7, (2016) 750-62.
10.21037/jgo.2016.06.10.

[4] Pellino A., Riello E., Nappo F., Brignola S., Murgioni S., Djaballah S.A., Lonardi S., Zagonel V.,
Rugge M., Loupakis F., Fassan M., Targeted therapies in metastatic gastric cancer: Current
knowledge and future perspectives, World journal of gastroenterology. 25, (2019) 5773-88.
10.3748/wjg.v25.i38.5773.

[5] Yan M., Schwaederle M., Arguello D., Millis S.Z., Gatalica Z., Kurzrock R., HER2 expression status
in diverse cancers: review of results from 37,992 patients, Cancer metastasis reviews. 34, (2015)
157-64. 10.1007/s10555-015-9552-6.

[6] Swain S.M., Shastry M., Hamilton E., Targeting HER2-positive breast cancer: advances and
future directions, Nature reviews Drug discovery. 22, (2023) 101-26. 10.1038/s41573-022-00579-0.

[7] Chen G., Zeng L., Bi B., Huang X., Qiu M., Chen P., Chen Z.Y., He Y., Pan Y., Chen Y., Zhao J.,
Engineering Bifunctional Calcium Alendronate Gene-Delivery Nanoneedle for Synergistic
Chemo/Immuno-Therapy Against HER2 Positive Ovarian Cancer, Advanced science (Weinheim,
Baden-Wurttemberg, Germany). 10, (2023) e2204654. 10.1002/advs.202204654.

[8] Ivanova M., Venetis K., Guerini-Rocco E., Bottiglieri L., Mastropasqua M.G., Garrone O., Fusco N.,
Ghidini M., HER2 in Metastatic Colorectal Cancer: Pathology, Somatic Alterations, and Perspectives
for Novel Therapeutic Schemes, Life (Basel, Switzerland). 12, (2022). 10.3390/life12091403.

[9] Han S.H., Ryu K.H., Kwon A.Y., The Prognostic Impact of HER2 Genetic and Protein Expression in
Pancreatic Carcinoma-HER2 Protein and Gene in Pancreatic Cancer, Diagnostics (Basel, Switzerland).
11, (2021). 10.3390/diagnostics11040653.

[10] Guan W.L., He Y., Xu R.H., Gastric cancer treatment: recent progress and future perspectives,
Journal of hematology & oncology. 16, (2023) 57. 10.1186/s13045-023-01451-3.

[11] Luo H., Xu X., Ye M., Sheng B., Zhu X., The prognostic value of HER2 in ovarian cancer: A
meta-analysis of observational studies, PloS one. 13, (2018) e0191972.
10.1371/journal.pone.0191972.

[12] Ajani J.A., D'Amico T.A., Bentrem D.J., Chao J., Cooke D., Corvera C., Das P., Enzinger P.C., Enzler T., Fanta P.J.J.o.t.N.C.C.N., Gastric cancer, version 2.2022, NCCN clinical practice guidelines in oncology. 20, (2022) 167-92.

[13] Gradishar W.J., Moran M.S., Abraham J., Aft R., Agnese D., Allison K.H., Anderson B., Burstein H.J., Chew H., Dang C., Elias A.D., Giordano S.H., Goetz M.P., Goldstein L.J., Hurvitz S.A., Isakoff S.J., Jankowitz R.C., Javid S.H., Krishnamurthy J., Leitch M., Lyons J., Mortimer J., Patel S.A., Pierce L.J., Rosenberger L.H., Rugo H.S., Sitapati A., Smith K.L., Smith M.L., Soliman H., Stringer-Reasor E.M., Telli M.L., Ward J.H., Wisinski K.B., Young J.S., Burns J., Kumar R., Breast Cancer, Version 3.2022, NCCN Clinical Practice Guidelines in Oncology, Journal of the National Comprehensive Cancer Network : JNCCN. 20, (2022) 691-722. 10.6004/jnccn.2022.0030.

[14] Malla M., Pedersen K.S., Parikh A.R., Updates in the Treatment of Metastatic Colorectal Cancer %J Journal of the National Comprehensive Cancer Network. 21, (2023) 567-71. 10.6004/jnccn.2023.5012.

[15] Maadi H., Soheilifar M.H., Choi W.-S., Moshtaghian A., Wang Z., Trastuzumab Mechanism of Action; 20 Years of Research to Unravel a Dilemma. 13, (2021) 3540.

[16] Liu Q., Kulak M.V., Borcherding N., Maina P.K., Zhang W., Weigel R.J., Qi H.H., A novel HER2 gene body enhancer contributes to HER2 expression, Oncogene. 37, (2018) 687-94. 10.1038/onc.2017.382.

[17] Duru N., Fan M., Candas D., Menaa C., Liu H.C., Nantajit D., Wen Y., Xiao K., Eldridge A., Chromy B.A., Li S., Spitz D.R., Lam K.S., Wicha M.S., Li J.J., HER2-associated radioresistance of breast cancer stem cells isolated from HER2-negative breast cancer cells, Clinical cancer research : an official journal of the American Association for Cancer Research. 18, (2012) 6634-47. 10.1158/1078-0432.Ccr-12-1436.

[18] Cao N., Li S., Wang Z., Ahmed K.M., Degnan M.E., Fan M., Dynlacht J.R., Li J.J., NF-kappaB-mediated HER2 overexpression in radiation-adaptive resistance, Radiation research. 171, (2009) 9-21. 10.1667/rr1472.1.

[19] Chang C.H., Scott G.K., Kuo W.L., Xiong X., Suzdaltseva Y., Park J.W., Sayre P., Erny K., Collins C., Gray J.W., Benz C.C., ESX: a structurally unique Ets overexpressed early during human breast

tumorigenesis, Oncogene. 14, (1997) 1617-22. 10.1038/sj.onc.1200978.

[20] Liu X., Go M.L., Antiproliferative properties of piperidinylchalcones, Bioorganic & medicinal
chemistry. 14, (2006) 153-63. 10.1016/j.bmc.2005.08.006.

[21] Modzelewska A., Pettit C., Achanta G., Davidson N.E., Huang P., Khan S.R., Anticancer activities
of novel chalcone and bis-chalcone derivatives, Bioorganic & medicinal chemistry. 14, (2006)
3491-5. 10.1016/j.bmc.2006.01.003.

[22] Ni L., Meng C.Q., Sikorski J.A., Recent advances in therapeutic chalcones, Expert Opinion on
Therapeutic Patents. 14, (2004) 1669-91. 10.1517/13543776.14.12.1669.

[23] Friis-Møller A., Chen M., Fuursted K., Christensen S.B., Kharazmi A., In vitro antimycobacterial
and antilegionella activity of licochalcone A from Chinese licorice roots, Planta medica. 68, (2002)
416-9. 10.1055/s-2002-32087.

[24] Mahapatra D.K., Bharti S.K., Asati V., Anti-cancer chalcones: Structural and molecular target
perspectives, European journal of medicinal chemistry. 98, (2015) 69-114.
10.1016/j.ejmech.2015.05.004.

[25] Mishra L., Sinha R., Itokawa H., Bastow K.F., Tachibana Y., Nakanishi Y., Kilgore N., Lee K.H.,
Anti-HIV and cytotoxic activities of Ru(II)/Ru(III) polypyridyl complexes containing 2,6-(2'-
benzimidazolyl)-pyridine/chalcone as co-ligand, Bioorganic & medicinal chemistry. 9, (2001) 1667-
71. 10.1016/s0968-0896(01)00074-8.

[26] Chen M., Brøgger Christensen S., Zhai L., Rasmussen M.H., Theander T.G., Frøkjaer S.,
Steffansen B., Davidsen J., Kharazmi A., The novel oxygenated chalcone, 2,4-dimethoxy-4'-
butoxychalcone, exhibits potent activity against human malaria parasite Plasmodium falciparum in
vitro and rodent parasites Plasmodium berghei and Plasmodium yoelii in vivo, The Journal of
infectious diseases. 176, (1997) 1327-33. 10.1086/514129.

[27] Burgess A.W., Cho H.S., Eigenbrot C., Ferguson K.M., Garrett T.P., Leahy D.J., Lemmon M.A.,
Sliwkowski M.X., Ward C.W., Yokoyama S., An open-and-shut case? Recent insights into the
activation of EGF/ErbB receptors, Molecular cell. 12, (2003) 541-52. 10.1016/s1097-2765(03)00350-
2.

[28] Md. Azizur R., Anees A.S., PYRAZOLINE DERIVATIVES: A WORTHY INSIGHT INTO THE RECENT ADVANCES AND POTENTIAL PHARMACOLOGICAL ACTIVITIES, International Journal of Pharmaceutical Sciences and Drug Research. 2, (2010).

[29] Hwang S.Y., Park S., Jo H., Hee Seo S., Jeon K.H., Kim S., Jung A.R., Song C., Ahn M., Yeon Kwak S., Lee H.J., Uesugi M., Na Y., Kwon Y., Interrupting specific hydrogen bonds between ELF3 and MED23 as an alternative drug resistance-free strategy for HER2-overexpressing cancers, J Adv Res. 47, (2023) 173-87. 10.1016/j.jare.2022.08.003.

[30] Kim H.L., Jeon K.H., Jun K.Y., Choi Y., Kim D.K., Na Y., Kwon Y., A-62176, a potent topoisomerase inhibitor, inhibits the expression of human epidermal growth factor receptor 2, Cancer letters. 325, (2012) 72-9. 10.1016/j.canlet.2012.06.004.

[31] Cheeseright T., Mackey M., Rose S., Vinter A., Molecular field extrema as descriptors of biological activity: definition and validation, J Chem Inf Model. 46, (2006) 665-76. 10.1021/ci050357s.

[32] Floresta G., Abbate V.J.R.a., Machine learning vs. field 3D-QSAR models for serotonin 2A receptor psychoactive substances identification. 11, (2021) 14587-95.

[33] Tedesco G., Cheeseright T., Mackey M., Summarizing and Understanding SAR and Activity Cliffs in GPCRs.

[34] NCCN, NCCN GUIDELINES FOR PATIENTS®, Available at: https://wwwnccnorg/patients/guidelines/content/PDF/stomach-patientpdf. (2019).

[35] Gradishar W.J., Anderson B.O., Abraham J., Aft R., Agnese D., Allison K.H., Blair S.L., Burstein H.J., Dang C., Elias A.D., Giordano S.H., Goetz M.P., Goldstein L.J., Isakoff S.J., Krishnamurthy J., Lyons J., Marcom P.K., Matro J., Mayer I.A., Moran M.S., Mortimer J., O'Regan R.M., Patel S.A., Pierce L.J., Rugo H.S., Sitapati A., Smith K.L., Smith M.L., Soliman H., Stringer-Reasor E.M., Telli M.L., Ward J.H., Young J.S., Burns J.L., Kumar R., Breast Cancer, Version 3.2020, NCCN Clinical Practice Guidelines in Oncology, Journal of the National Comprehensive Cancer Network : JNCCN. 18, (2020) 452-78. 10.6004/jnccn.2020.0016.

[36] Fiszman G.L., Jasnis M.A.J.I.j.o.b.c., Molecular mechanisms of trastuzumab resistance in HER2
overexpressing breast cancer. 2011, (2011).

[37] Ferrando-Díez A., Felip E., Pous A., Bergamino Sirven M., Margelí M., Targeted Therapeutic
Options and Future Perspectives for HER2-Positive Breast Cancer, Cancers. 14, (2022).
10.3390/cancers14143305.

[38] Tarantino P., Morganti S., Curigliano G., Targeting HER2 in breast cancer: new drugs and
paradigms on the horizon, Exploration of targeted anti-tumor therapy. 2, (2021) 139-55.
10.37349/etat.2021.00037.

[39] Kreutzfeldt J., Rozeboom B., Dey N., De P., The trastuzumab era: current and upcoming
targeted HER2+ breast cancer therapies, Am J Cancer Res. 10, (2020) 1045-67.

[40] Wang Z.H., Zheng Z.Q., Jia S.C., Liu S.N., Xiao X.F., Chen G.Y., Liang W.Q., Lu X.F., Trastuzumab
resistance in HER2-positive breast cancer: Mechanisms, emerging biomarkers and targeting agents,
Front Oncol. 12, (2022) 1006429. 10.3389/fonc.2022.1006429.

[41] Vivekanandhan S., Knutson K.L., Resistance to Trastuzumab, Cancers (Basel). 14, (2022).
10.3390/cancers14205115.

[42] Cresti N., Lee J., Rourke E., Televantou D., Jamieson D., Verrill M., Boddy A.V., Genetic variants
in the HER2 gene: Influence on HER2 overexpression and loss of heterozygosity in breast cancer,
Eur J Cancer. 55, (2016) 27-37. 10.1016/j.ejca.2015.10.066.

[43] Nam J.M., Jeon K.H., Kwon H., Lee E., Jun K.Y., Jin Y.B., Lee Y.S., Na Y., Kwon Y.,
Dithiiranylmethyloxy azaxanthone shows potent anti-tumor activity via suppression of HER2
expression and HER2-mediated signals in HER2-overexpressing breast cancer cells, Eur J Pharm Sci.
50, (2013) 181-90. 10.1016/j.ejps.2013.06.014.

[44] Shimogawa H., Kwon Y., Mao Q., Kawazoe Y., Choi Y., Asada S., Kigoshi H., Uesugi M., A
wrench-shaped synthetic molecule that modulates a transcription factor-coactivator interaction, J
Am Chem Soc. 126, (2004) 3461-71. 10.1021/ja038855+.

[45] Li Q., Meissner T.B., Wang F., Du Z., Ma S., Kshirsagar S., Tilburgs T., Buenrostro J.D., Uesugi M.,

Strominger J.L., ELF3 activated by a superenhancer and an autoregulatory feedback loop is
required for high-level HLA-C expression on extravillous trophoblasts, Proc Natl Acad Sci U S A.
118, (2021). 10.1073/pnas.2025512118.

744
745 [46] Asada S., Choi Y., Yamada M., Wang S.C., Hung M.C., Qin J., Uesugi M., External control of
746 Her2 expression and cancer cell growth by targeting a Ras-linked coactivator, Proc Natl Acad Sci U
747 S A. 99, (2002) 12747-52. 10.1073/pnas.202162199.

748
749 [47] Mackey M., 250th ACS National Meeting & Exposition, http://wwwcresset-groupcom/activity-
750 atlas/. (2015).

Chalcone

Pyrazoline

Licochalcone E

YK-1

Proposed compounds

Group 1

1	R$_1$ = CH$_3$, R$_2$ = H
2	R$_1$ = CH$_3$, R$_2$ = Propenyl
3	R$_1$ = CH$_3$, R$_2$ = 3-Butenyl
4	R$_1$ = CH$_3$, R$_2$ = 2-Butenyl
5	R$_1$ = CH$_3$, R$_2$ = 3-Methyl-but-2-enyl
6	R$_1$ = H, R$_2$ = CH$_3$
7	R$_1$ = Propenyl , R$_2$ = CH$_3$
8	R$_1$ = 3-Butenyl, R$_2$ = CH$_3$
9	R$_1$ = 2-Butenyl, R$_2$ = CH$_3$
10	R$_1$ = 3-Methyl-but-2-enyl, R$_2$ = CH$_3$

Group 2

11	R$_1$ = CH$_3$, R$_2$ = H
12	R$_1$ = CH$_3$, R$_2$ = Propenyl
13	R$_1$ = CH$_3$, R$_2$ = 3-Butenyl
14	R$_1$ = CH$_3$, R$_2$ = 2-Butenyl
15	R$_1$ = H, R$_2$ = CH$_3$
16	R$_1$ = Propenyl , R$_2$ = CH$_3$
17	R$_1$ = 2-Butenyl, R$_2$ = CH$_3$

Group 3

18	R$_1$ = CH$_3$, R$_2$ = H
19	R$_1$ = CH$_3$, R$_2$ = Propenyl
20	R$_1$ = CH$_3$, R$_2$ = 3-Butenyl
21	R$_1$ = CH$_3$, R$_2$ = 2-Butenyl
22	R$_1$ = CH$_3$, R$_2$ = 3-Methyl-but-2-enyl
23	R$_1$ = H, R$_2$ = CH$_3$
24	R$_1$ = Propenyl , R$_2$ = CH$_3$
25	R$_1$ = 2-Butenyl, R$_2$ = CH$_3$
26	R$_1$ = 3-Methyl-but-2-enyl, R$_2$ = CH$_3$

(A)

■ Activity Cliff Summary of Positive Electrostatics
■ Activity Cliff Summary of Negative Electrostatics

Comp. 3
%Inh: 58.7

Comp. 5
%Inh: 52.4

chalcone

Comp. 10
%Inh: 62.7

Comp. 12
%Inh: 0

Comp. 15
%Inh: 12.8

pyrazoline acetyl

Comp. 20
%Inh: 0

Comp. 24
%Inh: 0

pyrazoline propionyl

%Inh: Relative HER2 inhibitory activity

(B)

▦ Activity Cliff Summary of Favorable Hydrophobics
▦ Activity Cliff Summary of Unfavorable Hydrophobics

Comp. 3
%Inh: 58.7

Comp. 5
%Inh: 52.4

chalcone

Comp. 10
%Inh: 62.7

Comp. 12
%Inh: 0

Comp. 15
%Inh: 12.8

pyrazoline acetyl

%Inh: Relative HER2 inhibitory activity

Comp. 20
%Inh: 0

Comp. 24
%Inh: 0

pyrazoline propionyl

(A)

Propidium iodide

CON   5 μM   10 μM   15 μM

Annexin V-FITC

(B)

Comp. **10** (μM, 16 h)

CON 5 10 15

c-PARP

c-Caspase 3

GAPDH

Comp. **10** (h, 10 μM)

CON 4 8 16

c-PARP

c-Caspase 3

GAPDH

(C)

Comp. **10** (μM)

CON 1 5 10

(D)

(E)

CON

Comp. **10**

(F)

(G)

HER2   Ki67

CON

Comp. **10**

(A)

Growth inhibition (% of control)

parent   TR   ***

(B)

TZMB (10 µg/mL)   −   +   −
Comp.**10** (10 µM)   −   −   +

HER2
p-AKT
AKT
p-MAPK
MAPK
GAPDH

(C)

NCI-N87 TR

CON   TZMB   Comp. 10

Survival fraction (%)

CON   TZMB   Comp.10
ns   *

(D)

Comp. 10 (µM)

CON   5   10   20

Propidium iodide

Annexin V

Apoptotic cell fraction (relative ratio)

CON   5   10   20   ns   ***   ***
Comp. 10 (µM)