1

Cyclin A2 Induces Human Adult Cardiomyocyte Cytokinesis and Elicits Cardiomyocyte Reprogramming and Dedifferentiation

Esmaa Bouhamida[1], Sangeetha Vadakke-Madathil[1#], Prabhu Mathiyalagan[2#], Amaresh K. Ranjan[1,3#], Amir Khan[1], Cherrie D. Sherman[1], Paul E Miller[4], Andre Ghetti[4], Najah Abi-Gerges[4] and Hina W. Chaudhry[1*]

[1] Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
[2] Benthos Prime Central, TX, USA
[3] Pharmazz Inc., Willowbrook, IL, USA, 60527
[4] AnaBios Corporation, San Diego, CA 92109, USA
[#] Contributed equally

* Correspondence to:

Hina W. Chaudhry, M.D., Department of Cardiology, Cardiovascular Regenerative Medicine, Mount Sinai School of Medicine, One Gustave L. Levy Place, Box 1030, New York, NY 10029, Phone: 212-241-4104, Fax: 212-241-5510. hina.chaudhry@mssm.edu.

Abstract

Cyclin A2 (CCNA2), a master cell cycle regulator, is silenced in postnatal mammalian cardiomyocytes. We have previously demonstrated its ability to promote cardiac repair in small and large animals when delivered to the heart via a viral vector. However, the effect of CCNA2 gene delivery on cytokinesis in isolated cardiomyocytes from adult human hearts has not been explored. We designed a human gene therapy vector featuring a replication-deficient adenovirus encoding human CCNA2 driven by the cardiac Troponin T promoter to enable the expression of CCNA2 in freshly isolated human cardiomyocytes. Time-lapse live imaging of adult human primary cardiomyocytes from a 21-year-old male, a 41-year-old female, and a 55-year-old male demonstrated the induction of complete cytokinesis in human adult cardiomyocytes with preservation of sarcomere integrity in the resulting daughter cells with active calcium mobilization in redifferentiated cardiomyocytes. To elucidate the transcriptional mechanisms underlying this response, we conducted single-nucleus transcriptomics analysis of hearts isolated from adult transgenic mice that constitutively express CCNA2 in cardiomyocytes (CCNA2-Tg) and non-transgenic mice (nTg). This revealed a cardiomyocyte subpopulation enriched with cytokinesis, proliferative, and reprogramming genes in hearts obtained from CCNA2-Tg mice as compared to nTg mice. Ultra-deep bulk RNA sequencing of human adult and fetal hearts identified key reprogramming genes relevant to understanding the mechanisms of CCNA2-induced effects observed in our experimental models. These findings provide a promising path for the clinical development of CCNA2-based cardiac regenerative therapy.

43 **Graphical Abstract**

44
45
46 CCNA2 Induces Cytokinesis and Drives Proliferation and Reprogramming of Adult
47 Cardiomyocytes: An Integrative Transcriptomic Analysis across Human and Mouse Models.
48
49
50
51

Introduction

The limited proliferative capacity of adult human cardiomyocytes remains a major barrier to effective cardiac regeneration following injury, contributing to the significant morbidity and mortality associated with heart disease [1]. Scar formation via fibrosis is, therefore, the primary response to cardiac injury.   Multitudes of molecular and cellular approaches have been investigated over the past 20 years aimed at regenerating the myocardium in various states of heart disease [2-4]. The use of stem/progenitor cell therapies for myocardial regeneration has demonstrated limited efficacy in clinical trials, with no conclusive evidence of actual cardiomyocyte differentiation arising from the cell types used previously in these trials. Human embryonic stem (ES) cells and induced pluripotent (iPS) cells have been evaluated in non-human primate models of heart disease with the demonstrated ability of these cell types to differentiate, at the very least, to immature cardiomyocytes with the hope that they would undergo maturation *in vivo.* However, it remains to be seen whether these approaches may be translated for human therapy [3, 5]. Aside from the differentiation and generation of new cardiomyocytes, some technical concerns still need to be addressed, such as the propensity for arrhythmias noted in primate hearts treated with ES-derived and iPS-derived cardiomyocytes [5, 6].

There has been evidence of low-level cardiomyocyte turnover in the healthy human heart, but it is very limited, and this ability declines with age [7, 8]. Thus, cardiac regeneration in response to injury such as myocardial infarction (MI) remains a clinical challenge. Evolution may not have favored adult mammals in this regard, but certain members of metazoan species are capable of heart regeneration, and these examples may perhaps serve to enlighten us. Urodele amphibians, such as the newt, retain a remarkable capacity to replace lost anatomical tissues through epimorphic regeneration [9]. This process relies on the local plasticity of differentiated cells near the region of injury and involves reentry to the cell cycle with loss of differentiated characteristics to generate a 'local progenitor cell' of restricted potentiality [9]. The adult zebrafish heart can regenerate up to 20% of its volume via a mechanism largely dependent on the proliferation of cardiomyocytes adjacent to the area of injury [10, 11]. However, zebrafish harboring a temperature-sensitive mutation in *mps1*, a gene encoding a mitotic checkpoint kinase [10], cannot regenerate the heart, and scarring and fibrosis are noted in the excised areas in a manner similar to the response of the human myocardium after injury [10, 12]. In summary, the loss of a single gene can avert cardiomyocyte mitosis and mitigate the normal regenerative process, thus permitting fibrosis to proceed unimpeded.

Previously, we have reported that CCNA2 is a 'master regulator' of the cardiomyocyte cell cycle [13]. Unlike other cyclins, cyclin A2 complexes with its cyclin-dependent kinase partners to regulate both key transitions of the cell cycle: G1-S and G2-M [14-16], and is silenced shortly after birth in mammalian cardiomyocytes [1, 17, 18]. Subsequently, we have also shown that *Ccna2* mediates cardiac repair by inducing cardiomyocyte mitoses after MI in two small animal models of MI [19-21]. As molecular mechanisms may be widely divergent across species, we performed a therapeutic efficacy study of CCNA2-mediated cardiac repair in a porcine model of MI as it closely mimics human anatomy and physiology [22]. Therapeutic delivery of *CCNA2* one week after MI in the porcine heart induced cardiomyocyte cell cycle activation *in vivo* with marked

enhancement of cardiac function as noted with multimodality imaging, including magnetic
resonance imaging. The experimental pigs also exhibited significantly decreased fibrosis, lack of
cardiomyocyte hypertrophy, and a 55% increase in cardiomyocyte cell numbers in the peri-infarct
zones. Using the same viral vector encoding CCNA2 that was delivered *in vivo*, cultured porcine
adult cardiomyocytes were induced to undergo cytokinesis with preservation of sarcomere
structure and captured via live imaging [22].

For the purposes of clinical translation, we designed the next-generation gene therapy vector
using a cardiomyocyte-specific promoter, cardiac Troponin T (cTnT), driving human CCNA2
expression. We then sought to determine whether this cTnT-CCNA2 vector would induce human
adult cardiomyocytes to undergo cytokinesis. We employed a multiparametric approach
combining live-cell imaging with high-resolution transcriptomics to uncover mechanistic insights
underlying CCNA2-driven cytokinesis in adult cardiomyocytes. Live imaging microscopy enabled
dynamic tracking of sarcomere integrity and cellular architecture during cytokinesis in adult human
cardiomyocytes. To dissect the transcriptional programs and signaling pathways associated with
CCNA2 expression, we leveraged three complementary RNA sequencing datasets: (i) bulk RNA-
sequencing of CCNA2-Tg and nTg cardiomyocytes, (ii) ultra-deep bulk RNA-sequencing of adult
and fetal human hearts, and (iii) single-nucleus RNA sequencing (snRNA-seq, 10x Genomics).

This integrative approach defined the molecular framework by which CCNA2 reactivates cell cycle
networks, promotes cytokinesis, and induces a regenerative transcriptional state in postnatal
cardiomyocytes, offering insights for future cardiac regenerative therapies.

Results

cTnT-CCNA2 adenovirus vector designed for therapeutic use induces expression
of CCNA2 in cultured adult human cardiomyocytes

First, we designed a therapeutic grade human CCNA2 adenovirus vector to selectively express
human cyclin A2 (CCNA2) in cardiomyocytes by cloning human cDNA (NCBI Reference
Sequence: NM_001237.4; 374-1672 bp) downstream to the cTnT promoter **(Figure 1A)**. The
cultured adult human cardiomyocytes were transduced with cTnT-hCCNA2 (test) versus cTnT-
eGFP (control) adenoviruses with multiplicity of infection (MOI) of 100 each for assessing the
induced expression of CCNA2. We observed significantly increased expression of CCNA2 in the
cultured adult human cardiomyocytes transduced with the test compared to the control adenovirus
(**Figure 1B**). We utilized cTnT-H2B-GFP to confirm that the nuclei of the daughter cells belonged
to cardiomyocytes.

Adenoviral vector-mediated expression of CCNA2 induces cytokinesis in cultured
adult human cardiomyocytes

To assess the effect of induced CCNA2 expression on cell division in adult human
cardiomyocytes, we cultured adult human cardiomyocytes and studied cytokinesis *in vitro* by

using live cell epifluorescence time-lapse microscopy. The adult human cardiomyocytes were
plated at equal densities and transduced with cTnT-hCCNA2, cTnT-eGFP and/or cTnT-H2B-GFP
and CMV-α-act-mCherry adenoviruses (test) or with cTnT-eGFP and/or cTnT-H2B-GFP and
CMV-α-act-mCherry adenoviruses (control). cTnT-eGFP and/or cTnT-H2B-GFP adenovirus was
transduced in both groups to confirm the initial tracking of cardiomyocytes (green) during live cell
epifluorescence microscopy. For live visualization of sarcomeric structure in cardiomyocytes, cells
from both groups were co-transduced with adenovirus containing α-act-mCherry, which we had
constructed to allow for proper folding of the virally delivered α-actinin into the live cardiomyocyte
sarcomere (CMV-α-act-mCherry) and was successfully used in cultured adult porcine
cardiomyocytes [22]. This strategy enabled us to confirm cardiomyocyte identity by assessing the
expression of eGFP before and after cytokinesis and tracking sarcomere dynamics during live
cell imaging. This approach provides a significant advantage over antibody-based identification,
which is susceptible to artifacts and restricted to a single time point after cell fixation. We observed
co-expression of eGFP (green) and α-actinin (red) in cultured adult cardiomyocytes **(Figure 1C**
and **1H**; first panel). We performed time-lapse microscopic imaging of live cells to capture
cardiomyocyte cytokinesis **(Figure 1C, 1F**, and **1H;** still images from **movie S1)**. The cytokinetic
index of adult human cardiomyocytes was calculated by counting the cytokinetic events observed
in 42 regions of interest (ROIs) **(Source Data Figure 1I)**. The cytokinetic index was significantly
higher in the test samples with cTnT-CCNA2 adenovirus transduction compared to control
samples in the 41- and 55-year-olds **(Figure 1I).** In contrast, no significant change in the
cytokinetic events between test and control samples was observed in the 21-year-old patient's
cardiomyocytes **(Figure 1I)**. Most remarkably, sarcomere structure was preserved in the daughter
cells after cytokinesis **(Figure 1D** and **Figure 1G**; upon magnification of a daughter cell, the
presence of sarcomeric structure is easily noted). The daughter cells were further identified with
the expression of H2B-GFP (as they were originally also transduced with cTnT-eGFP and/or H2B-
GFP) and noted to be mononuclear after they had been fixed and stained with DAPI **(Figure 1E**
and **1H)**. Clusters of other cardiomyocytes with expression of eGFP could be seen adjacent to
the daughter cells **(Figure 1E)**.
167
Next, we assessed the functional differentiation of cardiomyocytes by measuring the Ca^{2+}
transients in CCNA2-expressing cardiomyocytes using live-cell calcium imaging under pacing
conditions (0.5 Hz and 1 Hz; **Figure 2A**). Cardiomyocytes transduced with Ad5-cTnT-hCCNA2
exhibited active Ca^{2+} transients **(Figure 2B)**. These findings demonstrate that CCNA2 drives
cytokinesis while preserving sarcomere integrity and calcium mobilization in adult human
cardiomyocytes suggesting functional differentiation.
174

CCNA2 reprogramming induces fetal gene signatures for calcium handling and sarcomere dynamics

177
To quantify the developmental and reprogramming landscape, we performed ultra-deep bulk
RNA-sequencing of fetal (n=3) and adult (n=4) human hearts. This approach allowed us to
compare the molecular signatures of CCNA2-induced cardiomyocyte reprogramming with
naturally occurring fetal-like states. Our ultra-deep sequencing approach enhanced our ability to
detect low-abundance transcripts, improve quantification accuracy, and provide a more

comprehensive transcriptomic profile. This comparison contextualizes how CCNA2 expression
modulates gene networks associated with dedifferentiation, proliferation, and calcium handling,
highlighting its role in cardiac plasticity and regeneration. Notably, there were overlapping
differentially expressed genes (DEGs) between adult CCNA2-Tg mouse cardiomyocytes and fetal
human hearts with downregulated genes in both that were associated with cardiomyocyte
maturation **(Figure 2C and 2D)**. Furthermore, we found that calcium handling and contractility-
related genes, as well as sarcomere assembly genes such as *RYR2, ATP2A2, SLC8A1, CALM1,*
CALM2, NEB, TTN, and *TNNC1* were differentially upregulated in adult human hearts compared
to fetal human hearts **(Figure 2E)**.
192
Additionally, we performed bulk RNA-sequencing of mouse cardiomyocytes with constitutive
CCNA2 expression in cardiomyocytes (CCNA-Tg mice) compared to non-Tg mice (n=6 in total).
Key calcium handling genes, including *Ryr2, Atp2a2, Slc8a1,* and sarcomere assembly genes as
well as *Neb, Ttn,* and *Tnnc1,* were significantly downregulated in CCNA2-Tg cardiomyocytes. In
contrast, genes involved in intracellular calcium release and store-operated calcium entry, such
as *Itpr1, Orai2,* and *Stim2*, were upregulated in CCNA2-Tg mouse hearts, further aligning with
the fetal gene expression profile **(Figure 2F and S1)**. This suggests that *CCNA2* may regulate
sarcomere assembly genes during cell division **(Figure S1)**, thereby facilitating cardiomyocyte
proliferation [23]. Furthermore, *CCNA2* appears to influence calcium mobilization post-
cytokinesis, suggesting a broader role beyond cell cycle regulation, potentially impacting
cardiomyocyte maturation and function.
204
Live-cell imaging of CCNA2-overexpression in adult human cardiomyocytes undergoing
cytokinesis revealed that daughter cells retain sarcomeric integrity post-division. Despite RNA-
sequencing data showing a significant downregulation of thin filament genes (*Neb, Tnn, Tnnc1*),
structural sarcomere proteins such as *Tnnt2* and *Actn2* remained stable **(Figure S1)**, suggesting
that sarcomere remodeling may occur transiently to accommodate cell division without leading to
complete dedifferentiation. Interestingly, transcriptomic profiling revealed upregulation of *ZEB1* in
both fetal hearts and CCNA2-Tg mouse hearts, suggesting a conserved role in cardiomyocyte
plasticity **(Figure S1)**. *SNAI1,* a gene implicated in epithelial to mesenchymal transition with roles
in cancer development and progression, was not consistently upregulated. However, *ZEB1*'s
expression aligns with transient mesenchymal activation, which has been implicated in neonatal
heart regeneration [24].
216

217 Single-nucleus transcriptomic profiling reveals CCNA2-driven reprogramming in
218 selective cardiomyocytes, supported by integrative bulk RNA-seq analysis

219
To investigate CCNA2-induced cardiomyocyte cytokinesis in an unbiased manner, we performed
snRNA-seq on hearts isolated from control non-transgenic (nTg) and *CCNA2* constitutively
expressing transgenic (CCNA2-Tg) adult mice (n=8 in total). Both male and female mice aged 8-
12 weeks were included. This approach enabled the identification of differentially expressed
genes across cardiac clusters in both nTg and CCNA2-Tg mice in the context of CCNA2
expression. **(Figure 3A**, **3B**, and **Figure S1)**. Notably, snRNA-seq also captured

underrepresented cell types often lost during enzymatic digestion for single cell RNA-seq,
providing a comprehensive view of cardiac cellular heterogeneity [23, 25].
228
t-distributed Stochastic Neighbor Embedding (t-SNE) of combined nuclei profiles from both nTg
and CCNA2-Tg mouse hearts revealed 16 major cardiac cell populations, including
cardiomyocytes, endothelial cells, fibroblasts, cardiac, immune cells, and neuronal cells,
distributed across 35 subclusters **(Figure 3C-E)**. Cell type clusters were annotated based on
expression of canonical marker genes and validated through the differential expression analysis
and comparison with a publicly available reference dataset (iRhythmics FairdomHub) [26, 27].
(Figure 3F, 3G).
236
237 We identified several common clusters between nTg and CCNA2-Tg mice **(Figure 3E, Figure**
238 **S3)**, with differential distribution across conditions   **(Figure 3H and 3I)**. Cardiomyocyte
239 populations were identified by the classical marker genes (*Myh7*, *Actn2,* and *Tnnc1*) in both nTg
240 and CCNA2-Tg hearts. Cardiomyocytes that express an enhanced level of contractility genes,
241 including the *Ryr2*, are annotated as mature cardiomyocytes [27].
242
243 SnRNA-seq identified 11 subclusters of the ventricular cardiomyocyte (vCM) population common
244 to both nTg and CCNA2-Tg hearts, with distinct differences in the proportion of nuclei across
245 subclusters. Notably, the vCM1 and vCM5 were more prevalent in nTg hearts, whereas the vCM2,
246 vCM3, vCM4, vCM6, vCM7, vCM8, vCM9, pro-vCM10, and the vCM11 were more abundant in
247 CCNA2-Tg hearts **(Figure 3J).**
248
Subclusters vCM1, vCM2, vCM4, vCM5, and vCM11 expressed cardiomyocyte maturation-
related genes, particularly sarcomere development genes, calcium (Ca^{2+}) handling, and cardiac
contractility genes (*Ttn*, *Pln, Ryr2*, and *Pkp2, Kcnd2*, and *Tnni3k),* regulating cardiomyogenesis
and function [28]. Notably, vCM5 also expressed non-cardiac lineage markers such as *Ebf2*,
implicated in brown adipocyte differentiation [29], and *Lsamp*, which guides neuronal connections
[30]. In contrast, vCM3 and vCM7 showed increased expression of ATP synthase genes (*Atp5b*
and *Atp5g3),* while vCM4 and vCM8 co-expressed endothelial-associated gene vCM9 was
uniquely marked by *Igfbp3*, consistent with a mid-differentiation state.
257
Moreover, the transcriptomic profile of the Pro-vCM10 displayed a robust cell cycle signature with
high expression of early-phase proliferative genes (*Top2a, Cdc20, Cdk2d, Cdk1, Prc1 Bub1,*
Mki67, and *Kif23)*, and mitotic-cytokinesis drivers (*Stmn1*) [31], alongside adhesion complex
assembly and immune markers (*Cd44, and C1qa*) [32]. This transcriptional profile indicates an
actively cycling, dedifferentiated cardiomyocyte population poised for regeneration. Remarkably,
the proportion of Pro-vCM10 was significantly higher in CCNA2-Tg mouse hearts compared to
nTg mouse hearts **(Figure 3J)**.
265
CCNA2 was expressed in cardiomyocytes, predominantly within the proliferative cluster identified
in CCNA2-Tg hearts, which also expressed key cell cycle regulators **(Figure 4A** and **4B)**. This
expression pattern remained consistent across the combined transcriptomic profiles of all
subclusters **(Figure S4)**. The pro-vCM subcluster in CCNA2-Tg mice co-expressed proliferative

markers (*Mki67* and *Kif23*) and exhibited reduced *Ryr2*, consistent with a less mature phenotype
(Figure 5A and **5B).** Compared to nTg cardiomyocytes, where *Ryr2* expression was higher,
CCNA2-Tg cardiomyocytes expressed a transcriptional signature consistent with a
dedifferentiated state that may facilitate cell cycle re-entry and regeneration [33].
274
275 Bulk RNA-sequencing further supported this profile, revealing coordinated downregulation of
276 sarcomere, contractility genes, and upregulation of proliferative programs in CCNA2-Tg
277 cardiomyocytes (**Figure S5A** and **S5B**), aligning with a dedifferentiated state, which facilitates
278 cell cycle re-entry and cytokinesis. This transcriptional 'immaturity' may enable the proliferative
279 potential necessary for myocardial regeneration.
280
To further examine CCNA2-induced cytokinesis events, we investigated whether distinct
cardiomyocyte subclusters expressed genes associated with cytokinesis and/or reprogramming
in CCNA2-Tg mice. Intriguingly, the pro-vCM population demonstrated upregulation of core
cytokinesis regulators such as *Aurkb, Anln, Sept7,* and *Rho,* among others, with *Rho* and *Aurka*
significantly elevated in CCNA2-Tg compared to nTg mice **(Figure 5C)**. This pro-vCM subcluster
also showed increased expression of reprogramming genes, including *Pou5f1, Klf4, Gata4, Myc,*
Sox2, and *Tbx5*, with *Pou5f1* and *Tbx5* notably increased in CCNA2-Tg hearts compared to nTg
hearts **(Figure 5D)**. These findings were consistent with bulk RNA-sequencing analysis, which
revealed increased expression of cytokinesis- and reprogramming-associated genes in CCNA2-
Tg cardiomyocytes compared to nTg cardiomyocytes **(Figure S5C** and **S5D)**.
291
Pathway enrichment analysis of CCNA2-Tg cardiomyocytes shows significant activation of cell
cycle, cytokinesis, and mitotic regulation pathways compared to nTg controls. Notably, Rho
GTPase signaling, spindle checkpoint regulation, and cytokinesis pathways were highly enriched
(Figure 5E), indicating increased proliferation and cytoskeletal remodeling in CCNA2-Tg hearts.
In contrast, nTg cardiomyocytes showed enrichment in extracellular matrix organization, collagen
biosynthesis, and muscle contraction pathways, reflecting a more differentiated state. These
findings indicate that CCNA2 enhances proliferation and cellular plasticity, promoting a
regenerative phenotype in cardiomyocytes.
300
Our bulk RNA-sequencing data from mouse nTg and CCNA2-Tg cardiomyocytes yielded
comparable findings. We observed upregulation of *Ccnb1* and *Ccnb2* in CCNA2-Tg
cardiomyocytes compared to nTg cardiomyocytes, along with slight increases in the expression
of *Cdk1* in CCNA2-Tg cardiomyocytes **(Figure 5F)**. *Cdk1* is the sole essential cyclin-dependent
kinase (*CDK*), when complexed to Cyclin A2 promotes entry into mitosis, whereas Cyclin A2/Cdk2
is critical for the G1/S transition [13, 34, 35]. Key cytokinesis and proliferation genes, including
Aurka, Aurkb, Anln, Mki67, and *Kif23,* were upregulated in CCNA2-Tg cardiomyocytes, reinforcing
the role of CCNA2 in cell cycle reactivation **(Figure 5F)**. Adult human hearts exhibited
downregulation of proliferative and cytokinesis-related genes **(Figure 5G, 5H)**, whereas CCNA2-
Tg cardiomyocytes retained a transcriptional profile favoring cell cycle progression and
cytokinesis.
312

Differential expression analysis further revealed that adult human hearts exhibit a greater number
of downregulated genes compared to the fetal heart (**Figure 5I),** consistent with widespread
transcriptional silencing during cardiomyocyte maturation. In contrast, CCNA2-Tg cardiomyocytes
displayed a higher number of upregulated genes relative to nTg controls **(Figure S5E)**, indicating
broad transcriptional reactivation. This inverse pattern supports the role of *CCNA2* in restoring a
fetal gene expression landscape.

Pathway enrichment analysis of these differentially expressed genes confirmed upregulation of
metabolism, Ca^{2+} mobilization, and ion homeostasis pathways in adult human heart, and
downregulation of pathways linked to mitotic progression and developmental gene programs
(Figure 5I). These findings support the hypothesis that CCNA2 reactivates a fetal-like
transcriptional program in adult cardiomyocytes, promoting cell cycle re-entry without complete
dedifferentiation.

Furthermore, pathway enrichment analysis revealed activation of key signaling pathways
associated with cardiomyocyte plasticity, proliferation, and reprogramming in CCNA2-Tg
cardiomyocytes. Specifically, the NOTCH signaling pathway and its downstream effectors
(HES/HEY) were upregulated, reinforcing CCNA2's role in progenitor-like gene activation and
cardiac reprogramming. Additionally, developmental pathways such as, TGF-beta regulation and
Wnt interactions suggest a shift toward a more plastic, regenerative state. Increased signaling
through MAPK, PDGF, and Jak-STAT pathways further supports enhanced proliferation and
cytoskeletal remodeling. Furthermore, upregulation of focal adhesion, integrin interactions, and
actin cytoskeleton regulation pathways indicate structural adaptations that facilitate
cardiomyocyte dedifferentiation and division. Conversely, mitochondrial metabolism and oxidative
phosphorylation, hallmarks of mature cardiomyocyte function, were significantly downregulated,
aligning with a metabolic reversion to a fetal-like, proliferative phenotype in CCNA2-Tg
cardiomyocytes. **(Figure S5F, S5G).** Moreover, adult CCNA2-Tg cardiomyocytes exhibit a greater
number of upregulated genes compared to nTg cardiomyocytes in mice **(Figure S5E)**.
339
340   In summary, our data highlight roles for *CCNA2* to enhance reprogramming and dedifferentiation,
341   which ultimately elicits cardiomyocyte cytokinesis. These results provide a compelling pathway
342   forward for the clinical development of cardiac regenerative therapy based on the manipulation of
343   CCNA2 expression in cardiomyocytes.
344

345   **Discussion**
346
Epimorphic regeneration is a conserved mechanism across evolutionary biology, permitting
organ-specific tissue regeneration in a variety of metazoan phylogeny, including early postnatal
mammals [36]. In fact, in human neonates, the ventricular myocardium responds to the pressure
overload of congenital aortic stenosis through hyperplastic growth, yet the degenerative, calcific
aortic stenosis of older patients elicits hypertrophic growth of the myocardium [37-39]. These
clinical observations imply that the cell cycle repertoire of human neonatal cardiomyocytes is also
intact, as cardiac growth is able to occur in a similar fashion to the embryonic heart. Moreover,
the adult human heart is also capable of a very low rate of turnover [8], with diminishing rates of

turnover past childhood and adolescence, with cardiomyocyte cytokinesis markers not found past age 20 [7].

Strategies to induce cardiomyocyte cytokinesis in adult human hearts are of critical need in order to face the growing public health crisis of congestive heart failure. To our knowledge, most cell therapy clinical trials in heart disease have provided inconsistent evidence of improvements in left ventricular ejection fraction, with a lack of evidence that the cell types utilized actually differentiate into functional cardiomyocytes. To this end, we previously explored the effects of cell cycle manipulation in small and large animal models and successfully induced cardiomyocyte proliferation *in vivo* with significant enhancement of cardiac function in animal models [19, 22]. In any translational plan to use cell cycle regulators for clinical use, two angles must be considered. Firstly, the wrong cell cycle regulators can induce cellular apoptosis [40-43] or be ineffective if the G2/M checkpoint is not navigated [13, 22, 44]. Secondly, precautions must be taken to prevent the expression of cell cycle activators in extra cardiac tissues due to the potential for oncogenic transformation [45]. This safety concern prompted us to explore methods of tissue-specific activation of cyclin A2. One such method was to design a next-generation viral vector in which a cardiomyocyte-specific promoter, cardiac Troponin T, drives the expression of human cyclin A2 (cTnT-hCCNA2). Human cardiomyocytes isolated from 21-, 41- and 55-year-olds who had died of noncardiac causes were cultured utilizing adult mammalian cardiomyocyte culture methods developed by our laboratory [22] and transduced with cTnT-hCCNA2. Cardiomyocytes from the 41- and 55-year-olds in the experimental wells underwent complete cytokinesis at a significantly higher frequency than the cardiomyocytes in the control wells. Based on our previous studies of rodent cardiomyocytes, we attribute the low rate of cytokinetic events we observed in the control wells to the reactivation of endogenous *CCNA2* with the prolonged culture of cardiomyocytes. This is a phenomenon we have previously observed and quantified in rodent cardiomyocytes, and likely not correlated with the low turnover seen in human hearts as measured by 14C dating [46] as cytokinesis markers were not noted at all in human hearts over the age 20 by Kühn and colleagues [7]. Similarly, the Frisén laboratory [46] used stereological and 14C quantification techniques to measure cardiomyocyte exchange. They found that most cardiomyocytes were never exchanged and that the substantial replacement of existing cardiomyocytes was generated in the first 10 years of life, while the second decade of life focused on DNA polyploidization.

It is noteworthy to mention that in the adult murine context, approximately 85–90% of cardiomyocytes exist in a binucleated state [47], indicating that the nuclear count does not precisely correspond to the cell count. The polyploidization of cardiomyocytes has been recognized as an impediment to heart regeneration [48]. It has been reported that binucleated cardiomyocytes re-entering the cell cycle are less likely to complete cytokinesis, thereby impeding the efficient regeneration of cardiac tissue following injury [36]. In adult mammalian hearts, the ratio of binucleated to mononucleated cardiomyocytes varies from one species to another. In rodent hearts, approximately 90% are binucleated while in human hearts, this percentage is markedly lower, varying from 25-60% [49]. Thus a greater percentage of mononuclear cardiomyocytes in the human heart would enable greater efficiency of CCNA2 gene therapy to induce cardiomyocyte proliferation as a repair strategy.

Our snRNA-Seq also confirms heightened cardiomyocyte plasticity of CCNA2 transgenic mice as
we note changes in gene expression involved in cell cycle progression and reprogramming. Ca^{2+}
signaling is a fundamental pathway in the regulation of cell division, cardiac contractility, and
remodeling [50-52]. The increase in the cytokinetic index in the test samples from 41- and 55-
year-olds shows the ability of *CCNA2* to stimulate intracellular Ca^{2+} mobilization. Of note, the
transduced cTnT-hCCNA2 (test samples) cultured cardiomyocytes isolated from a 21-year-old
male did not show a significant increase in cytokinetic index invoked by CCNA2 when compared
to the control samples, which is consistent with reports of previous investigators [7, 46] that cell
turnover and cell division can naturally occur at younger ages in humans.
408
SnRNA-seq has been used to identify transcriptomic changes occurring at the single-cell level,
shedding light on the precise mapping of transcriptomic alterations in a cell-type-specific manner
in response to cellular stimulation or gene delivery [53]. Employing snRNA-seq in intact tissues
enables the capturing of cell-specific transcriptomics changes in a more precise manner while
maintaining the integrity of tissue responses, compared to isolated *in vitro* transcriptomics
analyses. Our strategy of using snRNA-seq is to identify cardiomyocyte-specific changes in an *in*
vivo setting in a clinically relevant CCNA2-Tg model, which allowed us to map specific changes
to a subpopulation of cardiomyocytes. This would not have been captured by analyzing the
transcriptome of the bulk tissue of isolated cardiomyocytes using bulk RNA-sequencing
strategies. Live-cell imaging demonstrates that CCNA2-transfected adult human cardiomyocytes
successfully undergo cytokinesis, with daughter cells retaining what appear to be intact
sarcomeric structures. However, transcriptomic analyses (bulk RNA-sequencing and snRNA-seq)
revealed significant downregulation of sarcomere-related genes. These findings suggest a
dynamic remodeling process during cytokinesis, where transcriptional downregulation of
sarcomeric genes may occur transiently, while structural integrity is preserved through post-
transcriptional mechanisms or protein stability. This highlights the intricate balance between
cellular reprogramming and maintenance of cardiomyocyte identity during division.
426
Furthermore, the genes induced in the CCNA2-Tg model appear to represent a common
mechanism underlying cytokinesis in both male and female mice, consistent with observations in
human male and female hearts. Given that we identified a subset of cardiomyocytes that are
reprogrammed into a more proliferative state, we can envision targeting this subset in future
regenerative strategies. Unlike other regenerative approaches that separately target
cardiomyocyte proliferation or reprogramming, our findings suggest that *CCNA2* integrates both
processes, making it a unique therapeutic candidate for cardiac regeneration. Defining the
transcriptomic state of cyclin A2-expressing cardiomyocytes and their proliferative potential in
greater detail can enable us to recapitulate such proliferative cardiomyocytes both *in vitro* and *in*
vivo to determine whether targeting these specific subpopulations holds greater therapeutic
potential than targeting the entire cardiomyocyte population. The use of computational resources
can aid our comprehension of these molecular mechanisms for more precision-guided
regenerative strategies for human heart disease.
440
441 The ability of *CCNA2* to induce proliferation while maintaining cardiomyocyte function positions
442 *CCNA2* gene therapy as a promising clinical approach, minimizing risks associated with

exogenous cell transplantation, immune rejection, and off-target differentiation. Moreover, *CCNA2* appears to induce a transient mesenchymal-like state, as indicated by the upregulation of *Zeb1*, a key regulator of epithelial-to-mesenchymal transition (EMT). This suggests that CCNA2-Tg cardiomyocytes adopt a plastic phenotype to facilitate cell division, similar to mechanisms observed in neonatal heart regeneration. These findings highlight *CCNA2* as a novel and safe strategy for cardiac regeneration, offering a balance between proliferative potential and structural preservation. Ongoing studies by our laboratory have revealed silencing mechanims of endogenous CCNA2, with antisense approaches being utilized to reverse cell cycle exit and reactivate transcription of endogenous CCNA2. These approaches may offer alternative or complementary strategies to gene therapy with CCNA2.

Limitations

While the findings presented above are very promising, there are a few limitations to consider. Firstly, the use of freshly isolated human adult primary cardiomyocytes offers an ideal cellular model but presents several challenges. There is a high susceptibility to cell death in culture, which can hinder such experiments, and a fair amount of biological variability in getting the cells to adhere to the petri dishes. Adherence to the petri dishes is necessary to perform these cytokinesis experiments. Greater consistency with very rapid isolation from human cadavers and further refinements of the culture methods may overcome these challenges. Additionally, the use of adenovirus vectors is safe, and they are being utilized in other gene therapies that have been clinically tested. Immunogenicity is a concern, and therefore, there are ongoing investigations in our laboratory to explore alternative delivery strategies. Furthermore, *in vitro* models, though valuable, do not fully replicate the complexity of the *in vivo* cardiac environment. Nevertheless, this study is a valuable translational step forward from our previous preclinical report demonstrating that CCNA2 gene therapy induced a 55% increase in cardiomyocytes of the peri-infarct zones of the porcine heart [22] with significant enhancement of cardiac function. CCNA2-induced cytokinesis of human adult cardiomyocytes provides a compelling pathway toward human clinical trials of CCNA2 gene therapy for heart repair.

Methods
Culture of adult human cardiomyocytes and *in vitro* cytokinesis study
Cardiomyocytes from adult human (21-year-old male, 41-year-old female, and 55-year-old male) heart tissue were isolated after enzymatic digestion at Anabios, San Diego, CA, and were shipped to our laboratory within 24 hours of isolation. Adult human cardiomyocytes were cultured according to our previously published adult porcine cardiomyocyte culture technique [22] with slight modifications. In brief, upon arrival, cells were washed with serum-free Dulbecco's modified Eagle's medium (DMEM) media twice, and 10^5 cells were seeded in 100mm untreated polystyrene plates. Non-adherent cells were collected every 24 hours and centrifuged at 20Xg for 2 min at room temperature. Cell pellet was washed with serum-free DMEM and seeded on new polystyrene plates in modified Cardiomyocyte Culture Media (mod CMC) [22] formulated by adding 13% fetal bovine serum (FBS), 2.5% horse serum, 1X nonessential amino acid, 1mM sodium pyruvate, penicillin (100 U/ml), streptomycin (100 mg/ml), and fungizone (0.5 mg/ml) to DMEM/F12 (50:50). Cells were washed every day with serum-free DMEM, re-seeded in new

polystyrene plates and cultured for 3 days. On day 4, they were seeded in glass-bottom 24-well
tissue culture plates for 20 days, with the medium changed every 4th day. The wells with
cardiomyocytes exhibiting adhesion and spreading were selected. Cells in these wells were
trypsinized and counted, and 10^3 cells per well were seeded in new glass-bottom tissue culture
plates. After 2 days of culture, cells were divided into two groups (test and control) and were
transduced with adenoviruses. The test group was transduced with cTnT-hCCNA2 along with
cTnT-eGFP and/or cTnT-H2B-GFP and CMV-α-act-mCherry adenoviruses, while the control
group was transduced with only cTnT-eGFP and/or cTnT-H2B-GFP and CMV-α-act-mCherry
adenoviruses. MOI of adenoviruses were adjusted to 180 in each well of the test (with cTnT-
hCCNA2; MOI 100, CMV-α-act-mCherry; MOI 40, and cTnT-eGFP or cTnT-H2B-GFP; MOI 40)
and the control (cTnT-eGFP; MOI 140 and CMV-α-act-mCherry; MOI 40) group. After 48 hours
of incubation, transduction was confirmed by observing the desired fluorescence in live cell
imaging with Zeiss AxioVision Observer Z1 inverted microscope (Carl Zeiss). We also tested the
differentiation potential (Ca^{2+} flux) of cardiomyocytes after cytokinesis *in vitro* by culturing them in
an agarose-based semi-solid medium for 2 weeks. Cardiomyocytes were subjected to Ca^{2+} flux
imaging with pacing of 0.5 Hz and 1 Hz current.
503
Time-lapse microscopy
To capture cell division events in cardiomyocytes *in vitro*, we carried out live cell epifluorescence
time-lapse microscopy as described in [22]. In brief, Zeiss AxioVision Observer Z1 (Carl Zeiss,
Thornwood, NY, USA) inverted epifluorescence microscope in a humidified chamber in the
presence of 5% CO_2 at 37°C was used to carry out time-lapse microscopy. Multiple random
points with cells expressing eGFP (green) and mCherry (red) were selected in the test and control
groups. The positions were marked with the "position-list" tool in the AxioVision microscopy
software (AxioVision Release 4.7, Carl Zeiss). After the first cycle of imaging, only the channel for
Texas red was used (for detection of mCherry) to acquire images for 72 hours. The fluorescein
isothiocyanate (green) channel of the microscope was kept closed during the time-lapse imaging
to avoid cell death from exposure to ultraviolet rays in this channel. Images were taken at intervals
of 30 min. The 10X objective was used for all time-lapse imaging. Time-lapse movies were
generated after the end of each experiment and exported as MOV files. The time-lapse movies
were analyzed, and cells that underwent successful cytokinesis were enumerated in each group.
The percentage (%) of cytokinesis events was calculated for each position, and the graph was
plotted.
520
Cell fixation and nuclear staining
After time-lapse microscopy, cells in the glass-bottom plate were fixed with 4% paraformaldehyde
at room temperature for 20 min and were stored at 4°C. For nuclear staining, cells were washed
with 1XPBS then permeabilized with 0.5% Triton X-100 solution for 20 min at room temperature.
Cells were washed three times with 1XPBS and incubated in DAPI solution (2.5 µg/ml) for 5 min.
Cells were washed twice with 1X PBS, and imaging was carried out using a Zeiss AxioVision
Observer Z1 inverted epifluorescence microscope.
528
Real-time quantitative PCR

Quantitative PCRs were performed in cultured cardiomyocytes transduced with cTnT-hCCNA2
(test) or cTnT-eGFP (control) adenovirus with similar MOI (MOI 100). "SYBR Green quantitative
PCR protocol" on the "StepOnePlus" real-time PCR system (Applied Biosystems, CA) was used.
The PCR protocol consisted of one cycle at 95°C for 2 min, 39 cycles at 95°C for 15 sec + 60°C
for 1 min, and then 4°C on hold. Gene expression was determined by using the formula $2^{(39-\Delta\Delta CT)}$
$\Delta\Delta CT)$ with consideration of a CT value of 39 for a single transcript and with normalization to the
endogenous control *GAPDH*.
537

538　**Animal**
539　Cyclin A2 transgenic mice, adult males, and females 8-12 weeks old (n=4), were maintained in a
540　B6CBA background [13, 19]. Non-transgenic (wild-type) littermates were used as controls (n=4).
541

Collection of CCNA2-Tg mouse hearts and sample preparation for single-nucleus
RNA-sequencing (snRNA-seq)
Frozen heart tissues from CCNA2-Tg- and nTg adult mice, male and female, were minced and
transferred into a Dounce homogenizer containing ice-cold lysis buffer (Benthos Prime Central,
TX, USA). Samples were homogenized gently, and nuclei were isolated. Nuclei were ensured to
be free of clumps and debris by trypan blue staining under a microscope. Nuclei were then
counted, and concentration was adjusted. 10x library concentration, insert size, and quantification
were checked using Qubit, Agilent bioanalyzer, and q-PCR, respectively. For data Processing
and Quality Control of 10x Sequencing of snRNA-seq, the reads were processed with 10X
Genomics CellRanger software (v.3.1.0) with the default parameters for each sample. Cell Ranger
employs the STAR aligner for splicing-aware mapping of reads to the genome. It utilizes a
transcript annotation GTF file to categorize the reads into exonic, intronic, and intergenic regions
based on whether the reads align confidently with the genome. Only confidently mapped, non-
PCR duplicates with valid barcodes and UMIs were used to generate a gene-barcode matrix. This
involved various steps, including alignment to a reference, collapsing of unique molecular
identifiers (UMIs), UMI counting, and initial quality control procedures. The output of this process
was the generation of filtered gene expression matrices that exclusively contained cellular
barcodes. Subsequently, the Seurat R package (v.3.1.1) was used for data quality control and
downstream processing. The filtered_feature_bc_matrix generated by Cell Ranger was used for
Seurat processing. Post-sequencing quality control was performed using Seurat. Cells were
excluded from downstream processing in each sample if they met any of the following criteria:
fewer than 200 genes were detected per cell, genes had non-zero counts in at most three cells,
total feature counts exceeded 8000 (suggesting potential multiplets), more than 50% of the
feature count was attributable to mitochondrial genes, or more than 5% of the feature count was
attributable to hemoglobin-related genes, and doublets from all the clusters were filtered during
QC using DoubletFinder version 2.0.3 **(Figure S2)**. Raw read counts were log-normalized, cells
were clustered, and markers for each cluster were identified by using the Seurat FindMarkers and
FindAllMarker functions [54]. The cell clusters were annotated by using iRhythmics FairdomHub
instance (https://fairdomhub.org/studies/739) and prior knowledge.
571

Human fetal and adult hearts and RNA sequencing

Human fetal heart RNA (n=3 pooled, Cat #636532, Takara) and human adult heart RNA (n=4
pooled, Cat #636583, Takara) were used for bulk RNA sequencing with an increased read depth
(Ultra-deep sequencing), generating approximately 300 million reads per sample. This high
sequencing depth enabled the detection of rare transcripts, improved quantification accuracy, and
enhanced the resolution of transcript isoforms, including novel and unannotated RNAs. For library
preparation, we employed a ribosomal RNA (rRNA) depletion strategy instead of poly(A)
enrichment to capture both polyadenylated (polyA$^+$) and non-polyadenylated (polyA$^-$) RNAs.
While polyA$^+$ selection primarily enriches for mRNAs and some lncRNAs, rRNA depletion allows
for the inclusion of a broader spectrum of regulatory RNAs, including enhancer RNAs (eRNAs),
small nuclear RNAs (snRNAs), and uncharacterized lncRNAs that may lack poly(A) tails.
Additionally, this approach reduces transcript bias, preserves isoform diversity, and is better
suited for degraded RNA samples.
By combining rRNA depletion with ultra-deep sequencing, we achieved a comprehensive and
unbiased profile of the cardiac transcriptome. The sequence reads were mapped to the human
genome (GRCh38.110) using STAR [55] (version 2.7.9a). Using the mapped reads, the R
package edgeR [56] (version 3.30.3) was employed to calculate counts per million (CPM) values
and false discovery rate (FDR)-adjusted p-values obtained through the Benjamini–Hochberg
method (Benthos Prime Central, TX, USA).
591
Pathway enrichment analysis
Pathway enrichment analysis was performed using the Reactome pathway analysis tool [57]. The
differentially expressed genes among nTg and CCNA2-Tg conditions (log2FC > / < 1 and a P
value < 0.05) were used for pathway analysis. Fold enrichment was calculated to indicate the
gene expression observed in our list compared to the expected list. Categories with a fold
enrichment greater than 1 were considered overrepresented, while those with a fold enrichment
less than 1 were considered underrepresented in our experiment. Additionally, pathway
enrichment analysis was conducted using BioPlanet via the EnrichR suite [58], following a
previously described approach [59].
601
Statistical analysis
Statistical analyses were carried out using the R package (version 3.30.3), GraphPad Prism
(version 8.2.1, GraphPad), and Python (via Google Colab). Two-tailed unpaired t-tests or one-
/two-way analysis of variance (ANOVA) were used for normally distributed data. Chi-square tests
for proportions were performed using Python libraries, including *Scipy*. Data are presented as
means ± SEM unless otherwise indicated. Statistical significance was defined as P < 0.05.
608
Study approval
This study was conducted in accordance with the Declaration of Helsinki. Adult human
cardiomyocytes were obtained from Anabios (San Diego, CA) and were fully de-identified, and
therefore are exempt from ethics approval. For the mouse study, the protocol was approved by
the Institutional Animal Care and Use Committee (IACUC) of Mount Sinai Hospital, and all
procedures adhered to Institutional Animal Care and Use Guidelines.
615
616 **Data availability**

617 The processed single-nucleus data and the Deep bulk RNA sequencing, including raw expression
618 matrices and raw sequence files that support this research study's findings, are available on the
619 Gene Expression Omnibus GSE249433 and GSE256519, respectively.
620
621

Article Information
Author Contributions
E.B. wrote the original article, contributed to experiments, and analyzed and performed snRNA-
seq data with input from P.M. and A.K.; S.V.M and A.K.R contributed to cytokinesis experiments
and calcium measurements; P.M. contributed to processing snRNA-seq samples and initial
analysis; C.D.S. assisted with the cell biology experiments and organization of the manuscript;
P.E.M., A.G., N.A.G. provided cardiomyocytes through their proprietary technology. H.W.C. was
responsible for conceptualization, vector design, validation, supervision, reviewing, and funding
acquisition. Review & editing- E.B, H.W.C, P.M., and A.K.
631

632 **Acknowledgment**
633 We thank Dr. Bernhard Kühn for his contribution to the cTnT-H2B-GFP utilized in this study.
634

635 **Sources of funding**
636 This work is supported by the NIH R01 HL150345, the NYSTEM C029565, and NYSTEM
637 C32608GG.
638

639 **Disclosures**
640 H.W.C. is the inventor of multiple patents regarding cyclin A2-mediated cardiac repair and caudal-
641 related homeobox 2 cells for cardiac repair. She is in the midst of launching a new biotechnology
642 company to advance such therapies to clinical development. Other authors report no conflict.
643

Nonstandard Abbreviations and Acronyms	
Actn2	Actinin Alpha 2
Atp2a2	ATPase, SERCA2
Atp5b	ATP Synthase, Mitochondrial F1 Complex, Beta Polypeptide ATP
Atp5g3	Synthase, Mitochondrial C Subunit, Gamma Polypeptide 3
Bub1	BUB1 Mitotic Checkpoint Serine/Threonine Kinase
C1qa	Complement C1q A Chain
Ca^{2+}	Calcium ion
Calm1	Calmodulin 1
Calm2	Calmodulin 2
CCNA2	Cyclin A2
CCNA2-Tg	CCNA2- transgenic mice
Cdc20	Cell Division Cycle 20
Cdk1	Cyclin-Dependent Kinase 1
Cdk2	Cyclin-Dependent Kinase 2
CGC	Cardiac glial cells

CMs	Cardiomyocytes
CPM	Counts per million
CTnT	Cardiac Troponin T
CTnT-hCCNA2	Cardiac Troponin T- human cyclin A2
E1/E2	Early region 1/Early region 2
EC	Endothelial cells
EC-derived FB	Endothelial-derived fibroblasts
EDC	Endocardial cells
Efnb2	Ephrin B2
eGFP	Enhanced Green Fluorescent Protein
EPC	Epicardial cells
EP-derived FB	Epicardial derived fibroblasts
ES	Embryonic stem cells
FB	Fibroblasts
FDR	False discovery rate
Gata4	GATA Binding Protein 4
H2B	Histone 2B
Igfbp3	Insulin-Like Growth Factor Binding Protein 3
iPS	Induced pluripotent
Kcnd2	Potassium Voltage-Gated Channel Subfamily D Member 2
Kif23	Kinesin Family Member 23
Klf4	Kruppel Like Factor 4
MI	Myocardial infarction
Mki67	Marker of Proliferation Ki-67
MOI	Multiplicity of Infection
Mono	Monocytes
Myc	MYC Proto-Oncogene, BHLH Transcription Factor
Myh7	Myosin Heavy Chain 7
Myl2	Myosin Light Chain 2
Myob	Myoblasts
NC	Neuronal cells
Neb	Nebulin
nTg	Non-transgenic mice
PEC	Pericytes
Pkp2	Plakophilin 2
Pln	Phospholamban
Pou5f1	POU Class 5 Homeobox 1, Oct4
Prc1	Protein Regulator of Cytokinesis 1
Pro-vCM	Proliferative ventricular cardiomyocyte
Ryr2	Ryanodine Receptor 2
Scn5a	Sodium Voltage-Gated Channel Alpha Subunit 5
SMC	Smooth muscle cells
SnRNA-seq	Single nucleus RNA sequencing
TC	T-Cells

Tnnc1	Troponin C1
Tnni3k	Troponin I Interacting Kinase
Top2a	Topoisomerase (DNA) II Alpha
t-SNE	T-distributed Stochastic Neighbor Embedding
Ttn	Titin
UMI	Unique molecular identifier
UnkC	Unknown cells
vCM	Ventricular cardiomyocyte
VEC	Vascular endothelial cells
α-act	Alpha-actinin

644
645

References

1.   Yoshizumi, M., et al., *Disappearance of cyclin A correlates with permanent withdrawal of*
cardiomyocytes from the cell cycle in human and rat hearts. J Clin Invest, 1995. **95**(5): p.
2275-80.
2.   Miyamoto, K., et al., *Direct In Vivo Reprogramming with Sendai Virus Vectors Improves*
Cardiac Function after Myocardial Infarction. Cell Stem Cell, 2018. **22**(1): p. 91-103 e5.
3.   Doppler, S.A., et al., *Mammalian Heart Regeneration: The Race to the Finish Line.* Circ
Res, 2017. **120**(4): p. 630-632.
4.   Yoshida, Y. and S. Yamanaka, *Induced Pluripotent Stem Cells 10 Years Later: For*
Cardiac Applications. Circ Res, 2017. **120**(12): p. 1958-1968.
5.   Shiba, Y., et al., *Allogeneic transplantation of iPS cell-derived cardiomyocytes regenerates*
primate hearts. Nature, 2016. **538**(7625): p. 388-391.
6.   Chong, J.J., et al., *Human embryonic-stem-cell-derived cardiomyocytes regenerate non-*
human primate hearts. Nature, 2014. **510**(7504): p. 273-7.
7.   Mollova, M., et al., *Cardiomyocyte proliferation contributes to heart growth in young*
humans. Proc Natl Acad Sci U S A, 2013. **110**(4): p. 1446-51.
8.   Bergmann, O., et al., *Evidence for cardiomyocyte renewal in humans.* Science, 2009.
324(5923): p. 98-102.
9.   Brockes, J.P. and A. Kumar, *Plasticity and reprogramming of differentiated cells in*
amphibian regeneration. Nat Rev Mol Cell Biol, 2002. **3**(8): p. 566-74.
10. Poss, K.D., L.G. Wilson, and M.T. Keating, *Heart regeneration in zebrafish.* Science,
2002. **298**(5601): p. 2188-90.
11. Jopling, C., et al., *Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation*
and proliferation. Nature, 2010. **464**(7288): p. 606-9.
12. Furtado, M.B., et al., *View from the heart: cardiac fibroblasts in development, scarring*
and regeneration. Development, 2016. **143**(3): p. 387-97.
13. Chaudhry, H.W., et al., *Cyclin A2 mediates cardiomyocyte mitosis in the postmitotic*
myocardium. J Biol Chem, 2004. **279**(34): p. 35858-66.
14. Nurse, P., *A long twentieth century of the cell cycle and beyond.* Cell, 2000. **100**(1): p. 71-
8.
15. Pagano, M., et al., *Cyclin A is required at two points in the human cell cycle.* EMBO J,
1992. **11**(3): p. 961-71.

16.   Sherr, C.J. and J.M. Roberts, *Inhibitors of mammalian G1 cyclin-dependent kinases.* Genes
Dev, 1995. **9**(10): p. 1149-63.

17.   Wang, B.J. and H.W. Chaudhry, *Regulation of Cardiomyocyte Division During Cardiac*
Regeneration. Curr Cardiol Rep, 2023. **25**(7): p. 615-620.

18.   Vadakke-Madathil, S. and H.W. Chaudhry, *Cardiac Regeneration: Time to Revisit Nature.*
Circ Res, 2018. **123**(1): p. 24-26.

19.   Cheng, R.K., et al., *Cyclin A2 induces cardiac regeneration after myocardial infarction*
and prevents heart failure. Circ Res, 2007. **100**(12): p. 1741-8.

20.   Woo, Y.J., et al., *Therapeutic delivery of cyclin A2 induces myocardial regeneration and*
enhances cardiac function in ischemic heart failure. Circulation, 2006. **114**(1 Suppl): p.
I206-13.

21.   Woo, Y.J., et al., *Myocardial regeneration therapy for ischemic cardiomyopathy with*
cyclin A2. J Thorac Cardiovasc Surg, 2007. **133**(4): p. 927-33.

22.   Shapiro, S.D., et al., *Cyclin A2 induces cardiac regeneration after myocardial infarction*
through cytokinesis of adult cardiomyocytes. Sci Transl Med, 2014. **6**(224): p. 224ra27.

23.   Litvinukova, M., et al., *Cells of the adult human heart.* Nature, 2020. **588**(7838): p. 466-
472.

24.   Bak, S.T., et al., *Ploidy-stratified single cardiomyocyte transcriptomics map Zinc Finger*
E-Box Binding Homeobox 1 to underly cardiomyocyte proliferation before birth. Basic Res
Cardiol, 2023. **118**(1): p. 8.

25.   So, J., et al., *Robust single-nucleus RNA sequencing reveals depot-specific cell population*
dynamics in adipose tissue remodeling during obesity. Elife, 2025. **13**.

26.   Galow, A.M., et al., *Quality control in scRNA-Seq can discriminate pacemaker cells: the*
mtRNA bias. Cell Mol Life Sci, 2021. **78**(19-20): p. 6585-6592.

27.   Galow, A.M., et al., *Integrative Cluster Analysis of Whole Hearts Reveals Proliferative*
Cardiomyocytes in Adult Mice. Cells, 2020. **9**(5).

28.   Lai, Z.F., et al., *Overexpression of TNNI3K, a cardiac-specific MAP kinase, promotes*
P19CL6-derived cardiac myogenesis and prevents myocardial infarction-induced injury.
Am J Physiol Heart Circ Physiol, 2008. **295**(2): p. H708-16.

29.   Rajakumari, S., et al., *EBF2 determines and maintains brown adipocyte identity.* Cell
Metab, 2013. **17**(4): p. 562-74.

30.   Wang, L., et al., *Polymorphisms of the tumor suppressor gene LSAMP are associated with*
left main coronary artery disease. Ann Hum Genet, 2008. **72**(Pt 4): p. 443-53.

31.   Liu, J., et al., *Aberrantly high activation of a FoxM1-STMN1 axis contributes to*
progression and tumorigenesis in FoxM1-driven cancers. Signal Transduct Target Ther,
2021. **6**(1): p. 42.

32.   Zhao, L., et al., *CD44 regulates vascular gene expression in a proatherogenic*
environment. Arterioscler Thromb Vasc Biol, 2007. **27**(4): p. 886-92.

33.   Beisaw, A. and C.C. Wu, *Cardiomyocyte maturation and its reversal during cardiac*
regeneration. Dev Dyn, 2024. **253**(1): p. 8-27.

34.   Xing, Z., et al., *Expression and prognostic value of CDK1, CCNA2, and CCNB1 gene*
clusters in human breast cancer. J Int Med Res, 2021. **49**(4): p. 300060520980647.

35.   Santamaria, D., et al., *Cdk1 is sufficient to drive the mammalian cell cycle.* Nature, 2007.
448(7155): p. 811-5.

36.   Bersell, K., et al., *Neuregulin1/ErbB4 signaling induces cardiomyocyte proliferation and*
repair of heart injury. Cell, 2009. **138**(2): p. 257-70.

37.   Leeuwenburgh, B.P., et al., *Chronic right ventricular pressure overload results in a*
725   *hyperplastic rather than a hypertrophic myocardial response.* J Anat, 2008. **212**(3): p. 286-
726   94.

727   38.   Assey, M.E., et al., *Unexpected persistence into adulthood of low wall stress in patients*
728   *with congenital aortic stenosis: is there a fundamental difference in the hypertrophic*
729   *response to a pressure overload present from birth?* Circulation, 1987. **75**(5): p. 973-9.

730   39.   Torp, K.H., *Congenital aortic stenosis with hyperplasia of the endocardium in the left*
731   *ventricle.* Acta Pathol Microbiol Scand, 1951. **29**(2): p. 109-19.

732   40.   Kranenburg, O., A.J. van der Eb, and A. Zantema, *Cyclin D1 is an essential mediator of*
733   *apoptotic neuronal cell death.* EMBO J, 1996. **15**(1): p. 46-54.

734   41.   Freeman, R.S., S. Estus, and E.M. Johnson, Jr., *Analysis of cell cycle-related gene*
735   *expression in postmitotic neurons: selective induction of Cyclin D1 during programmed*
736   *cell death.* Neuron, 1994. **12**(2): p. 343-55.

737   42.   Sofer-Levi, Y. and D. Resnitzky, *Apoptosis induced by ectopic expression of cyclin D1 but*
738   *not cyclin E.* Oncogene, 1996. **13**(11): p. 2431-7.

739   43.   Kong, G., et al., *Functional analysis of cyclin D2 and p27(Kip1) in cyclin D2 transgenic*
740   *mouse mammary gland during development.* Oncogene, 2002. **21**(47): p. 7214-25.

741   44.   Liu, Y. and R.N. Kitsis, *Induction of DNA synthesis and apoptosis in cardiac myocytes by*
742   *E1A oncoprotein.* J Cell Biol, 1996. **133**(2): p. 325-34.

743   45.   Wang, J., et al., *Hepatitis B virus integration in a cyclin A gene in a hepatocellular*
744   *carcinoma.* Nature, 1990. **343**(6258): p. 555-7.

745   46.   Bergmann, O., et al., *Dynamics of Cell Generation and Turnover in the Human Heart.* Cell,
746   2015. **161**(7): p. 1566-75.

747   47.   Soonpaa, M.H., et al., *Cardiomyocyte DNA synthesis and binucleation during murine*
748   *development.* Am J Physiol, 1996. **271**(5 Pt 2): p. H2183-9.

749   48.   Gonzalez-Rosa, J.M., et al., *Myocardial Polyploidization Creates a Barrier to Heart*
750   *Regeneration in Zebrafish.* Dev Cell, 2018. **44**(4): p. 433-446 e7.

751   49.   Botting, K.J., et al., *Early origins of heart disease: low birth weight and determinants of*
752   *cardiomyocyte endowment.* Clin Exp Pharmacol Physiol, 2012. **39**(9): p. 814-23.

753   50.   Patergnani, S., et al., *Various Aspects of Calcium Signaling in the Regulation of Apoptosis,*
754   *Autophagy, Cell Proliferation, and Cancer.* Int J Mol Sci, 2020. **21**(21).

755   51.   Rosendo-Pineda, M.J., C.M. Moreno, and L. Vaca, *Role of ion channels during cell*
756   *division.* Cell Calcium, 2020. **91**: p. 102258.

757   52.   Liu, C., et al., *Alteration of calcium signalling in cardiomyocyte induced by simulated*
758   *microgravity and hypergravity.* Cell Prolif, 2020. **53**(3): p. e12783.

759   53.   Bode, D., et al., *Exploiting Single-Cell Tools in Gene and Cell Therapy.* Front Immunol,
760   2021. **12**: p. 702636.

761   54.   Koenig, A.L., et al., *Single-cell transcriptomics reveals cell-type-specific diversification in*
762   *human heart failure.* Nat Cardiovasc Res, 2022. **1**(3): p. 263-280.

763   55.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1):
764   p. 15-21.

765   56.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for*
766   *differential expression analysis of digital gene expression data.* Bioinformatics, 2010.
767   **26**(1): p. 139-40.

768   57.   Fabregat, A., et al., *Reactome pathway analysis: a high-performance in-memory approach.*
769   BMC Bioinformatics, 2017. **18**(1): p. 142.

770   58.   Xie, Z., et al., *Gene Set Knowledge Discovery with Enrichr.* Curr Protoc, 2021. **1**(3): p.
771   e90.
772   59.   Huang, R., et al., *The NCATS BioPlanet - An Integrated Platform for Exploring the*
773   *Universe of Cellular Signaling Pathways for Toxicology, Systems Biology, and Chemical*
774   *Genomics.* Front Pharmacol, 2019. **10**: p. 445.
775
776

778
Figure 1. CCNA2 expression induces cytokinesis in adult human cardiomyocytes *In Vitro*.
A) Clinical-use grade replication-deficient human adenovirus type 5 (Ad5) was used for this study.
The vector was designed with cardiac Troponin T (cTnT) promoter to express human cyclin A2
(hCCNA2) specifically in cardiomyocytes. **B)** The cTnT-hCCNA2 adenovirus was used to
transduce cultured adult human cardiomyocytes to induce the expression of CCNA2. Significantly
higher CCNA2 expression was observed in cells transduced with cTnT-hCCNA2 compared to the
control virus. **C-H)** Still images from representative time-lapse epifluorescence microscopy of
cultured cardiomyocytes isolated from adult humans **C**) 55-year-old and **F)** 41-year-old. Please
note that still images of 55 and 41-years-old have scale bars of 50µm and 100µm as it is shown in
the movies S1 and S2 respectively. Cells were transduced with cTnT-hCCNA2, cTnT-H2B-GFP,
cTnT-eGFP (to label cardiomyocytes), and CMV-α-actinin-m-Cherry adenoviruses prior to the
beginning of the time-lapse imaging. **C)** At time 0 hr, a cell (yellow) expressing both cTnT (green)
and α-actinin (red) was followed for 70 hrs via time-lapse microscopy. The green channel was
closed, and cells were only followed through the red channel to avoid UV photo-toxicity. The
observed human cardiomyocytes show the 1st cell division at 50 hrs of imaging, and one of the
daughter cells subsequently undergoes division at 70 hrs of imaging. **D** and **G)** At 70/72 hrs, a
daughter cell is partially magnified with a grayscale version to demonstrate the presence of an

intact sarcomere structure. **E, H)** 1 day after live imaging ended, these wells were fixed with
subsequent labeling of nuclei with DAPI. The green fluorescence of the original cTnT-eGFP and
cTnT-H2B-GFP transduction are visible, further confirming that these cells are cardiomyocytes. **I)**
The cytokinetic events were counted in the control and test samples from the 21-, 41-, and 55-
year-old subjects. A significantly higher rate of cytokinesis was observed in the test samples from
the 41-year-old (p=0.001) and 55-year-old (p<0.0001) compared to the control. On the other hand,
the control and test samples from the 21-year-old did not exhibit a significant difference in
cytokinetic events (p=0.47).
804

806
807
Figure 2. Differentiation of cultured adult human cardiomyocytes. Cultured adult human
cardiomyocytes typically undergo dedifferentiation and were then subjected to 3D redifferentiation
culture conditions and examined for Ca²⁺ flux. **A)** Overview of the 3D culture method of
differentiation, Ca²⁺ flux measurement, and fluorescence microscopy in adult human
cardiomyocytes. **B)** Representative Ca²⁺ flux graphs of re-differentiated cultured adult human
cardiomyocytes. Ad5-cTnT-hCCNA2 transduced adult human cardiomyocytes show Ca²⁺ flux with
pacing of 0.5 and 1 Hz of current, while no Ca²⁺ flux was detected in the non-Ad5-cTnT-hCCNA2
transduced cardiomyocytes. **C)** Venn diagram showing the overlap of differentially expressed
genes between CCNA2-Tg versus non-nTg mouse cardiomyocytes and human fetal versus adult
hearts. **D)** Venn diagram displaying the subset of downregulated genes shared between the
CCNA2-Tg mouse hearts and the fetal human heart. These genes include several associated
with cardiomyocyte maturation. **E)** Heatmap depicting the expression of Ca²⁺ handling,
contractility, and sarcomere-associated genes in human adult versus fetal hearts, **F)** Comparison
of gene expression in cardiomyocytes from CCNA2-Tg versus nTg mice, highlighting differences
in Ca²⁺ regulatory genes and contractile machinery. Positive values indicate upregulation in adult
hearts relative to fetal hearts. Data represent normalized RNA-sequencing expression values.
824

825
826
Figure 3. Single-nucleus RNA sequencing identified major cell types in CCNA2-Tg mice
A) Schematic depicting the design of the snRNA-seq experiments (n=2) for each condition nTg
female, CCNA2-Tg female, nTg male, and nTg male (Total: n=8). **B)** The analysis pipeline and
single-cell barcoded library generation (10X Genomics), sequence alignment (Cell Ranger), and
further analysis using R (SeuratR package). **C)** t-distributed Stochastic Neighbor Embedding (t-
SNE) clustering the nuclei reveals more than 16 distinct major cell types identified based on the
cell-specific markers dataset provided at iRhythmics FairdomHub instance, and **D)** the pie chart
showing the proportion of cells within the snRNA-seq. **E)** t-SNE of a total of 35 subclusters. **F)**
Dot plot depicting the expression levels of some of the key marker genes used to classify cell
types. Dot size represents the percentage of cells expressing each gene, while color intensity
reflects expression levels. **G)** Feature plots highlighting the expression of selected cell-type-
specific markers across the t-SNE space. **H)** Stacked bar plots representing the proportion of
nuclei classified into each major cell type (top) and **I)** each identified cluster (bottom) in both
CCNA2-Tg and nTg mice. **J)** Stacked bar plots displaying the proportion of nuclei in vCM clusters

(left). The absolute count comparison (right) highlights a significant increase in Pro-vCM10 in
CCNA2-Tg hearts relative to nTg controls (****p < 0.0001, Chi-square test), supporting the role of
CCNA2 in promoting cardiomyocyte proliferation. CM: Cardiomyocytes, vCM: ventricular
cardiomyocytes, EC: endothelial cells, FB: fibroblasts, EC-derived FB: endothelial-derived
fibroblasts, NC: neuronal, EDC: endocardial cells. PEC: pericytes, VEC: vascular endothelial
cells, Mono: monocytes, TC: T-Cells, SMC: smooth muscle cells, EPC: epicardial cells, UnkC:
unknown cells, EP-derived FB: epicardial derived fibroblasts, CGC: Cardiac glial cells, Myob:
myoblasts.
849
850
851

852
Figure 4. CCNA2 and cell cycle genes expression in CCNA2-Tg and nTg mice
cardiomyocytes delineated by snRNA-seq. A) CCNA2 was expressed in the vCM, specifically

in the pro-vCM subset of the cardiomyocyte population in CCNA2-Tg and cell cycle genes
projected on the t-SNE plot. **B)** Scatter plots showing the expression level of CCNA2 and cell
cycle genes in cardiomyocytes of CCNA2-Tg and nTg mice. Each point represents an individual
value; the mean value is represented by the horizontal line. Error bars represent SEM.

861
862
863
Figure 5. CCNA2 expression induces proliferation, cytokinesis, and reprogramming in
mouse cardiomyocytes. A) t-SNE projection showing *Ryr2* expression across cardiomyocyte
subpopulations in nTg and CCNA2-Tg mice. *Ryr2*+ cardiomyocytes correspond to a more mature
ventricular cardiomyocyte (vCM) state, whereas *Ryr2* low expression cardiomyocytes represent
less mature, or proliferative cardiomyocytes. Quantification of *Ryr2* expression levels among
cardiomyocyte subpopulations is shown in the accompanying violin plot, highlighting differences
in *Ryr2* expression across conditions. **B)** t-SNE plots and representative scatter plots and Violin
plots representing the expression of key proliferation, **C)** cytokinesis, and **D)** reprogramming gene
markers in both CCNA2-Tg and nTg mice. **E)** Representative dot plot of the differential gene
expression analysis showing the upregulation of proliferation, cytokinesis, and reprogramming
genes in CCNA2-Tg versus nTg cardiomyocytes. **F)** Heatmaps of log-transformed gene
expression changes in human fetal versus adult hearts, **G)** and in CCNA2-Tg versus nTg
cardiomyocytes. **H)** Venn diagram showing shared upregulated genes between CCNA2-

transgenic mouse hearts and human fetal hearts. The diagram displays the overlap of upregulated
genes from CCNA2-Tg versus non-nTg mouse cardiomyocytes and human fetal versus adult
heart RNA-seq datasets. These shared genes are further associated with cell cycle, cytokinesis,
and cardiac development, supporting a dedifferentiation and reprogramming phenotype induced
by CCNA2. **I)** Pathway enrichment analysis showing significantly enriched pathways in human
adult (top) and fetal (bottom) heart. In the adult heart, pathways related to metabolism and energy
are enriched, based on 4,348 downregulated genes and 3,713 upregulated genes ($P<0.05$). In
the fetal heart, cell cycle, DNA/RNA processing, and signaling pathways are enriched ($P<0.05$).
885

Figure 6. Translational roadmap and mechanistic model of CCNA2 gene therapy for cardiac regeneration. Building upon our previous *in vivo* studies in rodent and porcine myocardial infarction models, which demonstrated functional recovery via CCNA2 gene therapy (1-3) [19-22], this study provides direct evidence of complete cytokinesis in cultured human adult cardiomyocytes. Complementary transcriptomic analyses from fetal and adult human hearts, as well as CCNA2 transgenic models, reveal reactivation of fetal-like, proliferative, and reprogramming gene programs. The schematic illustrates the translational trajectory from preclinical to mechanistic insights and highlights CCNA2 as a driver of cardiomyocyte cell cycle reentry, cytokinesis, and regenerative gene expression.

Supplemental Material

 Figure S1. A) Representative scatter plots of sarcomere assembly genes *(Neb, Tnn, Tnnc1*
Actn2, and *Tnnt2),* and **B)** Ca²⁺ handling genes (*Ryr2, Atp2a2, Slc8a1*, and others) in nTg and
CCNA2-Tg cardiomyocytes as delineated from bulk RNA seq. Each point represents an individual
value; the mean value is represented by the horizontal line. Error bars represent SEM, with
statistical significance indicated (*P*-values). ns denotes non-significant differences. **C)** Heatmap
of log-normalized expression of mesenchymal transition markers (*Fn1, Vim, Zeb1, Snai1, and*
Twist1) in nTg and CCNA2-Tg cardiomyocytes, as well as **D)** human adult and fetal hearts. Gene
expression levels are log-transformed and normalized.

913
Figure S2. Quality control (QC) filtering and visualization in SnRNA-seq. **A)** "nFeature_RNA"
indicates the count of gene features; "nCount_RNA" represents the count of UMIs; "percent.MT"
signifies the proportion of counts originating from mitochondrial genes, and "percent.HB" signifies
the proportion of counts stemming from Hemoglobin-related genes in all conditions nTg and
CCNA2-Tg males and females. Each data point corresponds to a cell. **B)** The UMAP plots are
generated from the top principal components within the clustered dataset. Each data point
represents an individual cell and is color-coded based on whether it is categorized as a singlet or
doublet.
922
923

924
Figure S3. A) t-SNE plot clustering the nuclei of nTg and CCNA2-Tg conditions (n=8 total). **B)** 925
Pie chart representing the percentage of nuclei profile in all conditions of nTg and CCNA2-Tg 926
mice. 927
928

929
930 **Figure S4. A)** t-SNE plots and **B)** representative scatter plots of CCNA2 and cell cycle genes
931 expression across the combined transcriptomic profiles of all subclusters of nTg and CCNA2-Tg
932 mice. Bars represent mean ± SEM. Each point represents an individual value; the mean value
933 is represented by the horizontal line. Error bars represent SEM.

934
935

936
Figure S5. Representative scatter plots of **A)** cardiomyocytes maturation gene *Ryr2, B)*
proliferation, **C)** cytokinesis, and **D)** reprogramming gene expression in nTg and CCNA2-Tg
cardiomyocytes. Each dot represents an individual value, and bars indicate mean ± SEM. **E)**
Number of differentially expressed genes (Up or Down) in adult CCNA2-Tg mouse
cardiomyocytes as compared to adult nTg mouse cardiomyocytes. **F)** Significantly enriched
pathways in adult CCNA2-Tg and nTg mouse cardiomyocytes (P-value adjusted <0.05), analyzed
using BioPlanet. **G)** Gene set enrichment analysis (GSEA) in CCNA2-transgenic versus nTg
cardiomyocytesshowing a significant downregulation of oxidative phosphorylation and
mitochondrial respiratory chain complex assembly pathways in CCNA2-transgenic
cardiomyocytes, may suggest a metabolic shift consistent with dedifferentiation. Conversely,
pathways such as cardiac epithelial-to-mesenchymal transition (EMT) and Notch signaling were
upregulated, indicating activation of developmental and reprogramming pathways that facilitate a
more regenerative state.

969   **Movie S1.** The real-time, live-imaging movie of human adult cardiomyocytes (55-year-old male)
970   as described in **Figure 1.**
971   **Movie S2.** Real-time, live-imaging movie of human adult cardiomyocytes (41-year-old female) as
972   described in **Figure 1.**

979
980
981