Inferring stability and persistence in the vaginal microbiome: A stochastic model of ecological

dynamics

José M. Ponciano[1]*, Juan P. Gómez[2], Jacques Ravel[3] and Larry J. Forney[4]

[1]Department of Biology, University of Florida, Gainesville, FL

[2]Departamento de Química y Biología, Universidad del Norte, Barranquilla, Colombia.

[3]Institute for Genome Sciences and Department of Microbiology and Immunology, University of

Maryland School of Medicine, Baltimore, MD

[4]Institute for Interdisciplinary Data Science and Department of Biological Sciences, University

of Idaho, Moscow, ID

*Corresponding author: josemi@ufl.edu

Open Research statement: the data used here is available at Ravel et al (2013). The same data

is also available at the computer code repository, github.com/jmponciano/StochasticMicrobiome.

Upon acceptance, the data will be uploaded to Dryad and the code to Zenodo.

Keywords: stochastic population dynamics of the microbiome, environmental stochasticity,

stochastic stability, persistence probability, population viability monitoring, multivariate

autoregressive model of population dynamics, stochastic community population dynamics,

Abstract

The interplay of stochastic and ecological processes that govern the establishment and

persistence of host-associated microbial communities is not well understood. Here we illustrate

the conceptual and practical advantages of fitting stochastic population dynamics models to

multi-species bacterial time series data. We show how the stability properties, fluctuation

regimes and persistence probabilities of human vaginal microbial communities can be better

understood by explicitly accommodating three sources of variability in ecological stochastic

models of multi-species abundances: 1) stochastic biotic and abiotic forces, 2) ecological

feedback and 3) sampling error. Rooting our modeling tool in stochastic population dynamics

modeling theory was key to apply standardized measures of a community's reaction to

environmental variation that ultimately depends on the nature and intensity of the intra-specific

and inter-specific interaction strengths. Using estimates of model parameters, we developed a

Risk Prediction Monitoring (RPM) tool that estimates temporal changes in persistence

probabilities for any bacterial group of interest. This method mirrors approaches that are often

used in conservation biology in which a measure of extinction risks is periodically updated with

any change in a population or community. Additionally, we show how to use estimates of

interaction strengths and persistence probabilities to formulate hypotheses regarding the

molecular mechanisms and genetic composition that underpin different types of interactions.

Instead of seeking a definition of "dysbiosis" we propose to translate concepts of theoretical

ecology and conservation biology methods into practical approaches for the management of

human-associated bacterial communities.

 INTRODUCTION

For decades now, inferring the interplay between stochastic processes and the ecological and

evolutionary conditions that permit the establishment and persistence of host-associated

microbial communities has remained a topic laden with controversies and unresolved conceptual

and practical issues (Zaoli and Grilli 2021; Grilli 2020; Zhou and Ning 2017; Ferguson and

Ponciano 2014; Gudelj et al. 2010; Robinson, Bohannan, and Young 2010; Ponciano et al.

2007). The paucity of studies connecting extensive time-series data with population dynamics

models rooted in ecological principles has been at the center of the problems faced when

inferring processes from patterns in this area of research (Zhou and Ning 2017). This knowledge

gap is exemplified here for the human vaginal microbiome. Work done to characterize these

bacterial communities using experimental and quantitative analytical approaches (Ravel et al.

2011) has shown that idiosyncratic changes in species composition and wide temporal

fluctuations in the relative abundances of the different species are undeniably associated with

specific environmental variables like pH. However, even a basic understanding of the

mechanisms leading to these fluctuations remains elusive. Given that the structure and

composition of an ecological community often alternates between distinct, widely different states

(Shade et al. 2012; Gonze et al. 2017; 2018; Bardgett and Caruso 2020), the chances of dramatic

community shifts are better predicted using mechanistic, stochastic population dynamics models

(Schooler et al. 2011; Ives et al. 2003; Ponciano 2018; Ponciano, Taper, and Dennis 2018;

Auger-Méthé et al. 2021). Illustrating the conceptual and practical advantages of fitting

stochastic population dynamics models to multi-species bacterial time series data is the focus of

this paper.

Here we developed and tested a multi-species stochastic population modeling approach

(Ludwig 1976; Nisbet and Gurney 2003; Cushing et al. 2003; L. J. Allen 2010; Dennis et al.

2006; Ovaskainen and Meerson 2010; Dennis and Ponciano 2014; Ponciano 2018; Ponciano,

Taper, and Dennis 2018) to better understand how fluctuations in the environment ultimately

contribute to changes in species composition and abundances as well as to the overall community

stability. Our central hypothesis is that stability properties, diversity and fluctuation regimes of

human vaginal microbial communities can be better understood by explicitly accommodating the

following three sources of variability in time series models of multi-species abundances: 1)

stochastic biotic and abiotic forces, 2) ecological feedbacks and 3) sampling error. This modeling

framework translates tentative explanations of the sources of the temporal variation in bacterial

abundances into testable hypotheses that describe the interplay between ecological processes and

the dynamics of abiotic factors while taking sampling variability into account. This translation

was achieved by combining time series data of bacterial species composition with stochastic

models derived from basic ecological principles. This probabilistic approach results in a practical

statistical connection between biological hypotheses and time series data (Ponciano, Taper, and

Dennis 2018). Here we exemplify this process using 135 time series of human vaginal microbial

communities (Ravel et al. 2011).

In recent years considerable efforts have been made to characterize the composition of

vaginal bacterial communities found in healthy reproductive age women and to understand

interruptions to the homeostasis of this microbiome. Community compositions that widely differ

from these "normal" states are thought to be in a state of 'dysbiosis'. Indigenous bacterial

populations that reside in and on the human body constitute the first line of defense against

infection by preventing non-indigenous organisms from causing disease. In the context of the

vaginal microbiome, dysbiosis can reflect changes in the absolute numbers of microbes, the

species composition, or changes in the relative abundances of bacterial taxa or some combination

thereof. The bacterial communities of reproductive age women often vary over time in a

seemingly haphazard way, and investigators assert that certain community states reflect an

'imbalance' in the vaginal microbiome, and these are 'unhealthy' states. Some of these states,

like those depleted of *Lactobacillus* species are said to reflect 'dysbiosis' despite persisting for

extended periods of time in women who are asymptomatic and otherwise healthy.

The concept of biological community stability has motivated significant theoretical

advances and large empirical research efforts in ecology (McCann 2000; Loreau et al. 2001; May

2019; Ives and Carpenter 2007; Little et al. 2008; Loreau 2010). The disparity between the

theoretical predictions and empirical evidence concerning diversity-stability relationships has

generated historical controversies that remain unresolved (Loreau 2010). These can in part be

attributed to the multiple definitions of stability that have been used (Ives and Carpenter 2007),

and partly because diversity per se is rarely a primary driver of stability. Rather than being

immediately linked to stability, diversity commonly acts as a secondary driver, itself being

subject to the same anthropogenic and environmental drivers that affect stability via a variety of

mechanisms (Ives et al. 2008; Altizer et al. 2006). Studies are needed that reveal the ecological

processes and abiotic factors that link diversity to stability, particularly in microbial communities

(Arumugam et al. 2011; Faith et al. 2011; 2013; Gajer et al. 2012).

Combining mathematical, statistical and stochastic process tools to explicitly model the

mechanisms that underlie community dynamics on a temporal scale has long proved to be a

fruitful approach to fill knowledge gaps regarding the functioning of ecological communities

(Cushing et al. 2003; Ives et al. 2003). This approach has also been shown to reliably reproduce

the regular waxing and waning of natural population densities in single and multi-species

systems (Zeng et al. 1998; Ponciano et al. 2005; E. J. Allen, Allen, and Schurz 2005; Dennis et

al. 2006; Barger and Bunge 2008; Taper and Ponciano 2016; Ponciano 2018; Ponciano, Taper,

and Dennis 2018; Dennis et al. 2019). Here we approached the problem of estimating bacterial

community stability by explicitly modeling this property as resulting from the interaction of

ecological feedback and stochastic (randomly fluctuating) abiotic factors (Ives et al. 2003).

Biological communities are continuously buffeted by changing environments and abiotic

factors that induce temporal fluctuations in the growth rates of each species in the system

(Dennis 1989; Dennis and Taper 1994; Grenfell, Bjørnstad, and Kappey 2001; Ives et al. 2003).

Environmental changes are likely to affect the availability of resources and hence the rate at

which bacteria replicate. Furthermore, changes in the availability of limiting resources are

expected to be concomitant with changes in the nature and intensities of intra-specific and inter-

specific competition processes. Ultimately, these environmental changes are expected to be

translated into changes in population sizes. In the face of unpredictable environmental changes,

equilibrial states of ecological communities are better characterized by means, variances and

other statistical quantities instead of point equilibria derived from deterministic, Lotka-Volterra

like models (Ives et al. 2003; Grilli 2020).

Mathematical characterizations of how the mean and variance of population sizes change

over time can be obtained by formulating multi-species population growth as stochastic

processes (L. J. Allen 2010; Ferguson and Ponciano 2014; Ponciano 2018; 2018). These

mathematical expressions reveal the links between the patterns of population variation,

environmental variation and key ecological quantities like intrinsic growth rates and inter-

specific and intra-specific competition coefficients. In this work, we bring these mathematical

137 characterizations to life by fitting stochastic ecological population models to a large data set

138 consisting of 135 time series data sets that each spanned 70 days with daily samples. From the

139 abundance time series, we explicitly estimate the strength of intra- and inter-specific competition

140 and use these estimates to compute dynamic stability metrics that describe the system's behavior.

141 Furthermore, we show that it is possible to link changes in the persistence (or extinction)

142 probabilities of any given bacterial type of interest with changes in intra-specific and inter-

143 specific competition coefficients. By understanding the relationship between these coefficients

144 and the population fluctuations of vaginal bacteria of clinical interest, our analysis constitutes the

145 first step towards assessing the risk to diseases linked to either the fast growth, invasion, or

146 extinction of different species in vaginal bacterial communities. Our approach makes

147 fundamental restoration principles and modeling techniques accessible to applied research

148 programs that focus on predicting the tendencies of microbial communities.

150       STOCHASTIC MODELS OF POPULATION ABUNDANCE

151  Past decades have seen the theory and practice of statistical ecology merge into a unified,

152 coherent, and robust framework for scientific inquiry using time series of animal abundances

153 (Newman et al. 2014, Murray and Sandercock 2020). Stochastic models of the temporal

154 fluctuations of species' abundances aim to translate fundamental concepts in ecology and

155 evolution into testable hypotheses and predictions that can be confronted with abundance time

156 series datasets. These models decompose the changes in abundances of one, two or more species

157 over time into four main components (Lewontin and Cohen 1969; Athreya and Karlin 1971;

158 Ludwig 1976; Tier and Hanson 1981; Dennis and Taper 1994; Engen, Bakke, and Islam 1998;

159 Ives et al. 2003; Dennis and Ponciano 2014; Ferguson and Ponciano 2014; 2015; Ponciano 2018;

Ponciano, Taper, and Dennis 2018). These four components are: 1) basic demographic processes

of the study organisms (here bacteria) like reproduction and the effects of density dependence

and inter-specific interactions, all of which may depend on current and past abundances of the

species in the system; 2) chance variation and individual heterogeneities affecting births and

deaths, known as "demographic stochasticity" effects; 3) environmental stochasticity or temporal

variation in vital rates (e.g., birth and death rates) that reflect variation in environmental

conditions; and 4) observation error and sampling noise. If sampling error is not accounted for

then dynamics and processes may be grossly misrepresented. This caveat is particularly relevant

in microbial systems (Kareiva, Parker, and Pascual 1996; Dennis et al. 2006; Ferguson and

Ponciano 2014; Grilli 2020).

One of the most widespread applications of these stochastic population dynamics models is

the characterization of extinction processes and persistence dynamics of species of interest

(Lande and Orzack 1988; Dennis, Munholland, and Scott 1991; Boyce 1992; Lande 1993; Foley

1994; Staples, Taper, and Shepard 2005; Chaudhary and Oli 2020). Of particular interest for

studies of bacterial population dynamics are recent efforts that explicitly incorporate the effects

of interspecific interactions on extinction or persistence processes in experimental microcosms

(Ferguson and Ponciano 2014). By including the interactions between species, these models can

quickly become intractable. However, if the effects of environmental stochasticity are included,

simple models that forego some of the biological complexities can still provide accurate

characterizations of the fates of species in a community (Ferguson and Ponciano 2014).

The interaction between environmental stochasticity and intra-specific and inter-specific

competition coefficients determine how much population sizes will fluctuate over time (Ives et

al. 2003). Changes in the quality of the environment have historically been cast as agents that

change population growth rates in mathematical models of population dynamics. These are

directly expressed as increases in the mean of the progeny distribution in a population of interest,

with a concomitant improvement in the maximum growth rate. Simultaneously, temporal

fluctuations in the environment are modeled as a time dependent random variable that will

randomly improve or reduce the growth rates of a population (Lewontin and Cohen 1969;

Ludwig 1976). However, stochastic contributions to the quality of an environment and overall

population dynamics have been found to interact in important ways with ecological processes,

such as density-dependence and inter-specific competition (Tier and Hanson 1981; Engen,

Bakke, and Islam 1998). Ives et al (2003) showed that the overall effect of environmental

fluctuations in the growth of a population is modulated by ecological processes. These authors

showed that the growth rate of a population characterized by weak density-dependence was

easily affected by fluctuations in the quality of the environment whereas those populations

characterized by strong density-dependence were not. When presented with the same temporal

regime of environmental variation a population with strong density-dependence will fluctuate

much less than a population with weak density-dependence (see Figure 1). Ives et al (2003) went

on to show that for a single population, stability could be measured and conceptualized as the

ratio of the magnitude of environmental variation to the strength of density dependence. This

finding allows for a direct comparison of the reactions of two different populations to the same

environmental noise regime. This insight that was brought about by Ives et al. (2003) in the

context of community ecology, made it possible to compare different populations and

communities on the same level playing field.

In an ecological community, the influence of environmental noise variance is modulated by

the density dependence and the inter-specific interaction coefficients (Ives et al. 2003; Ferguson

and Ponciano 2015). This is illustrated in Figure 2. Here we show four different scenarios in

which the strength of intra-specific and inter-specific interactions varied while environmental

noise remained constant. In each of these scenarios, three species (1, 2 and 3) interact in the

following ways: species 1 and 2 and 2 and 3 are competitors and thus have a negative effect on

each other. Species 1 and 3 are mutualists and hence have a positive effect on each other (Figure

1). Finally, all the species show negative density dependence. In the first scenario, all the

interactions including intraspecific density dependence are weak. In the second scenario only

species 3 had strong density dependence while the rest of the interactions were weak. In the third

scenario species 2 has a strong negative effect over 1 and 3 but the rest of the interactions were

kept weak. Finally, in the fourth scenario we made intraspecific interactions strong while

keeping interspecific interactions weak. The coefficients used for each scenario are shown in

Table S1.

We show how the same amount of environmental variance may result in either large or

small growth rate variation, depending on the maximum growth rates and those specified

interaction strengths (Figure 2). In a community, the strength of the inter-specific and intra-

specific interactions and the overall architecture of its assembly is what ultimately modulates the

response to environmental variation. Just as in single species population dynamics, the same

level of observed variation in the growth rate can result from the populations in a community

over-reacting to mild exogenous fluctuations, or alternatively, from a community dampening

considerably unusually large environmental variability. Ives et al. (2003) showed that it was

possible, through the analysis of multi-species time series, to estimate four different statistics or

"stability metrics" (called VP, MR, VR and R in Figure 2 (and as explained below) that would

allow the comparison of multiple communities in the face of the same magnitude of

environmental noise. In essence, and without entering into mathematical details, these authors

showed that it was possible through these metrics to obtain a standardized measure of the

reaction of a community to such noise. These findings also imply that deeming a particular set of

time series as representative of "stable" or "unstable" dynamics just by its overall variability

might be misleading and conflate the fundamental processes governing the dynamics of an

ensemble of interacting populations.

THE "MAR" MULTI-SPECIES STOCHASTIC POPULATION DYNAMICS MODEL

Twenty years have elapsed since Ives et al. proposed their modeling approach (the "MAR"

model) yet its use in microbiology has seldom been considered. We believe that the MAR model

provides benefits in terms of understanding, classifying, and predicting the dynamics of bacterial

abundances that have seldom been clearly presented in the context of microbial communities.

What follows is an effort to explain these benefits.

The MAR model is a discrete-time Markov process that is deeply rooted in stochastic

population dynamics modeling theory (Ives et al. 2003). It jointly models three processes that

determine the variation in abundance of the species in a community through time: 1) a

deterministic density-dependent population growth for every species in the system on a log-

scale, 2) the effect of every species on the growth rate of any other species and 3) the effects of

environmental variation on the growth rate (see Ives et al 2003 for a full model description).

This stochastic model has as its deterministic counterpart, the multispecies Gompertz density-

dependent model, which has been widely applied to estimate bacterial growth (see Dennis and

Ponciano 2014 and citations therein). The MAR model is amenable to simulations via recursion

because the total abundance of any species in one time step only depends on the abundance of all

the species in the previous time step. Thus, the time series data can be modeled using its linear,

multivariate recursion and representation

$$X_t = A + BX_{t-1} + E_t.$$

X_t is a vector of the log-population abundances at time t, A is a vector whose elements give

the intrinsic rate of increase for each species in the system, B is a squared matrix, whose

elements b_{ij} denote the effect of the abundance of species j on the growth rate of species i.

Finally, E_t represents a vector of stochastic, environmental factors varying independently from

one time step to the next. These factors are modeled with a multivariate normal distribution with

mean 0 and variance-covariance matrix Σ. Through this variance-covariance matrix Σ the

modeler can specify whether the response to environmental variation is independent from one

species to the next or not, and if not, any covariance structure could be added. In macro-

ecological communities for instance the response to the environment from one species to the next

might be phylogenetically constrained. In bacterial communities, these phylogenetic constraints

likely directly translate into explicit functional constraints, since any given strain might be better

at doing something the others cannot do (Ma et al 2020). The MAR model can be viewed as a

linear, first approximation to a complex, multi-species population dynamics process of the form

$n_t = h(n_{t-1})$, where the species population abundances n_t at time t are given by some

transformation $h(n_{t-1})$ of the abundances on the previous time step. Specifically, it can be

shown that the community matrix of such a complex process has eigenvalues that are identical to

the matrix B of the MAR model. The diagonal elements of B, b_{ii}, which represent the intra-

specific, density-dependent effects also satisfy the three existing theoretical definitions of the

strength of density dependence (see Ponciano, Taper, and Dennis 2018): the marginal effect on

the per capita growth rate of an increase in density (Holt 1985; Holt and Barfield 2012), the

derivative of the recruitment map at equilibrium (Holt and Barfield 2012) and the negative

elasticity at equilibrium of the per capita population growth rate with respect to change in the

population (Lande et al. 2002). The latter measure is readily extendable to scenarios dealing

with more complex life histories (Lande et al. 2002).

Jointly, the model matrices $\boldsymbol{B}$ and $\boldsymbol{\Sigma}$ hold the key to formulate standardized measures of how

a community reacts to environmental variation. These measures ultimately depend on the nature

and intensity of the intra- and inter-specific interaction coefficients (Ives et al. 2003; Dennis et

al. 2006; Ferguson and Ponciano 2015; Ponciano, Taper, and Dennis 2018). Ives et al. (2003)

derived four standardized metrics based on the $\boldsymbol{B}$ and $\boldsymbol{\Sigma}$ matrices and their eigenvalues. Variance

Proportion (VP) quantifies how the long-run variance of the population compares to the variance

of the environmental noise process. It is a summary of how the environmental noise distribution

in blue in Figure 1 compares to the population size distribution in gray in Figure 1. As Figure 2

shows, differences in variability in the multi-species time series can be directly attributed to

species interactions. In a stable system the interactions among species that modulate changes in

population sizes in a community from one generation to the next will be such that they cause the

variance of the population abundances to be only slightly larger than the variance of the

environmental noise (see Figure 1 for an example with a one-species system). On the other

hand, in a less stable system the species interactions greatly amplify the environmental

variability thereby generating large population fluctuations (Ives et al. 2003). This amplification

can be directly measured by the eigenvalues of the matrix $\boldsymbol{B}$, namely, by $\det(\boldsymbol{B})^{(2/p)} =$

$\left(\lambda_1 \lambda_2 \ldots \lambda_p\right)^{2/p}$ where p is the number of species in the system (see Ives et al. 2003 eq. 24 and

subsequent paragraph). In the face of environmental variation, the growth rate of a population

will react. This reaction is modulated, or filtered, by the intra-specific and inter-specific

competition coefficients (Ives et al. 2003).

The Mean Return time (MR) and the Variance Return time (VR) refer to the amount of time

that it takes the system to return to its stationary distribution. It's the stochastic equivalent of the

deterministic return time (Ives et al 2003). Specifically, it refers to the rate at which the transition

distribution of the system converges to its stationary distribution. The shorter the time, the more

stable the community is. Finally, Reactivity (R) is a measurement of how far the system pushes

away from its equilibrium after it is perturbed and as Ives et al. argue, can be computed in two

different ways, giving a total of four metrics of stochastic stability.

307   METHODS

Fitting the MAR model to extensive time series of microbial abundances presents at least

three major methodological challenges: The first is determining whether there exists enough

information in the data to estimate the MAR model parameters. This question boils down to

determining which time series length is sufficient to provide statistically sound parameter

estimates. The second methodological task is separating the environmental process variability

from sampling noise. The third one is dealing with missing data points: incomplete time series

are commonplace in these ecological studies. In what follows, we detail our approach to these

three problems.

316   *Minimal sample size to fit a MAR model*

The quality of the statistical fit of the MAR model depends on the amount of information

present in the multi-species data set. This information can be measured through the statistical

properties and diagnostics related to the model parameter estimates. The statistical quality of the

parameter estimates is in turn related to how many data points per parameter, or "degrees of

freedom" one has available to do model fitting. Another way to think about quantifying this

information is by computing the ratio of data to the number of unknown parameters. Ives et al.

(2003) model is, however, quite data-hungry: Let p be the number of species in the data set. The

vector of maximum growth rates A has p unknown parameters. The matrix of interactions B and

the variance-covariance matrix of the environmental fluctuations Σ have each $p \times p$ unknown

parameters, thus, the total number of model unknowns is $2p^2 + p$. With 13 species this number

is 351. This number can be compared to the available number of independent data points in order

to gauge if one has enough "degrees of freedom" for estimation.

Because this model is Markovian, every time-step transition (change in population

abundance) is an independent data point. The likelihood function of the MAR model, from

where its parameter estimates are derived, is therefore computed as the product of all the

observed transitions (Ives et al. 2003). This likelihood is maximized to obtain the parameter

estimates. If n is the length of the time series (70 in our case, see below), then the number of

transitions that can be used for the maximization of the likelihood function is $n - 1$. If m is the

number of replicated samples per species per time point, then the number of data points available

for parameter estimation is simply $(n - 1)mp$. Consequently, for the estimation to be feasible,

one needs to verify that $(n - 1)mp > 2p^2 + p$. On the other hand, solving for n in this

inequality gives the minimum sample size (time series length and/or number of replicates per

time step) needed to ensure estimability as $\frac{2p+1}{m} + 1$, which is equal to $2(p + 1)$ in the common

case where $m = 1$. For example, with 13 species and one replicated time series with no

observation error, $2(p + 1) = 28$ and the ratio of observations to number of parameters is

$\frac{(n-1)\times1\times p}{2p^2+p} = \frac{897}{351}$. Finally, this thinking can be extended by including the parameters needed to

decompose biological (process) variation from sampling error variation.

345   *Statistical decomposition of the sources of temporal variation*

In this study, we decompose the changes in abundances of species over time into three of the

four main components mentioned above (Dennis et al. 2006; Dennis and Ponciano 2014;

Ferguson and Ponciano 2014): 1) Population growth, density dependence and inter-specific

interactions, or predictable changes in births and death due to current and past abundances of the

species in the system 2) environmental stochasticity or (random) temporal variation in vital rates

representing variation in environmental conditions (good/bad times for survival and reproduction

and 3) observation error and/or sampling noise which if left unaccounted can lead to grossly mis-

represented dynamics (Kareiva, Parker, and Pascual 1996; Dennis et al. 2006; Dennis and

Ponciano 2014). Demographic stochasticity, the fourth component, although not included in this

first phase of our studies can be accommodated in time series estimation methods (Newman et al.

2014).

State-space models, widely known as statistical hierarchical models, allow decomposing the

biological and sampling sources of variation using a one pass statistical fit (Ponciano 2004,

Dennis et al 2006, Ponciano et al 2009, Newman et al 2014). Stochastic population models with

added observation error are just one example of this wide class of models. Although these

models are routinely used (Auger-Méthé et al. 2021), it has long been known that their fitting

isn't without statistical difficulties due to parameter identifiability problems, among others

(Ponciano et al. 2005, 200; Dennis et al. 2006; Knape 2008; Lele, Nadeem, and Schmuland

2010). Recently, statistical ecologists have extensively documented and demonstrated such

challenges (Lele 2020; Auger-Méthé et al. 2021). Further studying the statistical and scientific

merits of different computer intensive approaches to obtain either the maximum likelihood

estimates (via Data Cloning, the Laplace approximation, the Geyer-Thompson likelihood ratio

algorithm, Monte Carlo integration to name a few) or the Bayesian posteriors as well as Bayes

Factors for these state-space models is a task that merits its own, separate efforts and goes well

beyond the conceptual scope of this manuscript. In any the case, our research group in

collaboration with microbial ecologists and evolutionary biologists has extensively compared

experimentally derived population dynamics parameter estimates with those obtained via

maximum likelihood fitting of multi-species/types models and thus verified experimentally and

theoretically the reliability of this approach (De Gelder et al 2004, 2007, Ponciano et al 2007,

2009, Loftie-Eaton et al 2016, 2017).

Our present approach to fit a multi-species population model to a bacterial community time

series data set was as follows: first, we estimated the most likely location of the true, unobserved

abundances with sampling error removed, along with their confidence intervals using the

Kalman estimation methodology developed by Dennis and Ponciano (2014). This methodology

simultaneously accounts for sampling error and missing data points in the time series of

abundances. The resulting observation-noise filtered time series of abundances were then used to

fit the MAR stochastic population dynamics model for the entire community. While doing this

second fit, the statistical uncertainty resulting from the first observation error step was

propagated via parametric bootstrap (Taper et al. 2021). Separating the estimation of the

observation error from the biological process error allowed us to be sure at each step that the

Mean Squared Error (MSE) of the model parameters were adequate via extensive simulations

(github.com/jmponciano/StochasticMicrobiome).

389   RESULTS

390   *From statistical ecology theory to practice: insights from a case study*

In what follows, we applied the theoretical insights described above to an extensive data set

of dynamic vaginal microbial communities. We then contrasted the resulting inference with the

traditional practice of using the presence of a particular bacterial species at certain abundances

from a snapshot of a bacterial community to imply etiology. We contend that such practice may

in the end obscure, rather than illuminate our understanding the effects of different bacterial

community compositions simply because population abundances can and often do vary widely

over time. Additionally, we show how the concepts explained above contribute to answer

questions of practical interest. For example: under which ecological scenarios (i.e,. set of inter-

specific and intra-specific interactions) will the abundances of species in a community quickly

return from their current state to one where variation and composition regimes imply low health

risks. How can the concept and measurement of "stochastic stability" contribute to estimate

persistence probabilities?

The data we analyzed to exemplify the application of statistical ecology concepts were part

of the Human Microbiome Project funded by the National Institutes of Health in which 135

women (see Clinical Study Methods in Supplementary Material). Women enrolled in this study

self-collected daily mid-vaginal swabs for 10 weeks. We examined temporal changes in the

composition of vaginal communities established using 16S rRNA gene sequencing. Every day

after swab collection each participant also measured vaginal pH (see Ravel et al. 2011 for pH

measurement methods). A simple examination of the temporal variation of pH in these samples

(Figure 3) clearly illustrate the fact that ample temporal variation in the dynamics of bacterial

populations and their metabolic activities were the rule, rather than the exception. Important

feedback loops between pH levels and bacterial metabolic activity are expected (Ravel et al.

2011) and these processes can be examined with our theoretical approach, as we explain below.

415   *Minimal sample sizes to fit a MAR model*

Using the diagnostic tool presented in the Methods section, we determined which time series

data sets had enough data to be able to estimate the MAR model parameters. Computing the ratio

of available data to the number of parameters to be estimated, we determined that 88 community

time series out of the 135 total available could be reliably used for a full, multi-species

population model-fitting analysis. The rest of the analyses presented here are based on these 88

community time series data sets.

STATISTICAL DECOMPOSITION OF THE SOURCES OF TEMPORAL VARIATION

Estimating the interaction coefficients: compositional data vs abundance data

We proceeded to fit the MAR model using the estimated population dynamics time

trajectories without sampling error for all species and all data sets (Fig 4). While doing this

second fit, the statistical uncertainty from the first step was propagated via parametric bootstrap.

However, for this second step microbiologists usually face a key practical decision: should they

only work with relative abundances of species or work with both, relative and estimated total

abundances. For our case, the total abundance is the number of 16S rRNA gene copies per

sample established using quantitative PCR. To make an informed decision as to which approach

to undertake, here we simulated time series community abundances under the four scenarios

shown in Figure 2. We then verified which interaction strength estimates were less biased

(whether those resulting from using the compositional data or those resulting from using the total

abundances). Although the relative abundances of species in a community (i.e., compositional

time series data) are sometimes the only time series data available, our simulations showed that

using compositional data leads to biased estimates of the interaction strengths specified in the

matrix $\boldsymbol{B}$ of the MAR model.

Our simulation approach was as follows: first, we selected the four community scenarios

described in Figure 2 and simulated for each case 1000 time series of the abundance of the three

taxa A, B and C. We then estimated the interaction coefficients using the MAR model described

above fitted to both the relative abundance time series and the absolute abundance time series.

The absolute abundances were estimated by anchoring the proportions into total abundances at

each time step. Next, we fitted the MAR model to estimate the interaction coefficients using both

the 1000 time series of relative abundances and the 1000 time series of total abundances. Then,

we calculated the ratio between the estimated and the true interaction strength in each case.

When the interaction coefficients were estimated appropriately, a boxplot centered at 1 with a

small variance resulted. We estimated if each ratio between estimated and true coefficients

departed from an expected value of 1. The results of this simulation experiment (see Figures S1-

S4) clearly show that when the total abundances are used, the relative bias boxplots are centered

around one. When compositional data is used, those boxplots have a much wider interquartile

range and most of the time, are not even centered around one. Thus, fitting the MAR model to

compositional data tends to lead to severely biased estimates of the interaction strengths.

Therefore, the best approach to estimate the interaction coefficients is to use total abundance

data.

In longitudinal studies of microbiomes, the number of 16S rRNA gene copies only provides

estimates of the absolute abundance of taxa and not the true abundance of each bacterial species.

Our simulations demonstrated that estimating the strength of intra-specific and inter-specific

interactions based on relative abundance data results in biased estimates of the interaction

strengths. Hence, we performed a pan-bacterial qPCR assay to quantify the total 16S rRNA gene

copies in each of the samples, which estimates the absolute bacterial abundance in each sample.

Estimates of true abundance were then calculated for each taxon by multiplying relative

abundance by total 16S rRNA gene copies. The qPCR assays were done in triplicate for each of

the 135 women to document the variability in the abundances of species due to observation error.

We fitted the three different multi-species population dynamics assemblies/model variants

using the MAR model of Ives et al. (2003) and the de-noised time series data sets. The first

model variant consisted of using all 13 species mentioned above. The second model variant

required fitting a three-species model where we grouped all four *Lactobacillus* species into a

single ecological species, *Gardnerella vaginalis* as the second species and the other eight species

grouped into a third species. For the third variant we fitted a simple 2-species model with all

four *Lactobacillus* species grouped as the first species and the other nine species grouped as the

second taxon.

With the MAR model parameter estimates for each model variant (13 species, 3 species and

2 species models), we computed Ives et al. stochastic stability metrics. For each one of the three

cases, we then classified the 88 vaginal bacterial communities into four different stability

categories using a Principal Components Analysis (PCA) on their estimated stability metrics.

Using k-means clustering on the resulting PCA scores for these women and the fact that for all

these metrics, lower values indicate higher stability, the 88 bacterial communities were classified

into four different categories: Highly stable, stable, unstable, highly unstable. The best

classification scheme out of the three different multi-species models corresponded to the two-

species MAR model where we pooled all the 4 *Lactobacillus* species into the first taxa and the

other 9 species into the second species (the performance criterion to pick a best classification

scheme was the amount of variance explained by the analysis). This classification scheme is

shown in Figure 5.

486   *Estimation of persistence dynamics from the MAR model parameter MLEs*

Alone, our stability classification scheme is an insufficient approach to understand multi-

species population dynamics because the "stable" and "unstable" attributes are given here to a

community without regard for the health risks associated with its composition. Stability, which

is a property of dynamic systems, should not be equated with desirable or undesirable behavior

in terms of health outcomes because one community can have stable population dynamics but

sustain a low relative abundance of a strain, thus bringing high health risk. Thus, considering

overall abundance and composition in addition to stability is needed in order to assess the

desirability of a particular community dynamics.   Indeed, Klatt et al (2017) show that when the

relative abundance of *Lactobacillus* dwindles down below a 0.5 proportion, the bacterial

community is under a high risk of infection by HIV. On the other hand, as the relative

abundance of *Lactobacillus* moves above 0.5, the risk of infection decreases. Seeking to

elucidate which type and magnitude of ecological interactions would lead to desirable dynamics

(i.e. fluctuations in relative abundance of *Lactobacillus* above 0.5) is a reachable target under our

analysis using the MAR model. If attaining a sustained high relative abundance of *Lactobacillus*

over time is a health-management target as in Klatt et al 2017 (see Figure S5), then we contend

that our approach described next should be used.

We developed a Risk Prediction Monitoring (RPM) tool that estimates the temporal changes

in persistence probabilities. This method mirrors conservation biology approaches for population

monitoring in which a metric measuring extinction risks is periodically updated with any change

in the population called Population Viability Monitoring, or VPM (Staples, Taper, and Shepard

2005). Before explaining and implementing our RPM tool, we first explain how the well-known

VPM method from Staples et al. works and apply it directly to one of our 88 data sets to

exemplify it. Immediately afterwards, we fully develop and implement our RPM method.

The VPM method consists of serially estimating the persistence probabilities with every

data point added to the current length of the time series of population abundances. For annually

reproducing species, with every year that passes a new total abundance is recorded. With it, an

updated estimate of extinction risk is computed. Repeating the same process for multiple years

yields a temporal trend of extinction risks. Consider the following example from conservation

biology: if population abundances of a threatened species are available for the past 30 years, and

if managers want to check whether as of late (e.g., for the past 10 years), the extinction risk of

the population has been increasing or decreasing, then the following is done (Staples, Taper, and

Shepard 2005): First, a stochastic population dynamics model is fit using the first 20 years of the

data. With the model parameter estimates and the data up to year 20, the probability that the

population will crash below a critical threshold within the near future, e.g., during next 5 years,

is computed. The resulting probability is recorded. Next, the observed population size for year

21 is added to the time series. The model parameters are then re-estimated and the probability

that the population will crash sometime during the next 5 years after year 21 is computed. That

probability is also recorded. Iterating this process for 10 more years yields a time series of the

extinction risks for the last 30 years.

Here we exemplify the conservation biology method with one of our 88 bacterial community

datasets. For our vaginal bacteria data set, our target was to track the probability that the

proportion of all the *Lactobacillus* species in a vaginal bacterial community drop below 50%.

Our time unit in this case is days, as new swabs were collected daily. The abundances for all

bacterial taxa, as well as for the proportion of *Lactobacillus* species were available for 70 time-

steps. In Figure 6, upper left panel, we first fitted the stochastic multi-species Gompertz model of

Ives et al. (2003) to a single time series of observed abundances and proportions of *Lactobacillus*

up to day 30 (black empty circles). We did so by placing all *Lactobacillus* taxa as one type in the

model and all other species were pooled together as a second type. We then used the model

parameters to project in the next ten days the *Lactobacillus* abundances and their proportion in

the population 50,000 times (grey lines). The proportion of such projected trajectories that

dropped below 50%, which was 0.4 in the upper left panel, is an estimate of the *Lactobacillus*

persistence probability above 50% during those ten days. With every passing day, this estimate

was updated. In the next three panels (upper right, lower left and then lower right) we show

these simulations for only days 40, 50 and 60, but daily changes in persistence probabilities for

days 30 to 70 were computed.

The VPM method illustrated above for our data set is essentially retrospective but here we

devised a prospective modified version of it, one that allows comparing the dynamics of multiple

communities in the near future. Furthermore, we switched the estimation focus from tracking the

probabilities of crashing below a population size or proportion threshold to follow their

complement, persistence probabilities. This modified method links our stability metrics with the

risk assessment task and is what we call the RPM tool. We developed the RPM tool because we

faced the problem of assessing the risk dynamics for all 88 communities and being able to

evaluate these under the same level playing field. To do such comparisons, we chose to evaluate

the risk dynamics all while answering the question: How would the risk of *Lactobacillus* spp.

falling below 45% change over the next 20 days if all communities were started with the same

proportion (50%) and then monitored over the next 20 days? We answered this question by

implementing these steps: First, we retrieved the MAR model parameter estimates for all 88

communities. Using these estimates, we computed the MAR model predicted mean abundance

of *Lactobacillus* at stationarity for every case. We then set these mean abundances as the starting

abundances for a 20-day projection in each case. Additionally, we assumed that the starting total

abundances for the non-*Lactobacillus* taxa in all these projections were equal to these

abundances. Thus, if in one case the mean abundance at stationarity of *Lactobacillus* was

predicted to be 3.5x108 16S rRNA gene copies per swab, the starting mean abundance of the

non-*Lactobacillus* species were assumed to be identical, 3.5x108 16S rRNA gene copies per

swab. With these starting values, we computed the mean projected abundances for the next 20-

day trajectories and used these to numerically estimate via simulations the probability that the

Lactobacillus taxa would remain above 45% on day t for $t = 1,2,3,...,20$. The resulting trends

in *Lactobacillus* persistence probabilities are shown in Figure 7.

Our RPM tool was used in conjunction with the estimated matrix ***B*** to identify which

interaction coefficient drove each persistence trend. We found that a decaying persistence trend

of a bacterial type of interest was explained by whether other bacteria impacted negatively or

positively its growth, which is the information contained in ***B***. We illustrated this finding using

the resulting decaying RPM trend for woman 60 (right panel of Figure 7). The estimated two-

by-two matrix of interactions B for woman 60 is as follows: the one-step total effect of non-

Lactobacillus species on the per capita growth rate of *Lactobacillus* species had a negative

coefficient, -0.39. On the other hand, *Lactobacillus* species had a small positive effect on non-

Lactobacillus, 0.001. Therefore, while the presence of non-*Lactobacillus* taxa had a negative

density dependence effect on the growth of *Lactobacillus*, while *Lactobacillus* had a positive

effect on the growth rate of non-*Lactobacillus*. In the end this asymmetry negatively affected the

growth of *Lactobacillus*. In both cases, the strength of the intra-specific density dependence was

weak (0.86 for *Lactobacillus* and 0.76 for non-*Lactobacillus* species). To verify whether that

asymmetry in the inter-specific growth rate effects was what drove the decay in persistence

probabilities for the *Lactobacillus* taxa, we did two numerical experiments: for the first

experiment, we simply switched the sign of the effect of one group on the other, so that non-

Lactobacillus had a positive effect of 0.39 in the growth rate of *Lactobacillus* and in turn,

Lactobacillus had a negative effect of -0.001 on the growth rate of non-*Lactobacillus* taxa. Next,

we re-computed the RPM trend in persistence probabilities using this modified B matrix of

interactions. The resulting RPM trend of persistence probabilities, plotted in pink in Figure 8,

remained at 1 for the next 20 days. The second numerical experiment consisted of artificially

increasing the maximum growth rate of the *Lactobacillus* taxa while leaving the B matrix of

interactions unchanged. Then, a restored trend in persistence probabilities was also obtained

(Figure 8).

590   DISCUSSION

This study constitutes an unprecedented integration of ecological, mathematical, statistical,

and conservation biology principles to understand and predict the dynamics of an extensive

microbial community's time series data set. In the past few years, the complex nature of

microbiome data has brought together an ever-growing number of multi-disciplinary research

teams (Qian, Lan, and Venturelli 2021). Yet, the fast pace of modern methodological research in

microbiome studies contrasts sharply with the paucity of population dynamics studies seeking to

understand from basic principles the benefits, or perhaps shortcomings, of novel data analysis

techniques. The main motivation of this study was the fact that by and large, variability in

microbial time-series data is still perceived as "statistical noise" rather than as an intrinsic

property of the growth of bacterial communities. Phrasing through the MAR model variability

over time as an intrinsic property of a growing population allows linking concepts like the

strength of intra-specific and inter-specific competition to the qualitative response of a

population in the face of uncertain environments. Not only can these competition coefficients be

estimated, and the stability of the system assessed by fitting the MAR model, but the chance of

persistence of bacteria taxa can be further assessed. To our knowledge, this is the first study that

demonstrates how persistence probabilities of bacteria of medical and ecological interest can be

estimated and even manipulated by identifying which interaction coefficient strengths are their

main drivers. We thus demonstrate how the apparently simple stochastic multi-species time

series model of Ives et al. (2003) can be used beyond its original applications to approach some

of the most pressing questions regarding the monitoring of bacterial communities (Bardgett and

Caruso 2020).

Vaginal communities dominated by species of *Lactobacillus* have been associated with

health and a reduced risk to diseases such as bacterial vaginosis or sexually transmitted

infections. The notion that dominance of *Lactobacillus* is associated with health is deeply

engrained in the field of women's urogenital health and strongly supported by the findings of

numerous studies (Chee, Chew, and Than 2020; Witkin and Linhares 2022). Regrettably, the

converse — that low proportions or the absence of *Lactobacillus* is unhealthy — has also

permeated the field's lexicon. This is a logical fallacy of denying the antecedent (Gaul 2018),

that essentially argues that if healthy women have vaginal communities dominated by

Lactobacillus, then the absence of *Lactobacillus* in vaginal communities is, of itself, unhealthy.

This claim is refuted by the findings of numerous studies on the species composition of healthy,

asymptomatic women that have shown that a significant proportion of healthy asymptomatic

women have vaginal communities with low proportions of *Lactobacillus* (Saraf et al. 2021;

Gosmann et al. 2017; Anahtar et al. 2018). Instead, they are dominated by various species of

strictly and facultatively anaerobic bacteria such as *Gardnerella vaginalis*, *Mobiluncus*,

Prevotella, *Brevibacterium*, *Peptoniphilus* and others (Onderdonk et al 2016). With that said, it

should also be recognized that low proportions of *Lactobacillus* in vaginal communities are

associated with an increased risk to disease (France et al 2022) though it is not a disease state per

se. Nonetheless, investigators have often referred to these communities as being either abnormal

(Green, Zarek, and Catherino 2015), out of balance (Olesen and Alm 2016), or in a state of

dysbiosis (Hooks and O'Malley 2017) that somehow needs to be corrected. We posit that except

for symptomatic bacterial infections, all other states are 'healthy' and in many instances they are

'normal' (meaning they are often observed) though they may differ in terms of risk to disease.

Most studies on the species composition of vaginal bacterial communities have employed

cross-sectional designs that yield point estimates of community composition. It seems to be

assumed that the species composition of communities is rather invariant over time in the absence

of some sort of natural or unnatural environmental disturbances such as menstruation or the use

of lubricants (Gajer et al. 2012; Wilkinson et al. 2019; O'Hanlon et al. 2021; Łaniewski et al.

2021). Contrary to this assumption, longitudinal studies have shown that the vaginal microbiota

of many women is dynamic and often transition through states in which *Lactobacillus* spp. are

lacking (Gajer et al. 2012; Lewis, Bernstein, and Aral 2017). These states vary in frequency and

duration and are therefore associated with varying levels of risk for urogenital infections and

other maladies. One could reasonably consider these to be windows of elevated risk that can

open and close, sometimes over very short periods of time.

Our PCA and MAR model-based stability classification scheme (Figure 5) takes a first,

admittedly imperfect step toward process-based management of bacterial community dynamics

and rigorous use of the term "stability". Although previous community classification schemes

using PCA relied on patterns of abundances, our approach relies on inferred ecological processes

from the time series of abundances. As theory and current practice in conservation biology show

(Murray and Sandercock 2020), the longer the multi-species time series data, the better the

information regarding species interactions in a community can be better teased apart. Here we

went one step further and estimated how these inferred interactions ultimately govern the

community response to environmental variability. The nature of such response was quantified

with Ives et al.'s (2003) four stability metrics and the PCA in Figure 5 separates bacterial

communities according to these metrics (see Supplementary material). Thus, the position of each

bacterial community in PCA space is determined by the strength of ecological interactions. If

one community is found to be largely unstable, an analyst can peer into the nature and intensity

of those estimated interactions and change them one by one to move the community in PCA

space from an unstable group into another classification group. In other words, an investigator

can test statistical hypotheses regarding which interactions are responsible for one or another

stability classification result. Identifying interactions that render a community stochastically

stable can be the first step in a research agenda that seeks to understand how to guarantee such

stability by modulating the strengths of interactions. Our RPM approach is a natural extension of

our stochastic stability inferences. It is an easy-to-understand approach to approximate the time-

dependent persistence probabilities of the bacterial species of interest. As Olesen and Alm (2016)

have argued, tools like our RPM approach that focus on prediction rather than simply the

detection of differences are needed, and here we deliver on that particular need.

Using this persistence probability methodology in studies of the vaginal microbiome would

mirror an approach called Population Viability Analyses that has been successfully used in

conservation biology for many years (Chaudhary and Oli 2020; Ponciano, Taper, and Dennis

2018). Unlike the majority of cases in conservation biology our model choice (the multivariate,

stochastic Gompertz with environmental stochasticity and added sampling error; Ives et al. 2003;

Dennis et al. 2006; Dennis, Ponciano, and Taper 2010) has been extensively tested in a recent

theoretical-simulation study (Ponciano, Taper, and Dennis 2018). Estimates of the strengths of

interactions can be used to formulate hypotheses regarding the molecular mechanisms and

genetic composition that underpin different types of interactions. By plotting the variability in

the sign (positive or negative) and intensity of interaction coefficients (for example, the effect of

Lactobacillus on the growth rate of *Gardnerella* or some other species) one can locate and

isolate cases where the sign of species interaction relations flip (say from positive to negative)

and eventually guide the laboratory determination of the genetic composition of strains

associated with interaction relationships in every quadrant (Figure S6).

Adopting statistical ecology theory and concepts reveals the inconsistencies of using terms

like "dysbiosis" to characterize a microbial community. Dysbiosis is commonly defined as a

change in the composition and function of a human microbial community that is typically driven

by environmental and host-related factors that exceed a community's resistance and resilience

(Kriss et al. 2018; Kindinger et al. 2016; Borgdorff et al. 2016; Levy et al. 2017). But this

definition doesn't seem to fully fit with what theoreticians in ecology understand as resilience

and resistance. Resilience, in one hand, is the rate at which a community returns to a state that

existed prior to a change. Resistance, on the other hand, is the magnitude of a community's

response to a given disturbance (Begon and Townsend 2021). Both, resilience, and resistance are

built into Ives' et al stability metrics. Instead of trying to frame a dysbiosis definition into these

concepts, it seems much more straightforward to use Ives' stability metrics directly to classify

the stability dynamics of a community, just as we do here. Additionally, in current practice,

investigators will often state that 'healthy' communities are 'in balance' (White et al. 2011;

Olesen and Alm 2016; Gupta 2021). This terminology reflects an erroneous assumption that the

composition of bacterial communities in healthy individuals is essentially invariant and that

changes in the relative abundances of species are necessarily bad and, in some cases, constitute

sufficient evidence to classify these variants as disease states. This classification is often done

based on pairwise comparisons of a microbiome at two points in time. Except for symptomatic

bacterial infections, it seems that all other states are 'healthy' and in many instances they are

'normal' (meaning they are often observed). These words and phrases are loosely defined and

inconsistently used, and this leads to confusion among non-experts. The literature is peppered

with examples (White et al. 2011; Olesen and Alm 2016; Gupta 2021). Instead of seeking a

definition of dysbiosis we assert that it might be better to translate concepts of theoretical

ecology into practical approaches for the management of human-associated bacterial

communities. This can be accomplished using concepts and methods that have come to be well

known in the fields of population dynamics and conservation biology.

713   CONCLUSION

Population dynamics as a field in ecology has long touted the theoretical and practical

advantages of jointly modeling demography and the influence of the environment and sampling

error (Cushing et al. 2003; Dennis et al. 2006), while conservation biology has taken advantage

of these ideas and modeling approaches to predict population persistence probabilities (Staples,

Taper, and Shepard 2005; Chaudhary and Oli 2020). Here we have shown that the same sort of

stochastic population dynamics equations can be used to re-phrase the concept of stability as the

magnitude of the reaction to a variable environment. Our work represents the first

comprehensive integration of theoretical stochastic population dynamics, unusually long time

series of bacterial community abundances and conservation biology principles. This integrated

approach resulted in two major steps towards a better understanding of human-associated

bacterial communities. First, through the estimation of each bacterial community's reaction to

exogenous variability we achieved a stability-based ecological community classification.

Second, we provide for the first time, estimates of the short-term persistence probabilities of

bacterial types of medical interest. This result is important because our estimated temporal trend

in persistence probabilities can be used to construct an evidence-based inference regarding the

fate of a pathogen, for example. Finally, we conclude that a comprehensive examination of the

reach of stochastic population dynamics modeling in the field of microbial community ecology is

beginning to take shape as a body of work. Our efforts provide a theoretical framework that can

very well represent microbial phenomena of interest in a simpler and unified way as effects of a

common cause: an alteration of the growth rate of a population by itself, by another population

or by the environment.

735   ACKNOWLEDGMENTS

736   We would like to thank the valuable input and advice at various stages of this work from Drs

737   Mark L Taper, Brian Dennis, and Robert D. Holt. Funding for this research was provided by the

738   National Institute of General Medical Sciences of the National Institutes of Health, under grant

739   number 1R01GM103604 to University of Florida, with JMP as PI. JMP was also supported by

740   NSF grant No. 2052372 to University of Florida.

742   CONFLICT OF INTEREST STATEMENT

JR is the cofounder of LUCA Biologics, a biotechnology company focusing on translating

microbiome research into live biotherapeutics drugs for women's health.

REFERENCES

Allen, Edward J, Linda JS Allen, and Henri Schurz. 2005. "A Comparison of Persistence-Time

Estimation for Discrete and Continuous Stochastic Population Models That Include

Demographic and Environmental Variability." Mathematical Biosciences 196 (1): 14–38.

Allen, Linda JS. 2010. An Introduction to Stochastic Processes with Applications to Biology.

CRC press.

Altizer, Sonia, Andrew Dobson, Parviez Hosseini, Peter Hudson, Mercedes Pascual, and Pejman

Rohani. 2006. "Seasonality and the Dynamics of Infectious Diseases." Ecology Letters 9

(4): 467–84.

Anahtar, Melis N, David B Gootenberg, Caroline M Mitchell, and Douglas S Kwon. 2018.

"Cervicovaginal Microbiota and Reproductive Health: The Virtue of Simplicity." Cell

Host & Microbe 23 (2): 159–68.

Arumugam, Manimozhiyan, Jeroen Raes, Eric Pelletier, Denis Le Paslier, Takuji Yamada,

Daniel R Mende, Gabriel R Fernandes, Julien Tap, Thomas Bruls, and Jean-Michel

Batto. 2011. "Enterotypes of the Human Gut Microbiome." Nature 473 (7346): 174–80.

Athreya, Krishna B, and Samuel Karlin. 1971. "On Branching Processes with Random

Environments: I: Extinction Probabilities." The Annals of Mathematical Statistics 42 (5):

1499–1520.

Auger-Méthé, Marie, Ken Newman, Diana Cole, Fanny Empacher, Rowenna Gryba, Aaron A

King, Vianey Leos-Barajas, Joanna Mills Flemming, Anders Nielsen, and Giovanni

Petris. 2021. "A Guide to State–Space Modeling of Ecological Time Series." Ecological

Monographs 91 (4): e01470.

Bardgett, Richard D, and Tancredi Caruso. 2020. "Soil Microbial Community Responses to Climate Extremes: Resistance, Resilience and Transitions to Alternative States." Philosophical Transactions of the Royal Society B 375 (1794): 20190112.

Barger, Kathryn, and John Bunge. 2008. "Bayesian Estimation of the Number of Species Using Noninformative Priors." Biometrical Journal: Journal of Mathematical Methods in Biosciences 50 (6): 1064–76.

Begon, Michael, and Colin R Townsend. 2021. Ecology: From Individuals to Ecosystems. John Wiley & Sons.

Borgdorff, Hanneke, Raju Gautam, Stuart D Armstrong, Dong Xia, Gilles F Ndayisaba, Nienke H van Teijlingen, Teunis BH Geijtenbeek, Jonathan M Wastling, and Janneke HHM van de Wijgert. 2016. "Cervicovaginal Microbiome Dysbiosis Is Associated with Proteome Changes Related to Alterations of the Cervicovaginal Mucosal Barrier." Mucosal Immunology 9 (3): 621–33.

Boyce, Mark S. 1992. "Population Viability Analysis." Annual Review of Ecology and Systematics 23 (1): 481–97.

Chaudhary, Vratika, and Madan K Oli. 2020. "A Critical Appraisal of Population Viability Analysis." Conservation Biology 34 (1): 26–40.

Chee, Wallace Jeng Yang, Shu Yih Chew, and Leslie Thian Lung Than. 2020. "Vaginal Microbiota and the Potential of *Lactobacillus* Derivatives in Maintaining Vaginal Health." Microbial Cell Factories 19 (1): 203.

Cushing, Jim Michael, Robert F Costantino, Brian Dennis, Robert Desharnais, and Shandelle Marie Henson. 2003. Chaos in Ecology: Experimental Nonlinear Dynamics. Vol. 1. Elsevier.

De Gelder, L., Ponciano, J.M., Abdo, Z., Joyce, P., Forney, L.J. and Top, E.M., 2004. Combining

mathematical models and statistical methods to understand and predict the dynamics of

antibiotic-sensitive mutants in a population of resistant bacteria during experimental

evolution. Genetics, 168(3): 1131-1144.

De Gelder, L., Ponciano, J.M., Joyce, P. and Top, E.M., 2007. Stability of a promiscuous plasmid

in different hosts: no guarantee for a long-term relationship. Microbiology, 153(2): 452-

463.

Dennis, Brian. 1989. "Allee Effects: Population Growth, Critical Density, and the Chance of

Extinction." Natural Resource Modeling 3 (4): 481–538.

Dennis, Brian, Patricia L Munholland, and J Michael Scott. 1991. "Estimation of Growth and

Extinction Parameters for Endangered Species." Ecological Monographs 61 (2): 115–43.

Dennis, Brian, and José Miguel Ponciano. 2014. "Density-dependent State-space Model for

Population-abundance Data with Unequal Time Intervals." Ecology 95 (8): 2069–76.

Dennis, Brian, José Miguel Ponciano, Subhash R Lele, Mark L Taper, and David F Staples.

2006. "Estimating Density Dependence, Process Noise, and Observation Error."

Ecological Monographs 76 (3): 323–41.

Dennis, Brian, José Miguel Ponciano, and Mark L Taper. 2010. "Replicated Sampling Increases

Efficiency in Monitoring Biological Populations." Ecology 91 (2): 610–20.

Dennis, Brian, José Miguel Ponciano, Mark L Taper, and Subhash R Lele. 2019. "Errors in

Statistical Inference under Model Misspecification: Evidence, Hypothesis Testing, and

AIC." Frontiers in Ecology and Evolution, 372.

Dennis, Brian, and Mark L Taper. 1994. "Density Dependence in Time Series Observations of

Natural Populations: Estimation and Testing." Ecological Monographs 64 (2): 205–24.

Engen, Steinar, Øyvind Bakke, and Aminul Islam. 1998. "Demographic and Environmental
Stochasticity-Concepts and Definitions." Biometrics, 840–46.

Faith, Jeremiah J, Janaki L Guruge, Mark Charbonneau, Sathish Subramanian, Henning Seedorf,
Andrew L Goodman, Jose C Clemente, Rob Knight, Andrew C Heath, and Rudolph L
Leibel. 2013. "The Long-Term Stability of the Human Gut Microbiota." Science 341
(6141): 1237439.

Faith, Jeremiah J, Nathan P McNulty, Federico E Rey, and Jeffrey I Gordon. 2011. "Predicting a
Human Gut Microbiota's Response to Diet in Gnotobiotic Mice." Science 333 (6038):
101–4.

Ferguson, Jake M, and José Miguel Ponciano. 2014. "Predicting the Process of Extinction in
Experimental Microcosms and Accounting for Interspecific Interactions in Single-species
Time Series." Ecology Letters 17 (2): 251–59.

Ferguson, Jake M, and José Miguel Ponciano. 2015. "Evidence and Implications of Higher-
Order Scaling in the Environmental Variation of Animal Population Growth."
Proceedings of the National Academy of Sciences 112 (9): 2782–87.

Foley, Patrick. 1994. "Predicting Extinction Times from Environmental Stochasticity and
Carrying Capacity." Conservation Biology 8 (1): 124–37.

Gajer, Pawel, Rebecca M Brotman, Guoyun Bai, Joyce Sakamoto, Ursel ME Schütte, Xue
Zhong, Sara SK Koenig, Li Fu, Zhanshan Ma, and Xia Zhou. 2012. "Temporal Dynamics
of the Human Vaginal Microbiota." Science Translational Medicine 4 (132): 132ra52-
132ra52.

Gaul, Brett. 2018. "Denying the Antecedent." In Bad Arguments, 46–47. John Wiley & Sons,
Ltd. https://doi.org/10.1002/9781119165811.ch3.

Gonze, Didier, Katharine Z Coyte, Leo Lahti, and Karoline Faust. 2018. "Microbial

Communities as Dynamical Systems." Current Opinion in Microbiology 44: 41–49.

Gonze, Didier, Leo Lahti, Jeroen Raes, and Karoline Faust. 2017. "Multi-Stability and the Origin

of Microbial Community Types." The ISME Journal 11 (10): 2159–66.

Gosmann, Christina, Melis N Anahtar, Scott A Handley, Mara Farcasanu, Galeb Abu-Ali,

Brittany A Bowman, Nikita Padavattan, Chandni Desai, Lindsay Droit, and Amber

Moodley. 2017. "*Lactobacillus*-Deficient Cervicovaginal Bacterial Communities Are

Associated with Increased HIV Acquisition in Young South African Women." Immunity

46 (1): 29–37.

Green, Katherine A, Shvetha M Zarek, and William H Catherino. 2015. "Gynecologic Health

and Disease in Relation to the Microbiome of the Female Reproductive Tract." Fertility

and Sterility 104 (6): 1351–57.

Grenfell, Bryan T, Ottar N Bjørnstad, and Jens Kappey. 2001. "Travelling Waves and Spatial

Hierarchies in Measles Epidemics." Nature 414 (6865): 716–23.

Grilli, Jacopo. 2020. "Macroecological Laws Describe Variation and Diversity in Microbial

Communities." Nature Communications 11 (1): 4743.

Gudelj, Ivana, Joshua S Weitz, Tom Ferenci, M Claire Horner-Devine, Christopher J Marx,

Justin R Meyer, and Samantha E Forde. 2010. "An Integrative Approach to

Understanding Microbial Diversity: From Intracellular Mechanisms to Community

Structure." Ecology Letters 13 (9): 1073–84.

Gupta, PD. 2021. "The Human Vaginal Microbiota: Boon or Bane." J of Obstetrics Gynecology

and Reproductive Sciences 5 (1).

Holt, Robert D. 1985. "Population Dynamics in Two-Patch Environments: Some Anomalous

Consequences of an Optimal Habitat Distribution." Theoretical Population Biology 28

(2): 181–208.

Holt, Robert D, and Michael Barfield. 2012. "Trait-Mediated Effects, Density Dependence and

the Dynamic Stability of Ecological Systems." Trait-Mediated Indirect Interactions:

Ecological and Evolutionary Perspectives 89.

Hooks, Katarzyna B, and Maureen A O'Malley. 2017. "Dysbiosis and Its Discontents." MBio 8

(5): e01492-17.

Ives, Anthony R, and Stephen R Carpenter. 2007. "Stability and Diversity of Ecosystems."

Science 317 (5834): 58–62.

Ives, Anthony R, Brian Dennis, Kathryn L Cottingham, and Stephen R Carpenter. 2003.

"Estimating Community Stability and Ecological Interactions from Time-series Data."

Ecological Monographs 73 (2): 301–30.

Ives, Anthony R, Árni Einarsson, Vincent AA Jansen, and Arnthor Gardarsson. 2008. "High-

Amplitude Fluctuations and Alternative Dynamical States of Midges in Lake Myvatn."

Nature 452 (7183): 84–87.

Jordt, H., Stalder, T., Kosterlitz, O., Ponciano, J.M., Top, E.M. and Kerr, B., 2020. Coevolution

of host–plasmid pairs facilitates the emergence of novel multidrug resistance. Nature

ecology & evolution, 4(6): 863-869.

Kareiva, Peter, Ingrid M Parker, and Miguel Pascual. 1996. "Can We Use Experiments and

Models in Predicting the Invasiveness of Genetically Engineered Organisms?" Ecology

77 (6): 1670–75.

Kindinger, Lindsay M, David A MacIntyre, Yun S Lee, Julian R Marchesi, Ann Smith, Julie AK

McDonald, Vasso Terzidou, Joanna R Cook, Christoph Lees, and Fidan Israfil-Bayli.

2016. "Relationship between Vaginal Microbial Dysbiosis, Inflammation, and Pregnancy

Outcomes in Cervical Cerclage." Science Translational Medicine 8 (350): 350ra102-

350ra102.

Knape, Jonas. 2008. "Estimability of Density Dependence in Models of Time Series Data."

Ecology 89 (11): 2994–3000.

Kriss, Michael, Keith Z Hazleton, Nichole M Nusbacher, Casey G Martin, and Catherine A

Lozupone. 2018. "Low Diversity Gut Microbiota Dysbiosis: Drivers, Functional

Implications and Recovery." Current Opinion in Microbiology 44: 34–40.

Lande, Russell. 1993. "Risks of Population Extinction from Demographic and Environmental

Stochasticity and Random Catastrophes." The American Naturalist 142 (6): 911–27.

Lande, Russell, S Engen, B-E Sæther, F Filli, E Matthysen, and H Weimerskirch. 2002.

"Estimating Density Dependence from Population Time Series Using Demographic

Theory and Life-History Data." The American Naturalist 159 (4): 321–37.

Lande, Russell, and Steven Hecht Orzack. 1988. "Extinction Dynamics of Age-Structured

Populations in a Fluctuating Environment." Proceedings of the National Academy of

Sciences 85 (19): 7418–21.

Łaniewski, Paweł, Kimberley A. Owen, Michael Khnanisho, Rebecca M. Brotman, and Melissa

M. Herbst-Kralovetz. 2021. "Clinical and Personal Lubricants Impact the Growth of

Vaginal Lactobacillus Species and Colonization of Vaginal Epithelial Cells: An in Vitro

Study." Sexually Transmitted Diseases 48 (1): 63.

https://doi.org/10.1097/OLQ.0000000000001272.

Lele, Subhash R. 2020. "How Should We Quantify Uncertainty in Statistical Inference?"

Frontiers in Ecology and Evolution 8: 35.

Lele, Subhash R, Khurram Nadeem, and Byron Schmuland. 2010. "Estimability and Likelihood

Inference for Generalized Linear Mixed Models Using Data Cloning." Journal of the

American Statistical Association 105 (492): 1617–25.

Levy, Maayan, Aleksandra A Kolodziejczyk, Christoph A Thaiss, and Eran Elinav. 2017.

"Dysbiosis and the Immune System." Nature Reviews Immunology 17 (4): 219–32.

Lewis, Felicia MT, Kyle T Bernstein, and Sevgi O Aral. 2017. "Vaginal Microbiome and Its

Relationship to Behavior, Sexual Health, and Sexually Transmitted Diseases." Obstetrics

and Gynecology 129 (4): 643.

Lewontin, Richard C, and Daniel Cohen. 1969. "On Population Growth in a Randomly Varying

Environment." Proceedings of the National Academy of Sciences 62 (4): 1056–60.

Little, Ainslie EF, Courtney J Robinson, S Brook Peterson, Kenneth F Raffa, and Jo

Handelsman. 2008. "Rules of Engagement: Interspecies Interactions That Regulate

Microbial Communities." Annu. Rev. Microbiol. 62: 375–401.

Loftie-Eaton, W., Yano, H., Burleigh, S., Simmons, R.S., Hughes, J.M., Rogers, L.M., Hunter,

S.S., Settles, M.L., Forney, L.J., Ponciano, J.M. and Top, E.M., 2016. Evolutionary paths

that expand plasmid host-range: implications for spread of antibiotic resistance. Molecular

biology and evolution, 33(4): 885-897.

Loftie-Eaton, W., Bashford, K., Quinn, H., Dong, K., Millstein, J., Hunter, S., Thomason, M.K.,

Merrikh, H., Ponciano, J.M. and Top, E.M., 2017. Compensatory mutations improve

general permissiveness to antibiotic resistance plasmids. Nature ecology & evolution, 1(9):

1354-1363

Loreau, Michel. 2010. "Linking Biodiversity and Ecosystems: Towards a Unifying Ecological

Theory." Philosophical Transactions of the Royal Society B: Biological Sciences 365

(1537): 49–60.

Loreau, Michel, Shahid Naeem, Pablo Inchausti, Jan Bengtsson, J Philip Grime, Andrew Hector,

DU Hooper, MA Huston, David Raffaelli, and Bernhard Schmid. 2001. "Biodiversity and

Ecosystem Functioning: Current Knowledge and Future Challenges." Science 294

(5543): 804–8.

Ludwig, Donald. 1976. "A Singular Perturbation Problem in the Theory of Population

Extinction." In , 10:87–104.

May, Robert M. 2019. Stability and Complexity in Model Ecosystems. Princeton university

press.

McCann, Kevin Shear. 2000. "The Diversity–Stability Debate." Nature 405 (6783): 228–33.

Murray, Dennis L, and Brett K Sandercock. 2020. Population Ecology in Practice. John Wiley &

Sons.

Newman, KB, ST Buckland, BJT Morgan, R King, DL Borchers, DJ Cole, P Besbeas, O

Gimenez, and L Thomas. 2014. "State-Space Models." In Modelling Population

Dynamics, 39–50. Springer.

Nisbet, Roger M, and William Gurney. 2003. Modelling Fluctuating Populations: Reprint of

First Edition (1982). Blackburn Press.

O'Hanlon, D. Elizabeth, Sarah E. Brown, Xin He, Christina A. Stennett, Sarah J. Robbins,

Elizabeth D. Johnston, Amelia M. Wnorowski, et al. 2021. "Observational Cohort Study

of the Effect of a Single Lubricant Exposure during Transvaginal Ultrasound on Cell-

Shedding from the Vaginal Epithelium." PLOS ONE 16 (5): e0250153.

https://doi.org/10.1371/journal.pone.0250153.

Olesen, Scott W, and Eric J Alm. 2016. "Dysbiosis Is Not an Answer." Nature Microbiology 1

(12): 1–2.

Ovaskainen, Otso, and Baruch Meerson. 2010. "Stochastic Models of Population Extinction."

Trends in Ecology & Evolution 25 (11): 643–52.

Ponciano, José Miguel. 2018. "A Parametric Interpretation of Bayesian Nonparametric Inference

from Gene Genealogies: Linking Ecological, Population Genetics and Evolutionary

Processes." Theoretical Population Biology 122: 128–36.

Ponciano, José Miguel, Leen De Gelder, Eva M Top, and Paul Joyce. 2007. "The Population

Biology of Bacterial Plasmids: A Hidden Markov Model Approach." Genetics 176 (2):

957–68.

Ponciano, José Miguel, Mark L Taper, and Brian Dennis. 2018. "Ecological Change Points: The

Strength of Density Dependence and the Loss of History." Theoretical Population

Biology 121: 45–59.

Ponciano, José Miguel, Frederik PJ Vandecasteele, Thomas F Hess, Larry J Forney, Ronald L

Crawford, and Paul Joyce. 2005. "Use of Stochastic Models to Assess the Effect of

Environmental Factors on Microbial Growth." Applied and Environmental Microbiology

71 (5): 2355–64.

Qian, Yili, Freeman Lan, and Ophelia S Venturelli. 2021. "Towards a Deeper Understanding of

Microbial Communities: Integrating Experimental Data with Dynamic Models." Current

Opinion in Microbiology 62: 84–92.

Ravel, Jacques, Pawel Gajer, Zaid Abdo, G Maria Schneider, Sara SK Koenig, Stacey L
McCulle, Shara Karlebach, Reshma Gorle, Jennifer Russell, and Carol O Tacket. 2011.
"Vaginal Microbiome of Reproductive-Age Women." Proceedings of the National
Academy of Sciences 108 (supplement_1): 4680–87.

Ravel, Jacques, Rebecca M. Brotman, Pawel Gajer, Bing Ma, Melissa Nandy, Douglas W.
Fadrosh, Joyce
Sakamoto, et al. 2013. "Daily Temporal Dynamics of Vaginal Microbiota before, during
and after Episodes of Bacterial Vaginosis." Microbiome 1 (1): 29.
https://doi.org/10.1186/2049-2618-1-29.

Robinson, Courtney J, Brendan JM Bohannan, and Vincent B Young. 2010. "From Structure to
Function: The Ecology of Host-Associated Microbial Communities." Microbiology and
Molecular Biology Reviews 74 (3): 453–76.

Saraf, Viqar Sayeed, Sohail Aslam Sheikh, Aftab Ahmad, Patrick M Gillevet, Habib Bokhari,
and Sundus Javed. 2021. "Vaginal Microbiome: Normalcy vs Dysbiosis." Archives of
Microbiology 203: 3793–3802.

Schooler, Shon S, Buck Salau, Mic H Julien, and Anthony R Ives. 2011. "Alternative Stable
States Explain Unpredictable Biological Control of Salvinia Molesta in Kakadu." Nature
470 (7332): 86–89.

Shade, Ashley, Hannes Peter, Steven D Allison, Didier L Baho, Mercè Berga, Helmut
Bürgmann, David H Huber, Silke Langenheder, Jay T Lennon, and Jennifer BH Martiny.
2012. "Fundamentals of Microbial Community Resistance and Resilience." Frontiers in
Microbiology 3: 417.

993   Staples, David F, Mark L Taper, and Bradley B Shepard. 2005. "Risk-based Viable Population

994   Monitoring." Conservation Biology 19 (6): 1908–16.

995   Taper, Mark L., Subhash R. Lele, José Miguel Ponciano, Brian Dennis, and Christopher L. Jerde.

996   2021. "Assessing the Global and Local Uncertainty of Scientific Evidence in the

997   Presence of Model Misspecification." Frontiers in Ecology and Evolution 9.

998   https://www.frontiersin.org/articles/10.3389/fevo.2021.679155.

999   Taper, Mark L, and José Miguel Ponciano. 2016. "Evidential Statistics as a Statistical Modern

1000   Synthesis to Support 21st Century Science." Population Ecology 58: 9–29.

1001   Tier, Charles, and Floyd B Hanson. 1981. "Persistence in Density Dependent Stochastic

1002   Populations." Mathematical Biosciences 53 (1–2): 89–117.

1003   White, Bryan A, Douglas J Creedon, Karen E Nelson, and Brenda A Wilson. 2011. "The Vaginal

1004   Microbiome in Health and Disease." Trends in Endocrinology & Metabolism 22 (10):

1005   389–93.

1006   Wilkinson, Ellen M, Paweł Łaniewski, Melissa M Herbst-Kralovetz, and Rebecca M Brotman.

1007   2019. "Personal and Clinical Vaginal Lubricants: Impact on Local Vaginal

1008   Microenvironment and Implications for Epithelial Cell Host Response and Barrier

1009   Function." The Journal of Infectious Diseases 220 (12): 2009–18.

1010   https://doi.org/10.1093/infdis/jiz412.

1011   Witkin, Steven S, and Iara Moreno Linhares. 2022. "The Medium Is the Message: Defining a

1012   'Normal' Vaginal Microbiome in Healthy Reproductive-Age Women." Reproductive

1013   Sciences, 1–6.

Zaoli, Silvia, and Jacopo Grilli. 2021. "A Macroecological Description of Alternative Stable

States Reproduces Intra-and Inter-Host Variability of Gut Microbiome." Science

Advances 7 (43): eabj2882.

Zeng, Zheng, Robert M Nowierski, Mark L Taper, Brian Dennis, and William P Kemp. 1998.

"Complex Population Dynamics in the Real World: Modeling the Influence of Time-

varying Parameters and Time Lags." Ecology 79 (6): 2193–2209.

Zhou, Jizhong, and Daliang Ning. 2017. "Stochastic Community Assembly: Does It Matter in

Microbial Ecology?" Microbiology and Molecular Biology Reviews 81 (4): e00002-17.

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

Figure 1. The abundances in stable (a) vs. unstable (b) populations. In both panels the grey lines

represent the log-population abundances at stationarity were simulated under the stochastic

Gompertz model of Ponciano et al. (2018) under the same environmental noise regime that are

shown in blue. The variance of the long run log-population abundances is equal to the ratio of the

environmental noise variance (here 0.11) to one minus the squared strength of density-

dependence c. This coefficient is stronger on the left than on the right. On the left $c = 0.75$ and

so the log-population size variance is $0.11/(1 - 0.75^2) = 0.2514$. On the right panel, density

dependence is much weaker, with a coefficient equal to 0.93. (Coefficients closer to 1 are close

to density-independence.) The variance of the population abundances under the same

environmental noise variance is approximately three times higher: $0.11/(1 - 0.93^2) = 0.8142$.

The magnitude of the response of a population to environmental noise, in terms of variability, is

modulated by c.

1050

Figure 2. This figure extends the simulation shown in Figure 1 to an instance with two or more

species for 70 days. It shows that changes in the nature and intensity of community interactions

directly affect the stochastic stability, measured using Ives' et al. four stability metrics:

VP,MR,VR and R which are defined and explained in the main text. In this case, the modulation

of a variable environment depends on the structure, as well as the nature and intensity of the

intraspecific and interspecific interactions. This figure shows the fluctuation in population sizes

of four different community structures (a-d) with three species, subject to the same

environmental noise regime. The upper row represents the four community structure types. The

intraspecific interactions (looped arrows) and interspecific interactions (straight arrows) change

in magnitude with weak interactions shown as thin arrows and strong interactions shown as thick

arrows. In the row directly below each of these interaction graphs we show the resulting

temporal dynamics of the abundances of each species. Since all four simulations were done

under the exact same environmental noise regime the differences in magnitude and fluctuation of

population abundances across community types can be directly attributed to differences in

structure. From left to right, it is shown that weaker interaction strengths lead to larger

fluctuations in populations under the same environmental variance. Note the different values of

the Y-axis.

1068

1069   **Figure 3.** Boxplots of all the pH measurements taken over 70 days for 88 women in our

1070   bacterial community time series data which were complete enough to estimate the MAR model

1071   parameters, as illustrated below. The boxplots here showcase the fact that the vaginal pH of 88

1072   healthy, asymptomatic women varied widely over 70 days, inside and outside what is considered

1073   to be a healthy vaginal pH region (shaded pink and region < 4.5).

1074

1075   **Figure 4.** Observed *Lactobacillus* species abundances with error (empty circles) vs. estimated

1076   abundances (black circles) after accounting for sampling error for one time series. The gray area

1077   shows the 95% confidence interval of the estimated true abundances.

1078

Figure 5. Principal component analysis (PCA) performed on the four stochastic stability metrics

estimated for the vaginal bacterial community time series data of 88 women. In this analysis the

samples (rows) correspond to each woman and the four columns (variables in the PCA analysis)

correspond to the four stability metrics estimated by fitting the MAR model of Ives et al. The

arrows' lengths and direction represent the strength of association of each one of these four

metrics with the principal component axes: the variance proportion, the mean return time and the

variance return time are highly associated with the first principal component while the reactivity

is highly associated with the second principal component. Using k-means clustering, the PCA

scores of these 88 bacterial communities were classified into four groups. Because lower values

in these stability metrics indicate higher stochastic stability, an examination of the magnitude of

these four metrics in each one of these four groups suggested the labeling of highly stable, stable,

unstable and highly unstable dynamics (see Supplementary material for details).

1091

1092   **Figure 6.** Population viability monitoring and estimating the temporally varying chances of

1093   *Lactobacillus* persistence. Illustrated is an example of Risk Prediction Monitoring (RPM) using

1094   stochastic population dynamics models.

1095

Figure 7. Projecting the probability of *Lactobacillus* persisting above 45% for two women,

starting at 50/50 from their carrying capacities for 20 days. The black dots are the projected

probabilities of persisting for woman 3 (on the left) and woman 60 (on the right), for 30 days.

As a background and in different colors, the same trend for all the other women in the study is

shown. The wide array of trajectories of all the trends for the other women emphasizes the wide

variability in predicted community dynamics.

1102

Figure 8. Probability of persisting above 45% of total abundances "t" days into the future. As

explained in the text, the changes in the estimated coefficients of the interactions matrix results

in restored dynamics and persistence probabilities. In panel A the effect of *Lactobacillus* on non-

Lactobacillus was changed from positive to negative and the effect of non-*Lactobacillus* on

Lactobacillus was switched from negative to positive. In panel B only the maximum growth rate

of the *Lactobacillus* species was increased by 25%.

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129   Figure 1.

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140   Figure 2.

1152   Figure 3.

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163   Figure 4.

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176   Figure 5.

1177

1178

1179

1180

1181

1182

1183   Figure 6.

1190   Figure 7.

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201   Figure 8.

W-60: Pr(persisting above 45% of total on day 't')

Days (t) since final observation

1202

1203