Focal cortical dysplasia type II-dependent maladaptive myelination in the human frontal lobe

Catharina Donkels[1], Susanne Huber[1], Theo Demerath[2], Christian Scheiwe[3], Mukesch J. Shah[3], Marcel Heers[4], Horst Urbach[2], Andreas Schulze-Bonhage[4,8], Marco Prinz[5,6], Ute Häussler[1,8], Andreas Vlachos[7,8], Jürgen Beck[3], Julia M. Nakagawa[3*], Carola A. Haas[1,8**]

[1] Experimental Epilepsy Research, Department of Neurosurgery, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

[2] Department of Neuroradiology, Medical Center-University of Freiburg, Faculty of Medicine, 79106 Freiburg, Germany

[3] Department of Neurosurgery, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

[4] Epilepsy Center Freiburg, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

[5] Institute of Neuropathology, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

[6] Signalling Research Centers BIOSS and CIBSS, University of Freiburg, Germany

[7] Department of Neuroanatomy, Institute of Anatomy and Cell Biology, Faculty of Medicine, University of Freiburg, Germany

[8] BrainLinks-BrainTools Center, University of Freiburg, Germany

*Equal contribution

Corresponding author: Dr. Catharina Donkels; email: catharina.donkels@uniklinik-freiburg.de

ORCID IDs: Catharina Donkels: 0000-0003-2636-3873; Theo Demerath: 0000-0002-5869-1110; Christian Scheiwe: 0000-0003-1695-8689; Mukesch J. Shah: 0000-0002-7224-9185; Marcel Heers: 0000-0001-5545-7014; Horst Urbach: 0000-0001-7264-4807; Andreas Schulze-Bonhage: 0000-0003-2382-0506; Marco Prinz: 0000-0002-0349-1955; Ute Häussler: 0000-0001-5601-9833; Andreas Vlachos: 0000-0002-2646-3770; Jürgen Beck: 0000-0002-7687-6098; Julia M. Nakagawa: 0000-0003-1071-5440; Carola A. Haas: 0000-0002-7022-4136

Abstract

Focal cortical dysplasias (FCDs) are local malformations of the human

neocortex and a leading cause of intractable epilepsy. FCDs are classified

into different subtypes including FCD IIa and IIb, characterized by a blurred

gray-white matter boundary or a transmantle sign indicating abnormal white

matter myelination. Recently, we have shown that myelination is also

compromised in the gray matter of FCD IIa of the temporal lobe. Since

myelination is key for brain function, we investigated whether deficient

myelination is a feature affecting also other FCD subtypes and brain areas.

Here, we focused on the gray matter of FCD IIa and IIb from the frontal lobe.

We applied *in situ* hybridization, immunohistochemistry and electron

microscopy to quantify oligodendrocytes, to visualize the myelination pattern

and to determine ultrastructurally the axon diameter and the myelin sheath

thickness. In addition, we analyzed the transcriptional regulation of myelin-

associated transcripts by real-time RT-qPCR and chromatin

immunoprecipitation (ChIP). We show that densities of myelinating

oligodendrocytes and the extension of myelinated fibers up to layer II were

unaltered in both FCD types but myelinated fibers appeared fractured mainly

in FCD IIa. Interestingly, both FCD types presented with larger axon

diameters when compared to controls. A significant correlation of axon

diameter and myelin sheath thickness was found for FCD IIb and controls,

whereas in FCD IIa large caliber axons were less myelinated. This was

mirrored by a down-regulation of myelin-associated mRNAs and by reduced

binding-capacities of the transcription factor MYRF to promoters of myelin-

associated genes. FCD IIb, however, had significantly elevated transcript

levels and MYRF-binding capacities reflecting the need for more myelin due

to increased axon diameters. These data show that FCD IIa and IIb are

characterized by divergent signs of maladaptive myelination which may

contribute to the epileptic phenotype and underline the view of separate

disease entities.

Keywords: neocortex, axon, oligodendrocyte, cortical malformation, epilepsy,

chromatin, transcriptional regulation, myelin sheath

Introduction

Focal cortical dysplasias (FCDs) are a frequent cause of pharmcoresistant focal epilepsy, especially in children [1, 2, 3]. Surgical resection of the lesion remains often the only therapeutic option, resulting in a seizure-free clinical outcome in many cases [4, 5]. FCDs comprise a heterogeneous group of cortical malformations morphologically characterized by disturbed lamination and cytoarchitecture including radial and/or horizontal alterations in lamination (FCD Ia or Ib) and the occurrence of dysmorphic neurons (FCD IIa) and/or balloon cells (FCD IIb) [6, 7]. Genetically, sporadic somatic mutations in the *MTOR* (mammalian target of rapamycin) gene and in mTOR pathway-related genes have been identified in FCD IIa and IIb cases leading to hyperactivation of mTOR signaling [8, 9, 10, 11, 12]. Recently, mutations in the GATOR1 complex, a GTPase-activating protein (GAP) complex that regulates TORC1 signaling by interacting with the Rag GTPase, have been found exclusively in FCD IIa of the frontal lobe [13], pointing out pathophysiological differences between FCD IIa and IIb. Accordingly, in the MRI many FCD IIa cases present with a blurring of the gray–white matter boundary [6, 7, 14], whereas FCD IIb shows a funnel-shaped formation through the white matter (WM) (transmantle sign) [15, 16], which is considered as abnormal WM myelination.

Correct myelination is very important for proper brain function and a highly dynamic process involved in memory consolidation in the adult brain [17]. There is recent evidence, that maladaptive myelination does also play a role in epileptic conditions [18, 19]. Indeed, recent reports show a loss of oligodendrocytes (OL) and myelination capacity in the white matter of FCD lesions [14, 20, 21, 22]. Likewise, deficits in myelination were found in the

gray matter of temporal lobe FCDs [23], where impaired proliferation of oligodendrocyte precursor cells (OPC) was part of the FCD pathology [24].

During the differentiation of OPCs into myelinating OLs, a number of transcription factors (TFs) are sequentially expressed to specify the OL fate [25, 26]. Depending on the differentiation stage, these TFs inhibit or stimulate the expression of myelin-associated genes [25]. Oligodendrocyte transcription factor 2 (OLIG2) is a prominent TF that initiates the differentiation of OPCs into myelinating OLs and is expressed across the entire lineage but with decreasing amounts toward the termination of differentiation [27, 28]. A key factor for the completion of OL differentiation is the myelin regulatory factor (MYRF) [29], [30], induced during terminal OL differentiation. MYRF positively regulates the transcription of myelin genes such as myelin basic protein (MBP), 2′, 3′-cyclic nucleotide-3′-phosphodiesterase (CNPase), myelin associated glycoprotein (MAG) and myelin oligodendrocyte glycoprotein (MOG) by binding to their promoters [29, 30].

In the current study, we focused on FCD IIa and IIb in the frontal lobe. There, we investigated the distribution of OLs, the myelination pattern, the ultrastructure of axons and myelin sheaths and the transcriptional regulation of myelin-associated genes in the gray matter. We present evidence for an FCD-type specific dysregulation of myelination: FCD IIa is characterized by a disorganization of the myelin fiber pattern, thinner myelin sheaths of large diameter axons and an attenuation of the myelin synthesis machinery. In contrast, FCD IIb presents with thicker axon diameters and myelin sheaths and an elevated transcriptional turnover of myelin-associated genes.

Methods

Patient Selection

A total of 22 surgically resected frontal lobe specimens from patients with

FCD IIa (n=9) [(age at surgery (years): mean: 16.78, range: 6 – 38), (epilepsy

duration (years): mean: 10.24, range: 3 – 34)], FCD IIb (n=9) [(age at surgery

(years): mean: 33.67, range: 12 – 48), (epilepsy duration (years): mean:

24.33, range: 9 – 41)] and epileptic but non-dysplastic controls (n=4)

[approach tissue from tumor resections (WHO grade I) or non-FCD lesions]

[(age at surgery (years): mean: 16.75, range: 10 – 26), (epilepsy duration

(years): mean: 8, range: 5 – 12)] were included in this study. Individual

patients were color-coded to facilitate their identification in the different assays

(Table 1).

All cases had undergone neurosurgical interventions due to

pharmacoresistant epilepsy or low-grade tumors and removal of cortical tissue

was clinically warranted to achieve seizure or tumor control. Pre-surgical

assessment included the documentation of a detailed history, neurological

examination, neuropsychological testing, MRI scanning and non-

invasive/invasive, long-term video EEG monitoring. Informed consent was

obtained from all individuals included in the study. All procedures received

prior approval by the institutional review board (Ethics Committee, Medical

Center - University of Freiburg) and were in accordance with the 1964

Helsinki declaration and its later amendments.

For histopathological diagnosis, all specimens were classified on

paraffin sections by the Institute of Neuropathology, Medical Center –

University of Freiburg) according to Blümcke et al. (2011) [6].

Tissue Preparation

Brain tissue was collected immediately after resection in ice-cold 0.1M phosphate buffer (PB), pH 7.4. For immunohistochemistry (IHC) and *in situ* hybridization histochemistry (ISHH), slices (~5 mm) were cut perpendicular to the brain's surface, fixed in 4% paraformaldehyde (PFA) [w/v] in PB (24 - 48h, 4°C), cryoprotected in 30% sucrose in PB (24 - 48h, 4°C) and finally stored at -80°C until further processing. For IHC and ISHH, cryostat sections (50 µm, orthogonal to the cortical surface) were prepared, collected in tissue culture dishes, and either rinsed in PB (IHC) or in 2× SSC (1× SSC = 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0) for ISHH.

For electron microscopy (EM), PFA-fixed frozen tissue was cut horizontally to the pia in a cryostat (60 µm) (Leica CM3050 S) and slices were post-fixed in 4% PFA/2.0% glutaraldehyde (GA) in PB (approx. 24h, 4°C) as described previously [23].

For molecular biological techniques the tissue was immediately frozen at -80°C and stored until further processing.

Immunohistochemistry

For NeuN, SMI32 and Vimentin immunolabeling a free-floating protocol was applied. Tissue slices were pretreated in 10% normal serum (NS) and 0.25% Triton X-100 for 30 min at RT and subsequently incubated in 0.1% Triton X-100 and 1% NS in PB with the following primary antibodies: mouse monoclonal anti-NeuN (1:1000; Millipore, #MAB377), anti-SMI32 (specific for non-phosphorylated neurofilament H, 1:1000; Biolegend #801701) and anti-

Vimentin (1:1000; Dako #M0725) at room temperature (RT) overnight. The

antibody binding was visualized by incubation with secondary antibodies

conjugated with Cyanine (Cy)-3 (1:400; Jackson ImmunoResearch

Laboratories) for 2h. Sections were mounted and coverslipped with Immu-

Mount (Thermo Scientific).

A modified IHC protocol published by Moreno-Jimerez et al. 2019 [31]

was used for the CNPase staining. Tissue sections were incubated in a 0.5%

sodium borohydride (NaBH$_4$, Sigma-Aldrich) solution for 30 min at RT. Next, a

heat-mediated 10 mM citrate buffer (pH 6.0) antigen retrieval step was

performed by boiling the tissue sections in a water bath for 20 min followed by

rinsing in 0.1 M PB and pretreatment in 0.5% bovine serum albumin (BSA)

and 0.5% Triton for 30 min at RT. Incubation with the CNPase antibody

(1:100, Cell Signaling Technology #5664S) was performed under gentle

shaking at 4 °C for 5 days followed by treatment with the Cy-3-labelled

secondary antibody (1:400; Jackson ImmunoResearch Laboratories) for 24h

at 4°C under constant shaking. Additionally, all sections were counterstained

for 10 min with 4,6-diamidino-2-phenylindole (DAPI,1:10000) to label nuclei.

Afterwards, slices were washed three times with PB, mounted on gelatinized

microscope slides and treated with an Autofluorescence Eliminator reagent

(EMD Millipore) for 5 min following the manufacturer's instructions. Finally, the

slices were washed three times with 70% ethanol and coverslipped in

aqueous mounting medium (ImmuMount; Thermo Scientific).

In situ Hybridization Histochemistry (ISHH)

Mbp mRNA expression was investigated by ISHH using digoxigenin (DIG)-labeled anti-sense riboprobes, generated from a pBluescript KS II$^+$ vector containing a 1.2 kb *mbp* cDNA as previously described [23].

In brief, cryostat sections (50 µm) were hybridized with a DIG-labeled anti-sense *mbp* cRNA riboprobe (100 ng/ml) at 55°C overnight followed by several stringent washing steps at 65 °C. DIG-labeled hybrids were detected with an anti-DIG antibody from sheep conjugated with alkaline phosphatase as recommended by the manufacturer (Roche) by incubating the slice overnight at 4°C. For signal detection, nitro blue tetrazolium chloride and 5-bromo-4-chloro-3-indolylphosphate were applied and the reaction was developed in the dark. Finally, sections were coverslipped and mounted in Kaiser's gelatin.

Microscopy and Cell counting

Fluorescence and bright field microscopy were performed with an *AxioImager2* microscope (Zeiss) using a 10x (NA 0.45) or 20x (NA 0.75) objective, (both Zeiss). Photomicrographs were taken as tile images with a digital camera (MR605) for fluorescence, AxioCam MRc5 for bright field (both Zeiss), and processed with *Zen* software (Zeiss).

Confocal images were taken using the *Fluoview* FV10i (Olympus) and a 63x oil objective applying the following parameters: image size 1024x1024 pixel, sensitivity 50% and laser power 20% for Cy3 and DAPI. The numerical aperture was selected 1, stacks were taken with 0.5 µm step sizes.

For the quantification *of mbp* mRNA-positive cells, bright field images

of three sections per patient were imported into *ImageJ* software. In each

image, two regions of interest (ROIs) covering all six cortical layers were

defined and within each ROI all digoxigenin-labeled cells were manually

counted using the integrated "cell counter" plugin (National Institutes of

Health).

For quantitative determination of OL densities, confocal image stacks

of CNPase-immunolabled sections were imported into *Image J*. OL cell bodies

were counted in three slices per patient and in each slice three ROIs of the

cortical layers V/VI (9 ROIs/patient) were manually counted by starting from

the first image, where cells appeared and stopped at the last countable

image. Cell numbers were related to the volume of the counted stacks and

given as a mean over all ROIs per patient (cell number/mm^3).

Electron Microscopy

For electron microscopy, slices were cut perpendicular to the pia from the

fixed, frozen block (60µm) and post-fixed in the presence of 4% PFA and 2%

glutaraldehyde for 24h at 4°C as described before [24]. Some sections were

stained with 1% cresyl violet to visualize the cortical layers, required for

orientation before embedding. The other slices were osmicated for 2 h with

1% osmium tetroxide in 6.86% sucrose in PB, subsequently contrasted with

1% uranyl acetate in 70% ethanol overnight at 4°C and dehydrated in graded

ethanol. The cortical gray matter was flat-embedded in epoxy resin (Durcupan

ACM, Sigma-Aldrich), which was polymerized overnight. Ultrathin sections (60

nm) in the horizontal plane were prepared with an ultramicrotome (LEICA

UC6) and collected on copper grids. Sections were analyzed with a TEM LEO

906E (Zeiss). Electron photomicrographs of layers V/VI were taken with the

CCD-Camera "sharp eye" (Tröndle) and visualized with ISP software

(Tröndle).

For quantitative analysis images were imported into *ImageSP* (TRS &

SYSPROG). With this software the outer and the inner diameter of myelinated

relation of the axon diameter to the total fiber diameter [32], was calculated.

RNA Isolation, Reverse Transcription and Real-Time RT-qPCR Analysis

The isolation of total RNA, reverse transcription, and real-time qPCR was

performed as described earlier [23]. In brief, frozen specimens were thawed

on ice and the white matter was carefully removed with a scalpel in *RNAlater*

(Qiagen). The total RNA was isolated from 30 mg cortical gray matter using

the *RNeasy Mini kit* (Qiagen) according to the manufacturer's guidelines.

Afterwards, a reverse transcription (RT) step was carried out with 1µg total

RNA per reaction using the *First Strand cDNA Synthesis Kit* (ThermoFisher

Scientific).

The expression of mRNAs was quantified by real-time RT-qPCR on a

CFX Connect^{TM} Real-Time PCR System (Bio-Rad Laboratories) in the

presence of *SsoAdvanced universal SYBR Green Mix* (Bio-Rad Laboratories).

The following human-specific primer pairs were used at 70 nM: *s12*

(mitochondrial ribosomal protein S12) (forward (f): 5'-

AGTTGGTGGAGGCCCTTTGT-3', reverse (r): 5'-

AGGCCTACCCATTCTCCTAGTTTC-3'), *mbp* (f: 5'-

AATTACTCACCGAGACACAC-3', r: 5'-GAAGGATATGGTCAGAAGAG-3'),

mag (f: 5'-CAACTGAGTCCAAGTTGTCT-3', r: 5'-
ATTCTGTACAGCGAGTGAGT-3'), *mog* (f: 5'-
CTTCGAGCAGAGATAGAGAA-3', r: 5'-TCGATGTAGCCAGTTGTAG-3'),
myrf (f: 5'-CTACCACATCCCTGTCAGT-3', r: 5'-GGAGGAGGAGTTCATCTG-
3'), CNP (f: 5'-ACACACAGCTTCTAGTGATTT-3', r: 5'-
ACAGAGAAGCGAGTTCAG-3'), *olig2* (f: 5'-AATCTCAATATCTGGGTC AA-
3', r: 5'-GGAACATCCACAGATTTATT-3'), *sip1* (Smad interacting protein 1) (f:
5'-AGCCTCTGTAGATGGTCCAGAAGAA-3', r: 5'-
GCGGTCTGGATCGTGGCTTC-3').

The PCR conditions were chosen as published before [23]. The
resulting Ct (cycle threshold) values (obtained by the CFX Manager™
Software, Bio-Rad Laboratories) were used to calculate relative expression
levels for genes of interest by normalization to the housekeeping gene S12.

Chromatin immunoprecipitation (ChIP)

The ChIP assays were performed using the *iDeal ChIP-seq* kit for
transcription factors (Diagenode) with fresh-frozen FCD and control tissue (30
mg per IP) according to manufacturer's guidelines. Firstly, the gray matter
was manually dissected and homogenized with 1% formaldehyde to cross-link
genomic DNA and proteins for 15 min at RT. The chromatin lysate was
sonicated and chromatin was fragmented to a size of approx. 500-1000 base
pairs using the *Bioruptor® Pico* (Diagenode) by running 10 cycles with
settings 30 sec. on, 30 sec. off per cycle. To reduce non-specific binding, we
included a preclearing step by incubating the chromatin lysate with *DiaMag*
Protein A-coated magnetic beads (Diagenode) at 4°C for 1.5 - 2 hours, Next,

the chromatin was precipitated with polyclonal rabbit MYRF (5 µg, Millipore),

and polyclonal goat OLIG2 (5 µg, R&D) antibodies, all pre-incubated for 4h at

4°C with Protein A-coated magnetic beads, and non-specific polyclonal rabbit

IgG (1 µg, Diagenode), overnight at 4°C under constant rotation. Afterwards,

the chromatin was stringently washed, eluted from the beads and the

crosslinking was reversed by incubating the eluate at 65°C overnight.

Subsequently, the DNA was purified using *IPure beads v2* (Diagenode) eluted

with H_2O and stored at -20°C until further use.

ChIP specificity was verified by real-time qPCR for known transcription

factor binding sites (e.g., *MAG*; *MOG*, *MBP*). The PCR primers were designed

around predicted transcription factor binding sites (BS), which were identified

by using the *MatInspector* tool form *Genomatix* software. Using this tool, we

detected the following transcription factor BS sequences: *MBP* (MYRF BS):

5'-TTTCCTGGTACAT-3', *SIP 1* (OLIG2 BS): 5'-AGGCCAGCTGTTTCA-3',

CNP (OLIG2 BS): 5'-AGCCCCAGCTGCTCTG-3' and *MBP* (OLIG2 BS): 5'-

AAAGCAGCTGCCTGG-3'. The following primers were used for ChIP-qPCR

analysis: *MBP* (MYRF ChIP): (f: 5'-GTCTGCTTGTCTGTTTCCCA-3', r: 5'-

TCCGGGGATTGACTTTCAGA-3'), *SIP1* (OLIG2 ChIP): (f: 5'-

TCCTCATGGAACTTGAGTCG-3', r: 5'-AAAACCCTGATATCCTGACAGC-3'),

CNP (OLIG2 ChIP): (f: 5'-GAA CCTGGACCAAAGCTGAC-3', r: 5'-

TCTGAGGTGTCTCCCCACTC-3') and *MBP* (OLIG2 ChIP): (f: 5'-

TCGTGTGCAGTGCTTAGACC-3', r: 5'-TTTCATTCAGCCTGTGCTTG-3').

The PCR was run with the following conditions: 1. Hotstart: 3 min 95°C, 2.

Amplification: 30 sec 95°C, 30 sec 63°C, 30 sec 72°C (40 cycles). ChIP-

qPCR was analyzed by calculating the relative amount of immunoprecipitated

DNA compared to INPUT DNA for the control regions (% of recovery) using

the following formula: % recovery = $2^{\wedge}[(Ct_{input} - 6.64) - Ct_{sample})] \times 100\%$.

Statistical analysis

Prism 8 software (GraphPad Software Inc.) was used for statistical analysis.

For all experimental values, mean and standard error of the mean (SEM) are

given. Shapiro-Wilk normality test was applied for distribution testing. Group

comparisons were performed with one-way ANOVA or Kruskal-Wallis test and

Tukey's or Dunn's multiple comparison post hoc test. Significance thresholds

were set to: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. Correlations of axon diameter

with myelin sheath thickness, g-ratio and mRNA expression levels with

transcription factor binding capacities were tested using Pearson's correlation.

Data for group comparisons for cumulative frequencies were tested for

normality and logarithmically transformed before applying the one-way

ANOVA and Tukey's multiple comparison post hoc test.

Results

MR imaging and histopathological analysis of FCD type II cases

All patients included in this study were scanned by MRI with either FLAIR- or

STIR-weighted sequences according to a standard protocol to detect potential

lesions [33, 34, 35]. The FCD IIa group was heterogeneous with respect to a

detectable lesion: five patients showed a thickening of the cortex and blurring

of the gray-white matter boundary (Fig. 1a, b), one case had a transmantle

sign and three were MRI-negative (Table 1). In contrast, all patients of the

FCD IIb cohort presented with a transmantle sign, a zone of increased signal

intensity tapering toward the lateral ventricle (Fig. 1c, d; Table 1).

312   Since FCDs can be variable in extensions and dimensions, we

characterized every resected specimen included in this study regarding the

lamination pattern and cytoarchitecture and determined the FCD type

according to the ILAE classification [6]. To this end, serial sections of all cases

were stained with cresyl violet and were immunolabeled for NeuN to visualize

all layers, or for SMI32, a monoclonal antibody specific for non-

phosphorylated neurofilament H [36], which labels pyramidal cells of layers III

and V.

320   In controls, all six cortical layers were present as shown by Nissl

staining or NeuN labeling (Fig. 2a, b). Pyramidal cells in layers III and V

appeared normally shaped with parallel-orientated apical dendrites, which

reached the border of layer I (Fig. 2c). In the FCD IIa cohort, Nissl and NeuN

stainings showed a mild hypercolumnization across layers II to V (Fig. 2d, e).

Moreover, the occurrence of dysmorphic neurons, characterized by

malorientation and the strong accumulation of non-phosphorylated

neurofilament was apparent, in particular, in layers III and V/VI (Fig. 2f, j).

FCD IIb cases were characterized by blurring of the cortical layers (Fig. 2g, h),

the presence of dysmorphic neurons in all layers and at the gray-white matter

boundary (Fig. 2i, k). Balloon cells, characteristic for FCD IIb, were

recognizable in the cortical white matter by vimentin labeling (Fig. 2l).

332   Altogether, by applying presurgical MRI and histological

333   characterization of all specimens, we ensured the characteristic phenotype of

334   the two FCD cohorts and of the control group.

Distribution of myelinating OLs in the gray matter of the frontal lobe

Inspired by our pervious results from the temporal lobe, where we had found FCD-associated myelination deficits and reduced numbers of myelinating OLs [23, 24], we started the analysis of myelination in frontal lobe FCD IIa and IIb with studying the distribution of mature OLs, the myelin-producing cells. To this end, we performed ISHH for *mbp* mRNA, which is exclusively expressed in mature, myelinating oligodendrocytes [37] (Fig. 3). In controls, *mbp* mRNA-expressing OLs were most numerous in the infragranular layers of the gray matter and thinned out gradually up to layer I (Fig. 3b, c). In FCD IIa, less OLs were detectable; some, located in the deep cortical layers, exhibited very strong *mbp* mRNA signals (Fig. 3e, f). In contrast, in FCD IIb *mbp* mRNA-expressing OLs appeared slightly more numerous across the entire cortex when compared to controls and FCD IIa cases (Fig. 3h, i). However, counting of *mbp*-positive cells did not reveal any significant differences between controls, FCD IIa and FCD IIb [controls (n=4): 178.9 ± 42.1 cells/mm²; FCD IIa (n=7) 129.8 ± 24.9 cells/mm²; FCD IIb (n=7): 246.9 ± 43.4 cells/mm^2; (one-way ANOVA F=2.64, p=0.106)] (Fig. 3j).

Altogether, in contrast to the temporal lobe, the density of myelinating OLs was unaltered in FCD type II of the frontal lobe.

Myelination pattern and ultrastructure of myelinated axons in the gray matter

To investigate the architecture of myelinated fibers in the gray matter of control and FCD type II specimens we used immunolabeling for CNPase, a marker for pre-OLs, mature OLs and myelin [38].

In control tissue, a dense, radially aligned network of CNPase-positive

myelinated fibers extended from the gray-white matter boundary up to layer

III, declining gradually from the white matter to the upper layers (Fig. 4a). In

FCD IIa and IIb, a similar pattern of CNPase-immunolabeled fibers was

observed, reaching up to layer III as in the non-dysplastic cortex (Fig. 4b, c).

High resolution confocal imaging in layers V/VI demonstrated radially aligned

fibers in the control case and in FCD IIb (Fig. 4d, f) but disorganized and

fractured strands in FCD IIa (Fig. 4e). Counting of CNPase-labelled pre-OL

and OL cell bodies across all layers did not reveal any significant differences

between controls, FCD IIa and FCD IIb cases [controls (n=4): 34380 ± 3259

cells/mm^3; FCD IIa (n=9) 38674 ± 5211 cells/mm^3; FCD IIb (n=9): 44580 ±

5457 cells/mm^3; (one-way ANOVA F=0.74, p=0.49)] (Fig. 4g).

Next, we compared the ultrastructure of myelinated axons of control

and FCD cases by transmission electron microscopy in layers V/VI (Fig. 5a-c),

where axons show a high and homogeneous myelin coverage [39]. We

determined the axon diameter, the myelin sheath thickness and the g-ratio,

defined as the ratio of the axon diameter to the total fiber diameter [32].

Mean axon diameters were slightly increased in both FCD cohorts

when compared to the control group [(axon diameters: control: 323.9 ± 27.7

nm; FCD IIa: 424.8 ± 23.8 nm; FCD IIb: 436.1 ± 55.5 nm); (one-way ANOVA

F=2.59, p=0.129)] (Fig. 5d). When we looked at the frequency distribution of

all axon diameters within each group, we found a significant shift towards

larger axon diameters in both FCD types [controls (n=4; 114 axons) 119.9.8 –

767.5 nm; FCD IIa (n=4; 192 axons) 129.5 – 1141.5 nm; FCD IIb (n=4; 116

axons) 151.4 – 1346.2 nm] (one-way ANOVA F=19.26, p<0.0001; Tukey's

post hoc test: control vs. FCD IIa p<0.0001, control vs. FCD IIb p<0.0001,

FCD IIa vs. FCD IIb p=0.691)] (Fig. 5e). The mean g-ratio of myelinated

axons was elevated in FCD IIb and significantly increased in FCD IIa when

compared to non-dysplastic controls [(g-ratios: control: 0.698 ± 0.008; FCD

IIa: 0.744 ± 0.017; FCD IIb: 0.728 ± 0.012); (one-way ANOVA F=4.511,

p=0.044; Tukey's post hoc test: control vs. FCD IIa p=0.039, control vs. FCD

IIb p=0.174, FCD IIa vs. FCD IIb p=0.611)] (Fig. 5f). This was even more

prominent when we looked at the relative frequency distribution of all g-ratio

values: both FCD II groups had significantly increased g-ratios in comparison

to the control group and FCD IIa showed significantly higher values than FCD

IIb [(one-way ANOVA F=14.69, p<0.0001; Tukey's post hoc test: control vs.

FCD IIa p<0.0001, control vs. FCD IIb p=0.019, FCD IIa vs. FCD IIb p=0.046)]

(Fig. 5g). Normally, an increased g-ratio indicates a reduced thickness of the

myelin sheath, based on the assumption that the axon diameters are not

altered. However, since our FCD cases had significantly larger axon

diameters, we investigated also the myelin sheath thicknesses. On average,

there was no difference between the three groups [(myelin sheath thickness:

control: 133.5 ± 12.39 nm; FCD IIa: 132.7 ± 4.312 nm; FCD IIb: 147.3 ± 9.828

nm); (one-way ANOVA F=0.746, p=0.501)] (Fig. 5h). However, plotting the

relative frequency distribution of all individual values revealed that FCD IIb

had significantly thicker myelin sheaths when compared to controls and FCD

IIa cases [(one-way ANOVA F=4.487, p=0.012; Tukey's post hoc test: control

vs. FCD IIa p= 0.941, control vs. FCD IIb p=0.023, FCD IIa vs. FCD IIb

p=0.024)] (Fig. 5i) but also thicker axons than controls.

So solve this controversy, we correlated all individual axon diameters

with the corresponding myelin sheath thickness for the three groups

separately: (in nm): controls (range: smallest axon diameter/myelin sheath

119.92/91.59; largest axon diameter/myelin sheath 767.5/130.77); FCD IIb

(range: smallest axon diameter/myelin sheath 151.41/103.22, largest axon

diameter/myelin 1346.19/146.09). We found significant correlations between

axon diameter and myelin sheath thickness in controls (Pearson's correlation:

r=0.329, p=0.0003) and FCD IIb (Pearson's correlation: r=0.309, p=0.0007)

(Fig. 5j, l), implying that the axon caliber determines the thickness of the

myelin sheath [40]. This was not the case for FCD IIa (Pearson's correlation:

r=0.138, p=0.056) (range: smallest axon diameter/myelin sheath

129.48/124.41, largest axon diameter/myelin sheath 1141.5/92.95), where

large caliber axons (above 700 nm) had thinner myelin sheaths than axons of

the same caliber in controls or FCD IIb (Fig. 5k) indicating that myelination

was selectively compromised for axons with a large diameter in FCD IIa.

Transcriptional regulation of OL differentiation factors and myelin-associated

genes

Since thinner myelin sheaths may indicate a disturbance in OL differentiation

and/or myelin synthesis, we next tested whether the transcriptional regulation

of OL differentiation and myelination was affected in our FCD cohorts. First,

we performed real-time RT-qPCR to measure the mRNA levels of

transcription factors either sequentially active at different stages of OL

maturation (SIP1, MYRF) or continuously expressed (OLIG2). In addition, we

quantified the expression levels of *mbp*, *mag* and *mog* mRNA, which encode

proteins that are all components of the myelin sheath.

Olig2 (Kruskal-Wallis p=0.793) and *sip1* (one-way ANOVA F=0.648, p=0.534)

mRNA levels were similar in all groups (Fig. 6a, b), whereas *myrf* mRNA

expression was significantly down-regulated in FCD IIa when compared to

FCD IIb (one-way ANOVA F=7.914, p=0.003; Tukey's post hoc test: control

vs. FCD IIa p=0.389, control vs. FCD IIb p=0.205, FCD IIa vs. FCD IIb

p=0.002) (Fig. 6c). Likewise, *mbp* (one-way ANOVA F=5.964, p=0.009;

Tukey's post hoc test: control vs. FCD IIa p= 0.672, control vs. FCD IIb

p=0.190, FCD IIa vs. FCD IIb p=0.008)*, mag* (one-way ANOVA F=11.46,

p=0.0005; Tukey's post hoc test: control vs. FCD IIa p=0.288, control vs. FCD

IIb p=0,099, FCD IIa vs. FCD IIb p=0.0004)*, and mog* (one-way ANOVA F=

7.551, p=0.004; Tukey's post hoc test: control vs. FCD IIa p=0.163, control vs.

FCD IIb p=0.515, FCD IIa vs. FCD IIb p=0.003)*,* mRNA levels were also

significantly decreased in FCD IIa in comparison to FCD IIb (Fig. 6d-f)

reflecting the regulatory action of MYRF on myelin genes.

To target the level of transcriptional regulation directly, we next

performed ChIP assays which give insight into transcription factor binding to

specific promoters. Chromatin was purified from gray matter of control, FCD

IIa and IIb specimens and immunoprecipitated with antibodies specific for

transcription factors which either determine the OL lineage (OLIG2) or the

advanced differentiation stage (MYRF). As transcription factor targets, we

chose either genes encoding transcription factors (*SIP1, SOX10*) or structural

proteins of the myelin sheath (*MBP*, *MAG*, *MOG*, *CNP*) and quantified the

OLIG2 and MYRF binding capacities to the predicted binding sites by real-

time qPCR.

We did not find any differences in the binding capacities of OLIG2 to

the *SIP1* (one-way ANOVA F=0.153, p=0,859), *MBP* (one-way ANOVA

F=0.247, p=0.784), *CNP* (one-way ANOVA F=0.654, p=0.531) and *MOG*

(Kruskal-Wallis p=0.475) promoters between control (n=4), FCD IIa (n=9) and

FCD IIb (n=9) cases indicating that OLIG2 is equally active in the different

groups (Fig. 7a-d). In contrast, MYRF showed a significantly decreased

affinity to the *SOX10* (one-way ANOVA F=10.51, p=0.0008; Tukey's post hoc

test: control vs. FCD IIa p=0.0009, control vs. FCD IIb p= 0.112, FCD IIa vs.

FCD IIb p= 0.024), *MBP* (one-way ANOVA F=6.55, p=0.007; Tukey's post hoc

test: control vs. FCD IIa p=0.032, control vs. FCD IIb p= 0.976, FCD IIa vs.

FCD IIb p= 0.011) and *MAG* (Kruskal-Wallis p=0.003; Dunn's multiple

comparison test: control vs. FCD IIa p=0.017, control vs. FCD IIb p>0.999,

FCD IIa vs. FCD IIb p=0.041) promoters in FCD IIa specimens when

compared to FCD IIb or controls (Fig. 7e-g) indicating an FCD IIa-specific

transcriptional dysregulation of myelin-associated genes.

Finally, we tested, whether there is a direct impact of the investigated

transcription factor binding capacities on the corresponding mRNA

expressions. We correlated the promoter binding capacities of OLIG2 and

MYRF with the corresponding *sip1*, *mbp*, *mog*, and *mag* mRNA levels. There

were no differences in the promoter binding capacities of OLIG2 to their target

promoters (data not shown). However, we found a significant correlation of

the MYRF binding capacity with expression levels of *mbp* (Pearson's

correlation: r=0.546, p=0.009) and *mag* (Pearson's correlation: r=0.539,

p=0.009) transcripts in FCD IIb but less in FCD IIa (Fig. 7h, i) mirroring the

low expression of myelin-associated genes in FCD IIa.

Discussion

Insulation of axons by myelin is important for fast and long-range signal

propagation in the brain. Myelin is constantly produced in the adult brain by

oligodendrocytes in a plasticity-dependent manner and necessary for memory

consolidation [17]. Myelination deficiencies have been found in several

neurologic disorders in particular in multiple sclerosis [41] and are often

associated with seizure occurrence [42]. There is increasing awareness that

also in epilepsy maladaptive myelination or axon demyelination are of high

relevance for seizure generation [19, 18].

In this study, we investigated maladaptive myelination and the

transcriptional regulation of myelin-associated genes in FCD specimens

obtained from patients with pharmacoresistant focal epilepsy. We focused on

the gray matter of FCD IIa and IIb cases from the human frontal lobe, all

thoroughly characterized by presurgical MR scans and *post hoc* histological

analysis. In the MRT, most FCD IIa patients showed a thickening of the

neocortex and blurring of the gray-white matter boundary, whereas all FCD IIb

cases displayed a transmantle sign [15]. Histological characterization ensured

that all specimens of our study fulfilled the morphological criteria proposed by

the ILAE [6, 7].

The extension of radially orientated, myelinated CNPase-positive

fibers, was similar in FCD cases and controls, reaching up to layer III as

described earlier for the temporal cortex [43, 24]. However, high resolution

confocal microscopy in layer V/VI revealed fractured and distorted myelinated

axons mainly in FCD IIa specimens, in line with our previous study in the gray

matter of the temporal lobe [23].

When we analyzed myelinated axons ultrastructurally and determined

axon diameter, myelin sheath thickness and g-ratio, we found a significantly

increased g-ratio for FCD IIa cases, pointing to a reduced thickness of the

myelin sheath as found in the temporal lobe [24]. However, the frequency

distribution of axon diameters indicated a significant increase of axon calibers

in both FCD cohorts in comparison to controls. Unfortunately, there is hardly

any information about the physiological range of axon diameters in the human

gray matter, since most studies focused on fiber tracts in the white matter.

Liewald et al. [44] reported a frequency distribution of 300 – 1500 nm for axon

diameters in the human white matter. We found much thinner axons (119 –

767 nm) in controls but both FCD cohorts approached values (FCD IIa: 130 –

1142 nm; FCD IIb: 151 – 1346 nm) similar to those reported by Liewald et al.

[44]. Since the axon diameter depends strongly on the soma size of the

parent neuron [45], it is tempting to speculate that the presence of large

dysmorphic neurons, being part of the FCD II pathology, could be the reason

for larger axon diameters in our FCD cases in comparison to control tissue.

Since the axon diameter is critical for the calculation of the g-ratio [32],

we performed a more refined analysis of all individual values and correlated

the corresponding axon diameters and myelin sheath thicknesses. We found

a clear correlation of both parameters in FCD IIb and controls, showing a

linear relationship between axon diameter and myelin sheath thickness as

reported in the literature [40]. This was not the case in FCD IIa specimens,

where large diameter axons (≥ 700 nm) had thinner myelin sheaths than

corresponding axons of FCD IIb and controls indicating a myelination

deficiency of large diameter axons only in the FCD IIa cohort.

What could be the explanation for thinner myelin sheaths in FCD IIa? It

might be that the differentiation of OLs, the myelin sheath-producing glial

cells, and their myelin-synthesizing capacities are compromised. In fact, a

loss of OPCs and OLs [14, 21] and decreased myelin synthesis, shown in an

in vitro myelination assay [22] has been demonstrated for the white matter of

FCD cases. Similarly, we had found significantly less OPCs and a reduced

proliferation capacity of OPCs purified from the gray matter of temporal FCD

IIa [24].

In the current study, we did not find altered numbers of *mbp* mRNA- or

CNPase-positive pre-OLs and OLs in both FCD II types suggesting that the

OL lineage is basically preserved in the dysplastic gray matter of the frontal

lobe. This was supported by constant levels of *olig2* and *sip1* mRNA, which

encode transcription factors required for the maturation of OLs [46, 47] and by

unaltered binding capacities of OLIG2 to genes encoding structural myelin

proteins as shown in our ChIP assays.

Alternatively, the mature OL population might have an impaired

myelination capacity in FCD type II. Indeed, we found significantly reduced

myrf mRNA levels in FCD IIa pointing to an attenuation of myelin production in

mature OLs of this FCD type. MYRF is a key component of the transcriptional

regulatory network responsible for the induction and maintenance of the

myelinating OL phenotype [48]. Accordingly, myelin-associated genes such as

MBP, *MAG* and *MOG* were significantly down-regulated in FCD IIa specimens

in comparison to FCD IIb. The latter had even higher mRNA levels of myelin

components than controls, in agreement with an elevated need for myelin

proteins due to thicker axons and myelin sheaths. This was also reflected on

the level of transcriptional regulation analyzed by our ChIP assays: all

investigated promotor-binding capacities of MYRF to the SOX10, MBP and

MAG promoters were significantly elevated in FCD IIb when compared to

FCD IIa indicating an enhanced myelin production. This was further supported

by the significant correlation between MYRF promotor binding capacities and

mag and *mbp* mRNA levels. Altogether, our results point to divergent forms of

maladaptive myelination in FCD IIa and IIb.

What are the functional effects of an impaired myelination? Under

physiological conditions, myelination depends strongly on neuronal activity

[49]. In rodents, there is recent evidence that neuronal activity increases OL

differentiation and myelin sheath thickness in a pathway-specific manner [50].

Accordingly, Knowles et al. [18] found increased oligodendrogenesis and

myelination in an animal model for absence seizures and demonstrated that it

contributes to epilepsy progression. In contrast, we found in our human FCD

IIa cases thinner myelin sheaths in layer V/VI of the temporal lobe [24] and, as

shown here, of the frontal lobe. We did not find any correlation with seizure

frequency or duration (data not shown) indicating that different epilepsy types

result in different forms of maladaptive myelination.

CONCLUSIONS

Our study demonstrates for the first time that there is a dysregulation of

myelination in the gray matter of frontal lobe FCD IIa and FCD IIb. Both FCD

types show divergent signs of maladaptive myelination: FCD IIa is

characterized by a disorganization of the myelin fiber pattern, thinner myelin

sheaths of large diameter axons and an attenuation of the myelin synthesis

machinery. In contrast, FCD IIb presents with thicker axon diameters and

myelin sheaths and an elevated transcriptional turnover of myelin-associated

genes. This imbalance in myelination may disrupt the equilibrium of signal

conduction and thereby contribute to the epileptic phenotype.

Abbreviations

ChIP: Chromatin immunoprecipitation

CNPase: 2',3'-cyclic nucleotide 3'-phosphodiesterase

EM: Electron microscopy

FCD: Focal cortical dysplasia

IHC: Immunohistochemistry

ISHH: *In situ* hybridization histochemistry

MAG: Myelin-associated glycoprotein

MBP: Myelin basic protein

MOG: Myelin oligodendrocyte glycoprotein

MRI: Magnetic resonance imaging

MTOR: mammalian target of rapamycin

MYRF: Myelin regulatory factor

OL: Oligodendrocyte

OLIG2: Oligodendrocyte transcription factor 2

OPC: Oligodendrocyte precursor cell

PB: phosphate buffer

SIP1: Smad interacting protein 1

SOX10: SRY-box transcription factor 10

TF: Transcription factor

Declarations

Ethics approval and consent to participate

All procedures received prior approval by the institutional review board (Ethics

Committee, Medical Center - University of Freiburg) and were in accordance

with the 1964 Helsinki declaration and its later amendments. Informed

consent was obtained from all individuals included in the study.

Availability of data and material

The data that support the findings of this study are available from the

corresponding author upon written request.

Competing interest

The authors declare no conflict of interest.

Funding

German Research Foundation (grant number: DO 2542/1-1), Research

Commission, Medical Faculty - University of Freiburg (grant DON 1207/19).

Authors' contributions

CD: Conceptualization; data curation; formal analysis; project administration;

visualization; writing - original draft; funding acquisition; SH: Visualization;

formal analysis; CS, MJS: Neurosurgical brain resection; TD, MH, ASB, HU,

MP, JB: Clinical data; UH: Statistical analysis; commenting manuscript; AV:

Electron microscopy; JMN: Brain resection, clinical data from patients; CAH:

Conceptualization; data interpretation, manuscript writing, resources.

Acknowledgement

We thank all the patients who donated tissue for this study. We are grateful to

Sigrun Nestel for excellent technical assistance and to Dr. Maria Stella Carro

for help with establishing the chromatin immunoprecipitation method.

References

1. Palmini A, Andermann F, Olivier A, Tampieri D, Robitaille Y, Andermann E,

et al. Focal neuronal migration disorders and intractable partial epilepsy: A

study of 30 patients. Ann. Neurol. 1991;30:741–9.

2. Fauser S, Schulze-Bonhage A, Zentner J. Focal cortical dysplasias:

surgical outcome in 67 patients in relation to histological subtypes and dual

pathology. Brain. 2004;127:2406–18.

3. Fauser S, Huppertz H-J, Schulze-Bonhage A. Clinical characteristics in

focal cortical dysplasia: a retrospective evaluation in a series of 120 patients.

Brain. 2006;129:1907–16.

4. Zhao B, Zhang C, Wang X, Wang Y, Liu C, Mo J, et al. Sulcus-centered

resection for focal cortical dysplasia type II: surgical techniques and

outcomes. J. Neurosurg. 2020;135:266–72.

5. Willard A, Antonic-Baker A, Chen Z, O'Brien TJ, Kwan P, Perucca P.

Seizure Outcome After Surgery for MRI-Diagnosed Focal Cortical Dysplasia:

A Systematic Review and Meta-analysis. Neurology [Internet]. 2022 [cited

2024   Jan   29];98.   Available   from:

https://www.neurology.org/doi/10.1212/WNL.0000000000013066

6. Blümcke I, Thom M, Aronica E, Armstrong DD, Vinters HV, Palmini A, et al.

The clinicopathologic spectrum of focal cortical dysplasias: A consensus

classification proposed by an ad hoc Task Force of the ILAE Diagnostic

Methods Commission1: The ILAE Classification System of FCD. Epilepsia.

2011;52:158–74.

7. Najm I, Lal D, Alonso Vanegas M, Cendes F, Lopes-Cendes I, Palmini A, et

al. The ILAE consensus classification of focal cortical dysplasia: An update

proposed by an ad hoc task force of the ILAE diagnostic methods

commission. Epilepsia. 2022;63:1899–919.

8. Abdijadid S, Mathern GW, Levine MS, Cepeda C. Basic Mechanisms of

Epileptogenesis in Pediatric Cortical Dysplasia. CNS Neurosci Ther.

2015;21:92–103.

9. Lim JS, Kim W, Kang H-C, Kim SH, Park AH, Park EK, et al. Brain somatic

mutations in MTOR cause focal cortical dysplasia type II leading to intractable

epilepsy. Nat Med. 2015;21:395–400.

10. Baldassari S, Ribierre T, Marsan E, Adle-Biassette H, Ferrand-Sorbets S,

Bulteau C, et al. Dissecting the genetic basis of focal cortical dysplasia: a

large cohort study. Acta Neuropathol. 2019;138:885–900.

11. Nguyen LH, Mahadeo T, Bordey A. mTOR Hyperactivity Levels Influence

the Severity of Epilepsy and Associated Neuropathology in an Experimental

Model of Tuberous Sclerosis Complex and Focal Cortical Dysplasia. J

Neurosci. 2019;39:2762–73.

12. Chung C, Yang X, Bae T, Vong KI, Mittal S, Donkels C, et al.

Comprehensive multi-omic profiling of somatic mutations in malformations of

cortical development. Nat Genet. 2023;55:209–20.

13. Honke J, Hoffmann L, Coras R, Kobow K, Leu C, Pieper T, et al. Deep

histopathology genotype–phenotype analysis of focal cortical dysplasia type II

differentiates between the GATOR1-altered autophagocytic subtype IIa and

MTOR-altered migration deficient subtype IIb. Acta Neuropathol. Commun.

2023;11:179.

14. Mühlebner A, Coras R, Kobow K, Feucht M, Czech T, Stefan H, et al.

Neuropathologic measurements in focal cortical dysplasias: validation of the

ILAE 2011 classification system and diagnostic implications for MRI. Acta Neuropathol. 2012;123:259–72.

15. Urbach H, Scheffler B, Heinrichsmeier T, Von Oertzen J, Kral T, Wellmer J, et al. Focal Cortical Dysplasia of Taylor's Balloon Cell Type: A Clinicopathological Entity with Characteristic Neuroimaging and Histopathological Features, and Favorable Postsurgical Outcome. Epilepsia. 2002;43:33–40.

16. Kimura Y, Shioya A, Saito Y, Oitani Y, Shigemoto Y, Morimoto E, et al. Radiologic and Pathologic Features of the Transmantle Sign in Focal Cortical Dysplasia: The T1 Signal Is Useful for Differentiating Subtypes. AJNR Am J Neuroradiol. 2019;40:1060–6.

17. Steadman PE, Xia F, Ahmed M, Mocle AJ, Penning ARA, Geraghty AC, et al. Disruption of Oligodendrogenesis Impairs Memory Consolidation in Adult Mice. Neuron. 2020;105:150-164.e6.

18. Knowles JK, Xu H, Soane C, Batra A, Saucedo T, Frost E, et al. Maladaptive myelination promotes generalized epilepsy progression. Nat Neurosci. 2022;25:596–606.

19. De Curtis M, Garbelli R, Uva L. A hypothesis for the role of axon demyelination in seizure generation. Epilepsia. 2021;62:583–95.

20. Shepherd C, Liu J, Goc J, Martinian L, Jacques TS, Sisodiya SM, et al. A

quantitative study of white matter hypomyelination and oligodendroglial

maturation in focal cortical dysplasia type II. Epilepsia. 2013;54:898–908.

21. Scholl T, Mühlebner A, Ricken G, Gruber V, Fabing A, Samueli S, et al.

Impaired oligodendroglial turnover is associated with myelin pathology in focal

cortical dysplasia and tuberous sclerosis complex. Brain Pathol. 2017;27:770–

80.

22. Gruber V, Lang J, Endmayr V, Diehm R, Pimpel B, Glatter S, et al.

Impaired myelin production due to an intrinsic failure of oligodendrocytes in

mTORpathies. Neuropathol. Appl. Neurobiol. 2021;47:812–25.

23. Donkels C, Pfeifer D, Janz P, Huber S, Nakagawa J, Prinz M, et al. Whole

Transcriptome Screening Reveals Myelination Deficits in Dysplastic Human

Temporal Neocortex. Cereb Cortex. 2016;bhv346.

24. Donkels C, Peters M, Fariña Núñez MT, Nakagawa JM, Kirsch M, Vlachos

A, et al. Oligodendrocyte lineage and myelination are compromised in the

gray matter of focal cortical dysplasia type IIa. Epilepsia. 2020;61:171–84.

25. Emery B. Regulation of Oligodendrocyte Differentiation and Myelination.

Science. 2010;330:779–82.

26. Sock E, Wegner M. Using the lineage determinants Olig2 and Sox10 to

explore transcriptional regulation of oligodendrocyte development. Dev.

Neurobiol. 2021;81:892–901.

27. Mei F, Wang H, Liu S, Niu J, Wang L, He Y, et al. Stage-Specific Deletion

of Olig2 Conveys Opposing Functions on Differentiation and Maturation of

Oligodendrocytes. J Neurosci. 2013;33:8454–62.

28. Meijer DH, Sun Y, Liu T, Kane MF, Alberta JA, Adelmant G, et al. An

Amino Terminal Phosphorylation Motif Regulates Intranuclear

Compartmentalization of Olig2 in Neural Progenitor Cells. J Neurosci.

2014;34:8507–18.

29. Emery B, Agalliu D, Cahoy JD, Watkins TA, Dugas JC, Mulinyawe SB, et

al. Myelin Gene Regulatory Factor Is a Critical Transcriptional Regulator

Required for CNS Myelination. Cell. 2009;138:172–85.

30. Bujalka H, Koenning M, Jackson S, Perreau VM, Pope B, Hay CM, et al.

MYRF Is a Membrane-Associated Transcription Factor That

Autoproteolytically Cleaves to Directly Activate Myelin Genes. ffrench-

Constant C, editor. PLoS Biol. 2013;11:e1001625.

31. Moreno-Jiménez EP, Flor-García M, Terreros-Roncal J, Rábano A, Cafini

F, Pallas-Bazarra N, et al. Adult hippocampal neurogenesis is abundant in

neurologically healthy subjects and drops sharply in patients with Alzheimer's

disease. Nat Med. 2019;25:554–60.

32. Rushton WAH. A theory of the effects of fibre size in medullated nerve. J.

Physiol. 1951;115:101–22.

33. Urbach H, Mast H, Egger K, Mader I. Presurgical MR Imaging in Epilepsy.

Clin Neuroradiol. 2015;25:151–5.

34. Bernasconi A, Cendes F, Theodore WH, Gill RS, Koepp MJ, Hogan RE, et

al. Recommendations for the use of structural magnetic resonance imaging in

the care of patients with epilepsy: A consensus report from the International

League   Against   Epilepsy   Neuroimaging   Task   Force.   Epilepsia.

2019;60:1054–68.

35. Wellmer J, Quesada CM, Rothe L, Elger CE, Bien CG, Urbach H.

Proposal for a magnetic resonance imaging protocol for the detection of

epileptogenic lesions at early outpatient stages. Epilepsia. 2013;54:1977–87.

36. Sternberger LA, Sternberger NH. Monoclonal antibodies distinguish

phosphorylated and nonphosphorylated forms of neurofilaments in situ. Proc

Natl Acad Sci USA. 1983;80:6126–30.

37. Baumann N, Pham-Dinh D. Biology of Oligodendrocyte and Myelin in the

Mammalian Central Nervous System. Physiol. Rev. 2001;81:871–927.

38. Vogel US, Thompson RJ. Molecular Structure, Localization, and Possible

Functions of the Myelin-Associated Enzyme 2′,3′-Cyclic Nucleotide

3′-Phosphodiesterase. J. Neurochem. 1988;50:1667–77.

39. Tomassy GS, Berger DR, Chen H-H, Kasthuri N, Hayworth KJ, Vercelli A,

et al. Distinct Profiles of Myelin Distribution Along Single Axons of Pyramidal

Neurons in the Neocortex. Science. 2014;344:319–24.

40. Stassart RM, Möbius W, Nave K-A, Edgar JM. The Axon-Myelin Unit in

Development and Degenerative Disease. Front Neurosci. 2018;12:467.

41. Nave K-A, Werner HB. Myelination of the Nervous System: Mechanisms

and Functions. Annu Rev Cell Dev Biol. 2014;30:503–33.

42. Rayatpour A, Farhangi S, Verdaguer E, Olloquequi J, Ureña J, Auladell C,

et al. The Cross Talk between Underlying Mechanisms of Multiple Sclerosis

and Epilepsy May Provide New Insights for More Efficient Therapies.

Pharmaceuticals. 2021;14:1031.

43. Zucca I, Milesi G, Medici V, Tassi L, Didato G, Cardinale F, et al. Type II

focal cortical dysplasia: Ex vivo 7 T magnetic resonance imaging

abnormalities and histopathological comparisons. Ann Neurol. 2016;79:42–

58.

44. Liewald D, Miller R, Logothetis N, Wagner H-J, Schüz A. Distribution of

axon diameters in cortical white matter: an electron-microscopic study on

three human brains and a macaque. Biol Cybern. 2014;108:541–57.

45. Tomasi S, Caminiti R, Innocenti GM. Areal Differences in Diameter and

Length of Corticofugal Projections. Cereb Cortex. 2012;22:1463–72.

46. Zhou Q, Anderson DJ. The bHLH Transcription Factors OLIG2 and OLIG1

Couple Neuronal and Glial Subtype Specification. Cell. 2002;109:61–73.

47. Lu QR, Sun T, Zhu Z, Ma N, Garcia M, Stiles CD, et al. Common

Developmental Requirement for Olig Function Indicates a Motor

Neuron/Oligodendrocyte Connection. Cell. 2002;109:75–86.

48. Hornig J, Fröb F, Vogl MR, Hermans-Borgmeyer I, Tamm ER, Wegner M.

The Transcription Factors Sox10 and Myrf Define an Essential Regulatory

Network Module in Differentiating Oligodendrocytes. Barres BA, editor. PLoS

Genet. 2013;9:e1003907.

49. Almeida RG, Lyons DA. On Myelinated Axon Plasticity and Neuronal

Circuit Formation and Function. J Neurosci. 2017;37:10023–34.

50. Gibson EM, Purger D, Mount CW, Goldstein AK, Lin GL, Wood LS, et al.

Neuronal Activity Promotes Oligodendrogenesis and Adaptive Myelination in

the Mammalian Brain. Science. 2014;344:1252304.

Figure Legends

Figure 1

Presurgical MRI of FCD IIa and IIb patients.

Representative 3T MRI of two patients with frontal FCD IIa and FCD IIb, illustrating their differing macroscopic phenotype. Subtle cortical thickening and blurring is seen at the gray-white matter junction in FCD IIa (arrows in a, b; a: coronal, b: axial, both FLAIR). FCD IIb presents with a transmantle and deep sulcus sign (arrows in c, d; c: FLAIR, d: T2 STIR; both coronal).

Figure 2

Histological characteristics of control, FCD IIa and FCD IIb specimens.

Representative photomicrographs of cresyl violet-stained (a, d, g) and NeuN- (all neurons) (b, e, h) or SMI32-immunolabeled (pyramidal neurons in layers III and V) tissue sections (c, f, i). The FCD IIa case shows a radial lamination disturbance with hypercolumnization (d, e). In controls and FCD IIa, SMI32-positive pyramidal neurons are densely arranged in layers III and V (c, f) with the appearance of dysmorphic neurons across all layers in FCD IIa but concentrated in supragranular layers in FCD IIb (l). Detailed images of SMI32-positive cells reveal the characteristic shape of dysmorphic neurons in FCD IIa (j) and FCD IIb (k). The hallmark of FCD IIb is the presence of Balloon cells which are visualized by vimentin staining (l).

WM: white matter; Scale bars: (a-i) 250 μm; (j-l) 20 μm. I-VI: cortical layers.

Figure 3

Distribution of *mbp* mRNA-expressing OLs in control and FCD II specimens.

Representative photomicrographs of cresyl violet-stained (a, d, g) and *mbp*-

ISHH-processed (b, c, e, f, h, i) tissue sections of controls (a-c), FCD IIa (d-f)

and FCD IIb (g-i) cases. ROIs in (b, e, h) indicate the magnified areas shown

on the right. In control and FCD IIb specimens, a gradient of *mbp* mRNA-

expressing cells with the highest densities in the deeper cortical layers and

the white matter can be observed. This gradient is less pronounced in FCD

IIa. Counting of *mbp* mRNA-expressing OLs in all cortical layers does not

reveal significant differences between the groups (one-way ANOVA F=2.64,

p=0.106) (j). Individual patients are color-coded: greenish: control, blueish:

FCD IIa, redish: FCD IIb.

WM, white matter; scale bars: (a, b, d, e, g, h) 250 µm; (c, f, i) 100 µm, I-VI,

cortical layers

Figure 4

Myelination pattern in control, FCD IIa and IIb cases from the frontal lobe.

Representative photomicrographs of tissue sections immunolabeled for

CNPase to identify myelinated fibers in the gray matter of control (a), FCD IIa

(b) and FCD IIb (c) cases. The radially aligned CNPase-positive fibers reach

the border of layer I in all cases. (d, e, f) CNPase-immunolabeled sections

(red) counterstained with DAPI (blue) of control (d), FCD IIa (e) and FCD IIb

(f) cases were imaged by confocal laser microscopy in layers V and VI.

Individual micrographs of confocal stacks are shown. In controls, a dense

network of horizontally and radially oriented CNPase-labeled fiber strands is

present. White dots (asterisks) represent pre-OL and OL cell bodies (double

labeled by CNPase and DAPI) (d). This pattern of myelinated fibers is rather

disorganized in FCD IIa (e) but similar to control in FCD IIb (f). (g)

Quantification of CNPase-expressing pre-OL and OL cell bodies across all six

layers reveals similar cell numbers in all cases (one-way ANOVA F=0.74,

p=0.49). Colored dots identify individual patients, greenish: control, blueish:

FCD IIa, redish: FCD IIb.

Scale bars: (a-c) 250 µm; (d-f) 20 µm; Asterisks in d, e, f: CNPase-positive

pre-OL and OL cell bodies. WM, white matter. I-VI: cortical layers.

Figure 5

Ultrastructure of myelinated axons in FCD II types of the frontal lobe.

Representative electron photomicrographs of myelinated axons in layers V/VI

of control (a), FCD IIa (b) and FCD IIb (c) specimens. (d) Bar graph

demonstrating that the mean axon diameters of FCD IIa and FCD IIb have on

average larger axon diameters than control cases (one-way ANOVA F=2.59,

p=0.129). (e) The cumulative frequency distribution of axon diameters shows

significantly increased axon diameters in both FCD II types (one-way ANOVA

F=19.26, p<0.0001; Tukey's post hoc test: control vs. FCD IIa p<0.0001,

control vs. FCD IIb p<0.0001)]. (f) The mean g-ratio value is significantly

increased in FCD IIa cases when compared to controls (one-way ANOVA

F=4.511, p=0.044; Tukey's post hoc test: control vs. FCD IIa p=0.039). (g)

The cumulative frequency distribution of g-ratios shows significant higher g-

ratios for FCD IIb and FCD IIa [(one-way ANOVA F=14.69, p<0.0001; Tukey's

post hoc test: control vs. FCD IIa p<0.0001, control vs. FCD IIb p=0.019, FCD

IIa vs. FCD IIb p=0.046)]. (h) Bar graph demonstrating the mean myelin

sheath thickness which is similar in the three analyzed groups (one-way

ANOVA F=0.746, p=0.501). (i) The cumulative frequency distribution of myelin

sheath thickness does not reveal differences between controls and FCD IIa

but a significant increase of FCD IIb in comparison to FCD IIa and controls

[(one-way ANOVA F=4.487, p=0.012; Tukey's post hoc tests: control vs. FCD

IIb p=0.023, FCD IIa vs. FCD IIb p=0.024)]. (j-l) Correlation plots of axon

diameters versus myelin sheath thickness. (j) Controls, (k) FCD IIa and (l)

FCD IIb. There is a significant correlation of axon diameters and the thickness

of myelin sheaths for controls (Pearson's correlation: r=0.329, p=0.0003) and

FCD IIb (Pearson's correlation: r=0.309, p=0.0007) but not for FCD IIa

(Pearson' correlation: r=0.138, p=0.056). Colored dots represent individual

patients, greenish: control, blueish: FCD IIa, redish: FCD IIb. Scale bars (a-c):

250 nm.

Figure 6

Expression levels of transcription factor and myelin-associated transcripts in

frontal lobe FCD IIa and IIb.

The mRNAs levels were quantified by real-time RT-qPCR. The expression of

the transcription factor mRNAs *olig2* (Kruskal-Wallis p=0.793) (a) and *sip1*

(one-way ANOVA F=0.648, p=0.534) (b) are similar in all groups. The mRNA

levels of *myrf* (c) and of the myelin components *mpb* (d), *mag* (e) and *mog* (f)

are significantly reduced in FCD IIa when compared to FCD IIb [(*myrf*: one-

way ANOVA F=7.914, p=0.003; Tukey's post hoc test: control vs. FCD IIa

p=0.389, control vs. FCD IIb p=0.205, FCD IIa vs. FCD IIb p=0.002) (*mbp*:

one-way ANOVA F=5.964, p=0.009; Tukey's post hoc test: control vs. FCD IIa

p= 0.672, control vs. FCD IIb p=0.190, FCD IIa vs. FCD IIb p=0.008)*, (mag*:

one-way ANOVA F=11.46, p=0.0005; Tukey's post hoc test: control vs. FCD

IIa p=0.288, control vs. FCD IIb p=0.099, FCD IIa vs. FCD IIb p=0.0004)*,

(*mog*: one-way ANOVA F= 7.551, p=0.004; Tukey's post hoc test: control vs.

FCD IIa p=0.163, control vs. FCD IIb p=0.515, FCD IIa vs. FCD IIb p=0.003)].

Colored dots represent individual patients, greenish: control, blueish: FCD IIa,

redish: FCD IIb.

Figure 7

Transcription factor binding capacities to promoters of myelin-associated

genes.

OLIG2 (a-d) and MYRF (e-g) binding sites were precipitated using ChIP

assays and quantified using real-time qPCR. The amounts of precipitated

DNA are presented as the percentage of input DNA. The binding capacities of

OLIG2 to the *SIP1* (a), *MBP* (b), *CNP* (c) and *MOG* (d) promoters are similar

in all analyzed groups [(OLIG2 on *SIP1*: one-way ANOVA F=0.153, p=0.859),

(OLIG2 on *MBP*: one-way ANOVA F=0.247, p=0.784), (OLIG2 on *CNP*: one-

way ANOVA F=0.654, p=0.531), (OLIG2 on *MOG*: Kruskal-Wallis p=0.475)].

However, MYRF has significantly lower binding potential to the *MBP* (f),

SOX10 (g) and *MAG* (h) promoters in FCD IIa when compared to FCD IIb

[(MYRF on *SOX10*: one-way ANOVA F=10.51, p=0.0008; Tukey's post hoc

test: control vs. FCD IIa p=0.0009, control vs. FCD IIb p= 0.112, FCD IIa vs.

FCD IIb p= 0.024), (MYRF on *MBP:* one-way ANOVA F=6.55, p=0.007;

Tukey's post hoc test: control vs. FCD IIa p=0.032, control vs. FCD IIb p=

0.976, FCD IIa vs. FCD IIb p= 0.011), (MYRF on *MAG:* Kruskal-Wallis

p=0.003; Dunn's multiple comparison test: control vs. FCD IIa p=0.017,

control vs. FCD IIb p>0.999, FCD IIa vs. FCD IIb p=0.041)]. Correlation of

MYRF binding capacities with *mpb* (i) and *mag* (j) mRNA expression levels.

Pearson's correlation was performed for the percentage of input values from

the ChIP assays with the relative expression values of the RT-qPCRs.

Significant correlations are present between MYRF binding capacities and

mpb (i) and *mag* (j) mRNA levels, in particular for FCD IIb (Pearson's

correlation: r=0.546, p=0.009) and *mag* (Pearson's correlation: r=0.539,

p=0.009). Colored dots represent individual patients, greenish: control,

blueish: FCD   IIa,   redish: FCD   IIb.

Table 1. Clinical data of FCD IIa (A), FCD IIb (B) and controls (C) selected for this study.

A. FCD IIa cases

ID	Color code	Sex	Age at epilepsy onset (years)	Age at surgery (years)	Epilepsy duration (years)	Seizure type	Seizure frequency/ day	Neuro-pathology	MRI findings	Brodmann area	Methods applied
1		m	0.2	6	5,8	FA	n.a., during night	IIa	FCD	8	IHC, ISHH, PCR, ChIP
2		m	8	22	14	FA, FIA, BTCS	2-3	IIa	TM-FCD	10	IHC, ISHH, PCR, ChIP
3		f	2	8	6	FA	>1	IIa	FCD	8	IHC, EM, ISHH, PCR, ChIP
4		f	18	23	5	FA, FIA, BTCS	0.07	IIa	Neg.	9	IHC, EM, ISHH, PCR, ChIP
5		f	10	18	8	FA, FIA	8-10	IIa	Neg.	9	IHC, EM, ISHH, PCR, ChIP
5		f	15	18	3	FA, FIA, BTCS	0.04	IIa	FCD	n. a.	IHC, ISHH, PCR, ChIP
7		m	0.6	10	9.4	FA, FIA	4.5	IIa	FCD	9	IHC, EM, PCR, ChIP
8		f	1	8	7	FIA	0.36	IIa	FCD	9	IHC, PCR, PCR, ChIP
9		m	4	38	34	FA, FIA	3	IIa	Neg.	44	IHC, PCR, PCR, ChIP

B. FCD IIb cases

ID	Color code	Sex	Age at epilepsy onset (years)	Age at surgery (years)	Epilepsy duration (years)	Seizure type	Seizure frequency/ day	Neuro-pathology	MRI findings	Brodmann area	Methods applied
1		m	7.0	48	41.0	FA, BTCS	0.1, during night	IIb	TM-FCD	6	IHC, EM, ISHH, PCR, ChIP
2		m	5	36	31	FA, FIA	6	IIb	TM-FCD,	8	IHC, EM, ISHH, PCR, ChIP
3		f	9	45	36	FA, BTCS	0.29	IIb	TM-FCD	6	IHC, ISHH, PCR, ChIP
4		f	9	27	18	FA	1	IIb	TM-FCD	n.a.	IHC, ISHH, PCR, ChIP
5		f	12	40	28	FA, BTCS	1	IIb	TM-FCD	8	IHC, EM, ISHH, PCR, ChIP
6		m	14	23	9	FA, FIA, BTCS	1	IIb	TM-FCD	8	IHC, EM, ISHH, PCR, ChIP
7		m	9	38	29	FA, BTCS	1.2	IIb	TM-FCD	6	ISHH, PCR, ChIP
8		f	3	12	9	FA, BTCS	multiple	IIb	TM-FCD	6	IHC, PCR, PCR, ChIP
9		f	16	34	18	FIA, BTCS	0.004	IIb	TM-FCD	44, 45	IHC, PCR, ChIP

C. Control cases

ID	Color code	Sex	Age at epilepsy onset (years)	Age at surgery (years)	Epilepsy duration (years)	Seizure type	Seizure frequency/ day	MRI findings	Brodmann area	Methods applied
1		f	19	26	7	FIA, BTCS	0.43	GG	10	IHC, EM, ISHH, PCR
2		m	12	17	5	FIA, BTCS	2-3	Cystic lesion	10	IHC, EM, ISHH, PCR
3		m	2	10	8	FIA, BTCS	2	Cystic lesion	9	IHC, EM, ISHH, PCR
4		f	2	14	12	FA	1	Blurring of GM-WM boundary	11	IHC, EM, ISHH, PCR

ChIP, chromatin immunoprecipitation; BTCS: focal to bilateral tonic-clonic seizures; EM, electron microscopy; FA: focal aware seizures; FCD, focal cortical dysplasia; FIA: focal impaired awareness seizures; GG, ganglioglioma; GM, gray matter; IHC, immunohistochemistry; ISHH, *in situ* hybridization histochemistry; n.a., not available; neg., negative; PCR, polymerase chain reaction; TM, transmantle sign; WM, white matter.

a Nissl   **b** *mbp*-ISHH   **c**

I, II, III, IV, V, VI, WM

Ctrl

d   **e**   **f**

I, II, III, IV, V, VI, WM

FCD IIa

g   **h**   **i**

I, II, III, IV, V, VI, WM

FCD IIb

j

mbp mRNA-positiv cells [mm²]

Ctrl   IIa   IIb

Transcription Factors

a — Rel. exp. of *olig2* mRNA — Ctrl, IIa, IIb

b — Rel. exp. of *sip1* mRNA — Ctrl, IIa, IIb

c — Rel. exp. of *myrf* mRNA — Ctrl, IIa, IIb **

Structural Myelin Proteins

d — Rel. exp. of *mbp* mRNA — Ctrl, IIa, IIb **

e — Rel. exp. of *mag* mRNA — Ctrl, IIa, IIb ***

f — Rel. exp. of *mog* mRNA — Ctrl, IIa, IIb **