Synthesis of versatile neuromodulatory molecules by a gut microbial glutamate decarboxylase

Authors. Pavani Dadi[1,2], Clint W. Pauling[1,3], Abhishek Shrivastava[1,2], and Dhara D. Shah*[1,3]

Affiliations. [1]Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University, Tempe, AZ 85281

[2]School of Life Sciences, Arizona State University, Tempe, AZ 85281

[3]School of Mathematical and Natural Sciences, Arizona State University, Glendale, AZ 85306

*Corresponding author: dhara.shah1@asu.edu

ABSTRACT

Dysbiosis of the microbiome correlates with many neurological disorders, yet

very little is known about the chemistry that controls the production of neuromodulatory

molecules by gut microbes. Here, we found that an enzyme glutamate decarboxylase

(*Bf*GAD) of a gut microbe *Bacteroides fragilis* forms multiple neuromodulatory

molecules such as γ-aminobutyric acid (GABA), hypotaurine, taurine, homotaurine, and

β-alanine. We evolved *Bf*GAD and doubled its taurine productivity. Additionally, we

increased its specificity towards the substrate L-glutamate. Here, we provide a chemical

strategy via which the *Bf*GAD activity could be fine-tuned. In future, this strategy could

be used to modulate the production of neuromodulatory molecules by gut microbes.

INTRODUCTION

Low levels of inhibitory neurotransmitter like γ-aminobutyric acid (GABA) have

been associated with many different neurological disorders such as epilepsy,

schizophrenia, autism spectrum disorder (ASD), ADHD, panic disorder, PTSD, major

depressive disorder, progressive multiple sclerosis, dementia, and Alzheimer's disease[1-

12]. Besides GABA, taurine is another amino acid that is found in abundance in the

brain[13]. Taurine is a GABA$_A$ receptor agonist and exerts downstream effects similar to

GABA[14]. Concentrations of taurine are altered in individuals with Alzheimer's disease,

where reduced levels of taurine are found in the brain[15-21] whereas, higher urinary

excretion of taurine is seen in elderly patients with dementia[22]. In a separate study,

gnotobiotic mouse transplanted with fecal microbiota from an Alzheimer's patient

showed lower abundance of GABA and taurine in the feces[23]. Based on these

observations, supplementation with taurine and its analogs has been investigated in

Alzheimer's disease models[24,25]. In one such study, Homotaurine, a molecule similar to

taurine is shown to restore cognitive functions in patients with Alzheimer's disease[26]. In

addition, it was also observed that taurine reverses cognitive deficits in APP/PS1 mouse

model[25] and improves learning and memory in mice[24,27].

Studies that correlate gut microbial dysbiosis to two of the prominent

neurodegenerative disorders – Dementia and Alzheimer's disease, often show

modulations in the abundance of microbes of the genus *Bacteroides*[28-33] . *Bacteroides*

are prominent members of the human gut[34,35] and have the ability to produce GABA[36,37].

All *Bacteroides* encode the gene for the enzyme glutamate decarboxylase (GAD)[38].

Glutamate decarboxylases (GADs) are PLP (pyridoxal phosphate) dependent enzymes

which catalyze conversion of an excitatory neurotransmitter glutamate to an inhibitory

neurotransmitter GABA. However, very little is known about the chemistry of the

glutamate decarboxylases encoded by *Bacteroides sp.*

Despite widespread prevalence of substrate promiscuity in glutamate

decarboxylases (GADs) across different domains of life, substrate specificity and

product formation landscape for most prokaryotic GADs, specifically GADs of

Bacteroides sp. are still not clearly understood. For this reason, we explored the role of

annotated GAD from *Bacteroides fragilis* in the production of neuromodulatory

molecules like GABA, taurine and its analogs, and β-alanine. Our biochemical

characterization of *Bf*GAD shows kinetic and functional divergence of this enzyme from

other studied prokaryotic GADs. Our study shows that *Bf*GAD is not only specific to L-

glutamate, but it can decarboxylate other substrates to produce multiple

neuromodulatory molecules. Even within the substrate mix containing L-glutamate, we

can detect products other than GABA. This points towards the ability of this enzyme to

function in a complex system with multiple substrates which can be a beneficial trait for

microbes present in the human gut. Through rational protein engineering, we have

evolved *Bf*GAD which is capable of producing two-fold more taurine compared to the

native *Bf*GAD. Additionally, we evolved *Bf*GAD to be more specific towards L-glutamate.

Based on our results of initial engineering with *Bf*GAD, the enzyme seems

resilient in nature and is able to tolerate changes to the active site very well. We think

that *Bf*GAD can be evolved with rational designing and engineering, either to produce

various neuromodulatory molecules in specific ratios, or to synthesize a particular

neuromodulatory molecule exclusively (Fig. 1a). This approach has the ability to

generate variants of *Bf*GAD that might provide a road to therapeutic interventions for

multiple neurodegenerative disorders.

MATERIALS AND METHODS

Materials. Kanamycin, IPTG, GABase from *Pseudomonas fluorescens*, β-ME (beta-

mercaptoethanol), α-ketoglutarate, D-glutamatic acid, L-cysteine sulfinic acid

monohydrate, L-cysteic acid monohydrate, GABA (γ-aminobutyric acid), hypotaurine,

taurine, L-aspartic acid sodium salt, β-alanine, HEPES, Imidazole, pyridoxal 5'-

phosphate monohydrate and TLC Silica gel 60 F_{254} (20 cm x 20 cm) were purchased

from Sigma-Aldrich. Sodium L- glutamate monohydrate, Luria-Bertani Broth (LB), buffer

components, Sodium chloride, Sodium acetate, ninhydrin and $NADP^+$ disodium salt

were purchased from Fisher Scientific. All restriction enzymes and competent cells of *E.*

coli BL21(DE3) and *E. coli* NEB5☐ were purchased from New England Biolabs (NEB).

Headspace vials were purchased from Chemglass Inc.

Experimental procedures

Gene cloning. A synthesized gene for *B. fragilis* glutamate decarboxylase (*Bf*GAD) was created with the help of Genewiz from Azenta life sciences. The gene was codon optimized for expression in *E. coli* cells, which was then subcloned into a pET28a expression vector between the NdeI and HindIII restriction sites to incorporate an N-terminal hexahistidine (His$_6$) Tag. The pET28a-*Bf*GAD construct was confirmed by agarose gel electrophoresis, restriction digestion, sanger sequencing (Genewiz), and full plasmid sequencing (Plasmidsaurus) and then transformed into *E. coli* NEB5☒ and *E. coli* BL21(DE3) competent cells by heat shock. Glycerol stocks of the cells harboring the construct pET28a-*Bf*GAD were stored at – 80 ºC.

Expression and purification of *Bf*GAD$_{WT}$ and engineered *Bf*GADs. *E. coli* BL21(DE3) cells containing the pET28a-*Bf*GADs were cultivated in LB medium containing kanamycin (50 μg/mL) with shaking (200 rpm) at 37 ºC until OD$_{600}$ reached 0.7. At this point, the *Bf*GAD expression was induced with isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 1 mM and growth was continued at 37 ºC for 3 hours to allow for protein expression with continuous shaking at 200 rpm. Cells were harvested via centrifugation at 4000g for 30 min at 4 ºC and cell pellets were stored at – 80 ºC. The frozen cell pellets were thawed on ice and resuspended in buffer A containing 20 mM Tris-HCl, 500 mM NaCl, 40 mM imidazole, 0.1 mM PLP, pH 7.8. All subsequent purification steps were carried out at 4 ºC. The cells were disrupted by sonication (20000 Hz for 10 cycles of 30 seconds each altering 1 minute on ice) and centrifuged at 13500g at 4 ºC for 30 min to separate supernatant from cell debris. The supernatant was then filtered through 0.2 μm PES filter membrane

and loaded onto a HisTrap HP column (5 ml, 1.6 x 2.5 cm, Ni Sepharose High-performance column, GE Healthcare, now Cytiva) pre-equilibrated with buffer A at 1 mL/min. The wash step was carried out by running 5-column volumes of lysis buffer A to elute contaminating proteins. *Bf*GAD was eluted with a linear gradient from buffer A (containing 40 mM imidazole) to an elution buffer B containing 20 mM Tris-HCl, 500 mM NaCl, 400 mM imidazole, 0.1 mM PLP, pH 7.8 at 2 mL/min. Eluted protein peaks were pooled, buffer exchanged and concentrated to a final buffer (50 mM HEPES, pH 7.2) using Vivaspin 50 kDa MWCO filters (Cytiva). Aliquots of purified proteins were stored at – 80 ºC. Engineered *Bf*GADs were purified similarly. SDS-PAGE analysis and activity assays were performed to confirm the purity and presence of functional *Bf*GAD proteins.

Gel filtration chromatography with *Bf*GAD$_{WT}$. To understand oligomeric state of *Bf*GAD$_{WT}$, gel filtration chromatography was performed at two different pH, 4.7 (50 mM sodium acetate, 150 mM NaCl) and pH 7.2 (50 mM HEPES, 150 mM NaCl) using HiPrep™ 16/60 Sephacryl® S-200 HR column. Equilibration and elution steps were carried out at a flow rate of 0.5 mL/min. High molecular weight (HMW) and Low molecular weight (LMW) calibration kits (Cytiva) were used for column calibration, to create a standard curve, and for molecular weight determination (Supplementary Fig. 5b, 5c).

Activity assays. *Bf*GAD activity assays were performed by a coupled enzyme assay with a GABase system to measure GABA production spectrophotometrically. Briefly, *Bf*GAD was incubated with 50 mM L-glutamate in 50 mM sodium acetate, pH 4.7 buffer or in buffers at other pH values for one hour at 25 ºC. Enzymatic reactions were stopped by boiling samples for 15 min. These samples were then centrifuged at 6000*g* for 5 min

and if necessary, dilutions were created in 50 mM Tris-HCl, pH 8.6. For GABA

measurement, 75 µL of samples (either diluted or undiluted) were added to 25 µL of

Gabase assay mix containing 10 mM BME, 2 mM α-ketoglutarate, 600 µM NADP$^+$, and

30 µg (0.015U/mL) of GABase in 50 mM Tris-HCl, pH 8.6[39-43]. In the GABase assay,

GABA is converted to succinic semialdehyde (SSA) and then to succinate with

subsequent production of NADPH which was measured at 340 nm. Using an extinction

coefficient of 6,220 M^{-1} cm^{-1} at 340 nm, NADPH concentrations were calculated which

will provide GABA concentrations in measured samples ([NADPH]=[GABA]).

BfGAD absorbance spectrophotometry and activity assays with pH variation.

Spectra of BfGAD$_{WT}$ were recorded from 200 nm - 800 nm in buffers at pH 4.7, 5.5, 6.7,

7.5, and 8.6 at 25 ºC on Agilent Cary 3500 UV-Vis spectrophotometer. From these

spectra, absorbance changes at 335 nm (enolimine) and 420 nm (ketoenamine) for

enzyme bound PLP cofactor were collected. A plot of pH versus absorbance at 420 nm

was generated and the curve fitting was carried out using equation (1)[44].

$$\frac{AbsHnE - AbsE}{Abs - AbsE} - 1 = \frac{[10^{-npK}]}{[10^{-npH}]} \hspace{3cm} (1)$$

Where pK is the acid dissociation constant, number of protons involved in the titration is

n and, AbsHnE and AbsE are the absorbances at acidic and basic pH which produce

protonated and deprotonated forms of the enzymes. The initial activity assays for the

BfGAD$_{WT}$ were performed in buffers at pH 4.7 - 8 with GABase assay system as

mentioned in the above method. Based on the results of these activity assays, all

subsequent enzymatic assays were carried out at pH 4.7.

Steady-state kinetics. Kinetic parameters of BfGAD$_{WT}$ were determined by varying

BfGAD concentrations (0.5 µM-2 µM) and by varying L-glutamate concentrations (0.125

mM-32 mM) in 50 mM sodium acetate, pH 4.7 at 25⁰ C. For enzyme concentration

variation experiments, reaction samples with each enzyme concentration were collected

at 1 min time point after starting the reaction and then stopped by boiling for 15 min.

Once cooled to room temperature, these samples were analyzed for GABA content with

GABase assay as mentioned above. For the substrate concentration variation

experiments, 100 µL aliquots of reactions with each substrate concentration were

collected at different time intervals (0.2 – 30 min). The reactions were stopped by boiling

the samples for 15 minutes. Once cooled to room temperature, these samples were

analyzed for GABA content with GABase assay as mentioned above. The initial

velocities of various reactions were calculated by fitting data of early timepoints to a

linear regression. These initial velocities were then plotted against substrate

concentrations and curve fitting was carried out using equation (2).

$$v = \frac{V_{max}[S]^h}{\left(k_{half}\right)^h + [S]^h}$$ (2)

Where v is the initial velocity, V_{max} is the maximum velocity, K_{half} is the concentration of

substrate at which the reaction velocity is half of V_{max}, [S] is the substrate concentration,

h is the hill coefficient.

Generation of engineered *Bf*GAD variants. The pET28a-*Bf*GAD was used as a

template for the generation of engineered *Bf*GAD variants. *Bf*GAD$_{D104N}$ and *Bf*GAD$_{F81W}$

were created by site-directed mutagenesis with Phusion DNA polymerase using the

following primer pairs incorporating codons for the specific amino acid substitution

(underlined).

For *Bf*GAD$_{D104N}$

F: 5' CCGCAATGCGCGGATATTCGGTTTC<u>ATT</u>AATATAGTTAATG 3'

R: 5' CATTAACTATATT<u>AAT</u>GAAACCGAATATCCGCGCATTGCGG 3'

For *Bf*GAD_{F81W}

F: 5' CATCCATATAGGTGGTCACCCAGGTCG<u>CC</u>AGGTTCAGGCGCG 3'

R: 5' CGCGCCTGAACCTGGCGACC<u>TGG</u>GTGACCACCTATATGGATG 3'

The variants were confirmed by gene sequencing (Genewiz) and full plasmid

sequencing with Plasmidsaurus.

$CO_{2(g)}$ detection by headspace gas chromatography. All enzymatic reactions and

controls were prepared in 6 mL vials and sealed prior to starting the reaction. The

reactions were started by the addition of *Bf*GAD (WT or engineered variants) with a

syringe into the reaction mixture containing 50 mM substrate in 50 mM sodium acetate,

pH 4.7. These vials were incubated at 37 ⁰C for 24 hours and 48 hours. Headspace GC

measurements were carried out on an Agilent 8890 gas chromatograph system

equipped with a flame ionization detector (FID) and a Hayesep Q packed column (1.8 m

x 2 mm x 3.17 mm) (Agilent) operating with an Argon carrier gas (flow rate= 5 mL/min).

The oven was programed to hold 30 °C for 6.5 min, ramp at 30 °C/min to 280 °C with a

hold for 4 min for a total run time of 18.83 min. The flame ionization detector was used

for the detection of CO_2 gas with a temperature setting of 275 °C, hydrogen flow of 60

mL/min, air flow of 400 mL/min and constant makeup gas (nitrogen) at 5 mL/min. The

retention time for CO_2 gas was 12.83 min. Multiple standards of CO_2 gas were analyzed

by this method to confirm retention time before injecting reaction samples

(Supplementary Fig. 15).

Thin layer chromatography (TLC). Silica gel plates (stationary phase) with a solvent

system (mobile phase) of 3:1:1 ratio of butanol: acetic acid: H_2O were used to separate

reaction products from substrates[45]. 2 µL of reaction mixtures were spotted on glass

silica plates along with 10 mM of various metabolites (L-glutamate, D-glutamate, L-CSA,

L-CA, L-HCA, L-Aspartate, GABA, hypotaurine, taurine, homotaurine and β-alanine) as

standards. Separation via TLC was carried out at 25 °C for 3-4 hours in an above-

mentioned mobile phase. Once mobile phase reached a sufficient height on the TLC

plate, chromatographic separation was discontinued. Plates were treated with 0.5%

ninhydrin in acetone (w/v) and heated minimally with a dryer for the color development.

The TLC plate images were initially inverted via ImageJ and then intensity of each spot

was quantified by a custom python code.

Liquid chromatography with tandem mass spectrometry (LC-MS/MS) detection of

decarboxylated products from reactions catalyzed by *Bf*GADs (WT and

engineered variants). Samples were analyzed by ESI-LC-MS/MS in positive (products)

and negative ion (substrates) mode using a Thermo vanquish LC and TSQ Altis Triple

Quadrupole Mass Spectrometer. Samples were separated by gradient elution using

Agilent Infinity lab Poroshell 120 HILIC-Z 2.1 x 100 mm, 2.7 µm column at 25 °C with 20

mM ammonium formate, pH 3.0 (Solvent A) and 20 mM ammonium formate in 9:1

acetonitrile: H_2O, pH 3.0 (Solvent B). Elution was initiated at 70% B for 11.5 min,

followed by gradient elution from 70 to 100% B over 11.5 min, and 100% B for 4 min at

a flow rate of 0.5mL/min. The precursor to product transitions of *m/z* $[M+H]^-$152$\rightarrow$

88.1(Cysteine sulfinic acid), $[M+H]^-$ 168$\rightarrow$ 81.1(Cysteic acid), $[M+H]^+$110.1$\rightarrow$

45.1(hypotaurine), $[M+H]^+$126.1$\rightarrow$ 44.1(taurine) were employed to monitor substrates

and products.

Multiple Sequence alignment and phylogenetic analyses of gut microbial GADs.

All the protein sequences were obtained from NCBI. Multiple sequence alignment was performed using Clustal Omega (1.2.4) tool with default parameters[46,47] to understand sequence similarities among various gut microbial glutamate decarboxylases (GADs). The phylogenetic tree was created for *Bacteroides* GADs using cyberinfrastructure for phylogenetic research (CIPRES)[48]. Within CIPRES, the MAFFT on XSEDE (7.505)[49] was utilized to create a separate multiple sequence alignment that goes through FastTreeMP on XSEDE(2.1.10)[50,51] to obtain the phylogenetic tree file which can be visualized and annotated with Interactive tree of life(iTOL v6.8.1)[52].

***Bf*GAD dimer generation with the AlphaFold2 and visualization, structural alignments, and mutagenesis via PyMOL.** The dimer structure of wildtype *Bf*GAD was generated using AlphaFold2 and visualized using PyMOL. Variants were then created using the mutagenesis wizard in PyMOL. Rotamers for the altered residues were selected based on their best fit to the position of wild-type residues in the *Bf*GAD protein. All structural alignments were carried out using PyMOL align method, involving five iteration cycles and a cutoff of 2 Å.

RESULTS

Glutamate Decarboxylases (GADs) are prevalent in the *Bacteroides* genus. A bioinformatics analysis showed that many species of the *Bacteroides* genus harbor genes annotated as GAD (Supplementary Fig. 1). We focused on a GAD (*Bf*GAD) from the model gut microbe, *Bacteroides fragilis*. Multiple sequence alignment performed with gut microbial GADs including *Bf*GAD revealed that multiple catalytic residues, either predicted to be involved in substrate binding (highlighted in cyan) or cofactor

(PLP) binding (highlighted in yellow) are highly conserved (Supplementary Fig. 2)[44,53].

Moreover, in a phylogenetic analysis of *Bf*GAD with other gut microbial GADs, we see

that the annotated glutamate decarboxylases from *Alistipes putredinis* and

Parabacteroides merdae are closely related to *Bf*GAD (Supplementary Fig. 3).

Wild-type *Bf*GAD (*Bf*GAD$_{WT}$) forms an oligomer. The recombinant wild-type *Bf*GAD

(*Bf*GAD$_{WT}$) with an N-terminal His-tag was purified by Ni-NTA affinity chromatography. A

protein band with an approximate expected mass of a *Bf*GAD$_{WT}$ monomer (calculated

mass 56 KDa) was observed with SDS PAGE, while a band indicating a tetrameric

oligomer was observed with a native PAGE (Supplementary Fig. 4). Oligomeric

composition of the *Bf*GAD$_{WT}$, as determined by gel filtration chromatography, is

presented in Supplementary Fig. 5a as chromatograms for the *Bf*GAD$_{WT}$ at an acidic pH

(blue) and a neutral pH (red). Peak 1 was in the void volume of the column indicating

protein aggregates. There were two peaks representing two different oligomeric

compositions (2 and 3) observed at each pH. Peak 2 was broad and corresponded to

approximately 351 KDa, indicating a hexamer. Peak 3 was the major peak and

exhibited an approximate molecular weight between a dimer and a tetramer at pH 4.7,

whereas at pH 7.2 the peak 3 appeared to be closer to a dimer than a tetramer

(Supplementary Fig. 5, a-c, 180 KDa at pH 4.7 and 150 KDa at pH 7.2) for *Bf*GAD$_{WT}$.

Our data indicated that *Bf*GAD$_{WT}$ is an oligomeric enzyme which can either be a dimer

or a tetramer. Furthermore, the addition of external PLP did not change the oligomeric

state of the *Bf*GAD$_{WT}$ (Supplementary Fig. 5d).

PLP cofactor bound to the *Bf*GAD$_{WT}$ undergoes pH dependent tautomeric

changes. Spectral studies with *Bf*GAD$_{WT}$ revealed that the enzyme was purified in its

holo form with the covalently bound PLP cofactor. There were two absorption maxima

observed for the BfGAD$_{WT}$ – 335 nm and 420 nm (Fig. 1b). These absorption maxima

result from two tautomeric forms of the enzyme bound PLP cofactor, enolimine form at

335 nm and ketoenamine form at 420 nm[54-56]. The proportions of these tautomers vary

depending on the pH (Fig. 1b). At lower pH, the species absorbing at 420 nm was more

dominant, whereas at higher pH, 340 nm absorbing form became more prevalent. With

increase in pH from 4.7 to 8.6, there was a decrease in 420 nm absorbing species with

concomitant increase in 340 nm absorbing species (Fig. 1b). These changes in the

absorbances of the bound PLP cofactor as the function of the change in pH with

BfGAD$_{WT}$, have been seen with other microbial GADs[57-59]. The analysis of the pH

versus absorbance curve for the 420 nm wavelength indicated that the transition in

absorbance in response to pH variation is influenced by the involvement of multiple

protons (Supplementary Fig. 6a). Previously with $E.\ coli$ GAD, Tramonti et al. and

Capitani et al. showed that the less hydrated active site at neutral pH drove the

formation of enolimine tautomer (340 nm) whereas more polar active site at acidic pH

predominantly produced ketoenamine tautomer (420 nm)[43,44]. Additionally, the

alterations in cofactor absorbance corresponding to different pH were in close

agreement with the BfGAD$_{WT}$'s activity profiles measured across the same range of pH

values. Particularly, at pH 4.7, where BfGAD$_{WT}$ with a bound PLP predominantly

absorbed at 420 nm, the enzyme also exhibited its maximum catalytic activity

(Supplementary Fig. 6b). Additionally, enzymatic activity was found to be minimal at a

pH other than 4.7 (Supplementary Fig. 6b). Therefore, all subsequent activity assays

were carried out at pH 4.7.

Evidence for allosteric regulation of *Bf*GAD_{WT}. The activity assays conducted with varying $BfGAD_{WT}$ concentrations exhibited a linear relationship between the initial reaction velocities and enzyme concentrations (Fig. 1c). From the plot of enzyme concentration variation, we found an optimal $BfGAD_{WT}$ concentration for subsequent kinetic assays. Interestingly, while varying L-glutamate concentrations, the plot of substrate concentration versus initial velocity displayed a sigmoidal curve instead of a typical hyperbolic dependence, suggestive of an allostery in $BfGAD_{WT}$. This is an example of a homotropic allosteric regulation. From the fit of this plot with the hill equation (eq. 2), the analyzed $K_{0.5}$ (K_{half}) was 8.6 ± 0.4 mM and V_{max} was 0.86 ± 0.04 mM/min with a hill coefficient of 1.7 (Fig. 1d). $(K_{half})^h$ is also known as K_{prime} (K'), which is equivalent to K_m from a hyperbolic kinetics of non-allosteric enzymes, when h=1. However, in cooperative systems where h $\neq$ 1 like presented here for $BfGAD_{WT}$ then, K' (K_{half}^{h}) no longer represents the substrate concentration required to achieve half of maximal velocity.

Engineered *Bf*GAD variants exhibit structural and catalytic perturbations. The earliest microbial GAD structures were solved for *E. coli* GADs (GAD*A*, PDB ID 1XEY and GAD*B*, PDB ID 1PMM) (Supplementary Fig. 7)[44,53]. Both structures were solved with the bound ligand in the active site. Based on the interactions of the acetate ion in the GAD*B* active site and glutarate (substrate analog) in the GAD*A* active site (Supplementary Fig. 7b), we selected residues for the evolution of the $BfGAD_{WT}$ via rational design[44,53,60]. Two such residues that are present in the active sites of both *Ec*GADs are phenylalanine 63 (corresponds to F81 in $BfGAD_{WT}$) and aspartate 86 (corresponds to D104 in $BfGAD_{WT}$) that make H-bonding interactions with one of the

carboxylates of the substrate analog and an acetate ion[44,53]. Specifically, in *E. coli*

GAD*B* ligand bound structure, the carboxylate group of acetate received a hydrogen

bond from the amide nitrogen of the F63 (F81 BfGAD$_{WT}$) and a side chain carboxylate

of D86 (D104 BfGAD$_{WT}$) of the neighboring subunit[44]. Whereas in *E. coli* GAD*A* ligand

bound structure, one of the carboxylate group of the glutarate (substrate analogue)

forms H-bond with the amide nitrogen of F63 (F81 BfGAD$_{WT}$) and with the side chain

carboxylate of D86 (D104 BfGAD$_{WT}$) of the neighboring subunit[53]. These residues are

completely conserved in annotated GADs from prominent gut microbes with the

exception of *Eggerthella lenta* that harbors shorter GAD with the absence of a

conserved phenylalanine (Supplementary Fig. 2). Most microbial GADs are functionally

active as dimers, with residues from both monomers contributing to the active

site[44,53,57,61] . Recently, structure of glutamate decarboxylase from *Bacteroides*

thetaiotamicron (*Bt*GAD), bound with a substrate analog (glutarate) was solved[57]. With

these available structures, we aligned the substrate analog (glutarate) bound structures

of *Ec*GAD*A* (PDB ID 1XEY) and *Bt*GAD (PDB ID 7X51) with BfGAD$_{WT}$ dimer created

with AlphaFold2 (Fig. 2). Based on these structural alignments we engineered two

enzymes by introducing single amino acid alterations at two separate positions to test if

the substrate preference of the BfGAD$_{WT}$ can be evolved. These changes were

Asp104Asn (D104N) and Phe81Trp (F81W). Figure 2a and 2b illustrate the locations of

these residues within the active site of BfGAD$_{WT}$ (green). These residues are depicted

in close proximity to glutarate and the cofactor PLP bound in the active site of *Ec*GAD*A*,

which has been superimposed on the BfGAD$_{WT}$ dimer (Fig. 2a, 2b). Similarly, Fig. 2c

and 2d display aligned BfGAD$_{WT}$ and *Bt*GAD structures with glutarate and PLP bound in

the active site of *Bt*GAD. In addition to *Bf*GAD$_{WT}$, the structures of variants, *Bf*GAD$_{D104N}$

(Fig. 2a, 2c) and *Bf*GAD$_{F81W}$ (Fig. 2b, 2d) were then superimposed with ligand bound

structures of *Ec*GAD and *Bt*GAD.

The overall active site architecture of *Bf*GAD is mostly consistent with that of

*Ec*GAD and *Bt*GAD; however, notable positional shifts were identified among the

residues of interest. Specially, aspartic acid (D104) showed a considerable difference in

*Ec*GAD versus *Bf*GAD active site. Additionally, we detected changes in the H-bonding

interactions between the substrate analog and residues of interest when substitutions

were made to create engineered *Bf*GAD variants. We were able to purify both variants,

*Bf*GAD$_{D104N}$ and *Bf*GAD$_{F81W}$ with the same conditions as the *Bf*GAD$_{WT}$. Both of these

variants were active and purified as holoenzymes (PLP bound form). Additionally, we

collected UV-Vis spectra of *Bf*GAD$_{WT}$ and both variants at pH 4.7 (Fig. 2e). While each

of the three enzyme preparations exhibits absorbance maxima at both 335 nm and 420

nm, the relative amounts of these two tautomeric forms differ across the enzymes.

Furthermore, we measured initial velocities in the presence of the native substrate L-

glutamate via progress curve analysis for engineered *Bf*GADs along with *Bf*GAD$_{WT}$.

With that, we detected variations in the decarboxylation reaction rates across these

enzymes (Fig. 2f). *Bf*GAD$_{D104N}$ showed 1.7-fold increase whereas *Bf*GAD$_{F81W}$ showed

2.5-fold decrease in the initial velocities compared to the *Bf*GAD$_{WT}$ for decarboxylation

reactions with L-glutamate. These observed differences in catalysis between the wild-

type and engineered variants of the *Bf*GAD due to the structural perturbations resulting

from amino acid substitutions at the active site, reinforce the role of F81 and D104 in

establishing interactions with the substrate.

Wild-type and engineered *Bf*GADs can decarboxylate substrates other than L-glutamate. As mentioned above, the purified *Bf*GAD$_{WT}$ is able to catalyze the conversion of L-glutamate to GABA. To test if it can utilize the D-form of the glutamate as a substrate, we conducted activity assays with D-glutamate using the GABase assay system and TLC. Our results show that both the wild-type and engineered *Bf*GADs were unable to decarboxylate D-glutamate (Supplementary Fig. 8). Most microbial GADs studied so far show high substrate specificity towards L-glutamate[61-64]. However, there are examples of archaeal GADs that prefer L-aspartate over L-glutamate[65,66]. An archaeal GAD from *P. horikoshii* also shows decarboxylation activity with L-cysteate[65]. In addition to archaeal GADs, certain mammalian glutamate decarboxylase (GAD) homologs are known to decarboxylate one or more of the non-native substrates such as L-aspartate, L-cysteate, and L-cysteine sulfinate[67,68]. Moreover, proteins similar to GADs have been demonstrated to form taurine in some marine microbes[69]. However, there are limited systematic studies investigating the production of taurine, its analogs, and β-alanine using L-CA (cysteate), L-CSA (cysteine sulfinate), L-HCA (homocysteate), and L-aspartate as substrates (Scheme 1), specifically for gut microbial GADs. Based on two key observations, we hypothesized that gut microbial GADs might utilize these molecules as substrates: 1. *E. coli* GAD can accept the phosphonated form of L-glutamate as a substrate[70]. and 2. Archaeal and eukaryotic GADs exhibit diverse substrate specificities, enabling them to catalyze the decarboxylation reactions of non-native substrates[45,65,66,69,71].

If *Bf*GAD$_{WT}$ or engineered *Bf*GADs are able to decarboxylate substrates other than the native substrate L-glutamate, then the detection of the common product CO_2

will be a positive test for the utilization of other substrates (Scheme 1). For this reason,

we decided to use a headspace GC to measure the CO_2 evolved from reactions

catalyzed by either BfGAD$_{WT}$ or engineered BfGADs during the decarboxylation of

various substrates. Fig. 3 shows data collected for BfGAD$_{WT}$ and engineered BfGADs

with five different substrates via headspace GC. Panels 3a, 3c, and 3e demonstrate

chromatograms with the CO_2 peaks, visible immediately after 12.8 min, produced by the

reactions of BfGAD$_{WT}$, BfGAD$_{D104N}$, and BfGAD$_{F81W}$ respectively with five different

substrates. Whereas 3b, 3d, and 3f depict CO_2 peak areas from the decarboxylation

reactions of the substrates after 24 h incubation with either BfGAD$_{WT}$ or engineered

BfGADs. Here, BfGAD$_{WT}$ showed significant production of CO_2 generated from the

decarboxylation of both L-glutamate (blue) and L-CSA (red) (Fig. 3a, 3b). This is the first

indication to our knowledge of a gut microbial GAD, specifically a GAD from

$Bacteroides,$ that can utilize L-CSA as a substrate. In addition to L-CSA, BfGAD$_{WT}$ was

capable of decarboxylating L-HCA (orange) and L-CA (green) to small extents as

evidenced by the small CO_2 peaks (Fig. 3a) in the chromatogram and small peak areas

for evolved CO_2 (Fig. 3b). Additionally, BfGAD$_{WT}$ was able to decarboxylate L-aspartate

(purple) (Fig. 3a, 3b).   For BfGAD$_{WT}$, if CO_2 production peak area with the native

substrate L-glutamate is considered to be 100% then we observed 54%, 3%, 6% and

20% of CO_2 production with L-CSA, L-CA, L-HCA, and L-Asp respectively within 24 h

which increased to 75% (L-CSA), 6% (L-CA), 7% (L-HCA), and 28% (L-Asp) within 48 h

compared to the native substrate L-glutamate (Supplementary Fig. 9a).

For BfGAD$_{D104N}$, CO_2 production is comparable to the BfGAD$_{WT}$ while using native

substrate L-glutamate. Compared to the peak area of CO_2 production for L-glutamate

(native substrate), we observed around 16%, 11%, 3% and 7% CO_2 production from L-

CSA, L-CA, L-HCA, and L-aspartate respectively within 24 h (Fig. 3c, 3d) that

increased to 18% for both L-CSA and L-CA, 7% for L-HCA and slightly decreased with

L-aspartate within 48 h (Supplementary Fig. 9b). Based on the data collected with

$BfGAD_{WT}$ and $BfGAD_{D104N}$, residue Asp104 (D104) from the neighboring monomer is

playing a crucial role in accommodating various substrates. This engineered enzyme

was also able to decarboxylate L-CSA, L-HCA, and L-Asp in addition to the native

substrate L-glutamate but less efficiently than the $BfGAD_{WT}$, specifically the activity

towards L-CSA was significantly impacted. It is likely that the negative charge of the

side chain carboxylate from D104 is important for the specificity towards alternative

substrates L-CSA, L-HCA and L-Asp but not for the native substrate L-glutamate. As a

result, when the charge was removed due to the substitution of Asp (D) to Asn (N) at

position 104, the catalysis with these alternate substrates was affected, but not with the

native substrate L-glutamate. Interestingly, we noticed a 2-2.5-fold increase in the

production of taurine with this evolved enzyme compared to the $BfGAD_{WT}$ (Fig. 3d),

indicating that the substitution from D to N was favorable for taurine production.

The second engineered enzyme, $BfGAD_{F81W}$ retained its activity with the native

substrate L-glutamate as well but as observed in the initial velocity experiments (Fig.

2f), it was slower compared to $BfGAD_{WT}$. If the CO_2 peak area of $BfGAD_{WT}$ or

$BfGAD_{D104N}$ with the native substrate L-glutamate is considered to be 100%, $BfGAD_{F81W}$

showed the peak area of around 93-94% with the same concentration of the substrate.

Additionally, there was minimal decarboxylation activity of $BfGAD_{F81W}$ with L-CA, L-

HCA, and L-ASP. However, this evolved enzyme still retained considerable

decarboxylation activity with L-CSA. It is interesting to note that a bulky substitution of

tryptophan (W) at position 81, instead of phenylalanine (F) (Fig. 2b, 2d), made the

enzyme more specific towards the native substrate L-glutamate. Based on the CO_2

evolution captured by our GC data, the substrate preference for BfGAD$_{WT}$ was L-Glu >

L-CSA > L-Asp > L-HCA > L-CA that changed to L-Glu > L-CSA ≥ L-CA > L-Asp > L-

HCA for BfGAD$_{D104N}$. Whereas for BfGAD$_{F81W}$, only L-glutamate and L-CSA showed

decarboxylation and L-glutamate was a much better substrate than L-CSA.

Wild-type and evolved BfGADs produce multiple neuromodulatory molecules. The

generation of CO_2 in reactions facilitated by BfGAD$_{WT}$ and its engineered variants

prompted us to detect and identify products resulting from decarboxylation with various

substrates. Both BfGAD$_{WT}$ and BfGAD$_{D104N}$ produced decarboxylated products –

hypotaurine, taurine, homotaurine, and β-alanine from substrates L-CSA, L-CA, L-HCA,

and L-Asp respectively (Fig. 4, a-d, Supplementary Fig. 10, a-d).   Intensity plots

generated from TLC plates showed that BfGAD$_{WT}$ decarboxylated L-CSA and L-Asp to

produce hypotaurine and β-alanine more efficiently than BfGAD$_{D104N}$ (Fig. 4a, 4c).

However, BfGAD$_{D104N}$ was better at decarboxylating L-CA to taurine than BfGAD$_{WT}$ (Fig.

4b). We do not observe much of decarboxylated product taurine with BfGAD$_{WT}$ from L-

CA decarboxylation during a 24 h incubation, whereas BfGAD$_{D104N}$ shows a significant

production of taurine during this time frame (Fig. 4b, Supplementary Fig. 10e). While our

CO_2 evolution experiments have shown that BfGAD$_{WT}$ can catalyze L-CA

decarboxylation in 24 h, the failure to observe decarboxylated product on TLC may be

attributed to the low concentration of taurine produced, making it undetectable on a TLC

plate. Both enzymes BfGAD$_{WT}$ and BfGAD$_{D104N}$ were able to catalyze the

decarboxylation of L-HCA to homotaurine to a very small extent (Fig. 4d). We also

verified that the product from the L-Asp decarboxylation was β-alanine and not L-

alanine. Although, the R_f (retention factor) values were almost similar for β-alanine and

L-alanine, the staining with the ninhydrin differs for these molecules. β-alanine exhibited

a purple color whereas L-alanine shows a brick red color with ninhydrin stain

(Supplementary Fig. 10c). We did not find any activity of $BfGAD_{F81W}$ with L-CA, L-Asp,

and L-HCA and were unable to detect taurine, β-alanine, and homotaurine even after 48

h incubation (Supplementary Fig. 11a, 11b). These results reinforce our observations

from CO_2 evolution experiments. Additionally, despite the detection of CO_2 during the

decarboxylation activity of $BfGAD_{F81W}$ with L-CSA (Fig. 3f), we could not observe

detectable hypotaurine spots on the TLC plate (Supplementary Fig. 11a). We

hypothesize that the amount of hypotaurine produced by $BfGAD_{F81W}$ may not be

sufficient to be detected via TLC, as a higher concentration of molecules (in mM range)

are needed for TLC analysis.

$BfGAD_{WT}$ and $BfGAD_{D104N}$ generate neuromodulatory molecules at differential

abundances, even when presented with a mixed substrate pool. In a complex gut

environment, the organism will encounter multiple substrates simultaneously. To

investigate how $BfGAD_{WT}$ functions in such a complex environment, we conducted

competition assays between various substrates. To this end, we tried to detect the

presence of decarboxylated products in mixtures containing the native substrate L-

glutamate with a specific alternate substrate. Our data with $BfGAD_{WT}$ indicate that when

the concentration ratio of the native substrate to the alternative substrate was 1:1,

GABA was predominantly seen as the primary decarboxylated product (Supplementary

Fig. 12f). However, when the concentration ratios were changed to 1:5 (L-Glu with L-

CSA/L-Asp) and 1:10 (L-Glu with L-CSA/L-CA/L-Asp), a gradual increase over time in

the formation of the alternative decarboxylated products - hypotaurine, taurine, and β-

alanine was observed (Fig. 5-7,). For the $BfGAD_{WT}$ catalyzed decarboxylation reactions

of L-CSA and L-Asp, the resulting products hypotaurine and β-alanine could be

observed when these substrates were in 5-fold excess to the native substrate L-

glutamate. However, the accumulation of products hypotaurine and β-alanine becomes

significant only after 9 hours (for β-alanine) – 24 h (for hypotaurine) (Supplementary Fig.

12d, 12e). In contrast, when L-CSA and L-Asp are used in 10-fold excess of L-

glutamate, the resulting products, hypotaurine and β-alanine (peaks highlighted in pink)

started accumulating significantly much earlier around 3 hours (Fig. 5, 7). We did not

observe the accumulation of taurine even when 10-fold excess of L-CA was mixed with

L-glutamate in the $BfGAD_{WT}$ catalyzed reaction (Fig. 6). However, as mentioned earlier,

we were able to detect CO_2 in the same time frame (48 h) in the reaction catalyzed by

$BfGAD_{WT}$ when L-CA was provided as the sole substrate (Supplementary Fig. 9a). The

inability to detect taurine might be because of the low concentrations produced under

the competitive environment of the mixed substrate pool. Unlike $BfGAD_{WT,}$ $BfGAD_{D104N}$

was able to decarboxylate and accumulate taurine (peaks highlighted in pink) to a small

extent when L-CA is mixed with L-glutamate at a 10-fold excess concentration, specially

between 24-48 h (Fig. 6). Interestingly, in all mixed substrate experiments, we observed

maximum accumulation of the native product GABA (peaks highlighted in blue) within 3

h. The intensity peaks showed no further increase after 3 h time point. This shows that

even in the presence of other substrates the native activity of L-glutamate

decarboxylation occurred at the highest velocity for all *Bf*GADs compared to the

decarboxylation reactions of other substrates.

Additionally, the decarboxylated products hypotaurine and taurine were analyzed

using LC-ESI-MS/MS in addition to TLC to confirm their presence and mass in mixed

substrate assays where L-glutamate to L-CSA or L-CA ratios were 1:10. The qualitative

analysis of these molecules was conducted using MRM mode by running mixed

standards (Supplementary Fig. 13). Figure 8a depicts the presence of the product

hypotaurine and the remaining L-CSA substrate in the mixed substrate reactions with a

1:10 ratio of L-glutamate:L-CSA catalyzed by *Bf*GAD$_{WT}$. Figure 8b depicts the presence

of taurine and remaining L-CA in the mixed substrate reactions with a 1:10 ratio of L-

Glutamate:L-CA catalyzed by *Bf*GAD$_{D104N}$. We did not analyze the native substrate and

product, L-glutamate and GABA in these samples using LC-MS/MS because the

presence of these molecules was confirmed by both TLC and GABase assay prior to

LC-MS/MS analysis.

DISCUSSION.

Prior research has implicated gut microbial contributions to the production of γ-

aminobutyric acid (GABA), specifically from *Bacteroides* genus[36,37]. GABA, a

neurotransmitter, is modulated in many neurodegenerative diseases, including

Alzheimer's and dementia, where lower GABA levels have been consistently reported[12].

Microbes from the genus *Bacteroides* are known to be fluctuated in individuals afflicted

by Alzheimer's and dementia[30-33]. Due to the prevalent nature of *Bacteroides* in the gut

and the modulation of Bacteroides during Alzheimer's, we decided to understand the

mechanism by which GABA is produced in these organisms. For this reason, an

annotated glutamate decarboxylase from *B. fragilis* (*Bf*GAD) was selected as a

candidate enzyme representing annotated GADs in all *Bacteroides*.

Most of the studied microbial GADs are oligomeric which include dimeric,

tetrameric or hexameric states[44,57,61,72-74] . It is known that the functional GAD unit is a

dimer as residues from both monomers make an intact active site and play an important

role during the catalysis. Therefore, an oligomeric composition that is not an even

number would be unusual in this context. Our gel filtration chromatography data with pH

4.7 buffer gave a molecular weight of the protein that is higher than a dimer but lower

than a tetramer. We hypothesize that the protein likely exists in a dynamic equilibrium

between these two states, with a tendency towards a tetrameric state. We observed an

intermediate molecular weight possibly due to the inability to resolve dimeric and

tetrameric states completely. However, at pH 7.2, there might be a shift in this

equilibrium favoring a more dimeric state which is observed with a molecular weight

closer to that of a dimer than a tetramer. This shift in oligomeric states has been

observed previously in *E. coli* GAD that undergoes significant changes in oligomeric

forms upon shift in the pH. Thus, purified BfGAD$_{WT}$ behaves similarly in the oligomeric

composition to other characterized prokaryotic GADs. Our data suggests that there

might be a dynamic equilibrium between dimeric and tetrameric forms of BfGAD$_{WT}$.

Unlike oligomeric conformations, the kinetic parameters of BfGAD$_{WT}$ vary from

previously investigated prokaryotic GADs[44,57,61,63,72,73]. We find that BfGAD$_{WT}$ is an

allosteric enzyme, displaying a positive cooperativity with the substrate L-glutamate.

This happens when binding of substrate on one site or monomer of the protein

promotes the binding of additional substrate to the other sites or monomers of the

protein, where instead of a hyperbolic dependence, a sigmoidal curve is observed for

the plot of initial velocity vs. substrate concentration. This phenomenon has not been

seen with any other prokaryotic GAD. The physiological consequences of this

cooperativity in *Bf*GAD need to be explored further. However, cooperative binding and

adaptability in the allosteric enzymes helps amplify the enzyme's response to changes

in substrate concentration, making it more sensitive to the physiological substrate

concentrations and conditions.

Although the kinetic parameters are different for the *Bf*GAD$_{WT}$ compared to other

studied microbial GADs, there are prominent structural similarities among all these

GADs. The AlphaFold model of *Bf*GAD predicts active site architecture that is very

similar to other prokaryotic GADs. Particularly, catalytic residues in the active site are

positioned similar to those in *Ec*GAD and *Bt*GAD, suggesting that corresponding

residues in *Bf*GAD might maintain similar interactions with ligands. While the hydrogen

bonding interactions of the residues with the substrate analog glutarate are predicted to

be within the expected limits of H-bond distances, in the superimposed structures of

*Ec*Gad and *Bf*GAD$_{WT}$, there are differences in the hydrogen bond lengths when amino

acid substitutions are made in the active site. The changes in these H-bonding

interactions could potentially be involved in some capacity in accommodating or

excluding alternative substrates in the active site of the engineered *Bf*GADs to evolve

new activity or specificity.

In contrast to GADs from *Bacteroides* sp., microbial GADs from various other

genera have been widely characterized[44,61,72,75,76]. Most of these studies show that

many annotated microbial glutamate decarboxylases prefer L-glutamate as a substrate.

However, substrate promiscuity has been seen in GADs from different domains of life.

For example, *E. coli* GAD was able to convert a phosphinic analog of glutamate to a

phosphinic analog of GABA[70]. In addition, human and other mammalian GADs are

known to catalyze decarboxylation of substrates other than L-glutamate. Specifically,

sulfinic acid and sulfonic acid derivatives of alanine (known as cysteine sulfinic

acid/CSA and cysteic acid/CA respectively) are known alternative substrates for

mammalian GADs[69,71,77]. The decarboxylation of these molecules by mammalian GADs

generate hypotaruine and taurine, from the respective sulfinic or sulfonic acids.

Additionally, mammals have a couple of other *de novo* pathways for the production of

taurine and hypotaurine. The primary pathway involves the enzyme cysteine sulfinic

acid decarboxylase (CSAD) that generates taurine and its intermediate hypotaurine by

decarboxylating either L-cysteic acid (CA) or L-cysteine sulfinic acid (CSA)[71,78]. In

addition to hCSAD, mammalian GADL1 (glutamic acid decarboxylase like 1) enzyme

that has very high sequence similarity with hCSAD, is capable of producing taurine[67,79].

Taurine, a sulfur containing non-proteinogenic amino acid whose production is

controlled primarily by the enzyme cysteine sulfinic acid decarboxylase (CSAD)[80,81].

However, except for a few marine microbes the major *de novo* pathway for taurine

formation with the help of CSAD has not been seen in other prokaryotes[69]. Interestingly,

in these microbes CSAD genes are present in the operon containing cysteine

dioxygenase (CDO) enzyme that catalyzes the conversion of L-cysteine to L-cysteine

sulfinic acid (L-CSA) which then can be converted to hypotaurine and taurine

(Supplementary Fig. 14). CDOs are not present in the members of the human gut

microbiome possibly due to the hypoxic and anaerobic conditions of the gut. Despite an

extensive bioinformatics search, we did not find any gut microbial genes annotated as

CSAD (cysteine sulfinic acid decarboxylase). Due to the lack of the annotated CSAD

enzymes responsible for the *de novo* taurine biosynthesis, we investigated the role of

*Bf*GAD in the formation of taurine and its derivatives in addition to GABA.

Apart from taurine, mammalian GADL1 is also able to generate β-alanine from L-

aspartate[67,79]. β-alanine is a precursor for the dipeptide beta-alanyl-L-histidine in

humans, commonly known as carnosine. Carnosine is found in muscles and brain

tissues at high concentrations[82,83]. In microbes, β-alanine is a precursor for coenzyme A

(CoA) biosynthesis, which is an important molecule in various metabolic pathways[84].

Both carnosine and β-alanine show protective effects in individuals with cognitive

deficits and Alzhiermer's[85-87]. Moreover, higher serum concentrations of β-alanine

prevent dementia[85]. Considering the structural similarities between L- glutamate and L-

aspartate and the ability of some GADs to use L-aspartate as a substrate[65,68,88], we

examined the possible role of *Bf*GAD in the production of β-alanine via the

decarboxylation of L-aspartate. The neuromodulatory molecules, GABA, taurine and β-

alanine generated by glutamate decarboxylases or enzymes similar to GADs, have the

ability to reverse cognitive deficits in neurodegenerative disorders like dementia and

Alzheimer's[12,13,24,80,85,86,89,90]. This prompted us to understand the modulation of these

molecules by the members of the human gut microbiome.

Our results show that the $Bf\text{GAD}_{WT}$ is promiscuous and is able to decarboxylate

four additional substrates structurally similar to L-glutamate to produce hypotaurine,

taurine, homotaurine, and β-alanine. There are distinct enzymes in microbes such as

aspartate1-decarboxylase (A1DC) and aspartate 4-decarboxylase (A4DC) that catalyze

decarboxylations of L-aspartate to produce β-alanine[91,92] and L-alanine[93,94] respectively.

While the genome of *B. fragilis* has genes annotated for both these enzymes, it is

interesting to observe that the glutamate decarboxylase from *B. fragilis* still shows

decarboxylation activity towards L-aspartate to produce β-alanine. Additionally,

prokaryotic A1DC and eukaryotic A1DC show evolutionary divergence where the former

uses pyruvoyl cofactor whereas the latter uses PLP cofactor[95,96]. Thus, *Bf*GAD wild-type

and variants harbor an activity that is mostly seen in eukaryotic organisms.

To understand the factors that drive substrate specificity in *Bf*GAD$_{WT}$ and to

evolve enzymes that potentially have switched substrate preferences or specificity, we

chose two active site residues that are known to make interactions with bound

ligands[44,53]. Headspace GC was used for the detection of evolved CO_2 which is the

common decarboxylation product for all enzyme catalyzed reactions (wild-type and

engineered enzymes). Absolute quantification of $CO_{2(g)}$ was challenging due to its high

solubility at acidic pH where we conducted our enzymatic reactions. In these conditions,

the CO_2 produced through the reaction might still be in a soluble form as a dissolved

CO_2. So, headspace GC was utilized as a primary tool to identify alternate substrates by

allowing detection of CO_2 peak that served as a positive indication of enzymatic

decarboxylation of the substrate molecules. From the identification provided by the GC

experiments, subsequent analysis was carried out with TLC to detect corresponding

decarboxylated products from the multiple substrates. Our headspace GC data

reinforces our TLC data in most cases. Both engineered enzymes *Bf*GAD$_{D104N}$ and

*Bf*GAD$_{F81W}$ retain preference for L-glutamate as a substrate. However, their preferences

for alternate substrates are different than the *Bf*GAD$_{WT}$. Additionally, significant spectral

perturbations are observed with evolved GADs indicating that the local environment of

the cofactor binding site is changed due to those amino acid substitutions. With this, we

show that there might be a *de novo* pathway in gut microbes to make taurine and

*Bf*GAD$_{WT}$ is able to produce multiple neuromodulatory molecules.

References

1   Murley, A. G. *et al.* GABA and glutamate deficits from frontotemporal lobar

degeneration are associated with disinhibition. *Brain* **143**, 3449-3462 (2020).

2   Nuss, P. Anxiety disorders and GABA neurotransmission: a disturbance of

modulation. *Neuropsychiatr Dis Treat* **11**, 165-175 (2015).

3   Cawley, N. *et al.* Reduced gamma-aminobutyric acid concentration is associated

with physical disability in progressive multiple sclerosis. *Brain* **138**, 2584-2595

(2015).

4   Goddard, A. W. *et al.* Reductions in occipital cortex GABA levels in panic

disorder detected with 1h-magnetic resonance spectroscopy. *Arch Gen*

Psychiatry **58**, 556-561 (2001).

5   Goddard, A. W. *et al.* Impaired GABA neuronal response to acute
benzodiazepine administration in panic disorder. *Am J Psychiatry* **161**, 2186-
2193 (2004).

6   Long, Z. *et al.* Decreased GABA levels in anterior cingulate cortex/medial
prefrontal cortex in panic disorder. *Prog Neuropsychopharmacol Biol Psychiatry*
44, 131-135 (2013).

7   Orhan, F. *et al.* CSF GABA is reduced in first-episode psychosis and associates
to symptom severity. *Mol Psychiatry* **23**, 1244-1250 (2018).

8   Sarlo, G. L. & Holton, K. F. Brain concentrations of glutamate and GABA in
human epilepsy: A review. *Seizure* **91**, 213-227 (2021).

9   Umesawa, Y. *et al.* Altered GABA Concentration in Brain Motor Area Is
Associated with the Severity of Motor Disabilities in Individuals with Autism
Spectrum Disorder. *J Autism Dev Disord* **50**, 2710-2722 (2020).

10   Porges, E. C. *et al.* Frontal Gamma-Aminobutyric Acid Concentrations Are
Associated With Cognitive Performance in Older Adults. *Biol Psychiatry Cogn*
Neurosci Neuroimaging **2**, 38-44 (2017).

11   Carello-Collar, G. *et al.* The GABAergic system in Alzheimer's disease: a
systematic review with meta-analysis. *Mol Psychiatry* (2023).

12   Solas, M., Puerta, E. & Ramirez, M. Treatment Options in Alzheimer´s Disease:
The GABA Story. *CPD* **21**, 4960-4971 (2015).

13   Jia, F. *et al.* Taurine is a potent activator of extrasynaptic GABA(A) receptors in
the thalamus. *J Neurosci* **28**, 106-115 (2008).

14   Ochoa-de la Paz, L., Zenteno, E., Gulias-Cañizo, R. & Quiroz-Mercado, H.

Taurine and GABA neurotransmitter receptors, a relationship with therapeutic

potential? *Expert Review of Neurotherapeutics* **19**, 289-291 (2019).

15   Alom, J., Mahy, J. N., Brandi, N. & Tolosa, E. Cerebrospinal fluid taurine in

Alzheimer's disease. *Ann Neurol* **30**, 735 (1991).

16   Aytan, N. *et al.* Fingolimod modulates multiple neuroinflammatory markers in a

mouse model of Alzheimer's disease. *Sci Rep* **6**, 24939 (2016).

17   Chiquita, S. *et al.* A longitudinal multimodal in vivo molecular imaging study of the

3xTg-AD mouse model shows progressive early hippocampal and taurine loss.

Hum Mol Genet **28**, 2174-2188 (2019).

18   Garcia-Serrano, A. M., Vieira, J. P. P., Fleischhart, V. & Duarte, J. M. N. Taurine

and N-acetylcysteine treatments prevent memory impairment and metabolite

profile alterations in the hippocampus of high-fat diet-fed female mice. *Nutr*

Neurosci **26**, 1090-1102 (2023).

19   Pomara, N. *et al.* Glutamate and other CSF amino acids in Alzheimer's disease.

Am J Psychiatry **149**, 251-254 (1992).

20   Vermeiren, Y. *et al.* Behavioral correlates of cerebrospinal fluid amino acid and

biogenic amine neurotransmitter alterations in dementia. *Alzheimers Dement* **9**,

488-498 (2013).

21   Csernansky, J. G., Bardgett, M. E., Sheline, Y. I., Morris, J. C. & Olney, J. W.

CSF excitatory amino acids and severity of illness in Alzheimer's disease.

Neurology **46**, 1715-1720 (1996).

22 Gao, R., Bae, M. A., Chang, K. J. & Kim, S. H. Comparison of Urinary Excretion

of Taurine Between Elderly with Dementia and Normal Elderly. *Adv Exp Med Biol*

975 Pt 1, 57-65 (2017).

23 Fujii, Y. *et al.* Fecal metabolite of a gnotobiotic mouse transplanted with gut

microbiota from a patient with Alzheimer's disease. *Biosci Biotechnol Biochem*

83, 2144-2152 (2019).

24 Jang, H. *et al.* Taurine Directly Binds to Oligomeric Amyloid-beta and Recovers

Cognitive Deficits in Alzheimer Model Mice. *Adv Exp Med Biol* **975 Pt 1**, 233-241

(2017).

25 Kim, H. Y. *et al.* Taurine in drinking water recovers learning and memory in the

adult APP/PS1 mouse model of Alzheimer's disease. *Sci Rep* **4**, 7467 (2014).

26 Manzano, S., Aguera, L., Aguilar, M. & Olazaran, J. A Review on Tramiprosate

(Homotaurine) in Alzheimer's Disease and Other Neurocognitive Disorders. *Front*

Neurol **11**, 614 (2020).

27 Singh, P. *et al.* Taurine deficiency as a driver of aging. *Science* **380**, eabn9257

(2023).

28 Borsom, E. M., Lee, K. & Cope, E. K. Do the Bugs in Your Gut Eat Your

Memories? Relationship between Gut Microbiota and Alzheimer's Disease. *Brain*

Sci **10** (2020).

29 Saji, N. *et al.* Analysis of the relationship between the gut microbiome and

dementia: a cross-sectional study conducted in Japan. *Sci Rep* **9**, 1008 (2019).

30 Zhuang, Z.-Q. *et al.* Gut Microbiota is Altered in Patients with Alzheimer's

Disease. *JAD* **63**, 1337-1346 (2018).

31   Guo, M. *et al.* Gut Microbiome Features of Chinese Patients Newly Diagnosed

with Alzheimer's Disease or Mild Cognitive Impairment. *JAD* **80**, 299-310 (2021).

32   Li, B. *et al.* Mild cognitive impairment has similar alterations as Alzheimer's

disease in gut microbiota. *Alzheimer's & Dementia* **15**, 1357-1366 (2019).

33   Jemimah, S., Chabib, C. M. M., Hadjileontiadis, L. & AlShehhi, A. Gut

microbiome dysbiosis in Alzheimer's disease and mild cognitive impairment: A

systematic review and meta-analysis. *PLoS ONE* **18**, e0285346 (2023).

34   Salyers, A. A. Bacteroides of the human lower intestinal tract. *Annu Rev*

Microbiol **38**, 293-313 (1984).

35   Wexler, H. M. Bacteroides: the good, the bad, and the nitty-gritty. *Clin Microbiol*

Rev **20**, 593-621 (2007).

36   Strandwitz, P. *et al.* GABA-modulating bacteria of the human gut microbiota. *Nat*

Microbiol **4**, 396-403 (2019).

37   Otaru, N. *et al.* GABA Production by Human Intestinal Bacteroides spp.:

Prevalence, Regulation, and Role in Acid Stress Tolerance. *Front. Microbiol.* **12**,

656895 (2021).

38   Horvath, T. D. *et al.* Bacteroides ovatus colonization influences the abundance of

intestinal short chain fatty acids and neurotransmitters. *iScience* **25**, 104158

(2022).

39   Cozzani, I. Spectrophotometric assay of l-glutamic acid decarboxylase. *Analytical*

Biochemistry **33**, 125-131 (1970).

40   Tramonti, A., De Biase, D., Giartosio, A., Bossa, F. & John, R. A. The roles of
His-167 and His-275 in the reaction catalyzed by glutamate decarboxylase from
Escherichia coli. *J Biol Chem* **273**, 1939-1945 (1998).

41   Tsukatani, T., Higuchi, T. & Matsumoto, K. Enzyme-based microtiter plate assay
for γ-aminobutyric acid: Application to the screening of γ-aminobutyric acid-
producing lactic acid bacteria. *Analytica Chimica Acta* **540**, 293-297 (2005).

42   De Biase, D., Tramonti, A., John, R. A. & Bossa, F. Isolation, overexpression,
and biochemical characterization of the two isoforms of glutamic acid
decarboxylase from Escherichia coli. *Protein Expr Purif* **8**, 430-438 (1996).

43   Tramonti, A., John, R. A., Bossa, F. & De Biase, D. Contribution of Lys276 to the
conformational flexibility of the active site of glutamate decarboxylase from
Escherichia coli. *European Journal of Biochemistry* **269**, 4913-4920 (2002).

44   Capitani, G. *et al.* Crystal structure and functional analysis of *Escherichia coli*
glutamate decarboxylase. *EMBO J* **22**, 4027-4037-4037 (2003).

45   Tramonti, A. *et al.* A Novel, Easy Assay Method for Human Cysteine Sulfinic Acid
Decarboxylase. *Life* **11**, 438 (2021).

46   Goujon, M. *et al.* A new bioinformatics analysis tools framework at EMBL-EBI.
Nucleic Acids Res **38**, W695-699 (2010).

47   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).

48   Miller, M. A., Pfeiffer, W. & Schwartz, T. in *2010 Gateway Computing*
Environments Workshop (GCE). 1-8.

776 49 Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid

777   multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res*

778   **30**, 3059-3066 (2002).

779 50 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: computing large minimum

780   evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* **26**, 1641-

781   1650 (2009).

782 51 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-

783   likelihood trees for large alignments. *PLoS ONE* **5**, e9490 (2010).

784 52 Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for

785   phylogenetic tree display and annotation. *Nucleic Acids Res* **49**, W293-W296

786   (2021).

787 53 Dutyshev, D. I. *et al.* Structure of Escherichia coli glutamate decarboxylase

788   (GADα) in complex with glutarate at 2.05 Å resolution. *Acta Crystallographica*

789   *Section D Biological Crystallography* **61**, 230-235 (2005).

790 54 Beattie, A. E. *et al.* The Pyridoxal 5′-Phosphate (PLP)-Dependent Enzyme Serine

791   Palmitoyltransferase (SPT): Effects of the Small Subunits and Insights from

792   Bacterial Mimics of Human hLCB2a HSAN1 Mutations. *BioMed Research*

793   *International* **2013**, 1-13 (2013).

794 55 Moore, P. S., Dominici, P. & Borri Voltattorni, C. Cloning and expression of pig

795   kidney dopa decarboxylase: comparison of the naturally occurring and

796   recombinant enzymes. *Biochemical Journal* **315**, 249-256 (1996).

56 Zhou, X. & Toney, M. D. pH Studies on the Mechanism of the Pyridoxal
Phosphate-Dependent Dialkylglycine Decarboxylase. *Biochemistry* **38**, 311-320
(1999).

57 Liu, S. *et al.* Coordinated regulation of Bacteroides thetaiotaomicron glutamate
decarboxylase activity by multiple elements under different pH. *Food Chem* **403**,
134436 (2023).

58 Pennacchietti, E. *et al.* Mutation of His465 Alters the pH-dependent
Spectroscopic Properties of Escherichia coli Glutamate Decarboxylase and
Broadens the Range of Its Activity toward More Alkaline pH. *Journal of Biological*
Chemistry **284**, 31587-31596 (2009).

59 Shukuya, R. & Schwert, G. W. Glutamic acid decarboxylase. II. The spectrum of
the enzyme. *J Biol Chem* **235**, 1653-1657 (1960).

60 De Biase, D. & Pennacchietti, E. Glutamate decarboxylase-dependent acid
resistance in orally acquired bacteria: function, distribution and biomedical
implications of the gadBC operon. *Mol Microbiol* **86**, 770-786 (2012).

61 Chang, C. *et al.* Purification and characterization of glutamate decarboxylase
from Enterococcus raffinosus TCCC11660. *J Ind Microbiol Biotechnol* **44**, 817-
824 (2017).

62 Fonda, M. L. Glutamate decarboxylase. Substrate specificity and inhibition by
carboxylic acids. *Biochemistry* **11**, 1304-1309 (1972).

63 Ueno, Y., Hayakawa, K., Takahashi, S. & Oda, K. Purification and
characterization of glutamate decarboxylase from Lactobacillus brevis IFO
12005. *Biosci Biotechnol Biochem* **61**, 1168-1171 (1997).

64   Fonda, M. L. L-Glutamate decarboxylase from bacteria. *Methods Enzymol* **113**,

11-16 (1985).

65   Kim, H. W., Kashima, Y., Ishikawa, K. & Yamano, N. Purification and

characterization of the first archaeal glutamate decarboxylase from Pyrococcus

horikoshii. *Biosci Biotechnol Biochem* **73**, 224-227 (2009).

66   Tomita, H., Yokooji, Y., Ishibashi, T., Imanaka, T. & Atomi, H. An archaeal

glutamate decarboxylase homolog functions as an aspartate decarboxylase and

is involved in beta-alanine and coenzyme A biosynthesis. *J Bacteriol* **196**, 1222-

1230 (2014).

67   Liu, P. *et al.* Role of glutamate decarboxylase-like protein 1 (GADL1) in taurine

biosynthesis. *J Biol Chem* **287**, 40898-40906 (2012).

68   Richardson, G. *et al.* An examination of aspartate decarboxylase and glutamate

decarboxylase activity in mosquitoes. *Mol Biol Rep* **37**, 3199-3205 (2010).

69   Agnello, G., Chang, L. L., Lamb, C. M., Georgiou, G. & Stone, E. M. Discovery of

a Substrate Selectivity Motif in Amino Acid Decarboxylases Unveils a Taurine

Biosynthesis Pathway in Prokaryotes. *ACS Chem. Biol.* **8**, 2264-2271 (2013).

70   Daniela De Biase, F. C., Eugenia Pennacchietti, Fabio Giovannercole, Antonio

Coluccia, Jouko Vepsäläinen & Alex Khomutov Enzymatic kinetic resolution of

desmethylphosphinothricin indicates that phosphinic group is a bioisostere of

carboxyl group. *Commnications chemistry* (2020).

71   Wu, J. Y. Purification and characterization of cysteic acid and cysteine sulfinic

acid decarboxylase and L-glutamate decarboxylase from bovine brain. *Proc Natl

Acad Sci U S A* **79**, 4270-4274 (1982).

72   Komatsuzaki, N., Nakamura, T., Kimura, T. & Shima, J. Characterization of

glutamate decarboxylase from a high gamma-aminobutyric acid (GABA)-

producer, Lactobacillus paracasei. *Biosci Biotechnol Biochem* **72**, 278-285

(2008).

73   Hiraga, K., Ueno, Y. & Oda, K. Glutamate decarboxylase from Lactobacillus

brevis: activation by ammonium sulfate. *Biosci Biotechnol Biochem* **72**, 1299-

1306 (2008).

74   Giovannercole, F. *et al.* On the effect of alkaline pH and cofactor availability in

the conformational and oligomeric state of Escherichia coli glutamate

decarboxylase. *Protein Eng Des Sel* **30**, 235-244 (2017).

75   Lee, J. Y. & Jeon, S. J. Characterization and immobilization on nickel-chelated

Sepharose of a glutamate decarboxylase A from Lactobacillus brevis BH2 and its

application for production of GABA. *Biosci Biotechnol Biochem* **78**, 1656-1661

(2014).

76   Nomura, M. *et al.* Lactococcus lactis contains only one glutamate decarboxylase

gene. *Microbiology (Reading)* **145 (Pt 6)**, 1375-1380 (1999).

77   Blindermann, J. M., Maitre, M., Ossola, L. & Mandel, P. Purification and some

properties of L-glutamate decarboxylase from human brain. *Eur J Biochem* **86**,

143-152 (1978).

78   Winge, I. *et al.* Mammalian CSAD and GADL1 have distinct biochemical

properties and patterns of brain expression. *Neurochemistry International* **90**,

173-184 (2015).

79 Mahootchi, E. *et al.* GADL1 is a multifunctional decarboxylase with tissue-specific

roles in beta-alanine and carnosine production. *Sci Adv* **6**, eabb3713 (2020).

80 Ripps, H. & Shen, W. Review: taurine: a "very essential" amino acid. *Mol Vis* **18**,

2673-2686 (2012).

81 De La Rosa, J. & Stipanuk, M. H. Evidence for a rate-limiting role of

cysteinesulfinate decarboxylase activity in taurine biosynthesis in vivo.

Comparative Biochemistry and Physiology Part B: Comparative Biochemistry **81**,

565-571 (1985).

82 Boldyrev, A. A., Aldini, G. & Derave, W. Physiology and pathophysiology of

carnosine. *Physiol Rev* **93**, 1803-1845 (2013).

83 Sale, C., Saunders, B. & Harris, R. C. Effect of beta-alanine supplementation on

muscle carnosine concentrations and exercise performance. *Amino Acids* **39**,

321-333 (2010).

84 Lopez-Samano, M. *et al.* A novel way to synthesize pantothenate in bacteria

involves beta-alanine synthase present in uracil degradation pathway.

Microbiologyopen **9**, e1006 (2020).

85 Hata, J. *et al.* Association Between Serum beta-Alanine and Risk of Dementia.

Am J Epidemiol **188**, 1637-1645 (2019).

86 Ostfeld, I. *et al.* Role of β-Alanine Supplementation on Cognitive Function,

Mood, and Physical Function in Older Adults; Double-Blind Randomized

Controlled Study. *Nutrients* **15** (2023).

87   Dhingra, D., Parle, M. & Kulkarni, S. K. β - Alanine protects mice from memory

deficits induced by ageing, scopolamine, diazepam and ethanol. *Indian Journal*

of Pharmaceutical Sciences **68** (2006).

88   Wu, J.-Y., Matsuda, T. & Roberts, E. Purification and Characterization of

Glutamate Decarboxylase from Mouse Brain. *Journal of Biological Chemistry*

248, 3029-3034 (1973).

89   Albrecht, J. & Schousboe, A. Taurine Interaction with Neurotransmitter Receptors

in the CNS: An Update. *Neurochem Res* **30**, 1615-1621 (2005).

90   Bae, M. A., Lee, E. S., Cho, S. M., Kim, S. H. & Chang, K. J. The Effects of

Dietary Taurine-Containing Jelly Supplementation on Cognitive Function and

Memory Ability of the Elderly with Subjective Cognitive Decline. *Adv Exp Med*

Biol **1370**, 395-403 (2022).

91   Cui, W., Liu, H., Ye, Y., Han, L. & Zhou, Z. Discovery and Engineering of a Novel

Bacterial L-Aspartate alpha-Decarboxylase for Efficient Bioconversion. *Foods* **12**

(2023).

92   Begley, T. P., Kinsland, C. & Strauss, E. The biosynthesis of coenzyme A in

bacteria. *Vitam Horm* **61**, 157-171 (2001).

93   Lima, S. *et al.* The crystal structure of the Pseudomonas dacunhae aspartate-

beta-decarboxylase dodecamer reveals an unknown oligomeric assembly for a

pyridoxal-5'-phosphate-dependent enzyme. *J Mol Biol* **388**, 98-108 (2009).

94   Jha, S., Speth, R. C. & Macheroux, P. The possible role of a bacterial aspartate

beta-decarboxylase in the biosynthesis of alamandine. *Med Hypotheses* **144**,

110038 (2020).

95   Yu, X. J. *et al.* Protein Engineering of a Pyridoxal-5'-Phosphate-Dependent l-
Aspartate-alpha-Decarboxylase from Tribolium castaneum for beta-Alanine
Production. *Molecules* **25** (2020).

96   Webb, M. E. *et al.* Structure of Escherichia coli aspartate alpha-decarboxylase
Asn72Ala: probing the role of Asn72 in pyruvoyl cofactor formation. *Acta*
Crystallogr Sect F Struct Biol Cryst Commun **68**, 414-417 (2012).

Acknowledgments. This research is supported by the seed grant from Edson initiative

for the dementia care and solutions to DDS. We acknowledge resources and support

from the Knowledge Enterprise Biosciences Core Facilities at Arizona State University.

We thank Dr. Rosa Krajmalnik-Brown and Christopher Muse from the Biodesign Centre

for Health Through Microbiomes for the help with headspace GC.

Author Contributions. PD performed majority of the experiments and data analysis.

CWP performed some of the TLC experiments. AS performed analysis and data fitting

for pH and Abs variation experiment and provided code and advice on the image

analysis of TLC plates, PD and DDS conceptualized the project and wrote the paper.

Data availability. All the data are available in the manuscript or with associated

supporting information. The python code used for TLC image analysis is available at

https://github.com/Pavani-dadi/TLC_profile.

Figures.

Fig. 1: Characterization of *Bf*GAD$_{WT}$. a Possible role of gut microbes in producing neuromodulatory molecules. **b** Spectra of PLP cofactor bound to the holoenzyme *Bf*GAD$_{WT}$ captured at various pH, depicting tautomers of internal aldimine at 335 nm and 420 nm. C) Plot of initial velocity as a function of *Bf*GAD$_{WT}$ concentrations (n=4). D) Plot of initial velocity vs. L-glutamate concentrations (n=2). The solid line is a sigmoidal fit to the data.

Fig. 2: Structural and catalytic perturbations of engineered *Bf*GADs. a Aligned

structures of *Ec*GAD (1XEY, pink), *Bf*GAD$_{WT}$ (model, green) illustrating the position of

the residue D104* (*residue from the neighbouring monomer), and *Bf*GAD$_{D104N}$

depicting the point substitution D104N*(model, gray). **b** Aligned structures of *Ec*GAD

(1XEY, pink), *Bf*GAD$_{WT}$ (model, green) illustrating the position of the residue F81, and

*Bf*GAD$_{F81W}$ depicting the point substitution F81W (model, gray). **c** Aligned structures of

*Bt*GAD (7X51, purple), *Bf*GAD$_{WT}$ (model, green) illustrating the position of the residue

D104* (*residue from the neighbouring monomer), and $BfGAD_{D104N}$ depicting the point

substitution D104N*(model, gray). **d** Aligned structures of BtGAD (7X51, purple),

$BfGAD_{WT}$ (model, green) illustrating the position of the residue F81, and $BfGAD_{F81W}$

depicting the point substitution F81W (model, gray).   The dashed lines indicate

hydrogen bond interactions of the amino acid residues with substrate analog, glutarate

(GUA) where pink bonds are for interactions between EcGAD-GUA, yellow bonds are

for interactions between BfGAD-GUA, and purple bonds are for interactions between

BtGAD-GUA. **e** Spectra of PLP cofactor bound to the holoenzymes $BfGAD_{WT}$ and

engineered variants, depicting tautomers of internal aldimine at 335 nm and 420 nm. **f**

Progress curves depicting initial velocities for reactions catalyzed $BfGAD_{WT}$ and variants

(n=3).

Fig. 3: Wild-type and engineered *Bf*GADs can decarboxylate multiple substrates.

Chromatograms showing the CO_2 peaks obtained from the decarboxylation reactions of multiple substrates after 24 h incubation with **a** *Bf*GAD$_{WT}$, **c** *Bf*GAD$_{D104N}$, and **e** *Bf*GAD$_{F81W}$ using headspace GC (n=3, only one representative CO_2 peak is depicted for every reaction). The CO_2 peak areas are depicted for the decarboxylation of various substrates after 24 h incubation with **b** BfGAD$_{WT}$, **d** BfGAD$_{D104N}$, and **f** BfGAD$_{F81W}$ (n=3). Various substrates are presented as specific colors which are consistent in all panels

where L-glutamate is blue, L-CSA is red, L-Asp is purple, L-HCA is orange and L-CA is

green.

Fig. 4: Production of multiple neuromodulatory molecules by Wild-type and

evolved *Bf*GADs. Intensity analysis of TLC plates depicting intensity plots for each

lane. The peak size is proportional to the intensity of the sample spot on the TLC plate.

A Production of hypotaurine from the decarboxylation of L-CSA catalyzed by $BfGAD_{WT}$

and $BfGAD_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks

highlighted in pink are for the substrate L-CSA and peaks highlighted in blue are for the

product hypotaurine. **b** Production of taurine from the decarboxylation of L-CA catalyzed

by $BfGAD_{WT}$ and $BfGAD_{D104N}$ in 24 h as indicated by TLC intensity analysis. For

intensity plots, peaks highlighted in pink are for the substrate L-CA and peaks

highlighted in blue are for the product taurine. **c** Production of β-alanine from the

decarboxylation of L-ASP catalyzed by $BfGAD_{WT}$ and $BfGAD_{D104N}$ in 24 h as indicated

by TLC intensity analysis. For intensity plots, peaks highlighted in pink are for the

substrate L-ASP and peaks highlighted in blue are for the product β-alanine. **d**

Production of homotaurine from the decarboxylation of L-HCA catalyzed by $BfGAD_{WT}$

and $BfGAD_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks

highlighted in pink are for the substrate L-HCA and peaks highlighted in blue are for the

product homotaurine. The original TLC plates are shown in Supplementary Fig. 10.

Fig. 5: *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ generate hypotaurine when presented with a

mixture of L-glutamate and L-CSA. Intensity analysis of TLC plates for various

timepoints of decarboxylation reactions catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ for

mixed substrates where the concentration ratio of L-glutamate to L-CSA is 1:10.

Standards of L-glutamate, L-CSA, GABA, and hypotaurine were included as references.

Production of hypotaurine (peaks highlighted in pink) can be seen as the peak area

increases over time. GABA peaks are highlighted in blue, peaks for the substrate L-CSA

are highlighted in beige, and peaks for the substrate L-glutamate are in brown. Spots

visible on TLC plates from top to bottom show corresponding intensity peaks from left to

right.

Fig. 6: *Bf*GAD$_{D104N}$ generate taurine when presented with a mixture of L-glutamate

and L-CA. Intensity analysis of TLC plates for various timepoints of decarboxylation

reactions catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ for mixed substrates where the

concentration ratio of L-glutamate to L-CA is 1:10. Standards of L-glutamate, L-CA,

GABA, and taurine were included as references. Production of taurine (peaks

highlighted in pink) can be seen as the peak area increases over time for *Bf*GAD$_{D104N}$

catalyzed reactions. GABA peaks are highlighted in blue, peaks for the substrate L-CA

are highlighted in beige, and peaks for the substrate L-glutamate are in brown. Spots

visible on TLC plates from top to bottom show corresponding intensity peaks from left to

right.

1015

1016

Fig. 7: *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ generate β-alanine when presented with a

mixture of L-glutamate and L-aspartate. Intensity analysis of TLC plates for various

timepoints of decarboxylation reactions catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ for

mixed substrates where the concentration ratio of L-glutamate to L-aspartate is 1:10.

Standards of L-glutamate, L-aspartate, GABA, and β-alanine were included as

references. Production of β-alanine (peaks highlighted in pink) can be seen as the peak

area increases over time. The increase over time for *Bf*GAD$_{D104N}$ is very clear from 3 h

to 9 h. Due to the differences in the background intensity of the TLC plate in areas

containing 3 h and 9 h samples vs areas containing 24 h, 30 h and 48 h samples from

*Bf*GAD$_{D104N}$ catalyzed reaction, the increase is the intensity peak is not uniform. GABA

peaks are highlighted in blue, peaks for the substrate L-aspartate (L-ASP) are

highlighted in beige, and peaks for the substrate L-glutamate are in brown. Spots visible

on TLC plates from top to bottom show corresponding intensity peaks from left to right.

1031

1032

1033
1034

Figure 8: **Hypotaurine and Taurine detection via LC-MS/MS from *Bf*GAD$_{WT}$ and**

***Bf*GAD$_{D104N}$ catalyzed mixed substrate reactions. a** Ion chromatograms of product

hypotaurine and the remaining L-CSA substrate in the mixed substrate reactions with a

1:10 ratio of L-glutamate:L-CSA catalyzed by *Bf*GAD_WT. **b** Ion chromatograms of taurine

and remaining L-CA in the mixed substrate reactions with a 1:10 ratio of L-Glutamate:L-

CA catalyzed by *Bf*GAD_D104N.

1041
1042
1043

1044
1045 **Scheme 1.** Predicted substrates and products of *Bf*GAD catalyzed decarboxylations.

1046
1047