Synthesis of versatile neuromodulatory molecules by a gut microbial glutamate decarboxylase

Authors. Pavani Dadi[1,2], Clint W. Pauling[1,3], Abhishek Shrivastava[1,2], and Dhara D. Shah*[1,3]

Affiliations. [1]Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University, Tempe, AZ 85281

[2]School of Life Sciences, Arizona State University, Tempe, AZ 85281

[3]School of Mathematical and Natural Sciences, Arizona State University, Glendale, AZ 85306

*Corresponding author: dhara.shah1@asu.edu

ABSTRACT

Dysbiosis of the microbiome correlates with many neurological disorders, yet very little is known about the chemistry that controls the production of neuromodulatory molecules by gut microbes. Here, we found that an enzyme glutamate decarboxylase (*Bf*GAD) of a gut microbe *Bacteroides fragilis* forms multiple neuromodulatory molecules such as γ-aminobutyric acid (GABA), hypotaurine, taurine, homotaurine, and β-alanine. We evolved *Bf*GAD and doubled its taurine productivity. Additionally, we increased its specificity towards the substrate L-glutamate. Here, we provide a chemical strategy via which the *Bf*GAD activity could be fine-tuned. In future, this strategy could be used to modulate the production of neuromodulatory molecules by gut microbes.

Keywords: GABA, Taurine, Bacteroide fragilis, Neuromodulatory molecules, Glutamate decarboxylase, microbiome, hypotaurine, homotaurine

INTRODUCTION

Low levels of inhibitory neurotransmitter like γ-aminobutyric acid (GABA) have been associated with many different neurological disorders such as epilepsy, schizophrenia, autism spectrum disorder (ASD), ADHD, panic disorder, PTSD, major depressive disorder, progressive multiple sclerosis, dementia, and Alzheimer's disease[1-12]. Besides GABA, taurine is another amino acid that is found in abundance in the brain[13]. Taurine is a GABA$_A$ receptor agonist and exerts downstream effects similar to GABA[14]. Concentrations of taurine are altered in individuals with Alzheimer's disease, where reduced levels of taurine are found in the brain[15-21] whereas, higher urinary excretion of taurine is seen in elderly patients with dementia[22]. In a separate study, gnotobiotic mouse transplanted with fecal microbiota from an Alzheimer's patient showed lower abundance

of GABA and taurine in the feces[23]. Based on these observations, supplementation with

taurine and its analogs has been investigated in Alzheimer's disease models[24,25]. In one

such study, Homotaurine, a molecule similar to taurine is shown to restore cognitive

functions in patients with Alzheimer's disease[26]. In addition, it was also observed that

taurine reverses cognitive deficits in APP/PS1 mouse model[25] and improves learning and

memory in mice[24,27].

Studies that correlate gut microbial dysbiosis to two of the prominent

neurodegenerative disorders – Dementia and Alzheimer's disease, often show

modulations in the abundance of microbes of the genus *Bacteroides*[28-33]. *Bacteroides* are

prominent members of the human gut[34,35] and have the ability to produce GABA[36,37]. All

Bacteroides encode the gene for the enzyme glutamate decarboxylase (GAD)[38].

Glutamate decarboxylases (GADs) are PLP (pyridoxal phosphate) dependent enzymes

which catalyze conversion of an excitatory neurotransmitter glutamate to an inhibitory

neurotransmitter GABA. However, very little is known about the chemistry of the

glutamate decarboxylases encoded by *Bacteroides sp.*

Despite widespread prevalence of substrate promiscuity in glutamate

decarboxylases (GADs) across different domains of life, substrate specificity and product

formation landscape for most prokaryotic GADs, specifically GADs of *Bacteroides sp.* are

still not clearly understood. For this reason, we explored the role of annotated GAD from

Bacteroides fragilis in the production of neuromodulatory molecules like GABA, taurine

and its analogs, and β-alanine. Our biochemical characterization of *Bf*GAD shows kinetic

and functional divergence of this enzyme from other studied prokaryotic GADs. Our study

shows that *Bf*GAD is not only specific to L-glutamate, but it can decarboxylate other

substrates to produce multiple neuromodulatory molecules. Even within the substrate mix

containing L-glutamate, we can detect products other than GABA. This points towards

the ability of this enzyme to function in a complex system with multiple substrates which

can be a beneficial trait for microbes present in the human gut. Through rational protein

engineering, we have evolved *Bf*GAD which is capable of producing two-fold more taurine

compared to the native *Bf*GAD. Additionally, we evolved *Bf*GAD to be more specific

towards L-glutamate.

Based on our results of initial engineering with *Bf*GAD, the enzyme seems resilient

in nature and is able to tolerate changes to the active site very well. We think that *Bf*GAD

can be evolved with rational designing and engineering, either to produce various

neuromodulatory molecules in specific ratios, or to synthesize a particular

neuromodulatory molecule exclusively (Fig. 1a). This approach has the ability to generate

variants of *Bf*GAD that might provide a road to therapeutic interventions for multiple

neurodegenerative disorders.

RESULTS

Glutamate Decarboxylases (GADs) are prevalent in the *Bacteroides* genus. A

bioinformatics analysis showed that many species of the *Bacteroides* genus harbor genes

annotated as GAD (Supplementary Fig. 1). We focused on a GAD (*Bf*GAD) from the

model gut microbe, *Bacteroides fragilis*. Multiple sequence alignment performed with gut

microbial GADs including *Bf*GAD revealed that multiple catalytic residues, either

predicted to be involved in substrate binding (highlighted in cyan) or cofactor (PLP)

binding (highlighted in yellow) are highly conserved (Supplementary Fig. 2)[39,40].

Moreover, in a phylogenetic analysis of *Bf*GAD with other gut microbial GADs, we see

that the annotated glutamate decarboxylases from *Alistipes putredinis* and

Parabacteroides merdae are closely related to *Bf*GAD (Supplementary Fig. 3).

Wild-type *Bf*GAD (*Bf*GAD$_{WT}$) forms an oligomer. The recombinant wild-type *Bf*GAD

(*Bf*GAD$_{WT}$) with an N-terminal His-tag was purified by Ni-NTA affinity chromatography. A

protein band with an approximate expected mass of a *Bf*GAD$_{WT}$ monomer (calculated

mass 56 KDa) was observed with SDS PAGE, while a band indicating a tetrameric

oligomer was observed with a native PAGE (Supplementary Fig. 4). Oligomeric

composition of the *Bf*GAD$_{WT}$, as determined by gel filtration chromatography, is presented

in Supplementary Fig. 5a as chromatograms for the *Bf*GAD$_{WT}$ at an acidic pH (blue) and

a neutral pH (red). Peak 1 was in the void volume of the column indicating protein

aggregates. There were two peaks representing two different oligomeric compositions (2

and 3) observed at each pH. Peak 2 was broad and corresponded to approximately 351

KDa, indicating a hexamer. Peak 3 was the major peak and exhibited an approximate

molecular weight between a dimer and a tetramer at pH 4.7, whereas at pH 7.2 the peak

3 appeared to be closer to a dimer than a tetramer (Supplementary Fig. 5, a-c, 180 KDa

at pH 4.7 and 150 KDa at pH 7.2) for *Bf*GAD$_{WT}$. Our data indicated that *Bf*GAD$_{WT}$ is an

oligomeric enzyme which can either be a dimer or a tetramer. Furthermore, the addition

of external PLP did not change the oligomeric state of the *Bf*GAD$_{WT}$ (Supplementary Fig.

5d).

PLP cofactor bound to the *Bf*GAD$_{WT}$ undergoes pH dependent tautomeric changes.

Spectral studies with *Bf*GAD$_{WT}$ revealed that the enzyme was purified in its holo form with

the covalently bound PLP cofactor. There were two absorption maxima observed for the

*Bf*GAD$_{WT}$ – 335 nm and 420 nm (Fig. 1b). These absorption maxima result from two

tautomeric forms of the enzyme bound PLP cofactor, enolimine form at 335 nm and

ketoenamine form at 420 nm[41-43]. The proportions of these tautomers vary depending on

the pH (Fig. 1b). At lower pH, the species absorbing at 420 nm was more dominant,

whereas at higher pH, 340 nm absorbing form became more prevalent. With increase in

pH from 4.7 to 8.6, there was a decrease in 420 nm absorbing species with concomitant

increase in 340 nm absorbing species (Fig. 1b). These changes in the absorbances of

the bound PLP cofactor as the function of the change in pH with BfGAD$_{WT}$, have been

seen with other microbial GADs[44-46]. The analysis of the pH versus absorbance curve for

the 420 nm wavelength indicated that the transition in absorbance in response to pH

variation is influenced by the involvement of multiple protons (Supplementary Fig. 6a).

Previously with $E.\ coli$ GAD, Tramonti et al. and Capitani et al. showed that the less

hydrated active site at neutral pH drove the formation of enolimine tautomer (340 nm)

whereas more polar active site at acidic pH predominantly produced ketoenamine

tautomer (420 nm)[39,47]. Additionally, the alterations in cofactor absorbance corresponding

to different pH were in close agreement with the BfGAD$_{WT}$'s activity profiles measured

across the same range of pH values. Particularly, at pH 4.7, where BfGAD$_{WT}$ with a bound

PLP predominantly absorbed at 420 nm, the enzyme also exhibited its maximum catalytic

activity (Supplementary Fig. 6b). Additionally, enzymatic activity was found to be minimal

at a pH other than 4.7 (Supplementary Fig. 6b). Therefore, all subsequent activity assays

were carried out at pH 4.7.

Evidence for allosteric regulation of BfGAD$_{WT}$. The activity assays conducted with

varying BfGAD$_{WT}$ concentrations exhibited a linear relationship between the initial

reaction velocities and enzyme concentrations (Fig. 1c). From the plot of enzyme

concentration variation, we found an optimal $BfGAD_{WT}$ concentration for subsequent

kinetic assays. Interestingly, while varying L-glutamate concentrations, the plot of

substrate concentration versus initial velocity displayed a sigmoidal curve instead of a

typical hyperbolic dependence, suggestive of an allostery in $BfGAD_{WT}$. This is an example

of a homotropic allosteric regulation. From the fit of this plot with the hill equation (eq. 2),

the analyzed $K_{0.5}$ (K_{half}) was 8.6 ± 0.4 mM and V_{max} was 0.86 ± 0.04 mM/min with a hill

coefficient of 1.7 (Fig. 1d). $(K_{half})^h$ is also known as K_{prime} (K'), which is equivalent to K_m

from a hyperbolic kinetics of non-allosteric enzymes, when h=1. However, in cooperative

systems where h $\neq$1 like presented here for $BfGAD_{WT}$ then, K' (K_{half}^h) no longer represents

the substrate concentration required to achieve half of maximal velocity.

Engineered BfGAD variants exhibit structural and catalytic perturbations. The

earliest microbial GAD structures were solved for *E. coli* GADs (GAD*A*, PDB ID 1XEY

and GAD*B*, PDB ID 1PMM) (Supplementary Fig. 7)[39,40]. Both structures were solved with

the bound ligand in the active site. Based on the interactions of the acetate ion in the

GAD*B* active site and glutarate (substrate analog) in the GAD*A* active site

(Supplementary Fig. 7b), we selected residues for the evolution of the $BfGAD_{WT}$ via

rational design[39,40,48]. Two such residues that are present in the active sites of both

*Ec*GADs are phenylalanine 63 (corresponds to F81 in $BfGAD_{WT}$) and aspartate 86

(corresponds to D104 in $BfGAD_{WT}$) that make H-bonding interactions with one of the

carboxylates of the substrate analog and an acetate ion[39,40]. Specifically, in *E. coli* GAD*B*

ligand bound structure, the carboxylate group of acetate received a hydrogen bond from

the amide nitrogen of the F63 (F81 $BfGAD_{WT}$) and a side chain carboxylate of D86 (D104

$BfGAD_{WT}$) of the neighboring subunit[39]. Whereas in *E. coli* GAD*A* ligand bound structure,

one of the carboxylate group of the glutarate (substrate analogue) forms H-bond with the

amide nitrogen of F63 (F81 BfGAD$_{WT}$) and with the side chain carboxylate of D86 (D104

BfGAD$_{WT}$) of the neighboring subunit[40]. These residues are completely conserved in

annotated GADs from prominent gut microbes with the exception of *Eggerthella lenta* that

harbors shorter GAD with the absence of a conserved phenylalanine (Supplementary Fig.

2). Most microbial GADs are functionally active as dimers, with residues from both

monomers contributing to the active site[39,40,44,49]. Recently, structure of glutamate

decarboxylase from *Bacteroides thetaiotamicron* (BtGAD), bound with a substrate analog

(glutarate) was solved[44]. With these available structures, we aligned the substrate analog

(glutarate) bound structures of EcGADA (PDB ID 1XEY) and BtGAD (PDB ID 7X51) with

BfGAD$_{WT}$ dimer created with AlphaFold2 (Fig. 2). Based on these structural alignments

we engineered two enzymes by introducing single amino acid alterations at two separate

positions to test if the substrate preference of the BfGAD$_{WT}$ can be evolved. These

changes were Asp104Asn (D104N) and Phe81Trp (F81W). Figure 2a and 2b illustrate

the locations of these residues within the active site of BfGAD$_{WT}$ (green). These residues

are depicted in close proximity to glutarate and the cofactor PLP bound in the active site

of EcGADA, which has been superimposed on the BfGAD$_{WT}$ dimer (Fig. 2a, 2b). Similarly,

Fig. 2c and 2d display aligned BfGAD$_{WT}$ and BtGAD structures with glutarate and PLP

bound in the active site of BtGAD. In addition to BfGAD$_{WT}$, the structures of variants,

BfGAD$_{D104N}$ (Fig. 2a, 2c) and BfGAD$_{F81W}$ (Fig. 2b, 2d) were then superimposed with ligand

bound structures of EcGAD and BtGAD.

The overall active site architecture of BfGAD is mostly consistent with that of

EcGAD and BtGAD; however, notable positional shifts were identified among the residues

of interest. Specially, aspartic acid (D104) showed a considerable difference in EcGAD

versus BfGAD active site. Additionally, we detected changes in the H-bonding interactions

between the substrate analog and residues of interest when substitutions were made to

create engineered BfGAD variants. We were able to purify both variants, BfGAD$_{D104N}$ and

BfGAD$_{F81W}$ with the same conditions as the BfGAD$_{WT}$. Both of these variants were active

and purified as holoenzymes (PLP bound form). Additionally, we collected UV-Vis spectra

of BfGAD$_{WT}$ and both variants at pH 4.7 (Fig. 2e). While each of the three enzyme

preparations exhibits absorbance maxima at both 335 nm and 420 nm, the relative

amounts of these two tautomeric forms differ across the enzymes. Furthermore, we

measured initial velocities in the presence of the native substrate L-glutamate via

progress curve analysis for engineered BfGADs along with BfGAD$_{WT}$. With that, we

detected variations in the decarboxylation reaction rates across these enzymes (Fig. 2f).

BfGAD$_{D104N}$ showed 1.7-fold increase whereas BfGAD$_{F81W}$ showed 2.5-fold decrease in

the initial velocities compared to the BfGAD$_{WT}$ for decarboxylation reactions with L-

glutamate. These observed differences in catalysis between the wild-type and engineered

variants of the BfGAD due to the structural perturbations resulting from amino acid

substitutions at the active site, reinforce the role of F81 and D104 in establishing

interactions with the substrate.

Wild-type and engineered BfGADs can decarboxylate substrates other than L-

glutamate. As mentioned above, the purified BfGAD$_{WT}$ is able to catalyze the conversion

of L-glutamate to GABA. To test if it can utilize the D-form of the glutamate as a substrate,

we conducted activity assays with D-glutamate using the GABase assay system and TLC.

Our results show that both the wild-type and engineered BfGADs were unable to

decarboxylate D-glutamate (Supplementary Fig. 8). Most microbial GADs studied so far

show high substrate specificity towards L-glutamate[49-52]. However, there are examples

of archaeal GADs that prefer L-aspartate over L-glutamate[53,54]. An archaeal GAD from *P.*

horikoshii also shows decarboxylation activity with L-cysteate[53]. In addition to archaeal

GADs, certain mammalian glutamate decarboxylase (GAD) homologs are known to

decarboxylate one or more of the non-native substrates such as L-aspartate, L-cysteate,

and L-cysteine sulfinate[55,56]. Moreover, proteins similar to GADs have been demonstrated

to form taurine in some marine microbes[57]. However, there are limited systematic studies

investigating the production of taurine, its analogs, and β-alanine using L-CA (cysteate),

L-CSA (cysteine sulfinate), L-HCA (homocysteate), and L-aspartate as substrates

(Scheme 1), specifically for gut microbial GADs. Based on two key observations, we

hypothesized that gut microbial GADs might utilize these molecules as substrates: 1. *E.*

coli GAD can accept the phosphonated form of L-glutamate as a substrate[58]. and 2.

Archaeal and eukaryotic GADs exhibit diverse substrate specificities, enabling them to

catalyze the decarboxylation reactions of non-native substrates[53,54,57,59,60].

If *Bf*GAD$_{WT}$ or engineered *Bf*GADs are able to decarboxylate substrates other than

the native substrate L-glutamate, then the detection of the common product CO_2 will be a

positive test for the utilization of other substrates (Scheme 1). For this reason, we decided

to use a headspace GC to measure the CO_2 evolved from reactions catalyzed by either

*Bf*GAD$_{WT}$ or engineered *Bf*GADs during the decarboxylation of various substrates. Fig. 3

shows data collected for *Bf*GAD$_{WT}$ and engineered *Bf*GADs with five different substrates

via headspace GC. Panels 3a, 3c, and 3e demonstrate chromatograms with the CO_2

peaks, visible immediately after 12.8 min, produced by the reactions of *Bf*GAD$_{WT}$,

*Bf*GAD$_{D104N}$, and *Bf*GAD$_{F81W}$ respectively with five different substrates. Whereas 3b, 3d,

and 3f depict CO_2 peak areas from the decarboxylation reactions of the substrates after

24 h incubation with either *Bf*GAD$_{WT}$ or engineered *Bf*GADs. Here, *Bf*GAD$_{WT}$ showed

significant production of CO_2 generated from the decarboxylation of both L-glutamate

(blue) and L-CSA (red) (Fig. 3a, 3b). This is the first indication to our knowledge of a gut

microbial GAD, specifically a GAD from *Bacteroides,* that can utilize L-CSA as a

substrate. In addition to L-CSA, *Bf*GAD$_{WT}$ was capable of decarboxylating L-HCA

(orange) and L-CA (green) to small extents as evidenced by the small CO_2 peaks (Fig.

3a) in the chromatogram and small peak areas for evolved CO_2 (Fig. 3b). Additionally,

*Bf*GAD$_{WT}$ was able to decarboxylate L-aspartate (purple) (Fig. 3a, 3b). For *Bf*GAD$_{WT}$, if

CO_2 production peak area with the native substrate L-glutamate is considered to be 100%

then we observed 54%, 3%, 6% and 20% of CO_2 production with L-CSA, L-CA, L-HCA,

and L-Asp respectively within 24 h which increased to 75% (L-CSA), 6% (L-CA), 7% (L-

HCA), and 28% (L-Asp) within 48 h compared to the native substrate L-glutamate

(Supplementary Fig. 9a).

For *Bf*GAD$_{D104N}$, CO_2 production is comparable to the *Bf*GAD$_{WT}$ while using native

substrate L-glutamate. Compared to the peak area of CO_2 production for L-glutamate

(native substrate), we observed around 16%, 11%, 3% and 7% CO_2 production from L-

CSA, L-CA, L-HCA, and L-aspartate respectively within 24 h (Fig. 3c, 3d) that increased

to 18% for both L-CSA and L-CA, 7% for L-HCA and slightly decreased with L-aspartate

within 48 h (Supplementary Fig. 9b). Based on the data collected with *Bf*GAD$_{WT}$ and

*Bf*GAD$_{D104N}$, residue Asp104 (D104) from the neighboring monomer is playing a crucial

role in accommodating various substrates. This engineered enzyme was also able to

decarboxylate L-CSA, L-HCA, and L-Asp in addition to the native substrate L-glutamate

but less efficiently than the BfGAD$_{WT}$, specifically the activity towards L-CSA was

significantly impacted. It is likely that the negative charge of the side chain carboxylate

from D104 is important for the specificity towards alternative substrates L-CSA, L-HCA

and L-Asp but not for the native substrate L-glutamate. As a result, when the charge was

removed due to the substitution of Asp (D) to Asn (N) at position 104, the catalysis with

these alternate substrates was affected, but not with the native substrate L-glutamate.

Interestingly, we noticed a 2-2.5-fold increase in the production of taurine with this evolved

enzyme compared to the BfGAD$_{WT}$ (Fig. 3d), indicating that the substitution from D to N

was favorable for taurine production.

The second engineered enzyme, BfGAD$_{F81W}$ retained its activity with the native

substrate L-glutamate as well but as observed in the initial velocity experiments (Fig. 2f),

it was slower compared to BfGAD$_{WT}$. If the CO_2 peak area of BfGAD$_{WT}$ or BfGAD$_{D104N}$

with the native substrate L-glutamate is considered to be 100%, BfGAD$_{F81W}$ showed the

peak area of around 93-94% with the same concentration of the substrate. Additionally,

there was minimal decarboxylation activity of BfGAD$_{F81W}$ with L-CA, L-HCA, and L-ASP.

However, this evolved enzyme still retained considerable decarboxylation activity with L-

CSA. It is interesting to note that a bulky substitution of tryptophan (W) at position 81,

instead of phenylalanine (F) (Fig. 2b, 2d), made the enzyme more specific towards the

native substrate L-glutamate. Based on the CO_2 evolution captured by our GC data, the

substrate preference for BfGAD$_{WT}$ was L-Glu > L-CSA > L-Asp > L-HCA > L-CA that

changed to L-Glu > L-CSA ≥ L-CA > L-Asp > L-HCA for BfGAD$_{D104N}$. Whereas for

*Bf*GAD$_{F81W}$, only L-glutamate and L-CSA showed decarboxylation and L-glutamate was

a much better substrate than L-CSA.

Wild-type and evolved *Bf*GADs produce multiple neuromodulatory molecules. The

generation of CO_2 in reactions facilitated by *Bf*GAD$_{WT}$ and its engineered variants

prompted us to detect and identify products resulting from decarboxylation with various

substrates. Both *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ produced decarboxylated products –

hypotaurine, taurine, homotaurine, and β-alanine from substrates L-CSA, L-CA, L-HCA,

and L-Asp respectively (Fig. 4, a-d, Supplementary Fig. 10, a-d). Intensity plots

generated from TLC plates showed that *Bf*GAD$_{WT}$ decarboxylated L-CSA and L-Asp to

produce hypotaurine and β-alanine more efficiently than *Bf*GAD$_{D104N}$ (Fig. 4a, 4c).

However, *Bf*GAD$_{D104N}$ was better at decarboxylating L-CA to taurine than *Bf*GAD$_{WT}$ (Fig.

4b). We do not observe much of decarboxylated product taurine with *Bf*GAD$_{WT}$ from L-

CA decarboxylation during a 24 h incubation, whereas *Bf*GAD$_{D104N}$ shows a significant

production of taurine during this time frame (Fig. 4b, Supplementary Fig. 10e). While our

CO_2 evolution experiments have shown that *Bf*GAD$_{WT}$ can catalyze L-CA decarboxylation

in 24 h, the failure to observe decarboxylated product on TLC may be attributed to the

low concentration of taurine produced, making it undetectable on a TLC plate. Both

enzymes *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ were able to catalyze the decarboxylation of L-HCA

to homotaurine to a very small extent (Fig. 4d). We also verified that the product from the

L-Asp decarboxylation was β-alanine and not L-alanine. Although, the R$_f$ (retention factor)

values were almost similar for β-alanine and L-alanine, the staining with the ninhydrin

differs for these molecules. β-alanine exhibited a purple color whereas L-alanine shows

a brick red color with ninhydrin stain (Supplementary Fig. 10c). We did not find any activity

of BfGAD$_{F81W}$ with L-CA, L-Asp, and L-HCA and were unable to detect taurine, β-alanine,

and homotaurine even after 48 h incubation (Supplementary Fig. 11a, 11b). These results

reinforce our observations from CO_2 evolution experiments. Additionally, despite the

detection of CO_2 during the decarboxylation activity of BfGAD$_{F81W}$ with L-CSA (Fig. 3f),

we could not observe detectable hypotaurine spots on the TLC plate (Supplementary Fig.

11a). We hypothesize that the amount of hypotaurine produced by BfGAD$_{F81W}$ may not

be sufficient to be detected via TLC, as a higher concentration of molecules (in mM range)

are needed for TLC analysis.

BfGAD$_{WT}$ and BfGAD$_{D104N}$ generate neuromodulatory molecules at differential

abundances, even when presented with a mixed substrate pool. In a complex gut

environment, the organism will encounter multiple substrates simultaneously. To

investigate how BfGAD$_{WT}$ functions in such a complex environment, we conducted

competition assays between various substrates. To this end, we tried to detect the

presence of decarboxylated products in mixtures containing the native substrate L-

glutamate with a specific alternate substrate. Our data with BfGAD$_{WT}$ indicate that when

the concentration ratio of the native substrate to the alternative substrate was 1:1, GABA

was predominantly seen as the primary decarboxylated product (Supplementary Fig. 12f).

However, when the concentration ratios were changed to 1:5 (L-Glu with L-CSA/L-Asp)

and 1:10 (L-Glu with L-CSA/L-CA/L-Asp), a gradual increase over time in the formation

of the alternative decarboxylated products - hypotaurine, taurine, and β-alanine was

observed (Fig. 5-7). For the BfGAD$_{WT}$ catalyzed decarboxylation reactions of L-CSA and

L-Asp, the resulting products hypotaurine and β-alanine could be observed when these

substrates were in 5-fold excess to the native substrate L-glutamate. However, the

accumulation of products hypotaurine and β-alanine becomes significant only after 9

hours (for β-alanine) – 24 h (for hypotaurine) (Supplementary Fig. 12d, 12e). In contrast,

when L-CSA and L-Asp are used in 10-fold excess of L-glutamate, the resulting products,

hypotaurine and β-alanine (peaks highlighted in pink) started accumulating significantly

much earlier around 3 hours (Fig. 5, 7). We did not observe the accumulation of taurine

even when 10-fold excess of L-CA was mixed with L-glutamate in the BfGAD$_{WT}$ catalyzed

reaction (Fig. 6). However, as mentioned earlier, we were able to detect CO_2 in the same

time frame (48 h) in the reaction catalyzed by BfGAD$_{WT}$ when L-CA was provided as the

sole substrate (Supplementary Fig. 9a). The inability to detect taurine might be because

of the low concentrations produced under the competitive environment of the mixed

substrate pool. Unlike BfGAD$_{WT,}$ BfGAD$_{D104N}$ was able to decarboxylate and accumulate

taurine (peaks highlighted in pink) to a small extent when L-CA is mixed with L-glutamate

at a 10-fold excess concentration, specially between 24-48 h (Fig. 6). Interestingly, in all

mixed substrate experiments, we observed maximum accumulation of the native product

GABA (peaks highlighted in blue) within 3 h. The intensity peaks showed no further

increase after 3 h time point. This shows that even in the presence of other substrates

the native activity of L-glutamate decarboxylation occurred at the highest velocity for all

BfGADs compared to the decarboxylation reactions of other substrates.

Additionally, the decarboxylated products hypotaurine and taurine were analyzed

using LC-ESI-MS/MS in addition to TLC to confirm their presence and mass in mixed

substrate assays where L-glutamate to L-CSA or L-CA ratios were 1:10. The qualitative

analysis of these molecules was conducted using MRM mode by running mixed

standards (Supplementary Fig. 13). Figure 8a depicts the presence of the product

hypotaurine and the remaining L-CSA substrate in the mixed substrate reactions with a

1:10 ratio of L-glutamate:L-CSA catalyzed by *Bf*GAD$_{WT}$. Figure 8b depicts the presence

of taurine and remaining L-CA in the mixed substrate reactions with a 1:10 ratio of L-

Glutamate:L-CA catalyzed by *Bf*GAD$_{D104N}$. We did not analyze the native substrate and

product, L-glutamate and GABA in these samples using LC-MS/MS because the

presence of these molecules was confirmed by both TLC and GABase assay prior to LC-

MS/MS analysis.

DISCUSSION

Prior research has implicated gut microbial contributions to the production of γ-

aminobutyric acid (GABA), specifically from *Bacteroides* genus[36,37]. GABA, a

neurotransmitter, is modulated in many neurodegenerative diseases, including

Alzheimer's and dementia, where lower GABA levels have been consistently reported[12].

Microbes from the genus *Bacteroides* are known to be fluctuated in individuals afflicted

by Alzheimer's and dementia[30-33]. Due to the prevalent nature of *Bacteroides* in the gut

and the modulation of Bacteroides during Alzheimer's, we decided to understand the

mechanism by which GABA is produced in these organisms. For this reason, an

annotated glutamate decarboxylase from *B. fragilis* (*Bf*GAD) was selected as a candidate

enzyme representing annotated GADs in all *Bacteroides*.

Most of the studied microbial GADs are oligomeric which include dimeric,

tetrameric or hexameric states[39,44,49,61-63] . It is known that the functional GAD unit is a

dimer as residues from both monomers make an intact active site and play an important

role during the catalysis. Therefore, an oligomeric composition that is not an even number

would be unusual in this context. Our gel filtration chromatography data with pH 4.7 buffer

gave a molecular weight of the protein that is higher than a dimer but lower than a

tetramer. We hypothesize that the protein likely exists in a dynamic equilibrium between

these two states, with a tendency towards a tetrameric state. We observed an

intermediate molecular weight possibly due to the inability to resolve dimeric and

tetrameric states completely. However, at pH 7.2, there might be a shift in this equilibrium

favoring a more dimeric state which is observed with a molecular weight closer to that of

a dimer than a tetramer. This shift in oligomeric states has been observed previously in

E. coli GAD that undergoes significant changes in oligomeric forms upon shift in the pH.

Thus, purified *Bf*GAD_WT behaves similarly in the oligomeric composition to other

characterized prokaryotic GADs. Our data suggests that there might be a dynamic

equilibrium between dimeric and tetrameric forms of *Bf*GAD_WT.

379   Unlike oligomeric conformations, the kinetic parameters of *Bf*GAD_WT vary from

previously investigated prokaryotic GADs[39,44,49,51,61,62]. We find that *Bf*GAD_WT is an

allosteric enzyme, displaying a positive cooperativity with the substrate L-glutamate. This

happens when binding of substrate on one site or monomer of the protein promotes the

binding of additional substrate to the other sites or monomers of the protein, where

instead of a hyperbolic dependence, a sigmoidal curve is observed for the plot of initial

velocity vs. substrate concentration. This phenomenon has not been seen with any other

prokaryotic GAD. The physiological consequences of this cooperativity in *Bf*GAD need to

be explored further. However, cooperative binding and adaptability in the allosteric

enzymes helps amplify the enzyme's response to changes in substrate concentration,

making it more sensitive to the physiological substrate concentrations and conditions.

Although the kinetic parameters are different for the *Bf*GAD$_{WT}$ compared to other

studied microbial GADs, there are prominent structural similarities among all these GADs.

The AlphaFold model of *Bf*GAD predicts active site architecture that is very similar to

other prokaryotic GADs. Particularly, catalytic residues in the active site are positioned

similar to those in *Ec*GAD and *Bt*GAD, suggesting that corresponding residues in *Bf*GAD

might maintain similar interactions with ligands. While the hydrogen bonding interactions

of the residues with the substrate analog glutarate are predicted to be within the expected

limits of H-bond distances, in the superimposed structures of *Ec*Gad and *Bf*GAD$_{WT}$, there

are differences in the hydrogen bond lengths when amino acid substitutions are made in

the active site. The changes in these H-bonding interactions could potentially be involved

in some capacity in accommodating or excluding alternative substrates in the active site

of the engineered *Bf*GADs to evolve new activity or specificity.

In contrast to GADs from *Bacteroides* sp., microbial GADs from various other genera

have been widely characterized[39,49,61,64,65]. Most of these studies show that many

annotated microbial glutamate decarboxylases prefer L-glutamate as a substrate.

However, substrate promiscuity has been seen in GADs from different domains of life.

For example, *E. coli* GAD was able to convert a phosphinic analog of glutamate to a

phosphinic analog of GABA[58]. In addition, human and other mammalian GADs are known

to catalyze decarboxylation of substrates other than L-glutamate. Specifically, sulfinic acid

and sulfonic acid derivatives of alanine (known as cysteine sulfinic acid/CSA and cysteic

acid/CA respectively) are known alternative substrates for mammalian GADs[57,59,66]. The

decarboxylation of these molecules by mammalian GADs generate hypotaruine and

taurine, from the respective sulfinic or sulfonic acids. Additionally, mammals have a

couple of other *de novo* pathways for the production of taurine and hypotaurine. The

primary pathway involves the enzyme cysteine sulfinic acid decarboxylase (CSAD) that

generates taurine and its intermediate hypotaurine by decarboxylating either L-cysteic

acid (CA) or L-cysteine sulfinic acid (CSA)[59,67]. In addition to hCSAD, mammalian GADL1

(glutamic acid decarboxylase like 1) enzyme that has very high sequence similarity with

hCSAD, is capable of producing taurine[55,68]. Taurine, a sulfur containing non-

proteinogenic amino acid whose production is controlled primarily by the enzyme cysteine

sulfinic acid decarboxylase (CSAD)[69,70]. However, except for a few marine microbes the

major *de novo* pathway for taurine formation with the help of CSAD has not been seen in

other prokaryotes[57]. Interestingly, in these microbes CSAD genes are present in the

operon containing cysteine dioxygenase (CDO) enzyme that catalyzes the conversion of

L-cysteine to L-cysteine sulfinic acid (L-CSA) which then can be converted to hypotaurine

and taurine (Supplementary Fig. 14). CDOs are not present in the members of the human

gut microbiome possibly due to the hypoxic and anaerobic conditions of the gut. Despite

an extensive bioinformatics search, we did not find any gut microbial genes annotated as

CSAD (cysteine sulfinic acid decarboxylase). Due to the lack of the annotated CSAD

enzymes responsible for the *de novo* taurine biosynthesis, we investigated the role of

*Bf*GAD in the formation of taurine and its derivatives in addition to GABA.

Apart from taurine, mammalian GADL1 is also able to generate β-alanine from L-

aspartate[55,68]. β-alanine is a precursor for the dipeptide beta-alanyl-L-histidine in humans,

commonly known as carnosine. Carnosine is found in muscles and brain tissues at high

concentrations[71,72]. In microbes, β-alanine is a precursor for coenzyme A (CoA)

biosynthesis, which is an important molecule in various metabolic pathways[73]. Both

carnosine and β-alanine show protective effects in individuals with cognitive deficits and

Alzhiermer's[74-76]. Moreover, higher serum concentrations of β-alanine prevent

dementia[74]. Considering the structural similarities between L- glutamate and L-aspartate

and the ability of some GADs to use L-aspartate as a substrate[53,56,77], we examined the

possible role of *Bf*GAD in the production of β-alanine via the decarboxylation of L-

aspartate. The neuromodulatory molecules, GABA, taurine and β-alanine generated by

glutamate decarboxylases or enzymes similar to GADs, have the ability to reverse

cognitive deficits in neurodegenerative disorders like dementia and

Alzheimer's[12,13,24,69,74,75,78,79]. This prompted us to understand the modulation of these

molecules by the members of the human gut microbiome.

Our results show that the *Bf*GAD$_{WT}$ is promiscuous and is able to decarboxylate four

additional substrates structurally similar to L-glutamate to produce hypotaurine, taurine,

homotaurine, and β-alanine. There are distinct enzymes in microbes such as aspartate1-

decarboxylase (A1DC) and aspartate 4-decarboxylase (A4DC) that catalyze

decarboxylations of L-aspartate to produce β-alanine[80,81] and L-alanine[82,83] respectively.

While the genome of *B. fragilis* has genes annotated for both these enzymes, it is

interesting to observe that the glutamate decarboxylase from *B. fragilis* still shows

decarboxylation activity towards L-aspartate to produce β-alanine. Additionally,

prokaryotic A1DC and eukaryotic A1DC show evolutionary divergence where the former

uses pyruvoyl cofactor whereas the latter uses PLP cofactor[84,85]. Thus, *Bf*GAD wild-type

and variants harbor an activity that is mostly seen in eukaryotic organisms.

To understand the factors that drive substrate specificity in *Bf*GAD$_{WT}$ and to evolve

enzymes that potentially have switched substrate preferences or specificity, we chose

two active site residues that are known to make interactions with bound ligands[39,40].

Headspace GC was used for the detection of evolved CO_2 which is the common

decarboxylation product for all enzyme catalyzed reactions (wild-type and engineered

enzymes). Absolute quantification of $CO_{2(g)}$ was challenging due to its high solubility at

acidic pH where we conducted our enzymatic reactions. In these conditions, the CO_2

produced through the reaction might still be in a soluble form as a dissolved CO_2. So,

headspace GC was utilized as a primary tool to identify alternate substrates by allowing

detection of CO_2 peak that served as a positive indication of enzymatic decarboxylation

of the substrate molecules. From the identification provided by the GC experiments,

subsequent analysis was carried out with TLC to detect corresponding decarboxylated

products from the multiple substrates. Our headspace GC data reinforces our TLC data

in most cases. Both engineered enzymes *Bf*GAD$_{D104N}$ and *Bf*GAD$_{F81W}$ retain preference

for L-glutamate as a substrate. However, their preferences for alternate substrates are

different than the *Bf*GAD$_{WT}$. Additionally, significant spectral perturbations are observed

with evolved GADs indicating that the local environment of the cofactor binding site is

changed due to those amino acid substitutions. With this, we show that there might be a

de novo pathway in gut microbes to make taurine and *Bf*GAD$_{WT}$ is able to produce multiple

neuromodulatory molecules.

Materials and Methods

Materials. Kanamycin, IPTG, GABase from *Pseudomonas fluorescens*, β-ME (beta-

mercaptoethanol), α-ketoglutarate, D-glutamic acid, L-cysteine sulfinic acid

monohydrate, L-cysteic acid monohydrate, GABA (γ-aminobutyric acid), hypotaurine,

taurine, L-aspartic acid sodium salt, β-alanine, HEPES, Imidazole, pyridoxal 5'-phosphate

monohydrate and TLC Silica gel 60 F$_{254}$ (20 cm x 20 cm) were purchased from Sigma-

Aldrich. Sodium L- glutamate monohydrate, Luria-Bertani Broth (LB), buffer components,

Sodium chloride, Sodium acetate, ninhydrin and NADP$^+$ disodium salt were purchased

from Fisher Scientific. All restriction enzymes and competent cells of *E. coli* BL21(DE3)

and *E. coli* NEB5α were purchased from New England Biolabs (NEB). Headspace vials

were purchased from Chemglass Inc.

Experimental procedures

Gene cloning. A synthesized gene for *B. fragilis* glutamate decarboxylase (*Bf*GAD) was

created with the help of Genewiz from Azenta life sciences. The gene was codon

optimized for expression in *E. coli* cells, which was then subcloned into a pET28a

expression vector between the NdeI and HindIII restriction sites to incorporate an N-

terminal hexahistidine (His$_6$) Tag. The pET28a-*Bf*GAD construct was confirmed by

agarose gel electrophoresis, restriction digestion, sanger sequencing (Genewiz), and full

plasmid sequencing (Plasmidsaurus) and then transformed into *E. coli* NEB5α and *E. coli*

BL21(DE3) competent cells by heat shock. Glycerol stocks of the cells harboring the

construct pET28a-*Bf*GAD were stored at –80 ⁰C.

Expression and purification of *Bf*GAD$_{WT}$ and engineered *Bf*GADs. *E. coli* BL21(DE3)

cells containing the pET28a-*Bf*GADs were cultivated in LB medium containing kanamycin

(50 µg/mL) with shaking (200 rpm) at 37 ⁰C until OD$_{600}$ reached 0.7. At this point, the

*Bf*GAD expression was induced with isopropyl β-D-1-thiogalactopyranoside (IPTG) to a

final concentration of 1 mM and growth was continued at 37 ⁰C for 3 hours to allow for

protein expression with continuous shaking at 200 rpm. Cells were harvested via

centrifugation at 4000*g* for 30 min at 4 ⁰C and cell pellets were stored at –80 ⁰C. The

frozen cell pellets were thawed on ice and resuspended in buffer A containing 20 mM

Tris-HCl, 500 mM NaCl, 40 mM imidazole, 0.1 mM PLP, pH 7.8. All subsequent

purification steps were carried out at 4 °C. The cells were disrupted by sonication (20000

Hz for 10 cycles of 30 seconds each altering 1 minute on ice) and centrifuged at 13500g

at 4 °C for 30 min to separate supernatant from cell debris. The supernatant was then

filtered through 0.2 µm PES filter membrane and loaded onto a HisTrap HP column (5 ml,

1.6 x 2.5 cm, Ni Sepharose High-performance column, GE Healthcare, now Cytiva) pre-

equilibrated with buffer A at 1 mL/min. The wash step was carried out by running 5-column

volumes of lysis buffer A to elute contaminating proteins. *Bf*GAD was eluted with a linear

gradient from buffer A (containing 40 mM imidazole) to an elution buffer B containing 20

mM Tris-HCl, 500 mM NaCl, 400 mM imidazole, 0.1 mM PLP, pH 7.8 at 2 mL/min. Eluted

protein peaks were pooled, buffer exchanged and concentrated to a final buffer (50 mM

HEPES, pH 7.2) using Vivaspin 50 kDa MWCO filters (Cytiva). Aliquots of purified

proteins were stored at –80 °C. Engineered *Bf*GADs were purified similarly. SDS-PAGE

analysis and activity assays were performed to confirm the purity and presence of

functional *Bf*GAD proteins.

Gel filtration chromatography with *Bf*GAD$_{WT}$. To understand oligomeric state of

*Bf*GAD$_{WT}$, gel filtration chromatography was performed at two different pH, 4.7 (50 mM

sodium acetate, 150 mM NaCl) and pH 7.2 (50 mM HEPES, 150 mM NaCl) using

HiPrep™ 16/60 Sephacryl® S-200 HR column. Equilibration and elution steps were

carried out at a flow rate of 0.5 mL/min. High molecular weight (HMW) and Low molecular

weight (LMW) calibration kits (Cytiva) were used for column calibration, to create a

standard curve, and for molecular weight determination (Supplementary Fig. 5b, 5c).

Activity assays. *Bf*GAD activity assays were performed by a coupled enzyme assay with

a GABase system to measure GABA production spectrophotometrically. Briefly, *Bf*GAD

was incubated with 50 mM L-glutamate in 50 mM sodium acetate, pH 4.7 buffer or in

buffers at other pH values for one hour at 25 ⁰C. Enzymatic reactions were stopped by

boiling samples for 15 min. These samples were then centrifuged at 6000*g* for 5 min and

if necessary, dilutions were created in 50 mM Tris-HCl, pH 8.6. For GABA measurement,

75 µL of samples (either diluted or undiluted) were added to 25 µL of Gabase assay mix

containing 10 mM BME, 2 mM α-ketoglutarate, 600 µM NADP⁺, and 30 µg (0.015U/mL)

of GABase in 50 mM Tris-HCl, pH 8.6[47,86-89]. In the GABase assay, GABA is converted to

succinic semialdehyde (SSA) and then to succinate with subsequent production of

NADPH which was measured at 340 nm. Using an extinction coefficient of 6,220 M^{-1} cm⁻

^{1} at 340 nm, NADPH concentrations were calculated which will provide GABA

concentrations in measured samples ([NADPH]=[GABA]).

*Bf*GAD **absorbance spectrophotometry and activity assays with pH variation.**

Spectra of *Bf*GAD$_{WT}$ were recorded from 200 nm - 800 nm in buffers at pH 4.7, 5.5, 6.7,

7.5, and 8.6 at 25 ⁰C on Agilent Cary 3500 UV-Vis spectrophotometer. From these

spectra, absorbance changes at 335 nm (enolimine) and 420 nm (ketoenamine) for

enzyme bound PLP cofactor were collected. A plot of pH versus absorbance at 420 nm

was generated and the curve fitting was carried out using equation (1)[39].

$\frac{AbsHnE - AbsE}{Abs - AbsE} - 1 = \frac{[10^{-npK}]}{[10^{-npH}]}$ (1)

Where pK is the acid dissociation constant, number of protons involved in the titration is

n and, AbsHnE and AbsE are the absorbances at acidic and basic pH which produce

protonated and deprotonated forms of the enzymes. The initial activity assays for the

*Bf*GAD$_{WT}$ were performed in buffers at pH 4.7 - 8 with GABase assay system as

mentioned in the above method. Based on the results of these activity assays, all

subsequent enzymatic assays were carried out at pH 4.7.

Steady-state kinetics. Kinetic parameters of *Bf*GAD$_{WT}$ were determined by varying

*Bf*GAD concentrations (0.5 µM-2 µM) and by varying L-glutamate concentrations (0.125

mM-32 mM) in 50 mM sodium acetate, pH 4.7 at 25º C. For enzyme concentration

variation experiments, reaction samples with each enzyme concentration were collected

at 1 min time point after starting the reaction and then stopped by boiling for 15 min. Once

cooled to room temperature, these samples were analyzed for GABA content with

GABase assay as mentioned above. For the substrate concentration variation

experiments, 100 µL aliquots of reactions with each substrate concentration were

collected at different time intervals (0.2 – 30 min). The reactions were stopped by boiling

the samples for 15 minutes. Once cooled to room temperature, these samples were

analyzed for GABA content with GABase assay as mentioned above. The initial velocities

of various reactions were calculated by fitting data of early timepoints to a linear

regression. These initial velocities were then plotted against substrate concentrations and

curve fitting was carried out using equation (2).

$$v = \frac{V_{max}[S]^h}{\left(k_{half}\right)^h + [S]^h}$$ (2)

Where v is the initial velocity, V$_{max}$ is the maximum velocity, K$_{half}$ is the concentration of

substrate at which the reaction velocity is half of V$_{max}$, [S] is the substrate concentration,

h is the hill coefficient.

Generation of engineered *Bf*GAD variants. The pET28a-*Bf*GAD was used as a

template for the generation of engineered *Bf*GAD variants. *Bf*GAD$_{D104N}$ and *Bf*GAD$_{F81W}$

were created by site-directed mutagenesis with Phusion DNA polymerase using the

following primer pairs incorporating codons for the specific amino acid substitution

(underlined).

For BfGAD$_{D104N}$

F: 5' CCGCAATGCGCGGATATTCGGTTTC<u>ATT</u>AATATAGTTAATG 3'

R: 5' CATTAACTATATT<u>AAT</u>GAAACCGAATATCCGCGCATTGCGG 3'

For BfGAD$_{F81W}$

F: 5' CATCCATATAGGTGGTCACCCAGGTCG<u>CCA</u>GGTTCAGGCGCG 3'

R: 5' CGCGCCTGAACCTGGCGACC<u>TGG</u>GTGACCACCTATATGGATG 3'

The variants were confirmed by gene sequencing (Genewiz) and full plasmid sequencing

with Plasmidsaurus.

CO$_{2(g)}$ detection by headspace gas chromatography. All enzymatic reactions and

controls were prepared in 6 mL vials and sealed prior to starting the reaction. The

reactions were started by the addition of BfGAD (WT or engineered variants) with a

syringe into the reaction mixture containing 50 mM substrate in 50 mM sodium acetate,

pH 4.7. These vials were incubated at 37 ºC for 24 hours and 48 hours. Headspace GC

measurements were carried out on an Agilent 8890 gas chromatograph system equipped

with a flame ionization detector (FID) and a Hayesep Q packed column (1.8 m x 2 mm x

3.17 mm) (Agilent) operating with an Argon carrier gas (flow rate= 5 mL/min). The oven

was programed to hold 30 °C for 6.5 min, ramp at 30 °C/min to 280 °C with a hold for 4

min for a total run time of 18.83 min. The flame ionization detector was used for the

detection of CO_2 gas with a temperature setting of 275 °C, hydrogen flow of 60 mL/min,

air flow of 400 mL/min and constant makeup gas (nitrogen) at 5 mL/min. The retention

time for CO_2 gas was 12.83 min. Multiple standards of CO_2 gas were analyzed by this

method to confirm retention time before injecting reaction samples (Supplementary Fig.

15).

Thin layer chromatography (TLC). Silica gel plates (stationary phase) with a solvent

system (mobile phase) of 3:1:1 ratio of butanol: acetic acid: H_2O were used to separate

reaction products from substrates[60]. 2 μL of reaction mixtures were spotted on glass silica

plates along with 10 mM of various metabolites (L-glutamate, D-glutamate, L-CSA, L-CA,

L-HCA, L-Aspartate, GABA, hypotaurine, taurine, homotaurine and β-alanine) as

standards. Separation via TLC was carried out at 25 °C for 3-4 hours in an above-

mentioned mobile phase. Once mobile phase reached a sufficient height on the TLC

plate, chromatographic separation was discontinued. Plates were treated with 0.5%

ninhydrin in acetone (w/v) and heated minimally with a dryer for the color development.

The TLC plate images were initially inverted via ImageJ and then intensity of each spot

was quantified by a custom python code.

Liquid chromatography with tandem mass spectrometry (LC-MS/MS) detection of

decarboxylated products from reactions catalyzed by *Bf*GADs (WT and engineered

variants). Samples were analyzed by ESI-LC-MS/MS in positive (products) and negative

ion (substrates) mode using a Thermo vanquish LC and TSQ Altis Triple Quadrupole

Mass Spectrometer. Samples were separated by gradient elution using Agilent Infinity lab

Poroshell 120 HILIC-Z 2.1 x 100 mm, 2.7 μm column at 25 °C with 20 mM ammonium

formate, pH 3.0 (Solvent A) and 20 mM ammonium formate in 9:1 acetonitrile: H_2O, pH

3.0 (Solvent B). Elution was initiated at 70% B for 11.5 min, followed by gradient elution

from 70 to 100% B over 11.5 min, and 100% B for 4 min at a flow rate of 0.5mL/min. The

precursor to product transitions of *m/z* [M+H]⁻152→ 88.1(Cysteine sulfinic acid), [M+H]⁻

168→ 81.1(Cysteic acid), [M+H]⁺110.1→ 45.1(hypotaurine), [M+H]⁺126.1→ 44.1(taurine)

were employed to monitor substrates and products.

Multiple Sequence alignment and phylogenetic analyses of gut microbial GADs. All

the protein sequences were obtained from NCBI. Multiple sequence alignment was

performed using Clustal Omega (1.2.4) tool with default parameters[90,91] to understand

sequence similarities among various gut microbial glutamate decarboxylases (GADs).

The phylogenetic tree was created for *Bacteroides* GADs using cyberinfrastructure for

phylogenetic research (CIPRES)[92]. Within CIPRES, the MAFFT on XSEDE (7.505)[93] was

utilized to create a separate multiple sequence alignment that goes through FastTreeMP

on XSEDE(2.1.10)[94,95] to obtain the phylogenetic tree file which can be visualized and

annotated with Interactive tree of life (iTOL v6.8.1)[96].

***Bf*GAD dimer generation with the AlphaFold2 and visualization, structural**

alignments, and mutagenesis via PyMOL. The dimer structure of wildtype *Bf*GAD was

generated using AlphaFold2 and visualized using PyMOL. Variants were then created

using the mutagenesis wizard in PyMOL. Rotamers for the altered residues were selected

based on their best fit to the position of wild-type residues in the *Bf*GAD protein. All

structural alignments were carried out using PyMOL align method, involving five iteration

cycles and a cutoff of 2 Å.

Acknowledgments. This research is supported by the seed grant from Edson initiative

for the dementia care and solutions to DDS. We acknowledge resources and support from

the Knowledge Enterprise Biosciences Core Facilities at Arizona State University. We

thank Dr. Rosa Krajmalnik-Brown and Christopher Muse from the Biodesign Centre for

Health Through Microbiomes for the help with headspace GC. This research is supported

by the seed grant from Edson initiative for the dementia care and solutions to DDS. AS

was supported by an NIH-NIGMS MIRA award R35GM147131.

Author Contributions. PD performed majority of the experiments and data analysis.

CWP performed some of the TLC experiments. AS performed analysis and data fitting for

pH and Abs variation experiment and provided code and advice on the image analysis of

TLC plates, PD and DDS conceptualized the project and wrote the paper.

Data availability. All the data are available in the manuscript or with associated

supporting information. The python code used for TLC image analysis is available at

https://github.com/Pavani-dadi/TLC_profile.

Declaration of interests. Aspects of this research are part of a pending patent

application.

References

1   Murley, A. G. *et al.* GABA and glutamate deficits from frontotemporal lobar degeneration
are associated with disinhibition. *Brain* **143**, 3449-3462 (2020).
https://doi.org/10.1093/brain/awaa305
2   Nuss, P. Anxiety disorders and GABA neurotransmission: a disturbance of modulation.
Neuropsychiatr Dis Treat **11**, 165-175 (2015). https://doi.org/10.2147/NDT.S58841
3   Cawley, N. *et al.* Reduced gamma-aminobutyric acid concentration is associated with
physical disability in progressive multiple sclerosis. *Brain* **138**, 2584-2595 (2015).
https://doi.org/10.1093/brain/awv209
4   Goddard, A. W. *et al.* Reductions in occipital cortex GABA levels in panic disorder detected
with 1h-magnetic resonance spectroscopy. *Arch Gen Psychiatry* **58**, 556-561 (2001).
https://doi.org/10.1001/archpsyc.58.6.556
5   Goddard, A. W. *et al.* Impaired GABA neuronal response to acute benzodiazepine
administration in panic disorder. *Am J Psychiatry* **161**, 2186-2193 (2004).
https://doi.org/10.1176/appi.ajp.161.12.2186
6   Long, Z. *et al.* Decreased GABA levels in anterior cingulate cortex/medial prefrontal cortex
in panic disorder. *Prog Neuropsychopharmacol Biol Psychiatry* **44**, 131-135 (2013).
https://doi.org/10.1016/j.pnpbp.2013.01.020
7   Orhan, F. *et al.* CSF GABA is reduced in first-episode psychosis and associates to symptom
severity. *Mol Psychiatry* **23**, 1244-1250 (2018). https://doi.org/10.1038/mp.2017.25
8   Sarlo, G. L. & Holton, K. F. Brain concentrations of glutamate and GABA in human epilepsy:
A review. *Seizure* **91**, 213-227 (2021). https://doi.org/10.1016/j.seizure.2021.06.028
9   Umesawa, Y. *et al.* Altered GABA Concentration in Brain Motor Area Is Associated with
the Severity of Motor Disabilities in Individuals with Autism Spectrum Disorder. *J Autism
Dev Disord* **50**, 2710-2722 (2020). https://doi.org/10.1007/s10803-020-04382-x
10   Porges, E. C. *et al.* Frontal Gamma-Aminobutyric Acid Concentrations Are Associated With
Cognitive Performance in Older Adults. *Biol Psychiatry Cogn Neurosci Neuroimaging* **2**, 38-
44 (2017). https://doi.org/10.1016/j.bpsc.2016.06.004
11   Carello-Collar, G. *et al.* The GABAergic system in Alzheimer's disease: a systematic review
with meta-analysis. *Mol Psychiatry* (2023). https://doi.org/10.1038/s41380-023-02140-w
12   Solas, M., Puerta, E. & Ramirez, M. Treatment Options in Alzheimer´s Disease: The GABA
Story. *Current Pharmaceutical Design* **21**, 4960-4971 (2015).
https://doi.org/10.2174/1381612821666150914121149
13   Jia, F. *et al.* Taurine is a potent activator of extrasynaptic GABA(A) receptors in the
thalamus. *J Neurosci* **28**, 106-115 (2008). https://doi.org/10.1523/JNEUROSCI.3996-
07.2008
14   Ochoa-de la Paz, L., Zenteno, E., Gulias-Cañizo, R. & Quiroz-Mercado, H. Taurine and GABA
neurotransmitter receptors, a relationship with therapeutic potential? *Expert Review of
Neurotherapeutics* **19**, 289-291 (2019). https://doi.org/10.1080/14737175.2019.1593827
15   Alom, J., Mahy, J. N., Brandi, N. & Tolosa, E. Cerebrospinal fluid taurine in Alzheimer's
disease. *Ann Neurol* **30**, 735 (1991). https://doi.org/10.1002/ana.410300518

16   Aytan, N. *et al.* Fingolimod modulates multiple neuroinflammatory markers in a mouse
model of Alzheimer's disease. *Sci Rep* **6**, 24939 (2016).
https://doi.org/10.1038/srep24939

17   Chiquita, S. *et al.* A longitudinal multimodal in vivo molecular imaging study of the 3xTg-
AD mouse model shows progressive early hippocampal and taurine loss. *Hum Mol Genet*
28, 2174-2188 (2019). https://doi.org/10.1093/hmg/ddz045

18   Garcia-Serrano, A. M., Vieira, J. P. P., Fleischhart, V. & Duarte, J. M. N. Taurine and N-
acetylcysteine treatments prevent memory impairment and metabolite profile
alterations in the hippocampus of high-fat diet-fed female mice. *Nutr Neurosci* **26**, 1090-
1102 (2023). https://doi.org/10.1080/1028415X.2022.2131062

19   Pomara, N. *et al.* Glutamate and other CSF amino acids in Alzheimer's disease. *Am J*
Psychiatry **149**, 251-254 (1992). https://doi.org/10.1176/ajp.149.2.251

20   Vermeiren, Y. *et al.* Behavioral correlates of cerebrospinal fluid amino acid and biogenic
amine neurotransmitter alterations in dementia. *Alzheimers Dement* **9**, 488-498 (2013).
https://doi.org/10.1016/j.jalz.2012.06.010

21   Csernansky, J. G., Bardgett, M. E., Sheline, Y. I., Morris, J. C. & Olney, J. W. CSF excitatory
amino acids and severity of illness in Alzheimer's disease. *Neurology* **46**, 1715-1720
(1996). https://doi.org/10.1212/wnl.46.6.1715

22   Gao, R., Bae, M. A., Chang, K. J. & Kim, S. H. Comparison of Urinary Excretion of Taurine
Between Elderly with Dementia and Normal Elderly. *Adv Exp Med Biol* **975 Pt 1**, 57-65
(2017). https://doi.org/10.1007/978-94-024-1079-2_5

23   Fujii, Y. *et al.* Fecal metabolite of a gnotobiotic mouse transplanted with gut microbiota
from a patient with Alzheimer's disease. *Biosci Biotechnol Biochem* **83**, 2144-2152 (2019).
https://doi.org/10.1080/09168451.2019.1644149

24   Jang, H. *et al.* Taurine Directly Binds to Oligomeric Amyloid-beta and Recovers Cognitive
Deficits in Alzheimer Model Mice. *Adv Exp Med Biol* **975 Pt 1**, 233-241 (2017).
https://doi.org/10.1007/978-94-024-1079-2_21

25   Kim, H. Y. *et al.* Taurine in drinking water recovers learning and memory in the adult
APP/PS1 mouse model of Alzheimer's disease. *Sci Rep* **4**, 7467 (2014).
https://doi.org/10.1038/srep07467

26   Manzano, S., Aguera, L., Aguilar, M. & Olazaran, J. A Review on Tramiprosate
(Homotaurine) in Alzheimer's Disease and Other Neurocognitive Disorders. *Front Neurol*
11, 614 (2020). https://doi.org/10.3389/fneur.2020.00614

27   Singh, P. *et al.* Taurine deficiency as a driver of aging. *Science* **380**, eabn9257 (2023).
https://doi.org/10.1126/science.abn9257

28   Borsom, E. M., Lee, K. & Cope, E. K. Do the Bugs in Your Gut Eat Your Memories?
Relationship between Gut Microbiota and Alzheimer's Disease. *Brain Sci* **10** (2020).
https://doi.org/10.3390/brainsci10110814

29   Saji, N. *et al.* Analysis of the relationship between the gut microbiome and dementia: a
cross-sectional study conducted in Japan. *Sci Rep* **9**, 1008 (2019).
https://doi.org/10.1038/s41598-018-38218-7

30   Zhuang, Z.-Q. *et al.* Gut Microbiota is Altered in Patients with Alzheimer's Disease. *Journal*
of Alzheimer's Disease **63**, 1337-1346 (2018). https://doi.org/10.3233/JAD-180176

31 Guo, M. *et al.* Gut Microbiome Features of Chinese Patients Newly Diagnosed with
Alzheimer's Disease or Mild Cognitive Impairment. *Journal of Alzheimer's Disease* **80**, 299-
310 (2021). https://doi.org/10.3233/JAD-201040

32 Li, B. *et al.* Mild cognitive impairment has similar alterations as Alzheimer's disease in gut
microbiota. *Alzheimer's & Dementia* **15**, 1357-1366 (2019).
https://doi.org/10.1016/j.jalz.2019.07.002

33 Jemimah, S., Chabib, C. M. M., Hadjileontiadis, L. & AlShehhi, A. Gut microbiome dysbiosis
in Alzheimer's disease and mild cognitive impairment: A systematic review and meta-
analysis. *PLOS ONE* **18**, e0285346 (2023). https://doi.org/10.1371/journal.pone.0285346

34 Salyers, A. A. Bacteroides of the human lower intestinal tract. *Annu Rev Microbiol* **38**, 293-
313 (1984). https://doi.org/10.1146/annurev.mi.38.100184.001453

35 Wexler, H. M. Bacteroides: the good, the bad, and the nitty-gritty. *Clin Microbiol Rev* **20**,
593-621 (2007). https://doi.org/10.1128/CMR.00008-07

36 Strandwitz, P. *et al.* GABA-modulating bacteria of the human gut microbiota. *Nat*
Microbiol **4**, 396-403 (2019). https://doi.org/10.1038/s41564-018-0307-3

37 Otaru, N. *et al.* GABA Production by Human Intestinal Bacteroides spp.: Prevalence,
Regulation, and Role in Acid Stress Tolerance. *Frontiers in Microbiology* **12**, 656895
(2021). https://doi.org/10.3389/fmicb.2021.656895

38 Horvath, T. D. *et al.* Bacteroides ovatus colonization influences the abundance of
intestinal short chain fatty acids and neurotransmitters. *iScience* **25**, 104158 (2022).
https://doi.org/10.1016/j.isci.2022.104158

39 Capitani, G. *et al.* Crystal structure and functional analysis of *Escherichia coli*
glutamate decarboxylase. *The EMBO Journal* **22**, 4027-4037-4037 (2003).
https://doi.org/https://doi.org/10.1093/emboj/cdg403

40 Dutyshev, D. I. *et al.* Structure of *Escherichia coli* glutamate decarboxylase (GADα)
in complex with glutarate at 2.05 Å resolution. *Acta Crystallographica Section D Biological*
Crystallography **61**, 230-235 (2005). https://doi.org/10.1107/S0907444904032147

41 Beattie, A. E. *et al.* The Pyridoxal 5'-Phosphate (PLP)-Dependent Enzyme Serine
Palmitoyltransferase (SPT): Effects of the Small Subunits and Insights from Bacterial
Mimics of Human hLCB2a HSAN1 Mutations. *BioMed Research International* **2013**, 1-13
(2013). https://doi.org/10.1155/2013/194371

42 Moore, P. S., Dominici, P. & Borri Voltattorni, C. Cloning and expression of pig kidney dopa
decarboxylase: comparison of the naturally occurring and recombinant enzymes.
Biochemical Journal **315**, 249-256 (1996). https://doi.org/10.1042/bj3150249

43 Zhou, X. & Toney, M. D. pH Studies on the Mechanism of the Pyridoxal Phosphate-
Dependent Dialkylglycine Decarboxylase. *Biochemistry* **38**, 311-320 (1999).
https://doi.org/10.1021/bi981455s

44 Liu, S. *et al.* Coordinated regulation of Bacteroides thetaiotaomicron glutamate
decarboxylase activity by multiple elements under different pH. *Food Chem* **403**, 134436
(2023). https://doi.org/10.1016/j.foodchem.2022.134436

45 Pennacchietti, E. *et al.* Mutation of His465 Alters the pH-dependent Spectroscopic
Properties of Escherichia coli Glutamate Decarboxylase and Broadens the Range of Its
Activity toward More Alkaline pH. *Journal of Biological Chemistry* **284**, 31587-31596
(2009). https://doi.org/10.1074/jbc.M109.049577

46   Shukuya, R. & Schwert, G. W. Glutamic acid decarboxylase. II. The spectrum of the
enzyme. *J Biol Chem* **235**, 1653-1657 (1960).
47   Tramonti, A., John, R. A., Bossa, F. & De Biase, D. Contribution of Lys276 to the
conformational flexibility of the active site of glutamate decarboxylase from
<i>Escherichia coli</i>: Role of Lys276 in <i>E. coli</i> glutamate decarboxylase.
European Journal of Biochemistry **269**, 4913-4920 (2002).
https://doi.org/10.1046/j.1432-1033.2002.03149.x
48   De Biase, D. & Pennacchietti, E. Glutamate decarboxylase-dependent acid resistance in
orally acquired bacteria: function, distribution and biomedical implications of the gadBC
operon. *Mol Microbiol* **86**, 770-786 (2012). https://doi.org/10.1111/mmi.12020
49   Chang, C. *et al.* Purification and characterization of glutamate decarboxylase from
Enterococcus raffinosus TCCC11660. *J Ind Microbiol Biotechnol* **44**, 817-824 (2017).
https://doi.org/10.1007/s10295-017-1906-3
50   Fonda, M. L. Glutamate decarboxylase. Substrate specificity and inhibition by carboxylic
acids. *Biochemistry* **11**, 1304-1309 (1972). https://doi.org/10.1021/bi00757a029
51   Ueno, Y., Hayakawa, K., Takahashi, S. & Oda, K. Purification and characterization of
glutamate decarboxylase from Lactobacillus brevis IFO 12005. *Biosci Biotechnol Biochem*
61, 1168-1171 (1997). https://doi.org/10.1271/bbb.61.1168
52   Fonda, M. L. L-Glutamate decarboxylase from bacteria. *Methods Enzymol* **113**, 11-16
(1985). https://doi.org/10.1016/s0076-6879(85)13005-3
53   Kim, H. W., Kashima, Y., Ishikawa, K. & Yamano, N. Purification and characterization of the
first archaeal glutamate decarboxylase from Pyrococcus horikoshii. *Biosci Biotechnol*
Biochem **73**, 224-227 (2009). https://doi.org/10.1271/bbb.80583
54   Tomita, H., Yokooji, Y., Ishibashi, T., Imanaka, T. & Atomi, H. An archaeal glutamate
decarboxylase homolog functions as an aspartate decarboxylase and is involved in beta-
alanine and coenzyme A biosynthesis. *J Bacteriol* **196**, 1222-1230 (2014).
https://doi.org/10.1128/JB.01327-13
55   Liu, P. *et al.* Role of glutamate decarboxylase-like protein 1 (GADL1) in taurine
biosynthesis. *J Biol Chem* **287**, 40898-40906 (2012).
https://doi.org/10.1074/jbc.M112.393728
56   Richardson, G. *et al.* An examination of aspartate decarboxylase and glutamate
decarboxylase activity in mosquitoes. *Mol Biol Rep* **37**, 3199-3205 (2010).
https://doi.org/10.1007/s11033-009-9902-y
57   Agnello, G., Chang, L. L., Lamb, C. M., Georgiou, G. & Stone, E. M. Discovery of a Substrate
Selectivity Motif in Amino Acid Decarboxylases Unveils a Taurine Biosynthesis Pathway in
Prokaryotes. *ACS Chemical Biology* **8**, 2264-2271 (2013).
https://doi.org/10.1021/cb400335k
58   Daniela De Biase, F. C., Eugenia Pennacchietti, Fabio Giovannercole, Antonio Coluccia,
Jouko Vepsäläinen & Alex Khomutov Enzymatic kinetic resolution of
desmethylphosphinothricin indicates that phosphinic group is a bioisostere of carboxyl
group. *Commnications chemistry* (2020).
59   Wu, J. Y. Purification and characterization of cysteic acid and cysteine sulfinic acid
decarboxylase and L-glutamate decarboxylase from bovine brain. *Proc Natl Acad Sci U S*
A **79**, 4270-4274 (1982). https://doi.org/10.1073/pnas.79.14.4270

60   Tramonti, A. *et al.* A Novel, Easy Assay Method for Human Cysteine Sulfinic Acid
Decarboxylase. *Life* **11**, 438 (2021). https://doi.org/10.3390/life11050438

61   Komatsuzaki, N., Nakamura, T., Kimura, T. & Shima, J. Characterization of glutamate
decarboxylase from a high gamma-aminobutyric acid (GABA)-producer, Lactobacillus
paracasei. *Biosci Biotechnol Biochem* **72**, 278-285 (2008).
https://doi.org/10.1271/bbb.70163

62   Hiraga, K., Ueno, Y. & Oda, K. Glutamate decarboxylase from Lactobacillus brevis:
activation by ammonium sulfate. *Biosci Biotechnol Biochem* **72**, 1299-1306 (2008).
https://doi.org/10.1271/bbb.70782

63   Giovannercole, F. *et al.* On the effect of alkaline pH and cofactor availability in the
conformational and oligomeric state of Escherichia coli glutamate decarboxylase. *Protein*
Eng Des Sel **30**, 235-244 (2017). https://doi.org/10.1093/protein/gzw076

64   Lee, J. Y. & Jeon, S. J. Characterization and immobilization on nickel-chelated Sepharose
of a glutamate decarboxylase A from Lactobacillus brevis BH2 and its application for
production of GABA. *Biosci Biotechnol Biochem* **78**, 1656-1661 (2014).
https://doi.org/10.1080/09168451.2014.936347

65   Nomura, M. *et al.* Lactococcus lactis contains only one glutamate decarboxylase gene.
Microbiology (Reading) **145 (Pt 6)**, 1375-1380 (1999).
https://doi.org/10.1099/13500872-145-6-1375

66   Blindermann, J. M., Maitre, M., Ossola, L. & Mandel, P. Purification and some properties
of L-glutamate decarboxylase from human brain. *Eur J Biochem* **86**, 143-152 (1978).
https://doi.org/10.1111/j.1432-1033.1978.tb12293.x

67   Winge, I. *et al.* Mammalian CSAD and GADL1 have distinct biochemical properties and
patterns of brain expression. *Neurochemistry International* **90**, 173-184 (2015).
https://doi.org/https://doi.org/10.1016/j.neuint.2015.08.013

68   Mahootchi, E. *et al.* GADL1 is a multifunctional decarboxylase with tissue-specific roles in
beta-alanine and carnosine production. *Sci Adv* **6**, eabb3713 (2020).
https://doi.org/10.1126/sciadv.abb3713

69   Ripps, H. & Shen, W. Review: taurine: a "very essential" amino acid. *Molecular Vision* **18**,
2673-2686 (2012).

70   De La Rosa, J. & Stipanuk, M. H. Evidence for a rate-limiting role of cysteinesulfinate
decarboxylase activity in taurine biosynthesis in vivo. *Comparative Biochemistry and*
Physiology Part B: Comparative Biochemistry **81**, 565-571 (1985).
https://doi.org/10.1016/0305-0491(85)90367-0

71   Boldyrev, A. A., Aldini, G. & Derave, W. Physiology and pathophysiology of carnosine.
Physiol Rev **93**, 1803-1845 (2013). https://doi.org/10.1152/physrev.00039.2012

72   Sale, C., Saunders, B. & Harris, R. C. Effect of beta-alanine supplementation on muscle
carnosine concentrations and exercise performance. *Amino Acids* **39**, 321-333 (2010).
https://doi.org/10.1007/s00726-009-0443-4

73   Lopez-Samano, M. *et al.* A novel way to synthesize pantothenate in bacteria involves beta-
alanine synthase present in uracil degradation pathway. *Microbiologyopen* **9**, e1006
(2020). https://doi.org/10.1002/mbo3.1006

74   Hata, J. *et al.* Association Between Serum beta-Alanine and Risk of Dementia. *Am J*
Epidemiol **188**, 1637-1645 (2019). https://doi.org/10.1093/aje/kwz116

75 Ostfeld, I. *et al.* Role of β-Alanine Supplementation on Cognitive Function, Mood,
and Physical Function in Older Adults; Double-Blind Randomized Controlled Study.
Nutrients **15** (2023).

76 Dhingra, D., Parle, M. & Kulkarni, S. K. β - Alanine protects mice from memory deficits
induced by ageing, scopolamine, diazepam and ethanol. *Indian Journal of Pharmaceutical*
Sciences **68** (2006). https://doi.org/10.4103/0250-474X.25718

77 Wu, J.-Y., Matsuda, T. & Roberts, E. Purification and Characterization of Glutamate
Decarboxylase from Mouse Brain. *Journal of Biological Chemistry* **248**, 3029-3034 (1973).
https://doi.org/https://doi.org/10.1016/S0021-9258(19)44004-0

78 Albrecht, J. & Schousboe, A. Taurine Interaction with Neurotransmitter Receptors in the
CNS: An Update. *Neurochemical Research* **30**, 1615-1621 (2005).
https://doi.org/10.1007/s11064-005-8986-6

79 Bae, M. A., Lee, E. S., Cho, S. M., Kim, S. H. & Chang, K. J. The Effects of Dietary Taurine-
Containing Jelly Supplementation on Cognitive Function and Memory Ability of the Elderly
with Subjective Cognitive Decline. *Adv Exp Med Biol* **1370**, 395-403 (2022).
https://doi.org/10.1007/978-3-030-93337-1_37

80 Cui, W., Liu, H., Ye, Y., Han, L. & Zhou, Z. Discovery and Engineering of a Novel Bacterial L-
Aspartate alpha-Decarboxylase for Efficient Bioconversion. *Foods* **12** (2023).
https://doi.org/10.3390/foods12244423

81 Begley, T. P., Kinsland, C. & Strauss, E. The biosynthesis of coenzyme A in bacteria. *Vitam*
Horm **61**, 157-171 (2001). https://doi.org/10.1016/s0083-6729(01)61005-7

82 Lima, S. *et al.* The crystal structure of the Pseudomonas dacunhae aspartate-beta-
decarboxylase dodecamer reveals an unknown oligomeric assembly for a pyridoxal-5'-
phosphate-dependent enzyme. *J Mol Biol* **388**, 98-108 (2009).
https://doi.org/10.1016/j.jmb.2009.02.055

83 Jha, S., Speth, R. C. & Macheroux, P. The possible role of a bacterial aspartate beta-
decarboxylase in the biosynthesis of alamandine. *Med Hypotheses* **144**, 110038 (2020).
https://doi.org/10.1016/j.mehy.2020.110038

84 Yu, X. J. *et al.* Protein Engineering of a Pyridoxal-5'-Phosphate-Dependent l-Aspartate-
alpha-Decarboxylase from Tribolium castaneum for beta-Alanine Production. *Molecules*
25 (2020). https://doi.org/10.3390/molecules25061280

85 Webb, M. E. *et al.* Structure of Escherichia coli aspartate alpha-decarboxylase Asn72Ala:
probing the role of Asn72 in pyruvoyl cofactor formation. *Acta Crystallogr Sect F Struct*
Biol Cryst Commun **68**, 414-417 (2012). https://doi.org/10.1107/S1744309112009487

86 Cozzani, I. Spectrophotometric assay of l-glutamic acid decarboxylase. *Analytical*
Biochemistry **33**, 125-131 (1970). https://doi.org/10.1016/0003-2697(70)90446-X

87 Tramonti, A., De Biase, D., Giartosio, A., Bossa, F. & John, R. A. The roles of His-167 and
His-275 in the reaction catalyzed by glutamate decarboxylase from Escherichia coli. *J Biol*
Chem **273**, 1939-1945 (1998). https://doi.org/10.1074/jbc.273.4.1939

88 Tsukatani, T., Higuchi, T. & Matsumoto, K. Enzyme-based microtiter plate assay for γ-
aminobutyric acid: Application to the screening of γ-aminobutyric acid-producing lactic
acid bacteria. *Analytica Chimica Acta* **540**, 293-297 (2005).
https://doi.org/10.1016/j.aca.2005.03.056

89 De Biase, D., Tramonti, A., John, R. A. & Bossa, F. Isolation, overexpression, and
biochemical characterization of the two isoforms of glutamic acid decarboxylase from
Escherichia coli. *Protein Expr Purif* **8**, 430-438 (1996).
https://doi.org/10.1006/prep.1996.0121
90 Goujon, M. *et al.* A new bioinformatics analysis tools framework at EMBL-EBI. *Nucleic*
Acids Res **38**, W695-699 (2010). https://doi.org/10.1093/nar/gkq313
91 Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence
alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).
https://doi.org/10.1038/msb.2011.75
92 Miller, M. A., Pfeiffer, W. & Schwartz, T. in *2010 Gateway Computing Environments*
Workshop (GCE). 1-8.
93 Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066
(2002). https://doi.org/10.1093/nar/gkf436
94 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: computing large minimum evolution trees
with profiles instead of a distance matrix. *Mol Biol Evol* **26**, 1641-1650 (2009).
https://doi.org/10.1093/molbev/msp077
95 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490 (2010).
https://doi.org/10.1371/journal.pone.0009490
96 Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Res* **49**, W293-W296 (2021).
https://doi.org/10.1093/nar/gkab301

Figures.

Fig. 1: Characterization of *Bf*GAD$_{WT}$. **a** Possible role of gut microbes in producing

neuromodulatory molecules. **b** Spectra of PLP cofactor bound to the holoenzyme

*Bf*GAD$_{WT}$ captured at various pH, depicting tautomers of internal aldimine at 335 nm and

420 nm. **c** Plot of initial velocity as a function of *Bf*GAD$_{WT}$ concentrations (n=4). **d** Plot of

initial velocity vs. L-glutamate concentrations (n=2). The solid line is a sigmoidal fit to the

data.

Fig. 2: Structural and catalytic perturbations of engineered *Bf*GADs. a Aligned

structures of *Ec*GAD (1XEY, pink), *Bf*GAD$_{WT}$ (model, green) illustrating the position of the

residue D104* (*residue from the neighbouring monomer), and *Bf*GAD$_{D104N}$ depicting the

point substitution D104N*(model, gray). **b** Aligned structures of *Ec*GAD (1XEY, pink),

*Bf*GAD$_{WT}$ (model, green) illustrating the position of the residue F81, and *Bf*GAD$_{F81W}$

depicting the point substitution F81W (model, gray). **c** Aligned structures of *Bt*GAD (7X51,

purple), *Bf*GAD$_{WT}$ (model, green) illustrating the position of the residue D104* (*residue

from the neighbouring monomer), and *Bf*GAD$_{D104N}$ depicting the point substitution

D104N*(model, gray). **d** Aligned structures of *Bt*GAD (7X51, purple), *Bf*GAD$_{WT}$ (model,

green) illustrating the position of the residue F81, and *Bf*GAD$_{F81W}$ depicting the point

substitution F81W (model, gray). The dashed lines indicate hydrogen bond interactions

of the amino acid residues with substrate analog, glutarate (GUA) where pink bonds are

for interactions between *Ec*GAD-GUA, yellow bonds are for interactions between *Bf*GAD-

GUA, and purple bonds are for interactions between *Bt*GAD-GUA. **e** Spectra of PLP

cofactor bound to the holoenzymes *Bf*GAD$_{WT}$ and engineered variants, depicting

tautomers of internal aldimine at 335 nm and 420 nm. **f** Progress curves depicting initial

velocities for reactions catalyzed *Bf*GAD$_{WT}$ and variants (n=3).

Fig. 3: Wild-type and engineered *Bf*GADs can decarboxylate multiple substrates.
Chromatograms showing the CO_2 peaks obtained from the decarboxylation reactions of multiple substrates after 24 h incubation with **a** *Bf*GAD$_{WT}$, **c** *Bf*GAD$_{D104N}$, and **e** *Bf*GAD$_{F81W}$ using headspace GC (n=3, only one representative CO_2 peak is depicted for every reaction). The CO_2 peak areas are depicted for the decarboxylation of various substrates after 24 h incubation with **b** BfGAD$_{WT}$, **d** BfGAD$_{D104N}$, and **f** BfGAD$_{F81W}$ (n=3). Various

997   substrates are presented as specific colors which are consistent in all panels where L-

998   glutamate is blue, L-CSA is red, L-Asp is purple, L-HCA is orange and L-CA is green.

1000

Fig. 4: Production of multiple neuromodulatory molecules by Wild-type and evolved *Bf*GADs. Intensity analysis of TLC plates depicting intensity plots for each lane. The peak size is proportional to the intensity of the sample spot on the TLC plate. **a** Production of hypotaurine from the decarboxylation of L-CSA catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks highlighted in pink are for the substrate L-CSA and peaks highlighted in blue are for the product hypotaurine. **b** Production of taurine from the decarboxylation of L-CA catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks highlighted in pink are for the substrate L-CA and peaks highlighted in blue are for the product taurine. **c** Production of β-alanine from the decarboxylation of L-ASP catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks highlighted in pink are for the substrate L-ASP and peaks highlighted in blue are for the product β-alanine. **d** Production of homotaurine from the decarboxylation of L-HCA catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ in 24 h as indicated by TLC intensity analysis. For intensity plots, peaks highlighted in pink are for the substrate L-HCA and peaks highlighted in blue are for the product homotaurine. The original TLC plates are shown in Supplementary Fig. 10.

Fig. 5: *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ generate hypotaurine when presented with a

mixture of L-glutamate and L-CSA. Intensity analysis of TLC plates for various

timepoints of decarboxylation reactions catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ for mixed

substrates where the concentration ratio of L-glutamate to L-CSA is 1:10. Standards of L-

glutamate, L-CSA, GABA, and hypotaurine were included as references. Production of

hypotaurine (peaks highlighted in pink) can be seen as the peak area increases over time.

GABA peaks are highlighted in blue, peaks for the substrate L-CSA are highlighted in

beige, and peaks for the substrate L-glutamate are in brown. Spots visible on TLC plates

from top to bottom show corresponding intensity peaks from left to right.

Fig. 6: *Bf*GAD$_{D104N}$ generate taurine when presented with a mixture of L-glutamate

and L-CA. Intensity analysis of TLC plates for various timepoints of decarboxylation

reactions catalyzed by *Bf*GAD$_{WT}$ and *Bf*GAD$_{D104N}$ for mixed substrates where the

concentration ratio of L-glutamate to L-CA is 1:10. Standards of L-glutamate, L-CA,

GABA, and taurine were included as references. Production of taurine (peaks highlighted

in pink) can be seen as the peak area increases over time for *Bf*GAD$_{D104N}$ catalyzed

reactions. GABA peaks are highlighted in blue, peaks for the substrate L-CA are

highlighted in beige, and peaks for the substrate L-glutamate are in brown. Spots visible

on TLC plates from top to bottom show corresponding intensity peaks from left to right.

1039

1040

Fig. 7: *Bf*GAD_{WT} and *Bf*GAD_{D104N} generate β-alanine when presented with a mixture

of L-glutamate and L-aspartate. Intensity analysis of TLC plates for various timepoints

of decarboxylation reactions catalyzed by *Bf*GAD_{WT} and *Bf*GAD_{D104N} for mixed substrates

where the concentration ratio of L-glutamate to L-aspartate is 1:10. Standards of L-

glutamate, L-aspartate, GABA, and β-alanine were included as references. Production of

β-alanine (peaks highlighted in pink) can be seen as the peak area increases over time.

The increase over time for *Bf*GAD_{D104N} is very clear from 3 h to 9 h. Due to the differences

in the background intensity of the TLC plate in areas containing 3 h and 9 h samples vs

areas containing 24 h, 30 h and 48 h samples from *Bf*GAD_{D104N} catalyzed reaction, the

increase is the intensity peak is not uniform. GABA peaks are highlighted in blue, peaks

for the substrate L-aspartate (L-ASP) are highlighted in beige, and peaks for the substrate

L-glutamate are in brown. Spots visible on TLC plates from top to bottom show

corresponding intensity peaks from left to right.

1055

1056

1057
1058

Figure 8: Hypotaurine and Taurine detection via LC-MS/MS from *Bf*GAD_{WT} and

***Bf*GAD_{D104N} catalyzed mixed substrate reactions. a** Ion chromatograms of product

hypotaurine and the remaining L-CSA substrate in the mixed substrate reactions with a

1:10 ratio of L-glutamate:L-CSA catalyzed by *Bf*GAD_WT. **b** Ion chromatograms of taurine

and remaining L-CA in the mixed substrate reactions with a 1:10 ratio of L-Glutamate:L-

CA catalyzed by *Bf*GAD_D104N.

1065
1066
1067

1068
1069   **Scheme 1.** Predicted substrates and products of *Bf*GAD catalyzed decarboxylations.