Rapid Identification of Genomic Alterations in Tumors affecting lymphocyte Infiltration

(RIGATonI)

Running title: Identification of genomic variants and altered immune phenotypes in cancer

Raven Vella[1,2,3,4], Emily L. Hoskins[1,2,4], Lianbo Yu[4,5], Julie W. Reeser[1,4], Michele R. Wing[1,4], Eric

Samorodnitsky[1,4], Leah Stein[1,2,4], Anoosha Paruchuri[1,4], ORIEN investigators[5], Michelle

Churchman[5], Nancy Single[4], Wei Chen[4,7], Aharon G. Freud[4,7], Sameek Roychowdhury[1,4]

[1] Department of Internal Medicine, Division of Medical Oncology, The Ohio State University

[2] Biological Sciences Graduate Program, The Ohio State University

[3] Medical Scientist Training Program, The Ohio State Univer

[4] Comprehensive Cancer Center, The Ohio State University

[5] Department of Biomedical Informatics, The Ohio State University

[6] Aster Insights

[7] Department of Pathology, The Ohio State University

Corresponding author: Sameek Roychowdhury, Department of Internal Medicine, Division of

Medical Oncology, Comprehensive Cancer Center and The James Cancer Hospital, Ohio State

University, Columbus, OH 43210, USA. Tel: +1 614-685-5842; email:

Sameek.roychowdhury@osumc.edu

Key words: Genomics, immune infiltration, computational staining, tumor immunity

Word count: 4471 words

Abstract

Recently identified tumor genomic alterations are associated with altered tumor immune

microenvironments and therapeutic outcomes. These studies beg the question: are there other

genomic variations altering the immune microenvironment in tumors that can provide insight into

mechanisms of immune evasion? Current approaches to estimate immunity in bulk RNAseq from

tumors include gene set enrichment and cellular deconvolution. In contrast, our new software

package RIGATonI utilizes a gene agnostic approach to classify immune cell infiltration in tumors.

Using pathologist reviewed histology slides and paired bulk RNA sequencing expression data, we

trained a machine learning algorithm to detect inflamed and non-inflamed immune

microenvironments. RIGATonI enables identifcation of candidate genomic alterations associated

with immune infilration in subsets of cancer through evaluation of various features including gene

expression, genomic alterations, pathologist-classified tumors, clinical outcomes, and protein

validation. In addition to immune infiltrate classification, RIGATonI leverages a novel algorithm

using known protein-protein interactions for prediction of gain-of-function and loss-of-function for

genomic alterations to improve filtering of genomic alteration candidats. Applying these two novel

methods, we analyzed all genomic alterations present in The Cancer Genome Altas and

visualized promising results with R shiny for public use. Thus, we present our R tool and online

database for Rapid Identification of Genomic Alterations in Tumors affecting lymphocyte

Infiltration (RIGATonI).

Introduction

Immune surveillance is crucial for the eradication of cancer and is driven through understanding factors that impact the tumor immune microenvironment[1]. The tumor microenvironment is composed of tumor cells, along with surrounding immune cells, stroma cells and tissue matrix, and microbiota[1]. Each of these elements vary across patients and tumor types producing a spectrum of immune phenotypes and consequently clinical outcomes[1]. Although scRNA-seq and spatial transcriptomic technologies have dramatically increased our understanding of tumor immunology, there are substantially more tumor samples with bulk RNA sequencing and clinical data available for analysis. Thus, there is a need to develop computational tools specifically designed to assess immunity in large databases of tumor transcriptomes. In particular, there is a growing need to better understand the interconnected relationships between tumor genomics and their associated microenvironments[2, 3].

Although there are many techniques to assess immunity in cancer using bulk RNA sequencing data, there are alternative approaches that have yet to be explored. Current approaches (**Figure 1A**) include gene signature analyses (ImSig[4], Thorsson immune landscape of cancer[5]), cellular deconvolution (MCP-counter[6], QuantiseqR[7], CIBERSORT[8]), and ensemble results provided in databases (TIMER2.0[9], TIMEDB[10]). Gene set enrichment involves assessing the relative expression of gene sets related to immunity and then clustering cancers based on these results. This technique is particularly useful for understanding the drivers of immunity within one cancer type. This is routinely employed in large studies of immunotherapy outcomes to reveal what tumor features underscore interpatient differences[11, 12]. On the other hand, cellular deconvolution tools attempt to emulate flow cytometry or immunohistochemistry by estimating the number of specific immune cells in the sample[6-8]. This technique is extremely valuable for small subsets of tumors and for researchers who have a good understanding of tumor immunity. Databases of cellular deconvolution results have emerged which include ensemble analyses

across The Cancer Genome Atlas (TCGA) and other sources[9, 10]. As an exploratory tool for

researchers without strong bioinformatic resources, this is exceptionally useful. We developed a

novel computational approach to connect functional genomic variants and altered immune

phenotypes in an unbiased manner. Our approach explores new frontiers in precision immuno-

oncology not addressed by current methods: 1) model training with pathologist-defined immune

infiltration as gold standard, 2) consideration for immune cell histologic features (tertiary lymphoid

structures, dispersion characteristics, and degree of infiltration), 3) rapid, robust, and precise

immune phenotype classification across big data resources, and 4) candidate filtering that

leverages protein function prediction rather than simply gene mutation status.

We developed rapid identification of genomic alterations in tumors affecting lymphocyte

Infiltration (RIGATonI) to accurately predict both the protein-level function and associated immune

phenotype of genomic alterations. To build RIGATonI, we combined histologic features found by

computational staining[13], pathologist-defined immune infiltration, protein-protein interaction

networks[14], genomic data, and transcriptomic data (**Figure 1B**). First, we trained a machine

learning algorithm to predict immune phenotypes using bulk RNAseq expression. We validated

and fine-tuned our approach using manually reviewed histology by pathologists, computational

staining, immunohistochemistry, and flow cytometry**.** We also built a novel modeling algorithm

which can accurately predict the function of genomic variants (copy number alterations, single

nucleotide variations, and structural variations) from bulk RNAseq expression. These two

methods were combined to uncover connections between all the genomic alterations in TCGA

and the immune phenotypes of samples harboring these alterations. We created a visualization

interface (https://rigatoni.osc.edu) to help researchers access our TCGA analysis results on a

gene-by-gene basis. Additionally, we built an R package which can be used to perform the

analyses described. Finally, we provide examples of how RIGATonI identified genomic alterations

associated with altered immune infiltration in subsets of cancer. In summary, RIGATONI will allow

researchers to investigate and shortlist genes of interest that associate with immune phenotypes

in cancer.

Results

Building RIGATonI algorithm based on histology and bulk RNAseq

We used 403 tumors which were classified in a blinded fashion by two pathologists for low, intermediate, or high immune infiltration (**Figure 2A**). Each H&E slide was manually reviewed for amount of space occupied by lymphocytes (not occupied by tumor or stroma cells) and the distribution of these lymphocytes throughout the tumor (deeply penetrating, semi-penetrating, or peripheral). Further the pathologist also considered the quality of inflammation such as evidence of tumor cell killing or presence of tertiary lymphoid structures. All bulk RNAseq data for these samples were extracted and subsequent analyses utilized only RNAseq expression. 334/403 tumors were used for training the algorithm with XgBoost[15] machine learning in R to classify tumors as low, medium, or high immune infiltration. We assessed the accuracy of the algorithm by using the remaining 69 pathologist reviewed slides (not used for training), through comparison to computational staining results from TCGA[13], and finally an independent cohort of paired multiplex immunohistochemistry (mIHC) and flow cytometry data[16] (**Figure 2B-E**).

Assessment of the RIGATonI algorithm on tumors reserved for testing

We evaluated the RIGATonI model's assessment of 69 tumors to classify tumors high, medium, low infiltration and compared to pathologist review as the ground truth (**Figure 2B**). The balanced accuracy for high was 82.04%, the sensitivity was 73.33%, and the specificity was 90.74%. The balanced accuracy for medium was 63.4%, the sensitivity was 44.44%, and the specificity was 82.35%. The balanced accuracy for low was 82.58%, the sensitivity was 83.33%, and the specificity was 81.82%. The overall accuracy of the model was 71.01% (95% confidence interval 58.84% - 81.31%). The accuracy was significantly different from the no information rate (p <.01) and we failed to reject the null hypothesis of Mcnemar's test which indicates there is not significant evidence that the predictions made by the algorithm are different from the true

phenotypes (p > .05). Detailed statistics as well as a confusion matrix of the results are available

in **Supplementary Table 1**.

Protein level validation of RIGATonI assessment of immune infiltrate

We compared RIGATonI-classified low, medium, and high immune phenotypes based on

RNAseq from a study of 32 gastric cancer tumors which also included matching mIHC and flow

cytometry[16]. First, we found that highly infiltrated tumors detected by RIGATonI display higher

counts of immune cells collectively detected by mIHC (**Figure 2C)** including T cells and

macrophages. In *post hoc* analysis of these results, we discovered this was predominantly due to

increased CD8+ T cells in high samples compared to low samples (**Figure 2D**). In flow cytometry

results from the same study, we found that RIGATonI classsified highly infiltrated tumors display

significantly increased lymphocytes compared to RIGATonI-low tumors (**Figure 2E**).

Comparison of RIGATonI annotations to those of a computational staining

We applied RIGATonI to 5,202 tumors from TCGA and used an orthogonal approach for

computational staining of the H&E images as previously described[13]. Saltz et al., developed a

convolutional neural network to detect the percentage and distribution of lymphocytes on digital

histology images. First, we evaluated 5 types of spatial attributes of tumors classified by RIGATonI

as high, medium, and low infiltrates (**Figure 2F**). There is a significant difference in the

composition of spatial arrangements of lymphocytes between these three groups. Highly

infiltrated tumors had a higher proportion of brisk diffuse lymphocytes than the population (p <.01).

Brisk diffuse arrangements indicate a broad distribution of many lymphocytes throughout the

histology slide. Medium infiltrated tumors display brisk band-like arrangements more often than

the population (p <.01). Brisk band-like infiltration patterns mean that the lymphocytes are

clustered in a band across the slide, but that there are many lymphocytes. Tumors with low

infiltration displayed a higher proportion of non-brisk focal lymphocyte arrangements (p <.01).

Non-brisk focal arrangements indicate that there are very few lymphocytes in a handful of

locations across the image. Finally, we compared the number of lymphocytes across each group:

tumors with RIGATonI high status have increased lymphocyte infiltration compared to both

medium and low, and medium tumors display increased lymphocytes percentages to low tumors

(Figure 2G).

Developing an algorithm for functional assessment and shortlisting of tumor genomic

alterations

RIGATonI seeks to connect genomic alterations with features of immune infiltration. There

are many genomic alterations to evaluate; therefore, a functional measure of these alterations

can help to prioritize alterations that are more likely to have a biologic impact. Currently, there is

no bioinformatic tool to assess the functional impact of a genomic alteration from corresponding

RNAseq. To address this gap, we developed RIGATonI's function assessment algorithm using

parallel modelling and machine learning approaches. In contrast to DNA-based annotation of

gene alteration pathogenicity, this approach has several benefits: 1) RIGATonI's assessment

method is blind to the cause of altered function, so although we apply this method to groups with

different DNA mutation status, RIGATonI could be applied to differing methylation status,

alternative splicing conditions, or even treatment. 2) RIGATonI does not assess the mutation

change itself, rather it investigates the RNA-based changes associated by that genomic alteration.

This allows RIGATonI to assess novel variants and novel mechanisms of altered expression. 3)

Not all patients with a genomic alteration experience the same molecular effects. RIGATonI can

make predictions about alterations as a whole and sample level predictions based on the sample's

specific molecular characteristics. This allows for users to investigate either by mutation status or

functional status.

We sought to annotate alterations as gain-of-function (GOF) or loss-of-function (LOF). First, using the STRING[14] protein-protein interaction database, the algorithm identifies proteins which have direct and validated interactions with the gene of interest (**Figure 3A**). Both proteins which act on the gene of interest (upstream proteins) and proteins on which the protein of interest acts (downstream proteins) are considered. These two gene sets serve different purposes: upstream genes allow us to assess alterations which impact the expression level of the gene of interest; downstream genes allow us to assess the impact of alterations which may change the activity level of the gene of interest. We build a generalized linear model over a Poisson distribution using the extracted RNAseq counts of upstream and downstream proteins using samples with no alteration in the gene of interest. These two models are then saved and used to predict the RNAseq counts of the gene of interest within the mutant samples. 95% prediction intervals are created for both the upstream and downstream models. If the true counts of a mutant sample fall below the 95% prediction interval of either model, we annotate the sample LOF. If the true counts fall above the 95% prediction interval of either model, we annotate the sample GOF. If the true counts are within the 95% prediction interval for both models, the sample is annotated "unknown" (**Figure 3A**). To assess the function algorithm, we analyzed genomic alterations from 1008 oncogenes and tumor suppressor genes annotated by OncoKB [17]. OncoKB is the premiere source for clinically actionable genomic variants in cancer that are annotated based on curation and review of publications with corresponded experimental validation[17]. When applied to 10,464 tumors from the TCGA, the algorithm successfully identified 400 GOF alterations and 966 LOF genomic alterations (SNVs, structural variations, and copy number variations) within these 1008 genes. Genomic alterations harbored by fewer than 5 tumors were not assessed. Fusions were excluded mainly due to the high potential for false positives for fusions as many tumors harbored multiple fusions and imprecise breakpoints.

Validation of algorithm to classify known GOF and LOF genomic alterations

Of the 400 candidate GOF alterations uncovered, 62 were annotated in the OncoKB[17]

database. The algorithm correctly categorized 60 of 62 as GOF, and 2 were incorrectly

categorized as LOF, yielding an accuracy for GOF calling of 96% (**Figure 3B**). Notably, 84.5% of

the alterations annotated by the algorithm had no information available in OncoKB[17]; most of

these variants were whole gene amplifications (**Supplementary Table 2).** Similarly, we validated

the accuracy of this algorithm for LOF alterations. Of the 966 LOF alterations uncovered, only 229

were annotated in the OncoKB[17] database. The algorithm correctly categorized 207 as LOF, while

22 were incorrectly categorized as GOF, yielding an accuracy of 90% (**Figure 3B**). Further, 76%

of the candidate LOF alterations annotated by the algorithm had no information available in

OncoKB[17]; most of these variants were whole gene deletions or premature terminations

(**Supplementary Table 2)**. Altogether, these results suggest that this algorithm can successfully

classify known GOF and LOF mutations and uncover novel candidate alterations that have not

been well described.

The TCGA analysis yielded 7410 genomic alterations with possible immune effects among

5746 genes

These genes have been visualized in the RIGATonI web app. The front page of the

website allows the user to choose a gene of interest and subset the analysis to only include certain

cancer types or alterations (**Figure 4A**). After setting these parameters, there are two pages which

allow for analysis of the mutant vs control tumors. First, the user may analyze the tumors'

expression of specific genes using the transcriptome tab (**Figure 4B**). The user can choose to

analyze using mutation status, function, or immune status. Axis labels, colors, and graph type can

be changed for the user's convenience (**Figure 4B**). Across the website, non-parametric statistical

tests are performed, and their p-values reported. We uncovered alterations of interest in 22 cancer

types; most alterations discovered were associated with a highly infiltrated immune

microenvironment (**Figure 4C**). To verify our results and allow for more detailed investigation, we

have analyzed these cohorts with quantiseqR[7], a popular cellular deconvolution tool, in the immune tab (**Figure 4D**). The user can analyze the same subsets as on the transcriptome tab and make the same adjustments to the graphs. Here we also show the composition of immune cells as a whole (left) and allow the user to select cell-types to investigate (right) (**Figure 4D**). To determine the number of novel results among the RIGATonI output, we investigated 226,093 original research works between 6/12/2010 and 6/12/2023 mentioning cancer and immunity in their abstracts via text mining. We discovered that 2773 (48%) of the RIGATonI output genes had not been previously connected to cancer immunity (**Figure 4E**). Only 72 genes (1%) had been mentioned in cancer immunity abstracts more than 100 times (**Figure 4E**).

RIGATonI identifies genomic alterations associated with altered immune infiltration in cancer

We sought to explore how RIGATonI can identify genomic alterations associated altered immune infiltration. We chose to investigate two genomic alterations of interest which had been previously reported in cancer immunity literature: *IDH1* and *MED16*. *MED16* has been recently identified as a potential immunotherapy target in melanoma via a CRISPR mouse screen [18]. Additionally, *MED16* has been connected to cancer progression and treatment resistance in breast cancer[19] and papillary thyroid cancer[20]. RIGATonI identified a connection between *MED16* copy number loss and immunity in melanoma. *MED16*-mutant tumors display a highly infiltrated phenotype more often than wildtype tumors (p < .05) (**Figure 5A**) and occurred in 12% of melanoma. These alterations were annotated LOF (p < .01). Using quantiseqR[7], it is evident that *MED16*-mutant tumors display increased immune infiltration compared to wildtype (**Figure 5B**). On *post hoc* analysis, this difference can be attributed to CD8+ T cell infiltration. *IDH1* alterations have been studied for their neoplastic, metabolic, or immunologic effects in glioma, cholangiocarcinoma, and leukemia[21]. RIGATonI identified a novel connection between *IDH1* copy number loss and immunity in lung adenocarcinoma. *IDH1*-mutant tumors display a highly

infiltrated phenotype more often than wildtype tumors (p < .05) (**Figure 5C**) and occurred in 2.5%

of lung adenocarcinoma. These alterations were annotated LOF (p < .01). Using quantiseqR[7], it

is evident that *IDH1*-mutant tumors display increased immune infiltration compared to wildtype

(**Figure 5D**). This difference can be attributed to CD8+ T cell infiltration. Thus, *MED16* and *IDH1*

alterations are examples of how RIGATonI can discover genomic alterations with altered immune

infiltrates in subsets of cancer.

Discussion

RIGATonI's immune phenotyping algorithm leverages tumor RNAseq with two key

approaches and offers a novel means to discover genetic subsets of cancer with altered immunity.

Currently, gene set enrichment, cellular deconvolution, and ensemble approaches for estimating

immunity from bulk RNAseq have advantages such as deeply understanding the immunological

features of small cohorts, specificity of immune cell predictions, and increased access to big data

for research teams lacking bioinformatic expertise, respectively. In contrast, RIGATonI utilizes a

different approach from these methods by incorporating histologic features into our algorithm's

development and validation. RIGATonI provides an additional layer of information not investigated

by any other bulk RNAseq method: spatial characteristics of lymphocyte infiltrates. Although

absolute infiltration is important, it is crucial to understand whether the infiltrates are deeply

penetrating the tumor itself, displaying signs of active cell killing, and organize into tertiary

lymphoid structures. This information, to our knowledge, is not captured in any other approach.

RIGATonI is specifically designed to analyze thousands of samples and to categorize genomic

alterations and their association with altered immune infiltrates.

RIGATonI's prediction of protein-level effects of genomic alterations from bulk RNAseq

expression data is also novel. To our knowledge, only one other method has been devised to

determine the function of genomic variants using bulk RNAseq: eVIP2[22]. Importantly, eVIP2[22] was

validated using only 2 alterations in the same gene of interest whereas our validation set

demonstrated remarkable accuracy across 291 different alterations across 207 different genes

(**Figure 3B**). Additionally, although we use the protein connections in STRING[14], if a user wanted

to use a list of their own protein connections, they could do so with our technique thus identifying

switch of function or specific pathways effected by a genomic alteration. This flexibility is not

offered by methods which rely on DNA sequencing or by eVIP2[22] itself. VIPER is an R package

which utilizes cell-type specific regulons built from bulk RNAseq expression data to predict the

protein activity of different samples[23]. However, VIPER was not developed to determine the

mutation level effects of genomic alterations themselves and was not validated against a robust

set of known genomic alterations with molecularly confirmed protein-level effects[23]. RIGATonI's

use of parallel modelling using two gene sets is highly innovative (**Figure 3A**). These two gene

sets provide different information to the user and can be exploited separately or together. The

upstream gene list uses expression counts of transcription factors which directly impact the

expression of the gene of interest. Using this model, we can uncover genomic alterations which

increase or decrease the expression of the gene directly. The downstream model however is build

using any proteins on which the gene of interest acts. Using this model, we can understand how

a genomic alteration impacts the function of the protein when the expression of the protein is not

affected. We attempted to combine these models but found that doing so worsens the

performance of the model. The reason for this is unclear and an area for future exploration.

Although user specific gene lists can be used, we believe the use of the STRING[14]

database is a powerful feature of our tool. Initially, our group attempted to perform traditional

pathway analysis (using both Kegg[24-26] and GO[27, 28]) to determine the functional outcomes of

genomic alterations; however, we were not successful. Protein-protein interaction networks from

STRING[14] proved the most effective choice because of the ability to filter for proteins with direct

interactions with the gene of interest (**Figure 3A**). Individual research groups with specific

knowledge of a particular protein can bypass the use of STRING[14] and input their own upstream

and downstream gene lists.

The only other approach we are aware of which systematically assesses protein effects

on cancer immunity is CRISPR screening in mouse models[29]. This approach also has various

benefits: the ability to assess proteins which are not frequently mutated in human cancer,

molecular validation of the immunologic effects, and the ability to test different treatment

protocols. The RIGATonI software suite is strictly an in-silico approach, and all results certainly

require experimental validation to confirm. Doing so is outside the scope of this analysis, but we

did investigate the results of *IDH1* copy number loss in lung adenocarcinoma (**Figure 5C-D**).

Mutant *IDH1* has been connected to cancer immunity in glioma with improved survival benefit in

numerous cancer types[21]. We see a pattern of increased immune cell infiltration in IDH1 mutant

tumors compared to wildtype (**Figure 5C-D**). Additionally, we identified 72 genes with

immunologic affects which have been discussed more than 100 times in the literature (**Figure**

4E). These results together are encouraging there is evidence that these alterations have

legitimate immunological impacts. Despite this clear limitation, RIGATonI does have key

advantages over a CRISPR screen. RIGATonI assesses gain of function variants, alterations that

already occur in human cancers, and immunity in human patients not mice. Furthermore,

RIGATonI is inexpensive and instantaneous. Coupling RIGATonI output and those of a CRISPR

screen would be a powerful combination to filter and assess which variants merit further

investigation. In fact, we explored this possibility via a target recently identified by a CRISPR

screen, *MED16*[18]. Loss of function alterations in *MED16* associate with increased immune

infiltration and increased CD8+ T cells in melanoma (**Figure 5A-B**). A CRISPR screen performed

in a melanoma cell line specifically mentioned *MED16* as a promising target[18]. This result not only

gives credence to RIGATonI's output, but also extends the CRISPR screen results to include a

rapid analysis of human tumors. We hope to explore candidates from this CRISPR screen and

other in the future.

RIGATonI is a powerful approach for identifying novel genomic alterations with immune

effects. That said, RIGATonI suffers from several limitations including sample size limitations,

discordance between pathologists' reviews, and finally exclusion of genomic fusions from the

function annotation algorithm validation.

By using pathologist review rather than computational staining as a ground truth for our

algorithm, we do limit the sample size of our training data. However, the quality of sample

annotations is far more important that the size of the training data (in this case 403 tumors). For

instance, although the Saltz et al approach considers the digital slides as a whole and does not

differentiate between lymphocytes deeply penetrating the tumor and lymphocytes outside the

tumor boundaries[13]. Our initial model trained with samples selected using Saltz et al annotations

was discordant with pathologist annotations. New approaches like Lunit SCOPE IO also use deep

learning techniques to estimate immune cells on digital pathology slides[30]. Lunit SCOPE IO was

trained on >17,000 H&E images across 24 cancer types[30]. This scale of training data would never

be possible using manual pathologist review. Additionally, Lunit SCOPE IO considers the

boundaries of the tumor on the histology slide[30]. Importantly, the goal of this software is to predict

immunotherapy response not to quantify or qualify inflammation more broadly[30]. Lunit SCOPE IO

is primarily a clinical tool and does not make available its software for research use on big data

repositories. In contrast, RIGATonI is a bioinformatic research tool developed specifically to

analyze   connections   between   genomic   alterations   and   their   associated   immune

microenvironments. Although it could be argued that Lunit SCOPE IO could be adapted for this

purpose, not all large databases of tumors make digital slides available. For these reasons,

RIGATonI offers a distinct benefit over other methods to estimate spatial features of immunity in

big data despite smaller sample size of the training data.

Pathologist assessment introduces some subjectivity into the training data annotations.

This subjectivity is clear from the heterogeneity of annotations made within the medium group

(**Figure 2B**). Although this may limit the calculated accuracy of our model, it reflects the reality

that independent physician judgement of the same slides may yield different results. In fact, two

pathologists initially reviewed slides for this study and disagreed with each other's annotations

~50% of the time. We built models using each pathologist's annotations separately and together

using two different machine learning approaches (**Supplementary Table 3**). The best model was

yielded using the annotations from pathologist two (**Supplementary Table 3**). We speculate that

pathologist two's annotations are more internally consistent and thus easier to model. It is widely

agreed that physician opinion is the gold standard against which AI and machine learning

approaches should be measured[31]; however, doing so demonstrates that human judgement is

more malleable than computational modelling. In this case, our extensive validation using two

orthogonal datasets with protein and spatial characteristics reveals that although the model

statistics may suffer from human error, qualitatively the model performs exceptionally well

regardless.

Validation of RIGATonI's functional analysis only took place on alterations with >5

instances of the alteration to ensure that statistical tests could be performed to determine the

functional status. The ability of RIGATonI to classify alterations declines with declining sample

count. Additionally, co-mutations are not considered in the analysis of each sample because,

again, of sample size concerns. Finally, in our pan cancer analysis of TCGA, we did not consider

whether samples contained more than one alteration in each gene. Unfortunately, such

distinctions would not be computationally scalable. Because of this sample size limitation, in the

pan-TCGA analysis, we annotate copy number alterations more frequently than any other

alteration type. Copy number alterations are extremely common compared to specific point

mutations, gene fusions, or structural variations. Additionally, we did not group together early

terminations or different point mutations occurring at the same locus out of an abundance of

caution (early terminations at different loci may have different effects; different point mutations at

the same locus may have different effects). Without detailed knowledge of all 17 thousand genes

analyzed, we approached the first version of RIGATonI conservatively.

We were not able to include gene fusions in our function annotation algorithm validation.

Gene fusions are genomic events wherein two genes translocate together and sometimes

produce a functional gene product which behaves differently than the parent genes[32]. Not all gene

fusions have functional impacts on the gene of interest[32]. Additionally, although two samples

might harbor a fusion of the same two genes, the breakpoints of this fusion could be vastly

different and the resulting protein product entirely distinct[32]. Fusion calling bioinformatic methods

are very accurate for determining the breakpoints and read support for a fusion event; however,

these tools are intended to be manually assessed by a genetics expert who can determine

whether the resulting gene product is likely to yield altered gene function[32]. Developing a

computational approach to carry out this manual assessment for any gene of interest and any

fusion partner with any breakpoint is outside the scope of this analysis. Without such a method to

provide ground truth for our validation, we were not able to perform validation with fusion samples.

Despite this limitation, the RIGATonI method is blind to the genomic cause of altered function,

and therefore we don't believe there is any reason fusion samples would perform differently

(Figure 3A).

Despite these limitations, RIGATonI is an innovative and novel approach for identifying

genomic alterations with immunological impacts in cancer. Overall, RIGATonI is a novel tool

specifically designed for the emerging field of precision immuno-oncology. We believe RIGATonI

can be used for a vast variety of purposes including biomarker discovery and identification of

novel drug targets. RIGATonI will assist researchers in their search for targetable genes which

impact tumor immunity.

Methods

Tumor sequencing data. All data shown here were previously published or made available by a large database of tumor RNA sequencing (RNAseq) and whole exome sequencing data. Secondary analyses were performed within Ohio Supercomputer Center. Data sources include The Cancer Genome Atlas and the Oncology Research Information Exchange Network. RNAseq count data were downloaded from TCGA using GenomicDataCommons[33], and batch corrected with ComBat-seq[34] using institution of origin as the batch. Genomic variant calling was downloaded in the form of combined .maf files. Copy number alterations were downloaded in the form of gene based raw copy number. Finally, all whole genome .bam files were downloaded from TCGA and then processed with Delly[35], Manta[36], and GRIDSS[37] to assess for structural variants. These results were combined using SURVIVOR[38] with default settings. RNA sequencing data was also obtained from ORIEN and batch corrected with ComBat-seq[34] according to the RNAseq batch information made available through Aster Insights. Copy number alterations were downloaded from ORIEN in the form of gene based raw copy number. Demographic information was downloaded from ORIEN as well. ORIEN data is managed by Aster Insights, requests for this data should be sent to Aster Insights.

RIGATONI immune phenotyping algorithm validation with pathologist review. Two pathologists independently reviewed 403 slides assessing lymphocyte infiltration characteristic. Pertinent characteristics included percentage of space not occupied by tumors or stroma which was occupied by lymphocytes, dispersion of lymphocytes within tumor, and presence or absence of tertiary lymphoid structures. Taking into consideration all these characteristics, each pathologist annotated the slide either high, medium, or low infiltration. Paired RNA sequencing was collected and used along with annotations to build a series of models predicting immune phenotypes from bulk RNAseq counts. First, important features were selected using multinomial ElasticNet[39] via the R package glmnet[40]. Next, these features were used as predicted for six different machine

learning models. Three models were built with XgBoost[15] via the R package with the same name

and three with the R package ordinalForest[41]. Two models were built considering both pathologist

predictions, two considering just pathologist one and two considering pathologist two. Each

algorithm uses the same cohort of 334 samples for training and 69 samples for testing. The

algorithm with the best performance on the testing data was selected to be used going forward.

Please see supplemental table 4 for model specific information. Accuracy of the testing data for

the model selected were visualized using ggplot2[42].

Gastric cancer multiplex immunohistochemistry (mIHC) and flow cytometry. RNAseq count

data was downloaded from Saito et al along with flow cytometry and IHC outputs. These include

30 patients with gastric cancer in Tokyo[16]. These results were processed as previously described

in Saito et al. Using this resource, we used a MANOVA[43] to compare the IHC counts to determine

if there were any significant differences between groups. Next, we used a Wilcox test[44] to assess

the counts of each subset of IHC-marked cells to find significant differences. We performed

pairwise comparisons of flow cytometry results using Wilcox tests[44]. Results were visualized using

ggplot2[42].

Determine resulting immune phenotype of each genomic alteration using RNAseq data. An

R function was created which converts RNAseq count data to TPM using the R packages

DESeq2[45] and DGE.obj.utils[46], and then filters the data down to only the genes selected by

ElasticNet[39]. The immune phenotype of each samples was predicted using XgBoost[15]. The

proportion of high and low tumors for each cancer type were calculated. To analyze a genomic

alteration, all mutant samples provided to the function are compiled, and a 1-proportion z-test is

performed where the null hypothesis is that the proportion of hot samples will be equal to the

population proportion within that cancer type and the alternative hypothesis is that the proportion

of high samples is greater than the population proportion. If the null hypothesis is rejected (P<.05),

the genomic alteration is annotated high. If not, a second 1-proportion z-test is performed where

the null hypothesis is that the proportion of low samples will be equal to the population proportion

and the alternative hypothesis is that the proportion of low samples is greater than the population

proportion within that cancer type. If this null hypothesis is rejected, the genomic alteration is

annotated low. If not, the alteration is annotated Unknown.

Determining functional status of each genomic alteration using RNA-seq data. The

STRING[14] database's protein action version 10.5 was downloaded. The list of protein actions is

subset to include only actions on or by the protein of interest (POI). These actions are further

filtered into two lists: an upstream protein list with only proteins that act to affect the expression

of the POI, and a downstream protein list including all genes the POI activates, inhibits, or alters

expression. The upstream gene list is used to model the RNAseq counts of the POI using

modulators of the POI's expression. The downstream gene list is used to model the RNAseq

counts of the POI using downstream genes as indicators of its activity.

To model typical expression patterns of the POI, all samples with no alteration (control samples)

in the gene of interest (GOI) are collected, and two generalized linear models are created over a

Poisson distribution to predict the RNA counts of the POI/GOI. One model uses the upstream

protein list as predictors, and another using the downstream protein list. Both models predict the

RNAseq counts of the POI.

Next, the RNAseq counts of the GOI within samples harboring mutations (mutant sample)

predicted separately with each regression model. If, in either model, the expression of the GOI is

lower than the lower bound of the 95% prediction interval (created by ciTools[47]), the mutant

sample is annotated loss of function (LOF). On the other hand, if, in either model, the expression

of the GOI is higher than the upper bound of the 95% prediction interval, the mutant sample is

annotated gain-of-function (GOF). Falling outside the bounds of these prediction intervals

indicates that the mutant sample's GOI expression or activity is more different than that of a

control sample than we would expect from random chance. If the mutant sample's GOI

expression falls within the bounds of both these prediction intervals, we can conclude that there

is not enough evidence to indicate the mutation is causing abnormal expression or activity in that

sample.

Next, all mutant samples provided are compiled, and a 2-proportion z-test is performed (null

hypothesis z = .5) comparing the proportion of LOF and GOF annotations. If the null hypothesis

is rejected (P<.05), the genomic alteration is annotated with the more frequent sample level

annotation. If not, the alteration is annotated Unknown.

Function annotation algorithm validation. OncoKB[17] provides a list of oncogenes with either

targeting drugs or known oncogenic mutations. All alterations in these 1008 genes were analyzed

in parallel. The results were filtered to only include GOF and LOF calls from the algorithm. Various

alterations are called Unknown due to low sample count, confounding variables, or lack of known

connections in STRING[14]. We did not include these samples in the pan-cancer analysis of TCGA,

however users can elect to include them in their own analysis. Additionally, gene fusions were

removed due to complexity of their calling. This left 1366 genomic alterations to investigate. We

manually searched OncoKB[17] for information about each alteration and, if available, recorded the

true function of the variant. Results were visualized with ggplot2[42].

Building the R package. Functions were created with the R package devtools[48] which create an

upstream and downstream gene list from STRING[17], determine the function of a group of samples

using RNA expression data from bulk RNAseq, and the sample level immune phenotype using

RNA expression data from bulk RNAseq. These functions are described in detail above. The R

package along with relevant documentation is available at https://github.com/OSU-

SRLab/RIGATONI.

Comprehensive analysis of genomic alterations in TCGA. All mutation information in TCGA was downloaded and compiled. A sample is said to have a copy number variation (CNV) if the total number of copies is >= 6 or <2. We also considered the sex of the patient in question if the gene of interest was on the X or Y chromosome. For male patients, genes on the X chromosome were said to be deleted if there were 0 copies. For female patients, no genes on the Y chromosome were considered deleted. For any patients without biological sex available, we excluded them from the copy number analysis of genes on the X or Y chromosome. Next, all genomic alterations were analyzed in parallel through RIGATonI R package described above. We ensured to analyze each alteration within each cancer type for both functional annotations and immune phenotyping. The population proportions used for immune phenotype were those of the cancer type being analyzed. Finally, we stored the significant results from TCGA to be visualized by our online tool. The RIGATonI website is located at https://rigatoni.osc.edu/ and is managed by the Roychowdhury lab group and the Ohio Supercomputing Center. The website is built with the R package shiny[49], all graphs are visualized with ggplot2[42]. The website input is a user-provided GOI, and the website compiles gene level copy number, fusion, and simple nucleotide variation results from the pan TCGA analysis. Next, we provide various visualizations to the user (quantiseqR[7] cell type proportions, RNA seq count data, and primary site prevalence) along with a table describing the different genomic alterations which were both functional and immunogenic within the GOI.

Text mining of PubMed abstracts to estimate novelty of RIGATONI TCGA output. PubMed abstracts containing the words "cancer" and "immunity" or "immunology" or "immune" since 6/12/2010 were downloaded using the R package pubmed.mineR[50]. The function gene_atomization was used to perform text mining annotation of each gene mentioned. The RIGATonI results were extracted and each gene was annotated with their frequency of

appearance. Preprint publications were excluded from this analysis. A total of 226,093 abstracts

were analyzed. Results were visualized with ggplot2[42].

Gene level analysis of *MED16* and *IDH1*. These genes were analyzed as part of the pan-TCGA

analysis described above. Results were extracted to highlight here.

Acknowledgements. The results published here are in whole or part based upon data generated

by the TCGA Research Network: https://www.cancer.gov/tcga.

1.   Binnewies M, Roberts EW, Kersten K, Chan V, Fearon DF, Merad M, et al. Understanding the
tumor immune microenvironment (TIME) for effective therapy. Nature Medicine. 2018;24(5):541-50.
2.   Haslam A, Prasad V. Estimation of the Percentage of US Patients With Cancer Who Are Eligible
for and Respond to Checkpoint Inhibitor Immunotherapy Drugs. JAMA network open.
2019;2(5):e192535.
3.   Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Ou Yang TH, et al. The Immune Landscape
of Cancer. Immunity. 2018;48(4):812-30.e14.
4.   Nirmal AJ, Regan T, Shih BB, Hume DA, Sims AA-O, Freeman TC. Immune Cell Gene Signatures
for Profiling the Microenvironment of Solid Tumors. (2326-6074 (Electronic)).
5.   Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Ou Yang TH, et al. The Immune Landscape
of Cancer. (1097-4180 (Electronic)).
6.   Becht E, Giraldo NA, Lacroix L, Buttard B, Elarouci N, Petitprez F, et al. Estimating the population
abundance of tissue-infiltrating immune and stromal cell populations using gene expression. Genome
biology. 2016;17(1):218.
7.   Finotello F, Mayer C, Plattner C, Laschober G, Rieder D, Hackl H, et al. Molecular and
pharmacological modulators of the tumor immune contexture revealed by deconvolution of RNA-seq
data. Genome Medicine. 2019;11(1):34.
8.   Chen B, Khodadoust MS, Liu CL, Newman AM, Alizadeh AA. Profiling Tumor Infiltrating Immune
Cells with CIBERSORT. Methods in molecular biology (Clifton, NJ). 2018;1711:243-59.
9.   Li T, Fu J, Zeng Z, Cohen D, Li J, Chen Q, et al. TIMER2.0 for analysis of tumor-infiltrating immune
cells. (1362-4962 (Electronic)).
10.   Wang X, Chen L, Liu W, Zhang Y, Liu D, Zhou C, et al. TIMEDB: tumor immune micro-environment
cell composition database with automatic analysis and interactive visualization. Nucleic acids research.
2023;51(D1):D1417-D24.
11.   Motzer RJ, Robbins PB, Powles T, Albiges L, Haanen JB, Larkin J, et al. Avelumab plus axitinib
versus sunitinib in advanced renal cell carcinoma: biomarker analysis of the phase 3 JAVELIN Renal 101
trial. Nature Medicine. 2020;26(11):1733-41.
12.   Motzer RJ, Powles T, Atkins MB, Escudier B, McDermott DF, Alekseev BY, et al. Final Overall
Survival and Molecular Analysis in IMmotion151, a Phase 3 Trial Comparing Atezolizumab Plus
Bevacizumab vs Sunitinib in Patients With Previously Untreated Metastatic Renal Cell Carcinoma. (2374-
2445 (Electronic)).
13.   Saltz J, Gupta R, Hou L, Kurc T, Singh P, Nguyen V, et al. Spatial Organization and Molecular
Correlation of Tumor-Infiltrating Lymphocytes Using Deep Learning on Pathology Images. (2211-1247
(Electronic)).

14.   Szklarczyk D, Kirsch R, Koutrouli MA-O, Nastou KA-O, Mehryary FA-O, Hachilif R, et al. The
STRING database in 2023: protein-protein association networks and functional enrichment analyses for
any sequenced genome of interest. (1362-4962 (Electronic)).
15.   Chen T, Guestrin C, editors. Xgboost: A scalable tree boosting system. Proceedings of the 22nd
acm sigkdd international conference on knowledge discovery and data mining; 2016.
16.   Saito N, Sato Y, Abe H, Wada I, Kobayashi Y, Nagaoka K, et al. Selection of RNA-based evaluation
methods for tumor microenvironment by comparing with histochemical and flow cytometric analyses in
gastric cancer. Scientific Reports. 2022;12(1):8576.
17.   Chakravarty D, Gao J, Phillips SM, Kundra R, Zhang H, Wang J, et al. OncoKB: A Precision
Oncology Knowledge Base. LID - 10.1200/PO.17.00011 [doi] LID - PO.17.00011. (2473-4284 (Print)).
18.   Dubrot J, Du PP, Lane-Reticker SK, Kessler EA, Muscato AJ, Mehta A, et al. In vivo CRISPR screens
reveal the landscape of immune evasion pathways across cancer. Nature Immunology.
2022;23(10):1495-506.
19.   Li H, Li K, Shu D, Shen M, Tan Z, Zhang W, et al. MED16 Promotes Tumour Progression and
Tamoxifen Sensitivity by Modulating Autophagy through the mTOR Signalling Pathway in ER-Positive
Breast Cancer. LID - 10.3390/life12101461 [doi] LID - 1461. (2075-1729 (Print)).
20.   Gao H, Bai P, Xiao L, Shen M, Yu Q, Lei Y, et al. Mediator complex subunit 16 is down-regulated
in papillary thyroid cancer, leading to increased transforming growth factor-β signaling and radioiodine
resistance. (1083-351X (Electronic)).
21.   Ni Y, Shen P, Wang X, Liu H, Luo H, Han X. The roles of IDH1 in tumor metabolism and immunity.
Future Oncology. 2022;18(35):3941-53.
22.   Thornton AM, Fang L, Lo A, McSharry M, Haan D, O'Brien C, et al. eVIP2: Expression-based
variant impact phenotyping to predict the function of gene variants. PLOS Computational Biology.
2021;17(7):e1009132.
23.   Alvarez MJ, Shen Y, Giorgi FM, Lachmann A, Ding BB, Ye BH, et al. Functional characterization of
somatic mutations in cancer using network-based inference of protein activity. Nature genetics.
2016;48(8):838-47.
24.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic acids research.
2000;28(1):27-30.
25.   Kanehisa M. Toward understanding the origin and evolution of cellular organisms. Protein
science : a publication of the Protein Society. 2019;28(11):1947-51.
26.   Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for taxonomy-
based analysis of pathways and genomes. Nucleic acids research. 2023;51(D1):D587-d92.
27.   Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, Ebert D, et al. The Gene Ontology
knowledgebase in 2023. Genetics. 2023;224(1).
28.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for
the unification of biology. The Gene Ontology Consortium. Nature genetics. 2000;25(1):25-9.
29.   Holcomb EA, Pearson AN, Jungles KM, Tate A, James J, Jiang L, et al. High-content CRISPR
screening in tumor immunology. (1664-3224 (Electronic)).
30.   Shen JA-O, Choi YL, Lee T, Kim H, Chae YK, Dulken BW, et al. Inflamed immune phenotype
predicts favorable clinical outcomes of immune checkpoint inhibitor therapy across multiple cancer
types. LID - 10.1136/jitc-2023-008339 [doi] LID - e008339. (2051-1426 (Electronic)).
31.   Davenport T, Kalakota R. The potential for artificial intelligence in healthcare. (2514-6645
(Print)).
32.   Taniue K, Akimitsu N. Fusion Genes and RNAs in Cancer Development. LID -
10.3390/ncrna7010010 [doi] LID - 10. (2311-553X (Electronic)).
33.   Martin Morgan SD, Marcel Ramos. GenomicDataCommons. 1.26.0 ed2023.

34.   Zhang Y, Parmigiani G, Johnson WE. ComBat-seq: batch effect adjustment for RNA-seq count
data. NAR Genomics and Bioinformatics. 2020;2(3):lqaa078.
35.   Rausch T, Zichner T, Schlattl A, Stütz AM, Benes V, Korbel JO. DELLY: structural variant discovery
by integrated paired-end and split-read analysis. Bioinformatics (Oxford, England). 2012;28(18):i333-i9.
36.   Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Källberg M, et al. Manta: rapid
detection of structural variants and indels for germline and cancer sequencing applications.
Bioinformatics (Oxford, England). 2016;32(8):1220-2.
37.   Cameron DL, Schröder J, Penington JS, Do H, Molania R, Dobrovic A, et al. GRIDSS: sensitive and
specific genomic rearrangement detection using positional de Bruijn graph assembly. Genome research.
2017;27(12):2050-60.
38.   Jeffares DA-O, Jolly C, Hoti M, Speed DA-O, Shaw L, Rallis C, et al. Transient structural variations
have strong effects on quantitative traits and reproductive isolation in fission yeast. (2041-1723
(Electronic)).
39.   Zou H, Hastie T. Regularization and Variable Selection Via the Elastic Net. Journal of the Royal
Statistical Society Series B: Statistical Methodology. 2005;67(2):301-20.
40.   Friedman J, Hastie T, Tibshirani R, Narasimhan B, Tay K, Simon N, et al. glmnet: Lasso and Elastic-
Net Regularized Generalized Linear Models. cran2023.
41.   Hornung R. Ordinal Forests. Journal of Classification. 2020;37(1):4-17.
42.   Wickham H. ggplot2: Elegant Graphics for Data Analysis: Springer-Verlag New York; 2016.
Available from: https://ggplot2.tidyverse.org.
43.   Smith H, Gnanadesikan R, Hughes JB. Multivariate Analysis of Variance (MANOVA). Biometrics.
1962;18(1):22-41.
44.   Wilcoxon F. Individual comparisons by ranking methods. Biom. Bull., 1, 80–83. 1945.
45.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome biology. 2014;15(12):550.
46.   Law CW, Alhamdoosh MA-O, Su S, Dong X, Tian LA-O, Smyth GA-O, et al. RNA-seq analysis is
easy as 1-2-3 with limma, Glimma and edgeR. LID - ISCB Comm J-1408 [pii] LID -
10.12688/f1000research.9005.3 [doi]. (2046-1402 (Electronic)).
47.   John Haman MA, Institute for Defense Analyses. ciTools: Confidence or Prediction Intervals,
Quantiles, and Probabilities for Statistical Models. 2020.
48.   Hadley Wickham JH, Winston Chang, Jennifer Bryan, RStudio. devtools: Tools to Make
Developing R Packages Easier. 2.4.5 ed2022.
49.   Winston Chang JC, JJ Allaire, Carson Sievert, Barret Schloerke, Yihui Xie, Jeff Allen, Jonathan
McPherson, Alan Dipert, Barbara Borges, Posit Software, PBC, jQuery Foundation, jQuery contributors,
jQuery UI contributors, Mark Otto, Jacob Thornton, Bootstrap contributors, Twitter, Inc, Prem Nawaz
Khan, Victor Tsaran, Dennis Lembree, Srinivasu Chakravarthula, Cathy O'Connor, PayPal, Inc, Stefan
Petre, Andrew Rowls, Brian Reavis, Salmen Bejaoui, Denis Ineshin, Sami Samhuri, SpryMedia Limited,
John Fraser, John Gruber, Ivan Sagalaev, R Core Team. shiny: Web Application Framework for R. 1.8.0
ed2023.
50.   Rani J, Shah Ab Fau - Ramachandran S, Ramachandran S. pubmed.mineR: an R package with
text-mining algorithms to analyse PubMed abstracts. (0973-7138 (Electronic)).

A

	Gene Set Enrichment (ImSig, Thorsson et al, JAVELIN)	Cellular Deconvolution (TIMER, CIBERSORT, quanTIseq)	RIGATONI immune phenotyping
Goal	Find subgroups of cancer	Predict number of each immune cells and their subtype	Determine degree of inflammation in tumor sample
Feature Selection	Known immunity gene sets	Known immunity gene sets	Agnostic gene selection with machine learning
Solid Tumor Validation	Immunotherapy response	Correlation with flow cytometry, liquid cell suspensions, and in-silico mixtures	Histology slide review

B

Figure 1: A. Table of current approaches to estimate immunity in big data bulk RNA seq. Unlike other big data approaches for measuring immunity, RIGATonI is not developed to quantify the number of cells or the activity of those cells. **B.** RIGATonI was developed using both the degree of infiltration and the spatial organization of the lymphocytes. Instead of using canonical markers of immune activation, we allowed unsupervised selection of predictors to best predict pathologist review.

Figure 2: A. ORIEN samples used for training were selected based on a preliminary algorithm developed to predict strongly hot and cold phenotypes. Medium samples were selected at random from the ORIEN dataset. A pathologist reviewed each slide and annotated high, medium or low degrees of infiltration. We built a machine learning model to predict these annotations using XgBoost. We validated the accuracy of the algorithm using computational staining, a subset of pathologist reviewed slides not used for training, multiplex IHC, and flow cytometry of an independent dataset. **B.** The accuracy of the model (Y-axis) was determined using a subset of pathologist reviewed slides (X-axis) not using for training data. **C/D/E.** To confirm our algorithm was robust outside of the ORIEN training and testing database, we investigated an independent dataset of gastric tumors with paired RNAseq, mIHC and Flow cytometry. **C.** The overal composition of the immune microenvironment of high and low samples. **D.** The counts of CD8+ T cells detected by mIHC are significantly increased in high samples. **E**. Flow cytometry of these tumors demonstrated an increase in lymphocytes in RIGATonI high vs. low samples. **F/G.** We also investigated our algorithms association with histologic features detected by convolutional neural networks in TCGA. The overal distribution of spatial characteristics is significantly different across low medium and high subsets **(F)**. High samples more often display brisk diffuse lymphocyte patterns shown in maroon **(F)**. Additionally, the percentage of cells on the slide estimated to be lymphocytes is significantly greater in Medium samples compared to Low samples, High samples compared to Medium samples, and High samples compared to low samples **(G)**.

Figure 3: A. The function annotation algorithm was developed using upstream regulators and downstream targets of a given gene of interest from the STRING database. Parrallel linear models over a poisson distribution are built. First the model of the upstream regulators is assessed. Next, if the expression of the gene of interest falls within the prediction interval, the expression gene predicted by the downstream targets is assessed. **B.** The function annotation algorithm was assessed using OncoKB. 291 genomic alterations with annotations in OncoKB were called, 96.8% of the gain of funciton calls (60/62) were accurate and 90.4% of the loss of function calls (207/229).

Figure 4. A. After analyzing all available alterations from TCGA across all cancer types, we provide a public website to visualize and explore these results. The user selects a gene of interest and can filter results to types of alterations and cancers. Only positive results will be displayed. **B/D.** Once an alteration is selected for analysis, there are two methods to explore the characteristics of the mutant group. The first is to explore the expression of specific genes under the transcriptome tab. We provide relevant statistical tests and the functionality to change characteristics of the graph such as colors, axis labels, and graph type **(B).** We also analyzed each group using the cellular deconvolution software quanitseqR. These results are available under immune tab. We provide the same functionalities as the transcriptome tab. Estimated proportions of specific cell types can be explored (right) or the wholistic immune microenvironment can be displayed (left) **(D). C.** We discovered alterations in 22 cancer types which are associated with altered immunity. Most of these alterations are associated with a highly infiltrated immune microenvironment. **E.** To assess the novelty of these results, we performed a text mining study of cancer immunity abstracts in PubMed. 84% of the genes with suspected immunologic impacts from TCGA have been mentioned in cancer immunity literature less than 10 times. 72 genes or 1% of genes discovered were mentioned over 100 times.

Figure 5. A/B. In TCGA, we evaluated the immune impact of *MED16* loss in melanoma. We see mutant patients display significantly enriched highly infiltrated phenotypes compared to wildtype tumors **(A).** Using quanitseqR, we see enrichment of CD8+ T cells and overall increase in immune cells in mutant samples compared to wildtype **(B). C/D.** In TCGA, we evaluated the immune impact of *IDH1* loss in lung adenocarcinoma. We see mutant patients display significantly enriched highly infiltrated phenotypes compared to wildtype tumors **(C).** Using quanitseqR, we see enrichment of CD8+ T cells and overall increase in immune cells in mutant samples compared to wildtype **(D).** We also see decreased NK cells and neutrophils in IDH1 mutant tumors **(D).**