Spatial attention alters visual cortical representation during target anticipation

Ekin Tünçok[1], Marisa Carrasco[1,2,^], Jonathan Winawer[1,2,^]

[1] Department of Psychology, New York University, New York, NY, 10003, USA

[2] Center for Neural Science, New York University, New York, NY, 10003, USA

[^] Equal contributors

Corresponding author: Ekin Tünçok, tuncokekin@gmail.com

Number of pages: 43

Number of figures: 8

Conflict of interest: The authors declare no conflict of interest.

Acknowledgments: This research was supported by NIH NEI Grant R01-EY027401 to MC and JW, and NYU Center for Brain Imaging pilot grants to ET, MC and JW. We thank Michael Jigo for help in piloting the experimental protocol, Ilona Bloem and Rania Ezzo for helpful discussions, Jan Kurzawski, Ilona Bloem and Nina Hanning for contributing to the experimental code, Sarah Master for assistance in sharing some anatomical MRI data, Ilona Bloem, Marc Himmelberg and Yuna Kwak for feedback regarding the manuscript, as well as the members of the Carrasco Lab and Winawer Lab for useful feedback. This work was supported in part through the NYU IT High Performance Computing resources, services, and staff expertise.

ABSTRACT

Attention enables us to efficiently and flexibly interact with the environment by prioritizing some
image features in preparation for responding to a stimulus. Using a concurrent psychophysics–
fMRI experiment, we investigated how covert spatial attention affects responses in human visual
cortex prior to target onset, and how it affects subsequent behavioral performance. Performance
improved at cued locations and worsened at uncued locations, relative to distributed attention,
demonstrating a selective tradeoff in processing. Pre-target BOLD responses in cortical visual
field maps changed in two ways: First, there was a stimulus-independent baseline shift, positive
in map locations near the cued location and negative elsewhere, paralleling the behavioral
results. Second, population receptive field centers shifted toward the attended location. Both
effects increased in higher visual areas. Together, the results show that spatial attention has
large effects on visual cortex prior to target appearance, altering neural response properties
throughout and across multiple visual field maps.

Keywords: spatial attention, anticipatory response, fMRI, population receptive fields,
psychophysics

INTRODUCTION

The visual system needs to prioritize behaviorally relevant locations because it has limited
capacity to fully process all neural inputs simultaneously[1-4]. Attention is a mechanism that
selectively prioritizes some aspects of information over others. In covert spatial attention, the
prioritized location can be varied while keeping the fixation point constant, which enables the
study of spatial properties of top-down attention in retinotopic coordinates across the visual field.
Covert spatial attention improves behavioral and neural sensitivity to a stimulus at the attended
location at the cost of worsening the processing at other locations[4-8].

Two effects of covert spatial attention on neural responses have been reliably found. First, it
increases response amplitude for neurons with receptive fields near the attended location,
measured in both animal single unit studies and human fMRI, in striate and extrastriate areas[9-
13]. Second, it modulates the preferred position and size of single neuron receptive fields
(RFs)[14,15] and population receptive fields (pRFs)[16-20]. These studies used experimental designs
in which the neural measurements were made in response to the attended stimulus, or to
another stimulus, such as a retinotopic mapping stimulus, while the attended stimulus was
viewed.

The effects of attention begin, however, before the appearance of an attended target, once the
task relevance of a particular location is established. For example, in non-human primates, V4
neurons increase their baseline activity prior to the attended target appearance[10]. In humans,
attention affects the BOLD signal in visual cortex prior to target appearance[21-23]. Moreover,
contralateral alpha oscillations prior to target appearance gate information flow to enhance the
visual cortical representation of an attended feature or location[24-27]. These pre-target attentional
effects are predictive of performance on the subsequently presented target[28-31, but see 32].

It is unknown to what degree similar circuits are involved during target anticipation, which is
purely top-down, and attentional modulation of stimulus evoked responses, in which top-down
and stimulus-driven effects interact. Moreover, studies of attention during stimulus presentation
have led to conflicting conclusions about whether spatial attention primarily works by changing
neural position tuning[19] or response amplitude[33]. The finding in macaque V4 that distinct
neuronal populations are modulated by pre- and post-stimulus attentional signals[34] suggests
that human fMRI evidence on the attentional modulation of responses to an attended stimulus
may not generalize to the anticipatory effects of attention. In particular, how receptive field
properties and amplitude modulations are affected by attention before stimulus onset is an open
question.

Here, we investigated the spatial organization of anticipatory attentional effects in human visual
cortex by implementing a concurrent fMRI/psychophysics experiment. We manipulated the
location of endogenous (voluntary) attention in each trial with an event-related design. During
the period between cue and target, a pRF mapping stimulus, irrelevant to the participant's task,
was briefly presented. This enabled us to measure how response amplitudes to the task-
irrelevant mapping stimuli were influenced by attention for the upcoming target. Because the
pattern of response amplitudes across different bar stimuli determines the pRF location, we

could also assess how attention affected pRF position. Moreover, because participants did a
task at the end of each trial, we were also able to measure performance benefits (valid cues)
and costs (invalid cues) of attention, relative to distributed attention (neutral cues). Together the
experimental results revealed robust effects of attention on behavior, and changes in BOLD
amplitude and pRF position throughout the entirety of multiple visual field maps (from V1
toLO1). Together, these results provide compelling evidence that anticipatory spatial attention
changes basic neural response properties throughout the cortical visual processing hierarchy.

RESULTS

Participants (*n*=8) performed a two-alternative forced-choice orientation discrimination task
while attention was manipulated in the scanner (**Figure 1**). On each trial, a central pre-cue
indicated the likely target location: one of the four cardinal meridians (focal attention) or all four
locations (distributed attention). Then, participants viewed a pRF mapping stimulus, followed by
an array of four Gabor stimuli. A response cue at the end of each trial indicated which of the four
stimuli to discriminate (CW/CCW of horizontal). For each of the five attention conditions, we
measured (a) behavioral sensitivity to the Gabor target to quantify the selective attentional effect
on performance and (b) the BOLD response to the mapping bar stimuli to quantify the
attentional effect on response amplitude and position tuning.

Figure 1. Concurrent psychophysics and pRF mapping protocol. Each trial started with a
fixation cross, which then turned into an attentional pre-cue indicating one of four likely target
locations (focal attention, 80% of total trials) or all four locations (distributed attention). The pre-
cue was followed by one of 49 mapping stimuli as participants sustained their attention at the
cued location. After a jittered ISI, the target and distractors, four Gabor patches, appeared.
Then, a response cue informed participants on which stimulus' tilt to report (clockwise vs
counterclockwise). In 60% of trials, the response cue indicated the same location as the pre-cue
(*valid trials);* in 20% of trials, it indicated a different location (*invalid trials).* The remaining 20%
were neutral trials where the attentional cue was not informative. Participants received accuracy
feedback by a color change in the fixation cross upon responding. The attentional cue (one of
five) and the mapping stimulus (one of 49) varied from trial to trial in a pseudo-randomized
design. Image properties are altered for increased visibility.

Attention improves performance at the attended location and impairs it at unattended locations

For each participant, we estimated the tilt angle threshold at each Gabor location to equate
baseline performance at 82% for each target (in an experimental room). Thresholds revealed
typical performance fields[35-37]; mean thresholded tilt angle across participants was 11.6° (68%
CI bootstrapped: 7.4°-15.8°) and 11.4° (7.3°-15.9°) at left and right horizontal meridians, and
15.8° (9.2°-22.7°) and 21.6° (16.8°-26.1°) at lower and upper vertical meridians. Individual
thresholds were used in subsequent psychophysics measurements during the scanning
sessions, to equate the baseline performance across locations within and across participants.

Attention had a large effect on performance. We calculated sensitivity (d′) at each target location
for valid, neutral, and invalid cue types (**Figure 2A**). The sensitivity for validly cued trials was
more than double the sensitivity for neutral pre-cue trials, indicating that attention improved
performance: across participants and the four locations, the mean d′ for valid trials was 4.0 (3.7-
4.4) vs 1.6 (1.3-1.8) for neutral trials. Further, sensitivity was lower for invalidly cued trials, with
a mean d′ of 0.5 (0.4-0.7), indicating that attention impaired performance at the unattended
locations, imposing a behavioral tradeoff. An ANOVA yielded a main effect of cue type
$(F_{(2,14)}=114, p<0.0001, \eta_p^2=0.94)$, no main effect of location $(F_{(3,21)}=1.63, p=0.23, \eta_p^2=0.19)$,
as expected, and no interaction between location and cue type $(F_{(6,42)}=2.26, p=0.10,$
$\eta_p^2=0.24)$. Further, reaction times ruled out speed-accuracy tradeoffs **(Figure 2B)**. Participants
were about twice as fast in the valid (*M*=0.30 sec [0.28-0.32]) than neutral (*M*=0.56 sec [0.53-
0.59]) or invalid (*M*=0.56 [0.50-0.61]) trials. Finally, the average gaze position was within 0.05°
of the fixation point in each of the five cue types. Good fixation is essential to measure covert
spatial attention and accurate pRF position during focal attention.

Figure 2. A) Behavioral sensitivity (d′) as a function of cue validity at four target locations (UVM:
upper vertical meridian, LVM: lower vertical meridian, LHM: left horizontal meridian, RHM: right
horizontal meridian). Sensitivity was higher for valid and lower for invalid than for neutral cue
trials. Filled circles indicate participant values. **B)** Reaction time (s) as a function of cue validity.
Valid cue trials yielded the fastest responses, ruling out speed-accuracy trade-offs. Reaction
time did not differ between invalid and neutral cues. Error bars indicate the 68% confidence
interval computed across bootstrapped participant means. **C)** Participant- (circles) and group-

level (triangles) averaged gaze position for different pre-cue types revealed only minor

deviations from fixation point (≤0.05 deg). This figure is produced by *fig2_A_B_dprime_RT.m*

and *fig2_C_averageGaze.m*.

pRF model fits are accurate

To conduct the behavioral and fMRI mapping experiments concurrently, we used an event-
related design with randomized mapping stimulus position across trials. This mapping sequence
differs from the more common sweeping stimulus design. We first checked the accuracy of the
pRF model fits. PRF models were estimated for each surface vertex in each retinotopic map
separately for the five attention cues (attend-up, attend-down, attend-left, attend-right, attend-
all) from the responses to the retinotopic bar stimuli that appeared before the Gabor target and
distractors. First, a single GLM was conducted for each participant, providing beta weights for
each bar position (48+1 blank) crossed with each cue type (5) **(Figure 3)**. PRF models,
parameterized by preferred center (x, y) and size (σ), were then fit to the GLM beta estimates,
rather than the fMRI time series.
168
The pRF model provided an accurate fit to the BOLD data from each of the five attention
conditions. We quantified the median variance explained by the pRF models across vertices

within an ROI for each participant and each attention condition. We then averaged this value

across participants, to obtain R^2 values per attention condition per visual field map. Model

accuracy was similar across the five attention conditions, with ranges of 39-40% for V1, 39-41%

for V2, 36-38% for V3, 31-34% for hV4, 34-38% for LO1, and 26-29% for V3A/B. We also

solved the pRF models for the data averaged across the five conditions. As expected, the

variance explained by this model was higher: V1–64%, V2-64%, V3-59%, hV4-56%, V3A/B-

48%, LO1-58%. The relatively high variance explained by all the pRF models is important for

the subsequent analyses, which make use of the pRF parameter estimates.

Figure 3. Trial-based analysis of functional time series data. **A)** In the first step, raw timeseries

data from each vertex were submitted to a general linear model. Vertex responses were

modeled as a weighted combination of the BOLD activity to distinct mapping stimulus bar

locations (48+1 blank) for four different focal attention conditions and the distributed attention

condition. **B)** Estimated beta weights were converted to percent BOLD change and reorganized

and concatenated to create a pseudo time series for each vertex separately for each attention

condition. **C)** Beta weights were submitted to the pRF model to estimate the pRF of each vertex

separately for each attention condition. **D)** Vertex-level pRF estimates were aggregated on each

participant's native surface map to draw the visual field map boundaries from V1 to LO1.

Retinotopy maps from an example participant (*wlsubj138*) estimated under the averaged

attention condition.

Attentional modulation of the BOLD amplitude is time-locked to the cue onset and sustained
throughout the trial

An anticipatory effect predicts a sustained increase in BOLD amplitude from cue onset until
target onset in cortical regions whose pRFs are close to the target. There are two other possible
ways that attention might impact the BOLD response. First, an attentional modulation might start
later in the trial, in response to the Gabor target, rather than to the cue. Second, an attentional
response might be transient, starting after the cue but not sustaining throughout the trial. The
GLM does not distinguish among the three possibilities because the Gabor target always
followed the cue and the BOLD signal pools over time; hence all predict increased BOLD
responses on trials when focal attention is directed toward a voxel's pRF. Nonetheless, we can
distinguish the three possibilities by analyzing the full within-trial time courses separately
depending on whether the duration of the bar stimulus was one or two seconds. Though
sluggish, the fMRI time series can resolve timing effects at the second or even sub-second time
scale[38]. The three accounts make distinct predictions for when attention-related modulations
rise and fall within a trial **(Figure 4A)**.
207
First, we ask whether there is a modulation of the BOLD amplitude from focal attention. As an
example, in **Figure 4B** we show the time course data from V3 vertices with pRFs near the cued
location (cortical target ROIs; see *Methods*), time-locked to either the cue (top) or the Gabor
target (bottom). The plots show that focal attention increased the BOLD amplitude: the time
course of Attend In minus Attend Out is a positive curve, for both 1-s and 2-s trials, irrespective
of how the data are time-locked. This pattern holds for all visual maps **(Supplementary Figure
1)**.
215
Second, we consider the *onset* of the attentional effect on the BOLD signal. When time locked
to the cue (**Figure 4B**, left), the attentional effect becomes apparent at about 2s, rising at the
same rate for the 1-s and 2-s bar trials, consistent with the interpretation that the attentional
modulation is locked to the cue rather than target. In contrast, when the time series data are
aligned to the Gabor target (**Figure 4B**, bottom), the rise time is about a second earlier for the 2-
s trials, inconsistent with the possibility that the attentional effect is time-locked to the target. To
quantify these observations for each visual area, we computed the rising latency of the
attentional modulation of BOLD separately for the 1- and 2-s bar trials, operationalized as the
time point where the attentional modulation reached 10% of the maximum response before the
peak. This point was inferred by fitting the rising portion of the attentional modulation curves
with a logistic function, bootstrapped 1,000 times across participants. We then subtracted the
estimated latency parameters of 1-s bar trials from 2-s bar trials **(Figure 4C; Supplementary
Table 1).** When time-locked to the cue onset, the difference in rising latency between 1- and 2-s

bar trials was close to 0 for all visual field maps (i.e., 0 was within the 68% CI of the

bootstrapped estimates). Had the attentional effect been time-locked to the Gabor target rather

than the cue, the difference measures would have been about +1s, inconsistent with the data in

most visual field maps (all except for V2). In contrast, when time-locked to target onset, the

difference was close to −1s, meaning the rise was one second earlier for the 2-s trials than 1-s

trials, relative to the target. These timing differences confirm that attentional resources were

allocated at an approximately fixed time following the attentional cue, rather than the Gabor

target, ruling out the "target-evoked" account **(Figure 4A).**

237
Figure 4. A) Three possibilities of the attentional modulation profile with the expected outcomes
for the cue- and target-locked time series analysis. Anticipatory and sustained account predicts
that participants deploy attention upon receiving the attentional cue, and sustain it until the
target onset; the rise time of attentional modulation is locked to the cue onset, and the fall time
is locked to the target onset. Target-evoked account predicts no anticipatory modulation, with
both the rise time and the fall time locked to the attentional target onset. Finally, the anticipatory
and transient account predicts that attentional modulation rises and falls prior to the target
appearance; both the rise and fall time of attentional modulation is locked to the cue onset. **B)**
An example of attentional modulation of BOLD in 1-s and 2-s mapping stimulus bar trials in V3,
cue-locked (left), and target-locked and shifted (right). Vertical rectangles indicate the time
points of interest representing the rising and falling latency of attentional modulation. **C)**
Estimated latency parameters from the best-fitting logistics functions to cue-locked (top) and
target-locked (bottom) BOLD activity for rising time. Latency of the rise time was around the
same for 1-s and 2-s bars in cue-locked responses in contrast to target-locked responses,
indicating that attentional modulation started around the cue onset. **D)** Estimated latency
parameters from the best-fitting logistics functions to cue-locked (top) and target-locked
(bottom) BOLD activity for falling time. Latency of the fall time was delayed in line with the

prolonged anticipatory period in target-locked responses, indicating that attentional allocation
was sustained until the target appearance. This figure is produced by
fig4_A_TTA_bar_duration.m and *fig4_B_TTA_latency.m*.
258
Third, we considered the attentional *offset*, and whether the anticipatory response was
sustained throughout the trial. Specifically, we asked whether the attentional modulation of the
BOLD signal lasted a second longer when the bar stimulus lasted a second longer. We
addressed this question with a similar analysis used for the onsets, but estimating the falling
latency as a measure of when the attentional responses started decaying. We fit a logistic
function, bootstrapped across participants, and identified the time point at which the response
declined 10% from the peak. When aligned to the cue, the decline was about a second later for
the 2-s trials in each visual area, indicating that attention was maintained throughout target
anticipation. When aligned to the target, the latency difference was mostly zero, indicating that
the decay period was aligned to the target appearance (**Figure 4D**), ruling out the "Anticipatory
and transient" account (**Figure 4A**)**.**
270
Together the analysis of rise time and fall time is consistent with an anticipatory account in
which focal attention was initiated following the cue onset and was sustained until the target
appeared, giving rise to an increased BOLD response throughout the trial. The variability in rise
times is about double the variability in the fall times (error bars in **Figure 4C** vs **4D**), suggesting
more temporal variability in engaging attention following the cue than in withdrawing attention
following the target. This difference is not surprising because in our protocol there were several
seconds between cue and Gabor target onset; participants were not pressed to deploy attention
as fast as possible.

Focal attention affects the BOLD amplitude in a spatially tuned manner

We next asked how the attentional modulation varied as a function of retinotopic position. Our
main observation is that in all six visual field maps[1], during the anticipatory period, there was an
increase in BOLD amplitude for regions of visual cortex with pRF centers near the attended
target (*target enhancement region*) and decreases for regions with pRF centers far from the
target location (*distractor suppression region*)**.** We visualize these effects in 2D stimulus space
by plotting each vertex's change in amplitude (focal cue minus distributed cue, averaged across
all 49 bar positions), as estimated by the GLM (**Figure 5A**). The 2D space is defined by pRF
centers computed from the data averaged across the 5 attention conditions. For higher SNR, we
rotated the 2D maps from each of the four focal conditions to align them at the upper vertical
meridian and averaged across the rotated responses and participants. A mixed ANOVA of 6
(visual field map: V1, V2, V3, hV4, V3A/B, LO1) x 2 (attention: attend in, attend out) x 4 (target
location: UVM, LVM, LHM, RHM) on the averaged amplitude change of pRFs within each
cortical target ROI revealed no interaction between attention condition and polar angle location

---

[1a] We made measurements in V1, V2, V3, hV4, V3A/B, and LO1. V3A and V3B are considered to be
distinct maps, but their boundary is not straightforward to identify. Hence for analysis purposes we group
them as "V3A/B"; we analyze data from seven maps, grouped as six regions (maps) of interest.

(F(15, 60)=0.96, p=0.51). The target enhancement region was a wedge shape (red) centered at
the cued meridian. The shape of the target enhancement regions–narrow near the fovea and
wide in the periphery–reflects increasing receptive field size as a function of eccentricity. For all
six maps, the rest of the visual field (outside the wedge) tends to show decreased responses,
comprising the distractor suppression field (blue). The amplitude and consistency of
suppression is greater in some maps (e.g., hV4, V3A/B) than others (V1, V2).
299

Figure 5. A) Change in BOLD response from distributed to focal attention projected onto the preferred baseline position of each vertex pRF. Attentional activity was projected onto the vertex pRF center for each polar angle target. These responses were rotated to align at the upper vertical meridian, and averaged across targets. Change in BOLD activity from distributed to focal attention was positive around the cued target location (red blobs), and negative away from the cued location (blue blobs), indicating distractor suppression fields. **B)** 2D focal attention responses collapsed onto one dimension, polar angle distance from target, representing the spatial tuning of attentional modulation around the iso-eccentric target configuration (dots indicate the measurement, solid line indicates the fits averaged across iterations) **C)** Estimated characteristics of spatial tuning of attention across visual field maps. Across maps, the peak of attentional BOLD was similar (top panel, red), but the distractor suppression magnitude varied (top panel, blue). Estimates of the attentional spread (bottom panel) showed a largely uniform

314 spread area from V1 to V3A/B, with an increase in the spread of attention at LO1. This figure is
315 produced by *fig5_A_2D_amplitude_plots.m* and *fig5_B_C_vonMises_fits.m*.
316

The pattern of enhancement and suppression varies across visual cortex

To quantify the effects of enhancement and suppression, we summarize the 2D maps as one-
dimensional polar angle tuning curves, fit by a difference of two von Mises functions **(Figure 5B;**
Table 1). A difference of two curves captures the clear center-surround organization: increased
amplitude near the attended location, decreased amplitude far from the attended location. The
increase in BOLD amplitude for target enhancement varied two-fold across visual field maps,
ranging from 0.04 in hV4 to 0.08 BOLD% in LO1. The magnitude of suppression varied six-fold
among areas, ranging from −0.02 (V1) to −0.12 (hV4) **(Figure 5C)**. V1 and V2 showed mostly
enhancement, whereas hV4 and V3A/B showed mostly suppression. The overall modulation in
BOLD amplitude due to focal attention (enhancement minus suppression) generally increased
along the visual hierarchy (V1<V2<V3<hV4,V3A/B). Visual areas also differed in the spatial
extent of attentional enhancement, which was largest in LO1 and smallest in hV4, quantified as
the degree width of the positive portion of the 1D tuning plots (**Figure 5B, C**). The small spatial
extent in hV4 is related to its large distractor suppression magnitude (most of the tuning curve is
below 0). Conversely, LO1 had the largest width and the smallest distractor suppression
modulation.
333

ROI	Peak (%BOLD change)		Trough (%BOLD change)		Width (°)	
	Mean	68% CI	Mean	68% CI	Mean	68% CI
V1	0.07	0.06, 0.09	-0.02	-0.04, -0.01	105	90, 120
V2	0.08	0.07, 0.09	-0.03	-0.04, -0.02	82	71, 92
V3	0.07	0.05, 0.08	-0.06	-0.09, -0.04	95	83, 107
hV4	0.04	0.03, 0.05	-0.12	-0.13, -0.09	74	64, 84
V3AB	0.06	0.04, 0.08	-0.09	-0.11, -0.09	89	75, 102
LO1	0.08	0.06, 0.1	-0.04	-0.05, -0.02	140	132, 142

334
Table 1. Estimated peak, trough and width of attentional spread within each visual field map.

The effect of focal attention is independent of stimulus location

Thus far, the pre-target effects of focal attention across cortical locations show that the
modulation of the fMRI signal was highly location dependent, with attentional increases in
BOLD% change for vertices near the cued locations and decreases for vertices far from the
cued location. Here we consider the complementary analysis, the effect of attentional
modulation as a function of the mapping stimulus location rather than the vertex. To do so, we

defined regions of interest, *cortical target ROIs*, comprising vertices whose pRFs are close to
one of the focal targets. As an example, we plot the pseudo-time series for the cortical target
ROI for the left target. The pseudo-time series consists of the response amplitudes to different
bar stimuli organized by bar position, separately for distributed attention, "attend in" (focal
attention to the left) and "attend out" (focal attention to one of the other three locations,
averaged across the three locations). As expected, we see two peaks, one when the horizontal
and one when the vertical bar stimuli overlap the receptive fields **(Figure 6A).** Across visual field
maps and stimulus positions, the distributed attention responses tended to fall in between the
attend in and attend out responses, consistent with the pattern of behavioral effects shown in
Figure 2.
352
Across the six visual field maps, the effect of focal attention was largely independent of the
mapping stimulus. First, even in the absence of a bar stimulus ("blank" condition), the BOLD
amplitude differs between attend-in and attend-out (**Figure 6A**, dots). Second, for most visual
field maps, there is a difference between attend-in and attend-out across the different stimulus
locations. Third, to more directly assess whether the attentional modulation depends on the
baseline response (the response amplitude to neutral cues), we plotted response to attend-in
minus response to attend-out against response to the distributed cue (**Figure 6B**). These plots
enable us to distinguish among several possible ways in which attention could modulate the
responses. A positive correlation would indicate multiplicative gain, a modulation amplitude
proportional to the baseline response. A flat line above the x-axis would indicate an additive shift
–an increase in response amplitude for attend-in irrespective of the bar position. The data are
most consistent with an additive shift: no correlation between the attentional modulation and the
responses to the mapping stimuli under distributed attention. Averaged Pearson's correlation
coefficients across participants were around zero in all visual field maps, except for a very small
positive correlation in hV4 (median and confidence intervals from bootstrapping in **Figure 6B**).
We interpret this baseline shift as an anticipatory effect of attention, meaning that anticipation of
the cued target increases the baseline BOLD amplitude at the appropriate retinotopic locations,
independent of the mapping stimuli.

Figure 6. A) Averaged %BOLD change of left target meridian selective vertices as a function of mapping stimulus position for attend-in (red), distributed (gray), and attend-out (blue) conditions. Filled dots represent the blank presentation. Bottom section (purple) shows the difference between attend in and attend out at each bar position. Error bars of attentional modulation were computed across participant means for 68% confidence interval. **B)** Correlation between attentional modulation (attend in–attend out) and baseline response levels (distributed), averaged across polar angle locations and participants. Each dot represents a mapping stimulus position. Purple dots represent the blank presentation. Purple line represents the best fit to the group data. Correlation coefficients and 68% confidence bounds were computed by averaging and correlating the subsampled data 1000 times. This figure is produced by *fig6_A_ROI_timeseries.m* and *fig6_B_ROI_timeseries_corr.m*.

pRF centers shift towards an attended location

We just showed that for vertices with pRF centers close to a cued location, the effect of attention was additive; an approximately fixed increment in BOLD amplitude irrespective of the

bar position. There are two reasons, however, why we might expect to be able to find shifts in
pRF centers. First, small differences in the effect of attention on BOLD responses to just a few
bar positions near the pRF center could shift the pRF center toward an attended location, as
observed previously[16,18]. Second, the conclusion about additive effects was based on analysis
of only those vertices with pRFs centers close to targets; for pRFs whose centers were further
away, other patterns might be observed. Here we quantified shifts in pRF centers as a function
of the attentional cue.
393
To estimate pRF shifts, within each visual field map, we computed the average pRF center
distance to an attentional target when focal attention was deployed to that target and compared
this to the distance of the pRF centers when attention was distributed. In all maps, the distance
decreased during focal attention, consistent with a shift toward the attended location (**Figure
7A)**. The decrease was slight in V1 ($M=-0.04°$, 68%CI= [$-0.07,0$]), and increased 10-fold along
the cortical hierarchy (V3A/B: $-0.46°$ [$-0.54,-0.38$]; LO1: $M=-0.37°$[$-0.41,-0.33$]). As a control
analysis, we randomly shuffled the attention cue on each trial and reran a GLM on the shuffled
design matrix, followed by a new set of pRF solutions estimated from the shuffled GLM weights.
For this analysis, the pRF center distance change to attentional targets was around 0 in all
visual field maps **(Figure 7A)**. This indicates that anticipatory attentional signals bias the
position preference of pRFs in visual cortex, and increasingly so up the cortical hierarchy.
405

406
Figure 7. A) pRF centers were attracted towards an attended target. For each visual field map,
averaged distance change to an attentional target from distributed to focal attention is plotted.
PRF center changes were quantified by comparing the distance of the distributed and the focal
attention pRF center from an attentional target. In all visual field maps, the distance to an
attentional target was smaller under focal attention condition, with the effect magnitude
increasing with cortical hierarchy. When pRFs were estimated from a shuffled GLM analysis, the
center shift effect disappeared (red data points). Error bars represent the 68% confidence
interval across bootstrapped participant means. Dots indicate participant data. This figure is
produced by *fig7_A_average_spatial_shifts.m.* **B)** PRF center shifts averaged across vertices
separately for each focal attention cue direction. The end point of vectors, representing the
average pRF center estimated for each focal attention cue, are aligned with the direction of focal
attention. **C)** PRF center shifts visualized along the horizontal (top) and vertical (bottom)
attentional shift axes in V3, hV4 and V3A/B/LO1 averaged. At each bin, the vector color
indicates whether the center shift occurred towards (black lines) or away (red lines) from the
attentional target. Overall, pRF center shifts are in the expected direction, in line with the
attentional shift. This figure is produced by *fig7_B_directional_vector_graphs_of_shifts.m.*
423
424
We next examined the topography of center shifts across the visual field in maps with the
largest shifts: V3, hV4, V3A/B, LO1. We first averaged the estimated pRF center across vertices
separately for each attention condition within each map. We then recentered average center

positions such that the distributed attention condition is represented in the origin, and plotted the
focal attention pRF centers as a vector end point. For each focal attention condition, the
average pRF center followed the direction of where attention was allocated **(Figure 7B)**. We
then examined the pattern of shifts as a function of pRF location. To do so, we visualized the
changes in pRF centers for paired cue conditions (upper vs lower, left vs right) for vertices
binned within regions of the visual field (annulus sectors; **Figure 7C**). Overall, the pRF center
shift direction was congruent with the attentional shift axis for both horizontal and vertical cue
locations. The magnitude of pRF center shift increased from fovea to periphery, and increased
from V3 to the higher maps. These patterns are not an artifact of binning, a problem that has
affected some reports on attention-related shifts of pRFs[39]. The same patterns hold whether the
binning of vertices is based on the average pRF location (attend-up and -down, or attend-left
and -right) or based on independent data, i.e., pRFs solved from the neutral cueing
(Supplementary Figure 2). Moreover, the results shown in Figure 7A are not binned.

DISCUSSION

We investigated the cortical specificity, stimulus specificity, and spatial tuning of anticipatory
attentional signals in human visual cortex by implementing an event-related pRF mapping
protocol that enabled simultaneous measurements of BOLD and behavior. Perceptual sensitivity
increased at the attended location and decreased elsewhere. Visual cortical responses across
the entirety of visual field representation changed in two main ways under focal attention in
anticipation of the target. First, in parallel with behavioral benefits and costs of attention, the
response amplitude of vertices whose pRFs were near the target location increased and far
from the target location decreased. While the magnitude of target enhancement was mostly
similar across visual field maps, that of suppression varied. Second, attentional modulation of
response amplitude was approximately uniform as a function of mapping stimulus position,
indicating a baseline shift in responses rather than a change in response gain. Third, the pRF
preferred centers were shifted towards the anticipated attentional target in all visual field maps,
and increasingly so up the cortical hierarchy.
455
Importance of the neutral condition in an attentional experiment
A novel feature of our experiment was measuring pRFs while manipulating the attended location
on a trial by trial basis. A disadvantage of the event-related design is that it is less efficient for
pRF mapping[40,] but we compensated for this by increasing the number of scans. The trial-by-
trial attention manipulation differs from previous fMRI studies on pRFs and attention, which used
either designs in which the cue was always valid and the task was always performed at the
cued location; the attended location was constant within a scan of several minutes[17,19] or
attention continuously tracked the mapping stimuli[16,18]. A benefit of our design is that it enabled
us to measure the performance differences among valid, neutral, and invalid cueing. A hallmark
of spatial attention is that it imposes tradeoffs: performance improves at the attended location
and worsens at unattended locations[3,5,41-43]. Our trial-by-trial design enabled verification of the
attentional task in our fMRI study. The selective nature of attention was confirmed by our
behavioral results, which showed large differences among the three conditions, valid better than
neutral (benefit) and invalid worse than neutral (cost). This provided a secure footing for
examining the positive and negative neural effects of attention on neural responses.

471
The selective nature was assessed neurally on pre-target BOLD by comparing the effect of focal
vs distributed cues and by comparing responses at the attended and unattended locations.
When assessing the benefits and costs of attention, it is important to consider what the neutral
reference point is. In psychophysical protocols, it is common to quantify attentional benefits and
costs against a distributed attention baseline condition[6, 41, 43, 44]. In contrast, in electrophysiology,
the comparison is often between "attend-in" and "attend-out," which both entail focal
attention[1,8,45]. Without a neutral condition, there is ambiguity about whether response
differences between the two conditions arise from a change at the attended location (enhanced
responses), at the unattended location (suppression), or a combination. Similarly, fMRI mapping
experiments on spatial attention tend to either compare the effects of attending two
locations[13,17,19,20], or use attend-fixation as baseline[16,19,46]. Attending fixation or attending a focal
location in the periphery is likely to result in withdrawal of resources from the rest of the visual
field, making it difficult to assess local target enhancement and distractor suppression effects.
Using distributed attention as a baseline removes these potential ambiguities. We illustrate how
the choice of the neutral condition can affect estimates of both attention-related enhancement
and suppression **(Supplementary Figure 3)**. Consistent with this schematic, our results
showed that the BOLD response amplitude to the mapping stimulus bars under distributed
attention primarily falls in between *attend-in* and *attend-out* states, confirming that it constitutes
an effective baseline measurement.
491
492   **Spatial attention induces extensive pre-target changes throughout the representation of**
493   **the visual field**
494
Anticipation enables people to successfully interact with the environment. Anticipation plays an
important role in many cognitive phenomena like working memory[47] and expectation from
stimulus probability[48,49]. Attention directs anticipation as to when, where or which information in
the environment needs to be prioritized and further processed than the rest, guiding action
production and memory[50,51]. Thus understanding how anticipatory responses in attention affect
the nervous system is an important goal in cognitive neuroscience. Doing so requires separating
the effects of anticipation from the effects of the onset of the attended target. The most
straightforward way to do so is to measure neural responses in the absence of a target by
having long delays between cue and target. And indeed, several fMRI studies have
characterized anticipatory modulation of BOLD response amplitude in the absence of visual
stimulation[21-23,28-30]. Such studies consistently reveal changes in BOLD amplitude during
anticipation. These studies cannot, however, measure either the effect of attention on position
tuning, which requires presentation of a stimulus at a variety of target locations.
508
Our design allowed us to characterize three systematic effects simultaneously. First, the
modulations in signal amplitude affected the entirety of visual field maps in each map studied
(increases near the cued location, decreases elsewhere, i.e. push-pull dynamics). Second, in
regions near the cued location, response increases were largely non-specific to the mapping
location, meaning that the attentional modulation was not limited to just mapping stimuli that
were close to the cued location. Third, pRF preferred positions shifted toward the focally

attended target. This effect was seen throughout each of the visual maps, with overall larger
shifts in later visual field maps. Together, these systematic patterns imply that attentional
anticipation reshapes responses throughout the visual maps.

A multiplicity of effects of attention on BOLD amplitude

Single neuron studies show that spatial attention can induce both additive[52, 53] and
multiplicative[54-57] effects. The simplest multiplicative effect is a scaling of the response
amplitude; however, more complicated patterns can emerge from gain changes if the gain
modulation acts on the inputs. For example, a shift in the contrast response function or in
receptive field location have each been interpreted as changes in *input* gain. For position shifts,
this manifests as a multiplication of an attentional gain field with a neural receptive field[15, 45]. If
the attention field and the receptive field differ in center location or in size or both, then input
gain causes a shift in the receptive field, rather than a response gain.

We found clear evidence for two of these three effects: a baseline shift and a position shift. As in
the single unit data[15] and computational models[15,45], we interpret the position shift as a gain
modulation on inputs. The shift effect is, by definition, stimulus dependent: the pRF center
location depends on which bar positions elicit the largest responses. We interpret the baseline
shift to indicate that anticipation of the target increased neural activity independent of the
mapping stimulus. This interpretation is supported by the observation that focal attention
increases the BOLD response during blank trials, when no mapping stimulus is present. The
two effects are measured in different units –the baseline change is measured in response
amplitude (percent BOLD) and the shift is measured in degrees of visual angle. We can,
however, make a comparison by estimating how much each effect influences the response
amplitude as a function of bar position (**Figure 8**). This shows that the baseline shift is the
dominant effect in V1, whereas both the position shift and the baseline shift are substantial in
hV4. Note that the effect of position shifts on the pattern of responses to the bars is not evident
when averaging across a region of interest (**Figure 8**) because the direction of the shift differs
across voxels.

Fig 8. A simulation of the BOLD response to the 49 mapping stimuli in V1 and hV4. The upper
panel shows separate traces for focal and distributed attention. The simulations assumed typical
baseline shifts from our datasets (V1:0.07%, hV4:0.04%) and typical position shifts (V1:0.04°,
hV4:0.3°). The bottom panel plots the difference between focal and distributed attention,
showing two effects: a mean above 0, indicating the baseline shift, and a modulation around the
mean caused by the position shift. This figure is produced by *fig8_simulateShifts.m*.

We did not observe a change in response gain. The most likely explanation lies in the type of
attention we studied. We deliberately isolated anticipatory responses from the stimulus-evoked
modulation. The task-relevant target (the Gabor) was low contrast, small and briefly presented
to minimize its effect on the BOLD. The mapping stimulus was designed to dominate the BOLD
signal but was not relevant to the task. It is likely that, had the mapping stimulus been task-
relevant, we would have observed a gain change. Indeed, attention-related changes in
response gain have been observed in an experiment in which participants performed a task on
the mapping stimulus [18]. A less likely explanation is the possibility that fMRI is generally
insensitive to the type of gain modulations caused by attention. This might seem plausible
because several fMRI studies measuring the effect of spatial attention on the contrast response
function found a baseline shift rather than a gain change[46,58-60, but see 61], in contrast to
electrophysiology studies which report both[52-55]. Some have speculated that the discrepancy
may be due to an inherent insensitivity of the BOLD signal to gain modulation[62, 63]. We think the
limitation is more likely due to the complexities of how gain changes affect the pooled response
of a diverse neural population[64]. This is supported by two observations. First, a recent fMRI
study on feature-based attention observed a gain change in the contrast response function[65].
Second, as noted above, several studies on spatial encoding (rather than spatial contrast) have

reported robust attention-related gain effects, either in terms of response gain[18], or position
shifts[17, 19]. Hence it seems likely that were there a large change in response gain, our protocol
could have detected it. Rather, anticipatory spatial attention seems to have large effects on
baseline response and input gain, with less effect on response gain.

Attentional enhancement and suppression may recruit cortical areas differentially
The amplitude of attentional modulations, particularly in terms of distractor suppression, varied
across cortical areas. Suppression magnitude increased along the cortical hierarchy from V1 to
hV4. The pattern of target enhancement was quite different. In fact, the map with the largest
suppression effect, hV4, had the least enhancement and conversely, the map with the largest
enhancement, LO1, had little suppression. If the two effects were mediated by the same
mechanism, we would expect that enhancement and suppression would covary across maps,
contrary to our results. Our results are more consistent with the idea that there are at least
partially independent circuits mediating target enhancement and distractor suppression in visual
cortical areas, as suggested by some psychophysical and neural results[66-69].
580
The spatial extent of the attentional effects was relatively similar across maps. With the
exception of LO-1, there was a sharp transition from enhancement to suppression, with about
one quarterfield of enhancement and three quarterfields of suppression. The sharp transition
indicates that the top-down signals mediating anticipatory target selection are represented with
high precision in visual cortex. The similar spatial extent across most maps is compatible with a
single attentional field modulating responses in multiple maps inferred by a model-based
analysis of how attention alters pRF location[17]. We confirm it with a direct measure of the spatial
extent of amplitude modulation.
589
Polar angle asymmetries.
We chose four target locations on the four cardinal meridians because visual performance is
known to differ at these locations. Specifically, performance is typically best along the
horizontal, intermediate along the lower vertical, and poorest along the upper vertical meridian[70].
Certain structural and functional differences in the retina[71,72] and cortex[73-76] might underlie these
asymmetries. We equated discriminability in a pre-scan experiment without focal attention by
choosing orientation deviations that yielded the same performance at all locations, which
showed the largest tilt angle on the upper vertical meridian, and smallest on the horizontal.
Once equated, performance for each neutral, valid and invalid trials did not differ across
meridians during the scanning experiment, consistent with the finding that endogenous attention
improves performance similarly at iso-eccentric locations[77]. In addition to replicating this
behavioral finding, we also show neural evidence that attentional modulations are similar at
different iso-eccentric locations, when baseline performance is equated.
603
CONCLUSIONS
Attention alters many properties of visual cortical responses across the representation of the
visual field in anticipation of a target, prioritizing the processing of the target to facilitate the
subsequent stimulus-evoked activity. These effects include shifts in pRFs toward the attended
location and stimulus-independent amplitude modulations that are positive near the cued

location and negative away from it. The positive and negative amplitude modulations parallel the
positive and negative effects on performance for valid and invalid trials, respectively. The
combination of a baseline shift in BOLD amplitude and a position shift in pRFs indicate two
processes by which anticipation affects activity in visual cortex: a change in input gain
underlying the position shift, and a sustained change in neural activation. An important
remaining question is how each of these two effects, measured in anticipation of a stimulus,
alter responses to the subsequent stimulus, measured both behaviorally and neurally.
616
617 **Data and code availability.** Code used to model, analyze, and visualize this data resides in
618 https://github.com/ekintuncok/Attention_pRF. Raw and processed data reside in
619 https://osf.io/g8b9v/.
620
621 **Author contributions.** Conceptualization and methodology: ET, MC, JW; Software: ET; Data
622 collection: ET; Analysis: ET; Visualization: ET, MC, JW; Writing – original draft: ET; review and
623 editing: ET, MC, JW.
624
Funding acquisition: NIH NEI Grant R01-EY027401 to MC, JW; NYU Center for Brain Imaging
Pilot Grant to ET, MC, JW.

MATERIALS AND METHOD

Participants

Nine human participants (M_{age} = 28.5, seven females, two males) with normal or corrected-to-
normal vision participated in the experiment. Following behavioral thresholding and training
sessions outside the MRI scanner (about 1.5 hours), participants took part in four scan sessions
(*n*=7), except for one participant who became unavailable after 3 sessions, and one participant
who was excluded from the study after 3 sessions due to poor quality retinotopic maps (After
participant exclusion, M_{age} = 29.1, with seven females, one male). Participants were
compensated $30/h for participation. They signed a consent form approved by the New York
University Institutional Review Board, and an MRI screening form prior to each scan session.

Apparatus

For both data collected inside and outside the scanner, stimuli were generated on an Apple
iMac using MATLAB (R2019) and Psychtoolbox-3[78]. Both inside and outside the scanner, the
gaze position of the participants' dominant eye was tracked at a sampling rate of 1 kHz using an
Eyelink 1000 (SR Research, Ottawa, Ontario, Canada). For the behavioral thresholding and
training sessions outside the MRI scanner, stimuli were displayed on a cathode-ray tube monitor
with a flat screen (1280 x 960 pixels, 40.5 x 30 cm; 60 Hz). A Konica Minolta LS-100
photometer was used to linearize the display luminance prior to data collection. Participants sat
57 cm away from the monitor. Behavioral responses were collected using a computer keyboard.
In the MRI scanner, stimuli were displayed on a ProPixx DLP LED projector in the MRI scanner
with a linear gamma table (VPixx Technologies Inc., Saint-Bruno-de- Montarville, QC, Canada).
The projected screen (60 cm x 36.2 cm) had a resolution of 1920 x 1080 with a refresh rate of
60Hz. The participants viewed the screen through an angled mirror mounted on the head coil.
The total viewing distance (eye to mirror plus mirror to screen) was 86.5 cm. Behavioral
responses were collected using two assigned keys of a 4-button fiber optic response box
(Current Designs). Participants were placed inside the scanner with positioner pads to reduce
head motion.

Stimuli for attentional task

All stimuli were presented on a gray background (~60 cd/m^2). A white (~120 cd/m^2) central
cross subtending 0.35° in visual angle was used for stabilizing gaze position. One or all four
arms of the cross changed from white to black (~0 cd/m^2) to cue participants to attend to one of
four locations, or to distribute attention to all locations, respectively. There were four target
stimuli presented at 6° of eccentricity on the four cardinal meridians (0°, 90°, 180° and 270°).
Each target was a tilted Gabor patch, constructed by windowing a sine wave grating (4 cycles
per degree, 10% Michelson contrast) by a Gaussian (1° full-width-at-half-maximum), truncated
within a square aperture at 3° per side. The Gabor patches were tilted slightly clockwise or
counterclockwise from horizontal. To eliminate position uncertainty, white placeholders (~120
cd/m^2), positioned just outside the corners of the four apertures, indicated the target positions
throughout the experiment.

Stimuli for pRF mapping

Between the attentional cue and target, a retinotopic mapping stimulus was viewed for 1 or 2
seconds. The mapping stimulus was a contrast pattern windowed within a bar aperture (3° in
width). All stimuli were limited to a circular window 12.4° in radius. The bar aperture was
presented either horizontally or vertically, at one of 24 positions per orientation. The 24 vertical
and the 24 horizontal bar apertures gridded the circular window in steps of 1° between two
adjacent locations, with an overlap of 2°. The contrast pattern within the bars was composed of
curved gray-scale contours (peak spatial frequency of 3 cpd) with Michelson contrast of 20%.
These textures were found to be effective in eliciting BOLD responses across multiple
retinotopic maps[79,80].

Task

Trial sequence and attentional cueing. Each trial began with the presentation of a white
central fixation cross for 300 ms **(Figure 1)**. The cross then changed to a pre-cue, which was
shown alone on the screen for 300 ms: On 80% of trials, the pre-cue consisted of one of the
four arms of the fixation turning black to cue *one* of the four target locations (focal attention;
equal probability at each location). On the other 20% of trials, the pre-cue consisted of all four
arms changing to black to cue *all* target locations (distributed attention). In the focal condition,
cue validity was 75% (i.e. a response cue matched the location of the pre-cue for the 75% valid
trials, but not for the 25% invalid trials; see below). Because the task was difficult (thresholded
in pre-scan testing) and the cue was informative, it was in the participant's best interest to
attend to the cued location for better performance in the task.
687
Following the 300 ms white cross and the 300ms pre-cue, a mapping stimulus was presented
for 1000 or 2000 ms. The two durations were used to reduce expectations of when the stimulus
would disappear, and the target would appear. The bar aperture did not move during this 1- or
2-s period, but the contrast pattern inside it updated to a new random sample of the texture
every 250 ms. In 10% of trials, the mapping stimulus was blank to estimate the baseline BOLD
response in the absence of mapping stimuli, consistent with prior pRF mapping protocols[81]. The
pre-cue remained on the screen during the 1s or 2-s bar stimulus. The prolonged presentation
of the pre-cue was to ensure that attention would be deployed and sustained while the mapping
stimulus was displayed. The mapping stimulus was followed by a jittered interstimulus interval
(ISI) of 50 ms (10% of trials), or 300, 400, or 500 ms (equiprobable at 30%). During the ISI, only
the white fixation cross was presented. The two durations of the mapping stimulus and the
jittered ISI imposed temporal uncertainty about the target appearance, to promote sustained
attentional deployment.
701
The jittered ISI was followed by the four Gabor targets, which were presented for 100 ms and
tilted around a horizontal reference angle in CW or CCW directions. The tilt amount was titrated
to 82% baseline performance for each participant at each target location prior to scanning (see
Thresholding session for details). Target display was followed by another ISI of 500 ms. Finally,
a response cue, a central cross with one black arm, informed the participant of which target to

make the 2AFC orientation judgment on. The participant reported the direction of the tilt by
making a keyboard press.
709
Participants had about 1500 ms to make a response, and then received feedback. The exact
maximum response window varied from 1350-1700 ms to ensure that the trial ended
synchronously with the MRI volume acquisition. This trial to trial variation had no effect on
performance, as responses were much faster than the maximum window (typically about 300 to
600 ms after the response cue). Participants received feedback through a color change in
fixation cross from white to green for correct answers, to red for incorrect answers, and to yellow
for late answers that were not registered (<1% of trials across participants).
717
718   **Psychophysical protocols.** Participants completed four experimental protocols, three prior to
719   scanning and one during scanning: (1) familiarization with task, (2) thresholding, (3) attentional
720   cueing (outside the scanner), and (4) attentional cueing (during scanning). All four protocols
721   used the same trial structure, described above. During all protocols, participants were instructed
722   to emphasize accuracy over speed.
723
724   *1. Familiarization.* The familiarization protocol consisted of 30 trials, which were the same as
725   those in the main experiment except that the Gabor targets were at 100% contrast and the
726   orientation difference was large (±15º from vertical). The purpose was to ensure that
727   participants understood the task. This was verbally confirmed before advancing to the second
728   protocol.
729
2. Thresholding. The thresholding protocol was conducted with *neutral cues only* to estimate the
tilt deviation from the reference angle needed to reach threshold performance at each of the
four locations. Threshold was defined as the tilt deviation corresponding to 82% accuracy on a
fitted Weibull function. The threshold was found using the best PEST adaptive procedure of "3-
down-1-up", interleaved at each target location, where the tilt angle got smaller after three
consecutive accurate answers and got larger after one inaccurate answer
(palamedestoolbox.org) [82]. Because the thresholding experiment only contained neutral trials,
the target location was identified by the response cue. At the three other locations, the tilt angle
was determined by the current staircase value for the location, and the angular deviation
(clockwise or counterclockwise) was random. The thresholding session incorporated one
additional type of feedback informing the participants if their gaze deviation was ≥2º from the
fixation cross by a change in the color of the fixation cross from white to blue. The thresholding
comprised about 65 trials at each target location (~260 trials) and took about 30 min.
743
3. Attentional cueing outside the scanner. After the threshold measurements, and prior to
moving to the scanner, participants completed one full session of the main experiment (~1hr,
520 trials, including all 5 cueing conditions) at the tilt angles measured in the thresholding
experiment. This served as additional training for the participants, as well as assessment of the

protocol to verify that the cueing protocol affected performance as expected (improved
performance at the cued locations). This benefit ensured that participants were in fact using the
pre-cue.
751
4. Attentional cueing inside the scanner. Upon the completion of the three protocols outside the
scanner, participants took part in multiple scan sessions, performing the 2AFC orientation
discrimination task described above. During each scan session, there were 10 functional scans
in which participants performed the psychophysical task. Each of these scans had 52 trials and
lasted 4 min 14 s; with a total of 520 trials per scanning session. Seven participants completed
2080 trials in total (across 4 scan sessions), and two participants completed 1560 trials (3 scan
sessions).

MRI acquisition

Functional and anatomical MRI data were collected on a 3T Siemens MAGNETOM Prisma MRI
scanner (Siemens Medical Solutions, Erlangen, Germany) located at the Center for Brain Imaging
at NYU using a Siemens 64-channel head/neck coil. Before the functional echo-planar images
(EPIs) were obtained, two distortion scans were conducted in anterior-to-posterior (AP) and
posterior-to-anterior (PA) phase encoding directions to correct for distortions in functional EPI
scans due to susceptibility. After the distortion scans, in each scan session, ten scans of functional
EPIs were obtained using the CMRR multiband sequence with a multiband factor of 6[83-85]. Each
scan had 244 volumes (TR = 1000 ms, TE = 37.6 ms, voxel size: 2mm^3 isotropic, flip angle 68°)
[83,85]. For each participant, T1-weighted magnetization-prepared rapid gradient echo images
(MPRAGE) (TR = 2400 ms, TE = 2.2 ms, voxel size: 0.8mm^3 isotropic, flip angle 8°) were
previously acquired for previous studies, and were reused in the current study.

MRI preprocessing

All DICOMS were anonymized and defaced using *pydeface*
(https://github.com/poldracklab/pydeface) prior to transfer from the acquisition computer to the
researcher storage space. The DICOMS were then converted to NIFTIs and organized according
to the Brain Imaging Data Structure (BIDS) convention[86] using *heudiconv*
(https://github.com/nipy/heudiconv). The Anatomical and functional data were then preprocessed
using fMRIPrep v.20.2.1[87]. T1-weighted (T1w) anatomical images were corrected for intensity
non-uniformity and skull-stripped. Brain tissue was segmented into cerebrospinal fluid (CSF),
white-matter (WM) and gray matter (GM) using *fast* based on both T1w and T2w input images.
Segmented images were reconstructed using *recon-all*[88]. For the preprocessing of the functional
data, a reference volume and its skull-stripped version were generated and corrected for
susceptibility distortions using the two distortion maps collected with opposing phase encoding
directions (AP and PA). Corrected functional reference images were co-registered to the T1w
anatomical reference using *bbregister*[89]. Head-motion parameters with respect to the corrected
BOLD reference were estimated and the functional images were slice-time corrected using
3dTshift[90]. Slice-time corrected functional data was resampled to the anatomical space using one-
shot interpolations (incorporating transformation matrices, distortion maps and the coregistration).
Preprocessed BOLD data was then resampled onto each participant's reconstructed cortical

surface using *fsnative,* and all further analyses were performed on the *fsnative* space, using the
surface-based fMRI data.

Behavioral sensitivity analysis

Each participant's behavioral data were concatenated across all scan sessions. The effect of
attention on behavioral sensitivity was calculated across four target locations as a function of
attentional pre-cue validity. For each target location and pre-cue type, d′ sensitivity index was
calculated using *z(hit rate) - z(false alarm rate)*. To map our 2AFC procedure into a common
signal detection framework, we arbitrarily assigned one direction (clockwise) as 'target present'
and the other direction as 'target absent'[43,44,91]. To correct for extreme values, prior to computing
d′, we added 0.5 to the number of hits and false alarms[92, 93]. Although we emphasized accuracy
over speed to the participants in the instructions, we measured reaction time to assess speed-
accuracy tradeoffs, specifically, the possibility that conditions with higher accuracy had slower
reaction times. To calculate the 68% confidence interval, participant-level sensitivity and reaction
time values at each cue type and target location were computed 1000 times by resampling the
data with replacement (bootstrapping). With two-way repeated measures ANOVAs with location
(UVM x LVM x RHM x LHM) and cue type (Valid x Neutral x Invalid) factors we assessed the
differences among cue types both for sensitivity and reaction time.

Gaze position analysis

Because we were interested in the effect of endogenous covert attention (a top-down process), it
was critical that the stimuli were matched across retinotopic coordinates, and that participants
maintained central fixation. Outside the scanner, trials began once participants were fixating for
500 ms, and trials were aborted and repeated when fixation was broken (criterion of 2º). During
scanning, trials were scheduled to match the fMRI acquisition, and hence were never aborted or
delayed. However, fixation breaks were rare (2% of sampled frames across participants, 68% CI
= [1% - 4%]). We also analyzed the gaze position during scanning to determine whether there
were systematic biases in gaze direction across different attention conditions (the five cue types).
Gaze position was recorded at 1,000 Hz in X, Y screen pixels. We converted the gaze position
data from *X* and *Y* positions in screen pixels to degrees in visual angle. Time points with blinks
were identified by the EyeLink software and removed from analysis. For every trial, we then
averaged the X, Y position across time points from the trial onset to target display onset (1.6 or
2.6 s, depending on the duration of the mapping stimulus). These time-averaged values were
then averaged across all trials with the same pre-cue, yielding 5 *X*, *Y* pairs per participant. Finally,
to correct for calibration errors, we demeaned the values for each participant.

fMRI Data Analysis

823 Functional time series data of each vertex were analyzed in two steps[80]: First, a general linear
824 model (GLM) was solved on the concatenated trial responses across multiple scan sessions.
825 Second, a pRF model was fit to the estimated GLM weights. All analyses were performed on the
826 NYU High Performance Computer cluster using MATLAB (Version 9.14.0, R2023a).
827

General linear model. Functional time series of each vertex on each participant's native surface
were modeled with a GLM. Preprocessed functional time series were concatenated across
multiple scan sessions and modeled using
GLMdenoise (https://github.com/cvnlab/GLMdenoise)[94]. GLMdenoise is a powerful analysis
toolbox that *1)* identifies and regresses out task-irrelevant signals in time series data, *2)* optimizes
a hemodynamic response function for each participant, and *3)* estimates GLM weights with
bootstrapping to obtain confidence intervals and cross-validation to quantify model accuracy.
835
A design matrix was constructed for each run with 250 predictors (columns). Of these, 245
columns were for the mapping stimuli: 5 attention conditions x 49 mapping stimuli (48 bar locations
+ 1 blank). For these columns, there was a '1' for each TR when the mapping stimulus was present
(either 1 TR or 2 TRs, because the mapping stimuli were either 1 or 2 s). There were 5 additional
columns for the Gabor targets, one for each of the 5 attention conditions. These were modeled
as a '1' during the TR that the target appeared. The predictors did not take into account cue
validity because the validity only becomes apparent after the mapping stimulus and Gabor target
are viewed.
844

845 $$\hat{y}_i = \left(b_{M_1} x_{M_1} + \cdots b_{M_{245}} x_{M_{245}} + b_{T_1} x_{T_1} + \cdots b_{T_5} x_{T_5} \right) * HRF + N$$ *(Eq 1)*

846
where y_i represents the predicted time series of voxel *i*, the *b* terms are the coefficients ('beta
weights'), and the *x* terms are indicator variables (either 0 or 1). The subscripts denote the
condition, with M_1 to M_{245} corresponding to the mapping stimulus, and *T* corresponding to the
Gabor targets and *N* represents the nuisance variables (polynomial regressors for detrending low
frequency fMRI drift, and task-irrelevant time series regressors identified by GLMdenoise). All
variables other than the nuisance regressors were convolved with a hemodynamic response
function (*HRF)* estimated for each participant (a 50-s finite impulse response function). Estimated
coefficients from the GLM were converted to percent BOLD change by dividing them by the mean
signal intensity. For visualization purposes, estimated coefficients from the GLM representing the
responses to mapping stimulus bars were reorganized as a function of mapping stimulus position,
and convolved the measured responses with a triangular function that was padded with the first
and the last values of the data, and then normalized by the total sum[95].
859
pRF models. We solved separate pRF models for each of the five attention conditions, using the
49 mapping stimulus coefficients for that condition. We also solved a pRF model for the average
response across the five attention conditions by averaging the five coefficients for each of the 49
mapping stimuli. This resulted in six different preferred position and size estimates per vertex:
one for each of the five attention conditions (4 focal and 1 distributed) and one for the average
across attention conditions. For visualization purposes, the 49 coefficients per attention condition
were ordered into horizontal and vertical sweeps (plus a blank), so that the estimated BOLD
response and model predictions could be viewed as pseudo time series.
868
PRF analysis was conducted using *vistasoft* software (https://vistalab.stanford.edu/software,
Vista Lab, Stanford University), with a custom wrapper function

(https://github.com/WinawerLab/prfVista, New York University). Each vertex's pRF was modeled
as a circular 2D Gaussian, parameterized by its center (x, y in deg) and size (σ, one standard
deviation of the Gaussian, in deg)[81]. The pRF model predicts the response amplitude to each
mapping stimulus by pointwise multiplication of the Gaussian pRF and the binarized contrast
aperture, scaled by a nuisance factor (the vertex-specific gain). Unlike many implementations of
pRF models, our pRF model did not explicitly account for the hemodynamic response, low
frequency detrending, or conversion to percent BOLD change, because these steps were already
computed by the GLM.
879
The pRF fitting software estimates the pRF parameters (x, y, σ) by minimizing the residual sum
of squared between the measured BOLD pseudo-time series (the 49 coefficients from the GLM)
and the predictions from the model. We implemented a coarse-to-fine search for pRF fits for both
computational efficiency and to reduce the chance of model solutions that correspond to local
rather than global minima. In this procedure, the results from an initial brute force grid search
were used as the starting point for a finer second stage fitting procedure[74]. The coefficient of
determination, R_2, was calculated as 1 - (RSS / TSS), where *RSS* is the sum of squares of
residuals and *TSS* is the total sum of squares (i.e., the sum of squares of the data).
888
ROI definitions. We visualized retinotopic data by projecting each vertex's preferred position in
polar coordinates onto flattened maps of the visual cortex, centered at the occipital pole, using
Neuropythy v0.12.11 (https://github.com/noahbenson/neuropythy)[96]. Boundaries of V1, V2, V3,
hV4, V3A/B and LO1 were drawn manually on each participant's native MRI surface, guided by
the eccentricity, polar angle, and cortical curvature maps. The criteria for identifying map
boundaries follow prior work: V1-V3[97], hV4[98]; V3A/B and LO1[99]. We defined V3A and V3B as a
single ROI because the boundary separating them was not clear in every participant.
896
897   **Bootstrapping procedure.** We used bootstrapping procedures throughout this paper to estimate
898   confidence intervals and to test for the statistical significance of most of our results[100]. We sampled
899   eight participants with replacement 1000 times and calculated the 68% confidence interval based
900   on the bootstrapped population distribution. The bootstrapped distribution was compared against
901   a null distribution to test for statistical significance.
902
Latency analysis of attentional modulation of BOLD. For each visual field map in each
participant, we estimated the latency of attentional modulation, meaning the time during a trial at
which the BOLD response differed as a function of attentional condition. To estimate the latency,
we first extracted the BOLD time series of cortical ROIs representing the target locations. We
converted the time series of each vertex to percent BOLD change, and detrended the converted
data by projecting out a first order polynomial within each scan. We then extracted a 10-s event-
triggered time series for each trial, aligned either to the attentional cue ("cue-locked") or to the
Gabor target onset ("target locked"). These event-triggered time series were averaged across
trials in which the mapping stimulus bar overlapped the target corresponding to that cortical target
ROI. The time series were then also averaged across vertices within the target ROI. These
averages were computed separately for trials with 1-s bar stimuli and 2-s bar stimuli, and for trials
in which focal attention was directed to the target corresponding to the ROI ("attend in") and trials

in which focal attention was directed to one of the other targets ("attend out"). Next, we subtracted
the average BOLD time series for attend-out from attend-in trials to quantify the attentional
modulation over time.

To quantify the latency of attentional modulation, we used the latency analysis method previously
described[101]. We first normalized the event-triggered responses such that the minimum attentional
response is equal to 0 and the maximum is equal to 1. We next extracted the time point (TRs) at
which the response peaked and then separated the response curves as before and after the peak
point, representing the rising and falling of attentional responses. We fit logistic functions to the
two portions of the attentional response, separately to the 1-s and 2-s mapping stimulus trials of
cue- and target-locked responses, using the following equation:

$$f(t; m, t_{50}) = 1/ \left(1 + \exp(-m * (t - t_{50}))\right)$$

927   *(Eq 2)*

where *t* represents time in seconds (independent variable), *m* represents the slope and t50
represents the mean. On the best-fitting functions, we defined the latency of the attentional
response with a t10 parameter, the time point at which the attentional modulation reaches the
10% of the maximum response. We lastly subtracted the estimated latency of the rising time of
attentional modulation in 2-s mapping stimulus trials from 1-s mapping stimulus trials. The fitting
procedure was repeated 1,000 times on the group-level responses averaged across sub-samples
of participant-level responses with replacement.

2D visualizations of attentional activity. To investigate the effects of focal attention on BOLD
amplitude, we constructed visualizations of attention effects in visual space. We did this by first
constructing separate visual field representations of each visual area for each of the four focal
attention conditions for each participant. For a given participant and visual field map, we assigned
each vertex to its preferred visual field coordinates, as measured by the pRF center from the
dataset averaged across attention conditions. Next, for a given focal attention condition, at each
vertex we calculated the change in BOLD amplitude from distributed to focal attention conditions
estimated by the GLM, averaged across the 49 mapping stimuli, and plotted this average value
at the pRF location using a colormap. This produces a map of the visual field with, typically,
several hundred points (each point is a surface vertex plotted at its pRF center). These data points
were then resampled to a square grid using linear interpolation. Finally, to increase SNR, we
averaged across these 2D visual field plots across the four attention conditions by rotating each
one to align so that the attention direction was upward (90º).

Von Mises fitting procedure. To quantify the 2D visualizations of attentional activity, we reduced
the images to one spatial dimension, polar angle, constructing attentional tuning curves as a
function of polar angle. For these analyses, we included the vertices whose preferred eccentricity
was between 4° and 9° (close to the 6° eccentricity of the Gabor targets). For each of the target
locations, we binned vertices by their polar angle distance from the target in 20° steps from -180°
to 180°. Within each bin, we computed the attentional modulation as focal minus distributed
response, averaging across the 49 mapping stimuli per vertex, and across vertices within the bin.

958 The bin centers were computed as the average polar angle distance between the target location
959 and the vertices' pRF centers. These calculations produce one attentional tuning curve per target
960 location for each visual area and each participant. Because the tuning curves did not differ
961 systematically across the target locations, we averaged the four tuning curves per visual field
962 map, and fit the averaged tuning curve with a difference of two von Mises functions, using the
963 following equation using MATLAB's *besseli* function:
964

$$\hat{y} = [V_{\mu, \kappa_1}(x) - V_{\mu, \kappa_2}(x)] + B,$$

$$V(x; \mu, \kappa) = S * \frac{\exp{(\kappa * \cos(x - \mu))}}{2\pi * besseli(0, \kappa)}$$

965  *(Eq 3)*
966
967
968 where A is a scalar, B is an offset, is the Von Mises center (assumed to be the same for the
969 positive and negative von Mises), and 1 and 2 are the concentrations of the two von Mises
970 functions. These five parameters were fit for each visual field map in each participant by nonlinear
971 least squares using MATLAB's *fit* function. We defined the spread of attention in degrees of visual
972 angle as the distance at which the difference of von Mises functions intercept the x-axis, meaning
973 the transition point from attentional enhancement (focal > distributed) to attentional suppression
974 (distributed > focal). To define the x-intercepts on either side of the center, we solved for f(x) = 0
975 twice, at [-60, 0] and [0, 60] lower and upper limits for the solution space. Lastly, we calculated
976 the distance of two x-intercepts from each other to quantify the width along the x-axis. The group
977 level estimates were obtained by averaging the estimated values across participants.
978
979 **Definition of cortical target ROIs.** To quantify the retinotopic effects of attention, we created
980 wedge-ROIs on the cortical surface by extracting the portions of cortex that represent four target
981 locations. We used the averaged pRF solutions to extract a preferred visual field position for each
982 vertex. In V1, V2, V3, hV4, V3A/B and LO1, we extracted all vertex pRF with a preferred
983 eccentricity between 4° and 8°, and an angular preference of ±30° centered on each of the
984 cardinal meridians (*UVM:* 60° - 120°, *LVM:* 240° - 300°, *LHM:* 150° - 210°, *RHM:* 330° - 360° &
985 0° - 30°).
986
987 **Calculating the spatial shift effect.** We tested whether pRFs shifted towards an attentional
988 target compared to their preferred position under distributed attention. For each vertex assigned
989 to a visual map within a participant, we computed the changes in the best fitting pRF center across
990 attention conditions. First, for each vertex, we extracted the estimated pRF center in *x* and *y*
991 coordinates in degrees of visual angle, and the variance explained (*R2*) of each pRF model
992 estimated for four focal and the distributed attention conditions. We only included vertices which
993 met two criteria in all five pRF models: eccentricity in the range of 0.5° to 5.5° and variance
994 explained greater than 0.25. Next, for each vertex and each of the four targets, we subtracted the
995 distance between the distributed pRF center and target from the distance between focal pRF
996 center and target. A negative value indicates that the pRF center got closer to a target when the

target was attended. Last, we averaged the change in distance across vertices within an ROI and
across attentional conditions. This analysis produces a summary metric of spatial shift per
ROI. As a control analysis, we computed the same distance metric but on pRFs estimated from
shuffled GLM coefficients. For this analysis, we first randomly shuffled the focal attention cue on
each trial and created a shuffled design matrix. We then ran a GLM on this shuffled design matrix,
followed by five pRF models fitted to the estimated coefficients representing mapping stimulus
positions to estimate a pRF for each vertex for each attention condition.
1004
Directional vector graphs of spatial shifts. To investigate the pattern of spatial shifts within an
ROI, we implemented a method from previous research[17], by plotting the direction the average
pRF centers within bins for two opposite focal attention conditions. For pairs of opposite focal
conditions, we computed the average pRF position for each vertex (average of attend up/attend
down or attend left/attend right). We then binned the vertex pRFs based on their averaged
eccentricity (from 0.5° to 6.25° in steps of 1°) and polar angle (from 0° to 360° in steps of 45°)
estimates, and averaged the center position across each bin separately for the two targets. We
then plotted these two points for each bin connected by a line, and color coded the line by the
congruence of the pRF center shift direction based on the changes in *x* coordinates for the
horizontal targets, and in *y* coordinates for the vertical targets. If the pRF center shifts toward the
attentional target, the attend left condition should have an *x* value that is smaller (or more
negative) than the attend right condition (and vice versa for the attend right condition); and attend
down condition should result in a smaller (or more negative) *y* than attend up condition (and vice
versa for the attend up condition).
1019

1020
1021
1022   References
1023

1024 1. Desimone, R., and Duncan, J. (1995). Neural mechanisms of selective visual attention.
1025 Annual Review of Neuroscience 18, 193-222. 10.1146/annurev.ne.18.030195.001205.
1026
1027 2. Lennie, P. (2003). The cost of cortical computation. Current Biology, 6, 493-497.
1028 10.1016/s0960-9822(03)00135-0.
1029
1030 3. Carrasco, M. (2011). Visual attention: The past 25 years. Vision Research, 13, 1484-1525.
1031 10.1016/j.visres.2011.04.012.
1032
1033 4. Anton-Erxleben, K., and Carrasco, M. (2013). Attentional enhancement of spatial resolution:
1034 linking behavioural and neurophysiological evidence. Nat Rev Neurosci. 14(3), 188-200.
1035 10.1038/nrn3443.
1036
1037 5. Dosher, B.A., and Lu Z. L. (2000). Noise exclusion in spatial attention. Psychological Science,
1038 11(2), 139-46. 10.1111/1467-9280.00229.
1039

1040   6. Pestilli, F., and Carrasco, M. (2005). Attention enhances contrast sensitivity at cued and
1041   impairs it at uncued locations. Vision Research, 45(14), 1867. 10.1016/j.visres.2005.01.019.
1042

1043   7. Ling, S., Jehee, J.F.M., and Pestilli, F. (2014). A review of the mechanisms by which
1044   attentional feedback shapes visual selectivity. Brain Struct Funct, 220(3), 1237–1250.
1045   10.1007/s00429-014-0818-5.
1046

1047   8. Maunsell, J. (2015). Neuronal mechanisms of visual attention. Annu Rev Vis Sci, 1(1), 373-
1048   391. 10.1146/annurev-vision-082114-035431.
1049

1050   9. Kastner, S., de Weerd, P., Desimone, R., and Ungerleider, L.G. (1998). Mechanisms of
1051   directed dttention in the human extrastriate cortex as revealed by functional MRI. Science, 282,
1052   108-111. 10.1126/science.282.5386.108.
1053

1054   10. Reynolds, J.H., Chelazzi, L., and Desimone, R. (1999). Competitive mechanisms subserve
1055   attention in macaque areas V2 and V4. Journal of Neuroscience, 19(5), 1736-1753.
1056   10.1523/JNEUROSCI.19-05-01736.1999.
1057

1058   11. Tootel, R.B.H., Hadjikhani, N., Hall, K.E., Marrett, S., Vanduffel, W., Vaughan, T.J., and
1059   Dale, A. M. (1998). The retinotopy of visual spatial attention, Neuron, 21(6), 1409-22.
1060   10.1016/s0896-6273(00)80659-5.
1061

1062   12. Bloem, I.M., and Ling, S. (2019). Normalization governs attentional modulation within human
1063   visual cortex. Nature Communications, 10(1), 5660. 10.1038/s41467-019-13597-1.
1064

1065   13. Dugué, L., Merriam, E., Heeger, D., and Carrasco, M. (2020). Differential impact of
1066   endogenous and exogenous attention on activity in human visual cortex. Sci Rep, 10(1), 21274.
1067   10.1038/s41598-020-78172-x.
1068

1069   14. Womelsdorf, T., Anton-Erxleben, K., Pieper, F., and Treue, S. (2006). Dynamic shifts of
1070   visual receptive fields in cortical area MT by spatial attention. Nature Neuroscience, 9(9), 1156-
1071   60. 10.1038/nn1748.
1072

1073   15. Womelsdorf, T., Anton-Erxleben, K., and Treue, S. (2008). Receptive field shift and
1074   shrinkage in macaque middle temporal area through attentional gain modulation. Journal of
1075   Neuroscience, 28(36), 8934-44. 10.1523/JNEUROSCI.4030-07.2008.
1076

1077   16. Sprague T.C., and Serences, J.T. (2013). Attention modulates spatial priority maps in the
1078   human occipital, parietal and frontal cortices. Nat Neurosci. 16(12), 1879-87. 10.1038/nn.3574.
1079

1080   17. Klein, B.P., Harvey, B., and Dumoulin, S. (2014). Attraction of position preference by spatial
1081   attention throughout human visual cortex. Neuron, 84(1), 227-237.
1082   10.1016/j.neuron.2014.08.047.
1083

1084 18. Kay, K.N., Weiner, K.S., and Grill-Spector, K. (2015). Attention reduces spatial uncertainty in
1085 human ventral temporal cortex. Current Biology, 25(5), 595-600. 10.1016/j.cub.2014.12.050.
1086
1087 19. Vo, V.A., Sprague, T. C., and Serences. J. T. (2017). Spatial tuning shifts increase the
1088 discriminability and fidelity of population codes in visual cortex. Journal of Neuroscience, 37(12),
1089 3386-3401. 10.1523/JNEUROSCI.3484-16.2017.
1090
1091 20. Klein, B.P., Fracasso, A., Van Dijk, J.A., Paffen, C.L.E., Te Pas, S.F. and Dumoulin, S.O.
1092 (2018). Cortical depth dependent population receptive field attraction by spatial attention in
1093 human V1. NeuroImage, 176, 301-312. 10.1016/j.neuroimage.2018.04.055.
1094
1095 21. Kastner, S., Pinsk, M.A., de Weerd, P., Desimone, R., and Ungerleider, L.G. (1999).
1096 Increased activity in human visual cortex during directed attention in the absence of visual
1097 stimulation. Neuron, 22(4), 751-61. 10.1016/s0896-6273(00)80734-5.
1098
1099 22. Serences, J. T., Yantis, S., Culberson, A., and Awh, E. (2004). Preparatory activity in visual
1100 cortex indexes distractor suppression during covert spatial orienting. Journal of
1101 Neurophysiology, 92(6), 3538-45. 10.1152/jn.00435.2004.
1102
1103 23. Silver, M.A., Ress, D., and Heeger, D.J. (2006). Neural correlates of sustained spatial
1104 attention in human early visual cortex. Journal of Neurophysiology, 97(1), 229-237.
1105 10.1152/jn.00677.2006.
1106
1107 24. Capotosto, P., Babiloni, C., Romani, G.L., and Corbetta, M. (2009). Frontoparietal cortex
1108 controls spatial attention through modulation of anticipatory alpha rhythms. Journal of
1109 Neuroscience, 29(18), 5863-5872. 10.1523/JNEUROSCI.0539-09.2009.
1110
1111 25. Worden MS, Foxe JJ, Wang N, and Simpson GV. Anticipatory biasing of visuospatial
1112 attention indexed by retinotopically specific a-band electroencephalography increases over
1113 occipital cortex. Journal of Neuroscience, 95, 3844–3851. 10.1523/JNEUROSCI.20-06-
1114 j0002.2000.
1115
1116 26. Gould, I.C., Rushworth, M.F., and Nobre A.C. (2011). Indexing the graded allocation of
1117 visuospatial attention using anticipatory alpha oscillations. Journal of Neurophysiology, 105(3),
1118 1318-26. 10.1152/jn.00653.2010.
1119
1120 27. Zumer, J.M., Scheeringa, R., Schoffelen, J.M., Norris, D.G., and Jensen, O. (2014).
1121 Occipital alpha activity during stimulus processing gates the information flow to object-selective
1122 cortex. PLoS Biology, 12(10), e1001965. 10.1371/journal.pbio.1001965.
1123
1124 28. Ress, D, Backus, B.T, and Heeger, D.J. (2000). Activity in primary visual cortex predicts
1125 performance in a visual detection task. Nature Neuroscience, 3, 940-5. 10.1038/78856.
1126

1127 29. Sapir, A, Giovanni, A., McAvoy, M., Shulman, G.L., and Corbetta, M. (2005). Brain signals
1128 for spatial attention predict performance in a motion discrimination task. Proceedings of the
1129 National Academy of Sciences, 102(49), 17810-5. 10.1073/pnas.0504678102.
1130
1131 30. Giesbrecht, B., Weissman, D.H., Woldorff, M.G., and Mangun, G.R. (2006). Pre-target
1132 activity in visual cortex predicts behavioral performance on spatial and feature attention tasks.
1133 Brain Research, 1080(1), 63-72. 10.1016/j.brainres.2005.09.068.
1134
1135 31. Stokes, M., Thompson, R., Nobre, A.C., and Duncan, J. (2009). Shape-specific preparatory
1136 activity mediates attention to targets in human visual cortex. Proceedings of the National
1137 Academy of Sciences, 106(46), 19569-74. 10.1073/pnas.0905306106.
1138
1139 32. Fannon, S.P., Saron, C.D., and Mangun, G.R. (2008). Baseline shifts do not predict
1140 attentional modulation of target processing during feature-based visual attention.
1141 Front Hum Neurosci, 1, 7. 10.3389/neuro.09.007.2007.
1142
1143 33. Fox, K.J., Birman, D., and Gardner, J.L. (2023). Gain, not concomitant changes in spatial
1144 receptive field properties, improves task performance in a neural network attention model. eLife,
1145 12:e78392. 10.7554/eLife.78392.
1146
34. Snyder, A.C., Yu, B.M., and Smith, M.A. (2018). Distinct population codes for attention in the
absence and presence of visual stimulation. Nature Communications, 9(1), 4382.
10.1038/s41467-018-06754-5.
35. Carrasco, M., Talgar, C.P., and Cameron, E.L. (2001). Characterizing visual performance
fields: effects of transient covert attention, spatial frequency, eccentricity, task and set size.
Spatial Vision, 15(1), 61-75. 10.1163/15685680152692015.
1153
1154 36. Himmelberg, M.M., Winawer, J., and Carrasco, M. (2020). Stimulus-dependent contrast
1155 sensitivity asymmetries around the visual field. Journal of Vision, 20(9), 1-19.
1156 10.1167/jov.20.9.18.
1157
1158 37. Barbot, A., Xue, S., and Carrasco, M. (2021). Asymmetries in visual acuity around the visual
1159 field. Journal of Vision, 21(1):2, 1-23. 10.1167/jov.21.1.2.
1160
1161 38. Lee, S., Blake, R., and Heeger, D.J. (2004). Traveling waves of activity in primary visual
1162 cortex during binocular rivalry. Nature Neuroscience, 8(1), 22-23. 10.1038/nn1365.
1163
1164 39. Stoll, S., Infanti, E., De Haas, B., and Schwarzkopf, D.S. (2022). Pitfalls in post hoc
1165 analyses of population receptive field data. NeuroImage, 8, 263, 119557.
1166 10.1016/j.neuroimage.2022.119557.
1167
1168 40. Lerma-Usabiaga, G., Benson, N., Winawer, J., and Wandell, B. A. (2020). A validation
1169 framework for neuroimaging software: The case of population receptive fields. PLoS

1170   computational biology, 16(6), e1007924.

1171

1172   41. Posner, M.I. (1980). Orienting of attention. The Quarterly Journal of Experimental
1173   Psychology, 32(1), 3-25. 10.1080/00335558008248231.

1174

1175   42. Pestilli, F., Ling, S., and Carrasco, M. (2009). A population-coding model of attention's
1176   influence on contrast response: Estimating neural effects from psychophysical data. Vision
1177   Research, 49(10), 1144–1153. 10.1016/j.visres.2008.09.018.

1178

1179   43. Herrmann, K., Montaser-Kouhsari, L., Carrasco, M., and Heeger, D.J. (2010). When size
1180   matters: attention affects performance by contrast or response gain. Nature Neuroscience,
1181   13(12), 1554-1559. 10.1038/nn.2669.

1182

1183   44. Fernández, A., Hanning, N.M., and Carrasco, M. (2023). Transcranial magnetic stimulation
1184   to frontal but not occipital cortex disrupts endogenous attention. Proceedings of the National
1185   Academy of Sciences, 120(10), e2219635120. 10.1073/pnas.2219635120.

1186

1187   45. Reynolds, J.H., and Heeger. D.J. (2009). The normalization model of attention. Neuron,
1188   61(2), 168-185. 10.1016/j.neuron.2009.01.002.

1189

46. Itthipuripat, S., Sprague, T.C., and Serences, J.T. (2019). Functional MRI and EEG index
complementary attentional modulations. Journal of Neuroscience, 39(31), 6162–6179.
10.1523/JNEUROSCI.2519-18.2019.
47. Zanto, T.P., Chadick, J.Z., and Gazzaley, A. (2013). Anticipatory alpha phase influences
visual working memory performance. NeuroImage, 85, 794-802.
10.1016/j.neuroimage.2013.07.048.

1196

1197   48. Kok, P., Jehee, J.F.M., and de Lange, F.P. (2012). Less is more: Expectation sharpens
1198   representations in the primary visual cortex. Neuron, 75(2), 265–270.
1199   10.1016/j.neuron.2012.04.034.

1200

1201   49. Kok, P., Mostert, P., and de Lange, F.P. (2017). Prior expectations induce prestimulus
1202   sensory templates. Proceedings of the National Academy of Sciences, 114(39), 10473–10478.
1203   10.1016/j.neuron.2012.04.034.

1204

1205   50. Summerfield, C., and Egner, T. (2009). Expectation (and attention) in visual cognition.
1206   Trends in Cognitive Sciences, 13(9), 403-9. 10.1016/j.tics.2009.06.003.

1207

1208   51. Bubic, A., von Cramon, Y.D., and Schubotz, R.I. (2010). Prediction, cognition and the brain.
1209   Front. Hum. Neurosci, 4, 25. 10.3389/fnhum.2010.00025.

1210

1211   52. Williford, T., and Maunsell, J.H.R. (2006). Effects of spatial attention on contrast response
1212   functions in macaque area V4. Journal of Neurophysiology, 96(1), 40-54.

1213   10.1152/jn.01207.2005.

1214

1215   53. Thiele, A., Pooresmaeili, A., Delicato, L.S., Herrero, J.H., and Roelfsema, P.R. (2009).
1216   Additive effects of attention and stimulus contrast in primary visual cortex. Cerebral Cortex,
1217   19(12), 2970-81. 10.1093/cercor/bhp070.

1218

1219   54. McAdams, C.J., and Maunsell, J.H.R. (1999). Effects of attention on orientation-tuning
1220   functions of single neurons in macaque cortical area V4. Journal of Neuroscience, 19(1), 431-
1221   41. 10.1523/JNEUROSCI.19-01-00431.1999.

1222

1223   55. Treue, S, and Martínez-Trujillo, J.C. (1999). Feature-based attention influences motion
1224   processing gain in macaque visual cortex. Nature, 399(6736), 575–579. 10.1038/21176.

1225

1226   56. Rabinowitz, N.C., Goris, R.L., Cohen, M., and Simoncelli, E.P. (2015). Attention stabilizes
1227   the shared gain of V4 populations. eLife, 4:e08998. 10.7554/eLife.08998.

1228

1229   57. Ruff, D.A., and Cohen, M.R. (2014). Attention can either increase or decrease spike count
1230   correlations in visual cortex. Nature Neuroscience, 17(11), 1591–1597. 10.1038/nn.3835.

1231

1232   58. Buracas, G.T. and Boynton, G.M. (2007). The effect of spatial attention on contrast
1233   response functions in human visual cortex. Journal of Neuroscience, 27(1), 93-97.
1234   10.1523/JNEUROSCI.3162-06.2007.
1235   59. Murray, S.O. (2008). The effects of spatial attention in early human visual cortex are
1236   stimulus independent. Journal of Vision, 8(10):2, 1-11. 10.1167/8.10.2.

1237

1238   60. Pestilli, F., Carrasco, M., Heeger, D.J., and Gardner, J.L. (2011). Attentional enhancement
1239   via selection and pooling of early sensory responses in human visual cortex. Neuron, 72(5),
1240   832-46. 10.1016/j.neuron.2011.09.025.

1241

1242   61. Li, X., Lu, Z., Tjan, B.S., Dosher, B.A., and Chu, W. (2008). Blood oxygenation level-
1243   dependent contrast response functions identify mechanisms of covert attention in early visual
1244   areas. Proceedings of the National Academy of Sciences, 105(16), 6202-7.
1245   10.1073/pnas.0801390105.

1246

1247   62. Itthipuripat, S, Ester, E.F., Deering, S., and Serences, J.T. (2014). Sensory gain outperforms
1248   efficient readout mechanisms in predicting attention-related improvements in behavior. Journal
1249   of Neuroscience, 34(40), 13384-98. 10.1523/JNEUROSCI.2277-14.2014.

1250

1251   63. Foster, J.J., Thyer, W., Wennberg, J.A., and Awh, E. (2021). Covert attention increases the
1252   gain of stimulus-evoked population codes. Journal of Neuroscience, 41(8), 1802-15.
1253   10.1523/JNEUROSCI.2186-20.2020.

1254

1255   64. Hara, Y., Pestilli, F., and Gardner, J.L. (2014). Differing effects of attention in single-units
1256   and populations are well predicted by heterogeneous tuning and the normalization model of

1257   attention. Front. Comput. Neurosci, 8, 12. 10.3389/fncom.2014.00012.
1258
1259   65. Foster, J.J., and Ling, S. (2022). Feature-based attention multiplicatively scales the fMRI-
1260   BOLD contrast-response function. Journal of Neuroscience, 42(36), 6894-906.
1261   10.1523/JNEUROSCI.0513-22.2022.
1262
1263   66. Carrasco, M, Penpeci-Talgar, C., and Eckstein, M. (2000). Spatial covert attention increases
1264   contrast sensitivity across the CSF: support for signal enhancement. Vision Research, 40(10-
1265   12), 1203–1215. 10.1016/s0042-6989(00)00024-9.
1266
1267   67. Suzuki, M., and Gottlieb, J. (2012). Distinct neural mechanisms of distractor suppression in
1268   the frontal and parietal lobe. Nature Neuroscience, 16(1), 98-104. 10.1038/nn.3282.
1269
1270   68. Noonan, M.P., Adamian, N., Pike, A., Printzlau, F., Crittenden, B.M., Stokes M. (2016).
1271   Distinct mechanisms for distractor suppression and target facilitation. Journal of Neuroscience,
1272   36(6), 1797-807. 10.1523/JNEUROSCI.2133-15.2016.
1273
69. Schneider, D., Herbst, S.K., Klatt, L.I., and Wöstmann, M. (2021). Target enhancement or
distractor suppression? Functionally distinct alpha oscillations form the basis of attention. Eur J
of Neuroscience, 55(11-12), 3256-3265. 10.1111/ejn.15309.
70. Himmelberg, M.M., Winawer, J., and Carrasco, M. (2023). Polar angle asymmetries in visual
perception and neural architecture. Trends in Neurosciences, 46(6), 445-458.
10.1016/j.tins.2023.03.006.
1280
1281   71. Kupers, E.R., Carrasco, M., and Winawer, J. (2019). Modeling visual performance
1282   differences 'around' the visual field: A computational observer approach. PLoS Computational
1283   Biology, 15(5), e1007063. 10.1371/journal.pcbi.1007063.
1284
1285   72. Kupers, E.R., Benson, N.C., Carrasco, M., and Winawer, J. (2022). Asymmetries around the
1286   visual field: From retina to cortex to behavior. PLoS Computational Biology, 18(1), e1009771.
1287   10.1371/journal.pcbi.1009771.
1288
1289   73. Liu, T., Heeger, D.J., and Carrasco, M. (2006). Neural correlates of the visual vertical
1290   meridian asymmetry. Journal of Vision, 6(11), 1294-1306. 10.1167/5.1.1.
1291
1292   74. Benson, N.C., Kupers, E.R., Barbot, A., Carrasco, M., and Winawer. J. (2021). Cortical
1293   magnification in human visual cortex parallels task performance around the visual field. eLife,
1294   10, e67685. 10.7554/eLife.67685.
1295
1296   75. Himmelberg, M.M., Winawer, J., and Carrasco, M. (2022). Linking individual differences in
1297   human primary visual cortex to contrast sensitivity around the visual field. Nature
1298   Communications, 13(1), 3309. 10.1038/s41467-022-31041-9.
1299

76. Himmelberg, M.M., Tünçok, E., Gomez, J., Grill-Spector, K., Carrasco, M., and Winawer, J.
(2023). Comparing retinotopic maps of children and adults reveals a late-stage change in how
V1 samples the visual field. Nature Communications, 14(1), 1561. 10.1038/s41467-023-37280-
8.

77. Purokayastha, S., Roberts, M., and Carrasco, M. (2021). Voluntary attention improves
performance similarly around the visual field. Atten Percept Psychophys, 83(7), 2784–2794.
10.3758/s13414-021-02316-y.

78. Kleiner, M., Brainard, D., and Pelli, D. (2007). What's new in Psychtoolbox-3?.

79. Kay, K.N., Winawer, J., Rokem, A., Mezer, A., and Wandell, B.A. (2013). A two-stage
cascade model of BOLD responses in human visual cortex. PLoS Comput Biol, 9(5), e1003079.
10.1371/journal.pcbi.1003079.

80. Kay, K.N., Winawer, J., Mezer, A., and Wandell, B.A. (2013). Compressive spatial
summation in human visual cortex. Journal of Neurophysiology, 110(2), 481-494.
10.1152/jn.00105.2013.

81. Dumoulin, S.O., and Wandell, B.A. (2007). Population receptive field estimates in human
visual cortex. NeuroImage, 39(2), 647-60. 10.1016/j.neuroimage.2007.09.034.

82. Prins, N., and Kingdom, F.A.A. (2018). Applying the model-comparison approach to test
specific research hypotheses in psychophysical research using the Palamedes Toolbox.
Frontiers in Psychology, 9, 1250. 10.3389/fpsyg.2018.01250.

83. Feinberg, D.A., Moeller, S., Smith, S.M., Auerbach, E., Ramanna, S., Glasser, M.F., Miller,
K.L., Ugurbil, K., and Yacoub, E. (2010). Multiplexed echo planar imaging for sub-second whole
brain fMRI and fast diffusion imaging. PLoS ONE, 6(9), e15710. 10.1371/journal.pone.0015710.

84. Moeller, S., Yacoub, E., Olman, C.A., Auerbach, E., Strupp, J., Harel, N., and Ugurbil, K.
(2010). Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel
imaging with application to high spatial and temporal whole-brain fMRI. Magnetic Resonance in
Medicine, 63(5), 1144-1153. 10.1002/mrm.22361.

85. Xu, J., Moeller, S., Auerbach, E.J., Strupp, J., Smith, S.M., Feinberg, D.A., Yacoub, E., and
Ugurbil, K. (2013). Evaluation of slice accelerations using multiband echo planar imaging at 3T.
NeuroImage, 83, 991-1001. 10.1016/j.neuroimage.2013.07.055.

86. Gorgolewski, K.J., Auer, T., Calhoun, V.D., Craddock, R.C., Das, S., Duff, E.P., … Poldrack,
R. A. (2016). The brain imaging data structure, a format for organizing and describing outputs of
neuroimaging experiments. Sci Data, 3(1), 160044. 10.1038/sdata.2016.44.
1342

1343   87. Esteban, O., Markiewicz, C.J., Blair, R.W., Moodie, C.A., Isik, A.I., Erramuzpe, A., …
1344   Gorgolewski, K. J. (2018). fMRIPrep: a robust preprocessing pipeline for functional MRI. Nature
1345   Methods, 16(1), 111-116. 10.1038/s41592-018-0235-4.
1346
1347   88. Dale, A.M., Fischl, B., and Sereno, M.I. (1999). Cortical surface-based analysis: I.
1348   Segmentation and surface reconstruction. NeuroImage, 9(2), 179-194.
1349   10.1006/nimg.1998.0395.
1350
1351   89. Greve, D.N., and Fischl, B. (2009). Accurate and robust brain image alignment using
1352   boundary-based registration. NeuroImage, 48(1), 63-72. 10.1016/j.neuroimage.2009.06.060.
1353
1354   90. Cox, R.W., and Hyde, J.S. (1997). Software tools for analysis and visualization of fMRI data.
1355   NMR Biomed, 10(4-5), 171-8. 10.1002/(sici)1099-1492(199706/08)10:4/5<171::aid-
1356   nbm453>3.0.co;2-l.
1357
1358   91. Recht, S., Mamassian, P., and de Gardelle, V. (2019). Temporal attention causes
1359   systematic biases in visual confidence. Sci Rep, 9(1), 11622. 10.1038/s41598-019-48063-x.
1360
1361   92. Hautus, M.J. (1995). Corrections for extreme proportions and their biasing effects on
1362   estimated values of d'. Behavior Research Methods, Instruments, & Computers, 27(1), 46–51.
1363   10.3758/BF03203619.
1364
1365   93. Brown, G.S., and White, K.G. (2005). The optimal correction for estimating extreme
1366   discriminability. Behavior Research Methods, 37(3), 436-49. 10.3758/BF03192712.
1367
1368   94. Kay, K.N., Rokem, A., Winawer, J., Dougherty, R.F., and Wandell, B.A. (2013).
1369   GLMdenoise: a fast, automated technique for denoising task-based fMRI data. Frontiers in
1370   Neuroscience, 7, 247. 10.3389/fnins.2013.00247.
1371
1372   95. Parent R. (2012). Computer Animation: Algorithms and Techniques (Morgan Kauffman)
1373
1374   96. Benson, N.C., and Winawer, J. (2018). Bayesian analysis of retinotopic maps. eLife, 7,
1375   e40224. 10.7554/eLife.40224.
1376
1377   97. Benson, N.C., Yoon, J.M.D., Forenzo, D., Engel, S.A., Kay, K.N., and Winawer, J. (2022).
1378   Variability of the surface area of the V1, V2, and V3 maps in a large sample of human
1379   observers. Journal of Neuroscience, 42(46), 8629-8646. 10.1523/JNEUROSCI.0690-21.2022.
1380
1381   98. Winawer, J., and Witthoft, N. (2015). Human V4 and ventral occipital retinotopic maps. Vis
1382   Neurosci, 32, E020. 10.1017/S0952523815000176.
1383
1384   99. Larsson, J., and Heeger, D.J. (2006). Two retinotopic visual areas in human lateral occipital
1385   cortex. Journal of Neuroscience, 26(51), 13128-13142. 10.1523/JNEUROSCI.1657-06.2006.
1386

1387   100. Efron, B. (1981). Non parametric estimates of standard Error: The jackknife, the bootstrap
1388   and other methods. Biometrika, 68(3), 589–599. 10.2307/2335441.
1389
1390   101. Sit, Y.F., Chen, Y., Geisler, W.S., Miikkulainen, R., and Seidemann, E. (2009). Complex
1391   dynamics of V1 population responses explained by a simple gain-control model. Neuron, 64(6),
1392   778-80.