Thymic dendritic cell-derived IL-27p28 promotes the establishment of functional bias against IFN-γ production in newly generated CD4[+] T cells through STAT1-related epigenetic mechanisms

Jie Zhang[1], Hui Tang[1], Haoming Wu[1], Xuewen Pang[1], Rong Jin[1*], Yu Zhang[1,2*]

[1] Department of Immunology, School of Basic Medical Sciences, NHC Key Laboratory of Medical Immunology, Medicine Innovation Center for Fundamental Research on Major Immunology-related Diseases, Peking University, Beijing, China

2 Institute of Life Sciences, Jinzhou Medical University, Jinzhou, Liaoning Province, China

Corresponding Author

Yu Zhang, zhangyu007@bjmu.edu.cn

Rong Jin, jinrong@bjmu.edu.cn

Keywords: IL-27p28, single positive thymocyte, Th2 bias, IFN-γ, autoimmune disease

Abstract

The newly generated CD4 single-positive (SP) T lymphocytes are featured by enhanced IL-4 but repressed IFN-γ production. The mechanisms underlying this functional bias remain elusive. Previous studies have reported that CD4$^+$ T cells from mice harboring DC-specific deletion of IL-27p28 display an increased capacity of IFN-γ production upon TCR stimulation. Here we demonstrated that similarly altered functionality occurred in CD4SP thymocytes, recent thymic emigrants (RTEs) as well as naive T cells from either *Cd11c-p28$^{f/f}$* mice or mice deficient in the α subunit of IL-27 receptor. Therefore, DC-derived IL-27p28-triggered, IL-27Rα-mediated signal is critically involved in the establishment of functional bias against IFN-γ production during their development in the thymus. Epigenetic analyses indicated reduced DNA methylation of the *Ifng* locus and increased trimethylation of H3K4 at both *Ifng* and *Tbx21* loci in CD4SP thymcoytes from *Cd11c-p28$^{f/f}$* mice. Transcriptome profiling demonstrated that *Il27p28* ablation resulted in coordinated up-regulation of STAT1-activated genes. Concurrently, STAT1 was found to be constitutively activated. Moreover, we observed increased accumulation of STAT1 at the *Ifng* and *Tbx21* loci and a strong correlation between STAT1 binding and H3K4me3 modification of these loci. Of note, *Il27p28* deficiency exacerbated the autoimmune phenotype of *Aire$^{-/-}$* mice. Collectively, this study reveals a novel mechanism underlying the functional bias of newly generated CD4$^+$ T cells and the potential relevance of such a bias in autoimmunity.

41   Graphic abstract

Introduction

The thymus plays a crucial role in the phenotypic development and functional maturation of T cells. After positive and negative selection, T cell progenitors become CD4 or CD8 single positive (CD4SP or CD8SP) thymocytes that are self-tolerant. These SP thymocytes then emigrate to the periphery, known as recent thymic emigrants (RTEs), and 2-3 weeks later, RTEs develop into mature naive T cells (*Ashby & Hogquist, 2023*). Upon activation, CD4SP thymocytes and RTEs exhibit a reduced proliferation capacity and decreased production of interleukin-2 (IL-2), interferon-γ (IFN-γ), and tumor necrosis factor α (TNF-α) compared to mature naive T cells. However, they show increased secretion of IL-4, which is known as T helper (Th) 2 bias in immature T cells (*Hendricks & Fink, 2011*). This Th2 bias in CD4SP thymocytes and RTEs is thought to assist newly emigrated self-reactive T cells to respond appropriately to self-antigens (*Fink & Hendricks, 2011*). Furthermore, the low level of IFN-γ production in RTEs renders them more prone to becoming regulatory T cells, which also helps to prevent the induction of autoimmune diseases (*Bhaumik et al., 2013*). The low level of DNA methylation at the *Il4* locus and reduced expression of *Dnmts* contribute to the Th2 bias in CD4SP thymocytes and RTEs (*Berkley et al., 2013*). However, the mechanism behind repressed IFN-γ expression in favor of the Th2 bias in these immature T cells is elusive.

The heterodimeric cytokine IL-27 is composed of IL-27p28 and Epstein-Barr virus-induced gene 3 (EBI3), which signals through the IL-27R consisting of the IL-27Rα (also named as WSX-1, TCCR) and the gp130 subunits. Elevated levels of IL-27 are associated with the pathogenesis of autoimmune diseases such as ankylosing

spondylitis (AS) (*Lin et al., 2015*), behcet's disease (BD) (*Shen et al., 2013*), and

experimental arthritis (*Cao et al., 2008*). The IL-27-regulated differentiation and

function of peripheral CD4[+] T cells subsets Th1, Th17 and Treg have been proposed as

an explanation for the pathogenesis of autoimmune disease (*Mei et al., 2021*), however,

conflicting reports have been demonstrated in different settings. For example, it is

consistent that IL-27 inhibits Th17 differentiation and function *in vitro* and *in vivo* in

Il27p28[-/-] and *Ebi3[-/-]* experimental autoimmune encephalomyelitis (EAE) mice models

(*Diveu et al., 2009; Kim et al., 2019; Liu et al., 2012*). Although IL-27 inhibits Treg

differentiation *in vitro* (*Huber et al., 2008; Neufert et al., 2007*), and *Ebi3[-/-]* EAE models

show increased numbers and suppressive functions of Tregs (*Liu et al., 2012*), Tregs in

Il27ra[−/−] and Treg-specific *Il27ra[−/−]* mice loss the suppressive function (*Kim et al., 2019;*

Nguyen et al., 2019; Y. J. Park et al., 2019). Furthermore, IL-27 was initially found to

promote Th1 differentiation of naive CD4[+] T cells under physiological conditions

(*Pflanz et al., 2002*). However, naive CD4[+] T cells derived from *Ebi3[-/-]* and *Il27p28[-/-]*

mice exhibit increased production of IFN-γ. In addition, the absence of *Il27ra[-/-]*

promotes pathological Th1 accumulation following *Trypanosoma cruzi* infection

(*Hamano et al., 2003; Pflanz et al., 2002; Villarino et al., 2003*). A subsequent study

demonstrates that both the absence and overexpression of IL-27p28 lead to increased

susceptibility to *T. gondii* infection, with IL-27p28 attributed as an independent

negative regulator of IFN-γ production in T cells (*J. Park et al., 2019*). Notably,

deficiency in gp130 does not alter IFN-γ production in CD4[+] T cells in a virus-infection

model (*Harker et al., 2015*). These results indicate the intricate functionality of each

subunit of IL-27 and IL 27R on peripheral CD4[+] T cells, which is insufficient to fully

explain the pathogenesis of autoimmune disease.

The structural analysis of IL-27 has demonstrated that IL-27p28, serving as a

central subunit, possesses the ability to bind with EBI3, as well as the two receptor

subunits IL-27Rα and gp130 (*Caveney et al., 2022*). Previous studies have reported IL-

27p28 as an independent antagonist of gp130-mediated signaling (*Stumhofer et al.,*

2010). In a ConA-induced liver damage model, CD4[+] T cells from mice with dendritic

cell (DC)-specific deletion of p28 (*Cd11c-p28*[f/f] mice) exhibit a propensity for rapid and

robust production of IFN-γ upon activation. This observation is attributed to the absence

of IL-27p28 in the thymic environment of these mice (*Zhang et al., 2013*). These

findings raise the intriguing possibility that thymic DC-derived IL-27p28 may

contribute to shaping the Th1/Th2 balance of newly generated CD4SP thymocytes.

In this paper, we have demonstrated that DC-derived IL-27p28 plays a crucial role in

maintaining the Th2 bias of CD4SP thymocytes. Our findings suggest that IL-27p28

regulates the DNA methylation status and histone modification of transcription

regulatory regions of *Ifng* and *Tbx21* genes. These epigenetic modifications were

further modulated by the low level of basal and phosphorylated signal transducer and

activator of transcription 1 (STAT1) protein. Furthermore, we observed that the

disruption of Th2 bias in CD4SP thymocytes due to IL-27p28 deficiency exacerbated

autoimmune diseases in *Aire*[-/-] mice. Our results provide insights into the molecular

mechanisms underlying the regulation of Th2 bias by DC-derived IL-27p28 and its

potential role in autoimmune diseases.

Results

1. DC-specific deletion of IL-27p28 endows the newly generated CD4$^+$ T cells with inherently enhanced capacity of IFN-γ production

Although IL-27 is widely reported to promote peripheral Th1 differentiation *in vitro*, CD4$^+$ T cells from *Cd11c-p28$^{f/f}$* mice exhibit enhanced IFN-γ production, a capacity seemingly acquired in their development in the thymus (*Zhang et al., 2013*). To investigate the possibility that thymic DC-derived IL-27 shapes the functionality of newly generated CD4$^+$ T cells, CD4SP thymocytes, CD4$^+$ RTEs, and CD4$^+$ naive T cells were purified from *Cd11c-p28$^{f/f}$* (cKO) mice and their wild-type (WT) littermates harboring a Rag2-GFP transgene, which allows easy tracking of newly generated thymocytes and RTEs (*Ashby & Hogquist, 2023*). The purified cells were then compared for cytokine expression in response to anti-CD3 and anti-CD28 stimulation under non-polarizing conditions. As expected, CD4$^+$ naive T cells from cKO mice exhibited a significantly higher level of *Ifng* mRNA expression compared to WT cells (Figure 1A). Intracellular staining (Figure 1B) and ELISA assay of the culture supernatant (Figure 1C) confirmed enhanced IFN-γ production at the protein level. Notably, this difference was already detectable at the SP thymocyte stage, indicating that it is developmentally regulated. Another prominent feature of CD4SP thymocytes is the hyperproduction of Th2 cytokines (*Fink & Hendricks, 2011; Makar et al., 2003*). Despite the profound impact of IL-27p28 deficiency on IFN-γ production, we observed no significant difference in IL-4 expression between cKO and WT T cells (Figure 1A and C). Moreover, there was no significant difference in the expression of IL-2 and

TNF-α between cKO and WT T cells (Figure 1A and Figure 1-figure supplement 1).

These results suggest that DC-specific deletion of IL-27p28 specifically enhances IFN-γ production in newly generated T cells without affecting Th2 cytokines.

Next, we investigated the expression of T-box transcription factor (T-bet) and GATA3, which are the master regulators of Th1 and Th2 differentiation, respectively (*Fang et al., 2022*). The differential impact of IL-27p28 deficiency on IFN-γ and IL-4 production was accompanied by an increase in Tbx21 expression but no significant alteration in Gata3 expression (Figure 1D). Western blotting confirmed the upregulation of T-bet protein expression (Figure 1E). Furthermore, we extended our analyses onto CD4$^+$ T cells at steady state or under polarizing conditions. Higher basal levels of *Ifng* and *Tbx21* transcripts were detected in CD4SP thymocytes and naive T cells from cKO mice than the counterparts from WT mice even without stimulation (Figure 1F), indicating the constitutive activation of these loci. When cultured under polarizing conditions, these cells were effectively induced to differentiate into Th1, Th2, Th17 and Treg lineages. In contrast to the previous report of different differentiation potentials of RTEs versus naive T cells (*Hendricks & Fink, 2011*), CD4SP thymocytes, RTEs and naive T cells showed similar capacity to differentiate into effector cells. Moreover, cKO and WT cells were equally positioned for the differentiation (Figure 1-figure supplement 2A-F), suggesting that IL-27p28 deficiency does not affect effector cell differentiation under optimal conditions. Taken together, these data support that IL-27p28 is actively involved in the functional maturation of developing CD4SP thymocytes, biasing them away from the Th1 lineage.

2. *Il27ra*-deficient CD4⁺ T cells are similarly predisposed for potent IFN-γ production

Apart from acting as a subunit of IL-27, IL-27p28 have been reported to possess activities independent of IL-27, by acting on its own or coupling with other proteins, such as IL-Y (p28/p40, binding to IL-27Rα/IL-12Rβ1 to activate STAT3) (*Flores et al., 2015*) and CLY (p28/CLF, binding to gp130/IL-6Rα to activate STAT1 and STAT3) (*Tormo et al., 2013*). To ascertain whether the phenotype in the *Cd11c-p28^(f/f)* mice results from defective IL-27Rα signaling, we investigated the functional characters of T lineage cells in mice deficient in *Il27ra*. Reminiscent to what was observed in *Il27p28*-deficient mice, increased expression of *Ifng* and *Tbx21* were detected in *Il27ra^(-/-)* CD4SP thymocytes and naive CD4⁺ T cells, whereas *Il4*, *Il2,* and *Gata3* transcripts were equally present in WT and *Il27ra^(-/-)* cells (Figure 2A). Intracellular staining confirmed the enhanced IFN-γ production by *Il27ra*-deficient cells stimulated under non-polarizing conditions (Figure 2B). IFN-γ production by *Il27ra^(-/-)* and WT cells, however, was comparable under Th1 polarizing conditions (Figure 1-figure supplement 2G). The similar phenotype shared by *Il27p28* and *Il27ra* knockout mice indicates that the altered T cell functionality in the absence of IL-27p28 is indeed a reflection of disrupted IL-27Rα signaling.

3. IL-27p28 deficiency induces permissive epigenetic changes at *Ifng* and *Tbx21* loci

175 Developmentally regulated transcriptional activation and repression of genes or

176 cell-type specific expression patterns are largely achieved by modifying the chromatin

177 template at a gene locus (*Ivashkiv, 2018*). Murine neonatal CD4$^+$ T cells have been

178 documented to be poised to rapid Th2 cytokine secretion due to the pre-existing

179 hypomethylation at a key Th2 cytokine regulatory region (*Rose et al., 2007*). In other

180 reports, Th2 polarization in adult SP thymocytes and RTEs is attributable to active

181 recruitment of DNA methyltransferases and increased H3K4 methylation at *Il4* locus

182 (*Berkley et al., 2013; Makar et al., 2003*). In addition, DNA methylation and histone

183 modification are found to be important for the control of *Ifng* and *Tbx21* expression

184 *(Fang et al., 2022; Friedman et al., 2023; Jones & Chen, 2006)*. Therefore, we next

185 evaluated the contribution of epigenetic modifications to the enhanced IFN-γ

186 production in the absence of IL-27p28. Firstly, we performed bisulfite genomic

187 sequencing on DNA of CD4 SP thymocytes from *Cd11c-cre p28^f/f* mice and WT

188 littermates. We focused on the nine CpG sites (Figure 3A) most proximal to the *Ifng*

189 transcription start site, among which the -53 CpG has previously been suggested to play

190 a key role in IFN-γ repression in Th2 effector cells (*Jones & Chen, 2006*). DNA

191 methylation was significantly reduced at three CpG sites (-53, -34 and +16 site) of the

192 *Ifng* locus in CD4SP thymocytes from*Cd11c-cre p28^f/f* mice (Figure 3B). On the

193 contrary, the five CpG sites in the *Il4* promoter region (Figure 3C, D), whose

194 demethylation is associated with high level IL-4 production (*Lee et al., 2002*), showed

195 no difference between cKO and WT CD4SP thymocytes.

196 As much as histone modifications are concerned, histone H3 lysine 27

trimethylation (H3K27me3) and H3 lysine 9 trimethylation (H3K9me3) are typically

associated with repressed *Ifng* and *Tbx21* expression in Th cells, whereas histone H3

lysine 4 trimethylation (H3K4me3) enhances gene expression (*Fang et al., 2022;*

Friedman et al., 2023). To evaluate the histone trimethylation modifications at *Ifng,*

Tbx21, Il4 and *Gata3* loci in CD4SP thymocytes from WT and *Cd11c-p28^{f/f}* mice,

chromatin immunoprecipitation was performed using antibodies against H3K4me3,

H3K9me3 and H3K27me3. H3K4me3 was found to be accumulated at *Ifng* and *Tbx21*

loci in cKO CD4SP thymocytes, while its level in *Gata3* and *Il4* loci was comparable

to that of wild type cells (Figure 3E). On the other hand, neither the occupancy of

H3K9me3 and H3K27me3 at *Ifng, Tbx21, Il4* and *Gata3* loci was altered (Figure 3F,

G), nor were the total protein levels of H3K4me3, H3K9me3 H3K27me3 or the mRNA

level of enzymes catalyzing the formation of these modifications (Figure 3-figure

supplement 1A-C). These results suggest a link between up-regulated *Ifng* and *Tbx21*

expression and altered epigenetic modifications in the absence of IL-27p28.

4.   Coordinated up-regulation of STAT1-activated genes in CD4SP thymocytes

from *Cd11c-p28^{f/f}* mice

    To better understand the molecular mechanisms underlying the altered

functionality, the transcriptional profile of CD4SP thymocytes from *Cd11c-p28^{f/f}* mice

was explored by RNA-Seq and compared with that of the WT counterparts. The two

sets of differentially expressed genes (DEGs) were subjected to analyses for gene

ontology (GO) biological process terms and Kyoto Encyclopedia of Genes and

Genomes (KEGG) pathways. The up-regulated DEGs showed significant enrichment

in genes related to host defense to viral infection, including interferon response and Jak-

STAT signaling (Figure 4A). No meaningful hit, on the other hand, was revealed for the

down-regulated gene set (data not shown).

Given that STAT1 is a key mediator downstream of IL-27 signaling (*Philips et al.,*

2022), we were particularly interested in the altered expression of STAT1-regulated

genes in the absence of IL-27p28. A previous study by Wan et al. (*Wan et al., 2015*) has

compared the transcriptomes of wild-type and STAT1-deficient CD4$^+$ T cells stimulated

by IL-21. Using their data (GSE63204), we compiled a list of genes potentially under

regulation of STAT1. The DEGs were then examined for overlaps with this gene set.

Surprisingly, over 40% of up-regulated DEGs (55 out of 137) fell into the category of

STAT1-activated genes, whereas only 2 of them belonged to STAT1-suppressed genes

(Figure 4B). To validate the RNA-Seq data, quantitative PCR was carried out for a

group of overlapping genes, including *Gm12250, Oasl2, Oas3, Parp14, Ifit3, Usp18,*

Igtp, Irf1, Ifi44, Oas2, Rsad2, Il12r. As shown in Figure 4C, all these genes showed

differential expression between WT and cKO CD4SP thymocytes. To further illustrate

modest but coordinate changes in the expression of STAT1-regulated genes, gene set

enrichment analysis (GSEA) was performed without imposing an arbitrary fold-change

cutoff. Indeed, significant enrichment of STAT1-activated genes was detected in the

transcriptome of CD4SP thymocytes from the knockout mice (NES=-2.27, NOM p-

val=0, Figure 4D). Therefore, IL-27p28 deficiency resulted in an enhanced STAT1

activity.

To reveal functionally important connections of the DEGs, protein-protein

interaction network was generated using Network Analyst (*Xia et al., 2015*). After

removal of isolated and loosely connected nodes, the rest of the proteins encoded by

the DEGs were imported into Network Analyst for network construction. As shown in

Figure 4E, STAT1 was centrally positioned in the network with the highest degree of

interactions. Taken together, these results indicate that IL-27p28 deficiency is strongly

associated with an altered transcriptional program featured by enhanced expression of

STAT1-activated genes.

5.   Constitutive activation of STAT1 in CD4SP thymocytes in the absence of IL-

27p28

The distinct transcriptional signature prompted us to examine the activation status

of STAT1 in freshly isolated CD4SP thymocytes from *Cd11c-p28^{f/f}* mice and the WT

littermates. Total thymocytes were stained for CD4 and CD8, followed by intracellular

staining with antibodies against phosphorylated STATs. Significant levels of STAT1

Y701 phosphorylation were observed in cKO but not in WT cells. On the other hand,

both cKO and WT cells were stained negative for phosphorylated STAT3 and STAT4

(Figure 5A). Phosphorylated and total STAT proteins were also analyzed by Western

blotting in purified CD4SP thymocytes, RTEs and naive CD4$^+$ T cells. While the total

protein of STAT1 was comparable between cKO and WT cells in all populations

examined, elevated levels of phosphorylated STAT1 Y701 were constantly detected in

cKO cells. Moreover, the phosphorylation of STAT1 Y701 appeared to be

developmentally regulated, increasing progressively from CD4SP thymocytes to naive T cells (Figure 5B). Another predominant phosphorylation site in STAT1 is serine 727, which is crucial for maximal STAT1 transcription activity (*Barnholt et al., 2009*). Basal levels of S727 phosphorylation were also observed in CD4SP thymocytes, but with no difference in WT and cKO cells (data not shown). As much as STAT3 and STAT4 were concerned, neither total nor phosphorylated STAT3 and STAT4 were found to be altered with IL-27p28 ablation (Figure 5B), indicating that the impact is specific for STAT1.

STAT1 plays an important role in the development of Th1 response by regulating *Tbx21* and *Ifng* expression (*Fang et al., 2022*). To explore the correlation of the increased STAT1 activation with the enhanced T-bet expression and IFN-γ production, we assessed STAT1 binding on the *Tbx21* and *Ifng* loci by ChIP assay. Possibly due to the relatively low abundance of activated STAT1, poor consistency was observed from one experiment to another with freshly isolated CD4SP thymocytes. Therefore, we chose to treat the cells with anti-CD3 and anti-CD28 for 3 days prior to the assay. Under such conditions, STAT1 was found to be markedly accumulated in the promoter and regulatory regions of both *Tbx21* and *Ifng* loci of cKO mice (Figure 5C). Notably, there were strong positive correlations between the levels of STAT1 binding and H3K4me3 (Pearson's correlation coefficient for *Ifng* locus r = 0.917, p = 0.010 and *Tbx21* locus r = 0.991, p <0.001, Figure 5D), indicating the implication of STAT1 in the epigenetic modifications of the loci. Together, these data suggest that IL-27p28-medaited signal normally imposes an inhibitory effect on STAT1 activation, which in turn helps the establishment of functional bias against the Th1 lineage.

6. Augmented autoimmune phenotype of *Aire*-deficient mice in the absence of IL-27p28

The physiological significance remains elusive of the bias against the Th1 effector functions for CD4SP thymocytes and RTEs. One hypothesis proposes that it favors the development of peripheral tolerance by avoiding a more inflammatory and potential detrimental response to self-antigens (*Cunningham et al., 2018; Fink & Hendricks, 2011; Hendricks & Fink, 2011*). The disruption of such a bias in *Cd11c-p28^{f/f}* mice provided a good model to test this hypothesis. Flow cytometric analysis of the peripheral T cell compartment demonstrated a significant increase of $CD44^{hi}CD62L^-$ activated T cells with a concomitant decrease of $CD44^{lo}CD62L^+$ naive T cells in the cKO mice compared to the WT littermates (Figure 6A). Nevertheless, the cKO mice displayed no obvious signs of autoimmunity up to the age of 24-30 weeks when assessed by auto-antibodies against double-strand DNA (Figure 6B) and tissue pathology (Figure 6C). Occasionally, small lymphoid foci were observed in the lung of cKO mice. But the $CD4^+$ T cells recovered from the lung tissues displayed a similar cytokine secretion profile to that of WT mice except for the intrinsic augmentation of IFN-γ production (Figure 6D). Therefore, IL-27p28 deficiency alone is insufficient to drive the development of autoimmunity.

We next sought to determine whether IL-27p28 deficiency would affect the autoimmune phonotype in the *Aire^{-/-}* mouse, which are predisposed to developing autoimmunity due to a defect in clonal deletion of autoreactive T cells (*van Laar et al.,*

2022; J. X. Wang et al., 2021). To this end, *Cd11c-p28$^{f/f}$* mice were crossed with *Aire$^{-/-}$*

mice to generate double knockout mice. Histological examination revealed increased

lymphocyte infiltration and more severe structural distortion in the lung and stomach

in double knockout mice in comparison to mice deficient for Aire only (Figure 6C). In

addition, the double knockout mice exhibited an elevated level of autoantibodies to

double-strand DNA, although the difference did not reach a statistical significance

(*p*=0.068) possibly due to the limited sample size (Figure 6B). When CD4$^+$ T cells

recovered from the lung tissues were analyzed for IFN-γ, IL-4 and IL-17 production,

the double knockout mice were virtually indistinguishable from the *Aire$^{-/-}$* mice, except

for increased IFN-γ production (Figure 6D).

Immunological anergy constitutes an important mechanism for peripheral

tolerance. Two studies have identified a subset of naturally occurring

Foxp3$^-$CD44^{hi}CD73^{hi}FR4^{hi} polyclonal CD4$^+$ T cells in healthy hosts. They are enriched

for self-antigen–specific T cell antigen receptors and represent functionally anergic

cells (*ElTanbouly & Noelle, 2021; Kalekar et al., 2016*). We wondered whether the Th1

bias induced by IL-27p28 deficiency would affect the generation of the anergic

population. As shown in Figure 6E, the CD4$^+$Foxp3$^-$CD44^{hi}CD73^{hi}FR4^{hi} anergic cells

were significantly reduced in the double KO mice, which was accompanied by an

increase in the CD4$^+$Foxp3$^-$CD44^{hi}CD73^{lo}FR4^{lo} effector/memory cells. Taken together,

these data support that the bias against the Th1 lineage in newly generated T cells

provides a unique mechanism to render autoreactive cells exquisitely sensitive to

tolerance induction.

Discussion

Th2 polarization in adult SP thymocytes and RTEs is believed to be maintained through epigenetic modification at the *Il4* locus. However, the regulation of the opposing cytokine IFN-γ in this context remains unclear. We reported here that thymic DC-derived IL-27p28 played a key role in the establishment of the repressive status of the *Ifng* locus in newly generated CD4$^+$ T cells. As such, the loss of IL-27p28 in DCs endowed these cells with increased capacity of INF-γ production upon TCR engagement. Transcriptome profiling demonstrated coordinated up-regulation of STAT1-activated genes in the absence of IL-27p28. Indeed, enhanced phosphorylation of STAT1 was observed in CD4SP thymocytes from *Cd11c-p28*$^{f/f}$ mice, which was accompanied by the accumulation of STAT1 at the promoter and enhancer regions of *Ifng* and *Tbx21*. Epigenetic analyses indicated reduced DNA methylation at the *Ifng* locus and increased trimethylation of H3K4 at both *Ifng* and *Tbx21* loci in the absence of IL-27p28. Moreover, the H3K4me3 modification was shown to be strongly correlated with STAT1 binding to these loci. These data support a hypothesis that the exposure of developing thymocytes to IL-27p28 induces repressive epigenetic changes at the *Ifng* and *Tbx21* loci, possibly through antagonizing STAT1 activation, ultimately leading to an attenuated IFN-γ response of newly generated CD4SP thymocytes.

The present study revealed a significantly elevated level of phosphorylated STAT1 and up-regulation of a variety of STAT1-regulated genes in CD4SP thymocytes from *Cd11c-p28*$^{f/f}$ mice, indicating that IL-27p28 plays an important role in the fine tuning of STAT1 signaling in the developing thymocytes. As IL-27 is known to activate

multiple STAT proteins, including STAT1, STAT3, STAT4, and STAT5 following

engagement of the gp130/IL-27Rα receptor (*Philips et al., 2022*), the hyperactivation

of STAT1 in the absence of IL-27p28 is somehow unexpected. We examined the

potential impact of IL-27 deficiency on the expression of suppressors of cytokine

signaling 3 (SOCS3), which has been shown to have a regulatory role in IL-27 signaling

(*Owaki et al., 2006*). Comparable levels of SOCS3 were detected in CD4SP thymocytes

and naive CD4$^+$ T cells from *Cd11c-p28$^{f/f}$* mice and their littermates (Figure 5-figure

supplement 1). We also explored the possibility that the constitutive activation of

STAT1 might be the secondary effect of enhanced IFN-γ production. Addition of IFN-

γ antibodies to the culture of CD4SP thymocytes and naive CD4$^+$ T cells, however,

showed no significant effect on phosphorylated STAT1 levels in knockout cells (Figure

5-figure supplement 2A-B). Thus, STAT1 hyperaction is unlikely due to loss of SOCS3

expression or positive feedback of IFN-γ signaling. Of note, it has been previously

demonstrated that IL-27p28 on its own can act as a natural antagonist of gp130-

mediated signaling triggered by IL-6, IL-11 and IL-27 (*Stumhofer et al., 2010*).

Consistent with this finding, Chong et al. reported that IL-27p28 was able to inhibit IL-

27-induced Th1 differentiation by reducing the phosphorylation of STAT1 and STAT3

(*Chong et al., 2014*). Therefore, we speculate that IL-27p28 in the thymus may

primarily functions to suppress STAT1 activation induced by other cytokines.

A recent study demonstrated that both recombinant murine and human IL-27

induced the expression of antiviral proteins *Oas1*, *Oas2*, *Oas3*, *Oasl1*, and *Oasl2* in

epidermal keratinocytes. IL-27 signaling leads to OAS2 expression in a manner

dependent on IL27Rα and STAT1, but independent of STAT2. Apart from their antiviral

activity, OAS proteins are also involved in cell growth, differentiation, and apoptosis

(*Castelli et al., 1997; Huang et al., 2022; Salzberg et al., 1997*). The genomic mutation

and methylation of most OAS genes have been shown to alter their expression levels,

which is associated with the levels of infiltrated $CD4^+$ T cells and $CD8^+$ T cells in the

tumor microenvironment (*Gao et al., 2022*). Secreted OAS2 has also been observed to

inhibit CD3 zeta chain expression in T cells (*Dar et al., 2016*). In our current study, we

observed an increase in the transcription of *Oasl2, Oas2*, and *Oas3* in CD4SP

thymocytes from *Cd11c-p28^{f/f}* mice, indicating that IL-27p28 signaling also involved

in the fine-tuning of these OAS protein expressions. Whether these molecules play a

role in thymocyte development remains to be clarified.

The biased functionality of newly generated $CD4^+$ T cells has been proposed to be

important for the induction of peripheral tolerance to self-antigens (*Cunningham et al.,*

2018; Fink & Hendricks, 2011; Hendricks & Fink, 2011). Despite the much enhanced

capacity of IFN-γ production by $CD4^+$ T cells and a significant increase of

$CD44^{hi}CD62L^-$ activated T cells, *Cd11c-p28^{f/f}* mice displayed no apparent signs of

autoimmunity under steady-state conditions. This is consistent with the result of our

previous study in which negative selection was found to proceed normally in *Cd11c-*

p28^{f/f} mice using the RIP-mOVA and OT-II models (*Tang et al., 2016*). We further

interrogated the impact of the disrupted functional bias in *Aire^{-/-}* mice. Aire induces

ectopic expression of self-antigens in medullary thymic epithelial cells to mediate

negative selection of autoreactive thymocytes. Its mutation causes a rare life-

threatening autoimmune disease affecting multiple organs (*van Laar et al., 2022; J. X.*

Wang et al., 2021). In comparison to *Aire*[-/-] mice, *Cd11c-p28*[f/f] *Aire*[-/-] mice showed

elevated levels of double-strand DNA autoantibodies, increased lymphocytes

infiltration, and reduced anergic cells. Therefore, although disruption of the functional

bias in newly generated T cells is not sufficient to drive autoimmunity, it accelerates

disease development in hosts poised for autoimmunity.

In further support of a direct link between IL-27 and Aire, a recent study revealed

that IL-27 production was inhibited in Aire-overexpressing murine dendritic cells (*Zou*

et al., 2021). The two subunits IL-27p28 and EBI3 of human IL-27 are usually

coordinately expressed in antigen-presenting cells, while mouse IL-27p28 can be

secreted independently to inhibit the activities of IL-27 signaling (*Pflanz et al., 2002;*

Stumhofer et al., 2010). The distinct production profile of IL-27p28 between humans

and mice and our finding of the critical role of IL-27p28 in the maintenance of

functional bias in newly generated CD4[+] T cells may explain the much weaker

autoimmune phenotype in *Aire*[-/-] mice than *AIRE*-mutated human subjects. In addition,

recent studies have reported elevation of serum IL-27p28 (also known as IL-30) in

patients with prostate cancer (*Sorrentino et al., 2019*), psoriasis (*Omar et al., 2021*),

and obesity (*Q. Wang et al., 2021*). It would be interesting to explore the involvement

of altered functionality of newly generated CD4[+] T cells under these pathological

conditions.

Taken together, our data demonstrate that DC-derived IL-27p28 serves as an

endogenous inhibitor for STAT1 hyper-activation in developing thymocytes. By

regulating DNA and H3K4 methylation levels at the promoter and transcription

regulatory regions of *Ifng* and *Tbx21*, it is critically involved in the establishment of the

functional bias against IFN-γ production by newly generated CD4$^+$ T cells. Disruption

of this mechanism exacerbates autoimmune phenotypes in hosts predisposed for

autoimmunity.

Materials and methods

Mice

C57BL/6 mice were obtained from the Vital River Laboratories (Beijing, China). *Cd11c-p28*$^{f/f}$ mice and *Il27ra*$^{-/-}$ mice on C57BL/6 background were kindly provided by Dr. Zhinan Yin from Jinan University (Guangzhou, China). FVB-Tg (Rag2-EGFP) 1Mnz/J mice were purchased from Jackson Laboratory (Bar Harbor, ME) and were backcrossed for 10 generations onto the C57BL/6 background (termed as RAG2p-EGFP in this paper). RAG2p-EGFP mice were bred with *Cd11c-p28*$^{f/f}$ mice to generate *Cd11c-p28*$^{f/f}$ RAG2p-EGFP mice. *Aire*$^{-/-}$ mice were generously provided by Yangxin Fu (University of Chicago, IL) and were bred with *Cd11c-p28*$^{f/f}$ mice to generate *Aire*$^{-/-}$ *Cd11c-p28*$^{f/f}$ mice. Mice were used at 6-8 weeks of age unless stated otherwise. All the animal procedures were conformed to the Chinese Council on Animal Care Guidelines and the study was approved by the ethics committee of Peking University Health Science Center with an approval number of LA2014178.

Antibodies and reagents

PE-Cy7-conjugated anti-mouse CD4 (RM4-5), PE- and APC-conjugated anti-mouse CD8a (53–6.7), APC-conjugated anti-mouse IL-2 (JES6-5H4), PE-Cy7-conjugated anti-mouse TNF-α (MP6-XT22), PE-conjugated anti-mouse Stat1 (pY701) (4a), PerCP-Cy5.5-conjugated anti-mouse Stat3 (pY705) (4/P-STAT3), Alexa Fluor 488-conjugated anti-mouse Stat4 (pY693) (38/p-Stat4) were purchased from BD Biosciences (San Diego, CA). PE- and APC-conjugated anti-mouse CD25 (PC61.5)

and CD44 (IM7), FITC-conjugated anti-mouse FR4 (eBio12A5), PerCP-eFluor710-

conjugated anti-mouse CD73 (eBioTY/11.8), FITC-conjugated anti-mouse IL-4

(BVD6-24G2), PE-conjugated anti-mouse IL-17A (eBio17B7), APC-conjugated anti-

mouse FOXP3 (3G3), biotin-conjugated anti-mouse CD8a (53–6.7) and anti-phospho-

STAT-1(S727) were obtained from eBioscience (Waltham, MA). PE-conjugated anti-

mouse IFN-γ (XMG1.2) and PerCP-Cy5.5-conjugated anti-mouse T-bet (4B10) were

purchased from BioLegend (San Diego, CA).

Functional grade monoclonal antibodies for murine CD3 (145-2C11) and CD28

(37.51) and neutralizing anti-IL4 (11B11), neutralizing anti-IFN-γ (clone XMG1.2)

were obtained from eBioscience (Waltham, MA). Recombinant murine IL-2, IL-4, IL-

6, IL-12 and recombinant human TGF-β1 were purchased from R&D Systems

(Abingdon, UK).

Anti-phospho-STAT1(Tyr701), anti-STAT1, anti-phospho-STAT3 (Tyr705), anti-

STAT3, anti-phospho-STAT4 (Tyr693), anti-STAT4, anti-SOCS3, anti-Actin and HRP-

labeled goat-anti-rabbit or anti-mouse IgGs for Western blot assay were purchased from

Cell Signaling Technology (Danvers, MA). Phorbol 12-myristate 13-acetate (PMA) and

ionomycin were obtained from Sigma-Aldrich.

Cell sorting and CD4$^+$ T-cell differentiation *in vitro*

To enrich CD4SP thymocytes, CD8$^-$ thymocytes were obtained by negative

selection using anti-CD8 (Ly-2) MicroBeads (Miltenyi Biotec). The cells were then

stained with fluorescently labeled antibodies to CD4, CD8 and CD44. CD4$^+$SP

thymocytes with the phenotype of $GFP^+CD4^+CD8^-CD44^{lo}$ were then sorted. For the

isolation of $CD4^+$ RTEs and mature naive $CD4^+$ T cells, $GFP^+CD4^+CD8^-CD25^-NK1.1^-$

(RTEs) and $GFP^-CD4^+CD8^-CD25^-CD44^{lo}$ (naive T) cells were sorted from mesenteric

lymph nodes. All these cells were sorted to > 99% purity with a FACS Aria II (BD

Biosciences, San Diego, CA).

For *in vitro* activation, CD4SP thymocytes, RTEs and mature naive T cells were

cultured at a density of $2×10^6$/mL with plate-coated anti-CD3 (clone 2C11; 2 μg/mL)

and soluble anti-CD28 (clone 37N; 1 μg/mL) in RPMl 1640 medium supplemented

with 10% fetal bovine serum (Biochrom Ag, Berlin), penicillin, streptomycin and 50

μM 2-mercaptoethanol. Conditions for $CD4^+$ T-cell differentiation were as follows: Th0

cells, IL-2 (2 ng/mL); Th1 cells, IL-12 (10 ng/mL), IL-2 (2 ng/mL), anti-IL-4 (10

μg/mL); Th2 cells, IL-4 (20 ng/mL), IL-2 (2 ng/mL), anti-IFN-γ (10 μg/mL); Th17

cells, recombinant human TGFβ1 (10 ng/mL), IL-6 (10 ng/mL), anti-IL-4 (10 μg/mL),

anti-IFN-γ (10 μg/mL); and Treg cells, TGFβ1 (1 ng/mL), IL-2 (10 ng/mL), anti-IL-4

(10 μg/mL), anti-IFN-γ (10 μg/mL). After 3 days of differentiation, supernatants were

collected and cells were subjected to intracellular cytokine staining and flow cytometry

analyses.

Intracellular staining and flow cytometry

For detection of surface molecules, T cells were labelled with the appropriate

fluorescent mAbs on ice for 30 min. For detection of cytoplasmic molecules, T-bet, and

FOXP3, T cells were collected, stained with surface molecules, fixed, permeabilized

with the Foxp3/Transcription Factor Staining Buffer (eBioscience), and stained with

fluorochrome-conjugated mouse antibodies on ice for 30 min. For intracellular cytokine

staining, five hours before harvest, T cells were stimulated with PMA (50 ng/mL) plus

ionomycin (1 mg/mL) in the presence of a protein transport inhibitor, monensin (2 μM)

or BFA (3 μg/mL) (eBioscience). Cells were collected, washed, fixed, permeabilized

(FIX AND PERM, Invitrogen) and stained with IFN-γ, IL-4, IL-2, TNF-α, and IL-17

antibodies according to the manufacturer's instructions. For detection of

phosphorylated STATs, 1×10^7 thymocytes were fixed for 10 min at 37°C with 2%

(wt/vol) paraformaldehyde. After fixing, cells were permeabilized for 30 min on ice

with 90% (vol/vol) methanol, and stained with the appropriate antibodies. Flow

cytometry was conducted on a Galios (Beckman Coulter) and data analysis was

performed using Kaluza software.

Enzyme-Linked Immunosorbent Assay (ELISA)

Supernatants of *in vitro* cell cultures were obtained at 72 hr. IFN-γ and IL-4

production were determined using ELISA kits (BioLegend, San Diego, CA) according

to the manufacturer's instructions.

For the analysis of anti-dsDNA antibodies, serum was collected from 24-30-week-

old *Aire$^{-/-}$Cd11c-Cre p28$^{f/f}$*, *Aire$^{-/-}$*, *Cd11c-Cre p28$^{f/f}$* and WT mice. Nunc MaxiSorp

ELISA plates were precoated with dsDNA (100 mg/mL, Sigma, St. Louis, MO) in

phosphate-buffered saline (PBS) at 4°C overnight. Plates were blocked with 5% BSA

for 1 hour at 37°C, then washed and incubated with 1/100 dilutions of mouse sera for 2

hours at 37˚C. Plates were washed, and anti-dsDNA antibodies were detected with a

1/1000 dilution of alkaline phosphatase–conjugated goat anti-mouse IgG (BioLegend)

for 1 hour at 37˚C and developed with a phosphatase substrate for 30 min at 37˚C.

Quantitative PCR

RNA was purified from various T cell subsets cultured under Th0 conditions for

12 h using Trizol reagent (Invitrogen). cDNA was synthesized using reverse

transcription kit (Progema). Quantitative PCR was performed using FastStart Universal

SYBR Green Master mix (Roche, Basel, Switzerland) on an iCycler real-time PCR

system (Bio-Rad Laboratories, Hemel Hempstead, U.K.), with each sample in triplicate.

The quantification was based on delta delta CT calculations and was normalized to β-

actin as loading controls. The primers used in the study were listed in Supplementary

Table 1.

DNA methylation

Bisulfite modification of genomic DNA from FACS purified cells was performed

using the EZ DNA methylation kit (Zymo Research). For methylation analysis on *Ifng*

gene promoter CpG sites, bisulfite-treated DNA was amplified in semi-nested PCR

using primers: 5'-GGTGTGAAGTAAAAGTGTTTTTAGAGAATTTTAT-3' and 5'-

CAATAACAACCAAAAACAACCATAAAAAAAAACT-3', then 5'-

GGTGTGAAGTAAAAGTGTTTTTAGAGAATTTTAT-3' and 5'-

CCATAAAAAAAAACTACAAAACCAAAATACAATA-3'. For methylation analysis

on *Il4* gene promoter CpG sites, bisulfite-treated DNA was amplified in PCR using

primers: 5-GGATCCACACGGTGCAAAGAGAGACCC-3'   and   5′-

TCGGCCTTTCAGACTAATCTTATCAGC-3' The PCR products were gel purified

and cloned into the pGEM-T vector (Promega; Madison, WI, USA). The inserted PCR

fragments of individual clones were sequenced by Tsing KE Biological Technology,

Beijing, China. For all samples, 10 reads per CpG site were used to determine the

average percentage of methylated CpG.

Chromatin immunoprecipitation (ChIP)

ChIP assays were carried out by using Pierce Agarose Chip Kit (Pierce

Biotechnology). Briefly, cells were cross-linked for 10 min with 1% formaldehyde and

lysed with Nuclease. 10% of the digested chromatin was preserved as input control and

the rest of the digested chromatin were incubated with specific antibody or normal IgG

(as a control). The purified DNA from immunoprecipitation and the input samples were

analyzed by quantitative PCR using primers specific listed in Supplementary Table 1.

Data is presented as a percent input of each IP sample relative to input chromatin. The

following antibodies were used in ChIP analysis: H3K4me3 (Millipore), H3K27me3

(Millipore), H3K9me3 (Abcam), and STAT1 (Cell Signaling Technology).

Western blotting

For short term TCR stimulation, CD4SP thymocytes and mature naive T cells were

incubated with 2 μg/mL anti-mouse CD3 and 1 μg/mL anti-mouse CD28 on ice for 20

min, followed by cross-linking for 10 min at 37°C with 5 μg/mL goat-anti-hamster IgG.

Cells were then washed with PBS. Freshly isolated T cells or stimulated T cells were

lysed in RIPA buffer for 30 minutes. Whole-cell lysates (5–10 μg per sample) were

separated by SDS/PAGE and analyzed by immunoblotting with antibodies to phospho-

STAT-1 (Tyr701), phospho-STAT-1 (Ser727), STAT-1, phospho-STAT-3 (Tyr705),

STAT-3, phospho-STAT-4 (Tyr693), STAT-4, SOCS3 and Actin. HRP-conjugated anti-

Rabbit IgG was used as the detection antibody. The bands were quantified using ImageJ

software.

RNA-seq

Total RNA was extracted from approximately 2×10^6 CD4SP thymocytes from WT

and *Cd11c-p28*$^{f/f}$ mice using Trizol reagent (Invitrogen). Each sample was a mixture of

equal amounts of CD4SP thymocytes from three mice. RNA underwent quality control

testing using a bionalayser followed by cDNA library preparation. Library construction

and sequencing on Illumina HiSEq 2000 were performed by BIOPIC, Peking

University.

H&E

Organs from 24-30-week-old *Aire*$^{-/-}$*Cd11c-p28*$^{f/f}$ mice, *Aire*$^{-/-}$ mice and WT mice

were collected and fixed overnight in 10% formalin, embedded in paraffin, sectioned

and stained with hematoxylin and eosin (H&E). The degree of lymphocytic infiltrates

was analyzed in a blinded fashion. In general, 0, 1, 2 and 3 indicate no, mild, moderate,

or severe lymphocytic infiltration, respectively.

Statistical analysis

Data are reported as the mean ± Standard deviation (SD). Differences between

groups were analyzed by Student's t test or two-way Anova (for multiple variant

comparisons) using GraphPad Prism software (GraphPad). Throughout the text, figures,

and figure legends, the following terminology is used to denote statistical significance:

* $P<0.05$, ** $P<0.01$. *** $P<0.001$. NS, no significance.

Ashby, K. M., & Hogquist, K. A. (2023). A guide to thymic selection of T cells. *Nat Rev Immunol*.
doi:10.1038/s41577-023-00911-8

Barnholt, K. E., Kota, R. S., Aung, H. H., & Rutledge, J. C. (2009). Adenosine blocks IFN-gamma-
induced phosphorylation of STAT1 on serine 727 to reduce macrophage activation.
Journal of Immunology, 183(10), 6767-6777. doi:10.4049/jimmunol.0900331

Berkley, A. M., Hendricks, D. W., Simmons, K. B., & Fink, P. J. (2013). Recent Thymic Emigrants and
Mature Naive T Cells Exhibit Differential DNA Methylation at Key Cytokine Loci. *Journal of*
Immunology, 190(12), 6180-6186. doi:10.4049/jimmunol.1300181

Bhaumik, S., Giffon, T., Bolinger, D., Kirkman, R., Lewis, D. B., Weaver, C. T., & Randolph, D. A. (2013).
Retinoic Acid Hypersensitivity Promotes Peripheral Tolerance in Recent Thymic Emigrants.
Journal of Immunology, 190(6), 2603-2613. doi:10.4049/jimmunol.1200852

Cao, Y. X., Doodes, P. D., Glant, T. T., & Finnegan, A. (2008). IL-27 induces a Th1 immune response
and susceptibility to experimental arthritis. *Journal of Immunology, 180*(2), 922-930.
doi:DOI 10.4049/jimmunol.180.2.922

Castelli, J. C., Hassel, B. A., Wood, K. A., Li, X. L., Amemiya, K., Dalakas, M. C., Torrence, P. F., &
Youle, R. J. (1997). A study of the interferon antiviral mechanism: apoptosis activation by
the 2-5A system. *J Exp Med, 186*(6), 967-972. doi:10.1084/jem.186.6.967

Caveney, N. A., Glassman, C. R., Jude, K. M., Tsutsumi, N., & Garcia, K. C. (2022). Structure of the
IL-27 quaternary receptor signaling complex. *Elife, 11*. doi:10.7554/eLife.78463

Chong, W. P., Horai, R., Mattapallil, M. J., Silver, P. B., Chen, J., Zhou, R., Sergeev, Y., Villasmil, R.,
Chan, C. C., & Caspi, R. R. (2014). IL-27p28 inhibits central nervous system autoimmunity
by concurrently antagonizing Th1 and Th17 responses. *J Autoimmun, 50*, 12-22.
doi:10.1016/j.jaut.2013.08.003

Cunningham, C. A., Helm, E. Y., & Fink, P. J. (2018). Reinterpreting recent thymic emigrant function:
defective or adaptive? *Curr Opin Immunol, 51*, 1-6. doi:10.1016/j.coi.2017.12.006

Dar, A. A., Pradhan, T. N., Kulkarni, D. P., Shah, S. U., Rao, K. V., Chaukar, D. A., D'Cruz, A. K., &
Chiplunkar, S. V. (2016). Extracellular 2'5'-oligoadenylate synthetase 2 mediates T-cell
receptor CD3-zeta chain down-regulation via caspase-3 activation in oral cancer.
Immunology, 147(2), 251-264. doi:10.1111/imm.12560

Diveu, C., McGeachy, M. J., Boniface, K., Stumhofer, J. S., Sathe, M., Joyce-Shaikh, B., Chen, Y., Tato,
C. M., McClanahan, T. K., Malefyt, R. D., Hunter, C. A., Cua, D. J., & Kastelein, R. A. (2009).
IL-27 Blocks RORc Expression to Inhibit Lineage Commitment of Th17 Cells. *Journal of*
Immunology, 182(9), 5748-5756. doi:10.4049/jimmunol.0801162

ElTanbouly, M. A., & Noelle, R. J. (2021). Rethinking peripheral T cell tolerance: checkpoints across
a T cell's journey. *Nature Reviews Immunology, 21*(4), 257-267. doi:10.1038/s41577-020-
00454-2

Fang, D., Healy, A., & Zhu, J. (2022). Differential regulation of lineage-determining transcription
factor expression in innate lymphoid cell and adaptive T helper cell subsets. *Front*
Immunol, 13, 1081153. doi:10.3389/fimmu.2022.1081153

Fink, P. J., & Hendricks, D. W. (2011). Post-thymic maturation: young T cells assert their individuality.
Nature Reviews Immunology, 11(8), 544-549. doi:10.1038/nri3028

Flores, R. R., Kim, E., Zhou, L., Yang, C., Zhao, J., Gambotto, A., & Robbins, P. D. (2015). IL-Y, a
synthetic member of the IL-12 cytokine family, suppresses the development of type 1
diabetes in NOD mice. *European Journal of Immunology, 45*(11), 3114-3125.

doi:10.1002/eji.201445403

Friedman, M. J., Lee, H., Lee, J. Y., & Oh, S. (2023). Transcriptional and Epigenetic Regulation of
Context-Dependent Plasticity in T-Helper Lineages. *Immune Netw, 23*(1), e5.
doi:10.4110/in.2023.23.e5

Gao, L., Ren, R., Shen, J., Hou, J., Ning, J., Feng, Y., Wang, M., Wu, L., Sun, Y., Wang, H., Wang, D.,
& Cao, J. (2022). Values of OAS gene family in the expression signature, immune cell
infiltration and prognosis of human bladder cancer. *BMC Cancer, 22*(1), 1016.
doi:10.1186/s12885-022-10102-8

Hamano, S., Himeno, K., Miyazaki, Y., Ishii, K., Yamanaka, A., Takeda, A., Zhang, M., Hisaeda, H.,
Mak, T. W., Yoshimura, A., & Yoshida, H. (2003). WSX-1 is required for resistance to
infection by regulation of proinflammatory cytokine production. *Immunity, 19*(5), 657-667.
doi:Doi 10.1016/S1074-7613(03)00298-X

Harker, J. A., Wong, K. A., Dolgoter, A., & Zuniga, E. I. (2015). Cell-Intrinsic gp130 Signaling on CD4
T Cells Shapes Long-Lasting Antiviral Immunity. *Journal of Immunology, 195*(3), 1071-
1081. doi:10.4049/jimmunol.1402402

Hendricks, D. W., & Fink, P. J. (2011). Recent thymic emigrants are biased against the T-helper type
1 and toward the T-helper type 2 effector lineage. *Blood, 117*(4), 1239-1249.
doi:10.1182/blood-2010-07-299263

Huang, Y. Z., Zheng, Y. X., Zhou, Y., Xu, F., Cui, Y. Z., Chen, X. Y., Wang, Z. Y., Yan, B. X., Zheng, M.,
& Man, X. Y. (2022). OAS1, OAS2, and OAS3 Contribute to Epidermal Keratinocyte
Proliferation by Regulating Cell Cycle and Augmenting IFN-1−Induced Jak1−Signal
Transducer and Activator of Transcription 1 Phosphorylation in Psoriasis. *J Invest Dermatol,*
142(10), 2635-2645 e2639. doi:10.1016/j.jid.2022.02.018

Huber, M., Steinwald, V., Guralnik, A., Brüstle, A., Kleemann, P., Rosenplänter, C., Decker, T., &
Lohoff, M. (2008). IL-27 inhibits the development of regulatory T cells via STAT3. *Wiener*
Klinische Wochenschrift, 120, 45-45.

Ivashkiv, L. B. (2018). IFNgamma: signalling, epigenetics and roles in immunity, metabolism, disease
and cancer immunotherapy. *Nat Rev Immunol, 18*(9), 545-558. doi:10.1038/s41577-018-
0029-z

Jones, B., & Chen, J. (2006). Inhibition of IFN-gamma transcription by site-specific methylation
during T helper cell development. *EMBO J, 25*(11), 2443-2452.
doi:10.1038/sj.emboj.7601148

Kalekar, L. A., Schmiel, S. E., Nandiwada, S. L., Lam, W. Y., Barsness, L. O., Zhang, N., Stritesky, G. L.,
Malhotra, D., Pauken, K. E., Linehan, J. L., O'Sullivan, M. G., Fife, B. T., Hogquist, K. A.,
Jenkins, M. K., & Mueller, D. L. (2016). CD4(+) T cell anergy prevents autoimmunity and
generates regulatory T cell precursors. *Nat Immunol, 17*(3), 304-314. doi:10.1038/ni.3331

Kim, D., Le, H. T., Nguyen, Q. T., Kim, S., Lee, J., & Min, B. (2019). Cutting Edge: IL-27 Attenuates
Autoimmune Neuroinflammation via Regulatory T Cell/Lag3-Dependent but IL-10-
Independent Mechanisms In Vivo. *Journal of Immunology, 202*(6), 1680-1685.
doi:10.4049/jimmunol.1800898

Lee, D. U., Agarwal, S., & Rao, A. (2002). Th2 lineage commitment and efficient IL-4 production
involves extended demethylation of the IL-4 gene. *Immunity, 16*(5), 649-660.
doi:10.1016/s1074-7613(02)00314-x

Lin, T. T., Lu, J., Qi, C. Y., Yuan, L., Li, X. L., Xia, L. P., & Shen, H. (2015). Elevated serum level of IL-

27 and VEGF in patients with ankylosing spondylitis and associate with disease activity.
Clinical and Experimental Medicine, 15(2), 227-231. doi:10.1007/s10238-014-0281-x

Liu, J. Q., Liu, Z. Z., Zhang, X. J., Shi, Y., Talebian, F., Carl, J. W., Yu, C., Shi, F. D., Whitacre, C. C.,
Trgovcich, J., & Bai, X. F. (2012). Increased Th17 and Regulatory T Cell Responses in EBV-
Induced Gene 3-Deficient Mice Lead to Marginally Enhanced Development of
Autoimmune Encephalomyelitis. *Journal of Immunology, 188*(7), 3099-3106.
doi:10.4049/jimmunol.1100106

Makar, K. W., Perez-Melgosa, M., Shnyreva, M., Weaver, W. M., Fitzpatrick, D. R., & Wilson, C. B.
(2003). Active recruitment of DNA methyltransferases regulates interleukin 4 in
thymocytes and T cells. *Nat Immunol, 4*(12), 1183-1190. doi:10.1038/ni1004

Mei, Y. W., Lv, Z., Xiong, L. L., Zhang, H. W., Yin, N. L., & Qi, H. B. (2021). The dual role of IL-27 in
CD4+T cells. *Molecular Immunology, 138*, 172-180. doi:10.1016/j.molimm.2021.08.001

Neufert, C., Becker, C., Wirtz, S., Fantini, M. C., Weigmann, B., Galle, P. R., & Neurath, M. F. (2007).
IL-27 controls the development of inducible regulatory T cells and Th17 cells via
differential effects on STAT1. *European Journal of Immunology, 37*(7), 1809-1816.
doi:10.1002/eji.200636896

Nguyen, Q. T., Jang, E., Le, H. T., Kim, S., Kim, D., Dvorina, N., Aronica, M. A., Baldwin, W. M.,
Asosingh, K., Comhair, S., & Min, B. (2019). IL-27 targets Foxp3 Tregs to mediate
antiinflammatory functions during experimental allergic airway inflammation. *Jci Insight,*
4(2). doi:10.1172/jci.insight.123216

Omar, N. S., Xinxin, L., Jiayi, X., Hussain, S. G., & Biao, P. X. (2021). Serum level of cytokine IL-30 in
psoriatic patients and correlation with disease severity. *J Popul Ther Clin Pharmacol, 28*(1),
24-33. doi:10.15586/jptcp.2021.v28i1.823

Owaki, T., Asakawa, M., Fukai, F., Mizuguchi, J., & Yoshimoto, T. (2006). IL-27 induces Th1
differentiation via p38 MAPK/T-bet- and intercellular adhesion molecule-1/LFA-
1/ERK1/2-dependent pathways. *Journal of Immunology, 177*(11), 7579-7587.
doi:10.4049/jimmunol.177.11.7579

Park, J., DeLong, J. H., Knox, J. J., Konradt, C., Wojno, E. D. T., & Hunter, C. A. (2019). Impact of
Interleukin-27p28 on T and B Cell Responses during Toxoplasmosis. *Infection and*
Immunity, 87(12). doi:10.1128/IAI.00455-19

Park, Y. J., Ryu, H., Choi, G., Kim, B. S., Hwang, E. S., Kim, H. S., & Chung, Y. (2019). IL-27 confers a
protumorigenic activity of regulatory T cells via CD39. *Proceedings of the National*
Academy of Sciences of the United States of America, 116(8), 3106-3111.
doi:10.1073/pnas.1810254116

Pflanz, S., Timans, J. C., Cheung, J., Rosales, R., Kanzler, H., Gilbert, J., Hibbert, L., Churakova, T.,
Travis, M., Vaisberg, E., Blumenschein, W. M., Mattson, J. D., Wagner, J. L., To, W., Zurawski,
S., McClanahan, T. K., Gorman, D. M., Bazan, J. F., de Waal Malefyt, R., Rennick, D., &
Kastelein, R. A. (2002). IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein,
induces proliferation of naive CD4+ T cells. *Immunity, 16*(6), 779-790. doi:10.1016/s1074-
7613(02)00324-2

Philips, R. L., Wang, Y., Cheon, H., Kanno, Y., Gadina, M., Sartorelli, V., Horvath, C. M., Darnell, J. E.,
Stark, G. R., & O'Shea, J. J. (2022). The JAK-STAT pathway at 30: Much learned, much more
to do. *Cell, 185*(21), 3857-3876. doi:10.1016/j.cell.2022.09.023

Rose, S., Lichtenheld, M., Foote, M. R., & Adkins, B. (2007). Murine neonatal CD4+ cells are poised

for rapid Th2 effector-like function. *Journal of Immunology, 178*(5), 2667-2678. doi:10.4049/jimmunol.178.5.2667

Salzberg, S., Hyman, T., Turm, H., Kinar, Y., Schwartz, Y., Nir, U., Lejbkowicz, F., & Huberman, E. (1997). Ectopic expression of 2-5A synthetase in myeloid cells induces growth arrest and facilitates the appearance of a myeloid differentiation marker. *Cancer Res, 57*(13), 2732-2740.

Shen, H., Xia, L. P., & Lu, J. (2013). Elevated levels of interleukin-27 and effect on production of interferon-γ and interleukin-17 in patients with Behcet's disease. *Scandinavian Journal of Rheumatology, 42*(1), 48-51. doi:10.3109/03009742.2012.704391

Sorrentino, C., Yin, Z., Ciummo, S., Lanuti, P., Lu, L. F., Marchisio, M., Bellone, M., & Di Carlo, E. (2019). Targeting Interleukin(IL)-30/IL-27p28 signaling in cancer stem-like cells and host environment synergistically inhibits prostate cancer growth and improves survival. *J Immunother Cancer, 7*(1), 201. doi:10.1186/s40425-019-0668-z

Stumhofer, J. S., Tait, E. D., Quinn, W. J., Hosken, N., Spudy, B., Goenka, R., Fielding, C. A., O'Hara, A. C., Chen, Y., Jones, M. L., Saris, C. J. M., Rose-John, S., Cua, D. J., Jones, S. A., Elloso, M. M., Grötzinger, J., Cancro, M. P., Levin, S. D., & Hunter, C. A. (2010). A role for IL-27p28 as an antagonist of gp130-mediated signaling. *Nature Immunology, 11*(12), 1119-U1118. doi:10.1038/ni.1957

Tang, H., Zhang, J., Sun, X., Qian, X., Zhang, Y., & Jin, R. (2016). Thymic DCs derived IL-27 regulates the final maturation of CD4(+) SP thymocytes. *Sci Rep, 6*, 30448. doi:10.1038/srep30448

Tormo, A. J., Meliani, Y., Beaupré, L. A., Sharma, M., Fritz, J. H., Elson, G., Crabé, S., & Gauchat, J. F. (2013). The Composite Cytokine p28/Cytokine-Like Factor 1 Sustains B Cell Proliferation and Promotes Plasma Cell Differentiation. *Journal of Immunology, 191*(4), 1657-1665. doi:10.4049/jimmunol.1201595

van Laar, G. G., Hamburg, J. P. V., & Tas, S. W. (2022). Extrathymic AIRE-expressing cells: Friends or foes in autoimmunity and cancer? *Autoimmunity Reviews, 21*(9). doi:10.1016/j.autrev.2022.103141

Villarino, A., Hibbert, L., Lieberman, L., Wilson, E., Mak, T., Yoshida, H., Kastelein, R. A., Saris, C., & Hunter, C. A. (2003). The IL-27R (WSX-1) is required to suppress T cell hyperactivity during infection. *Immunity, 19*(5), 645-655. doi:Doi 10.1016/S1074-7613(03)00300-5

Wan, C. K., Andraski, A. B., Spolski, R., Li, P., Kazemian, M., Oh, J., Samsel, L., Swanson, P. A., 2nd, McGavern, D. B., Sampaio, E. P., Freeman, A. F., Milner, J. D., Holland, S. M., & Leonard, W. J. (2015). Opposing roles of STAT1 and STAT3 in IL-21 function in CD4+ T cells. *Proc Natl Acad Sci U S A, 112*(30), 9394-9399. doi:10.1073/pnas.1511711112

Wang, J. X., Lareau, C. A., Bautista, J. L., Gupta, A. R., Sandor, K., Germino, J., Yin, Y., Arvedson, M. P., Reeder, G. C., Cramer, N. T., Xie, F., Ntranos, V., Satpathy, A. T., Anderson, M. S., & Gardner, J. M. (2021). Single-cell multiomics defines tolerogenic extrathymic Aire-expressing populations with unique homology to thymic epithelium. *Science Immunology, 6*(65). doi:10.1126/sciimmunol.abl5053

Wang, Q., Li, D., Cao, G., Shi, Q., Zhu, J., Zhang, M., Cheng, H., Wen, Q., Xu, H., Zhu, L., Zhang, H., Perry, R. J., Spadaro, O., Yang, Y., He, S., Chen, Y., Wang, B., Li, G., Liu, Z., Yang, C., Wu, X., Zhou, L., Zhou, Q., Ju, Z., Lu, H., Xin, Y., Yang, X., Wang, C., Liu, Y., Shulman, G. I., Dixit, V. D., Lu, L., Yang, H., Flavell, R. A., & Yin, Z. (2021). IL-27 signalling promotes adipocyte thermogenesis and energy expenditure. *Nature, 600*(7888), 314-318.

764    doi:10.1038/s41586-021-04127-5

765 Xia, J., Gill, E. E., & Hancock, R. E. (2015). NetworkAnalyst for statistical, visual and network-based

766    meta-analysis of gene expression data. *Nat Protoc, 10*(6), 823-844.

767    doi:10.1038/nprot.2015.052

768 Zhang, S., Liang, R. F., Luo, W., Liu, C., Wu, X. L., Gao, Y. A., Hao, J. L., Cao, G. C., Chen, X., Wei, J.,

769    Xia, S. Y., Li, Z., Wen, T., Wu, Y. Y., Zhou, X. L., Wang, P. Y., Zhao, L. Q., Wu, Z. Z., Xiong, S.

770    D., Gao, X. M., Gao, X., Chen, Y. Y., Ge, Q., Tian, Z. G., & Yin, Z. N. (2013). High susceptibility

771    to liver injury in IL-27 p28 conditional knockout mice involves intrinsic interferon-γ

772    dysregulation of CD4 T cells. *Hepatology, 57*(4), 1620-1631. doi:10.1002/hep.26166

773 Zou, X. Y., Wang, S., Zhang, Y., Wang, X. Y., Zhang, R. C., Yang, W., & Li, Y. (2021). AIRE-

774    overexpressing BMDCs suppress TFH cells through ICOSL to prevent and attenuate

775    autoimmune diabetes in NOD mice. *International Immunopharmacology, 99*.

776    doi:10.1016/j.intimp.2021.107979

Abbreviations

AS, ankylosing spondyliti; BD, behcet disease; ChIP, chromatin immunoprecipitation; cKO, conditional knockout; DC, dendritic cell; DEGs, differentially expressed genes; EAE, experimental autoimmune encephalomyelitis; EBI3, Epstein-Barr virus-induced gene 3; ELISA, Enzyme-Linked Immunosorbent Assay; GO, gene ontology; GSEA, gene set enrichment analysis; H3K4me3, histone H3 lysine 4 trimethylation; H3K27me3, histone H3 lysine 27 trimethylation; H3K9me3, histone H3 lysine 9 trimethylation; IFN-γ, interferon-γ; IL-2, interleukin 2; KEGG, Kyoto Encyclopedia of Genes and Genomes; PBS, phosphate-buffered saline; RTE, recent thymic emigrants; SOCS3, suppressors of cytokine signaling 3; SP, single-positive thymocyte; STAT, signal transducer and activator of transcription; T-bet, T-box transcription factor; Th, T helper; TNF-α, tumor necrosis factor α; WT, wild-type.

Author Contributions

JZ and RJ designed and carried out the study, collected and analyzed the data and drafted the manuscript. JZ and HT performed the phenotype analysis, epigenetic assay and established the mouse model. HMW analyzed the RNA-seq data. XWP performed the cell sorting assay. YZ conceived and designed the study, analyzed the data, and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We thank Dr. Zhinan Yin from Jinan University (Guangzhou, China) for kindly

provided *Cd11c-cre p28*^{flox/flox} mice and *Il27ra*^{-/-} mice.

Conflicts of Interest

The authors declare no conflict of interest.

Funding

This work was supported by grants from the National Natural Sciences Foundation of

China (32230037; 82394412; 32071178; 31872733), National Key Research and

Development Program of China (2023YFB3507000).

Figure 1. Elevated IFN-γ production and T-bet expression in *Cd11c-p28*$^{f/f}$ mice initiated

from CD4SP thymocytes stage.

(A) CD4SP (GFP$^+$CD4$^+$CD8$^-$CD44^{lo}) thymocytes, CD4$^+$ RTEs (GFP$^+$CD4$^+$CD8$^-$CD25$^-$

CD44^{lo}) and CD4$^+$ naive (GFP$^-$CD4$^+$CD8$^-$CD25$^-$CD44^{lo}) T cells were sorted from 6-8-

week-old *Cd11c-p28*$^{f/f}$ mice and WT littermates and stimulated with anti-CD3 (2

μg/mL, plate coated) and anti-CD28 (1 μg/mL) for 12 hours. Quantitative PCR analysis

was employed to determine the mRNA levels of cytokines, including *Ifng, Il4,* and *Il2*.

The experiments were conducted four times with duplicates for each sample, and the

data are presented as mean±SD.

(B) Sorted CD4SP thymocytes, CD4$^+$ RTEs and CD4$^+$ naive T cells from *Cd11c-*

p28^{f/f} mice and WT littermates were cultured under Th0 conditions for 3 days. The

frequency of IFN-γ-producing CD4$^+$ T cells were measured by intracellular staining.

Representative dot plots were showed on the left. The statistical data collected from

three pairs of mice were shown on the right as mean±SD.

(C) Cells sorted as described in (B) were cultured under Th0 conditions for 3 days, and

the concentrations of IFN-γ and IL-4 in the supernatants of CD4SP thymocytes, RTEs,

and naive T cells were measured by ELISA. The statistical data collected from three

pairs of mice were shown as mean±SD.

(D) Sorted cells were treated as described in (A). The mRNA levels of transcription

factors *Tbx21* and *Gata3* were determined by quantitative PCR. The experiments were

repeated 4 times with duplicates for each sample and the data are presented as mean±SD.

(E) Western blotting was performed after a 3-day culture under Th0 conditions. The

experiments were repeated three times with consistent results. The statistical data

showed the relative protein expression levels of T-bet, quantified by densitometry and

normalized to β-actin levels (mean±SD).

(F) Freshly sorted cells were dissolved in Trizol. The mRNA levels of *Ifng* and *Tbx21*

were determined by quantitative PCR. The experiments were repeated four times with

triplicates for each sample and the data are presented as mean±SEM. Statistical

differences between groups were determined by the Student's *t*-test. * $P < 0.05$, ** $P <$

0.01, *** $P < 0.001$.

Figure 2. Enhanced IFN-γ production and T-bet expression in *Il27ra^-/-^* mice initiated

from CD4SP thymocytes stage.

(A) CD4SP thymocytes and naive CD4$^+$ T cells were isolated from *Il27ra^-/-^* mice and

WT littermates and stimulated with anti-CD3 (2 μg/mL, plate coated) and anti-CD28

(1 μg/mL) for 12 hours. The mRNA levels of cytokines including *Ifng, Il4* and *Il2,* and

transcription factors *Tbx21* and *Gata3* were determined by quantitative PCR. The

experiments were conducted three times with duplicates for each sample and the data

are presented as mean±SEM.

(B) CD4SP thymocytes and CD4$^+$ naive T cells were cultured under Th0 conditions for

3 days. The frequency of IFN-γ-producing CD4$^+$ T cells were determined by

intracellular staining. Representative dot plots were showed on the left. The statistical

data collected from three pairs of mice were shown on the right as mean±SD. Statistical

differences between groups were determined by the Student's *t*-test. * $P< 0.05$, ** $P<$

858   0.01.

Figure 3. Distinct methylation patterns of DNA and H3K4 at *Ifng* and *Tbx21* promoter

regions in CD4SP thymocytes from IL27p28-deficient mice.

(A) DNA methylation analysis of nine sites in the *Ifng* promoter with sodium bisulfite-

treated genomic DNA from GFP⁺CD4⁺CD8⁻CD44^lo CD4SP thymocytes. Each row

represents a cloned and sequenced allele. Analyses included 10 clones from one of the

three independent experiments. Filled circles (●) represent methylated cytosine, and

open circles (○) the unmethylated residue.

(B) The left panel graphs show the percent of methylation at each individual site from

one representative experiment, while the right panel graphs show the average

percentage of methylation of three adjacent sites (group1: -205, -190, -170; group2: -

53, -45, -34; group3: +16, +96, +120) and all CpG sites from the three independent

experiments. The statistical data were shown as mean±SEM.

(C) DNA methylation analysis for five CpG site upstream of the transcription start site

of *Il4* with sodium bisulfite-treated genomic DNA from GFP$^+$CD4$^+$CD8$^-$CD44^{lo}

CD4SP thymocytes. Each row represents a cloned and sequenced allele. Analyses

included 10 clones from the two independent experiments. Filled circles (●) represent

methylated cytosine, and open circles (○) the unmethylated residue.

(D) Graphs show the percentage of methylation at each individual site (left panel) or all

CpG sites (right panel).

(E-G) Histone trimethylation analysis on freshly isolated CD4SP thymocytes from

IL27p28-deficient mice and littermates. ChIP-qPCR was performed using H3K4me3

(E), H3K27me3 (F) and H3K9me3 (G) specific antibodies. Quantitative PCR was

conducted using primers specific for the promoters and trans-regulatory regions of *Ifng*,

Tbx21, *Il4,* and *Gata3*. The experiments were repeated three times with duplicates for

each sample and the data are presented as mean±SEM. * $P<0.05$; ** $P<0.01$ determined

by Student's *t*-test. Abbreviation: pro., promoter.

Figure 4. Increased expression of STAT1-activated genes in CD4SP thymcoytes from *Cd11c-p28^{f/f}* mice.

RNA-seq was performed to probe the transcriptome of CD4SP thymcoytes from *Cd11c-p28^{f/f}* and WT mice.

(A) The top 10 hits from Gene Ontology (GO) biological process and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses are shown for Differentially Expressed Genes (DEGs) up-regulated in *Cd11c-p28^{f/f}* mice.

(B) Comparison of genes up- or down-regulated in the absence of IL27p28 (*Cd11c-p28^{f/f}*) and those activated or suppressed by STAT1. The numbers of overlapping genes in different categories are indicated.

(C) Validation of the expression levels of representative genes up-regulated by RNA-Seq using quantitative PCR. The experiments were repeated three times with duplicates and data are presented as fold changes in cKO versus WT mice (mean±SEM).

(D) Coordinated upregulation of STAT1-activated genes in CD4SP thymcoytes from

Cd11c-p28^{f/f} mice revealed by Gene Set Enrichment Analysis (GSEA).

(E) Interaction network of the proteins encoded by DEGs. Nodes represent individual

proteins and edges between two nodes represent interactions between proteins. Larger

nodes indicate higher degrees of interaction. Red and green nodes represent up-

regulated and down-regulated genes, respectively. Gray nodes were not in our DEG

datasets but were inserted by Network Analyst as they are connected to the network.

Figure 5. Heightened activation of STAT1 in CD4SP thymocytes from *Cd11c-p28^{f/f}*

mice.

(A) Intracellular staining was performed with freshly isolated thymocytes from *Cd11c-p28^{f/f}* mice and WT littermates using antibodies against phosphorylated STAT1 (Y701), STAT3 (Y705) and STAT4 (Y693). Representative histograms are shown for CD4SP thymocytes (*left*). The mean fluorescence intensity (MFI) from three independent experiments are presented as mean±SD (*right*).

(B) Cell lysate was prepared from purified CD4SP thymocytes, CD4$^+$ RTEs and naive

CD4$^+$ T cells from WT and *Cd11c-p28$^{f/f}$* mice. The levels of total and phosphorylated

STAT1 (Y701), STAT3 (Y705) and STAT4 (Y693) were examined using Western

blotting. The experiments were repeated three times with similar results. Representative

blots are shown on the left, and the relative expression levels of phosphorylated and

total STATs are presented as mean±SD following quantification by densitometry and

normalization to β-actin.

(C) Increased binding of STAT1 onto the promoter and regulatory regions of *Tbx21*

and *Ifng* loci.

(D) Correlation of the levels of STAT1 binding and H3K4me3 at the *Tbx21* and *Ifng*

loci. * $P<0.05$; ** $P< 0.01$.

Figure 6. Exacerbated autoimmune responses in *Aire*[-/-] mice in the absence of IL27p28.

(A) Splenocytes from 6-8 week-old WT (n=6) and *Cd11c-p28*[f/f] (n=6) mice were

analyzed for CD44 and CD62L expression after gating on CD4[+] T cells. Representative

dot plots are shown on the *left*. The percentage of CD44[lo]CD62L[+] naive and

CD44[hi]CD62L[-] activated T cells are presented as mean±SD on the *right*.

(B-D) WT, *Cd11c-p28*[f/f], *Aire*[-/-], and *Cd11c-p28*[f/f] and *Aire*[-/-] double knockout mice

were sacrificed at 24-30 weeks of age and subjected to various analyses. Each symbol

represents an individual mouse. The horizontal bars denote the average and SD.

(B) The serum levels of anti-dsDNA antibodies as measured by ELISA.

(C) Representative images of H&E staining (*left*) and histological scores (*right*) of the

lung and stomach. The arrows point to lymphocytic infiltrates. Scale bar equals 100 μm.

(D) The percentage of IFN-γ^+, IL-4$^+$, and IL-17A$^+$ cells in CD4$^+$ T cells isolated from

the lung tissue as determined by intracellular staining following stimulation with

PMA/ionomycin.

(E) Splenocytes from 12-week-old mice were stained for the expression of various

cellular markers. CD44 and Foxp3 expression was analyzed by gating on CD4$^+$ T cells,

and Foxp3$^-$CD44^{hi} population was further analyzed for CD73 and FR4 staining. The

percentage   of   anergic   (CD4$^+$Foxp3$^-$CD44^{hi}CD73^{hi}FR4^{hi}),   effector/memory

(CD4$^+$Foxp3$^-$CD44^{hi}CD73^{lo}FR4^{lo}), and regulatory (CD4$^+$Foxp3$^+$) T cells are presented

on the *right*. Each symbol represents an individual mouse. The horizontal bars denote

the average and SD. * $P<0.05$; ** $P<0.01$; ns, not significant.