Title

Immunogenicity and safety of a live-attenuated SARS-CoV-2 vaccine candidate based on multiple

attenuation mechanisms

Mie Suzuki-Okutani,[1,2,8] Shinya Okamura,[1,2,8] Tanggis,[1] Hitomi Sasaki,[1] Suni Lee,[1] Akiho Yoshida,[1,2] Simon Goto,[1]

Tatsuya Nakagawa,[3] Masahito Ikawa,[3,4,7] Wataru Kamitani,[5] Shiro Takekawa,[1] Koichi Yamanishi,[1] Hirotaka

Ebina[1,2,4,6,7]

[1] The Research Foundation for Microbial Diseases of Osaka University, Suita, Osaka, Japan

[2] Virus Vaccine Group, BIKEN Innovative Vaccine Research Alliance Laboratories, Institute for Open and

Transdisciplinary Research Initiatives, Osaka University, Suita, Osaka, Japan.

[3] Department of Experimental Genome Research, Research Institute for Microbial Diseases, Osaka University,

Suita, Osaka, Japan

[4] Center for Advanced Modalities and DDS (CAMaD), Osaka University, Suita, Osaka, Japan.

[5] Department of Infectious Diseases and Host Defense, Gunma University Graduate School of Medicine, Maebashi,

Gunma, Japan

[6] Virus Vaccine Group, BIKEN Innovative Vaccine Research Alliance Laboratories, Research institute for

Microbial Diseases, Osaka University, Suita, Osaka, Japan.

[7] Center for Infectious Disease Education and Research (CiDER), Osaka University, Suita, Osaka, Japan.

[8] These authors contributed equally.

*Correspondence to: hebina@biken.osaka-u.ac.jp

Abstract

mRNA vaccines against SARS-CoV-2 were rapidly developed and effective during the pandemic. However, some

limitations remain to be resolved, such as the short-lived induced immune response and certain adverse effects.

Therefore, there is an urgent need to develop new vaccines that address these issues. While live-attenuated vaccines

are a highly effective modality, they pose a risk of adverse effects, including virulence reversion. In the current

study, we constructed a live-attenuated vaccine candidate, BK2102, combining naturally occurring

virulence-attenuating mutations in the *NSP14*, *NSP1*, spike and *ORF7-8* coding regions. Intranasal inoculation with

BK2102 induced humoral and cellular immune responses in Syrian hamsters without apparent tissue damage in the

lungs, leading to protection against a SARS-CoV-2 D614G and an Omicron BA.5 strains. The neutralizing

antibodies induced by BK2102 persisted for up to 364 days, which indicated that they confer long-term protection

against infection. Furthermore, we confirmed the safety of BK2102 using transgenic (Tg) mice expressing human

ACE2 (hACE2), that are highly susceptible to SARS-CoV-2. BK2102 did not kill the Tg mice, even when virus

was administered at a dose of 10^6 plaque-forming units (PFU), while 10^2 PFU of the D614G strain or an attenuated

strain lacking the furin cleavage site (FCS) of the spike was sufficient to kill mice. These results suggest that

BK2102 is a promising live-vaccine candidate strain that confers long-term protection without significant virulence.

Introduction

mRNA- and adenovirus vector-based vaccines have been successfully developed and used in clinical practice

against SARS-CoV-2, the pathogen that caused the COVID-19 pandemic. These vaccines were highly effective,

inducing robust humoral and cellular immunity (Baden et al., 2021; Polack et al., 2020). Nevertheless, certain

drawbacks of SARS-CoV-2 vaccines remain to be addressed, including adverse effects, such as thrombosis, fever,

and fatigue (Yasmin et al., 2023). Further, several boosters of the mRNA vaccines have been required to reactivate

the immune response and increase efficacy against variants. New and improved vaccine modalities are therefore

required for SARS-CoV-2 infection.

Live-attenuated vaccines are among the most effective vaccine modalities, as they induce humoral and cellular

immunity, both systemically and locally (e.g., within mucosal surfaces), conferring protection against various

infectious diseases (Hoft et al., 2017). For example, the live-attenuated poliovirus vaccine developed in 1962 was

highly effective in reducing the spread of the disease (Sabin, 1985). Nevertheless, this modality has certain

disadvantages, the most concerning being the risk of reversion to a virulent state as a result of mutations generated

during viral replication *in vivo*. In fact, vaccine-derived paralytic polio was reported 38 years after the vaccine had

been introduced, representing a threat to uninfected populations (Macklin et al., 2020). Advances in our basic

knowledge of viruses have enabled the design of attenuated strains, such as that in the new type 2 oral polio vaccine

(nOPV2), which is associated with a reduced risk of reversion due to genome modification (Yeh et al., 2020; Yeh et

al., 2023). nOPV2 was recently approved by the World Health Organization, indicating that live-attenuated

vaccines that overcome the risk of reversion are still in demand.

Various mechanisms leading to reduced pathogenicity of SARS-CoV-2 have been reported. A cold-adapted

SARS-CoV-2 strain was isolated through passaging at low temperatures, eventually showing an attenuated

phenotype (Seo and Jang, 2020). We also reported a temperature sensitive (TS) strain with low pathogenicity and

sufficient immunogenicity *in vivo*, achieved through the introduction of multiple TS-related mutations (Yoshida et

al., 2022). In addition to temperature sensitivity, several naturally occurring mutants exhibit attenuated phenotypes.

Passaging SARS-CoV-2 in Vero cells facilitates isolation of strains with deletions in the FCS (PRRAR) at the S1/S2

junction within the spike protein, resulting in low proliferation *in vivo* ((Davidson et al., 2020; Peacock et al., 2021).

Furthermore, a mutant that lacks four amino acids (PRRA) within the FCS did not induce weight loss in hamsters

during challenge experiments(Hoffmann et al., 2020a; Hoffmann et al., 2020b; Johnson et al., 2021; Lau et al.,

2020). Reportedly, S1/S2 cleavage is essential for TMPRSS2-mediated entry into host cells, and the deletion in the

FCS is thought to reduce viral growth in lung cells, which express more TMPRSS2 than cells of the upper

respiratory tract (Hoffmann *et al.*, 2020a; Hoffmann *et al.*, 2020b; Johnson *et al.*, 2021; Lau *et al.*, 2020).

Furthermore, partial loss of *NSP1*, which correlates with a lower viral load and less severe symptoms of infection in

SARS-CoV-2-infected patients, or of *ORF8*, which has been reported as associated with milder infection in humans,

were also associated with attenuated phenotypes (Lin et al., 2021; Young et al., 2020). Recovery of lost segments of

viral genomes carrying deletions is more difficult than the reversion of amino acid substitutions (Bull, 2015). In this

study, we designed and constructed live-attenuated vaccine candidates through a combination of substitutions and

deletions involved in attenuation and reduced risk of virulent reversion. We then evaluated candidate

immunogenicity and safety in animal models providing evidence that BK2102 is a promising vaccine candidate that

confers prolonged protection.

Results

Construction of the live-attenuated vaccine candidate strains

Several genomic alterations are involved in the attenuation of SARS-CoV-2. In this study, we focused on deletions

at three different sites within the viral genome: FCS within the spike protein, NSP1, and ORF7-8 (Supplementary

Fig. 1A). While loss of the FCS inhibits virus-cell fusion mediated by TMPRSS2 in lung cells, partial deletion of

NSP1 has been shown to impair viral proliferation *in vitro*, and the lack of ORF8 has been associated with milder

symptoms and disease outcomes (Johnson et al., 2021; Lin et al., 2021; Young et al., 2020; Zinzula, 2021). We

previously obtained SARS-CoV-2 TS strains showing diverse attenuated phenotypes, and revealed that NSP3

L445F, NSP14 G248V, G416S, and A504V as well as NSP16 V67I were substitutions responsible for such

phenotypes (Yoshida et al., 2022). Each substitution conferred some advantage for the development of an

attenuated vaccine candidate with restricted proliferative capacity in deep regions of the body, such as lungs and

brain. In addition, the presence of deletions is generally considered to confer a lower risk of reversion to a

wild-type genotype when compared to amino acid substitutions. To this end, we designed three candidates by

combining several of the above-described mutations to design a safe live-attenuated vaccine strain (Supplementary

Fig. 1A). The three candidates were inoculated locally into hamsters via the nasal route to mimic a natural infection.

The immunogenicity of candidates 1 and 3 was much greater than that of candidate 2 (Supplementary Fig. 1B).

Candidate 2 has TS-related substitutions in both NSP3 and NSP14 as well as three deletions in the viral genome.

Thus, we hypothesized that this candidate would be excessively attenuated. Candidate 1, which had three deletions

and three TS-responsible substitutions in NSP14, induced neutralizing antibodies when inoculated at a dose of 10^3

PFU. Candidate 3 was observed to induce neutralizing antibodies at a dose of only 10^2 PFU, but the attenuated

phenotypes of this candidate were conferred by only two deletions. Therefore, we speculated that there might be a

higher risk of virulent reversion. Taking these observations into consideration, we selected Candidate 1 for the

vaccine, hereafter referred to as BK2102. The growth dynamics of BK2102 were evaluated at 32 °C and 37 °C (Fig.

1). It proliferated similarly to the wild-type B-1 strain at 32 °C (Fig. 1A), but the infectious virus titer was

significantly lower compared to that of the wild type B-1 strain one day post-infection at 37 °C (Fig. 1B). Therefore,

BK2102 showed a severe TS phenotype but could be amplified by incubating infected cells at 32 °C, which would

also facilitate the manufacturing process.

BK2102 induced humoral and cellular immune responses

To evaluate immunogenicity, BK2102 was intranasally inoculated into Syrian hamsters (10^3 and 10^4 PFU/dose).

Four weeks post-inoculation, serum spike-specific IgG levels were measured using ELISA (Fig. 2A). Neutralizing

antibodies (Fig. 2B) against the D614G strain were detected in 9 of the 10 hamsters in each dose group, with titers

ranging between 2^5 and 2^9 (Fig. 2B, left). Cross-reactivity of the neutralizing antibodies against the delta variant

was also detected in 9 of 10 hamsters (titer range: 2^5–2^8), but that against the BA.5 variant was below the limit of

detection in all hamsters (Fig. 2B, middle and right, respectively). Furthermore, we performed BK2102

immunization of cynomolgus monkeys at a dose of 10^7 PFU, and the serum-neutralizing titer against the D614G

strain was detected in two of the four monkeys (titer range: 2^4–2^8) (Fig. 2C).

A short-lived immune response has been reported for current mRNA vaccines against SARS-CoV-2. For example, a

reduction in neutralizing antibodies was observed in humans after six months (Zhang et al., 2022). In hamsters,

these were undetectable after 250 days (Machado et al., 2023). Therefore, we evaluated the persistence of the immune response induced by BK2102 using a hamster model. We measured neutralizing antibody titers for up to 364 days after inoculation with BK2102. The titer peaks were observed 28 days after the first inoculation and slightly decreased, but were maintained until 364 days post-inoculation with a dose of 10^3 or 10^4 PFU. For example, the neutralizing antibody titer in the sera of hamsters inoculated with 10^3 PFU was 2^8 at 28 days and 2^5 at 364 days post-inoculation. Two of the ten hamsters inoculated at a dose of 10^3 PFU showed neutralizing antibody titers below the detection limit from the beginning, and those of another hamster in the same dose group began to decrease gradually from day 224 and fell below the detection limit on day 364 (Supplementary Fig. 2). However, the neutralizing antibody titers in hamsters inoculated with 10^4 PFU did not exhibit such a decrease during the evaluation period. Remarkably, a single dose of BK2102 was sufficient to induce long-lasting immunity, and there was no need for booster immunization 28 days after the first inoculation (Fig. 2D).

Furthermore, we evaluated cellular immune responses following inoculation with BK2102. Antigen-specific IFN-γ and IL-4 production in spleen cells from inoculated hamsters was measured via ELISA after *in vitr*o re-stimulation with spike or nucleocapsid peptides. As shown in Figure 2E, spike peptide-specific IFN-γ production significantly increased in the splenocytes of BK2102-inoculated hamsters, as did the nucleocapsid peptide-specific IFN-γ production, although in this case it did not reach statistical significance. IL-4 production was significantly increased by spike-peptide stimulation (Fig. 2F). We also analyzed the spike-specific IgG subclass levels (IgG1 and IgG2/3) in hamsters to evaluate the nature of the immune response induced by BK2102 (Supplementary Fig. 3). As other studies have demonstrated that aluminum adjuvant preferentially induces a Th2 response (Marrack et al., 2009), we also administered recombinant spike protein with alum adjuvant as a control. Total IgG levels were elevated in both groups. BK2102-inoculated hamsters showed high levels of IgG2/3, but we could not detect IgG1. IL-4 production and the presence of IgG1 reflects a Th2 response, while IFN-γ production and IgG2/3 are indicative of a Th1 response in hamsters (Kushawaha et al., 2011; Ploquin et al., 2013). Our results therefore suggest that BK2102 mainly induced a Th1 immune response in hamsters.

BK2102 induced protective immunity against SARS-CoV-2 infection

Next, we performed challenge experiments with the SARS-CoV-2 D614G strain or BA.5 variant in order to evaluate whether the immune responses induced by BK2102 would protect against infection. All hamsters inoculated with BK2102 did not lose weight, whereas the naïve hamsters lost approximately 10% of their total body

weight on day four post-challenge with the D614G strain (Fig. 3A). When challenged with the BA.5 variant, all hamsters pre-inoculated with a dose of 10^4 PFU of BK2102 and three of five hamsters pre-inoculated with a dose of 10^3 PFU did not lose weight. However, the rest of the animals in the 10^3 PFU dose group lost 5% of their total body weight, similarly to the naïve group (Fig. 3B).

In addition to body weight change, we determined infectious virus titers in lung homogenates and nasal wash specimens four days-post infection. The number of infectious viruses was significantly lower in hamsters inoculated with BK2102 than in naïve hamsters after challenge with the D614G strain (Fig. 3C and D). Although the cross-reactivity of neutralizing antibodies against the BA.5 variant was below the limit of detection in all hamsters (Fig. 2B, right), no infectious virus was detected following challenge with the BA.5 variant in most of the vaccinated animals (Fig. 3F and G, except for one animal in the 10^4 PFU dose group). In addition, one hamster in the 10^3 PFU dose group showed detectable levels of infectious virus following D614G challenge (Fig. 3C). This result was consistent with the undetectable levels of neutralizing antibodies in this animal (Fig. 2B). Lung tissue damage after the viral challenge was also evaluated in the hamsters. The inflammation score of hamsters inoculated with BK2102 was lower than that of naïve hamsters, regardless of the strain/variant used for the challenge (Fig. 3E and H, respectively). These results suggest that the immune response induced by BK2102 confers protection against infection that is not limited to the SARS-CoV-2 D614G strain, but also includes the BA.5 variant.

Furthermore, to evaluate whether the protection conferred after a full vaccination protocol would persist over time, hamsters with confirmed persistent immunity at day 364 post-inoculation (Fig. 2D) were challenged at day 420 post-inoculation. In elderly naïve hamsters, a body weight loss of approximately 15% was observed seven days after infection with the D614G strain, whereas the BK2102-inoculated hamsters showed a lower weight decrease, with significant differences noted at this time point (Fig. 3I). Therefore, BK2102 induced a prolonged humoral immune response, which contributed to the protection against viral infection in hamsters.

We then evaluated whether BK2102 could inhibit onward transmission, as a previous report of a live-attenuated vaccine generated through codon-pair deoptimization (sCPD9) had suggested that an effective immune response within the nasal cavity would likely prevent it (Nouailles et al., 2023). The naïve group, the spike protein-inoculated (intra-muscularly) group, and the BK2102 intranasal inoculation groups were challenged with the SARS-CoV-2 D614G strain and co-housed with another group of naïve hamsters one day later (Supplementary Fig. 4A). Naïve animals co-housed with hamsters in the naïve or intramuscular vaccine groups showed slight weight loss. However, no weight loss was observed in hamsters co-housed with the hamsters that had been

intranasally inoculated with BK2102 (Supplementary Fig. 4B). Therefore, intranasal inoculation of the BK2102

live-attenuated vaccine effectively prevented onward transmission, in line with a previous report (Nouailles *et al.*,

2023).

BK2102 caused localized tissue damage and conferred a low risk of transmission

Next, we evaluated the safety of BK2102 by assessing the tissue damage during acute infection. The lungs and

whole heads of hamsters at day three post-infection were extracted and fixed with formalin (Fig. 4A). We evaluated

inflammation and detected viral antigens in the lungs and multiple-depth sections of the nasal cavity (Fig. 4B, 4C

and Supplementary Fig. 5). The D614G strain caused broad inflammation within the nasal cavity (from level 1 to 3)

and lungs. Viral antigens were detected in the same areas, consistent with our previous report (Yoshida *et al.*, 2022).

However, in the BK2102-infected hamsters, viral antigens and weak-to-mild tissue damage were observed only in

the anterior area of the nasal cavity, whereas no tissue damage or viral antigens were detected in the posterior area

or lungs.

The replication of BK2102 at the tip of the nasal cavity may facilitate transmission because infectious viruses are

shed into the nasal fluid. In addition, virulent reversion may occur during replication *in vivo*. Therefore, we

evaluated the risk of transmission and reversion to virulence by passaging *in vivo* using hamsters (Fig. 4D).

SARS-CoV-2 A50-18 is a previously isolated TS strain, in which substitutions within the NSP14 protein alone

account for the TS phenotype, without the need for deletions, such as those in NSP1, spike, or other accessory

proteins (Yoshida et al., 2022). The viral genome was detected in all nasal wash specimens from A50-18

strain-infected hamsters during primary infection, and an increase in this amount was observed in subsequently

passaged samples (Fig. 4E), which correlated with progressive weight loss (Supplementary Fig. 6). When we

confirmed the sequence of the viruses detected in samples, we observed that the TS-responsible substitutions in

NSP14 had reverted to the wild-type sequence (Table 3). The viral genome was detected in the nasal wash

specimens from BK2102 infected hamsters during primary infection, but we could not detect it in the samples from

subsequent hamsters, except for in one case (Fig. 4E). In this individual animal, the viral genome was not detected

in the samples from later passages. No weight loss was observed in any of the primary or subsequently infected

hamsters, and we did not detect changes to the wild-type sequence (Table 3, Supplementary Fig. 6). Overall, our

results suggest that BK2102 is a safe live-attenuated vaccine candidate with a low risk of virulent reversion.

BK2102 showed a favorable safety profile in Tg mice

hACE2 Tg mice are also used as animal models of SARS-CoV-2 infection (Asaka et al., 2021; Bao et al., 2020). We established a mouse line expressing hACE2 driven by the CAG promoter, and these mice were used to evaluate the safety of BK2102 live-attenuated vaccine candidate. hACE2 expression was detected not only in the respiratory tract, but also in various tissues such as the central nervous system, heart, skeletal muscle, digestive system (except the small intestine), spleen, and testis (Supplementary Fig. 7). We evaluated the survival rates of Tg mice infected with various SARS-CoV-2 strains. All of the mice died by infection with the D614G strain and even with the FCS deleted B-1 (B-1 ΔFCS) strain, previously established attenuated phenotype, within six days after receiving a dose of 10^2 PFU (Fig. 5A). Meanwhile, a higher survival rate was observed in mice infected with the L50-33 and A50-18 strains, which were previously isolated TS and live-attenuated strains (Yoshida *et al.*, 2022), even at a dose of 10^5 PFU. However, one mouse in each group infected with 10^4 and 10^5 PFU of the L50-33 strain died 10 days post-infection that is four days later than those in the D614G strain-infected group. This time lag before death was expected as the virus could have reverted during replication *in vivo*, being able to grow in deeper regions of the body. Thus, we evaluated the presence of infectious virus in the lungs and brains of mice that died following infection with the D614G, B-1 ΔFCS, and L50-33 strains. Infectious virus titers in the lungs were approximately 2.40 to 5.64 $\log_{10}$ PFU/g, while those in the brains were higher, at approximately 5.75 to 8.69 $\log_{10}$ PFU/g (Table 4). We noticed that mice exhibiting head nodding, intense running, jumping and repeated falling, died despite a generally mild inflammation in the lungs at necropsy (data not shown). These results suggest that Tg mice were killed due to replication of SARS-CoV-2 in the brain rather than in the lungs. Sanger sequencing analysis of viruses in the lungs and brains of mice that died following infection with the L50-33 strain revealed that the 445F substitution in NSP3, responsible for the TS phenotype of this strain, had reverted to the wild-type amino acid leucine (TTT → TTG), which is the same in the D614G and B-1 ΔFCS strains (CTT, Table 4). These results suggest that the mice died due to viral proliferation in the brain, where a small virus population lost its temperature sensitivity, becoming virulent. This mouse is a highly susceptible model of SARS-CoV-2 virus infection able to detect a few temperature-sensitive revertant viruses. In contrast to L50-33 with the NSP3-based TS phenotype, A50-18, harboring three TS-responsible substitutions in NSP14, did not kill any mice. Moreover, in the case of BK2102, no mice died following infection, even at a dose of 10^6 PFU (Fig. 5B). Therefore, BK2102 is considered to have a low risk of virulent reversion, thus representing a suitable candidate for a safe live-attenuated vaccine.

Discussion

In the present study, we evaluated candidates for a new live-attenuated SARS-CoV-2 vaccine. The candidate harboring more mutations showed lower immunogenicity than those harboring less mutations (Supplementary Fig. 1A). Presumably, too many mutations would lead to little viral replication in the body, resulting in weaker immunogenicity. Deletions within viral genes are less prone to mutate back to the wild-type genotype, and TS-associated substitutions control the transfer of virus to deep or warmer regions of the body, including the brain, which makes them an attractive backbone for the development of safe live-attenuated vaccines. We designed our BK2102 vaccine candidate to harbor three deletions in addition to the TS-associated substitutions.

We previously reported that TS strains replicate only in the anterior regions of the nasal cavity and cannot proliferate in the posterior areas or lungs of hamsters. However, these isolated TS strains entailed a risk of reversion to virulence (Yoshida et al., 2022). K18-hACE2 Tg mice were reported to die after infection with several SARS-CoV-2 strains due to viral proliferation in the brain (Natekar et al., 2022). So, to better assess the safety of BK2102, we generated and used CAG-hACE2 Tg mice in addition to hamsters. This mouse model proved to be highly susceptible because the B-1 ΔFCS strain killed the mice, despite other S1/S2 cleavage-deficient strains showing only an attenuated phenotype in wild-type hamsters (Fig. 5A). However, the TS strains (L50-33 and A50-18) showed an attenuated phenotype even in this mouse model, highlighting the superior characteristics of TS in limiting viral replication and the replication-permissive regions by temperature. The TS-responsible substitutions in NSP14 were more stable than those in NSP3, as L50-33 killed one mouse in each of the dose groups tested, whereas A50-18 did not. Further, infectious viruses without TS mutations were detected in the central nervous system and respiratory tracts of dead mice following infection with the L50-33 strain (Table 4). These results suggest that the mouse model we employed is particularly suitable for the assessment of safety and the evaluation of TS strains' entry into the nervous system, because it enables the detection of even a small population of revertant viruses. In addition, there is concern regarding SARS-CoV-2 infection via the intranasal route causing damage to the central nervous system (Jha et al., 2021; Kumar et al., 2020). The use of this animal model could address this concern.

BK2102, which did not proliferate in the brains of Tg mice even at a dose of 10^6 PFU, which was 10,000 times higher than that killed mice in the case of the D614G or B-1 ΔFCS strains (Fig. 5B), was not detected during passaging *in vivo* using naïve hamsters. The A50-18 strain shares almost the same genetic background to that of BK2102 but without three deletion mutations. This TS strain showed an attenuated phenotype during primary

infection, but facilitated its reversion to virulence during viral replication in hamsters, leading to the transmission of

the virulent strain to naive animals, as shown in Fig. 4E. In contrast, BK2102, which combines multiple defective

mutations, controls replication in host animals. Even when infected with a dose of 5×10^5 PFU, BK2102 viruses did

not proliferate enough to be transmitted among hamsters. We hypothesized that even if TS-related substitutions are

lost at the time of primary infection, the remaining deletion mutations ensure low replication efficiency, preventing

transmission. Indeed, mutation in the FCS coding region was reported to lead to lower proliferation (Lau *et al.*,

2020; Sasaki et al., 2021; Wang et al., 2021a) and transmission *in vivo* (Peacock *et al.*, 2021).

Our vaccine candidate BK2102 induced humoral and cellular immune responses in hamsters, and those inoculated

with a high dose were protected from challenge with the heterologous BA.5 variant, even though the neutralizing

titer in serum was below the limit of detection (Fig. 2B, right). The nucleocapsid proteins of many coronaviruses

are highly immunogenic and expressed abundantly in infected cells, making them easy targets for antigen-specific

T cells (Cong et al., 2020; Dutta et al., 2020; Hasanpourghadi et al., 2023). Other studies have reported the benefit

of vaccination with the nucleocapsid protein of SARS-CoV-2, showing protective immunity in animal models

receiving the nucleocapsid protein alone (Primard et al., 2023) or in combination with the spike protein (Chiuppesi

et al., 2022; Hasanpourghadi et al., 2023). In our hamster study, we detected cellular immune responses against the

spike protein, but not against the nucleocapsid protein. BK2102 may induce a cellular immune response against

various structural proteins of SARS-CoV-2, thereby conferring protection against several variants. The potential

induction of mucosal immunity was also considered. Live-attenuated influenza vaccines have been reported as

effective against a broad range of variants owing to the efficient humoral and cellular immune responses they

induce, even in mucosal tissues (Thwaites et al., 2023). Furthermore, the SARS-CoV-2 live-attenuated vaccine

candidate sCPD9, constructed through codon-pair deoptimization, induced neutralizing antibodies in the nasal fluid

of hamsters as well (Nouailles et al., 2023). BK2102 was also inoculated via the intranasal route, and inhibition of

onward transmission was observed, as reported for the sCPD9 candidate (Supplementary Fig. 4). Therefore, we

assumed that BK2102 also induced a mucosal immune response in addition to systemic humoral and cellular

immunity, contributing to protection against infection with the BA.5 variant.

The key to developing live-attenuated vaccines is a balance between immunogenicity and safety. Most

live-attenuated vaccine candidates, including the sCPD9 mentioned above and CoviLiv[TM], which has entered a

phase III clinical trial, are evaluated with an emphasis on immunogenicity in basic research using animal models

(Wang et al., 2021b). In this study, we carefully evaluated not only immunogenicity, but also safety. Whether our

candidate will be the best for human use remains to be confirmed in human trials. In fact, we inoculated this

candidate into monkeys, but these were not as susceptible as the hamsters and required a very high titer of the virus

for inducing immunity. Therefore, considering the possible differences in susceptibility, we may need to optimize

dose. Notably, our vaccine candidate, BK2102, was difficult to passage *in vivo* due to its TS substitutions that

enabled to control or limit proliferation and thus was not transferred to the central nervous system. We found that,

while attenuated strains with amino acid substitutions carry a risk of reversion to virulence, it is still possible to

construct safer live-attenuated vaccine candidate strains by combining multiple modifications related to diverse

attenuated phenotypes. Among these modifications, deletions are especially useful for their effect on reducing the

risk of reversion to virulence, and TS-responsible substitutions significantly decrease viral replication in deep

regions of the body. Numerous viruses, such as the Japanese encephalitis and influenza viruses, have been reported

to replicate in the brain and lung (Desai et al., 1995; Weinheimer et al., 2012). The strategy utilizing multiple

modifications may thus be effective for the development of live-attenuated vaccines against these viruses.

Material and methods

Cells and viruses

Vero cells were purchased from ATCC and maintained in D-MEM supplemented with 10% FBS, penicillin (100

U/mL), and streptomycin (0.1 mg/mL). VeroE6/TMPRSS2 cells were obtained from the Japanese Collection of

Research Bioresources (JCRB) cell bank and cultured in D-MEM supplemented with 10% FBS, penicillin (100

U/mL), streptomycin (0.1 mg/mL), and G-418 (1 mg/mL). We previously constructed baby hamster kidney (BHK)

cells constitutively expressing the human angiotensin-converting enzyme 2 (hACE2) (BHK/hACE2 cells)

(Okamura et al., 2023), and these cells were maintained in MEM, supplemented with 10% FBS,

penicillin-streptomycin, and puromycin (3 μg/mL). The SARS-CoV-2 strains used in this study are listed in Table 1.

The SARS-CoV-2 B-1 (D614G) strain was isolated from a clinical specimen, and TS derivative strains were

obtained through random mutagenesis of this clinical isolate, as previously reported (Yoshida et al., 2022).

SARS-CoV-2 delta and omicron variant were obtained from the Research Foundation for Microbial Diseases of

Osaka University and the National Institute of Infectious Disease of Japan, respectively.

Construction of viruses through circular polymerase extension reaction (CPER)

SARS-CoV-2 live-attenuated vaccine candidate strains and B-1 ΔFCS strain has genetic backgrounds similar to that

of B-1 strain, in combination with following naturally occurring virulence-attenuating mutations. The mutations in the ORF7a-8, *NSP3*, *NSP14*, and *NSP16* coding regions are the same as those in the genomes of temperature-sensitive virus strains (L50-33, A50-18 and H50-11), described in Yoshida et al., 2022. The mutations in the spike FCS ($_{679}$NSPRRARSV$_{687}$ → I) and the NSP1 ($_{32}$GDSVEEVL$_{39}$) are the same as those in the genomes of a laboratory strain described in Davidson *et al.,*2020 and of a clinical isolate (accession: LC521925), respectively.

For construction of the strains we used the CPER method, in which 11 PCR-generated cDNA fragments covering the viral full genome plus another DNA fragment with controlling sequences are stitched into a circular DNA that can produce viral genomic RNA upon introduction to cells (Okamura et al., 2023; Torii et al., 2021). Primers used to prepare PCR fragments are listed in Table 2. The PCR and CPER reaction were performed using PrimeSTAR GXL DNA polymerase (Takara Bio, Cat# R050A). The CPER product was transfected into BHK/hACE2 cells using Lipofectamine LTX (Thermo Fisher Scientific). The cells were incubated at 32 °C in a CO_2 incubator for one week. At this point, the supernatants were collected and transferred to 6-well plates which had been pre-seeded with VeroE6/TMPRSS2 cells. The viruses contained in the supernatants of these cells were collected when cytopathic effects (CPEs) were clearly observed.

Virus titration

The infectious titer of SARS-CoV-2 was determined based on the median tissue culture infectious dose (TCID$_{50}$) or plaque formation assay (PFA). In order to obtain the TCID$_{50}$, virus-containing samples were serially diluted with D-MEM supplemented with 2% FBS. Fifty μL of each diluted sample were used to infect Vero cells in 96-well plates. The cells were fixed with 10% formalin after incubating at 32 °C for four days and stained with crystal violet solution. The TCID$_{50}$ was calculated using the Behrens-Karber method. For the calculation of PFU, 500 μL of the diluted samples were added to confluent Vero cells in 6-well plates and incubated at 32 °C for 3 h to allow virus adsorption. The supernatant was removed, and the cells were washed with D-PBS. Subsequently, 2 mL of 1% SeaPlaque agarose dissolved in D-MEM and supplemented with 2% FBS were layered on the cells. The plates were incubated at 32 °C in a CO_2 incubator for three days, fixed with 10% formalin, and stained with crystal violet solution. Visible plaques were counted to calculate PFU.

Viral proliferation assay at various temperatures

The SARS-CoV-2 B-1 (D614G) strain or vaccine candidate strain was used to infect Vero cells at a multiplicity of

infection (MOI) of 0.01. Infected cells were cultured at 32 °C or 37 °C, and a part of the supernatant was collected

daily and stored at -80 °C. The viral titers of these samples were determined using the $TCID_{50}$ assay described

above. Each experiment was performed in triplicates.

Quantitative RT-PCR

Viral RNA was extracted from various samples using the IndiSpin Pathogen Kit (INDICAL BIOSCIENCE)

following the manufacturer's instructions. RNA was quantified using a DetectAmp SARS-CoV-2 RT-PCR kit

(Sysmex). Five µL of a 100-fold dilution of the extracted RNA were used to perform the reverse transcription and

subsequent qPCR reactions in a 7500 Fast Real-time PCR System (Applied Biosystems). The reactions were

performed in triplicate.

Neutralization assay

The neutralizing activity of sera was evaluated using authentic SARS-CoV-2. One hundred PFU of authentic virus

were mixed with serially diluted serum samples and incubated at 37 °C for 1 h. The mixtures were then transferred

to confluent Vero cells in 96-well plates. The infected cells were fixed with 10% formalin and stained with crystal

violet solution after an incubation period of three days at 32 °C. Neutralization titers were calculated as the inverse

of the maximum dilution that prevented CPE formation.

Enzyme-linked immune-sorbent assay (ELISA)

Half-well protein high-binding 96-well plates (Greiner) were coated with recombinant SARS-CoV-2 (D614G)

spike or nucleocapsid proteins (SinoBiologicals) dissolved in PBS (50 ng/100 µL/well) and incubated at 4 °C

overnight. 1% BSA PBS was used as blocking solution, and 1% BSA PBS-T was used to dilute the sera, antibodies,

and streptavidin. Antigen-specific IgG antibodies in hamster sera were detected using horseradish peroxidase

(HRP)-labeled goat anti-Syrian hamster IgG H+L Ab (Abcam, 1/30000 dilution). IgG1 and IgG2/3 subclass Abs

were analyzed using biotinylated subclass-specific antibodies (Southern Biotech, 1/100 and 1/200000 dilutions,

respectively) and detected with HRP-labeled streptavidin (Abcam, 1/1000 dilution). Mouse IgG was detected using

horseradish peroxidase-labeled goat anti-mouse IgG H&L (Abcam, 1/100000 dilution). TMB one-component

substrate and stopping solutions (Thermodics) were used for the chromogenic reaction, and the absorbance at 450

nm was measured using a 450 nm microplate reader.

Evaluation of immunogenicity in hamsters

Four-week-old male Syrian hamsters were purchased from Japan SLC Inc. After a one-week housing period, they were first anesthetized via inhalation with 3% isoflurane, followed by intraperitoneal anesthesia with a combination of medetomidine, midazolam, and butorphanol (0.3, 4, and 5 mg/kg, respectively). One thousand or 1×10^4 PFU of BK2102 were administered intra-nasally in a volume of 20 μL to confine administration to the upper respiratory tract. To evaluate the humoral immune response, blood samples were collected from the facial vein using a lancet (MEDIpoint), and neutralizing antibody titers in the sera were measured as described above. To evaluate the cellular immune response, BK2102-inoculated hamsters were euthanized one week post-inoculation, and spleens were collected. The spleens were mechanically crushed with the piston of a syringe and passed through a cell strainer (100 μm). Cell suspensions were treated with RBC lysis buffer (Biolegend) to remove red blood cells, as per manufacturer instructions. The spleen cells were then suspended in RPMI medium supplemented with 10% FBS and antibiotics at a concentration of 1×10^6 cells/mL and stimulated by SARS-CoV-2 spike or nucleocapsid peptide pools, consisting mainly of 15-mer sequences with 11-amino-acid (aa) overlap (Miltenyi Biotec). Negative controls for non-stimulated cells were included in the test. Stimulated cells were incubated at 37 °C for three days, and supernatants were collected and stored at -80 °C until use. Interferon-gamma (IFN-γ) and interleukin-4 (IL-4) were quantified with commercially-available kits (MABTECH AB and FineTest, respectively) following the manufacturer's protocol. We also performed a challenge assay to investigate whether these immune responses were effective in protecting against other variants. Four weeks post-inoculation, 3×10^5 PFU of SARS-CoV-2 D614G strain or TY41-702 strain (Omicron variant BA.5) were used to challenge the hamsters through the intra-nasal route to target the lower respiratory tract. The weights of the hamsters were monitored daily, and they were euthanized four days after the challenge. Lungs were divided, cut into small pieces, and homogenized with a biomasher II device in 500 μL of D-MEM. Supernatants were collected as lung homogenates after centrifugation at 300 ×g for 5 min at 4 °C. Nasal wash specimens were obtained by flushing one mL of D-PBS into the nasal cavity from the trachea in the direction of the nose where it was collected. Infectious virus titers in these samples were evaluated using a plaque formation assay, as described above. To evaluate immune response persistence, hamsters inoculated with the BK2102 vaccine candidate were maintained for 364 days post-infection. Blood samples were collected from the facial vein using a lancet (MEDIpoint), and neutralizing antibody titers were measured. At 420 days after

inoculation, a viral challenge assay was performed to evaluate whether the immunity contributing to infection

protection was maintained. Hamsters were infected with 3×10^5 PFU of the SARS-CoV-2 D614G strain, and weight

changes were monitored daily for 10 days post-infection.

Evaluation of immunogenicity in monkeys

Two-to three-year-old male and female cynomolgus monkeys were purchased from Hamri Inc. After a one-week

acclimatization housing period, they were sedated with a mixture of ketamine (5 mg/kg) and xylazine (2 mg/kg)

administered intramuscularly. Ten million PFU of BK2102 were administered intranasally at a volume of 1.6 mL.

To evaluate the humoral immune response, blood samples were collected from the radial vein, femoral vein, or

abdominal vena cava using a syringe, and neutralizing antibody titers in the sera were measured as described above.

Evaluation of BK2102 pathogenicity in hamsters

Four-week-old male Syrian hamsters were obtained from Japan SLC Co. Ltd. After acclimatization, 1×10^4 PFU of

BK2102 or 1×10^5 PFU of the SARS-CoV-2 D614G strain were used for infection, as described above. These and

naïve hamsters were euthanized three days post-infection. The heads and lungs were collected and fixed in 10%

formalin. Sections of the head were prepared to expose different regions of the nasal cavity and were stained with

hematoxylin-eosin or an immunohistochemistry (IHC) staining kit using a SARS-CoV-2 spike RBD-specific

antibody (Sino Biological). The damage score of each section was defined as 0: not remarkable (< 10%); 1:

minimal (10-50%); and 2: mild (50-70%).

***In vivo* passage of BK2102 in hamsters**

Four-week-old male Syrian hamsters were obtained from Japan SLC Co. Ltd. After a one-week acclimatization

period, 1×10^4 PFU of BK2102 or A50-18 strain (a TS mutant isolated in a previous report) in a volume of 20 μL

were used to infect the hamsters intranasally. The hamsters were observed daily, and their body weights were

measured at 0 and 3 days post-infection. At this point, hamsters were euthanized, and nasal wash specimens were

collected with 500 μL D-PBS. After filtration through 0.45-μm and 0.22-μm filters, 100 μL were used to infect a

new group of naive hamsters. The passage of inoculum was repeated three times, and nasal wash samples were

collected at every passage. The viral genome in these nasal wash samples was quantified via qPCR, as described

above. Sanger sequencing was performed to analyze the mutations introduced to generate an attenuated phenotype.

Generation of human ACE2-transgenic mice

The transgene was prepared as described previously (Ikawa et al., 1995; Okabe et al., 1997). Briefly, the hACE2-coding sequence was amplified via PCR with the following primers: 5′-aatctagagccgccgccgccatgtcaagctcttcctggctccttc-3′ and 5′-aaactcgagctaaaaggaggtctgaacatcatca-3′, using human testis cDNA as the template. The *Xba*I and *Xho*I sites included in the PCR primers were used to introduce the amplified hACE2 cDNA into a pCAG1.1 expression vector (https://www.addgene.org/173685/) containing the chicken *beta-actin* promoter and cytomegalovirus enhancer, the *beta-actin* intron, and the rabbit *globin* poly-adenylation signal. The transgene fragment was excised using *Sac*I and *Pac*I and gel-purified. Transgenic mouse lines were generated by injecting purified transgene fragments into C57BL/6N×C57BL/6N fertilized eggs. A total of 350 DNA-injected eggs were transplanted into pseudopregnant mice, resulting in 32 newborn pups. Three of these were transgenic, and the first line was established as Tg (CAG-hACE2)01Osb. The Tg mice were kept of a B6D2F1 background. Expression of hACE2 in each organ was confirmed via western blotting. Briefly, organ homogenates were prepared in radioimmunoprecipitation assay buffer containing a proteinase inhibitor cocktail (Thermo Scientific) using a bead mill homogenizer (Fischer Scientific). Protein concentrations were measured via BCA assay (Pierce), and 10 μg of protein were subjected to SDS-PAGE and subsequent western blotting. hACE2 and beta-actin were detected using a rabbit polyclonal antibody against hACE2 (Abcam: ab15348) and a mouse monoclonal antibody to beta-actin (Abcam: ab8226) with Can Get Signal solutions (TOYOBO).

Evaluation of safety in transgenic mice

Seven- to nine-week-old Tg mice were anesthetized via inhalation of 3% isoflurane, then receiving a combination of medetomidine, midazolam, and butorphanol (0.3, 4, and 5 mg/kg, respectively) intraperitoneally. Mice were infected with SARS-CoV-2 in a volume of 20 μL, and their weights were measured daily. Mice that reached humane endpoints, such as difficulty walking or rapid weight loss, were euthanized by bleeding under isoflurane anesthesia. The brain, olfactory bulbs, nasal turbines, and lungs were collected. Mice infected with live-attenuated strains that did not reach the humane endpoint were also euthanized 14 days post-infection.

Statistical analysis

Two-way analysis of variance (ANOVA) was used for comparing body weight, and one-way ANOVA was used for comparing infectious titer, cytokine levels, and inflammation score. These data were analyzed using GraphPad Prism 9.4.1 software. Statistical significance was set at a p-value < 0.05.

ACKNOWLEDGMENTS

We appreciate the assistance of Paola Miyazato, Manae Morishima, and Kaori Yamamoto from The Research Foundation for Microbial Diseases of Osaka University (BIKEN). The authors thank Shiho Torii, Chikako Ono, and Yoshiharu Matsuura for their CPER technical support. We would like to thank Mitsuko Mori for generation of human ACE2-transgenic mice. The authors also acknowledge the NGS Core facility of the Genome Information Research Center at the Research Institute for Microbial Diseases of Osaka University for their support with next-generation sequencing analyses. This work was conducted as part of "The Research Foundation for Microbial Diseases of Osaka University Project for Infectious Disease Prevention".

FUNDING

This work was supported by BIKEN, Japan Agency for Medical Research and Development (AMED) grants (JP20pc0101047, JP23fa627002) and Central institute for experimental animals grant (JP23fa627006).

AUTHOR CONTRIBUTIONS

M.S. and S.O. conducted and controlled most of the experiments and prepared the manuscript. W.K. performed the *in vivo* passage assays. T.N. and M.I. established Tg-mice. A.Y., and S.G. constructed various recombinant viruses. T.G., and H.S. supported the experiments related to prolonged protection and *in vivo* passage assays. L.S. conducted the cellular immunity assays. S.T., K.Y., and H.E. designed and managed the study.

DECLARATION OF INTERESTS

M.S., S.O., T.G., S.G., H.S., S.L., A.Y., S.T., K.Y., and H.E. are employed by BIKEN. We report that S.O., A.Y., and H.E. are named as the inventors on a patent application that describes the use of the TS mutants and S.T., H.E., S.O., and A.Y. in another patent application relating to the attenuated strain. S.T. and H.E. are managers of BIKEN. K.Y. is the director-general of BIKEN.

 Figure legends

Fig. 1 Growth dynamics of the vaccine candidate strain at different temperatures

Vero cells were infected with the wild-type parent B-1 (D614G) or the BK2102 vaccine candidate strains at a

multiplicity of infection (MOI) = 0.01, and virus titers in the supernatants were determined for samples harvested

every day, after incubating at 32 °C (A) or 37 °C (B). Infectious virus titers were determined using the $TCID_{50}$

method. Symbols indicate the average of three independent experiments, and error bars represent the SD. Days

post-infection are indicated on the x-axis.

Fig. 2 Immunogenicity of the vaccine candidate in hamsters

(A) Spike-specific IgG in the sera of BK2102-inoculated hamsters and mock-treated hamsters. Spike-specific IgG

in sera obtained four weeks post-inoculation was detected by ELISA. Symbols depict data of individual hamsters,

and bars correspond to the median value. The limit of dilution is indicated in the x- axis.

(B) Neutralizing antibodies in the sera were induced by BK2102-inoculated hamsters and mock-treated hamsters.

Neutralizing antibodies in the sera were measured at day 28 post-inoculation using the following authentic

SARS-CoV-2 strains: wild-type D614G (left), Delta (middle), and BA.5 (right). Symbols represent titers of

individual animals, and the bars indicate the median. The dotted line represents the assay's limit of detection

(LOD).

(C) Immune response in monkeys. Neutralizing antibodies in the sera of BK2102-inoculated monkeys was

measured at the indicated time points post-inoculation. The data for individual monkeys are shown.

(D) Neutralizing antibodies persist in hamsters for at least 364 days. The neutralizing antibody titer against the

authentic D614G wild-type strain was measured periodically in the sera of hamsters inoculated with BK2102 (once

or twice at the indicated doses) for about a year. Symbols represent the mean of 9 to 10 animals, and error bars

represent the SD.

(E and F) Evaluation of the cellular immune response in BK2102-inoculated hamsters. Splenocytes were collected

at two weeks post-inoculation and were stimulated *in vitro* with spike or nucleocapsid peptide pools. IFN-γ (E) and

IL-4 (F) in the supernatants were measured with commercially available ELISA kits (MABTECH AB and FineTest,

respectively). Symbols represent titers of individual animals, and bars indicate the median. For statistical analysis,

one-way ANOVA with Tukey's multiple comparison test was performed (ns, not significant; [*], $p < 0.05$; [**], $p <$

0.01).

Fig. 3 BK2102 induces protective immunity.

(A and B) BK2102 protects hamsters against homologous and heterologous virus challenges. Hamsters that received a full vaccination protocol with the indicated doses of BK2102 were challenged with wild-type D614G (A) or BA.5 (B) strains, and their body weight was monitored for four days. Body weight is expressed as a percentage of the initial weight. Two-way ANOVA with Tukey's multiple comparison test was performed for statistical analysis ([**], $p < 0.01$; [****], $p < 0.0001$).

(C, D, F and G) The infectious virus titer in the lungs and nasal wash specimens taken on day four post-challenge was measured via a plaque assay for the wild-type D614G strain (C and D) and for the BA.5 strain (F and G). One-way ANOVA with Dunnett's multiple comparison test was performed for statistical analysis (ns, not significant; [*], $p < 0.05$; [**], $p < 0.01$; [***], $p < 0.001$).

(E and H) Lung inflammation scores were determined via H&E staining of D614G- (E) and BA.5-challenged (H) hamsters. The percentage of the disrupted area in the entire visual field was classified as 0: not remarkable ($< 10\%$); 1: minimal ($10–50\%$); and 2: mild ($50–70\%$). One-way ANOVA with Tukey's multiple comparison test was performed for statistical analysis (ns, not significant; [*], $p < 0.05$).

(I) Weight changes after the challenge assay one year post-inoculation with BK2102. Hamsters inoculated with BK2102 were challenged with the wild-type D614G strain at 3×10^5 PFU on 420 days. Nine-month-old elder hamsters were used as the naïve group. The symbols represent the average weight of the hamsters, and error bars indicate the mean SD. Two-way ANOVA with Tukey's multiple comparison test was performed for statistical analysis ([****], $p < 0.0001$).

Fig. 4 BK2102 caused localized tissue damage and posed as low risk of transmission.

(A) Scheme for the evaluation of tissue damage in acute infection with BK2102 in a hamster model. The wild-type D614G strain was used as a positive control.

(B) Inflammation score of nasal cavity sections and lungs determined via H&E. The percentage of the disrupted area in the entire visual field was classified as 0: not remarkable ($< 10\%$); 1: minimal ($10–50\%$); 2: mild ($50–70\%$), respectively.

(C) SARS-CoV-2 spike protein staining in the nasal cavity sections and lungs determined via immunohistochemistry using a SARS-CoV-2 spike RBD-specific antibody. The proportion of positive cells in the

entire visual field was classified as 0: not remarkable (< 10%); 1: minimal (10–50%); and 2: mild (50–70%),

respectively.

(D) Scheme for the evaluation of BK2102 transmission via *in vivo* passage in hamsters. The TS-strain A50-18 was

used as a positive control.

(E) Ct values obtained for the RT-PCR performed using RNA extracted from the nasal wash specimens.

Fig. 5 BK2102 showed a highly safe phenotype in Tg mice

(A and B) Survival rate of Tg mice infected with the wild-type D614G, B-1 ΔFCS, L50-33, and A50-18 TS strains

(A) and BK2102 (B).

Supplementary figure legends

Supplementary fig. 1 Characteristics of the vaccine candidates.

(A) Candidate vaccine constructs. The abbreviations shown in the figure are respectively, ORF: Open Reading

Frame, NSP: Non-Structural Protein, FCS: Furin Cleavage Site.

(B) Immunogenicity of vaccine candidates in hamsters. Neutralizing antibody titers in the sera were measured 21

days post-inoculation.

Supplementary fig. 2 Persistence of the neutralizing antibodies induced by BK2102 in each group.

Results for individual mice in the group used in the experiment of Figure 2D are shown. Light blue circles and

dotted line correspond to mice inoculated with one dose of 10^3 PFU; light blue circles and solid line: 10^3 PFU two

doses; dark blue circles and dotted-line: one dose of 10^4 PFU one dose; dark blue circles and solid line: two doses

of 10^4 PFU. ¶ Hamsters died due to aging, fighting, or mishandling.

Supplementary fig. 3 Analysis of the IgG subclasses induced by BK2102

SARS-CoV-2 spike-specific total IgG and IgG subclasses (IgG1 and IgG2/3) were measured via ELISA of serum

from inoculated hamsters, 28 days post-inoculation. Recombinant SARS-CoV-2 spike protein mixed with an alum

adjuvant was used as a positive control for the induction of both IgG1 and IgG2/3.

Supplementary fig. 4 Evaluation of BK2102 onward transmission in hamsters

(A) Scheme of the onward transmission experiment. Hamsters that received the full vaccination protocol with the

indicated doses of BK2102 and recombinant SARS-CoV-2 spike protein mixed with an alum adjuvant were

challenged with the wild-type D614G strain. Vaccinated and infected animals were co-housed with naïve hamsters

at one day post-infection.

(B) Body weight was monitored for six days.

Supplementary fig. 5 Evaluation of the tissue damage induced by BK2102

H&E staining of nasal cavity sections and lungs is shown for hamsters infected with the wild-type D614G strain or

the BK2102 vaccine candidate. IHC results using a SARS-CoV-2 spike RBD-specific antibody are shown in for the

same sections. Scale bar: 200 μm.

Supplementary fig. 6 BK2102 showed a low risk of transmission

The body weights of hamsters inoculated with the A50-18 TS and BK2102 strains obtained in the *in vivo* passage

experiment detailed in Figure 4D are shown. For statistical analysis, two-way ANOVA with Holm-Sidak's multiple

comparison test was performed (ns, not significant; **, $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

Supplementary fig. 7 Expression of hACE2 in Tg mice

The expression of hACE2 in various tissues of Tg mice was detected via western blotting using an anti-human

ACE2 antibody (ab15348).

Asaka, M. N., Utsumi, D., Kamada, H., Nagata, S., Nakachi, Y., Yamaguchi, T., Kawaoka, Y., Kuba, K.,
and Yasutomi, Y. (2021). Highly susceptible SARS-CoV-2 model in CAG promoter-driven hACE2-transgenic
mice. JCI Insight *6*. 10.1172/jci.insight.152529.
Baden, L. R., El Sahly, H. M., Essink, B., Kotloff, K., Frey, S., Novak, R., Diemert, D., Spector, S. A.,
Rouphael, N., Creech, C. B., et al. (2021). Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine.
N Engl J Med *384*, 403-416. 10.1056/NEJMoa2035389.
Bao, L., Deng, W., Huang, B., Gao, H., Liu, J., Ren, L., Wei, Q., Yu, P., Xu, Y., Qi, F., et al. (2020).
The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. Nature *583*, 830-833.
10.1038/s41586-020-2312-y.
Bull, J. J. (2015). Evolutionary reversion of live viral vaccines: Can genetic engineering subdue it?
Virus Evol *1*. 10.1093/ve/vev005.
Chiuppesi, F., Nguyen, V. H., Park, Y., Contreras, H., Karpinski, V., Faircloth, K., Nguyen, J., Kha,
M., Johnson, D., Martinez, J., et al. (2022). Synthetic multiantigen MVA vaccine COH04S1 protects
against SARS-CoV-2 in Syrian hamsters and non-human primates. NPJ Vaccines *7*, 7.
10.1038/s41541-022-00436-6.
Cong, Y., Ulasli, M., Schepers, H., Mauthe, M., V'Kovski, P., Kriegenburg, F., Thiel, V., de Haan,

C.A.M., and Reggiori, F. (2020). Nucleocapsid Protein Recruitment to Replication-Transcription
Complexes Plays a Crucial Role in Coronaviral Life Cycle. J Virol *94*. 10.1128/JVI.01925-19.
Davidson, A.D., Williamson, M.K., Lewis, S., Shoemark, D., Carroll, M.W., Heesom, K.J., Zambon, M.,
Ellis, J., Lewis, P.A., Hiscox, J.A., and Matthews, D.A. (2020). Characterisation of the transcriptome
and proteome of SARS-CoV-2 reveals a cell passage induced in-frame deletion of the furin-like cleavage
site from the spike glycoprotein. Genome Med *12*, 68. 10.1186/s13073-020-00763-0.
Desai, A., Shankar, S.K., Ravi, V., Chandramuki, A., and Gourie-Devi, M. (1995). Japanese encephalitis
virus antigen in the human brain and its topographic distribution. Acta Neuropathol *89*, 368-373.
10.1007/BF00309631.
Dutta, N.K., Mazumdar, K., and Gordy, J.T. (2020). The Nucleocapsid Protein of SARS-CoV-2: a Target
for Vaccine Development. J Virol *94*. 10.1128/JVI.00647-20.
Hasanpourghadi, M., Novikov, M., Ambrose, R., Chekaoui, A., Newman, D., Ding, J., Giles-Davis, W.,
Xiang, Z., Zhou, X.Y., Liu, Q., et al. (2023). Heterologous chimpanzee adenovirus vector immunizations
for SARS-CoV-2 spike and nucleocapsid protect hamsters against COVID-19. Microbes Infect *25*, 105082.
10.1016/j.micinf.2022.105082.
Hoffmann, M., Kleine-Weber, H., and Pohlmann, S. (2020a). A Multibasic Cleavage Site in the Spike Protein
of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Mol Cell *78*, 779-784 e775.
10.1016/j.molcel.2020.04.022.
Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S., Schiergens, T.S.,
Herrler, G., Wu, N.H., Nitsche, A., et al. (2020b). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2
and Is Blocked by a Clinically Proven Protease Inhibitor. Cell *181*, 271-280 e278.
10.1016/j.cell.2020.02.052.
Hoft, D.F., Lottenbach, K.R., Blazevic, A., Turan, A., Blevins, T.P., Pacatte, T.P., Yu, Y., Mitchell,
M.C., Hoft, S.G., and Belshe, R.B. (2017). Comparisons of the Humoral and Cellular Immune Responses
Induced by Live Attenuated Influenza Vaccine and Inactivated Influenza Vaccine in Adults. Clin Vaccine
Immunol *24*. 10.1128/CVI.00414-16.
Ikawa, M., Kominami, K., Yoshimura, Y., Tanaka, K., Nishimune, Y., and Okabe, M. (1995). A rapid and
non-invasive selection of transgenic embryos before implantation using green fluorescent protein (GFP).
FEBS Lett *375*, 125-128. 10.1016/0014-5793(95)01162-8.
Jha, N.K., Ojha, S., Jha, S.K., Dureja, H., Singh, S.K., Shukla, S.D., Chellappan, D.K., Gupta, G.,
Bhardwaj, S., Kumar, N., et al. (2021). Evidence of Coronavirus (CoV) Pathogenesis and Emerging Pathogen
SARS-CoV-2 in the Nervous System: A Review on Neurological Impairments and Manifestations. J Mol
Neurosci *71*, 2192-2209. 10.1007/s12031-020-01767-6.
Johnson, B.A., Xie, X., Bailey, A.L., Kalveram, B., Lokugamage, K.G., Muruato, A., Zou, J., Zhang,
X., Juelich, T., Smith, J.K., et al. (2021). Loss of furin cleavage site attenuates SARS-CoV-2
pathogenesis. Nature *591*, 293-299. 10.1038/s41586-021-03237-4.
Kumar, A., Pareek, V., Prasoon, P., Faiq, M.A., Kumar, P., Kumari, C., and Narayan, R.K. (2020). Possible
routes of SARS-CoV-2 invasion in brain: In context of neurological symptoms in COVID-19 patients. J
Neurosci Res *98*, 2376-2383. 10.1002/jnr.24717.
Kushawaha, P.K., Gupta, R., Sundar, S., Sahasrabuddhe, A.A., and Dube, A. (2011). Elongation factor-2,
a Th1 stimulatory protein of Leishmania donovani, generates strong IFN-gamma and IL-12 response in
cured Leishmania-infected patients/hamsters and protects hamsters against Leishmania challenge. J
Immunol *187*, 6417-6427. 10.4049/jimmunol.1102081.
Lau, S.Y., Wang, P., Mok, B.W., Zhang, A.J., Chu, H., Lee, A.C., Deng, S., Chen, P., Chan, K.H., Song,
W., et al. (2020). Attenuated SARS-CoV-2 variants with deletions at the S1/S2 junction. Emerg Microbes
Infect *9*, 837-842. 10.1080/22221751.2020.1756700.
Lin, J.W., Tang, C., Wei, H.C., Du, B., Chen, C., Wang, M., Zhou, Y., Yu, M.X., Cheng, L., Kuivanen,
S., et al. (2021). Genomic monitoring of SARS-CoV-2 uncovers an Nsp1 deletion variant that modulates
type I interferon response. Cell Host Microbe *29*, 489-502 e488. 10.1016/j.chom.2021.01.015.
Machado, R.R.G., Walker, J.L., Scharton, D., Rafael, G.H., Mitchell, B.M., Reyna, R.A., de Souza, W.M.,
Liu, J., Walker, D.H., Plante, J.A., et al. (2023). Immunogenicity and efficacy of vaccine boosters
against SARS-CoV-2 Omicron subvariant BA.5 in male Syrian hamsters. Nat Commun *14*, 4260.
10.1038/s41467-023-40033-2.

Macklin, G.R., O'Reilly, K.M., Grassly, N.C., Edmunds, W.J., Mach, O., Santhana Gopala Krishnan, R.,
Voorman, A., Vertefeuille, J.F., Abdelwahab, J., Gumede, N., et al. (2020). Evolving epidemiology of
poliovirus serotype 2 following withdrawal of the serotype 2 oral poliovirus vaccine. Science *368*,
401-405. 10.1126/science.aba1238.
Marrack, P., McKee, A.S., and Munks, M.W. (2009). Towards an understanding of the adjuvant action of
aluminium. Nat Rev Immunol *9*, 287-293. 10.1038/nri2510.
Natekar, J.P., Pathak, H., Stone, S., Kumari, P., Sharma, S., Auroni, T.T., Arora, K., Rothan, H.A.,
and Kumar, M. (2022). Differential Pathogenesis of SARS-CoV-2 Variants of Concern in Human
ACE2-Expressing Mice. Viruses *14*. 10.3390/v14061139.
Nouailles, G., Adler, J.M., Pennitz, P., Peidli, S., Teixeira Alves, L.G., Baumgardt, M., Bushe, J.,
Voss, A., Langenhagen, A., Langner, C., et al. (2023). Live-attenuated vaccine sCPD9 elicits superior
mucosal and systemic immunity to SARS-CoV-2 variants in hamsters. Nat Microbiol *8*, 860-874.
10.1038/s41564-023-01352-8.
Okabe, M., Ikawa, M., Kominami, K., Nakanishi, T., and Nishimune, Y. (1997). 'Green mice' as a source
of ubiquitous green cells. FEBS Lett *407*, 313-319. 10.1016/s0014-5793(97)00313-x.
Okamura, S., Yoshida, A., Miyazato, P., Matsumoto, M., and Ebina, H. (2023). Protocol to isolate
temperature-sensitive SARS-CoV-2 mutants and identify associated mutations. STAR Protoc *4*, 102352.
10.1016/j.xpro.2023.102352.
Peacock, T.P., Goldhill, D.H., Zhou, J., Baillon, L., Frise, R., Swann, O.C., Kugathasan, R., Penn,
R., Brown, J.C., Sanchez-David, R.Y., et al. (2021). The furin cleavage site in the SARS-CoV-2 spike
protein is required for transmission in ferrets. Nat Microbiol *6*, 899-909. 10.1038/s41564-021-00908-w.
Ploquin, A., Szecsi, J., Mathieu, C., Guillaume, V., Barateau, V., Ong, K.C., Wong, K.T., Cosset, F.L.,
Horvat, B., and Salvetti, A. (2013). Protection against henipavirus infection by use of recombinant
adeno-associated virus-vector vaccines. J Infect Dis *207*, 469-478. 10.1093/infdis/jis699.
Polack, F.P., Thomas, S.J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L., Perez
Marc, G., Moreira, E.D., Zerbini, C., et al. (2020). Safety and Efficacy of the BNT162b2 mRNA Covid-19
Vaccine. N Engl J Med *383*, 2603-2615. 10.1056/NEJMoa2034577.
Primard, C., Monchatre-Leroy, E., Del Campo, J., Valsesia, S., Nikly, E., Chevandier, M., Boue, F.,
Servat, A., Wasniewski, M., Picard-Meyer, E., et al. (2023). OVX033, a nucleocapsid-based vaccine
candidate, provides broad-spectrum protection against SARS-CoV-2 variants in a hamster challenge model.
Front Immunol *14*, 1188605. 10.3389/fimmu.2023.1188605.
Sabin, A.B. (1985). Oral poliovirus vaccine: history of its development and use and current challenge
to eliminate poliomyelitis from the world. J Infect Dis *151*, 420-436. 10.1093/infdis/151.3.420.
Sasaki, M., Toba, S., Itakura, Y., Chambaro, H.M., Kishimoto, M., Tabata, K., Intaruck, K., Uemura,
K., Sanaki, T., Sato, A., et al. (2021). SARS-CoV-2 Bearing a Mutation at the S1/S2 Cleavage Site Exhibits
Attenuated Virulence and Confers Protective Immunity. mBio *12*, e0141521. 10.1128/mBio.01415-21.
Seo, S.H., and Jang, Y. (2020). Cold-Adapted Live Attenuated SARS-Cov-2 Vaccine Completely Protects
Human ACE2 Transgenic Mice from SARS-Cov-2 Infection. Vaccines (Basel) *8*. 10.3390/vaccines8040584.
Thwaites, R.S., Uruchurtu, A.S.S., Negri, V.A., Cole, M.E., Singh, N., Poshai, N., Jackson, D., Hoschler,
K., Baker, T., Scott, I.C., et al. (2023). Early mucosal events promote distinct mucosal and systemic
antibody responses to live attenuated influenza vaccine. Nat Commun *14*, 8053.
10.1038/s41467-023-43842-7.
Torii, S., Ono, C., Suzuki, R., Morioka, Y., Anzai, I., Fauzyah, Y., Maeda, Y., Kamitani, W., Fukuhara,
T., and Matsuura, Y. (2021). Establishment of a reverse genetics system for SARS-CoV-2 using circular
polymerase extension reaction. Cell Rep *35*, 109014. 10.1016/j.celrep.2021.109014.
Wang, P., Lau, S.Y., Deng, S., Chen, P., Mok, B.W., Zhang, A.J., Lee, A.C., Chan, K.H., Tam, R.C.,
Xu, H., et al. (2021a). Characterization of an attenuated SARS-CoV-2 variant with a deletion at the
S1/S2 junction of the spike protein. Nat Commun *12*, 2790. 10.1038/s41467-021-23166-0.
Wang, Y., Yang, C., Song, Y., Coleman, J.R., Stawowczyk, M., Tafrova, J., Tasker, S., Boltz, D., Baker,
R., Garcia, L., et al. (2021b). Scalable live-attenuated SARS-CoV-2 vaccine candidate demonstrates
preclinical safety and efficacy. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2102775118.
Weinheimer, V.K., Becher, A., Tonnies, M., Holland, G., Knepper, J., Bauer, T.T., Schneider, P.,
Neudecker, J., Ruckert, J.C., Szymanski, K., et al. (2012). Influenza A viruses target type II

pneumocytes in the human lung. J Infect Dis *206*, 1685-1694. 10.1093/infdis/jis455.
Yasmin, F., Najeeb, H., Naeem, U., Moeed, A., Atif, A.R., Asghar, M.S., Nimri, N., Saleem, M.,
Bandyopadhyay, D., Krittanawong, C., et al. (2023). Adverse events following COVID-19 mRNA vaccines:
A systematic review of cardiovascular complication, thrombosis, and thrombocytopenia. Immun Inflamm
Dis *11*, e807. 10.1002/iid3.807.
Yeh, M.T., Bujaki, E., Dolan, P.T., Smith, M., Wahid, R., Konz, J., Weiner, A.J., Bandyopadhyay, A.S.,
Van Damme, P., De Coster, I., et al. (2020). Engineering the Live-Attenuated Polio Vaccine to Prevent
Reversion to Virulence. Cell Host Microbe *27*, 736-751 e738. 10.1016/j.chom.2020.04.003.
Yeh, M.T., Smith, M., Carlyle, S., Konopka-Anstadt, J.L., Burns, C.C., Konz, J., Andino, R., and Macadam,
A. (2023). Genetic stabilization of attenuated oral vaccines against poliovirus types 1 and 3. Nature
619, 135-142. 10.1038/s41586-023-06212-3.
Yoshida, A., Okamura, S., Torii, S., Komatsu, S., Miyazato, P., Sasaki, H., Ueno, S., Suzuki, H.,
Kamitani, W., Ono, C., et al. (2022). Versatile live-attenuated SARS-CoV-2 vaccine platform applicable
to variants induces protective immunity. iScience *25*, 105412. 10.1016/j.isci.2022.105412.
Young, B.E., Fong, S.W., Chan, Y.H., Mak, T.M., Ang, L.W., Anderson, D.E., Lee, C.Y., Amrun, S.N.,
Lee, B., Goh, Y.S., et al. (2020). Effects of a major deletion in the SARS-CoV-2 genome on the severity
of infection and the inflammatory response: an observational cohort study. Lancet *396*, 603-611.
10.1016/S0140-6736(20)31757-8.
Zhang, G.F., Meng, W., Chen, L., Ding, L., Feng, J., Perez, J., Ali, A., Sun, S., Liu, Z., Huang, Y.,
et al. (2022). Neutralizing antibodies to SARS-CoV-2 variants of concern including Delta and Omicron
in subjects receiving mRNA-1273, BNT162b2, and Ad26.COV2.S vaccines. J Med Virol *94*, 5678-5690.
10.1002/jmv.28032.
Zinzula, L. (2021). Lost in deletion: The enigmatic ORF8 protein of SARS-CoV-2. Biochem Biophys Res
Commun *538*, 116-124. 10.1016/j.bbrc.2020.10.045.
747

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Table 1. SARS-CoV-2 strains

REAGENT or RESOURCE	SOURCE	IDENTIFIER
SARS-CoV-2: pre-alpha type, D614G, B-1 strain	Yoshida. et al.,2022	NCBI: LC603286
A50-18 strain	Yoshida. et al.,2022	NCBI: LC603287
L50-33 strain	Yoshida. et al.,2022	NCBI: LC603289
B-1 ΔFCS strain	Current study	N/A[a]
Candidate 1 (BK2102)	Current study	N/A[a]
Candidate 2	Current study	N/A[a]
Candidate 3	Current study	N/A[a]
SARS-CoV-2: delta variant, BK325 strain	Research Foundation for Microbial Diseases of Osaka University	N/A[a]
SARS-CoV-2: omicron variant, TY41-702 strain	National Institute of Infectious Diseases	GISAID ID: EPI_ISL_13241867

[a] Not applicable

Table 2. Primer list

Primer ID	Oligonucleotides	SOURCE
F1_F	CTATATAAGCAGAGCTCGTTTAGTGAACCGTattaaaggtttataccttcccaggtaac	Torii, et al. (2021)
F1_R	cagattcaacttgcatggcattgttagtagccttatttaaggctcctgc	Torii, et al. (2021)
F2_F	gcaggagccttaaataaggctactaacaatgccatgcaagttgaatctg	Torii, et al. (2021)
F2_R	ggtaggattttccactacttcttcagagactggttttagatcttcgcaggc	Torii, et al. (2021)
F3_F	gcctgcgaagatctaaaaccagtctctgaagaagtagtggaaaatcctacc	Torii, et al. (2021)
F3_R	ggtgcacagcgcagcttcttcaaaagtactaaagg	Torii, et al. (2021)
F4_F	caccactaattcaacctattggtgctttggacatatcagcatctatagtagctggtgg	Torii, et al. (2021)
F4_R	gtttaaaaacgattgtgcatcagctgactg	Torii, et al. (2021)
F5_F	cacagtctgtaccgtctgcggtatgtggaaaggttatggctgtagttgtgatc	Torii, et al. (2021)
F5_R	gcggtgtgtacatagcctcataaaactcaggttcccaataccttgaagtg	Torii, et al. (2021)
F6_F	cacttcaaggtattgggaacctgagttttatgaggctatgtacacaccgc	Torii, et al. (2021)
F6_R	catacaaactgccaccatcacaaccaggcaagttaaggttagatagcactctag	Torii, et al. (2021)
F7_F	ctagagtgctatctaaccttaacttgcctggttgtgatggtggcagtttgtatg	Torii, et al. (2021)
F7_R	ctagagactagtggcaataaaacaagaaaaacaaacattgttcgtttagttgttaac	Torii, et al. (2021)
F8_F	gttaacaactaaacgaacaatgtttgtttttcttgttttattgccactagtctctag	Torii, et al. (2021)
F8_R	gcagcaggatccacaagaacaacagcccttgagacaactacagcaactgg	Torii, et al. (2021)
F9_F	ccagttgctgtagttgtctcaagggctgttgttcttgtggatcctgctgc	Torii, et al. (2021)
F9_R	caatctccattggttgctcttcatc	Torii, et al. (2021)
F10_F	gatgaagagcaaccaatggagattg	Torii, et al. (2021)
F10_R	GGAGATGCCATGCCGACCCtttttttttttttttttttttttttttgtcattctcctaag	Torii, et al. (2021)
Linker_F	cttaggagaatgacaaaaaaaaaaaaaaaaaaaaaaaaaaaGGGTCGGCATGGCATCTCC	Torii, et al. (2021)
Linker_R	gttacctgggaaggtataaacctttaatACGGTTCACTAAACGAGCTCTGCTTATATAG	Torii, et al. (2021)
TS_F6_R	catacaaactgccacTatcacaaccaggcaagttaaggttagatagcactctag	Yoshida, et al. (2022)
TS_F7_F	ctagagtgctatctaaccttaacttgcctggttgtgatAgtggcagtttgtatg	Yoshida, et al. (2022)

Table 3. Genetic variations of viruses passaged *in vivo*

	Mutations					
	ΔG359 - A382 (*NSP1*)	G18782T (*NSP14*)	G19285A (*NSP14*)	C19550T (*NSP14*)	ΔA23598-G23624 (*Spike* FCS)	ΔC27549-T28251 (*ORF7a-8*)
A50-18	N/A[a]	G	G/A	T	N/A[a]	N/A[a]
	N/A[a]	G/T	G/A	T	N/A[a]	N/A[a]
	N/A[a]	G/T	G/A	T	N/A[a]	N/A[a]
	N/A[a]	G/T	G	T	N/A[a]	N/A[a]
	N/A[a]	G/T	G/A	T	N/A[a]	N/A[a]
BK2102	○[b]	T	A	T	○[b]	○[b]
	○[b]	T	A	T	○[b]	○[b]
	○[b]	T	A	T	○[b]	○[b]
	○[b]	T	A	T	○[b]	○[b]
	○[b]	T	A	T	○[b]	○[b]

[a] Not applicable
[b] Same sequence as the inoculated virus

Table 4. NSP3 genetic variations in viruses recovered from infected Tg mice

| | n | Virus titer (Log10 PFU/g) | | TS substitution |
		Brain	Lung	NSP3
D614G (10^2 **PFU infected**)	5	6.94 ± 0.44[a]	3.67 ± 0.26[a]	N/A[b]
B-1 ΔFCS (10^2 **PFU infected**)	5	8.69 ± 0.33[a]	5.64 ± 0.62[a]	N/A[b]
L50-33 (10^4 **PFU infected**)	1	5.75	< 2.40	F445L (TTT → TTG)
L50-33 (10^5 **PFU infected**)	1	7.24	2.40	F445L (TTT → TTG)

[a] Average ± SD
[b] Not applicable