1

Immunomagneto-activated hepatocyte-specific tissue vesicle profiling

reveals message diversification from regenerating liver

Si-Qi Ying[1,#], Yuan Cao[1,2,#], Lu Liu[1,2,#], Ze-Kai Zhou[1,3,#], Xin-Yan Luo[1,3], Cheng-Han Li[1,3],

Shu-Yan Chen[1,3], Xiao Lei[1,2], Shu-Juan Xing[1,4], Ke Shi[1,5], Ji-Yu Qiu[6], Yuan-Yuan Li[1],

Chen-Xi Zheng[1], Fang Jin[1,2], Yan Jin[1,6,*], Bing-Dong Sui[1,*]

[1] Research and Development Center for Tissue Engineering, The Fourth Military Medical University, Xi'an, Shaanxi 710032, China.

[2] Department of Orthodontics, School of Stomatology, The Fourth Military Medical University, Xi'an, Shaanxi 710032, China.

[3] School of Basic Medicine, The Fourth Military Medical University, Xi'an, Shaanxi 710032, China.

[4] College of Life Science, Northwest University, Xi'an, Shaanxi 710069, China.

[5] Xi'an Institute of Tissue Engineering and Regenerative Medicine, Xi'an, Shaanxi 710032, China.

[6] Department of VIP Dental Care, School of Stomatology, The Fourth Military Medical University, Xi'an, Shaanxi 710032, China.

[#] These authors contributed equally to this study.

* Corresponding authors

Prof. Bing-Dong Sui. Research and Development Center for Tissue Engineering, The

Fourth Military Medical University, 145 West Changle Road, Xi'an, Shaanxi 710032, China.

E-mail: bingdong@fmmu.edu.cn.

Prof. Yan Jin. Research and Development Center for Tissue Engineering, The Fourth

Military Medical University, 145 West Changle Road, Xi'an, Shaanxi 710032, China. E-mail:

yanjin@fmmu.edu.cn.

Abstract

Background & Aims: The liver possesses a remarkable regenerative capacity which involves intricate intercellular communication. However, the mechanisms underlying the complex intercellular interactions remain incompletely understood. Extracellular vesicles (EVs) are emerging as important messengers in diverse pathophysiological conditions. Nevertheless, tissue-derived, cell-specific functional EV populations in regeneration have not been robustly analyzed.

Methods: Single-cell RNA sequencing (scRNA-seq) analysis of the regenerating liver after partial hepatectomy (PHx), ultrastructural examinations of *in situ* liver tissue extracellular vesicles (LT-EVs), and nanoscale and proteomic profiling of hepatocyte-specific EVs were integrated. Especially, a feasible approach for investigating antigen-specific endogenous EVs was developed after hepatic enzymatic digestion followed by differential centrifugation and immunomagnetic sorting of LT-EVs.

Results: scRNA-seq analysis revealed hepatocytes as the critical source of intercellular communication in the regenerating liver. In particular, hepatocytes are programmed to be enriched for biogenesis and release of EVs in liver regeneration. By analyzing *in situ* EVs using transmission electron microscope (TEM), LT-EVs were discovered to be emergingly present in the extracellular interstitial space adjacent to hepatocytes in the regenerating liver. Moreover, asialoglycoprotein receptor (ASGPR)-marked, magnetic microbead-sorted LT-EVs (named ASGPR$^+$mLT-EVs) were collected. These hepatocyte-specific LT-EVs were further demonstrated with strengthened release and distinct protein signatures during liver regeneration. Importantly, the EVs demonstrated parental hepatocyte-derived information

diversification, which not only inherited regulated proliferative and metabolic signals at the

cellular level but also showed vesicular enrichment of organelle components.

Conclusions: These findings provide the first knowledge that hepatocyte-specific tissue

vesicles are indeed phenotypically and functionally involved in liver regeneration. Our study

paves an avenue for in-depth biological and mechanistic research of regenerative LT-EVs

and sheds light on extensive dissection of antigen-marked, physiological, and endogenous

tissue EV populations.

Key words

Liver regeneration; hepatocytes; extracellular vesicles; single-cell RNA sequencing;

immunomagnetic sorting; proteomic analysis

1. Introduction

The liver is the largest solid organ in the body which critically maintains the metabolism and serves as the central hub for detoxification. In order to cope with daily challenges, the liver has developed a potent, robust, and fine-tuned regenerative capability, which can fully regenerate up-to-2/3 loss of the total parenchyma [1, 2]. The process of liver regeneration involves a complex cellular interplay primarily centered on hepatocytes but also including biliary cells and various non-parenchymal cells (NPCs) [3]. Particularly, in the healthy liver, hepatocytes are mitotically quiescent; whereas following toxic damage or surgical resection, hepatocytes can rapidly enter the cell cycle to proliferate or transdifferentiate from biliary epithelial cells (BECs) with orchestrating immune, endothelial, and stromal components to synergistically restore liver mass and function [4-8]. This hepatocyte expansion-correlated cellular coordination is acknowledged to be mediated by diverse paracrine cytokines, which safeguard efficient liver regeneration as a prerequisite for proper body functioning [4, 5, 9]. However, the liver regeneration capacity can be overwhelmed or suppressed under large injury or progressive damages, resulting in hepatic diseases remaining as a global medical challenge [10, 11]. Deciphering in-depth the mechanisms underlying liver regeneration will help to devise feasible strategies for hepatic disease therapy.

Extracellular vesicles (EVs) represent an emerging and alternative form of intercellular communication regulating a variety of biological processes, which are functionalized with surface molecules and encapsulated cargoes, such as proteins, nucleic acids, and lipids [12, 13]. Particularly, solid tissues are increasingly recognized as natural reservoirs of endogenous EVs, and tissue-derived EVs (Ti-EVs) play a putative role in reflecting and

regulating the tissue health and disease statuses [14]. A recent study has isolated Ti-EVs

from the kidney and the skin, which have been used to promote target tissue repair in a

donor tissue-specific manner after allogeneic transplantation [15]. It has also been reported

that liver tissue EVs (denoted LT-EVs) isolated from both normal and carbon tetrachloride

(CCl_4)-induced damaged liver accelerate the recovery of liver tissues from CCl_4-induced

hepatic necrosis after administration [16]. However, owing to the heterogeneous nature of

tissue resident cells and the imprecision property of the bulk isolation method of Ti-EVs, it

remains an unmet need to clarify and characterize the detailed endogenous EV populations

that contribute to tissue physiology and pathology [17-19]. With regard to liver regeneration,

small EVs derived from the cultured hepatocyte supernatant have been shown to promote

liver regeneration after infusion [20]. Nevertheless, whether LT-EVs from hepatocytes or

any other liver cell types specifically participate in the liver regeneration process has not

been evaluated.

In this study, we aimed to investigate *in situ* tissue-derived, cell-specific functional EV

population(s) in liver regeneration. By performing an unbiased examination of the hepatic

cell landscape *via* single-cell RNA sequencing (scRNA-seq), we revealed that hepatocytes

serve as the key source of intercellular communication in the regenerating liver after partial

hepatectomy (PHx), which are programmed to be enriched for biogenesis and release of

EVs. Furthermore, we first conducted the ultrastructural observation of LT-EVs *in situ* and

established an immunomagnetic sorting protocol analyzing the hepatocyte-specific LT-EVs

marked by the featured asialoglycoprotein receptor (ASGPR). These ASGPR$^+$ magnetic

microbead-sorted LT-EVs (hereby denoted mLT-EVs) were thoroughly characterized for the

physicochemical properties of morphology, size, particle content, and protein composition

at the nanoscale or high throughput level, and the transcriptome of hepatocytes and the

proteome of ASGPR$^+$mLT-EVs were integrated to understand their contributions to liver

regeneration. We discovered that hepatocytes release a specific LT-EV population which

transmits diversified messages and hallmarks liver regeneration. Our study thus paves an

avenue for in-depth biological and mechanistic research of regenerative LT-EVs and sheds

light on extensive investigation of antigen-marked, physiological, and endogenous tissue

EV populations.

2. Results

2.1 scRNA-seq analysis reveals hepatocytes as the key source of intercellular

communication in the regenerating liver

To begin, we select the PHx model which enables the study of liver regeneration in a

setting without enormous necrosis and associated inflammation [2]. A 2/3 PHx surgery was

performed in mice with laparotomy only as the Sham control [21]. Liver regeneration with

hepatocyte proliferation was confirmed at 72 h after PHx by histological examination and

immunofluorescence (IF) staining of phospho-histone H3 (PHH3), a mitosis marker [23]

(Figure S1A and B). We then collected and reanalyzed two scRNA-seq data of the mouse

liver respectively from Sham and PHx conditions (GSM4572241; GSM4572244). Cell

clustering using principal component analysis and t-distributed stochastic neighbor

embedding (tSNE) plot visualization identified 8 distinct liver cell types, in which the

proportion of hepatocytes decreased after PHx (Figure 1A and B). Intriguingly, despite

decreased proportion, enrichment of intercellular communication originated from hepatocytes was revealed prominently after PHx by the Cellchat analysis (Figure 1C). Moreover, the number of differentially expressed genes (DEGs) in hepatocytes over other liver cell types was high (Figure 1D). These DEGs were further demonstrated by the Gene Ontology (GO) analysis to be associated with multiple terms, notably the "cellular anatomical entity" in the "Cellular Component" domain and the "cellular process" and "metabolic process" in the "Biological Process" domain (Figure 1E). Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis also exhibited that hepatocyte DEGs were particularly associated with "Transport and catabolism" and "cell growth and death" in the "Cellular Processes" domain and were linked with the "Metabolism" domain (Figure 1F). These data were in accordance with hepatocyte turnover behaviors and metabolic function in liver homeostasis and regeneration, which additionally indicated the cellular transport-based information transmission as a characteristic of hepatocytes. Taken together, these results suggest that hepatocytes serve as the key source of intercellular communication in the regenerating liver.

Figure 1

Figure 1. scRNA-seq analysis reveals hepatocytes as the key source of intercellular

communication in the regenerating liver. **(A)** tSNE plots displaying the distribution of

different cell types in Sham and PHx livers. **(B)** Cell type proportion in Sham and PHx livers.

(C) Bubble maps showing intercellular interactions among Sham and PHx liver cells. **(D)**

The number of upregulated DEGs in respective liver cell clusters. **(E)** GO analysis of

upregulated DEGs in hepatocytes. **(F)** KEGG analysis of upregulated DEGs in hepatocytes.

Source data were derived from the GSM4572241 and GSM4572243 series in the GEO

database.

2.2 Hepatocytes are programmed to be enriched for biogenesis and release of EVs in

liver regeneration

Next, we investigated the detailed gene expression profiles of hepatocytes in Sham

and PHx livers using the scRNA-seq data. In the pseudotime analysis, hepatocytes were

ordered into seven stages (Figure 2A). Particularly, hepatocytes from Sham and PHx livers

fell into distinct stage branches with a few transitioning cells, indicating a cellular state shift

(Figure 2B). Accordingly, 3524 DEGs between Sham and PHx hepatocytes were identified,

with respectively 1232 upregulated DEGs and 2292 downregulated DEGs in PHx

hepatocytes compared to their Sham counterparts (Figure 2C). These DEGs, based on the

functional annotations of the GO database, were associated with "cellular anatomical entity"

in the "Cellular Component" domain and the "cellular process" and "metabolic process" in

the "Biological Process" domain, which were consistent with the overall analysis of

hepatocyte DEGs compared to other liver cells (Figure 2D). Notably, the PHx over Sham

hepatocyte DEGs were associated with the "biological regulation", "regulation of biological

process" and "response to stimulus" terms in the "Biological Process" domain, indicating

functional relationship with liver regeneration after injury (Figure 2D). Also, these DEGs

corresponded with the overall analysis of hepatocyte DEGs in the KEGG analysis, which

were associated with the "Transport and catabolism" and "cell growth and death" in the

"Cellular Processes" domain and were linked with the "Metabolism" domain (Figure 2E).

Rapid entry of hepatocytes into the cell cycle with promotion of mitosis was further

confirmed by the Cellcycle analysis (Figure S1C). To examine in-depth functional

implications of the DEGs, enrichment analyses were performed based on GO and KEGG

information, which showed the organelle-related terms, such as "mitochondrion" and

"ribosome", and detailed metabolic terms, were among the top enriched terms in PHx

hepatocytes (Figure 2F and G). Particularly, considering above findings on the hepatocyte

role of intercellular communication in liver regeneration, we focused on EV-related terms in

the PHx over Sham hepatocyte DEGs, and the results demonstrated remarkable

enrichment of potential EV function in PHx hepatocytes, including EV assembly and

secretion (Figure 2H). Taken together, these results suggest that hepatocytes are

programmed to be enriched for biogenesis and release of EVs in liver regeneration.

Figure 2

Figure 2. Hepatocytes are programmed to be enriched for biogenesis and release of

EVs in liver regeneration. (A and B) Pseudotime analysis plots indicating cellular

trajectories of hepatocytes in PHx and Sham livers. **(C)** Volcano plot showing DEGs

between PHx and Sham hepatocytes. **(D)** GO analysis of DEGs between PHx and Sham

hepatocytes. **(E)** KEGG analysis of DEGs between PHx and Sham hepatocytes. **(F)** GO

enrichment analysis of DEGs between PHx and Sham hepatocytes shown by the circle

diagram. **(G)** KEGG enrichment analysis of DEGs between PHx and Sham hepatocytes

shown by the circle diagram. **(H)** EV-related terms are enriched in PHx hepatocytes over

Sham hepatocytes. Source data were derived from the GSM4572241 and GSM4572243

series in the GEO database.

2.3 LT-EVs are emergingly present in the extracellular interstitial space adjacent to hepatocytes in the regenerating liver

Next, we intended to investigate that whether EVs are indeed involved in the liver

regeneration process. Ti-EVs putatively exist in the tissue extracellular space and reflect

pathophysiological features of the tissue [14, 15]. However, current studies examine Ti-EVs

mainly *via* enzymatic digestion and isolation, resulting in loss of spatiotemporal information

and potential compromise of EV amount and integrity [17-19]. Existing literatures are also

limited on the evaluation of *in situ* EVs in tissue regeneration. Therefore, to detect the *bona*

fide EVs in the liver, we conducted transmission electron microscopy (TEM) analysis

(Figure 3). Notably, in tissue sections of PHx, compared to those of Sham, hepatocytes

were enlarged and the interstitial space was narrowed, in accordance with the hypertrophic

phenotype in regeneration (Figure 3) [24]. Particularly, in the extracellular space adjacent to

hepatocytes, vesicle-like nanosized particles was detected, which were presence in the

Sham liver and demonstrated a dramatic increase in abundance in the PHx liver (Figure 3).

These findings first indicate the active involvement of *in situ* EVs in liver regeneration.

Figure 3

Liver tissue TEM

Figure 3. LT-EVs are emergingly present in the extracellular interstitial space adjacent to hepatocytes in the regenerating liver. Sham and PHx liver tissues were prepared and analyzed by TEM with showing representative images. Hepatocytes were identified with featured morphology. White arrows indicating EVs in the extracellular interstitial space. Bars: 1 μm (left) and 200 nm (right).

2.4 Protocol establishment for collecting and characterizing hepatocyte-specific LT-EVs

The above results inspired us to specifically investigate hepatocyte-derived EVs in liver

regeneration. To fulfill this aim, we established a protocol for collecting and characterizing

antigen-specific Ti-EVs by integrating our previous methods of immunomagneto-activated

sorting of the marked EV population from cultured cells and isolation of traceable EVs from

the tissue (Figure 4) [22, 25]. Particularly, ASGPR is primarily expressed on the surface of

hepatocytes [26], which was utilized for sorting and collecting hepatocyte-specific LT-EVs

(Figure 4). Briefly, liver tissues were disassociated, digested with Liberase, and subjected

to differential centrifugation to pellet LT-EVs (Figure 4, left). LT-EVs were then incubated for

the PE-conjugated ASGPR antibody followed by magnetic microbead incubation, which

were subsequently sorted via the LD columns and collected by centrifugation (Figure 4,

right). The hepatocyte-specific LT-EVs were accordingly denoted ASGPR$^+$ magnetic

microbead-sorted LT-EVs (mLT-EVs), which can be characterized by TEM for the

morphology, nanoparticle tracking analysis (NTA) for the size distribution and number, flow

cytometric and Western blot assays for the surface and cargo profiles, and the mass

spectrometry for multi-omic analyses (Figure 4, bottom). As expected, flow cytometric

experiment data confirmed that the ASGPR$^+$ percentage of LT-EVs was 33.79% before

magnetic sorting, while the percentage decreased to 0.10% in ASGPR$^-$LT-EVs and

increased to 98.34% in ASGPR$^+$mLT-EVs, indicating the successful application of this

approach (Figure 5A). The negative controls and gating strategy were additionally shown

(Figure S2A and B). Together, we suggest that this approach will help assess the properties

of hepatocyte-specific tissue EVs to enhance our understanding of liver regeneration.

Figure 4

Figure 4. Protocol establishment for collecting and characterizing hepatocyte-specific LT-EVs. Schematic overview of the protocol employing enzymatic digestion, gradient centrifugation, immunomagnetic separation, and multiscale analysis for the isolation and characterization of hepatocyte-specific EVs (hereby denoted ASGPR⁺mLT-EVs) from liver tissues.

2.5 Hepatocytes strengthen EV release in liver regeneration

After validation of the efficacy of immunomagnetic isolation, we adopted this approach to investigate LT-EVs and mLT-EVs derived from Sham and PHx mice, with the quality of isolated EVs evaluated according to the Minimal information for studies of extracellular vesicles 2018 (MISEV2018) [27]. TEM observation showed that LT-EVs exhibited typical cup-shape morphology with membranous structure in Sham and PHx (Figure 5B). Moreover, both LT-EVs and mLT-EVs from Sham and PHx mice expressed representative EV markers, such as the tetraspanin surface molecule CD63, the endosomal sorting complex required for transport (ESCRT)-associated protein tumor suppressor gene 101 (TSG101), and the lipid raft protein Flotillin-1 (Figure 5C). Meanwhile, the Golgi component protein Golgin84, a negative EV marker, was not detected in either LT-EVs or mLT-EVs (Figure 5C). Further measurement by NTA demonstrated that the diameters of LT-EVs from both Sham and PHx mice ranged from 50 to 500 nm and peaked at 150-200 nm (Figure 5D). Notably, compared to Sham mice, PHx mice produced significantly more amount of LT-EVs, as quantified by calculating the particle number by NTA (Fig. 5E) and the protein content by BCA assay (Figure 5F). After immunomagnetic sorting, ASGPR$^+$mLT-EVs from both Sham and PHx mice still showed characteristic particle distribution in NTA (Figure 5G), and PHx ASGPR$^+$mLT-EVs demonstrated higher yield than Sham ASGPR$^+$mLT-EVs (Figure 5H and I). Together, our results suggest that hepatocytes strengthen EV release in liver regeneration.

Figure 5

Figure 5. Hepatocytes strengthen EV release in liver regeneration. (A) Flow cytometric

analysis showing ASGPR surface expression on LT-EVs before and after immunomagnetic

sorting. **(B)** Representative TEM image of collected LT-EVs from Sham and PHx livers.

Bars, 250 nm. **(C)** Western blot analysis of protein marker expression of collected LT-EVs

and ASGPR⁺mLT-EVs from Sham and PHx livers. **(D and E)** Size distribution and particle

number quantification of LT-EVs determined by NTA. **(F)** Quantification of protein amount of

LT-EVs. **(G and H)** Size distribution and particle number quantification of ASGPR⁺mLT-EVs

determined by NTA. **(I)** Quantification of protein amount of ASGPR⁺mLT-EVs. n = 3 per

group. **, $p < 0.01$; ***, $p < 0.001$; two-tailed Student's unpaired t tests.

2.6 Proteomic analysis identifies liver regeneration-associated cargo profiles of

hepatocyte-specific LT-EVs

Next, we investigated the protein cargo profiles of hepatocyte-specific LT-EVs from

Sham and PHx by performing the liquid chromatography with tandem mass spectrometry

(LC-MS/MS) analysis (Table S1). A total of over 3000 proteins were identified (Table S2).

The Venn diagram illustrated the presence of 3357 shared proteins between Sham and

PHx mLT-EVs, with respective 56 and 129 distinct proteins in each group (Figure 6A).

Among the shared proteins, 387 proteins were identified as the differentially expressed

proteins (DEPs; fold change > 1.5 and p value < 0.05), and 193 DEPs were upregulated in

PHx ASGPR⁺mLT-EVs with 184 DEPs downregulated (Figure 6B). Heatmap also showed

clear separation and distinct protein signatures between Sham and PHx ASGPR⁺mLT-EVs

(Figure 6C). The top 30 DEPs of PHx over Sham ASGPR⁺mLT-EVs was exhibited in the

radar diagram (Figure 6D). With respect to their subcellular localization, most of the DEPs

existed in the cytoplasm region (32.1%) or located on the plasma membrane (20.42%),

indicating transferred cargoes or potential surface markers (Figure 6E). GO analysis of

DEPs further demonstrated that annotation related to metabolic and biosynthetic processes,

as well as binding and transporter activities, were particular regulated terms in hepatocyte

EVs during liver regeneration (Figure 6F). With regard to the functionality of DEPs, based

on the KEGG database categories, DEPs were notably associated with "Transport and

catabolism" and "Cell growth and death" in the "Cellular Processes" domain, as well as the

"Environmental Information Processing" and the "Metabolism" domains (Figure 6G).

Together, these findings identify liver regeneration-associated cargo profiles of

hepatocyte-specific LT-EVs.

Figure 6

Figure 6. Proteomic analysis identifies liver regeneration-associated cargo profiles

of hepatocyte-specific LT-EVs. (A) Venn analysis of mLT-EVs from PHx and Sham livers.

(B) Volcano plot of DEPs between PHx and Sham mLT-EVs. **(C)** Hierarchical clustering of

DEPs between PHx and Sham mLT-EVs. **(D)** Radar diagram of DEPs between PHx and

Sham mLT-EVs. **(E)** Distribution of DEPs in subcellular location categories retrieved from

UniProt database according to GO annotation. **(F)** GO analysis of DEPs between PHx and

Sham mLT-EVs. **(G)** KEGG analysis of DEPs between PHx and Sham mLT-EVs. n = 3 per

group.

2.7 EVs diversify hepatocyte-derived information in liver regeneration

Next, we kept focusing on these DEPs and performed detailed enrichment analysis.

GO enrichment analysis revealed that DEPs of PHx over Sham ASGPR$^+$mLT-EVs were

particularly related to metabolic processes, which, however, were largely downregulated

(Figure 7A). To further elucidate the functional disparities of DEPs, we classified the DEPs

into four segments: Q1 (upregulated more than two folds), Q2 (upregulated less than two

folds), Q3 (downregulated less than two folds), and Q4 (downregulated more than two

folds). Each segment was then subjected to GO enrichment analysis, followed by functional

cluster analysis. Notably, the "Biological Process" enrichment-based clustering analysis

revealed that PHx mLT-EVs exhibited remarkable upregulated DEPs in the Q1 segment

related to hepatocyte proliferation, including "telomere maintenance via recombination" and

"mitotic recombination" (Figure 7B). On the contrary, downregulated DEPs in the Q4

segment were related to the metabolic processes, such as "gluconeogenesis" (Figure 7B).

The "Cellular Component" enrichment-based clustering analysis also showed

proliferation-related DEP upregulation in the Q1 segment, such as "chromosomes", with

organelle-related terms, including "endoplasmic reticulum" (Figure 7C). The

enrichment-based clustering analysis in "Molecular Function" subsequently identified

specific upregulated DEPs in the Q1 segment linked to "epidermal growth factor receptor

binding", while demonstrating a significant DEP downregulation in "antioxidant activity"

(Figure 7D). KEGG enrichment analysis of DEPs with functional clustering confirmed the

trend of metabolic repression in PHx ASGPR$^+$mLT-EVs (Figure 7E and F). In the Q1

segment of KEGG enrichment-based DEP clustering analysis, regenerative

proliferation-related terms were further notable, such as "DNA replication" and "Cell cycle"

(Figure 7F). The correlations among the DEPs were discovered to be substantial (Figure

S3A).

We intended to take a final step to dissect the potential relationship between proteomic

alterations of hepatocyte-specific LT-EVs with transcriptomic changes of their parental cells.

By integrating the scRNA-seq data and the proteomic data, we performed a nine-quadrant

analysis to distinguish cellular and vesicular expression patterns under liver regeneration.

The graph depicted the same expression patterns in particular interested quadrants of P3

(upregulation in both cases) and P7 (downregulation in both cases), meanwhile displaying

different expression patterns in other quadrants, such as P6 (upregulation only in proteomic

analysis) and P9 (upregulation in proteins despite downregulation in mRNAs) (Figure 7G).

For P3, GO enrichment analysis revealed association of terms with cellular activity,

indicating that EVs inherited active regulation from parental hepatocytes in liver

regeneration (Figure 7H). It is especially notable that the term "epidermal growth factor

receptor binding" term was enriched in P3, suggesting proliferation-related functionality

(Figure 7H). Intriguingly, for P6 plus P9, where the mLT-EV proteins were uniquely

upregulated irrespective of the cellular transcriptome, plenty of organelle-related terms was

enriched (Figure 7I). Expectedly, metabolism-related terms were enriched in P7,

suggesting consistent downregulation in hepatocytes and their EVs during liver

regeneration (Figure S3B). To verify the bioinformatic results, we additionally performed

Western blot analysis on several organelle and cell cycle markers in mLT-EVs.

Accumulation of Calnexin, an endoplasmic reticulum (ER) protein, and the presence of

Cyclin-dependent kinase 1 (Cdk1), a cell cycle regulator, was detected in ASGPR$^+$mLT-EVs

from the PHx mice (Figure S3C). Taken together, these results suggest that EVs diversify

hepatocyte-derived information in liver regeneration.

Figure 7

Figure 7. EVs diversify hepatocyte-derived information in liver regeneration. (A) GO

enrichment analysis of DEPs between PHx and Sham mLT-EVs shown by the circle

diagram. **(B-D)** GO enrichment analysis of DEPs between PHx and Sham mLT-EVs shown

by functional cluster analysis. Q categories: Q1, upregulated more than two folds; Q2.

upregulated less than two folds; Q3, downregulated less than two folds; Q4, downregulated

more than two folds. **(E)** KEGG enrichment analysis of DEPs between PHx and Sham

mLT-EVs shown by the circle diagram. **(F)** KEGG enrichment analysis of DEPs between

PHx and Sham mLT-EVs shown by functional cluster analysis. Q categories: Q1,

upregulated more than two folds; Q2. upregulated less than two folds; Q3, downregulated

less than two folds; Q4, downregulated more than two folds. **(G)** Nine-quadrant diagram

showing integrated analysis of hepatocyte transcriptome from scRNA-seq and mLT-EV

proteome. **(H)** GO enrichment analysis of the P3 quadrant showing consistent upregulation

between the hepatocyte genes and the mLT-EV proteins. **(I)** GO enrichment analysis of the

P6 plus P9 quadrants showing upregulation of the mLT-EV proteins despite unchange or

downregulation of hepatocyte genes.

3. Discussion

The liver is responsible for maintaining metabolism and detoxification of the body and

is highly regenerative [1, 2]. The process of liver regeneration involves complex intercellular

communication between parenchymal cells and NPCs [3]. However, the mechanism of this

elaborate functional interaction has not been thoroughly understood. EVs are biologically

active paracrine messengers which act dynamically in various pathophysiological states [12,

13]. In the present study, we mapped the transcriptomic landscape of liver regeneration at

the single cell level, which highlighted the crucial role of hepatocyte-centered intercellular

communication based on EVs. Our investigation further uncovered that liver regeneration

promoted the release of LT-EVs, particularly hepatocyte-derived ASGPR$^+$mLT-EVs, which

can be specifically collected from the liver tissue following a multi-step protocol involving

enzymatic treatment, immunomagnetic separation, and differential centrifugation. Notably,

ASGPR$^+$mLT-EVs exhibited diversified proteomic cargoes with their parental hepatocytes,

which not only inherited regulated regenerative and metabolic signals at the cellular level

but also showed vesicular enrichment of organelle components. For the first time, this study

provides an integrated characterization of the hepatocyte transcriptome in related to the EV

proteome in liver regeneration, offering insights into the biological and mechanistic aspects

of the regenerating liver.

The liver possesses the remarkable ability to fully regenerate after undergoing partial

resection [2]. Studies have documented that liver regeneration involves a combination of

subsequent hepatocyte division, hypertrophy, and proliferation [8, 24]. In our scRNA-seq,

data further suggested significant influences of liver regeneration on hepatocyte metabolic

processes, which largely suppressed in the regenerating liver. Indeed, hepatic metabolic

reprogramming after PHx represent an important event, and the transcriptome sequencing

combined with metabolomic analysis has indicated detailed metabolic changes during liver

regeneration [28, 29]. Our findings, together with others' findings and despite preliminary,

suggest that hepatocytes might sacrifice metabolic activity for safeguarding regenerative

expansion, and that hepatocytes are equipped with flexible metabolic machinery able to

adapt dynamically to tissue regeneration [29]. It is also noteworthy that scRNA-seq analysis

revealed a substantial involvement of the immune system in liver regeneration, which is in

accordance with the current understanding [3, 10]. Our data further demonstrated that the

proportions of certain immune cell types, such as B cells and neutrophils, were increased

after PHx, and that the communication between hepatocytes and various immune cells

potentially contributed to liver regeneration. Future research is needed to examine in-depth

the reciprocal regulation mechanisms between hepatocytes and detailed immune cells that

safeguard liver regeneration.

Researchers worldwide have recently dedicated significant efforts to investigating EVs.

These membranous nanoparticles are discovered to escape from the autophagy-lysosomal

pathway and be release into the circulation and the extracellular space of tissues [30, 31].

Particularly, studies have shown the potential of Ti-EVs in promoting tissue regeneration,

and LT-EVs have been employed to expedite the hepatic disease recovery process [15, 16].

However, the specific cellular origin of Ti-EVs responsible for the action remains unknown.

In this study, we established an immunomagnetic microbead assay to specifically isolate

LT-EVs derived from hepatocytes in the regenerating liver. As assessed by TEM and NTA,

the isolation procedure did not exert significant impact on the LT-EV morphology and size

distribution, suggesting that the properties of these Ti-EVs were largely maintained. Indeed,

with advantages of potent specificity and efficiency, the immunomagnetic separation assay

based on immunomagnetic particles has been acknowledged as a powerful system for the

enrichment of target marked cells [32, 33]. Utilization of this approach in the EV research

will benefit the in-depth biological and mechanistic investigations of endogenous Ti-EVs, as

well as t diagnostic and therapeutic applications. In addition, this approach can be applied

to gain profound insights into crucial subpopulations of EVs not only derived from specific

cell sources but also possessing featured membrane proteins. Future examinations should

elucidate the distinct functionalities of LT-EVs from diverse cellular sources in liver health

and disease. Moreover, the communications between hepatocytes and other liver cell types

potentially mediated by ASGPR⁺ LT-EVs and functionally regulating liver regeneration will

require experimental dissection.

The pivotal role EVs play in regulating liver health and disease has been suggested,

but the function of endogenous LT-EVs remains not fully understood [34]. It is worth noting

that our integrated analysis of hepatocyte transcriptome and ASGPR⁺mLT-EV proteome

indicated two aspects of findings: on one hand, ASGPR⁺mLT-EVs inherited proliferative

messages and reduced metabolic cues from donor hepatocytes in liver regeneration; on

the other hand, ASGPR⁺mLT-EVs were particularly enriched for organelle components with

no significant changes of hepatocyte transcription. For the former aspect, epidermal growth

factor receptor (EGFR) is well known as a crucial regulator safeguarding liver regeneration,

and multiple biological enzymes in the mitochondrial respiratory electron transport chain

contribute to the cell cycle and proliferative regulation of hepatocytes [35, 36]. For the latter

aspect, interestingly, ER represents the major site of protein synthesis in the cell and form

the endosomal secretory pathway with Golgi apparatus [37]. Moreover, emerging studies

have documented that the EV biogenesis and secretion can be dependent on ER, and that

ER stress can regulate the release of EVs [38-40]. Additionally, ER proteostasis is essential

to adapt to cell proliferation and regeneration in development [41]. It has recently been

reported that the EV cargoes are not randomly but specifically selected [42]. Therefore,

future research is needed to further investigate the function of ER in regulating hepatic

regenerative EVs and orchestrating liver regeneration through EV-mediated intercellular

communication.

461   In summary, our findings provide the first knowledge that hepatocyte-specific tissue

462 vesicles are indeed phenotypically and functionally involved in liver regeneration. Our study

463 paves an avenue for in-depth biological and mechanistic research of regenerative LT-EVs

464 and sheds light on extensive dissection of antigen-marked, physiological, and endogenous

465 tissue EV populations.

References

1. Michalopoulos, G. K.; Bhushan, B., *Nat Rev Gastroenterol Hepatol* **2021,** *18* (1), 40-55. DOI 10.1038/s41575-020-0342-4.

2. Michalopoulos, G. K., *Am J Pathol* **2010,** *176* (1), 2-13. DOI 10.2353/ajpath.2010.090675.

3. Campana, L.; Esser, H.; Huch, M.; Forbes, S., *Nat Rev Mol Cell Biol* **2021,** *22* (9), 608-624. DOI 10.1038/s41580-021-00373-7.

4. Ait Ahmed, Y.; Fu, Y.; Rodrigues, R. M.; He, Y.; Guan, Y.; Guillot, A.; Ren, R.; Feng, D.; Hidalgo, J.; Ju, C.; Lafdil, F.; Gao, B., *Cellular & molecular immunology* **2021,** *18* (9), 2165-2176. DOI 10.1038/s41423-021-00731-7.

5. Shimizu, H.; Miyazaki, M.; Wakabayashi, Y.; Mitsuhashi, N.; Kato, A.; Ito, H.; Nakagawa, K.; Yoshidome, H.; Kataoka, M.; Nakajima, N., *J Hepatol* **2001,** *34* (5), 683-9. DOI 10.1016/s0168-8278(00)00055-6.

6. Xiao, T.; Meng, W.; Jin, Z.; Wang, J.; Deng, J.; Wen, J.; Liu, B.; Liu, M.; Bai, J.; Liu, F., *Communications biology* **2022,** *5* (1), 771. DOI 10.1038/s42003-022-03714-0.

7. Song, J.; Ma, J.; Liu, X.; Huang, Z.; Li, L.; Li, L.; Luo, L.; Ni, R.; He, J., *NPJ Regenerative medicine* **2023,** *8* (1), 20. DOI 10.1038/s41536-023-00294-3.

8. Yanger, K.; Knigin, D.; Zong, Y.; Maggs, L.; Gu, G.; Akiyama, H.; Pikarsky, E.; Stanger, B. Z., *Cell Stem Cell* **2014,** *15* (3), 340-349. DOI 10.1016/j.stem.2014.06.003.

9. Mei, Y.; Thevananther, S., *Hepatology* **2011,** *54* (5), 1777-89. DOI 10.1002/hep.24560.

10. Forbes, S. J.; Newsome, P. N., *Nat Rev Gastroenterol Hepatol* **2016,** *13* (8), 473-85. DOI 10.1038/nrgastro.2016.97.

11. Stravitz, R. T.; Lee, W. M., *Lancet* **2019,** *394* (10201), 869-881. DOI

10.1016/S0140-6736(19)31894-X.

12. S, E. L. A.; Mager, I.; Breakefield, X. O.; Wood, M. J., *Nat Rev Drug Discov* **2013,** *12* (5),

347-57. DOI 10.1038/nrd3978.

13. Yanez-Mo, M.; Siljander, P. R.; Andreu, Z.; Zavec, A. B.; Borras, F. E.; Buzas, E. I.;

Buzas, K.; Casal, E.; Cappello, F.; Carvalho, J.; Colas, E.; Cordeiro-da Silva, A.; Fais, S.;

Falcon-Perez, J. M.; Ghobrial, I. M.; Giebel, B.; Gimona, M.; Graner, M.; Gursel, I.; Gursel,

M.; Heegaard, N. H.; Hendrix, A.; Kierulf, P.; Kokubun, K.; Kosanovic, M.; Kralj-Iglic, V.;

Kramer-Albers, E. M.; Laitinen, S.; Lasser, C.; Lener, T.; Ligeti, E.; Line, A.; Lipps, G.;

Llorente, A.; Lotvall, J.; Mancek-Keber, M.; Marcilla, A.; Mittelbrunn, M.; Nazarenko, I.;

Nolte-'t Hoen, E. N.; Nyman, T. A.; O'Driscoll, L.; Olivan, M.; Oliveira, C.; Pallinger, E.; Del

Portillo, H. A.; Reventos, J.; Rigau, M.; Rohde, E.; Sammar, M.; Sanchez-Madrid, F.;

Santarem, N.; Schallmoser, K.; Ostenfeld, M. S.; Stoorvogel, W.; Stukelj, R.; Van der Grein,

S. G.; Vasconcelos, M. H.; Wauben, M. H.; De Wever, O., *J Extracell Vesicles* **2015,** *4*,

27066. DOI 10.3402/jev.v4.27066.

14. Qin, B.; Hu, X. M.; Su, Z. H.; Zeng, X. B.; Ma, H. Y.; Xiong, K., *Pathol Res Pract* **2021,**

226, 153604. DOI 10.1016/j.prp.2021.153604.

15. Lou, P.; Liu, S.; Wang, Y.; Lv, K.; Zhou, X.; Li, L.; Zhang, Y.; Chen, Y.; Cheng, J.; Lu, Y.;

Liu, J., *Adv Mater* **2023,** *35* (33), e2300602. DOI 10.1002/adma.202300602.

16. Lee, J.; Kim, S. R.; Lee, C.; Jun, Y. I.; Bae, S.; Yoon, Y. J.; Kim, O. Y.; Gho, Y. S., *J*

Extracell Vesicles **2021,** *10* (10), e12133. DOI 10.1002/jev2.12133.

17. Verweij, F. J.; Balaj, L.; Boulanger, C. M.; Carter, D. R. F.; Compeer, E. B.; D'Angelo, G.;

El Andaloussi, S.; Goetz, J. G.; Gross, J. C.; Hyenne, V.; Kramer-Albers, E. M.; Lai, C. P.; Loyer, X.; Marki, A.; Momma, S.; Nolte-'t Hoen, E. N. M.; Pegtel, D. M.; Peinado, H.; Raposo, G.; Rilla, K.; Tahara, H.; Thery, C.; van Royen, M. E.; Vandenbroucke, R. E.; Wehman, A. M.; Witwer, K.; Wu, Z.; Wubbolts, R.; van Niel, G., *Nat Methods* **2021,** *18* (9), 1013-1026. DOI 10.1038/s41592-021-01206-3.

18. Crescitelli, R.; Lasser, C.; Lotvall, J., *Nat Protoc* **2021,** *16* (3), 1548-1580. DOI 10.1038/s41596-020-00466-1.

19. Vella, L. J.; Scicluna, B. J.; Cheng, L.; Bawden, E. G.; Masters, C. L.; Ang, C. S.; Willamson, N.; McLean, C.; Barnham, K. J.; Hill, A. F., *J Extracell Vesicles* **2017,** *6* (1), 1348885. DOI 10.1080/20013078.2017.1348885.

20. Nojima, H.; Freeman, C. M.; Schuster, R. M.; Japtok, L.; Kleuser, B.; Edwards, M. J.; Gulbins, E.; Lentsch, A. B., *J Hepatol* **2016,** *64* (1), 60-8. DOI 10.1016/j.jhep.2015.07.030.

21. Mitchell, C.; Willenbring, H., *Nat Protoc* **2008,** *3* (7), 1167-70. DOI 10.1038/nprot.2008.80.

22. Cao, Y.; Qiu, J. Y.; Chen, D.; Li, C. Y.; Xing, S. J.; Zheng, C. X.; Liu, X.; Jin, Y.; Sui, B. D., *J Vis Exp* **2022,** (188). DOI 10.3791/63990.

23. Hans, F.; Dimitrov, S., *Oncogene* **2001,** *20* (24), 3021-7. DOI 10.1038/sj.onc.1204326.

24. Miyaoka, Y.; Ebato, K.; Kato, H.; Arakawa, S.; Shimizu, S.; Miyajima, A., *Curr Biol* **2012,** *22* (13), 1166-75. DOI 10.1016/j.cub.2012.05.016.

25. Zheng, C.; Sui, B.; Zhang, X.; Hu, J.; Chen, J.; Liu, J.; Wu, D.; Ye, Q.; Xiang, L.; Qiu, X.; Liu, S.; Deng, Z.; Zhou, J.; Liu, S.; Shi, S.; Jin, Y., *J Extracell Vesicles* **2021,** *10* (7), e12109. DOI 10.1002/jev2.12109.

26. D'Souza, A. A.; Devarajan, P. V., *Journal of controlled release : official journal of the*

Controlled Release Society **2015,** *203*, 126-39. DOI 10.1016/j.jconrel.2015.02.022.

27. Thery, C.; Witwer, K. W.; Aikawa, E.; Alcaraz, M. J.; Anderson, J. D.; Andriantsitohaina,

R.; Antoniou, A.; Arab, T.; Archer, F.; Atkin-Smith, G. K.; Ayre, D. C.; Bach, J. M.; Bachurski,

D.; Baharvand, H.; Balaj, L.; Baldacchino, S.; Bauer, N. N.; Baxter, A. A.; Bebawy, M.;

Beckham, C.; Bedina Zavec, A.; Benmoussa, A.; Berardi, A. C.; Bergese, P.; Bielska, E.;

Blenkiron, C.; Bobis-Wozowicz, S.; Boilard, E.; Boireau, W.; Bongiovanni, A.; Borras, F. E.;

Bosch, S.; Boulanger, C. M.; Breakefield, X.; Breglio, A. M.; Brennan, M. A.; Brigstock, D. R.;

Brisson, A.; Broekman, M. L.; Bromberg, J. F.; Bryl-Gorecka, P.; Buch, S.; Buck, A. H.;

Burger, D.; Busatto, S.; Buschmann, D.; Bussolati, B.; Buzas, E. I.; Byrd, J. B.; Camussi, G.;

Carter, D. R.; Caruso, S.; Chamley, L. W.; Chang, Y. T.; Chen, C.; Chen, S.; Cheng, L.; Chin,

A. R.; Clayton, A.; Clerici, S. P.; Cocks, A.; Cocucci, E.; Coffey, R. J.; Cordeiro-da-Silva, A.;

Couch, Y.; Coumans, F. A.; Coyle, B.; Crescitelli, R.; Criado, M. F.; D'Souza-Schorey, C.;

Das, S.; Datta Chaudhuri, A.; de Candia, P.; De Santana, E. F.; De Wever, O.; Del Portillo,

H. A.; Demaret, T.; Deville, S.; Devitt, A.; Dhondt, B.; Di Vizio, D.; Dieterich, L. C.; Dolo, V.;

Dominguez Rubio, A. P.; Dominici, M.; Dourado, M. R.; Driedonks, T. A.; Duarte, F. V.;

Duncan, H. M.; Eichenberger, R. M.; Ekstrom, K.; El Andaloussi, S.; Elie-Caille, C.;

Erdbrugger, U.; Falcon-Perez, J. M.; Fatima, F.; Fish, J. E.; Flores-Bellver, M.; Forsonits, A.;

Frelet-Barrand, A.; Fricke, F.; Fuhrmann, G.; Gabrielsson, S.; Gamez-Valero, A.; Gardiner,

C.; Gartner, K.; Gaudin, R.; Gho, Y. S.; Giebel, B.; Gilbert, C.; Gimona, M.; Giusti, I.;

Goberdhan, D. C.; Gorgens, A.; Gorski, S. M.; Greening, D. W.; Gross, J. C.; Gualerzi, A.;

Gupta, G. N.; Gustafson, D.; Handberg, A.; Haraszti, R. A.; Harrison, P.; Hegyesi, H.;

Hendrix, A.; Hill, A. F.; Hochberg, F. H.; Hoffmann, K. F.; Holder, B.; Holthofer, H.;

Hosseinkhani, B.; Hu, G.; Huang, Y.; Huber, V.; Hunt, S.; Ibrahim, A. G.; Ikezu, T.; Inal, J. M.;

Isin, M.; Ivanova, A.; Jackson, H. K.; Jacobsen, S.; Jay, S. M.; Jayachandran, M.; Jenster,

G.; Jiang, L.; Johnson, S. M.; Jones, J. C.; Jong, A.; Jovanovic-Talisman, T.; Jung, S.;

Kalluri, R.; Kano, S. I.; Kaur, S.; Kawamura, Y.; Keller, E. T.; Khamari, D.; Khomyakova, E.;

Khvorova, A.; Kierulf, P.; Kim, K. P.; Kislinger, T.; Klingeborn, M.; Klinke, D. J., 2nd; Kornek,

M.; Kosanovic, M. M.; Kovacs, A. F.; Kramer-Albers, E. M.; Krasemann, S.; Krause, M.;

Kurochkin, I. V.; Kusuma, G. D.; Kuypers, S.; Laitinen, S.; Langevin, S. M.; Languino, L. R.;

Lannigan, J.; Lasser, C.; Laurent, L. C.; Lavieu, G.; Lazaro-Ibanez, E.; Le Lay, S.; Lee, M.

S.; Lee, Y. X. F.; Lemos, D. S.; Lenassi, M.; Leszczynska, A.; Li, I. T.; Liao, K.; Libregts, S.

F.; Ligeti, E.; Lim, R.; Lim, S. K.; Line, A.; Linnemannstons, K.; Llorente, A.; Lombard, C. A.;

Lorenowicz, M. J.; Lorincz, A. M.; Lotvall, J.; Lovett, J.; Lowry, M. C.; Loyer, X.; Lu, Q.;

Lukomska, B.; Lunavat, T. R.; Maas, S. L.; Malhi, H.; Marcilla, A.; Mariani, J.; Mariscal, J.;

Martens-Uzunova, E. S.; Martin-Jaular, L.; Martinez, M. C.; Martins, V. R.; Mathieu, M.;

Mathivanan, S.; Maugeri, M.; McGinnis, L. K.; McVey, M. J.; Meckes, D. G., Jr.; Meehan, K.

L.; Mertens, I.; Minciacchi, V. R.; Moller, A.; Moller Jorgensen, M.; Morales-Kastresana, A.;

Morhayim, J.; Mullier, F.; Muraca, M.; Musante, L.; Mussack, V.; Muth, D. C.; Myburgh, K.

H.; Najrana, T.; Nawaz, M.; Nazarenko, I.; Nejsum, P.; Neri, C.; Neri, T.; Nieuwland, R.;

Nimrichter, L.; Nolan, J. P.; Nolte-'t Hoen, E. N.; Noren Hooten, N.; O'Driscoll, L.; O'Grady,

T.; O'Loghlen, A.; Ochiya, T.; Olivier, M.; Ortiz, A.; Ortiz, L. A.; Osteikoetxea, X.; Ostergaard,

O.; Ostrowski, M.; Park, J.; Pegtel, D. M.; Peinado, H.; Perut, F.; Pfaffl, M. W.; Phinney, D.

G.; Pieters, B. C.; Pink, R. C.; Pisetsky, D. S.; Pogge von Strandmann, E.; Polakovicova, I.;

Poon, I. K.; Powell, B. H.; Prada, I.; Pulliam, L.; Quesenberry, P.; Radeghieri, A.; Raffai, R.

L.; Raimondo, S.; Rak, J.; Ramirez, M. I.; Raposo, G.; Rayyan, M. S.; Regev-Rudzki, N.;

Ricklefs, F. L.; Robbins, P. D.; Roberts, D. D.; Rodrigues, S. C.; Rohde, E.; Rome, S.;

Rouschop, K. M.; Rughetti, A.; Russell, A. E.; Saa, P.; Sahoo, S.; Salas-Huenuleo, E.;

Sanchez, C.; Saugstad, J. A.; Saul, M. J.; Schiffelers, R. M.; Schneider, R.; Schoyen, T. H.;

Scott, A.; Shahaj, E.; Sharma, S.; Shatnyeva, O.; Shekari, F.; Shelke, G. V.; Shetty, A. K.;

Shiba, K.; Siljander, P. R.; Silva, A. M.; Skowronek, A.; Snyder, O. L., 2nd; Soares, R. P.;

Sodar, B. W.; Soekmadji, C.; Sotillo, J.; Stahl, P. D.; Stoorvogel, W.; Stott, S. L.; Strasser, E.

F.; Swift, S.; Tahara, H.; Tewari, M.; Timms, K.; Tiwari, S.; Tixeira, R.; Tkach, M.; Toh, W. S.;

Tomasini, R.; Torrecilhas, A. C.; Tosar, J. P.; Toxavidis, V.; Urbanelli, L.; Vader, P.; van

Balkom, B. W.; van der Grein, S. G.; Van Deun, J.; van Herwijnen, M. J.; Van

Keuren-Jensen, K.; van Niel, G.; van Royen, M. E.; van Wijnen, A. J.; Vasconcelos, M. H.;

Vechetti, I. J., Jr.; Veit, T. D.; Vella, L. J.; Velot, E.; Verweij, F. J.; Vestad, B.; Vinas, J. L.;

Visnovitz, T.; Vukman, K. V.; Wahlgren, J.; Watson, D. C.; Wauben, M. H.; Weaver, A.;

Webber, J. P.; Weber, V.; Wehman, A. M.; Weiss, D. J.; Welsh, J. A.; Wendt, S.; Wheelock,

A. M.; Wiener, Z.; Witte, L.; Wolfram, J.; Xagorari, A.; Xander, P.; Xu, J.; Yan, X.; Yanez-Mo,

M.; Yin, H.; Yuana, Y.; Zappulli, V.; Zarubova, J.; Zekas, V.; Zhang, J. Y.; Zhao, Z.; Zheng, L.;

Zheutlin, A. R.; Zickler, A. M.; Zimmermann, P.; Zivkovic, A. M.; Zocco, D.; Zuba-Surma, E.

K., *J Extracell Vesicles* **2018,** *7* (1), 1535750. DOI 10.1080/20013078.2018.1535750.

28. Li, Z.; Peng, B.; Chen, S.; Li, J.; Hu, K.; Liao, L.; Xie, Q.; Yao, M.; Liang, L.; Tomlinson,

S.; Yuan, G.; He, S., *BMC genomics* **2023,** *24* (1), 532. DOI 10.1186/s12864-023-09647-0.

29. Caldez, M. J.; Van Hul, N.; Koh, H. W. L.; Teo, X. Q.; Fan, J. J.; Tan, P. Y.; Dewhurst, M.

R.; Too, P. G.; Talib, S. Z. A.; Chiang, B. E.; Stunkel, W.; Yu, H.; Lee, P.; Fuhrer, T.; Choi, H.; Bjorklund, M.; Kaldis, P., *Developmental cell* **2018,** *47* (4), 425-438 e5. DOI 10.1016/j.devcel.2018.09.020.

30. Cao, Y.; Ying, S. Q.; Qiu, X. Y.; Guo, J.; Chen, C.; Li, S. J.; Dou, G.; Zheng, C. X.; Chen, D.; Qiu, J. Y.; Jin, Y.; Sui, B. D.; Jin, F., *Neurobiol Stress* **2023,** *22*, 100513. DOI 10.1016/j.ynstr.2023.100513.

31. Perrin, P.; Janssen, L.; Janssen, H.; van den Broek, B.; Voortman, L. M.; van Elsland, D.; Berlin, I.; Neefjes, J., *Curr Biol* **2021,** *31* (17), 3884-3893 e4. DOI 10.1016/j.cub.2021.06.022.

32. Tang, P.; Thongrom, B.; Arora, S.; Haag, R., *Adv Healthc Mater* **2023,** *12* (26), e2300842. DOI 10.1002/adhm.202300842.

33. Shin, H. S.; Park, J.; Lee, S. Y.; Yun, H. G.; Kim, B.; Kim, J.; Han, S.; Cho, D.; Doh, J.; Choi, S., *Small* **2023**, e2302809. DOI 10.1002/smll.202302809.

34. Balaphas, A.; Meyer, J.; Sadoul, R.; Morel, P.; Gonelle-Gispert, C.; Buhler, L. H., *Liver Int* **2019,** *39* (10), 1801-1817. DOI 10.1111/liv.14189.

35. López-Luque, J.; Caballero-Díaz, D.; Martinez-Palacián, A.; Roncero, C.; Moreno-Càceres, J.; García-Bravo, M.; Grueso, E.; Fernández, A.; Crosas-Molist, E.; García-Álvaro, M.; Addante, A.; Bertran, E.; Valverde, A. M.; González-Rodríguez, Á.; Herrera, B.; Montoliu, L.; Serrano, T.; Segovia, J. C.; Fernández, M.; Ramos, E.; Sánchez, A.; Fabregat, I., *Hepatology* **2016,** *63* (2), 604-19.

36. Purhonen, J.; Banerjee, R.; Wanne, V.; Sipari, N.; Morgelin, M.; Fellman, V.; Kallijarvi, J., *Nat Commun* **2023,** *14* (1), 2356. DOI 10.1038/s41467-023-38027-1.

37. Tang, V. T.; Ginsburg, D., *J Clin Invest* **2023,** *133* (1). DOI 10.1172/JCI163838.

38. Verweij, F. J.; Bebelman, M. P.; George, A. E.; Couty, M.; Becot, A.; Palmulli, R.;

Heiligenstein, X.; Sires-Campos, J.; Raposo, G.; Pegtel, D. M.; van Niel, G., *The Journal of*

cell biology **2022,** *221* (12). DOI 10.1083/jcb.202112032.

39. Barman, B.; Sung, B. H.; Krystofiak, E.; Ping, J.; Ramirez, M.; Millis, B.; Allen, R.;

Prasad, N.; Chetyrkin, S.; Calcutt, M. W.; Vickers, K.; Patton, J. G.; Liu, Q.; Weaver, A. M.,

Developmental cell **2022,** *57* (8), 974-994 e8. DOI 10.1016/j.devcel.2022.03.012.

40. Ye, J.; Liu, X., *Front Immunol* **2022,** *13*, 955419. DOI 10.3389/fimmu.2022.955419.

41. Yang, S.; Jiang, H.; Bian, W.; Xu, W.; Guo, Y.; Song, S.; Zheng, J.; Kuang, X.; Wu, C.;

Ding, X.; Guo, X.; Xue, L.; Yu, Z.; Zhang, Y.; Ryoo, H. D.; Li, X.; Ma, X., *Proc Natl Acad Sci*

U S A **2022,** *119* (42), e2202133119. DOI 10.1073/pnas.2202133119.

42. Dixson, A. C.; Dawson, T. R.; Di Vizio, D.; Weaver, A. M., *Nat Rev Mol Cell Biol* **2023,**

24 (7), 454-476. DOI 10.1038/s41580-023-00576-0.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (82371020 to B.D.S., 82301028 to C.X.Z., 81930025 to Y.J. and 82170988 to F.J.) and the Young Science and Technology Rising Star Project of Shaanxi Province (2023KJXX-027 to B.D.S.). We are grateful for the assistance of the National Experimental Teaching Demonstration Center for Basic Medicine (AMFU). We thank PTM Biolabs Co., Ltd (Hangzhou, China) for help in analyzing the proteomic data. We thank Mr. Tao Liu (Gene Denovo Biotechnology Co, Ltd, 790 Guangzhou, China) for his assistance in analyzing the single-cell sequencing data.

Author contributions

S.Q.Y., Y.C., L.L., and Z.K.Z. contributed equally to the design and conduction of the study and drafted the manuscript. X.Y.L. analyzed and interpreted data. C.H.L. performed partial hepatectomy. S.Y.C. contributed to the scRNA-seq experiment. X.L. contributed to the proteome experiment. S.J.X. contributed to H&E and IF experiments. K.S. contributed to TEM and flow cytometry analysis. J.Y.Q. and Y.Y.L. contributed to NTA and WB experiments. C.X.Z. and F.J. contributed to the discussion and interpretation of data. Y.J. and B.D.S. conceived the project, designed and supervised the experiments. All authors revised the manuscript and approved the final version.

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

All data included in this study are available upon reasonable request of corresponding authors.

Abbreviations

ASGPR, asialoglycoprotein receptor; BECs, biliary epithelial cells; BSA, bovine serum albumin; CCl_4, carbon tetrachloride; Cdk1, cyclin-dependent kinase 1; DEGs, differentially expressed genes; DEPs, differentially expressed proteins; EGFR, epidermal growth factor receptor; ER, endoplasmic reticulum; EVs, extracellular vesicles; GEO, Gene Expression Omnibus; GO, Gene Ontology; HRP, horseradish peroxidase; IF, immunofluorescence; KEGG, Kyoto Encyclopedia of Genes and Genomes; LC-MS/MS, Liquid chromatography with tandem mass spectrometry; LT-EVs, liver tissue extracellular vesicles; mLT-EVs, magnetic microbead-sorted liver tissue extracellular vesicles; NPCs, non-parenchymal cells; NTA, nanoparticle tracking analysis; PFA, paraformaldehyde; PHx, partial hepatectomy; PVDF, polyvinylidene difluoride; SD, standard deviation; TEM, transmission electron microscopy; Ti-EVs, tissue-derived EVs; TSG101, tumor suppressor gene 101; tSNE, t-distributed stochastic neighbor embedding.

A

LT-EVs

ASPGR⁻LT-EVs

ASGPR⁺mLT-EVs

Counts (%)

ASGPR

B TEM

Sham^LT-EVs   PHx^LT-EVs

C

CD63
TSG101
Flotillin-1
Golgin84

D NTA

E NTA

F BCA

G

H NTA

I BCA

A
PPI network analysis of DEPs

B
GO enrichment of P7

oxoacid metabolic process

carboxylic acid metabolic process

organic acid metabolic process

small molecule metabolic process

small molecule catabolic process

carboxylic acid catabolic process

organic acid catabolic process

alpha-amino acid metabolic process

monocarboxylic acid metabolic process

small molecule biosynthetic process

catabolic process

cellular amino acid metabolic process

alpha-amino acid catabolic process

Gene number
- 18
- 27.25
- 36.5
- 45.75
- 55

$-\log_{10}(FDR)$
- 25
- 22.5
- 20
- 17.5

Gene Ratio

0.02   0.04   0.06   0.08

C

Sham$^{mLT-EVs}$ PHx$^{mLT-EVs}$

Calnexin

Cdk1

CD63

A tSNE of total liver cells

Hepatocytes
B cells
T cells
Erythrocytes
Endothelial cells
Neutrophils
Macrophages
Plasma B cells

B

Hepatocytes
T cells
Erythrocytes
B cells
Endothelial cells
Neutrophils
Macrophages
Plasma B cells

C Sham

PHx

D

956 394 64 163 1186 1365 1619 1464

E GO analysis of hepatocyte DEGs

F KEGG analysis of hepatocyte DEGs

Liver tissue TEM

Specific Tissue EV Collection

Sample preparation

Dissociation

Liver tissue dissociation

Liberase TM
37°C 30 min

Digestion

Enzymatic incubation

Filtration

300 g, 10 min, 4°C
2,000 g, 20 min, 4°C

16,800 g,
30 min, 4°C

Differential centrifugantion

Liver tissue EVs
(LT-EVs)

Magnetic isolation

Antibody incubation

LT-EVs

Add ASGPR-PE antibody

4 h, 4°C, in dark place

Resuspension

ASGPR-PE incubation

Magnetic bead incubation

Add anti-PE magnetic microbeads

15 min, 4°C, in dark place

Beads incubation

Magnetic sorting

Passing *via* the LD column

Magnetic microbead-incubated LT-EVs

ASGPR⁻ LT-EVs

Collection

16,800 g, 30 min, 4°C

Magnetic microbead-sorted LT-EVs, mLT-EVs, ASGPR⁺)

Specific Tissue EV Characterization

EV morphology	EV assay	EV profiling
TEM NTA	Flow cytometer Western blot	Mass spectrometry