Comparative analysis of *Wolbachia* maternal transmission and localization in host ovaries

Michael T.J. Hague[a,#], Timothy B. Wheeler[b], Brandon S. Cooper[b]

[a]Biology Department
University of Scranton
800 Linden Street
Scranton, PA 18510

[b]Division of Biological Sciences
University of Montana
32 Campus Dr.
Missoula, MT 59812

[#]Corresponding Author
Michael T.J. Hague
Biology Department, Loyola Science Center
800 Linden Street, University of Scranton
Scranton, PA 18510
(570) 941-7962
michael.hague@scranton.edu

ABSTRACT

Many insects and other animals carry microbial endosymbionts that influence their

reproduction and fitness. These relationships only persist if endosymbionts are reliably

transmitted from one host generation to the next. *Wolbachia* are maternally transmitted

endosymbionts found in most insect species, but transmission rates can vary across

environments. Maternal transmission of *w*Mel *Wolbachia* depends on temperature in natural

Drosophila melanogaster hosts and in transinfected *Aedes aegypti*, where *w*Mel is used to block

pathogens that cause human disease. In *D. melanogaster*, *w*Mel transmission declines in the cold

as *Wolbachia* become less abundant in host ovaries and at the posterior pole plasm (the site of

germline formation) in mature oocytes. Here, we assess how temperature affects maternal

transmission and underlying patterns of *Wolbachia* localization across 10 *Wolbachia* strains

diverged up to 50 million years—including strains closely related to *w*Mel—and their natural

Drosophila hosts. Many *Wolbachia* maintain high transmission rates across temperatures, despite

highly variable (and sometimes low) levels of *Wolbachia* in the ovaries and at the developing

germline in late-stage oocytes. Identifying strains like closely related *w*Mel-like *Wolbachia* with

stable transmission across variable environmental conditions may improve the efficacy of

Wolbachia-based biocontrol efforts as they expand into globally diverse environments.

KEYWORDS: *w*Mel, *Drosophila*, host-microbe interaction, vertical transmission

INTRODUCTION

Microbes form diverse relationships with host organisms that span the tree of life, including animals, plants, and protists. Endosymbiosis is an especially close relationship where microbes occupy eukaryotic host cells. Heritable endosymbionts are particularly common in insects[1–3], and these relationships alter fundamental aspects of host biology, including reproduction[4,5], protection for natural enemies[6–9], and nutrient acquisition[10–12].

Many endosymbionts rely on vertical-maternal transmission to spread and persist in insect populations[13–15]. In obligate relationships, these endosymbionts must be maternally transmitted to ensure host survival (e.g., *Buchnera* and aphids[16]). For facultative endosymbionts like *Wolbachia*, imperfect maternal transmission decreases the prevalence of *Wolbachia*-positive individuals within the host population each generation[17–21]. Thus, maternal transmission is a fundamental determinant of how endosymbionts spread, persist, and evolve in host populations[17,22,23]. In addition, the efficacy of *Wolbachia*-based programs to control human diseases[24–26] and agricultural pests[27,28] explicitly requires efficient maternal transmission. This includes introducing the pathogen-blocking *w*Mel *Wolbachia* strain from *Drosophila melanogaster* into mosquito populations to reduce the incidence of dengue and other diseases on multiple continents[29–32]. Nonetheless, it is largely unknown why maternal transmission may breakdown under certain circumstances, especially for non-model systems[15,21,33–36].

Maternal transmission is predicted to depend on endosymbiont density (titer) in female reproductive tissues, and specifically at the site of germline formation during oogenesis[14]. In insects, endosymbionts often achieve vertical transmission by occupying the female ovaries. Endosymbionts are observed in germline stem cells in female hosts where they maintain a tight association with the germline throughout oogenesis, thereby ensuring a germline-to-germline route of vertical transmission through the host matriline[14,15,36,37]. In a more circuitous route, some endosymbionts may also colonize the germline from neighboring somatic cells via cell-to-cell migration through a soma-to-germline route of transmission[4,14,15,38,39]. Our canonical understanding of these cellular processes—and their contributions to maternal transmission—are based almost entirely in the context of standard laboratory conditions where maternal transmission rates are near perfect or perfect[18,20,21,40]. However, maternal transmission of facultative endosymbionts is often imperfect in natural host populations[19,20,41,42] and the cellular processes that lead to transmission breakdown under natural conditions are generally

unresolved[21,36]. Understanding why transmission breaks down under certain conditions is critical

for predicting endosymbiont spread in nature and improving *Wolbachia*-based biocontrol

programs[25,33,34,43].

Maternally transmitted *Wolbachia* are the most common endosymbionts on earth,

associating with about half of all insect species, as well as other arthropods and nematodes[4,44–46].

Wolbachia are primarily maternally (vertically) transmitted via the host germline within host

species; although, host switching via horizonal transfer between host species is common on

evolutionary timescales[47–54]. *Wolbachia* generally form facultative relationships (from the host

perspective) in insects, with some individuals in the host population that do not carry *Wolbachia*

due to imperfect maternal transmission. Maternal transmission is perhaps the most important

determinant of *Wolbachia* prevalence in host populations: theory demonstrates that *Wolbachia*

frequency dynamics and equilibria are approximated by the degree of imperfect maternal

transmission (μ), the relative fitness (e.g., fecundity) of females with *Wolbachia* (F), and the

strength of *Wolbachia*-induced cytoplasmic incompatibility (s_h)[17,55]. Some *Wolbachia* strains

(e.g., *w*Ri in *D. simulans*) cause cytoplasmic incompatibility (CI), a crossing incompatibility that

generates a frequency-dependent advantage for females with *Wolbachia*[5,56].

The prevalence of different *Wolbachia* strains varies widely among host systems, which

may be partially explained by differences in maternal transmission rates. For instance, the strong

CI-causing *w*Ri strain is observed at high frequencies in global *D. simulans* populations[17,18,57],

whereas the weak CI-causing *w*Mel strain is found at intermediate, fluctuating frequencies in *D.*

melanogaster[58]. *w*Ri frequencies can be plausibly explained by strong CI that causes spread to

high frequencies, whereas lower, variable *w*Mel frequencies can be plausibly explained by

maternal transmission breakdown as temperatures decrease, in addition to a minimal influence of

CI[21]. The thermal sensitivity of maternal transmission may be a feature of *w*Mel, given that heat

stress also reduces *w*Mel transmission in transinfected *Aedes aegypti* hosts[33,34,43]. Identifying

Wolbachia variants that are efficiently transmitted across environmental conditions could

potentially contribute to improving the efficacy of *Wolbachia* based biocontrol of human

diseases and agricultural pests[25,26].

What might underlie variable maternal transmission rates? Maternal transmission

breakdown is predicted to stem from reductions in *Wolbachia* abundance and localization

throughout host development and specifically at the critical stages of oogenesis thought to

facilitate germline-to-germline maternal transmission[14,15,21,36,59–63]. *w*Mel cells are present in the

germline stem cells of adult females, and over the course of oogenesis, *w*Mel localize via

kinesin-mediated transport to the pole plasm—the site of germline formation—at the posterior

cortex of the developing oocyte. In late oogenesis (stage 10), *w*Mel *Wolbachia* display a pattern

of posterior localization in oocytes, which is considered to be a key step in the process of

germline-to-germline maternal transmission under standard laboratory conditions[14,50,59–63]. In

support of this, we recently demonstrated that declining *w*Mel transmission in the cold covaries

with a significant reduction in cellular *Wolbachia* abundance at the posterior cortex of stage 10

oocytes, which can plausibly explain why maternal transmission is thermally sensitive[21].

Outside of the *w*Mel strain, it is unclear how temperature influences *Wolbachia* maternal

transmission and posterior localization in stage 10 oocytes for other diverse *Wolbachia*-host

associations[15,20]. Under standard 25°C conditions, divergent *Wolbachia* strains exhibit highly

variable patterns of localization to the pole plasm at the posterior cortex of stage 10 oocytes of

Drosophila species, deviating from our canonical understanding of *w*Mel[15]. These patterns range

from strongly localized (e.g., *w*Ana in *D. anannassae*) to a complete lack of localization (e.g.,

*w*Tri in *D. triauraria*) at the site of germline formation. Some strains, particularly the divergent

B-group *w*Mau strain, also exhibit evidence for an alternative soma-to-germline route of

transmission, whereby *Wolbachia* in somatically derived follicle cells may invade the oocyte via

cell-to-cell migration. It is entirely unknown whether the diverse localization patterns of these

Wolbachia strains influence rates of maternal transmission and/or are sensitive to thermal

perturbation like *w*Mel.

Here, we examine temperature effects on *Wolbachia* maternal transmission and

localization in host tissues across 10 divergent *Wolbachia* strains naturally found in eight

Drosophila host species within the *melanogaster* species group (Figure 1, Table S1). These

Wolbachia comprise nine A-group *Wolbachia*—including closely related *w*Mel-like strains

(*w*Mel, *w*MelCS, *w*Seg in *D. seguyi*, and *w*Cha in *D. chauvacae*) and closely related *w*Ri-like

strains (*w*Ri, *w*Ana in *D. ananassae*, *w*Aura in *D. auraria*, and *w*Tri in *D. triauraria*)[20,51,53]—and

divergent B-group *w*Mau in *D. mauritiana* that diverged up to 46 million years ago from the

other A-group strains[40]. We reared flies at 25° and 20°C and then measured maternal

transmission rates in conjunction with *Wolbachia* densities (titer) at the tissue level in female

ovaries and the remaining somatic tissues. We then used confocal microscopy to examine how

temperature impacts the cellular abundance of *Wolbachia* at the site of germline formation at the

posterior cortex of stage 10 oocytes during oogenesis. Our results reveal diverse patterns of

localization in ovary tissues and late-stage oocytes that are temperature-dependent, but only

weakly predictive of maternal transmission rates. We also identify *Wolbachia* strains that are

transmitted efficiently regardless of temperature.

RESULTS

We estimated the mean rate of maternal transmission ($\pm$ BC$_a$ confidence intervals) for

each *Wolbachia*-host genotype at 25° and 20°C by pairing individual virgin females with

Wolbachia-free males (see Methods), allowing the females to lay for eight days, and then

measuring maternal transmission ($1 - \mu$) to the newly emerged offspring of each subline (Figure

2, Table S2)[20,21]. The final dataset comprised 365 F0 females and 3,612 F1 offspring. Overall, we

found a significant interaction effect between the *Wolbachia*-host genotype and temperature

(GLMM LRT, $\chi^2_{(9)} = 23.49$, $P = 0.005$), such that temperature effects on maternal transmission

varied depending on the *Wolbachia*-host genotype. For most *Wolbachia*-host genotypes, mean

rates of maternal transmission were perfect (e.g., *w*Ri-*D. simulans*) or near-perfect (e.g., *w*Mel-

like *w*Seg-*D. seguyi*) and invariant across the two temperatures (Figure 2, Table S3), with the

exception of a few specific *Wolbachia*-host genotypes. Maternal transmission of *w*Mel by *D.*

melanogaster ($W = 251.5$, $P = 0.003$) and *w*Aura by *D. auraria* ($W = 154$, $P = 0.038$) decreased

significantly at 20°C; although, *w*Mel experienced a much greater decline in the average rate of

transmission (14.9%) than *w*Aura (2.8%). In contrast, B-group *w*Mau transmission by *D.*

mauritiana increased slightly by 3.4% in the cold at 20°C ($W = 117$, $P = 0.019$). Notably, our

analysis also revealed a significant effect of sex (GLMM LRT, $\chi^2_{(1)} = 63.71$, $P < 0.001$), whereby

rates of maternal transmission tended to be lower to female offspring than males (Table S2). This

finding is consistent with our previous findings for *w*Mel[21] and the *w*Mel-like *w*Yak strain in *D.*

yakuba[20], but counter to theoretical expectations that selection should favor faithful transmission

to female offspring[64,65].

To test for temperature effects on *Wolbachia* at the tissue level, we dissected out the

ovaries from a subset of the same F0 females assessed for maternal transmission above ($N =$

183). For this analysis we used qPCR to measure *Wolbachia* densities in the ovaries and the

remaining somatic carcass tissue (Figure 3). We found a significant interaction effect between

the *Wolbachia*-host genotype and temperature on *Wolbachia* density in ovaries (Two-way

ANOVA; $F_{(9,161)} = 4.286$, $P < 0.001$) and the remaining carcasses ($F_{(9,161)} = 10.612$, $P < 0.001$),

such that temperature effects on *Wolbachia* density varied depending on the *Wolbachia*-host

genotype. While we uncovered considerable variation depending on the *Wolbachia*-host

genotype and temperature, *Wolbachia* ovary densities generally tended to decline at 20°C for the

majority of systems (Figure 3, Table S4). Five of the *Wolbachia* strains (*w*Mel, *w*Ri, *w*Ana, *w*Ha,

and *w*Mau) decreased in density in the ovaries in the cold. In contrast, carcasses exhibited

variation in the directionality of changes in *Wolbachia* density. *w*Mel, *w*MelCS, *w*Seg, and

*w*Aura densities decreased significantly in the cold, whereas *w*Cha and *w*Mau significantly

increased. *w*Cha densities in carcasses were roughly an order of magnitude lower than all other

strains, regardless of temperature, suggesting a relatively high degree of localization to the

ovaries. *w*Mel, the strain that experienced the most dramatic decline in transmission in the cold

(Figure 2), was also the only *Wolbachia* strain that experienced a significant decline in

Wolbachia density in both the ovaries and the carcasses in the cold. This included a particularly

large decline in density in the carcasses (a 4.7-fold reduction) compared to the other strains.

For the three *Wolbachia*-host systems with temperature-dependent transmission (*w*Mel,

*w*Aura, and *w*Mau), we also tested for a correlation between transmission rates and *Wolbachia*

density in the ovaries and carcasses of the same individual females (Figure S1). Declining *w*Mel

transmission in the cold was significantly correlated with *Wolbachia* density in the ovaries

(Spearman's $\rho = 0.537$, $P = 0.022$), but not the carcasses ($\rho = 0.392$, $P = 0.108$), providing

additional support that reduced *w*Mel density in reproductive tissues contributes to transmission

breakdown in the cold. Maternal transmission of *w*Aura by *D. auraria* did not strongly correlate

with *w*Aura density in the ovaries ($\rho = -0.244$, $P = 0.380$) or the carcasses ($\rho = 0.035$, $P = 0.902$).

Finally, increasing maternal transmission of *w*Mau by *D. mauritiana* in the cold did not correlate

with *w*Mau density in the ovaries ($\rho = -0.119$, $P = 0.649$), but was significantly correlated with

density in the carcasses ($\rho = 0.549$, $P = 0.022$).

Finally, we examined how temperature impacts cellular *Wolbachia* abundance at the site

of germline formation in stage 10 oocytes during host oogenesis. At this stage, *w*Mel *Wolbachia*

localization at the posterior cortex of the oocyte is considered to be a critical step in the process

of germline-to-germline *Wolbachia* maternal transmission[14,15,21,36,59–63]. We imaged a total of 283

stage 10 oocytes from eight-day-old females reared at the two different temperatures and

quantified cellular *Wolbachia* abundance (measured as corrected total cell fluorescence; CTCF)

in the whole oocytes, the posterior region, and the posterior cortex (Figure S2)[14,15,21]. We defined

the posterior region as the posterior $1/8^{th}$ portion of the oocyte and the posterior cortex as the

narrow cortical region of Vasa expression (a germline protein component)[15,63]. For the sake of

clarity and consistency with the literature, we refer to our *Wolbachia* quantifications with

confocal microscopy here as "cellular *Wolbachia* abundance" in oocytes, as opposed to the

quantification of "*Wolbachia* density" at the tissue level using qPCR (described above).

Cellular *Wolbachia* abundance in whole oocytes (Two-way ANOVA; $F_{(9,263)} = 2.9$, $P =$

0.003), at the posterior region ($F_{(9,263)} = 4.16$, $P < 0.001$), and at the posterior cortex ($F_{(9,263)} =$

2.41, $P = 0.012$) depended on significant two-way interactions between the *Wolbachia*-host

genotype and temperature. As with *Wolbachia* density, cellular *Wolbachia* abundance in oocytes

varied considerably depending on the *Wolbachia*-host genotype and temperature (Figures 4, S3).

The cellular abundance of six of the *Wolbachia* strains (*w*MelCS, *w*Seg, *w*Ana, *w*Aura, and *w*Tri,

and *w*Ha) declined significantly throughout whole oocytes in the cold (Table S5). Temperature

affected fewer strains (only *w*Mel, *w*MelCS, *w*Ri, and *w*Aura) at the posterior region and the

posterior cortex, the location of the developing germline. *w*Mel and *w*Aura, the two strains

transmitted at significantly lower rates in the cold, both declined significantly in abundance in

the posterior oocyte region ($W = 113$, $P = 0.015$; $W = 172$, $P = 0.041$). *w*Mel cellular abundance

also significantly declined at the posterior cortex ($W = 122$, $P = 0.002$). Interestingly, *w*MelCS

also exhibited a significant decline in cellular abundance in the posterior region ($W = 203$, $P <$

0.001) and the cortex ($W = 205$, $P < 0.001$), despite the fact that *w*MelCS transmission by *D.*

melanogaster was not altered in the cold (Figure 2). This can perhaps be explained by the fact

that, of all 10 strains, *w*MelCS had the highest cellular abundance at the posterior region and

cortex, regardless of temperature. Finally, *w*Ri was perfectly transmitted by *D. simulans* at both

temperatures, but *w*Ri abundance increased at the oocyte posterior region ($W = 45$, $P = 0.004$)

and the cortex ($W = 49$, $P = 0.008$) in the cold. Also of note, *w*Aura and *w*Tri occurred at

particularly low abundances in the oocyte posterior region and the posterior cortex, regardless of

temperature, despite both being transmitted at relatively high rates by their female hosts ($1 - \mu >$

0.967).

Our analyses revealed that the diverse *Wolbachia* strains and *Drosophila* host species

exhibit extensive variation in *Wolbachia* density in host tissues (Figure 3) and abundance in host

oocytes (Figure 4). We hypothesized that this diversity may be explained by factors in the

Wolbachia or host genomes (or both). If so, we expect closely related *Wolbachia* strains (or host

species) to exhibit similar patterns of *Wolbachia* localization in host tissues[15,66,67]. We used the

Wolbachia and host trees (Figure 1) to test whether patterns of *Wolbachia* localization in tissues

and oocytes exhibit phylogenetic signal using Pagel's λ[68]. A value of 1 is consistent with trait

evolution that entirely agrees with the *Wolbachia* (or host) phylogeny, suggesting factors in the

Wolbachia genome (or host) contribute to *Wolbachia* localization patterns (i.e., strong

phylogenetic signal). In contrast, a value of 0 is consistent with trait evolution that occurs

independently of phylogenetic relationships[68,69]. We found no evidence that *Wolbachia* density

in the ovaries and carcasses, nor cellular abundance in the oocytes, exhibit phylogenetic signal

on the *Wolbachia* phylogeny (Table S6, Figure S4), implying that factors in the *Wolbachia*

genome do not account for observed diversity. In contrast, we found that *Wolbachia* density in

the ovaries at 25°C exhibits strong, significant phylogenetic signal on the host phylogeny ($\lambda =$

1.000, $P = 0.001$), implying that factors in the host genome may help determine observed

diversity of *Wolbachia* densities in host ovaries. In this instance, related species in the *montium*

species group tended to have lower *Wolbachia* densities in the ovaries relative to the

melanogaster subgroup (see Figure S5). Divergent *D. ananassae* also had a higher *Wolbachia*

density than all the other host species. We also found that cellular *Wolbachia* abundance in the

oocyte posterior region ($\lambda = 1.000$, $P = 0.037$) and the posterior cortex ($\lambda = 1.000$, $P = 0.036$)

exhibited a significant signature of phylogenetic signal at 20°C on the host phylogeny. This

pattern is largely driven by the relatively low abundance of *Wolbachia* in the posterior oocytes of

the two closely related species *D. triauraria* and *auraria*.

DISCUSSION

Maternal transmission rates are a key determinant of *Wolbachia* prevalence in host

populations[17,55]. For many *Wolbachia* strains and host species, we found that maternal

transmission is generally high and stable across two ecologically relevant temperatures[21,70]. Two

Wolbachia strains (*w*Aura and B-group *w*Mau) exhibited minor temperature-dependent changes

in transmission, but *w*Mel stood out with a relatively large decrease in transmission as

temperature dropped from 25° to 20°C (Figure 2). Maternal transmission is generally expected to

depend on *Wolbachia* density in host reproductive tissue[20,34,43,71], and specifically within

developing oocytes[14,15,21,36,59–63]. Our analyses uncovered considerable, significant variation in

Wolbachia densities at the tissue level and cellular abundances in oocytes, which also often

varied depending on temperature. With the exception of *w*Mel, *Wolbachia* densities in ovary

tissue generally did not predict maternal transmission rates (Figures 3, S1). In stage 10 oocytes,

declines in the cellular abundance of *w*Mel and *w*Aura at the posterior region coincided with

declining transmission in the cold (Figure 4), but the *w*MelCS and *w*Ri strains also experienced

changes in cellular abundance that did not correlate with differences in maternal transmission.

Together, these results suggest that maternal transmission rates cannot be accurately predicted

from measurements of *Wolbachia* quantities in ovary tissue or late-stage oocytes. First, we

explore the diverse patterns of *Wolbachia* localization in relation to maternal transmission in our

comparative analyses. Then, we examine why maternal transmission of some strains, particularly

*w*Mel, may be impacted by temperature.

At the tissue level, we found that *Wolbachia* densities in the ovaries generally did not

predict rates of maternal transmission, with the exception of *w*Mel. *w*Mel density in the ovaries

of individual females was tightly correlated with declining transmission rates in the cold (ρ =

0.537, Figure S1). Temperature did not alter *w*Aura density in the ovaries despite declining

*w*Aura transmission in the cold, and *w*Mau density counterintuitively declined at 20°C despite

transmission increasing. Other strains (e.g., *w*Ri and *w*Ha in *D. simulans*) experienced large

declines in *Wolbachia* ovary density at 20°C with no impact on maternal transmission.

Wolbachia densities in the remaining somatic carcass tissue also generally did not predict

declining transmission in the cold; however, *w*Mau densities in the carcasses of individual

females were strongly correlated with increased transmission in the cold (ρ = 0.549; discussed

further below). Notably, *w*Cha was exceptional in terms of its very low density in carcass tissue

relative to other strains, despite perfect transmission at both temperatures.

At the cellular level, *Wolbachia* abundance in stage 10 oocytes may help explain why

*w*Mel and *w*Aura experienced declining transmission in the cold. Many strains experienced

declines in cellular *Wolbachia* abundance throughout the whole oocyte at 20°C (Figure 4A), but

this was generally unrelated to transmission rates. *w*Mel was one of the few strains that had a

significant reduction in cellular abundance at the posterior region (Figure 4B) and cortex (Figure

4C), the site of germline formation. Similarly, *w*Aura had a reduced abundance in the posterior

region at 20°C (but not at the cortex). *w*MelCS (closely related to *w*Mel) was the only additional

strain with significant reductions in abundance at the posterior region and the cortex. However,

wMelCS occurred at a much higher absolute abundance than wMel, wAura, and all other

Wolbachia strains, which may explain why transmission remained high at 20°C (discussed

further below). We conjecture that there is presumably a minimum cellular abundance required

for efficient transmission, and *Wolbachia* strains like wMelCS that are generally very abundant

are less likely to fall below this threshold. A reduction in cellular abundance at the oocyte

posterior implies that fewer *Wolbachia* cells will be incorporated into the germline later during

host development, perturbing germline-to-germline maternal transmission[21]. The fact that both

wMel and wAura had significantly reduced abundances in the posterior region at 20°C implies

that perturbations of cellular *Wolbachia* abundance in this region may help explain why some

Wolbachia strains have reduced transmission in the cold. wRi was the only other *Wolbachia*

strain with an altered abundance at the oocyte posterior region and cortex, but cellular abundance

increased at 20°C. This may help explain why maternal transmission of wRi remained perfect at

20°C. Interestingly, a low absolute abundance of *Wolbachia* was not necessarily associated with

declining maternal transmission. The wTri strain (closely related to wAura) occurred at a lower

posterior abundance than all other strains, despite maintaining a high rate of transmission at both

temperatures. It is important to note that our analysis of stage 10 oocytes is only a snapshot of

host development. Other unexamined stages of development, including embryogenesis, could

also be involved mediating maternal transmission. For instance, cool temperatures could impact

Wolbachia localization at the newly formed germline of late blastoderm and cellularized

embryos, which could influence maternal transmission[15,60].

Our comparative analysis highlights how wMel seems to be especially susceptible to

cooling temperatures[21], which is consistent with our previous work suggesting the strain is less

prevalent in temperate host populations due to declining transmission in the cold[21,58]. In

transinfected mosquitoes, wMel maternal transmission and densities are also more susceptible to

heat stress relative to another *Wolbachia* strain, wAlbB[25,26,33,34,43]. A number of factors could

plausibly help explain why wMel maternal transmission is especially impacted by the cool

temperature, relative to other strains. At the tissue level, wMel was the only *Wolbachia* we

examined that exhibited a significant decline in *Wolbachia* density in both the ovaries and the

carcasses at 20°C (Figure 3). Declining transmission was highly correlated with declining wMel

density in the ovaries, plus wMel stood out with a dramatic decline in *Wolbachia* density in

carcasses at 20°C compared to other strains. While *w*Mel is generally thought to follow a

germline-to-germline route of maternal transmission during oogenesis, our recent work and

others suggests that the strain may also rely on a soma-to germline mode of transmission[15,72,73].

*w*Mel *Wolbachia* are capable of cell-to-cell migration[73] and *w*Mel injected directly into the

abdomen of adult *D. melanogaster* females can migrate to and occupy the germline and follicle

stem cells[72]. The declines of *w*Mel densities throughout host tissues could potentially impact

routes of transmission originating in somatic tissue. At the cellular level, *w*Mel is also one of the

few strains (in addition to *w*Aura) to experience a decline in abundance at the posterior of stage

10 oocytes, which is expected to impact a germline-to-germline route of transmission. The

concurrent reduction in *Wolbachia* density in female tissues and oocytes may be a one-two

punch that impacts both soma- and germline-to-germline routes of *w*Mel transmission.

The juxtaposition between *w*Mel and *w*MelCS is interesting, because the two strains

diverged recently in only the last 3,000-14,000 years[74]. The *w*Mel and *w*MelCS strains we used

in our study are highly similar (0.007% third-position pairwise differences). Despite the close

relationship, *w*MelCS transmission was not impacted by temperature and the strain occurred at a

higher cellular abundance at the oocyte posterior than any of the other strains that we examined,

regardless of temperature. Some oocytes contained especially large quantities of *w*MelCS

throughout the whole oocyte and at the posterior (Figure S6), as compared to all the other

oocytes we examined (e.g., Figure S3). *w*MelCS is only found at low frequencies in a few

populations of *D. melanogaster*[74,75], which is attributed to the fact that ancestral *w*MelCS was

largely replaced by *w*Mel in global populations of *D. melanogaster* in roughly the last 50 to 100

years[76–79]. Perhaps deleterious host effects associated with a high cellular *w*MelCS abundance in

oocytes may have contributed to the global replacement of *w*MelCS by *w*Mel. To ensure

maternal transmission, *Wolbachia* cells must maintain an association with the host germline

without perturbing highly conserved processes of germline and oocyte development[15,62,63,80].

Migration of *w*Mel to the posterior germ plasm is coincident with recruitment of host factors

required from germline formation and anterior/posterior and dorsal/ventral axis formation[62] and

an excessively high *Wolbachia* abundance can disrupt dorsal/ventral axis determination[60,62,80,81].

Perhaps similar deleterious effects may help explain why the highly abundant *w*MelCS strain

was replaced by *w*Mel, despite a higher rate of transmission in the cold. Pervious works has also

shown that higher densities of *w*MelCS in *D. melanogaster* often covaries with a reduced

lifespan[75]. Of course, we cannot rule out the possibility that the host genome also contributes to

some of the differences we observed here between wMelCS and wMel[20,21]. Indeed, our

phylogenetic analyses suggest the host genome is an important determinant of *Wolbachia*

abundance in oocytes (Table S6).

The divergent B-group wMau strain was the only strain where transmission significantly

increased at 20°C (Figure 2). Transmission was high at 25°C and increased to perfect at 20°C.

Our previous work suggests the wMau strain may also be an outlier in its mode of transmission

during oogenesis relative to A-group *Wolbachia*, which tend to exhibit evidence of a germline-

to-germline route of transmission[15]. wMau occurs at relatively low abundance at the posterior

cortex of stage 10 oocytes (Figure 4), but wMau cells are frequently found within the somatically

derived follicle cells surrounding the oocytes (Figure S6)[15], which is consistent with a soma-to-

germline route of transmission. The presence of *Wolbachia* in the follicle cells is generally not

the case for A-group *Wolbachia* (with the exception of wMel, discussed above). We found that

wMau was also one of the few strains (in addition to wCha) where *Wolbachia* density in the

somatic carcass tissues significantly increased at 20°C (Figure 3), which may help explain why

transmission increased slightly at 20°C. wMau transmission was also strongly correlated with

Wolbachia densities in the carcasses of individual females. Notably, our previous analysis of two

other wMau-*D. mauritiana* genotypes using three- to five-day old females found perfect rates of

transmission at 25°C[40], unlike our results here (Figure 2). This suggests that differences in fly

age[82], the *Wolbachia* and host genomes[21], or other unknown factors may also contribute to

variation in rates of maternal transmission.

The wMau findings also may inform our previous work on the thermoregulatory behavior of

Drosophila host species. We found that *Drosophila* species carrying A-group *Wolbachia*,

including wRi and wHa, tend to prefer cooler temperatures on a thermogradient (relative to flies

without *Wolbachia*), whereas *D. mauritiana* flies carrying wMau prefer warmer temperatures[66].

Taken together, these results suggest that fly movement to cool temperatures reduces the density

of A-group *Wolbachia* in host tissues, whereas movement to warm temperatures reduces B-

group wMau density. This is consistent with the inference that flies may thermoregulate as a

behavioral response to ameliorate negative effects of infection and a high *Wolbachia* density[67,83–

86]. Movement to cooler temperatures with A-group *Wolbachia* (i.e., behavioral chill) and to

warmer temperatures with B-group wMau (behavioral fever) are both behaviors that would

reduce *Wolbachia* density in host tissues according to our results here (Figure 3). This finding

motives further work dissecting the relationships among thermoregulatory behavior, *Wolbachia*

density in host tissues, and maternal transmission.

Our phylogenomic analyses revealed that *Wolbachia* density in the ovaries at 25°C and

cellular *Wolbachia* abundance in the oocyte posterior region and cortex at 20°C all exhibit

phylogenetic signal on the host phylogeny (Table S6, Figure S5), implying that factors in the

host genome help explain the observed diversity in *Wolbachia* localization across species. For

example, interspecific variation in the host proteins that *Wolbachia* engage during oogenesis

could plausibly influence *Wolbachia* localization patterns at the posterior cortex in stage 10

oocytes[15]. Unfortunately, it is not yet known how *Wolbachia* engage host factors like kinesin

during oogenesis[63]. Other work also suggests the host genome can influence *Wolbachia*

abundance in oocytes[21] and embryos[60]. Similarly, genome-wide screens reveal that host factors

play an important role in determining cellular *Wolbachia* abundance[87,88]. The findings here

complement our previous comparative analyses that suggest the *Wolbachia* genome also

contributes to variation in *Wolbachia* localization patterns in oocytes[15] and effects on host

thermoregulatory behavior[66]. While we did not identify any traits that exhibit phylogenetic signal

on the *Wolbachia* phylogeny, it is worth noting that relatively few *Wolbachia* genomic changes

(e.g., in the ampliconic "Octomom" gene region) are required to influence density and tissue

distributions[75,89–92]. This highlights the potential for rapid changes in these traits and their effects

on transmission.

The rate of *Wolbachia* maternal transmission, in conjunction with *Wolbachia* effects on host

fitness and cytoplasmic compatibility (CI), ultimately determine *Wolbachia* prevalence in host

populations[17,55]. In this study, we uncovered limited variation in maternal transmission rates

across temperatures, despite dramatic variation in *Wolbachia* localization patterns within host

tissues. Nonetheless, a few *Wolbachia* strains, particularly *w*Mel, had variable transmission rates

that are likely to impact *Wolbachia* prevalence in natural host populations. For instance *w*Mel

causes weak CI[82,93,94] and occurs at lower frequencies globally in temperature host populations,

which can plausibly be explained by declining transmission in the cold (Figure 2)[21]. Similarly,

*w*Mau does not cause CI and occurs at intermediate frequencies in *D. mauritiana* on

Mauritius[40,95]. Our results suggest temperature-related variation in *w*Mau transmission may

perhaps help contribute to intermediate frequencies. In contrast, *Wolbachia* strains like *w*Ri[17–

[19,96], wHa[97–99], and wAna[100,101] cause strong CI and tend to occur at high equilibrium frequencies

in nature. In addition to strong CI, our results suggest that stable transmission rates may help

these strains maintain high frequencies in the face of fluctuating conditions (Figure 2). However,

this may not always be the case, as we found that maternal transmission the strong CI-causing

strain wAura[51,100] can be perturbed by temperature. Unlike CI and *Wolbachia* maternal

transmission, the role of *Wolbachia* effects on host fitness in natural *Drosophila* populations is

poorly understood, although recent work suggests *Wolbachia* block viruses in their native

Drosophila hosts[8,9,102,103].

Understanding how temperature influences *Wolbachia* maternal transmission, especially in

relation to other key factors like CI and host fitness effects, is critical for explaining global

Wolbachia prevalence in host populations. Research on this front will also inform *Wolbachia*-

based biocontrol programs, as thermally stable *Wolbachia* strains can be leveraged to improve

the efficacy of *Wolbachia*-based biocontrol applications in mosquito populations that experience

extreme environments[25,104]. In this respect, temperature and the environment are emerging as key

factors that mediate interactions between *Wolbachia*, their hosts, and pathogens.

METHODS

Fly lines

We evaluated 10 different *Wolbachia* strains that naturally occur in eight different

Drosophila species (Table S1). For two of these host species, we tested multiple *Wolbachia*-host

genotypes: wMel and wMelCS in *D. melanogaster* and wRi and wHa in *D. simulans*. With the

exception of the wMelCS-*D. melanogaster* line, all the *Wolbachia*-host genotypes were sampled

from nature to form isofemale lines, such that single gravid females were collected from the field

and placed individually into vials (see [66] for further discussion of the wMelCS genotype). For the

maternal transmission experiments explained below, we paired each *Wolbachia*-positive female

with *Wolbachia*-free males of the same host species (Table S1). Here, we used males from

naturally *Wolbachia*-free isofemale lines whenever possible; however, in some cases, we were

unable to obtain natural genotypes without *Wolbachia*. For these species, *Wolbachia*-free

genotypes were created by treating the *Wolbachia*-positive genotype with 0.03% tetracycline for

at least four generations. After the fourth generation, we used quantitative PCR (qPCR) to

confirm that flies were cleared of *Wolbachia*[66]. We then reconstituted the gut microbiome of the

tetracycline-cleared flies by rearing them on food where *Wolbachia*-positive males of the same

genotype had been fed and defecated for the prior 48 hours. Tetracycline-cleared flies were given

at least three more generations before we conducted experiments to avoid detrimental effect of

the antibiotic treatment on mitochondrial function[105].

Maternal transmission

We tested how temperature influences maternal transmission and *Wolbachia* distributions

in host tissues by rearing females of each genotype at a moderate (25°C) and low (20°C)

temperatures. Prior to experiments, the isofemale lines were reared for at least two generations in

incubators set to each temperature and a 12L:12D light cycle (Percival Model I-36LL) on a

standard food diet[66]. We estimated maternal transmission at each temperature by aspirating

individual virgin females into vials with two 3-5 day old virgin males of the *Wolbachia*-free

genotype and then allowing females to lay eggs for eight days[20,21]. We crossed *Wolbachia*-

positive females to males without *Wolbachia*, because in crosses with *Wolbachia*-positive males,

Wolbachia-free ova produced by *Wolbachia*-positive females could be lost if they are susceptible

to cytoplasmic incompatibility, potentially leading to an overestimation of maternal

transmission[18]. The males in our experiments were all collected from the *Wolbachia*-free

genotypes that were maintained at 25°C.

Newly emerged adult F1 offspring were immediately preserved in 95% ethanol and

stored at -20°C for later analysis of *Wolbachia* status. For each subline, we measured maternal

transmission by screening five female and five male F1s for *Wolbachia* using PCR. We extracted

DNA from individual F1s using 96-well plates and a "squish" extraction buffer (10 mL Tris-HCl

[1M], 0.0372 g EDTA, 0.1461 g NaCl, 90 mL dH20, followed by 150 mL Proteinase K after

autoclaving) that allows for high-throughput and cost-effective DNA extraction. We screened

each F1 for *Wolbachia* status using previously described PCR primers for the *Wolbachia* surface

protein (*wsp*) and a second set of primers for the arthropod-specific 28S rDNA that served as a

positive control[20,40,106].

All statistical analyses were performed in R[107]. We computed maternal transmission for

each genotype at each temperature as the mean proportion of *Wolbachia*-positive offspring

produced by *Wolbachia*-positive mothers in each subline. We then estimated 95% bias-corrected

and accelerated bootstrap (BC$_a$) confidence intervals using the "boot.ci" function and 5000

iterations in the *boot* package in R[108]. We analyzed whether maternal transmission rates vary by

Wolbachia-host genotype (e.g., *w*Mel-*D. melanogaster*) and temperature using a generalized

linear mixed model (GLMM) logistic regression with the "glmer" function in the *lme4* R

package[109]. We treated the *Wolbachia* status of F1s as the dependent variable and then included

sex, the *Wolbachia*-host genotype, temperature, and a two-way interaction between the

Wolbachia-host genotype and temperature as independent variables. We included sex in the

model because we previously found that *w*Mel-like *Wolbachia* (*w*Mel and *w*Yak) are transmitted

more efficiently to male offspring[20,21]. Finally, we included the subline of each F1 as a random

effect. We assessed significance of the fixed effects with a likelihood ratio test (LRT) and type

III sum of squares using the "mixed" function in the *afex* package[110]. Lastly, we used Wilcoxon

rank-sum tests to evaluate whether transmission rates differ between 25° and 20°C for each

strain.

Wolbachia density in host ovaries

After each female was allowed to lay for eight days, we immediately dissected out her

ovaries in chilled 1X PBS. Both the ovaries and remaining carcass were then frozen at -80°C.

We then used qPCR to quantify *Wolbachia* density in the dissected ovaries and carcasses of each

individual female. DNA was first extracted using a DNeasy Blood and Tissue Kit (Qiagen) and

then amplified the *Wolbachia*-specific locus *ftsZ* and the *Drosophila*-specific locus *nAcRalpha-*

34E. Preliminary analyses indicated that separate primers were needed for *Wolbachia* in the

divergent A-Group (F: 5'-ATCCTTAACTGCGGCTCTTG-3', R: 5'-

TTCATCACAGCAGGAATGGG-3') and B-Group lineages (F: 5'-

CAGAGAAGCAAGAGCGGTAG-3', R: 5'-TCTTCAAGTCCAAGCTCTGC -3'). We were

able to use a single set of primers for all the *Drosophila* hosts (F: 5'-

CTATGGTCGTTGACAGACT-3', R: 5'-GTAGTACAGCTATTG TGGC-3'). We generated

efficiency curves to confirm that each primer pair amplified with adequate efficiency for each

Wolbachia strain and host species. All qPCR reactions were amplified using the following

cycling conditions: 50°C for 2 min, 95°C for 2 min, and then 40 cycles, with one cycle

consisting of 95°C for 15 s, 58°C for 15 s, and 72°C for 1 min. We used the average cycle

threshold (Ct) value of three technical replicates for each sample. We then estimated relative

Wolbachia density as $2^{\Delta Ct}$, where $\Delta Ct = Ct_{nAcRalpha-34E} - Ct_{ftsZ}$[111]. We used a two-way

ANOVA and type III sums of squares to test whether log-transformed *Wolbachia* density varied

among *Wolbachia*-host genotypes and temperature for both the ovaries and the carcasses. We

also used Wilcoxon rank-sum tests to test whether densities differed between 25° and 20°C for

each tissue and *Wolbachia*-host genotype.

Cellular Wolbachia abundance in host oocytes

492   We used previously described protocols and confocal imaging to quantify cellular

Wolbachia in host oocytes at each temperature[15,21]. Briefly, we paired 20 newly emerged virgin

Wolbachia-positive females *en masse* with ten 3-5 day old virgin *Wolbachia*-negative males in

individual vials for eight days to stimulate oocyte production. Females were then separated using

CO_2 and ovaries were dissected in a chilled dish of 1X PBS. Ovaries were fixed in 200 μl of

devitellinizing solution (2% paraformaldehyde and 0.5% v/v NP40 in 1X PBS) mixed with 600

μl of heptane for 20 min on a shaker at room temperature. We then removed the organic layer

using a brief centrifugation and washed the ovaries three times with PBS-T (0.1% Triton X-100

in 1X PBS), followed by three additional five min washes. Samples were treated with RNAse A

(10 mg/ml) and left overnight at room temperature. The samples were then washed in PBS-T

multiple times over two hours and then stained with a dilute solution of Alexa 488 phalloidin on

a shaker for two hours for actin staining. Subsequently, the samples were washed again multiple

times over the course of two more hour. Finally, the wash solution was removed and we added

60 μl of propidium iodide (PI) in mounting media (VECTASHIELD) and left again overnight.

Ovaries were then mounted and carefully separated out again for ease of imaging. Slides were

sealed with a coat of nail polish and stored at -20°C until imaging.

508   We used a Zeiss Olympia confocal microscope and the 40x Plan-Apochromat oil

objective for image acquisition. Alexa 488 was imaged using the 488 nm laser line and emissions

were collected from 493 to 584 nm. Propidium iodide was imaged using the 488 and 561 nm

lines and emissions were collected from 584 to 718 nm. Stage 10 oocytes were identified based

on two criterion. First, the egg chamber had to have two compartments, one with nurse cells and

the other with the developing oocyte. Second, the oocyte had to account for approximately 50%

of the egg chamber. The top and bottom of the oocyte were identified by scanning through the

egg chamber and then a 0.8 μm slice was taken around the midplane. The pixel dwell time was

8.19 μsec.

Image analysis was conducted in the program Fiji[112] following Russell *et al.*[63]. We first

manually adjusted the contrast of each image to increase the threshold so that only white puncta

corresponding to *Wolbachia* cells inside the oocyte were retained and all background noise was

rendered black. We then used the polygon selection tool to select three different regions of the

oocyte (see Figure S1 in Russell *et al.*): the whole oocyte, the posterior region, and the posterior

cortex. We defined the posterior region as the posterior $1/8^{th}$ of the oocyte and the posterior

cortex as the narrow cortical region where the pole plasm component Vasa is expressed[15,63]. The

"area" and "integrated density" were measured for each region. Additionally, the average of a

quadruplicate measure of "mean gray value" in background selections was measured to account

for any remaining background fluorescence. The corrected total cell fluorescence (CTCF) was

calculated as integrated density – (area x background mean gray value) for each region. We used

two-way ANOVAs and type III sums of squares to test whether log-transformed CTCF values

differed among *Wolbachia*-host genotypes and temperature in each region of the oocyte (whole

oocyte, posterior region, and posterior cortex). We also used Wilcoxon rank-sum tests to test

whether CTCF values differed between 25° and 20°C for each oocyte regions and *Wolbachia*-

host genotype.

Phylogenomic analyses

We conducted phylogenomic analyses to characterize the evolutionary relationships

among *Wolbachia* strains and *Drosophila* host species. We obtained *Wolbachia* sequences from

publicly available genome assemblies for wMel[21], wMelCS[66], wSeg, wCha[15], wRi[113], wAna[114],

wAura, wTri[51], wHa[115], and wMau[40]. We used Prokka 1.11 to identify homologs to known

bacterial genes in the assemblies[116]. We avoided pseudogenes and paralogs in our phylogenetic

analyses by selecting only single copy genes that uniquely matched a bacterial reference gene

identified by Prokka. We removed loci with indels by requiring all homologs to have identical

length in all *Wolbachia* genomes. A total of 170 full length single copy genes met these criteria,

totaling 135,105 bp. We then estimated a Bayesian phylogram using RevBayes 1.0.8 under the

GTR + Γ + I model partitioned by codon position[117]. Four independent runs were performed,

which all converged on the same topology. All nodes were supported with Bayesian posterior

probabilities >0.99.

We used similar methods to generate a phylogram for the *Drosophila* host species.

Publicly available sequences were obtained for *D. melanogaster*[118], *D. simulans*[119], *D.*

mauritiana[40], *D. seguyi, D. chauvacae*[120], *D. auraria, D. triauraria*[51], and *D. ananassae*[121]. A

host phylogeny was generated using the same nuclear genes implemented in Turelli *et al.*[51]:

aconitase, aldolase, bicoid, ebony, enolase, esc, g6pdh, glyp, glys, ninaE, pepck, pgi, pgm, pic,

ptc, tpi, transaldolase, white, wingless, and *yellow*. We used BLAST with the *D. melanogaster*

coding sequences to extract orthologs from the genomes of each host species. Sequences were

then aligned with MAFFT 7[122]. Finally, we used RevBayes and the GTR + Γ + I model

partitioned by codon position and gene to accommodate potential variation in the substitution

process among genes. All nodes were supported with Bayesian posterior probabilities of 1.

The resulting phylograms were used to test whether *Wolbachia* density and oocyte

cellular abundance exhibit phylogenetic signal on either the *Wolbachia* or host phylogenies. For

the host phylogeny, we pooled estimates of *Wolbachia* density and oocyte abundance by host

species, because two of the hosts (*D. melanogaster* and *D. simulans*) had data for multiple

genotypes carrying different *Wolbachia* strains. We used our estimates of *Wolbachia* density

($2^{\Delta Ct}$) and cellular oocyte abundance (log-transformed CTCF) to test for phylogenetic signal

using Pagel's lambda (λ)[68]. A Pagel's λ of 0 indicates that character evolution occurs

independently of phylogenetic relationships, whereas λ = 1 is consistent with a Brownian motion

model of character evolution. We used the "phylosig" function and a likelihood ratio test (LRT)

in the *phytools* package[123] to compare the fitted value to a model assuming no phylogenetic

signal (λ = 0). We also used a Monte Carlo-based method to generate 95% confidence intervals

surrounding our estimates using 1,000 bootstrap replicates in the *pmc* package[124].

ACKNOWLEDGEMENTS

We are especially thankful to W. Sullivan for very useful discussions that improved the

quality of this manuscript. We thank W. Conner for assistance with bioinformatic analyses. J.

Statz provided thoughtful feedback on the manuscript. We thank the Genomics Core, the

Environmental Control for Organismal Research (ECOR) facility, and the Center for

Biomolecular Structure and Dynamics at the University of Montana. This study was funded

by National Science Foundation CAREER (2145195) and National Institute of General Medical

Sciences of the National Institutes of Health MIRA (R35GM124701) awards to B.S.C.

AUTHOR CONTRIBUTIONS

M.T.J.H. designed the project, collected data, performed analyses, and prepared the manuscript. T.B.W. collected data. B.S.C. acquired funding, designed the project, prepared the manuscript, and provided leadership. All authors reviewed and edited the manuscript prior to submission.

COMEPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY

All newly generated data will be made publicly available upon publication.

    FIGURES

Figure 1. Divergent *Wolbachia* strains found in *Drosophila* host species. (A) Estimated

Bayesian phylogram of the 10 A- and B-group *Wolbachia* strains included in the study. The

phylogram was estimated using 170 single-copy genes of identical length in all genomes,

spanning 135,105 bp. **(B)** Estimated Bayesian phylogram of the eight *Drosophila* species using

20 single-copy genes. All nodes on both trees are supported with Bayesian posterior probabilities

of 1. Estimates of *Wolbachia* and *Drosophila* divergence reported in millions of years ago

(MYA) are reproduced from Meany *et al.*[40] and Suvorov *et al.*[125], respectively. The *Wolbachia*

and host trees are generally discordant, as expected with frequent *Wolbachia* host

switching[47,48,51,53,54].

Figure 2. Maternal transmission is high and stable across temperatures for many

***Wolbachia* strains.** Mean maternal transmission rates (±BC$_a$ confidence intervals) of different

Wolbachia strains in their naturally associated host species. Asterisks indicate the rates of

maternal transmission differ between 25° and 20°C for a given *Wolbachia* strain according to a

Wilcoxon rank-sum test at $P < 0.05$. Below, the cladogram depicts evolutionary relationships

among *Wolbachia* strains.

Figure 3. *Wolbachia* densities in host tissues vary by *Wolbachia*-host system and

temperature. *Wolbachia* densities are shown for dissected female ovaries and the remaining

somatic carcass tissue. Asterisks indicate the *Wolbachia* density differs between 25° and 20°C

for a given *Wolbachia* strain according to a Wilcoxon rank-sum test at *P* < 0.05.

Figure 4. Cellular *Wolbachia* abundance in late-stage oocytes generally does not align with maternal transmission rates.

Cellular *Wolbachia* abundance in stage 10 oocytes (measured as fluorescence due to propidium iodide; CTCF), measured in **(A)** the

whole oocyte, **(B)** the posterior region, and **(C)** the posterior cortex. Asterisks indicate that *Wolbachia* abundance differs between 25°

and 20°C for a given *Wolbachia* strain according to a Wilcoxon rank-sum test at $P < 0.05$. **(D)** Representative confocal images of

Wolbachia strains that have decreased abundance at the posterior cortex in the cold (*w*Mel), increased abundance at the cortex in the

cold (*w*Ri), and no change in the cold (*w*Cha). Confocal micrographs are DNA-stained with PI (red) and actin-stained with phalloidin

(green). The second column depicts a single channel image of the PI stain and the third column depicts an enlarged PI-stained image

at the posterior cortex of each oocyte. Scale bars are set to 25 µm.

REFERENCES

1. Sudakaran, S., Kost, C. & Kaltenpoth, M. Symbiont acquisition and replacement as a source of ecological innovation. *Trends Microbiol.* **25**, 375–390 (2017).

2. McCutcheon, J. P., Boyd, B. M. & Dale, C. The life of an insect endosymbiont from the cradle to the grave. *Curr. Biol.* **29**, R485–R495 (2019).

3. McCutcheon, J. P. The genomics and cell biology of host-beneficial intracellular infections. *Annu. Rev. Cell Dev. Biol.* **37**, 115–142 (2021).

4. Werren, J. H., Baldo, L. & Clark, M. E. *Wolbachia:* master manipulators of invertebrate biology. *Nat. Rev. Microbiol.* **6**, 741–751 (2008).

5. Shropshire, J. D., Leigh, B. & Bordenstein, S. R. Symbiont-mediated cytoplasmic incompatibility: what have we learned in 50 years? *Elife* **9**, e61989 (2020).

6. Oliver, K. M., Russell, J. A., Moran, N. A. & Hunter, M. S. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proc. Natl. Acad. Sci.* **100**, 1803–1807 (2003).

7. Jaenike, J., Unckless, R., Cockburn, S. N., Boelio, L. M. & Perlman, S. J. Adaptation via symbiosis: recent spread of a *Drosophila* defensive symbiont. *Science* **329**, 212–215 (2010).

8. Cogni, R., Ding, S. D., Pimentel, A. C., Day, J. P. & Jiggins, F. M. *Wolbachia* reduces virus infection in a natural population of *Drosophila*. *Commun. Biol.* **4**, 1–7 (2021).

9. Bruner-Montero, G. & Jiggins, F. M. *Wolbachia* protects *Drosophila melanogaster* against two naturally occurring and virulent viral pathogens. *Sci. Rep.* **13**, 8518 (2023).

10. Baumann, P. Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu. Rev. Microbiol.* **59**, 155–189 (2005).

11. Moran, N. A., McCutcheon, J. P. & Nakabachi, A. Genomics and evolution of heritable bacterial symbionts. *Annu. Rev. Genet.* **42**, 165–190 (2008).

12. Douglas, A. E. The microbial dimension in insect nutritional ecology. *Funct. Ecol.* **23**, 38–47 (2009).

13. Fisher, R. M., Henry, L. M., Cornwallis, C. K., Kiers, E. T. & West, S. A. The evolution of host-symbiont dependence. *Nat. Commun.* **8**, 1–8 (2017).

14. Russell, S. L., Chappell, L. & Sullivan, W. A symbiont's guide to the germline. *Curr. Top. Dev. Biol.* **135**, 315–351 (2019).

15. Radousky, Y. A. *et al.* Distinct *Wolbachia* localization patterns in oocytes of diverse host species reveal multiple strategies of maternal transmission. *Genetics* **224**, (2023).

16. Dunbar, H. E., Wilson, A. C., Ferguson, N. R. & Moran, N. A. Aphid thermal tolerance is governed by a point mutation in bacterial symbionts. *PLoS Biol.* **5**, (2007).

17. Hoffmann, A. A., Turelli, M. & Harshman, L. G. Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126**, 933–948 (1990).

664   18. Turelli, M. & Hoffmann, A. A. Cytoplasmic incompatibility in *Drosophila simulans*:
665   dynamics and parameter estimates from natural populations. *Genetics* **140**, 1319–1338
666   (1995).

667   19. Carrington, L. B., Lipkowitz, J. R., Hoffmann, A. A. & Turelli, M. A re-examination of
668   *Wolbachia*-induced cytoplasmic incompatibility in California *Drosophila simulans*. *PLoS*
669   *One* **6**, e22565 (2011).

670   20. Hague, M. T., Mavengere, H., Matute, D. R. & Cooper, B. S. Environmental and genetic
671   contributions to imperfect *w*Mel-like *Wolbachia* transmission and frequency variation.
672   *Genetics* **215**, 1117–1132 (2020).

673   21. Hague, M. T. J. *et al.* Temperature effects on cellular host-microbe interactions explain
674   continent-wide endosymbiont prevalence. *Curr. Biol.* **32**, 878-888.e8 (2022).

675   22. Wade, M. J. The co-evolutionary genetics of ecological communities. *Nat. Rev. Genet.* **8**,
676   185–195 (2007).

677   23. Weeks, A. R., Turelli, M., Harcombe, W. R., Reynolds, K. T. & Hoffmann, A. A. From
678   parasite to mutualist: Rapid evolution of *Wolbachia* in natural populations of *Drosophila*.
679   *PLoS Biol.* **5**, e114 (2007).

680   24. Ross, P. A., Turelli, M. & Hoffmann, A. A. Evolutionary ecology of *Wolbachia* releases for
681   disease control. *Annu. Rev. Genet.* **53**, (2019).

682   25. Ross, P. A. *et al.* Developing *Wolbachia*-based disease interventions for an extreme
683   environment. *PLoS Pathog.* **19**, e1011117 (2023).

684   26. Hien, N. T. *et al.* Environmental factors influence the local establishment of *Wolbachia* in
685   *Aedes aegypti* mosquitoes in two small communities in central Vietnam. *Gates Open Res.* **5**,
686   (2021).

687   27. Nikolouli, K. *et al.* Sterile insect technique and *Wolbachia* symbiosis as potential tools for
688   the control of the invasive species *Drosophila suzukii*. *J. Pest Sci.* **91**, 489–503 (2018).

689   28. Gong, J.-T. *et al.* Stable introduction of plant-virus-inhibiting *Wolbachia* into planthoppers
690   for rice protection. *Curr. Biol.* **30**, 4837–4845 (2020).

691   29. McMeniman, C. J. *et al.* Stable introduction of a life-shortening *Wolbachia* infection into
692   the mosquito *Aedes aegypti*. *Science* **323**, 141–144 (2009).

693   30. Hoffmann, A. A. *et al.* Successful establishment of *Wolbachia* in *Aedes* populations to
694   suppress dengue transmission. *Nature* **476**, 454–457 (2011).

695   31. Walker, T. *et al.* The *w*Mel *Wolbachia* strain blocks dengue and invades caged *Aedes*
696   *aegypti* populations. *Nature* **476**, 450–453 (2011).

697   32. Utarini, A. *et al.* Efficacy of *Wolbachia*-infected mosquito deployments for the control of
698   dengue. *N. Engl. J. Med.* **384**, 2177–2186 (2021).

699   33. Ross, P. A. *et al. Wolbachia* infections in *Aedes aegypti* differ markedly in their response to
700   cyclical heat stress. *PLOS Pathog.* **13**, e1006006 (2017).

701   34. Ross, P. A., Ritchie, S. A., Axford, J. K. & Hoffmann, A. A. Loss of cytoplasmic
702   incompatibility in *Wolbachia*-infected *Aedes aegypti* under field conditions. *PLoS Negl.*
703   *Trop. Dis.* **13**, e0007357 (2019).

35. Funkhouser-Jones, L. J., van Opstal, E. J., Sharma, A. & Bordenstein, S. R. The maternal
effect gene Wds controls *Wolbachia* titer in *Nasonia*. *Curr. Biol.* **28**, 1692–1702 (2018).

36. Porter, J. & Sullivan, W. The cellular lives of *Wolbachia*. *Nat. Rev. Microbiol.* 1–17 (2023).

37. Bright, M. & Bulgheresi, S. A complex journey: transmission of microbial symbionts. *Nat.*
Rev. Microbiol. **8**, 218–230 (2010).

38. Landmann, F. *et al.* Both asymmetric mitotic segregation and cell-to-cell invasion are
required for stable germline transmission of *Wolbachia* in filarial nematodes. *Biol. Open* **1**,
536–547 (2012).

39. Correa, C. C. & Ballard, J. *Wolbachia* associations with insects: winning or losing against a
master manipulator. *Front. Ecol. Evol.* **3**, 153 (2016).

40. Meany, M. K. *et al.* Loss of cytoplasmic incompatibility and minimal fecundity effects
explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*. *Evolution* **73**,
1278–1295 (2019).

41. Olsen, K., Reynolds, K. T. & Hoffmann, A. A. A field cage test of the effects of the
endosymbiont *Wolbachia* on *Drosophila melanogaster*. *Heredity* **86**, 731–737 (2001).

42. Hamm, C. A. *et al. Wolbachia* do not live by reproductive manipulation alone: Infection
polymorphism in *Drosophila suzukii* and *D. subpulchrella*. *Mol. Ecol.* **23**, 4871–4885
(2014).

43. Ross, P. A. *et al.* Heatwaves cause fluctuations in *w*Mel *Wolbachia* densities and
frequencies in *Aedes aegypti*. *PLoS Negl. Trop. Dis.* **14**, e0007958 (2020).

44. Ferri, E. *et al.* New insights into the evolution of *Wolbachia* infections in filarial nematodes
inferred from a large range of screened species. *PLoS One* **6**, e20843 (2011).

45. Zug, R. & Hammerstein, P. Still a host of hosts for *Wolbachia*: Analysis of recent data
suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7**, e38544 (2012).

46. Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z. & Welch, J. J. The incidence of bacterial
endosymbionts in terrestrial arthropods. *Proc. R. Soc. B Biol. Sci.* **282**, 20150249 (2015).

47. O'Neill, S. L., Giordano, R., Colbert, A. M., Karr, T. L. & Robertson, H. M. 16S rRNA
phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic
incompatibility in insects. *Proc. Natl. Acad. Sci.* **89**, 2699–2702 (1992).

48. Raychoudhury, R., Baldo, L., Oliveira, D. C. & Werren, J. H. Modes of acquisition of
Wolbachia: Horizontal transfer, hybrid introgression, and codivergence in the *Nasonia*
species complex. *Evolution* **63**, 165–183 (2009).

49. Conner, W. R. *et al.* Genome comparisons indicate recent transfer of *w*Ri-like *Wolbachia*
between sister species *Drosophila suzukii* and *D. subpulchrella*. *Ecol. Evol.* **7**, 9391–9404
(2017).

50. Gerth, M. & Bleidorn, C. Comparative genomics provides a timeframe for *Wolbachia*
evolution and exposes a recent biotin synthesis operon transfer. *Nat. Microbiol.* **2**, 16241
(2017).

51. Turelli, M. *et al.* Rapid global spread of *w*Ri-like *Wolbachia* across multiple *Drosophila*.
Curr. Biol. **28**, 963-971.e8 (2018).

52. Cooper, B. S., Vanderpool, D., Conner, W. R., Matute, D. R. & Turelli, M. *Wolbachia*
acquisition by *Drosophila yakuba*-clade hosts and transfer of incompatibility loci between
distantly related *Wolbachia*. *Genetics* **212**, 1399–1419 (2019).

53. Shropshire, D. *et al.* Rapid turnover of pathogen-blocking *Wolbachia* and their
incompatibility loci. *bioRxiv* 2023–12 (2023).

54. Vancaester, E. & Blaxter, M. Phylogenomic analysis of *Wolbachia* genomes from the
Darwin Tree of Life biodiversity genomics project. *PLoS Biol.* **21**, e3001972 (2023).

55. Caspari, E. & Watson, G. On the evolutionary importance of cytoplasmic sterility in
mosquitoes. *Evolution* **13**, 568–570 (1959).

56. Turelli, M., Katznelson, A. & Ginsberg, P. S. Why *Wolbachia*-induced cytoplasmic
incompatibility is so common. *Proc. Natl. Acad. Sci.* **119**, e2211637119 (2022).

57. Turelli, M. & Hoffmann, A. A. Rapid spread of an inherited incompatibility factor in
California *Drosophila*. *Nature* **353**, 440–442 (1991).

58. Kriesner, P., Conner, W. R., Weeks, A. R., Turelli, M. & Hoffmann, A. A. Persistence of a
Wolbachia infection frequency cline in *Drosophila melanogaster* and the possible role of
reproductive dormancy. *Evolution* **70**, 979–997 (2016).

59. Hadfield, S. J. & Axton, J. M. Germ cells colonized by endosymbiotic bacteria. *Nature* **402**,
482 (1999).

60. Veneti, Z., Clark, M. E., Karr, T. L., Savakis, C. & Bourtzis, K. Heads or tails: host-parasite
interactions in the *Drosophila-Wolbachia* system. *Appl. Environ. Microbiol.* **70**, 5366–5372
(2004).

61. Ferree, P. M. *et al. Wolbachia* utilizes host microtubules and Dynein for anterior
localization in the *Drosophila* oocyte. *PLoS Pathog.* **1**, e14 (2005).

62. Serbus, L. R. & Sullivan, W. A cellular basis for *Wolbachia* recruitment to the host
germline. *PLoS Pathog.* **3**, e190 (2007).

63. Russell, S. L., Lemseffer, N. & Sullivan, W. T. *Wolbachia* and host germline components
compete for kinesin-mediated transport to the posterior pole of the *Drosophila* oocyte.
PLoS Pathog. **14**, e1007216 (2018).

64. Prout, T. Some evolutionary possibilities for a microbe that causes incompatibility in its
host. *Evolution* **48**, 909–911 (1994).

65. Turelli, M. Evolution of incompatibility-inducing microbes and their hosts. *Evolution* **48**,
1500–1513 (1994).

66. Hague, M. T., Caldwell, C. N. & Cooper, B. S. Pervasive effects of *Wolbachia* on host
temperature preference. *mBio* **11**, e01768-20 (2020).

67. Hague, M. T. J., Woods, H. A. & Cooper, B. S. Pervasive effects of *Wolbachia* on host
activity. *Biol. Lett.* **17**, 20210052 (2021).

68. Pagel, M. Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884
(1999).

69. Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative data: a
test and review of evidence. *Am. Nat.* **160**, 712–726 (2002).

70. Rajpurohit, S. & Schmidt, P. S. Measuring thermal behavior in smaller insects: A case
study in *Drosophila melanogaster* demonstrates effects of sex, geographic origin, and
rearing temperature on adult behavior. *Fly (Austin)* **10**, 149–161 (2016).

71. Funkhouser-Jones, L. J., van Opstal, E. J., Sharma, A. & Bordenstein, S. R. The maternal
effect gene Wds controls *Wolbachia* titer in *Nasonia. Curr. Biol.* **28**, 1692–1702 (2018).

72. Frydman, H. M., Li, J. M., Robson, D. N. & Wieschaus, E. Somatic stem cell niche tropism
in *Wolbachia. Nature* **441**, 509 (2006).

73. White, P. M. *et al.* Mechanisms of horizontal cell-to-cell transfer of *Wolbachia* spp. in
Drosophila melanogaster. Appl Env. Microbiol **83**, e03425-16 (2017).

74. Richardson, M. F. *et al.* Population genomics of the *Wolbachia* endosymbiont in
Drosophila melanogaster. PLoS Genet. **8**, e1003129 (2012).

75. Chrostek, E. *et al. Wolbachia* variants induce differential protection to viruses in
Drosophila melanogaster: a phenotypic and phylogenomic analysis. *PLoS Genet.* **9**,
e1003896 (2013).

76. Riegler, M., Sidhu, M., Miller, W. J. & O'Neill, S. L. Evidence for a global *Wolbachia*
replacement in *Drosophila melanogaster. Curr. Biol.* **15**, 1428–1433 (2005).

77. Ilinsky, Y. Coevolution of *Drosophila melanogaster* mtDNA and *Wolbachia* genotypes.
PLoS One **8**, e54373 (2013).

78. Early, A. M. & Clark, A. G. Monophyly of *Wolbachia pipientis* genomes within *Drosophila*
melanogaster: Geographic structuring, titre variation and host effects across five
populations. *Mol. Ecol.* **22**, 5765–5778 (2013).

79. Strunov, A. *et al.* Historic museum samples provide evidence for a recent replacement of
Wolbachia types in European *Drosophila melanogaster. Mol. Biol. Evol.* **40**, msad258
(2023).

80. Serbus, L. R. *et al.* A feedback loop between *Wolbachia* and the *Drosophila* gurken mRNP
complex influences *Wolbachia* titer. *J. Cell Sci.* **124**, 4299–4308 (2011).

81. Poinsot, D., Bourtzis, K., Markakis, G., Savakis, C. & Merçot, H. Wolbachia transfer from
Drosophila melanogaster into *D. simulans*: Host effect and cytoplasmic incompatibility
relationships. *Genetics* **150**, 227–237 (1998).

82. Shropshire, J. D., Hamant, E. & Cooper, B. S. Male age and *Wolbachia* dynamics:
investigating how fast and why bacterial densities and cytoplasmic incompatibility
strengths vary. *mBio* **12**, e02998-21 (2021).

83. Hart, B. L. Biological basis of the behavior of sick animals. *Neurosci. Biobehav. Rev.* **12**,
123–137 (1988).

84. De Roode, J. C. & Lefèvre, T. Behavioral immunity in insects. *Insects* **3**, 789–820 (2012).

85. Curtis, V. A. Infection-avoidance behaviour in humans and other animals. *Trends Immunol.*
35, 457–464 (2014).

86. Fedorka, K. M., Kutch, I. C., Collins, L. & Musto, E. Cold temperature preference in
bacterially infected *Drosophila melanogaster* improves survival but is remarkably
suboptimal. *J. Insect Physiol.* **93**, 36–41 (2016).

87. White, P. M. *et al.* Reliance of *Wolbachia* on high rates of host proteolysis revealed by a
genome-wide RNAi screen of *Drosophila* cells. *Genetics* **205**, 1473–1488 (2017).

88. Grobler, Y. *et al.* Whole genome screen reveals a novel relationship between *Wolbachia*
levels and *Drosophila* host translation. *PLoS Pathog.* **14**, e1007445 (2018).

89. Chrostek, E. & Teixeira, L. Mutualism breakdown by amplification of *Wolbachia* genes.
PLoS Biol. **13**, e1002065 (2015).

90. Chrostek, E. & Teixeira, L. Within host selection for faster replicating bacterial symbionts.
PLoS One **13**, e0191530 (2018).

91. Duarte, E. H., Carvalho, A., López-Madrigal, S., Costa, J. & Teixeira, L. Forward genetics
in *Wolbachia*: Regulation of *Wolbachia* proliferation by the amplification and deletion of
an addictive genomic island. *PLOS Genet.* **17**, e1009612 (2021).

92. Woolfit, M. *et al.* Genomic evolution of the pathogenic *Wolbachia* strain, *w*MelPop.
Genome Biol. Evol. **5**, 2189–2204 (2013).

93. Hoffmann, A. A., Hercus, M. & Dagher, H. Population dynamics of the *Wolbachia*
infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148**,
221–231 (1998).

94. Reynolds, K. T. & Hoffmann, A. A. Male age, host effects and the weak expression or non-
expression of cytoplasmic incompatibility in *Drosophila* strains infected by maternally
transmitted *Wolbachia*. *Genet. Res.* **80**, 79–87 (2002).

95. Giordano, R., O'Neill, S. L. & Robertson, H. M. *Wolbachia* infections and the expression
of cytoplasmic incompatibility in *Drosophila sechellia* and *D. mauritiana*. *Genetics* **140**,
1307–1317 (1995).

96. Hoffmann, A. & Turelli, M. Unidirectional incompatibility in *Drosophila simulans*:
inheritance, geographic variation and fitness effects. *Genetics* **119**, 435–444 (1988).

97. Rousset, F. & Solignac, M. Evolution of single and double *Wolbachia* symbioses during
speciation in the *Drosophila simulans* complex. *Proc. Natl. Acad. Sci.* **92**, 6389–6393
(1995).

98. James, A. C. & Ballard, J. W. O. Expression of cytoplasmic incompatibility in *Drosophila*
simulans and its impact on infection frequencies and distribution of *Wolbachia pipientis*.
Evolution **54**, 1661–1672 (2000).

99. Ballard, J. W. O. Sequential evolution of a symbiont inferred from the host: *Wolbachia* and
Drosophila simulans. *Mol. Biol. Evol.* **21**, 428–442 (2004).

100. Bourtzis, K., Nirgianaki, A., Markakis, G. & Savakis, C. *Wolbachia* infection and
cytoplasmic incompatibility in *Drosophila* species. *Genetics* **144**, 1063–1073 (1996).

101. Choi, J. Y., Bubnell, J. E. & Aquadro, C. F. Population genomics of infectious and
integrated *Wolbachia pipientis* genomes in *Drosophila ananassae*. *Genome Biol. Evol.* **7**,
2362–2382 (2015).

102. Osborne, S. E., Leong, Y. S., O'Neill, S. L. & Johnson, K. N. Variation in antiviral protection mediated by different *Wolbachia* strains in *Drosophila simulans*. *PLOS Pathog.* **5**, e1000656 (2009).

103. Martinez, J. *et al.* Symbionts commonly provide broad spectrum resistance to viruses in insects: A comparative analysis of *Wolbachia* strains. *PLOS Pathog.* **10**, e1004369 (2014).

104. Gu, X. *et al.* A *w*Mel *Wolbachia* variant in *Aedes aegypti* from field-collected *Drosophila melanogaster* with increased phenotypic stability under heat stress. *Environ. Microbiol.* **24**, 2119–2135 (2022).

105. Ballard, J. & Melvin, R. Tetracycline treatment influences mitochondrial metabolism and mtDNA density two generations after treatment in *Drosophila*. *Insect Mol. Biol.* **16**, 799–802 (2007).

106. Cooper, B. S., Ginsberg, P. S., Turelli, M. & Matute, D. R. *Wolbachia* in the *Drosophila yakuba* complex: Pervasive frequency variation and weak cytoplasmic incompatibility, but no apparent effect on reproductive isolation. *Genetics* **205**, 333–351 (2017).

107. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for Statistical Computing, Vienna, Austria, 2023).

108. Canty, C. & Ripley, B. boot: Bootstrap R (S-Plus) functions. (2017).

109. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

110. Singmann, H., Bolker, B., Westfall, J., Aust, F. & Ben-Shachar, M. S. *Afex: Analysis of Factorial Experiments*. (2020).

111. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT–PCR. *Nucleic Acids Res.* **29**, e45–e45 (2001).

112. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675 (2012).

113. Klasson, L. *et al.* The mosaic genome structure of the *Wolbachia w*Ri strain infecting *Drosophila simulans*. *Proc. Natl. Acad. Sci.* **106**, 5725–5730 (2009).

114. Salzberg, S. L. *et al.* Serendipitous discovery of *Wolbachia* genomes in multiple *Drosophila* species. *Genome Biol.* **6**, R23 (2005).

115. Ellegaard, K. M., Klasson, L., Näslund, K., Bourtzis, K. & Andersson, S. G. Comparative genomics of *Wolbachia* and the bacterial species concept. *PLoS Genet.* **9**, e1003381 (2013).

116. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).

117. Höhna, S. *et al.* RevBayes: Bayesian phylogenetic inference using graphical models and an interactive model-specification language. *Syst. Biol.* **65**, 726–736 (2016).

118. Hoskins, R. A. *et al.* The Release 6 reference sequence of the Drosophila melanogaster genome. *Genome Res.* **25**, 445–458 (2015).

119. Hu, T. T., Eisen, M. B., Thornton, K. R. & Andolfatto, P. A second-generation assembly of
the Drosophila simulans genome provides new insights into patterns of lineage-specific
divergence. *Genome Res.* **23**, 89–98 (2013).

120. Conner, W. R. *et al.* A phylogeny for the *Drosophila montium* species group: A model
clade for comparative analyses. *Mol. Phylogenet. Evol.* **158**, 107061 (2021).

121. Clark, A. G. *et al.* Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature*
450, 203–218 (2007).

122. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

123. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

124. Boettiger, C., Coop, G. & Ralph, P. Is your phylogeny informative? Measuring the power
of comparative methods. *Evolution* **66**, 2240–2251 (2012).

125. Suvorov, A. *et al.* Widespread introgression across a phylogeny of 155 *Drosophila*
genomes. *Curr. Biol.* **32**, 111–123 (2022).

913