Similarities Between Antibiotic-Resistant *Escherichia coli* from Raw Meat, Commercial Raw Dog Food and Those Causing Extraintestinal Human Infections: A Contemporaneous Geographically Focussed Genomic Epidemiology Study

Jordan E. Sealey[1], Beth Astley[1], Katie L. Sealey[1], Aimee M. Daum[1], Defne Kiziltan[1], Laura Wright[1], Philip Williams[2] & Matthew B. Avison[1]

[1]School of Cellular & Molecular Medicine, University of Bristol. UK

[2]University Hospitals Bristol and Weston NHS Foundation Trust, Bristol. UK

Abstract

Objectives

There is strong evidence linking raw meat or raw dog food (RDF) feeding with dogs shedding antibiotic-resistant (ABR) *Escherichia coli*. As well as potential risks of ingesting ABR *E. coli* when handling/consuming meat, therefore, people who raw-feed pets may face additional risks. However, phylogenetic evidence closely linking *E. coli* found on raw meat/RDF with those causing opportunistic human infections is lacking, partly due to non-contemporaneous sampling. Here, we monitored ABR *E. coli* in meat/RDF in Bristol, UK in 2022/23, and compared *E. coli* isolates with *E. coli* causing human urinary tract and bloodstream infections in Bristol in 2023.

Methods

We monitored ABR *E. coli* positivity in raw chicken, beef, pork and lamb meat from 15 large-chain stores and chicken-based RDF from 15 pet stores and selected 200 ABR *E. coli* for Illumina WGS. *E. coli* causing 1182 human infections were sequenced, irrespective of resistance phenotype.

Results

Chicken/RDF were most likely to carry amoxicillin-, spectinomycin-, streptomycin-, cefotaxime-, and ciprofloxacin-resistant *E. coli*. Phylogroup A/B1 and B2 *E. coli* dominated meat/RDF and human isolates, respectively. Resistance to antibiotics used predominantly in farmed animals dominated meat/RDF isolates; resistance to antibiotics extensively used in humans dominated human isolates. Nineteen meat/RDF-infection *E. coli* pairs differed by <50 SNPs across the core-genome, with four pairs (three ST69 and one ST117) differing by <20 SNPs and carrying identical resistance genes, suggestive of recent sharing.

Conclusions

Using contemporaneous, geographically focussed sampling, we have confirmed

close phylogenetic relationships between meat/RDF and human infection-causing

ABR *E. coli*. Despite representing a small number of infections, these findings

support the consideration of tighter microbiological standards for RDF given that it is

explicitly sold to be fed uncooked. Improved risk communication to pet owners and

the inclusion of RDF within integrated ABR surveillance is warranted.

Introduction

Escherichia coli is an almost ubiquitous commensal in the gastrointestinal tracts of humans and other warm-blooded animals, and an important pathogen (1). Enteropathogenic *E. coli* causes food-borne diarrhea outbreaks, while extraintestinal pathogenic *E. coli* are leading causes of urinary tract infection (UTI) and bloodstream infection (BSI); these infections are opportunistic and predominantly arise from commensal gastrointestinal flora (2). Antibiotic resistant (ABR) *E. coli*, therefore, pose a significant health threat and are an important indicator for One Health ABR transmission (3).

Meat can become contaminated with fecal bacteria, including *E. coli,* during slaughter and processing (4). In the UK, regulations ensure that raw meat sold for consumption by people and domestic pets meet the same bacteriological limits. For commensal *E. coli*, acceptable limits may reach hundreds of bacteria per gram. Therefore, poor hygiene when handling, storing and preparing raw meat as well as consuming undercooked meat can lead to the ingestion of *E. coli*, including ABR strains (5).

Raw meat feeding of pets has increased in popularity, largely driven by owners' perceptions of enhanced pet wellbeing (6). A recent UK survey found that around 50% of owners who raw feed their dogs feed commercial raw dog food (RDF) and 50% prepare it themselves, using raw meat sold for human consumption (6). Like raw meat, RDF is frequently contaminated with ABR *E. coli*, including those resistant to the critically important third-generation cephalosporins (3GCs) or fluoroquinolones (FQs) (7-12).

We have demonstrated that dogs who are fed raw meat in the city of Bristol, UK, are

more likely to excrete 3GC-resistant (3GC-R) or FQ-resistant (FQ-R) *E. coli* (13,14).

Other studies performed in other geographical regions support our findings (15-18).

Therefore, people feeding raw meat face multiple ABR *E. coli* exposure risks:

handling meat contaminated with ABR *E. coli* and interacting with dogs which are

more likely to excrete them (7). However, there remains a scarcity of high-resolution,

contemporaneous, geographically focused studies that phylogenetically compare

ABR *E. coli* from raw meat and human clinical infections, with none involving RDF.

Here we compared 200 ABR *E. coli* from raw meat purchased from large-chain

grocery stores and commercial RDF sold in the city of Bristol, UK in 2022-23 with

1,182 *E. coli* isolates from human UTI and BSI in Bristol in 2023.

Materials and Methods

Sample collection

We sampled five locations, one per week during October and November 2022

(approximately 25,000 population in each location), within Bristol, a city in South

West England. For each location, three large chain grocery stores were selected,

and one packet each of raw minced beef, minced lamb, chicken thighs and pork

chops were purchased at each store. In total, fifteen samples per meat type were

collected, except pork (13 samples due to availability issues). Samples were kept at

4°C and processed within the "use by" date.

Five Bristol-based RDF stores were also identified; locations overlapped with the

sampled large-chain grocery stores. Three brands of chicken-based RDF were

purchased in each store during September 2023, totaling 15 RDF samples across 11

brands. As RDF is sold frozen, samples were defrosted at 4°C on the day of

purchase and processed the next day.

Microbiology

Under sterile conditions, 200 g meat was bagged (Stomacher® 400 Classic

Standard) alongside 200 mL of *E. coli* Enrichment Broth (Oxoid LTD). Sealed bags

were placed into screw-top plastic jars for incubation for 5 h at 37°C and 180 rpm.

Enriched culture (500 µL) was mixed with glycerol (25% v/v final), twenty microliters

of each mixture was spread onto Tryptone Bile X-Glucuronide agar without antibiotic,

or with amoxicillin (8 mg/L), spectinomycin (32 mg/L), streptomycin (32 mg/L),

amoxicillin/clavulanate (8/2 mg/L), cefotaxime (2 mg/L), and ciprofloxacin (0.5 mg/L).

Culture pates were incubated at 37°C for 18 h. Up to 10 blue colonies per plate were

tested against the same antibiotics to determine resistance phenotype. One isolate

per resistance phenotype per sample was selected for whole genome sequencing

(WGS).

Human infection isolates

Isolates were from blood or urine samples deduplicated by patient. They were

collected in 2023 at Severn Pathology, a diagnostic laboratory within Bristol. BSI

isolates (n=795) were collected throughout 2023 from hospitalised patients.

Community UTI isolates (n=387) were collected for one week in December 2023. All

isolates were sequenced, irrespective of antibiotic susceptibility profile.

WGS and phylogenetic analysis

WGS was performed by MicrobesNG to achieve a minimum 30-fold coverage using

standard protocols (https://www.microbesng.com/). WGS used an Illumina NovaSeq

6000 (Illumina, San Diego, USA) with a 250 bp paired end protocol. Data were

analyzed using ResFinder 4.7.2 (19), and MLST 2.0.9 (20). WGS data were

excluded due to quality control issues if >500 contigs, <50 or >51% GC or <4.5

million or >6 million nucleotides total length. Multiple genomes with the same

sequence type (ST) and antibiotic resistance gene (ARG) complement from a single

meat/RDF sample were deduplicated.

Phylogenetic analysis used the Bioconda environment. Alignments (**Table S1** shows

references) used Snippy and Snippy-Core (https://github.com/tseemann/snippy);

SNP distances were determined using SNP-dists (https://github.com/tseemann/snp-

dists). Maximum likelihood trees were produced using RAxML with the GTRGAMMA

model of rate of heterogeneity (21) and illustrated using Microreact (22).

Results

Sample-level positivity for ABR E. coli *in raw meat*

Eighty-one percent of meat from large-chain grocery stores tested positive for *E. coli*.

Positivity was highest in chicken (100%) and almost all chicken samples contained

streptomycin-, spectinomycin- and amoxicillin-resistant *E. coli*, significantly greater

(Fisher's Exact $p<0.05$) than that found in beef, lamb or pork. Notably, 47% of

chicken samples carried FQ-R *E. coli*, significantly higher than lamb and pork, but

not beef. Seven percent of chicken and 13% of beef samples contained 3GC-R *E.*

coli; no lamb or pork samples were positive (**Tables 1, S2**).

We found no statistically significant difference in ABR *E. coli* positivity among

chicken-based RDF samples, compared with grocery store chicken (Fisher's Exact

p>0.05; **Tables 1, S2, S3**).

Molecular epidemiology of ABR E. coli *from raw meat*

In total, WGS for 200 unique ABR *E. coli* isolates were retained: 90 from RDF, 66

(chicken), 17 (beef), 13 (lamb), and 14 (pork). A wide range of *E. coli* STs were

detected (**Table S4**). ST10, ST162, ST69, ST88, ST117 and ST1718 were found in

multiple meat types, but none in all types. Isolates from ST131, a common human

extraintestinal pathogenic ST, were found in one beef (serotype O16:H5 *fimH41*) and

one pork (O15:H4 *fimH22*) sample. Chicken-based RDF shared 13 STs with chicken

meat, with a statistically significant ST clustering between chicken versus chicken-

based RDF compared with all other meat sources combined (Fisher's Exact

p<0.001) (**Table S4**).

Following a core genome alignment, a difference of <20 SNPs between two core

genomes was used to define clonality, suggestive of recent transmission (23). Thirty

clones were identified of which 15 included isolates from more than one meat type

(**Table S5**). Considering all pairwise core genome alignments, pork and lamb were

significantly less likely to share clonal isolates with chicken-based RDF than chicken

(Fisher's Exact *p*=0.004) and beef (*p*=0.0001), with chicken and beef not being

significantly different in this regard (*p*=0.11). Though chicken/RDF sharing was seen

in 12 clones and beef/RDF sharing in only five (**Table S5**).

Across the 15 RDF samples, three were from the same brand, and across seven STs

we detected multiple isolate pairs <20 SNPs apart between these three samples.

Two further brands were each sampled twice, although no <20 SNP links were

observed. The remaining eight brands were each represented by a single sample.

Comparison of ABR E. coli *from raw meat sold in Bristol with* E. coli *causing human*

infections in the same region

Twenty-one STs were shared between meat and human infection isolates with 47/90,

33/60 and 20/44 isolates, respectively from RDF, chicken and red meat (beef, pork

and lamb combined) being of STs that are also seen among human clinical isolates

(**Table S6**). These proportions were not significantly different based on type of meat

(Fisher's Exact p>0.05).

When considering *E. coli* phylogroups (**Table S7**), phylogroup B2 was significantly

more common among human infection isolates than among isolates from poultry

(chicken or RDF) or red meat (Fisher's Exact p<0.00001 for each). In contrast,

phylogroups A/B1 isolates were significantly less abundant in human isolates

compared with poultry or red meat (Fisher's Exact p<0.00001 for each). Phylogroup

D was also significantly less abundant in human isolates than in poultry isolates

(Fisher's Exact p=0.012) but the difference (13.8% versus 22.4%) was much less

than for A/B1 isolates (13.5% versus 66.0%).

We identified 19 incidences where *E. coli* from meat were <50 SNPs different from a

paired isolate causing a human infection, with four of these meat/infection pairs

being <20 SNPs apart (**Table S8**). Overlaps between meat and human infection

isolates from ST162, ST88 and ST10 are illustrated in **Figures S1-S3** with STs ST69

and ST117 – having three and one <20 SNP distanced pairs, respectively – being

illustrated in **Figures 1 and 2**.

ARG Characterization of Resistant E. coli *from Meat and Human Infections*

The average number of ARGs per isolate was 5.5 in RDF, 4.7 in chicken, 5.2 in beef, 5.0 in pork, and 3.2 in lamb. Common ARGs detected across most meat sources included *aadA, aph(3")-Ib and aph(6)-Id (strAB), sul1/2, dfrA, bla*$_{TEM-1}$, *tetA/B* and *qnrS1* (**Table S9**). It was very rare to find meat isolates within one of the 30 clones defined above having identical ARG complements (**Table S5**). However, the two chicken meat/BSI ST69 pairs differing by 12 and 13 SNPs, respectively all carried *aac(3)-VIa*, *aadA1*, *sul1*, *sul2* and the beef meat/UTI ST69 pair differing by 16 SNPs both carried *strAB*, *bla*$_{TEM-1}$, *sul1*, *sul2*, *tetA*, *dfrA7*.

FQ resistance was predominately due to chromosomal mutations in meat isolates, with the most common being the triple combination of *gyrA*D87N*, gyrA*S83L *and parC*S80I. ARGs conferring 3GC resistance, were *bla*$_{CTX-M-1}$, *bla*$_{CTX-M-15}$, and *bla*$_{SHV-12}$ with one being a chromosomal *ampC* hyperproducer (**Table S9**).

In the UK, many antibiotics used in livestock are also used in human medicine, e.g. beta-lactams, trimethoprim, sulfonamides, gentamicin, tobramycin and FQs (though the last three are used very rarely in livestock). However, other antibiotics, e.g. florfenicol, neomycin, streptomycin, spectinomycin, and tetracycline are widely used in livestock while use in human medicine is low (though tetracycline derivatives are widely used). By comparing the number of ARGs relevant for each of these main groups of antibiotics in *E. coli* isolates from poultry (chicken and RDF) or red meat (beef, lamb and pork) with those from human infections, we noted that for the antibiotics widely used in human medicine there was either no significant difference in abundance between meat and human infections, or there was a significant

208 overabundance in human isolates. However, for antibiotics predominantly used to

209 treat farmed animals, in most cases there was a significant relative overabundance

210 in isolates from meat (**Table 2**).

Discussion

To our knowledge, this is the first study to phylogenetically compare ABR *E. coli* from

retail meats sold for consumption by human and canine populations within a

confined geographical location with contemporaneously isolated locally sourced

clinical isolates.

Of meat sold at large chain grocery stores, chicken had the highest positivity for ABR

E. coli, particularly for FQs, which, despite methodological differences between

studies, is consistent with the literature (10,12,24-26). For RDF, our findings align

with a recent report in the UK (27). Other international studies report similar trends,

although positivity rates for 3GC-R *E. coli* vary by region (13,14,28). RDF is

commonly sold frozen, and many owners incorrectly believe that freezing eliminates

pathogens (7). We therefore add to existing evidence that ABR *E. coli* remain viable

post-thawing of RDF (7,27,28).

Our phylogenetic analysis of ABR *E. coli* from chicken meat and chicken-based RDF

suggested a shared source, consistent with the fact that UK RDF is derived from the

same meat production chain as meat sold for human consumption (29). Genetic

relatedness was also observed between RDF and beef meat, possibly due to mixed

protein content in the RDF.

We have previously reported that excretion of FQ-R ST10, ST162 and ST744 *E. coli*

by dogs in Bristol is associated with them being fed raw meat (13,14). The fact that

we commonly found these same STs contaminating raw meat and RDF being sold in

Bristol provides a potential explanation.

We identified four incidences where *E. coli* isolates from meat were within 20 SNPs

of human clinical isolates, the closest being two ST69 chicken isolates with 12 and

13 SNPs different from a BSI isolate, with an ST69 beef isolate differing from a UTI

isolate by only 16 SNPs. In each case both members of the meat/infection pair

carried identical ARGs, further suggestive of isolate sharing. To our knowledge,

these represent the closest genomic relationships of *E. coli* from meat and human

extraintestinal infection isolates reported. This likely reflects the scarcity of high-

resolution phylogenetic comparisons between human and meat/livestock isolates

within the same region and time frame. One UK study (30) of similar scale to ours,

reported the closest human infection to livestock relationship as 85 SNPs. However,

the farm isolates were from an approximately 100 x 100 km region of England, the

meat isolates (n=20) were from one city within that region, but the human infection

isolates were sourced from across England, collected four to fourteen years prior to

the farm/meat isolates (30). Our findings, therefore, based on a contemporaneously

collected sample of meat/RDF (n=200) and human clinical isolates from one city

(approximately 10 x 10 km) provide insights into potential farm-to-fork transmission

routes and highlight the importance of locally integrated, high-resolution One Health

genomic surveillance for their investigation.

Limitations of our study include the cross-sectional design, which cannot establish

causality or transmission direction, and the relatively small sample size of meat

isolates, which may underrepresent the full diversity of *E. coli* in these sources. Our

study also does not capture the commensal *E. coli* present in the gastrointestinal

tracts of healthy humans.

Nonetheless, our findings have important implications for public health and food safety policy. First, they underscore the potential benefits of tighter microbiological standards and regulatory oversight of RDF, which, unlike meat sold for human consumption, is explicitly sold to be fed uncooked. Second, they support including RDF in ABR bacterial surveillance as part of an integrated One Health approach. Third, public health messaging should better inform pet owners of the risks of raw feeding, and about hygiene precautions to limit household transmission.

Funding

This work was funded by grant BB/X012670/1 to M.B.A. from the Biotechnology and Biological Sciences Research Council and grant MR/T005408/1 to P.W. and M.B.A. from the Medical Research Council. J.E.S. was supported by a scholarship from the Medical Research Foundation National PhD Training Programme in Antimicrobial Resistance Research (MRF-145-0004-TPG-AVISO).

Transparency declaration

M.B.A. is married to the owner of a veterinary practice that sells various mass-manufactured dog foods amounting to a value less than 5% of total turnover. Otherwise, the authors declare no competing interests.

Author Contributions

Conceived the Study: J.E.S., M.B.A.; Funding Acquisition: P.W., M.B.A.; Collection of Data: J.E.S., B.A., K.L.S., A.M.D, L.W., D.K., P.W.; Cleaning and Analysis of Data:

280   J.E.S., M.B.A.; Initial Drafting of Manuscript: J.E.S., M.B.A.; Corrected and Approved

281   Manuscript: All authors

282 **Table 1**. Sample-level positivity for *E. coli* in meat sold primarily for human

283 consumption at large chain grocery stores and in chicken-based RDF sold at

284 specialist pet stores. Total samples tested in brackets.

	Samples positive for *E. coli* in								
Selection	**Beef (15)**		**Chicken (15)**		**Lamb (15)**		**Pork (13)**		**RDF (15)**	
TBX only	12	80%	15	100%	11	73%	9	69%	13	87%
AMX	5	33%	14	93%	5	33%	5	38%	12	80%
SPC	4	27%	15	100%	4	27%	5	38%	12	80%
STR	7	47%	15	100%	6	40%	5	38%	13	87%
AMC	2	13%	5	33%	3	20%	2	15%	4	27%
CTX	2	13%	1	7%	0	0%	0	0%	4	27%
CIP	2	13%	7	47%	1	7%	1	8%	7	47%

286 **Table 2.** ARGs from meat and human sources conferring resistance to different

287 antibiotic classes.

Antibiotic	Poultry	Red Meat	Human	Human versus Poultry	Human versus Red Meat
Beta-lactam	139	37	588	0.038	NS
Fluoroquinolone	17	3	76	NS	NS
Gentamicin/ Tobramycin	19	2	90	NS	NS
Trimethoprim	71	13	353	0.006	0.01
Sulphonamide	130	30	482	NS	NS
Tetracycline	88	33	290	NS	0.007
Florfenicol	41	6	31	<0.00001	0.019
Neomycin/ Spectinomycin/ Streptomycin	241	66	744	0.023	0.048
Total	746	190	2654		

Orange highlights mean genes are more common in isolates from meat than human

infections, green highlights mean more common in isolates from human infections

than meat.

Figure 1 Mid-rooted phylogenetic tree of *E. coli* ST69 isolated from raw meat products (beef (blue), poultry (chicken and RDF,
yellow)) and from human clinical infection (HCI; bloodstream and urinary tract infections, blue). Single nucleotide polymorphisms
<150 SNPs in red are SNPs between raw meat source and human source, SNPs between isolates from same source are in black.

Figure 2 Mid-rooted phylogenetic tree of *E. coli* ST117 isolates from human clinical infections (BSI or UTI, blue), one from raw dog
food (RDF, yellow) and another from lamb (grey). Single nucleotide polymorphisms in red are SNPs between raw meat source and
human source, SNPs between isolates from one source in black.

References

1. Kaper JB, Nataro JP, Mobley HL. Pathogenic Escherichia coli. Nat Rev Microbiol. 2004;2:123-40. doi: 10.1038/nrmicro818.

2. Jang J, Hur HG, Sadowsky MJ, Byappanahalli MN, Yan T, Ishii S. Environmental Escherichia coli: ecology and public health implications-a review. J Appl Microbiol. 2017;123:570-581. doi: 10.1111/jam.13468.

3. Poirel L, Madec JY, Lupo A, Schink AK, Kieffer N, Nordmann P, et al. Antimicrobial Resistance in Escherichia coli. Microbiol Spectr. 2018;6:10.1128/microbiolspec.arba-0026-2017

4. FSA. 2022. Clean Beef Cattle for slaughter: A guide for producers. https://www.food.gov.uk/business-guidance/clean-beef-cattle-for-slaughter-introduction Accessed 6[th] February 2024.

5. FSA 2023. Cooking your food. https://www.food.gov.uk/safety-hygiene/cooking-your-food Accessed 5th February 2024

6. Morgan G, Williams N, Schmidt V, Cookson D, Symington C, Pinchbeck G. A Dog's Dinner: Factors affecting food choice and feeding practices for UK dog owners feeding raw meat-based or conventional cooked diets. Prev Vet Med. 2022;208:105741. doi: 10.1016/j.prevetmed.2022.105741.

7. Davies RH, Lawes JR, Wales AD. Raw diets for dogs and cats: a review, with particular reference to microbiological hazards. J Small Anim Pract. 2019;60:329-339. doi: 10.1111/jsap.13000.

8. Nüesch-Inderbinen M, Treier A, Zurfluh K, Stephan R. Raw meat-based diets for companion animals: a potential source of transmission of pathogenic and antimicrobial-resistant Enterobacteriaceae. R Soc Open Sci. 2019;6:191170. doi: 10.1098/rsos.191170.

9. Baede VO, Broens EM, Spaninks MP, Timmerman AJ, Graveland H, Wagenaar

JA, et al. Raw pet food as a risk factor for shedding of extended-spectrum beta-

lactamase-producing Enterobacteriaceae in household cats. PLoS One.

2017;12:e0187239. doi: 10.1371/journal.pone.0187239.

10. Lee KY, Lavelle K, Huang A, Atwill ER, Pitesky M, Li X. Assessment of

Prevalence and Diversity of Antimicrobial Resistant Escherichia coli from Retail

Meats in Southern California. Antibiotics (Basel). 2023;12:782. doi:

10.3390/antibiotics12040782.

11. Schlesinger DP, Joffe DJ. Raw food diets in companion animals: a critical review.

Can Vet J. 2011;52:50-4.

12. Randall LP, Lodge MP, Elviss NC, Lemma FL, Hopkins KL, Teale CJ, et al.

Evaluation of meat, fruit and vegetables from retail stores in five United Kingdom

regions as sources of extended-spectrum beta-lactamase (ESBL)-producing and

carbapenem-resistant Escherichia coli. Int J Food Microbiol. 2017;241:283-290.

doi: 10.1016/j.ijfoodmicro.2016.10.036.

13. Sealey JE, Hammond A, Mounsey O, Gould VC, Reyher KK, Avison MB.

Molecular ecology and risk factors for third-generation cephalosporin-resistant

Escherichia coli carriage by dogs living in urban and nearby rural settings. J

Antimicrob Chemother. 2022;77:2399-2405. doi: 10.1093/jac/dkac208.

14. Sealey JE, Hammond A, Reyher KK, Avison MB. One health transmission of

fluoroquinolone-resistant Escherichia coli and risk factors for their excretion by

dogs living in urban and nearby rural settings. One Health. 2023;17:100640. doi:

10.1016/j.onehlt.2023.100640.

15. Mounsey O, Wareham K, Hammond A, Findlay J, Gould VC, Morley K, et al.

Evidence that faecal carriage of resistant Escherichia coli by 16-week-old dogs in

the United Kingdom is associated with raw feeding. One Health. 2022;14:100370.

doi: 10.1016/j.onehlt.2022.100370.

16. Groat EF, Williams NJ, Pinchbeck G, Warner B, Simpson A, Schmidt VM. UK

dogs eating raw meat diets have higher risk of Salmonella and antimicrobial-

resistant Escherichia coli faecal carriage. J Small Anim Pract. 2022;63:435-441.

doi: 10.1111/jsap.13488.

17. Runesvärd E, Wikström C, Fernström LL, Hansson I. Presence of pathogenic

bacteria in faeces from dogs fed raw meat-based diets or dry kibble. Vet Rec.

2020;187:e71. doi: 10.1136/vr.105644.

18. Lefebvre SL, Reid-Smith R, Boerlin P, Weese JS. Evaluation of the risks of

shedding Salmonellae and other potential pathogens by therapy dogs fed raw

diets in Ontario and Alberta. Zoonoses Public Health. 2008;55:470-80. doi:

10.1111/j.1863-2378.2008.01145.x.

19. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al.

ResFinder 4.0 for predictions of phenotypes from genotypes. J Antimicrob

Chemother. 2020;75:3491-3500. doi: 10.1093/jac/dkaa345.

20. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, et al.

Multilocus sequence typing of total-genome-sequenced bacteria. J Clin Microbiol.

2012;50:1355-61. doi: 10.1128/JCM.06094-11.

21. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 2014;30:1312-3. doi:

10.1093/bioinformatics/btu033.

22. Argimón S, Abudahab K, Goater RJE, Fedosejev A, Bhai J, Glasner C, et al.

Microreact: visualizing and sharing data for genomic epidemiology and

phylogeography. Microb Genom. 2016;2:e000093. doi: 10.1099/mgen.0.000093.

23. Duval A, Opatowski L, Brisse S. Defining genomic epidemiology thresholds for

common-source bacterial outbreaks: a modelling study. Lancet Microbe.

2023;4:e349-e357. doi: 10.1016/S2666-5247(22)00380-9.

24. Innes GK, Patton AN, Nachman KE, Casey JA, Stapleton GS, Abraham AG, et al.

Distance and destination of retail meat alter multidrug resistant contamination in

the United States food system. Sci Rep. 2023;13:21024. doi: 10.1038/s41598-

023-48197-z.

25. Díaz-Jiménez D, García-Meniño I, Fernández J, García V, Mora A. Chicken and

turkey meat: Consumer exposure to multidrug-resistant Enterobacteriaceae

including mcr-carriers, uropathogenic E. coli and high-risk lineages such as

ST131. Int J Food Microbiol. 2020;331:108750. doi:

10.1016/j.ijfoodmicro.2020.108750.

26. Guo S, Aung KT, Leekitcharoenphon P, Tay MYF, Seow KLG, Zhong Y, et al.

Prevalence and genomic analysis of ESBL-producing Escherichia coli in retail

raw meats in Singapore. J Antimicrob Chemother. 2021;76:601-605. doi:

10.1093/jac/dkaa461.

27. Morgan G, Pinchbeck G, Taymaz E, Chattaway MA, Schmidt V, Williams N. An

investigation of the presence and antimicrobial susceptibility of

Enterobacteriaceae in raw and cooked kibble diets for dogs in the United

Kingdom. Front Microbiol. 2024;14:1301841. doi: 10.3389/fmicb.2023.1301841.

28. Cole SD, Healy I, Dietrich JM, Redding LE. Evaluation of canine raw food

products for the presence of extended-spectrum beta-lactamase- and

carbapenemase-producing bacteria of the order Enterobacterales. Am J Vet Res.

2022;83:ajvr.21.12.0205. doi: 10.2460/ajvr.21.12.0205.

29. FSA. 2024. Chapter 2.8 Animal By-Products. https://www.food.gov.uk/business-

guidance/chapter-28-animal-by-products Accessed 22nd February 2024.

30. Ludden C, Raven KE, Jamrozy D, Gouliouris T, Blane B, Coll F, et al. One Health

Genomic Surveillance of Escherichia coli Demonstrates Distinct Lineages and

Mobile Genetic Elements in Isolates from Humans versus Livestock. mBio.

2019;10:e02693-18. doi: 10.1128/mBio.02693-18.