Title:

Prevalent fast evolution of genes involved in heterochromatin functions

Leila Lin[1], Yuheng Huang[1], Jennifer McIntyre[1], Ching-Ho Chang[2], Serafin Colmenares[3], Yuh

Chwen G. Lee[1*]

1. Department of Ecology and Evolutionary Biology, University of California, Irvine

2. Division of Basic Sciences, Fred Hutchinson Cancer Center, United States

3. Department of Cell and Molecular Biology, University of California, Berkeley

*Author for Correspondence:

Grace Yuh Chwen Lee

Department of Ecology and Evolutionary Biology

Center for Complex Biological Systems

University of California, Irvine

grylee@uci.edu

Abstract

Heterochromatin is a gene-poor and repeat-rich genomic compartment universally found in eukaryotes. Despite its low transcriptional activity, heterochromatin plays important roles in maintaining genome stability, organizing chromosomes, and suppressing transposable elements (TEs). Given the importance of these functions, it is expected that the genes involved in heterochromatin regulation would be highly conserved. Yet, a handful of these genes were found to evolve rapidly. To investigate whether these previous findings are anecdotal or general to genes modulating heterochromatin, we compile an exhaustive list of 106 candidate genes involved in heterochromatin functions and investigate their evolution over short and long evolutionary time scales in *Drosophila.* Our analyses find that these genes exhibit significantly more frequent evolutionary changes, both in the forms of amino acid substitutions and gene copy number change, when compared to genes involved in Polycomb-based repressive chromatin. While positive selection drives amino acid changes within both structured domains with diverse functions and intrinsically disordered regions (IDRs), purifying selection may have maintained the proportions of IDRs of these proteins. Together with the observed negative associations between evolutionary rates of these genes and genomic TE abundance, we propose an evolutionary model where the fast evolution of genes involved in heterochromatin functions is an inevitable outcome of the unique functional roles of heterochromatin, while the rapid evolution of TEs may be an effect rather than cause. Our study provides an important global view of the evolution of genes involved in this critical cellular domain and provides insights into the factors driving the distinctive evolution of heterochromatin.

1 Introduction

Heterochromatin, first discovered as the darkly stained chromosomal regions that remain

condensed throughout the cell cycle (Heitz 1928), is a distinct cytological domain that is

conserved across eukaryotic cells (Liu et al. 2020). While a small fraction of heterochromatin

was found to be cell-type specific, or "facultative", the majority of these chromosomal blocks

remain condensed across different cell types and are known as "constitutive" heterochromatin

(Heitz 1928). Constitutive heterochromatin (referred to as "heterochromatin" for simplicity

hereafter) is usually located around centromeres and telomeres (Janssen et al. 2018), and its

underlying DNA is depleted of functional genes. Instead, heterochromatin is mainly composed

of repetitive sequences, including satellite repeats (Peacock et al. 1978) and transposable

elements (TEs) (Hoskins et al. 2007; Hoskins et al. 2015). Accordingly, heterochromatin is

oftentimes assumed to be functionally inert and nicknamed the "dark matter" of the genome.

Yet, studies have found heterochromatin playing critical roles in many chromosomal functions,

such as maintaining genome stability, mediating chromosome segregation, and ensuring proper

DNA repair of repeat-rich sequences, across eukaryotes (reviewed in (Feng and Michaels 2015;

Allshire and Madhani 2018; Janssen et al. 2018; Kendek et al. 2021)). Not surprisingly,

disruption of heterochromatin functions has been linked to various diseases (Hahn et al. 2010),

aging progression (Villeponteau 1997; J.-H. Lee et al. 2020), and cancer (Janssen et al. 2018).

The critical functions and inter-species conservation of heterochromatin domains would

naturally lead to the expectation that genes modulating the function of heterochromatin should

be highly conserved. Surprisingly, however, a handful of these genes show non-neutral

evolution of their amino acid sequence and rapid turnover of gene copy number, both of which

are consistent with a history of positive selection. The evolution of some of these genes was of

interest due to them being core structural components of heterochromatin, such as the

Heterochromatin Protein (HP1) family (Levine et al. 2012; Helleu and Levine 2018) and its

paralogs (Vermaak et al. 2005; Ross et al. 2013). The rest of the genes were observed to have

roles in the evolution of small RNA pathways (Obbard et al. 2006; Obbard et al. 2009),

transcription factors (Kasinathan et al. 2020), or germline stem cells (Flores et al. 2015), as well

as speciation (Barbash et al. 2003). Because satellite repeats (Henikoff et al. 2001) and TEs

(Cosby et al. 2019) are enriched in heterochromatic sequence and can defy Mendel's law to

increase their transmission to the next generation, an arms race between heterochromatic

sequence and host genes suppressing such selfish behaviors has been a common theme

explaining the rapid evolution of these genes (e.g., (Vermaak et al. 2005; Satyaki et al. 2014;

Helleu and Levine 2018; Brand and Levine 2021)). Further supporting such conjecture, the

composition, abundance, and location of repetitive sequences in heterochromatin are found to

diverge rapidly even between closely related species (Wei et al. 2014; Wei et al. 2018; Kim et

al. 2021; de Lima and Ruiz-Ruano 2022).

However, this handful of genes represents only a small fraction of genes involved in modulating

heterochromatin functions. The unique and essential functions of heterochromatin depend on its

specific enrichment of repressive di- and tri-methylation at histone H3 Lysine 9 (H3K9me2/3,

(Allshire and Madhani 2018)) and the "reader" of these marks, HP1a, which serves as the

foundational structural component of heterochromatin (Eissenberg and Elgin 2014) and leads to

DNA compaction (Verschure et al. 2005) and transcriptional suppression (Li et al. 2003).

Interestingly, the binding of HP1a to H3K9me2/3 further recruits "writers" of H3K9me (histone

methyltransferase, (Schotta 2002)), and this positive feedback mechanism between readers and

writers leads to the propagation of such repressive epigenetic marks independent of the

underlying DNA sequence (Allshire and Madhani 2018). This unique ability of heterochromatin

to "spread" is best exemplified by the Position Effect Variegation (PEV) first discovered in

Drosophila (Muller 1930)—the mosaic expression of euchromatic genes translocated to

heterochromatin-proximal regions due to the stochastic spreading of repressive H3K9me2/3

from heterochromatin (Girton and Johansen 2008; Elgin and Reuter 2013). By studying mutants

that either enhance or weaken PEV, many genes involved in heterochromatin functions were

identified. These include not only the structural (e.g., (Shaffer et al. 2006)) and enzymatic (e.g.,

(Czermin et al. 2001)) components for heterochromatin, but also those that antagonize the

initiation and/or maintenance of heterochromatin (e.g., writers of antagonizing histone

modifications; (Bao et al. 2007)). Still, other genes were discovered through their co-localization

with heterochromatin domain either cytologically (e.g., (Swenson et al. 2016)) or epigenomically

(e.g., (Alekseyenko et al. 2014)) and were subsequently identified to be involved in

heterochromatin function.

Given the large numbers and vastly diverse functional roles of genes involved in

heterochromatin function, we conducted an expansive survey to determine whether the

previously reported rapid evolution of a small subset of these genes is anecdotal or a common

feature of proteins involved in the functions of this unique chromatin environment. To do so, we

compiled an exhaustive list of candidate genes that have been shown, or are likely, to be

involved in heterochromatin functions, including PEV modifiers, histone-modifying enzymes

influencing H3K9me2/3 enrichment, and genes whose protein products localize to

heterochromatin. Our investigation finds that these genes evolve exceptionally fast, a pattern

that is not general to genes interacting with other repressive chromatin marks and suggests

selective pressure unique to constitutive heterochromatin. We further dissect the domains and

protein properties targeted by positive selection and specifically test the premise that the rapid

evolution of genes involved in heterochromatin functions could be driven by fast-changing

repetitive sequences. Based on our findings, we propose an evolutionary model where the rapid

changes in these genes are the unavoidable consequence of the unique functional roles of

heterochromatin, and that the evolution of repetitive sequences may be the consequence,

instead of the cause. Our study provides an important global view of the evolution of genes

involved in this critical cellular domain and sheds light on the drivers behind the unique evolution

of heterochromatin.

Results

Identification of candidate genes involved in heterochromatin functions

With the goal of identifying the global evolutionary patterns for genes involved in the functions of heterochromatin, we selected candidate genes based on three criteria to maximize inclusivity (**Table S1**). The first category contains genes demonstrated to modulate PEV (Girton and Johansen 2008; Elgin and Reuter 2013; Swenson et al. 2016). Genes with mutations weaken PEV and, therefore, heterochromatin functions are known as Suppressors of Variegation (*Su(var)*), while those that augment PEV are Enhancers of Variegation (*E(var)*). Here, we included 59 and seven previously identified *Su(var)s* and *E(var)s,* respectively.

The second category focuses on histone-modifying enzymes that can influence the enrichment levels of H3K9me2/3. Enrichment of H3K9me2/3 requires "eraser" proteins that first remove antagonizing active marks, such as H3K9ac (deacetylases, (Czermin et al. 2001)) and H3K4me3 (demethylase, (Rudolph et al. 2007)). This is followed by writer proteins depositing H3K9me2/3 (H3K9 methyltransferases, (Schotta 2002)). We consider these histone-modifying enzymes to enhance H3K9me2/3 enrichment and, thus, heterochromatin functions. On the contrary, erasers for H3K9me2/3 (H3K9 demethylase, (Herz et al. 2014)), as well as writers for acetylation (acetyltransferase, (Kuo and Andrews 2013)) and S10 kinase (Deng et al. 2008), are known to antagonize the deposition and maintenance of H3K9me2/3. H4K20me3 is another conserved hallmark of heterochromatin (Schotta et al. 2004), and we included the corresponding methyltransferases. In total, our list contains 16 and seven histone-modifying enzymes that enhance or weaken heterochromatin functions, respectively.

The last category includes genes whose protein products co-localize with heterochromatin, suggesting they play roles in this subnuclear compartment. We surveyed the literature for cytology-based evidence of co-localization with HP1a through either immunofluorescence (e.g., (Greil et al. 2007)) or live imaging (e.g., (Swenson et al. 2016)) as

well as epigenomics through Chromatin Immunoprecipitation (ChIP, e.g., (Alekseyenko et al. 2014; Kasinathan et al. 2020)). Because a previous comprehensive survey found that some HP1a-binding proteins identified through Immunoprecipitation-Mass Spectrometry (IP-Mass Spec) are not necessarily enriched in heterochromatin cytologically (Swenson et al. 2016), genes implicated in HP1a-binding but lacking localization evidence are not included in our analysis. Hereafter, we referred to this largest category with 63 genes as "heterochromatin proteins."

In total, we studied the evolution of 106 genes involved in heterochromatin functions, which we termed "heterochromatin-related genes" hereafter (see **Figure 1A** for the number of genes in each category and **Table S1** for a full list of genes and references). It is worth noting that 43 candidate genes belong to more than one category because these three categorizations are not mutually exclusive (**Table S1**). We consider genes in categories of *Su(var)s*, histone modifying enzymes enhancing H3K9me2/3 or H4K20me3 enrichment, or heterochromatin proteins as "mediating" heterochromatin functions, while those defined as *E(var)s* or histone modifying enzymes weakening H3K9me2/3 enrichment as "antagonizing" heterochromatin functions. To determine whether the evolutionary histories of heterochromatin-related genes are exceptional, we compared them with Polycomb group genes (Kassis et al. 2017), whose protein products are enriched at facultative heterochromatin. Even though also generally associated with transcriptional suppression, facultative heterochromatin is restricted to developmentally regulated genes in euchromatin and is enriched for another type of repressive histone modification (H3K27me3) (Bannister and Kouzarides 2011; Bell et al. 2023). Polycomb genes thus allowed us to determine whether the exceptional evolutionary histories, if any, of heterochromatin-related genes are due to their involvement in a repressive chromatin environment or specifically for heterochromatin structure.

Heterochromatin-related genes experience pervasive positive selection on protein

sequences across short and long-time scales

We performed evolutionary genetic tests to identify heterochromatin-related genes with

signatures of positive selection at two time scales—contrasting the polymorphism within *D.*

melanogaster and divergence between *D. melanogaster* and *D. simulans* (~4 million years

divergence between lineages) under the framework of unpolarized McDonald-Kreitman tests

(MK tests, (McDonald and Kreitman 1991)), and phylogenetic analysis by maximum likelihood

(PAML) tests (Yang 2007) using 16 *Drosophila* species with an estimated divergence time of

~25 million years (Suvorov et al. 2022) (see **Figure 1B** for studied species). In addition to

comparing the evolution of heterochromatin-related genes to that of polycomb genes, we made

the contrast to the evolution of a thousand randomly sampled genes to represent genome-wide

estimates (referred to as "random genes" hereafter).

On a short evolutionary time scale, we first estimated the proportion of amino acid

substitutions that might have been driven by positive selection (α, (Smith and Eyre-Walker

2002)). Heterochromatin-related genes, as a group, have significantly larger α than those of

Polycomb (*Mann-Whitney U tests, p* = 0.0244*, median* = 0.0847 (heterochromatin-related

genes) and -0.424 (Polycomb)) and random genes (*Mann-Whitney U tests, p* < 10^{-8}*, median* = -

0.516 (random genes)). Breaking down our candidate genes according to functions revealed

that most categories have larger α than Polycomb control or random genes, even though the

comparisons are statistically significant only for *Su(var)* and heterochromatin protein (**Figure**

1C). These observations suggest the possibility that heterochromatin-related genes experienced

frequent positive selection. Consistently, we identified that 21 out of 104 (20.2%)

heterochromatin-related genes with sufficient polymorphism data to perform MK tests have

evidence of adaptive evolution (rejection of the null hypothesis and an excess of

nonsynonymous fixed differences between species). This is a significantly larger proportion than

that of the Polycomb genes (5.56%; *Fisher's Exact Test*, *p* = 0.191; *Binomial test*, *p* < 10^{-6}, odds

ratio = 4.30) and random genes(6.94%, *Fisher's Exact Test*, $p < 10^{-3}$; *Binomial test*, $p < 10^{-5}$,

odds ratio = 2.90), while we found no difference in this proportion between Polycomb genes and

random genes (*Fisher's Exact Test* and *Binomial test*, $p = 1$). Specifically, *Su(var)* (22.0%) and

heterochromatin proteins (26.2%), both of which are expected to mediate heterochromatin

functions, have much larger proportions of genes with evidence of positive selection than either

Polycomb or random genes (**Figure 1D**). Indeed, candidate genes mediating heterochromatin

functions, as a group, show significantly more evidence of adaptive evolution than the Polycomb

(20.2%, *Fisher's Exact Test*, $p = 0.190$; *Binomial test*, $p < 10^{-6}$, odds ratio = 4.30) as well as

random genes (*Fisher's Exact Test*, $p < 10^{-3}$; *Binomial test*, $p < 10^{-4}$, odds ratio = 2.91). It is

worth noting that the number of codons involved in the MK tests is similar across categories of

genes (*Mann-Whitney U tests, p > 0.05* for all comparisons), suggesting that these observations

are unlikely driven by differences in the statistical power of the MK tests.

13   On a long evolutionary time scale, we estimated the relative rates of amino acid

substitution (nonsynonymous divergence/synonymous divergence, *dN/dS*) across 16 studied

Drosophila species. There is an overall trend that heterochromatin-related genes have larger

dN/dS ratios than the Polycomb control genes (medians = 0.0974 (heterochromatin-related

genes) *v.s.* 0.0852 (Polycomb controls)) and random genes (medians = 0.0802), and this

difference is statistically significant for comparisons to random genes (*Mann-Whitney U test, p =*

0.167 (compared to Polycomb control) and 0.0032 (compared to random genes). Nevertheless,

only the category of "heterochromatin protein," the largest category, has a statistically larger

dN/dS ratio than that of the Polycomb genes (*Mann-Whitney U test, p = 0.0204*) and random

genes (*Mann-Whitney U test, p < 10^{-5,}* **Figure 1E**). To test whether the changes in *dN/dS* ratios

of heterochromatin-related genes over the phylogenetic tree might have been driven by positive

selection, we compared the log-likelihood for two alternative models estimating *dN/dS* at each

site*:* the null model that assumes *dN/dS* ratio is beta-distributed and not greater than one (M8a)

and the alternative model in which *dN/dS* ratio is allowed to be greater than one (M8 models;

(Yang 2007)). A significantly larger likelihood of the alternative model is consistent with a history

of positive selection acting on the gene (Swanson et al. 2003). We found that the substitution

patterns of 42 heterochromatin-related genes (39.6%) better fit the alternative model,

suggesting frequent positive selection acting on them. This proportion is significantly larger than

that of Polycomb genes (26.3%; *Fisher's Exact Test*, p = 0.314; *Binomial test*, p = 0.00276,

odds ratio = 1.84) and the random genes (24.8%; *Fisher's Exact Test*, p = 0.00159; *Binomial*

test, p = 0.000975, odds ratio = 1.99), while there was no difference in this proportion between

the Polycomb and random genes (*Fisher's Exact Test*, p = 0.795; *Binomial test*, p = 0.796).

Similar to observations made on a short evolutionary time scale, *Su(var)s* and heterochromatin

proteins have stronger evidence of positive selection compared to Polycomb and random genes

(**Figure 1F**), as do all genes mediating heterochromatin functions (41.4%, *Fisher's Exact Test*, p

= 0.305; *Binomial test*, p = 0.00125, odds ratio = 1.97 (compared to Polycomb control) and

Fisher's Exact Test, p = 0.0115; *Binomial test*, p = 0.000272, odds ratio = 1.67 (compared to

random genes)). Interestingly, while histone-modifying enzymes that enhance heterochromatin,

as a group, do not exhibit exceptional rates of protein evolution, likely due to the low statistical

power associated with its small number of genes, *all* methyltransferases for H3K9me2/3

(*Su(var)3-9, egg, G9a*) show accelerated rates of protein evolution (**Table S1**), which provides

clues for the possible source of positive selection acting on heterochromatin-related genes (see

Discussion). Overall, we found that heterochromatin-related genes, especially those mediating

heterochromatin functions, experienced frequent positive selection on their protein sequence at

both short- and long-time scales, even when compared to Polycomb genes. This observation

suggests that the widespread adaptive evolution of heterochromatin-related genes is likely

driven by selection acting on their functions specific to the heterochromatin environment, rather

than by general selective pressure acting on genes interacting with repressive epigenetic marks.

Figure 1. Positive selection on protein sequences of heterochromatin-related genes

found by evolutionary tests at both short- and long- evolutionary time scales. (A) Barplot

showing the number of genes in each category of heterochromatin-related genes, the Polycomb

control genes, and a set of one thousand random genes. Those expected to mediate or

antagonize heterochromatin functions are in hues of green and orange, respectively. -HC

enzyme: histone-modifying enzymes weakening H3K9me2/3 enrichment; +HC enzyme: histone-

modifying enzymes enhancing H3K9me2/3 enrichment. (B) Phylogenetic tree for the species

included in the study. Note that the branch lengths are not to scale. (C and E) Violin plots for α,

the proportion of amino acid substitutions fixed by positive selection inferred by the MK test (C)

and *dN/dS* ratio, the relative rates of nonsynonymous substitutions (E), for different categories

of genes. Numbers of significant genes are in parentheses. (D and F) Barplots showing the

proportion of genes with significant MK tests (D) and accelerated rates of *dN/dS* driven by

positive selection (F). The number of genes in each category is either on top of the barplot (A)

or in parenthesis (D and F). *Mann-Whitney Test* (C and E) and *Binomial test* (D and F):

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$

Rapid evolution of heterochromatin-related genes also manifests as changes in gene copy number

In addition to changes in amino acid sequences, the rapid evolution of genes can be in the form of changes in gene copy number (Hastings et al. 2009), and dramatic turnover of gene copy number for a small set of heterochromatin-related genes was previously reported (Levine et al. 2012; Lewis et al. 2016; Lee et al. 2017; Helleu and Levine 2018). Changes in gene copy number may hold particular significance for heterochromatin functions—several enzymatic and structural proteins of heterochromatin have been shown to exhibit dosage-dependent effects (Elgin and Reuter 2013). Accordingly, changes in gene copy number of heterochromatin-related genes could have immediate functional consequences.

We performed reciprocal *BLAST* searches using *D. melanogaster* amino acid sequences as queries to identify homologs and paralogs in other species, followed by manual curations (see Materials and Methods). With these, we found 18 heterochromatin-related genes having differences in gene copy number among 16 *Drosophila* species studied (17.0% of the heterochromatin-related genes), with 17 genes having gains of copies and one gene loss when compared to *D. melanogaster.* Similar to our analyses for the evolution of amino acid sequences (**Figure 1**), the proportion of heterochromatin-related genes with copy number variation (CNV) is significantly more than that of Polycomb genes (5.26%, *Fisher's Exact Test*, *p* = 0.302, *Binomial test*, $p < 10^{-4}$, odds ratio = 3.68). *Su(var)* genes and heterochromatin proteins again exhibit exceptionally large proportions of genes with CNV (**Figure 2A**). CNV of several of these genes have been studied before (e.g., *HP1* (Levine et al. 2012), *AGO2* (Lewis et al. 2016)*, Cav* and *Nap-1* (Lee et al. 2017), and *mh* (Brand et al. 2024)) and our following discussions mainly focused on those first identified by our study.

By using synteny to infer the orthologous relationships between duplicates in different species (see Materials and Methods), we found that most observed CNVs for heterochromatin-related genes involve a single or two duplication events (**Figure 2B, Figure S1, and Table S2**).

The most notable exception is the nucleolin homolog, *mod,* which has 18 duplications in *D.*

eugracilis alone. *mod* plays important roles in morphogenesis (Graba et al. 1994) and

spermatogenesis (Park et al. 2023) and has been shown as a dosage-dependent *Su(var)*

(Garzino et al. 1992), suggesting these identified CNVs could readily alter heterochromatin

functions. In addition, we observed complex duplication/loss events leading to the CNV

observed in *mof* (**Figure S2**), an acetyltransferase that influences H3K9me2/3 enrichment

(Feller et al. 2015) and is known to be involved in dosage compensation (Hilfiker et al. 1997). In

addition to the duplication happening on the lineage leading to *D. kikkawai,* there is likely a

duplication event of *mof* in the common ancestor of the oriental lineage (see **Figure 1B**),

followed by a subsequent loss in the subsets of lineages that leads to the paraphyletic presence

of a *mof* duplicate (**Figure S2**)*.*

Duplicated genes are commonly found to have male-biased expression, which is

suggested as a resolution to sexual genetic conflicts (Gallach and Betrán 2011). We are

interested in examining whether duplicates of heterochromatin-related genes exhibit similar

trends of sex-biased expression. By using publicly available transcriptome data for the species

studied, we categorized the parental ortholog and duplicated paralog (with respect to *D.*

melanogaster) as male-biased, female-biased, or unbiased in expression (see Materials and

Methods). With the exception of *MTA-like* and *Su(var)2-10*, most identified duplicates indeed

show male-biased expression, irrespective of whether the parental copy is male or female-

biased in expression (**Figure 2B**). Interestingly, for *mod* in *D. eugracilis*, the parental copy has a

female-biased expression, while the 18 duplicates exhibit a mixture of male-biased, female-

biased, and unbiased expression. Another notable case is *del,* which is part of a germline

complex enabling the transcriptions of TE-targeting small RNAs from loci in pericentromeric

heterochromatin (Mohn et al. 2014) and shows female-biased expression. While the *del*

duplicate in *D. simulans* shows male-biased expression, the *orthologous* duplicated copy in the

sister species *D. sechellia* has no biases in expression. In summary, the rapid evolution of

heterochromatin-related genes, when compared to Polycomb genes, is also reflected in their

changes in gene copy number.

Figure 2. Rapid evolution of heterochromatin-related genes manifested in multiple forms.

(A) Barplot showing the proportion of heterochromatin-related genes and Polycomb control

genes with CNV. The numbers of genes with CNV are in parentheses. (B) Summary of the

duplication and loss events as well as sex-biased expression for heterochromatin-related genes

identified to have CNV. Each column represents one orthologous copy. The number of

duplication events is noted at the bottom. The colors of each square represent sex-biased

expression, except for the 19 copies of *mod* duplicates in *D.eugracilis,* which is shown at the top

left of figure B. (C) Venn diagram depicting the number of genes identified to have exceptional

evolution with one of the three evolutionary tests as well as their overlaps, with those identified

by more than one test listed. a.a.: amino acid. *Binomial tests*: **p < 0.01; ***p < 0.001.

Heterochromatin-related genes with diverse functions are recurrent targets of positive selection

The three evolutionary tests we conducted detect signals of positive selection at different time scales and in different forms (amino acid sequences *v.s.* gene copy number). We are interested in investigating whether some heterochromatin-related genes may exhibit rapid evolution with multiple tests. Consistent with analyses conducted separately for different evolutionary tests, *Su(var)s* and heterochromatin proteins have an especially large proportion of genes with evidence of rapid evolution in any one of the tests (*Binomial test, p* < 0.001 for both comparisons; **SFigure 3A**), and this difference becomes even more prominent when considering genes with exceptional evolution for *more than one test* (*Binomial test, p* < 0.001 for both comparisons; **SFigure 3B**). Three genes show evidence of rapid evolution for *all* three tests (**Figure 2C**): *AGO2*, the core effector in endogenous siRNA silencing pathway that silences TEs in somatic cells (Ghildiyal et al. 2008), *Su(var)2-10*, which critically links piRNA targeting and the transcriptional silencing of TEs (Ninova et al. 2020), and *mh*, whose evolution is the focus of a recent study (Brand et al. 2024). Other heterochromatin-related genes with more than one evidence of rapid evolution (**Figure 2C**) include those related to silencing of TEs (e.g., *del, moon, sov, Trf2, wde*), dosage compensation (e.g., *Su(var)3-7*), maintenance of chromatin structure (e.g., *Nipped-B*), and structural components of heterochromatin (e.g., *HP5*). While most of these genes are either *Su(var)* or heterochromatin proteins, several histone-modifying enzymes were also found to undergo positive selection with multiple evolutionary tests, such as the H3K4me3 demethylase *NO66* and acetyltransferase *mof*. These observations suggest that positive selection recurrently acts on heterochromatin-related genes with diverse functions.

1 **Positive selection targets both ordered domains and intrinsically disordered regions of**

2 **proteins encoded by heterochromatin-related genes**

3 To identify the potential evolutionary drivers for the observed rapid evolution of heterochromatin-

4 related genes, we investigated the locations of positive selection within windows of each rapidly

5 evolving gene. We identified windows with evidence of adaptive evolution by performing sliding

6 MK tests and located sites with high Bayes Empirical Bayes (BEB) posterior probability of

7 positive selection under the PAML framework (Yang et al. 2005). Locations of these windows or

8 sites with evidence of positive selection were then contrasted with the annotated and/or

9 predicted domains of heterochromatin-related genes (see Materials and Methods). We found

10 that signatures of positive selection are present in domains with diverse functions (**Figure 3**),

11 and several of them are especially pertinent to heterochromatin functions. These include

12 chromo domains that directly interact with histone methylation (in *HP6, rhi, Kis*) and Jmjc

13 demethylase domain in *NO66*, the H3K4me3 demethylase. We also found signatures of positive

14 selection located within various nucleic acid binding domains (e.g., C2H2 type and UBZ4-type

15 Zinc-Finger DNA binding domain) and domains mediating protein-protein interactions (e.g.,

16 BESS domains).

17 In addition to structured domains with well-characterized functions, proteins also contain

18 regions that lack fixed three-dimensional structures, known as Intrinsically Disordered Regions

19 (IDRs). IDRs are increasingly appreciated for playing critical roles in protein functions (Forman-

20 Kay and Mittag 2013) and are frequently found in proteins enriched in phase-separated cellular

21 compartments (Nott et al. 2015; Pak et al. 2016; Lin et al. 2017). Indeed, IDR-mediated phase

22 properties have been suggested to be critical for heterochromatin functions (Larson et al. 2017;

23 Strom et al. 2017). Accordingly, we investigated whether IDRs in heterochromatin-related genes

24 may also have signatures of positive evolution by using flDPnn (Hu et al. 2021) to predict IDRs

25 in *D. melanogaster* protein sequences. Interestingly, we found that signatures of rapid evolution

26 also frequently fall within IDRs (**Figure 3**), and, for multiple rapidly evolving heterochromatin-

related genes, fast-evolving sites/windows only fall within IDRs, but not other structured

domains (**Figure 3**, right column).

We also investigated whether varying IDR properties of heterochromatin-related genes

contribute to the evolutionary differences between gene categories by estimating the

percentage of amino acids falling within predicted IDR domains (% of IDRs). A higher % of IDRs

has been found in proteins involved in the formation of phase-mediated cellular domains (e.g.,

membrane-less organelles (Sawyer et al. 2019) and the heterochromatin domain (Guthmann et

al. 2019)). We found that the % of IDR for heterochromatin-related genes in *D. melanogaster* is

significantly greater than that of random genes (*Mann-Whitney U test, p* = 0.00214), which is

especially true for *Su(var)* and heterochromatin proteins (*Mann-Whitney U test, p* = 0.0448

(*Su(var)s*) and 0.0141 (heterochromatin proteins); **SFigure 4A**). There is a lack of difference in

% of IDR between our candidate genes and Polycomb control (*Mann-Whitney U test, p* > 0.05

for all tests, **SFigure 4A**), which may be due to the fact that Polycomb proteins were shown to

also undergo phase separation (Tatavosian et al. 2019). Surprisingly, despite previous reports

on the reduced evolutionary constraints in IDRs (Brown et al. 2002; Khan et al. 2015), the % of

IDRs does not differ between heterochromatin-related genes with or without evidence of positive

selection (*Mann-Whitney U test, p > 0.05*; **SFigure 4B**).

To further investigate the possible associations between % of IDRs and the evolution of

candidate genes, we estimated the phylogenetic signals (Blomberg and Garland Jr 2002) of %

of IDRs across studied species. A small *Blomberg's K* (Blomberg et al. 2003), our chosen index

for phylogenetic signal, suggests % of IDR evolves faster than that of random expectation

(Brownian motion of trait evolution; (Kamilar and Cooper 2013)). Intriguingly, we found no

difference in *K* values between heterochromatin-related genes and control genes, among

categories of heterochromatin-related genes that showed different rates of adaptive evolution,

and between heterochromatin-related genes with and without evidence of positive selection

over long evolutionary time scales (**SFigure 4C and 4D;** *Mann-Whitney U test, p* > 0.05 for all

comparisons). These observations indicate that, despite the observed pervasive adaptive

evolution of heterochromatin-related genes, their % of IDRs evolves similarly to that of other

genes in the genome. In summary, positive selection acts on both structured domains with

diverse functions as well as IDRs of heterochromatin-related genes, while stabilizing selection

might have maintained the IDR content of these genes in the face of rapidly changing amino

acid sequences.

Figure 3. Signatures of positive selection fall within domains with diverse functions and

IDRs of heterochromatin-related genes. Genes with evidence of positive selection on protein

sequences are categorized according to which types of domains/regions their signatures of

positive selection fall within. Annotated and/or predicted structured domains are shown as

horizontal lines with the following functional categories: nucleic acid binding (orange), protein-

protein interaction (yellow), histone modification binding (pink), and other domains (purple).

IDRs are marked with gray. Highlighted vertical windows represent those under positive

selection as identified by sliding MK tests (green) and PAML inferences (blue). Note that genes

with no positively selected windows/sites overlapping with either annotated domains or IDRs

were not shown.

Rates of protein evolution of heterochromatin-related genes significantly associate with genomic TE abundance

The antagonistic interaction with repetitive sequences enriched in heterochromatin was suggested as the driver for the rapid evolution of a handful of heterochromatin-related genes (Vermaak et al. 2005; Satyaki et al. 2014; Helleu and Levine 2018). To test whether this conjecture may be broadly applicable to heterochromatin-related genes, we estimated the correlation between the rates of amino acid substitution (dN/dS) of heterochromatin-related genes and the changes in abundance of heterochromatic repetitive sequences using methods developed in (Lartillot and Poujol 2011). This approach corrects for the phylogenetic nonindependence of the quantitative traits of interest (here, repeat abundance) and models their evolution following the Brownian process. To estimate repeat abundance, we performed Illumina sequencing with PCR-free library preparation to avoid the sequencing bias against AT-rich sequences, which is commonly found in repetitive sequences (Wei et al. 2018). Using these sequencing data, we quantified the genomic abundance of satellite repeats (both simple and complex satellites) and TEs, both which should be dominated by the DNA content of heterochromatin (see Materials and Methods).

Among the 42 heterochromatin-related genes with evidence of positive selection over the long evolutionary time scale, nearly forty percent (38.1%) of them have rates of protein evolution tracking changes in TE abundance across species, with significant ($p < 0.05$) phylogenetically controlled correlations between dN/dS and TE abundance (**Table S3**). This proportion is much higher than that of the 248 random genes with accelerated rates of protein evolution (20.5%, *Fisher's Exact test*, $p = 0.0173$, *Binomial test*, $p = 0.0112$, **Figure 4A**). Intriguingly, 93.8% of these correlations of the heterochromatin-related genes are in the negative direction, which indicates that these genes evolved faster in species with lower TE abundance (see **Figure 4C** for an example). We also examined this proportion separately for each category of heterochromatin-related genes (**Figure 4B**). Quite strikingly, even though only

a small number of histone-modifying enzymes were found to have evidence of positive selection

(26.1%), *all* of their rates of protein evolution were significantly associated with TE abundance

(**Figure 4B**). These histone-modifying enzymes include all three H3K9me2/3 methyltransferase

(*egg, G9a*, and *Su(var)3-9*), *NO66* (H3K4me3 demethylase), *KDM4B* (H3K9me2/3

demethylase), and *HDAC6* (zinc-dependent deacetylase) (**Table S3**). In stark contrast to

analysis focusing on TE abundance, only 7.14% of heterochromatin-related genes with

evidence of positive selection show a significant correlation between *dN/dS* and the abundance

of total satellite repeats, a proportion that is not significantly different from that of randomly

sampled genes (**Figure 4A**, *Fisher's Exact Test p* = 0.322 and *Binomial test p* = 0.360).

Analysis focusing on the abundance of only simple satellites gave consistent results (**Figure S5,**

p > 0.05 for both *Fisher's Exact Test* and *Binomial test*). Overall, we found that the rates of

protein evolution of heterochromatin-related genes negatively correlated with the abundance of

TEs, but not total repeats, across species, suggesting a possible source of selective force

shaping the evolution of these genes (see Discussions).

Figure 4. Extensive negative associations between *dN/dS* of heterochromatin-related

genes and genomic TE abundance. (A) Stacked bar plots showing the proportion of positively

selected genes whose *dN/dS* correlates with repeat abundance (left) or TE abundance (right)

for heterochromatin-related genes and randomly sampled genes. The color of the genes in the

side box represents the gene's category: *E(var)s* (yellow), *Su(var)s* (green), histone-modifying

enzyme enhancing heterochromatin (light green), and heterochromatin protein (dark green). (B)

Bar plots showing the proportion of positively selected heterochromatin-related genes whose

dN/dS significantly coevolve with TE abundance for different gene categories. (C) X-Y plots

showing the associations between *dN/dS* (X-axis) and normalized TE abundance for an

example gene (*sov*). Blue points are estimates for the extant species, and gray points are for

internal nodes. Lines across each dot denote 95% confidence intervals for *dN/dS* (extant

species and internal nodes) and normalized TE content (internal nodes). *Binomial test: n.s. p >*

0.05, **p < 0.05.*

Discussion

Heterochromatin is a highly conserved cellular compartment with essential functions across complex eukaryotes (Allshire and Madhani 2018; Janssen et al. 2018). Nevertheless, our evolutionary analyses revealed that genes involved in heterochromatin function are highly labile, exhibiting pervasive evidence of rapid evolution both in the forms of amino acid substitutions and gene copy number changes at both short and long evolutionary time scales. Importantly, the rapid evolution of these genes is likely driven by functions specific to constitutive heterochromatin, instead of mechanisms general to proteins interacting with repressive chromatin. Evidence of positive selection on protein evolution is especially prominent for heterochromatin-related genes that should enhance heterochromatin function and, strikingly, *all* three methyltransferases responsible for the enrichment of H3K9me2/3, the characteristic histone modification of heterochromatin, are under positive selection. Our further characterization of the various aspects of the evolution of heterochromatin-related genes provided an important avenue to identify the possible source of selective forces acting on heterochromatin, which we discussed below.

Close examinations of the signatures of positive selection of heterochromatin-related genes found them not only located within structured domains with well-known functions, but also inside unstructured IDRs (**Figure 3**). IDRs were recently found to be critical for the phase properties and thus functions of heterochromatin (Larson et al. 2017; Strom et al. 2017), and our observation suggests that positive selection could also act on heterochromatin functions mediated by such properties. Interestingly, despite the previous suggestions that IDRs are evolutionarily less constrained (Brown et al. 2002; Khan et al. 2015), the proportion of IDR sequences in positively selected heterochromatin-related genes is similar to other candidate genes (**Figure 4B**). Even more, we identified strong phylogenetic signals that are consistent

with the presence of stabilizing selection preserving the % of IDRs even for heterochromatin-related genes under strong positive selection. Such findings may indicate that the % of IDRs of a protein, but not the underlying amino acid sequence, is evolutionarily constrained and plays an essential role in the function of studied heterochromatin-related genes. Indeed, a few IDRs with rapidly evolving sequences were found to possess conserved molecular features (Moesa et al. 2012; Zarin et al. 2019), with some of them experimentally demonstrated to be functionally equivalent (Zarin et al. 2017). This raises another question of *why* the underlying amino acid sequences rapidly evolve, considering the need to maintain the IDR content of heterochromatin-related genes (see below).

Several of our findings suggest that TEs, but not satellite repeats, must be involved in the pervasive rapid evolution of heterochromatin-related genes. First, the associations between rates of protein evolution of heterochromatin-related genes and repeat abundance were mainly observed for TEs. In addition, many heterochromatin-related genes under positive selection have functions related to TE suppression. In particular, the interactions between the protein products of several of these genes (e.g., *del, moon, rhi, Trf2*, and *sov* (Klattenhoff et al. 2009; Mohn et al. 2014; Andersen et al. 2017; Andreev et al. 2022)) and repressive H3K9me2/3 are responsible for licensing piRNA clusters, which are TE-enriched loci generating the majorities of piRNAs targeting TEs and located within pericentromeric heterochromatin. Similar to the rapid evolution of the DNA and protein components of centromeres (i.e., the centromeric drive hypothesis, (Henikoff et al. 2001; Malik et al. 2002)), changes in heterochromatic TE sequences may alter their interactions with proteins encoded by heterochromatin-related genes. Consequently, selection may favor evolutionary changes in proteins that revert the strength of DNA-protein interaction to that prior to the alteration of TE sequences. Subsequent changes in TE sequences could initiate another cycle of this interaction and ultimately drive the fast evolution of genes involved. However, only some of the identified signatures of positive

selection fall within nucleic acid binding domains (**Figure 3**), and the majority of the positively

selected heterochromatin-related genes lack sequence specificity. Moreover, unlike viruses, the

"success" of TEs is tightly intertwined with host fitness, owing to the fact that they propagate in

the host germline and are inherited vertically (Haig 2016). Accordingly, an arms race between

TEs and host proteins suppressing them was suggested to be unlikely to drive the pervasive

adaptive evolution of host genes (Blumenstiel et al. 2016; Cosby et al. 2019).

What might have driven the fast evolution of heterochromatin-related genes then? The genomic

autoimmunity hypothesis (Blumenstiel et al. 2016) provides a plausible explanation—molecular

mechanisms suppressing TE activities are expected to be constantly juggling between

maximizing TE silencing while minimizing inadvertent off-target suppression of functional genes,

and this alternating selective pressure could drive the rapid evolution of genes involved. Under

this model, for example, variants of heterochromatin-related genes that enhance the generation

of piRNAs from piRNA clusters, and thus strengthen TE silencing, may lead to the inadvertent

production of piRNAs that target other functional elements, a possibility with empirical support

(Andersen et al. 2017; Kelleher 2021). Selection will then instead favor variants increasing the

stringency of this licensing to avoid fitness costs of the off-target effect. The repeated alternation

of selective targets could accordingly drive the rapid evolution of heterochromatin-related genes.

The genomic autoimmunity hypothesis is even more suitable to explain the rapid evolution of

heterochromatin-related genes involved in modulating the intrinsic and unique molecular

properties of heterochromatin—a tendency to "spread" to nearby loci in a sequence-

independent manner. The positive feedback between writers and readers of H3K9me2/3

promotes the propagation of repressive chromatin marks (Allshire and Madhani 2018; Bell et al.

2023), and the extent of this spreading depends on the concentrations of readers and writers at

the suppressed loci (Locke et al. 1988). Heterochromatin-mediated silencing thus needs to be

carefully balanced to prevent inadvertent silencing of functional elements while maintaining

sufficient suppression at the euchromatin-heterochromatin boundaries. Interestingly, such a

balance needs to be maintained not only at the ends of chromosomes (around pericentromeric

and subtelomeric heterochromatin), but also around epigenetically silenced, H3K9me2/3-

enriched TEs in the euchromatic genome. The spreading of repressive marks from silenced

euchromatic TEs into functional genes is widely documented (reviewed in (Choi and Lee 2020))

and has been inferred to impair individual fitness (Lee 2015; Lee and Karpen 2017; Huang et al.

2022). Consistently, several genes known to directly mediate TE epigenetic silencing are found

to be under positive selection in our analysis (e.g., *wde, Su(var)2-10,* (Ninova et al. 2020)). In

addition, histone-modifying enzymes, which are directly involved in the reader-writer positive

feedback loops, should frequently be caught in cycles of alternating selection for enhanced or

weakened TE epigenetic silencing. Indeed, we found that *all* three writers for H3K9me2/3 show

evidence of rapid evolution (*Su(var)3-9, egg, G9a*) and *all* histone-modifying enzymes with

accelerated rates of protein evolution have evidence of coevolving with genomic TE abundance

(**Figure 4**). Furthermore, the deleterious off-target effects of TE silencing could result from the

long-range spatial interactions between H3K9me2/3-enriched euchromatic TEs and

pericentromeric heterochromatin, which is mediated by phase separation mechanisms and is

selected against (Y.C.G. Lee et al. 2020). Repeated alteration in selective pressure to ensure

these proteins confer sufficient phase properties for proper heterochromatin functions while

avoiding such off-target effects may similarly drive rapid changes in protein sequences while

preserving the % of IDR for proteins encoded by heterochromatin-related genes, as we have

observed.

Intriguingly, we found predominantly negative associations between TE abundance and rates of

protein evolution of heterochromatin-related genes, which may initially seem counterintuitive.

However, this pattern can be explained by our proposed model that augments the genomic

autoimmunity hypothesis by incorporating how the alternating selective pressure on

heterochromatin-mediated silencing may concurrently influence genomic TE abundance (**Figure**

5A). Strong heterochromatin-mediated silencing should lead to rampant off-target effects, which

not only select *for* variants that weaken the strength of silencing (Huang and Lee 2024), but also

select *against* individual TE copies due to the deleterious spreading of repressive marks to TE-

adjacent functional sequences. TE abundance should accordingly decrease (**Figure 5A - a,**

(Huang et al. 2022)). When variants that reduce heterochromatin-mediated silencing become

fixed, the maintenance of heterochromatin at epigenetically silenced TEs weakens, resulting in

increased TE replication and more new TE insertions (**Figure 5A - b**). The consequentially

increased TE abundance should then drive selection *for* enhanced heterochromatin-mediated

silencing, returning to the initial state of strong silencing (**Figure 5B**). Accordingly, as

heterochromatin-related genes experience cycles of alternating selection targets and gaining

amino acid substitutions, genomic TE abundance also fluctuates. Yet, whether TE abundance

eventually increases or decreases depends on the relative strength of selection against TEs

with off-target effects and the changes in TE replication rates (**Figure 5B**). Selection coefficients

for the harmful off-target effects of TE epigenetic silencing are yet to be estimated, but they

could be strong if such effects perturb the expression of nearby vital genes (e.g., (Coronado-

Zamora and González 2023)) or disrupt global 3D genome organization (e.g., (Y.C.G. Lee et al.

2020)). On the other hand, replication rates of *Drosophila* TEs are generally low ($10^{-5}\sim10^{-4}$;

(Charlesworth and Langley 1989; Adrion et al. 2017; Wang et al. 2023)). Changes in these rates

are likely weaker than selection removing TEs through their off-target effects, leading to

decreased TE abundance over cycles of alternating selective pressure on the strength of

heterochromatin-mediated silencing and thus giving rise to negative associations between

genomic TE abundance and rates of protein evolution between species. It is worth noting that

many other processes, such as recent demographic changes (e.g., (Mérel et al. 2021)), also

contribute to between-species differences in TE abundance. If these forces systematically

influence genomic TE abundance in the species studied (e.g., correlate with the evolution of

heterochromatin-related genes), similar associations could arise in the absence of the proposed

model.

Our proposed model for how the need of maintaining a balanced TE silencing drives the fast

evolution of heterochromatin-related genes might help explain some intriguing patterns we

observed. Majorities of our identified duplicates of heterochromatin-related genes (48.78%)

have male-biased expression (**Figure 2**), indicating their potential functional significance in male

germlines. While the resolution of sexual genetic conflict (Gallach and Betrán 2011; Wyman et

al. 2012), a common explanation for male-biased expression of newly duplicated genes, may

underlie such an observation, it may also stem from the need to properly silence the male-

specific TE landscape (Chen et al. 2021), resulting in sexually dimorphic chromatin

environment. Another intriguing observation is the limited evidence of positive selection for

heterochromatin-antagonizing genes (**Figure 1**), which is in stark contrast to the pervasive

signatures of adaptive evolution of genes mediating heterochromatin functions. Protein products

of heterochromatin-antagonizing genes are not only found at heterochromatin-euchromatin

boundaries, but broadly distributed throughout the euchromatic genome (Dimova et al. 2003;

Kellner et al. 2012), playing diverse roles in transcriptional regulation (Meignin and Davis 2008;

Regnard et al. 2011; Crona et al. 2013). The pleiotropic functional roles of heterochromatin-

antagonizing proteins, in both euchromatin and heterochromatin, should constrain their

evolution, making them less likely to be caught up in alternating selection for the sensitivity or

the specificity of TE silencing. Nevertheless, the small sample size of heterochromatin-

antagonizing genes (only 12 genes) could limit our ability to detect positive selection among

these genes.

Our observed frequent positive selection acting on heterochromatin-related genes points out the

high lability of the molecular feature of this functionally conserved genomic compartment.

Instead of the usually assumed "arms race" with the changing repeatome, we proposed that the

selective pressure of heterochromatin-related genes might mainly come from a need to maintain

a delicate balance of its unique ability to "suppress" and to "spread," which also consequently

influence the evolutionary dynamics of TEs. Signatures of positive selection identified here

could serve as an entry point to further investigate how the delicate balance of heterochromatin-

mediated silencing may be conferred by vastly changing components between species (e.g.,

(Rosin and Mellone 2016; Parhad et al. 2017)), providing a fruitful opportunity to further dissect

the molecular mechanisms shaping heterochromatin functions and evolution.

Figure 5. Model for the correlated evolution of heterochromatin-related genes and genomic TE abundance. (A) The predicted impacts of different strengths of heterochromatin-mediated silencing on TE epigenetic silencing and genomic TE abundance. **(A-a)** Strong heterochromatin-mediated silencing results in strong off-target effects due to the spreading of repressive marks from silenced euchromatic TEs, leading to positive selection for variants of heterochromatin-related genes that weaken the strength of silencing. In addition, because of the low rates of TE replication from strong silencing and negative selection against TEs with off-target effects, TE abundance should reduce. **(A-b)** On the other hand, when heterochromatin-mediated silencing is weak, selection against TE-mediated off-target effects is weak and rates of TE replication should be high, leading to increased TE abundance. At the same time, selection may favor variants that enhance TE silencing to reduce rates of TE replication. **(B)** Proposed sequences of events that lead to correlated evolution of heterochromatin-related genes and genomic TE abundance. The model should be applicable to any of the states and the following description starts with strong heterochromatin-mediated silencing (i). TE abundance is expected to reduce due to the suppressed rates of TE replication and selection against TEs with off-target effects (Phase 1). Concurrently, there should be selection for variants of heterochromatin-related genes that weaken heterochromatin-mediated silencing to reduce deleterious off-target effects. When variants that weaken silencing are fixed, the maintenance of TE epigenetic silencing decreases (ii), leading to both reduced occurrence of off-target effects and increased rates of TE replication. TE abundance should thus increase (Phase 2). The high TE abundance and replicative activity would select for variants of heterochromatin-related genes that enhance TE silencing, going back to the initial state of strong heterochromatin-mediated silencing (e.g., iii here). The relative strength of a decrease in TE abundance through selection against off-target effects and increase in TE abundance through TE replication determines whether there are negative (scenario 1) or positive (scenario 2) associations between TE abundance and the evolution of heterochromatin-related genes across species.

A

B

Scenario 1: negative correlation between rates of gene evolution & TE abundance

Scenario 2: positive correlation between rates of gene evolution & TE abundance

Materials and Methods

Evolutionary analyses of protein sequences

Coding sequences and genome annotations for 16 studied species (**Figure 1B**) were

downloaded from NCBI, with GenBank ID listed in **Table S4**. Because we compiled the list of

candidate genes based on *D. melanogaster* literature, we used orthologous information from

OrthoDB (Kuznetsov et al. 2023)(last accessed 12/10/2022) to retrieve one-to-one orthologs for

D. melanogaster candidate genes. For genes without one-to-one ortholog according to

OrthoDB, we performed *BLAST* search (see below). The retrieved coding sequences were

translated to amino acid sequences, aligned using Clustal Omega (version 1.2.4; (Sievers et al.

2011)), and converted back to codon alignments using PAL2NAL (version 14; (Suyama et al.

2006)).

We performed unpolarized McDonald-Kreitman (MK) tests (McDonald and Kreitman 1991)

using polymorphism within a *D. melanogaster* Zambian population (197 strains, (Lack et al.

2015)) and divergence between *D. melanogaster* and *D. simulans.* Following the

recommendation of (Lack et al. 2015), we treated genomic regions with non-African ancestry or

identity-by-descent as missing data, and only included sites with at least 100 non-missing

alleles. To count the number of nonsynonymous and synonymous changes, we used the

mutational paths that minimize the number of amino acid changes. Codons with more than two

alleles, considered both within species polymorphism and between species divergence, were

excluded. Genes with fewer than 100 codons were excluded from the analysis due to

insufficient statistical power. A gene is deemed under positive selection if the MK test, whose

significance was assessed using *Fisher's Exact* test, rejected the null hypothesis and the ratio of

nonsynonymous to synonymous changes is greater for between-species substitutions than for

within-species polymorphism. Sliding window MK tests were performed with windows of 100

codons and 10-codon steps.

We conducted phylogenetic analysis by maximum likelihood (PAML) (Yang 2007) using 16

species to identify candidate genes experiencing positive selection over a long evolutionary time

scale. We compared two site models, M8a (*dN/dS* ratio is beta-distributed and not greater than

one) and M8 (*dN/dS* > 1), and determined the significance using likelihood ratio tests. The

species tree reported in (Suvorov et al. 2022) was used. Sites with > 0.95 BEB posterior

probability of coming from the site class with *dN/dS* > 1 (Yang et al. 2005) are considered under

recurrent adaptive evolution.

Evolutionary analysis for gene copy number

To identify genes with varying gene copy numbers, we first used a genome-wide, high

throughput search with liberal parameters to identify many potential candidates, followed by

careful manual curations. We first used tblastn and reciprocal blastx (Camacho et al. 2009) to

search for homologs and paralogs of candidate genes in studied species using *D. melanogaster*

amino acid sequence as queries, with the following parameter: e-value < 10^{-2}, amino acid

identity > 20%, and blast score > 10. We required the best reciprocal blastx hit to be the original

D. melanogaster query. Each potential CNV was manually validated using reciprocal best blast

with more stringent criteria (e-value < 10^{-5}), and orthologs and paralogs were distinguished

using synteny of flanking genes (**Figure S1**). We further examined the expression levels of

candidate CNVs using RNA-seq exon coverage tracks of NCBI Data Viewer and removed those

with no expression. Several duplicates identified in *D. suzukii* were filtered due to redundant

contigs. For gene loss, we followed the procedures detailed in (King et al. 2019) to confirm the

true absence of a gene.

To examine the sex-biased expression, we deemed a gene male-biased if the log2 fold change

of the ratio of male and female expression (TPM or FPKM) is >1, female-biased if such value is

< -1, and otherwise unbiased. For *D. melanogaster,* we used Insect Genome database

(http://www.insect-genome.com/Sexdb/, last retrieved November 2023) and FlyAtlas2 RNA-seq

data (Krause et al. 2022) for whole adult males and whole adult females. For the other 15

species, we mapped publicly available transcriptome datasets (**Table S5**) to publicly available

genome assemblies with gene annotations using HiSAT2 (v2.2.1 with parameters –exon and –

ss to specify the exon positions and splice sites; RRID:SCR_015530; (Kim et al. 2019)), and

estimated the expression levels usingStringtie (v2.1.4 with parameters -dta -G to specify

annotation files; RRID:SCR_016323; (Kovaka et al. 2019)). Candidate duplicates identified by

manual curation but have no annotation or show no expression were excluded from the

analysis.

Domain annotations

We used UniProt (The UniProt Consortium 2023) to annotate known structured domains within

the *D. melanogaster* allele of heterochromatin-related genes. We used flDPnn (Hu et al. 2021),

which performed superior in the previous benchmark study (Necci et al. 2021), and the

predicted binary index for IDRs (disorder propensity cutoff = 0.3) to annotate IDRs for all studied

species. To detect the phylogenetic signal of the % of IDR, we computed *Blomberg's K* statistic

using the phylosig function from phytools R package (Blomberg et al. 2003; Revell 2012). The

tree structure and branch lengths were obtained from (Suvorov et al. 2022) and generated by

treeio R package (Wang et al. 2020).

Analyses of repetitive sequences and their coevolution with candidate genes

DNA for each studied species was extracted from 40 females using DNeasy Blood & Tissue Kit

(Qiagen), following the manufacturer's protocol. To avoid PCR amplification bias during the

preparation of sequencing libraries, which was found to skew the quantification of repeats (Wei

et al. 2018), extracted DNA was prepared into Illumina sequencing library with PCR-free

protocol and sequenced with 150bp paired-end reads by Novogene (Sacramento, CA).

We used Satellite Repeat Finder (Zhang et al. 2023) to estimate the abundance of total satellite

repeat in the Illumina short-read sequencing data. Following the suggested procedures, we first

counted K-mers (K=21) in each sample using KMC (ver. 3.2.4; (Kokot et al. 2017)). Contigs of

satellites were then generated and mapped back to the source Illumina sequences to estimate

the abundance (in bases) of each satellite using minimap (ver. 2.24; (Li 2018)) and Satellite

Repeat Finder. We also used K-seek (Wei et al. 2014) to estimate the abundance of simple

repeats and reached similar conclusions (see Results).

TE abundance was estimated as the total TE reads from the Illumina sequencing data, with the

assumption that TEs in the heterochromatin mainly originated from jumping events of

euchromatic TEs and TE abundances in the heterochromatin and euchromatin are highly

correlated. To minimize TE annotation bias across species, repetitive sequences from each of

the 16 genomes (**Table S4**) were identified using RepeatMasker (version 4.1.0;

http://www.repeatmasker.or/) with the Dfam database (Storer et al. 2021), using the command

"RepeatMasker -q [genome sequence file] -species drosophila -e hmmer". TE sequences

annotated as LTR, LINE, DNA element, and Unknown categories were obtained from all 16

genomes to create a master TE library. Illumina reads from each species were then mapped to

the library with bwa-mem (version 0.7.17; (Li and Durbin 2009)) and viewed by samtools

(version 1.15.1, (Li 2011)). The total number of reads mapped to the library, regardless of

mapping quality, was considered the number of TE reads. It is worth noting that very few TEs

were classified as Unknown category, and inclusion/exclusion of such TEs did not change the

results. To compare the abundance of satellite repeats and TEs across samples/species, these

estimates were normalized. Illumina reads from each species were mapped to its repeat-

masked genome using bwa-mem, with sites with mapping quality lower than 30 filtered (using

samtools -q 30). The median read depth for the unmasked regions for each sample was then

used to normalize the number of bases for satellite repeats and TEs.

We used Coevolve (Lartillot and Poujol 2011) to estimate the correlation between *dN/dS* and

repeat abundance (satellite repeats or TEs) given the tree structure. For each gene, the

analysis was performed in duplicate to assess convergence (relative difference < 0.1) and a

burn-in of 300 with at least 3,000 MCMC chains to get the final estimated correlation between

dN/dS and repeat abundance. Because the number of Polycomb genes with significant PMAL

tests is small, we only compared the coevolution results of candidate genes to those of 248

randomly chosen genes with accelerated rates of amino acid evolution (i.e., significant for PAML

analysis).

Data and script availability

PCR-free Illumina data has been deposited to SRA under the accession number

PRJNA1113679. Scripts used in this study can be found at

https://github.com/YuhengHuang87/HC_gene_evo and

https://github.com/hilynano/Homology_detection.

Acknowledgments

We thank the High Performance Cluster at UC Irvine for computational resources and Harsh

Shukla for helping with the flDPnn software. We appreciate Aniek Janssen, Andrea Betancourt,

and Adriana Ludwig for their helpful comments on the manuscript. CHC was supported by

Damon-Runyon Cancer Research Foundation postdoctoral fellowship DRG 2438-21 and NIH

1   R01-GM74108 (to Hamit S Malik), SUC was supported by NIH R35GM139653 (to Gary H.

2   Karpen), YH, LL, JM, and YCGL were supported by NIH R35-GM14292 (to YCGL).

Supplementary Materials

Table S1. List of candidate and control genes

Names, FBgn, categories, and references of candidate heterochromatin-related genes and

Polycomb control genes. Results of evolutionary tests of each of the genes are also included.

Table S2. Duplicates of heterochromatin-related genes identified in this study

Table S3. Correlation between repeat abundance and dN/dS ratio for positively selected

heterochromatin-related genes

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$*

Table S4. NCBI assembly number of reference genomes used in this study

Table S5. Transcriptome resources and information used in this study

Figure S1. Synteny information for genes with duplicates found in more than one species

The following figures show the NCBI genome browser tracks for genes with duplicates (red

boxes) found in more than one species. Genes flanking the duplicates (either dark or light blue

boxes) were used to infer the synteny of duplicates between species.

Figure S2. Evolutionary history of *mof* duplicates

The generation of *mof* CNVs likely involved a duplication in the common ancestor of the oriental

linage (green), followed by a subsequent loss in the subsets of lineages (red) and a lineage-

specific duplication event (green leading to *D. kikkawai*), resulting in the paraphyletic presence

of a *mof* duplicate. Numbers next to species indicate the number of gene copies; species

without labeled numbers have one gene copy.

Figure S3. Proportion of genes with more than one significant evolutionary tests

Barplots showing the proportion of genes found to be under positive selection and/or fast evolve

with at least one (A) or at least two (B) conducted evolutionary tests. The numbers of genes in

each category are shown in parentheses. -HC enzyme: histone-modifying enzymes weakening

H3K9me2/3 enrichment; +HC enzyme: histone-modifying enzymes enhancing H3K9me2/3

enrichment. *Binomial test*: ****p* < 0.001.

Figure S4. The percentage of intrinsically disordered regions (% of IDRs) among gene categories.

(A and B) Violin plots comparing the % IDR of *D. melanogaster* proteins for heterochromatin-

related genes and the Polycomb control (A) and for heterochromatin-related genes with and

without evidence of positive selection over both long and short evolutionary time scales (B). (C

and D) Violin plots comparing *Blomberg's K* for % of IDR across species for heterochromatin-

related genes and the Polycomb control (C) and between heterochromatin-related genes with

and without evidence of positive selection over a long evolutionary time scale (D). *Mann-Whitney U test*: **p* < 0.05 and *n.s. p* > 0.05.

Figure S5. Associations between *dN/dS* of heterochromatin-related genes and the abundance of simple repeats.

Stacked bar plots showing the proportion of positively selected genes whose *dN/dS* correlates

with the abundance of simple satellite repeats for heterochromatin-related genes and randomly

sampled genes. The numbers of genes in each category are in parentheses. *Binomial test: n.s. p* > 0.05.

References

Adrion JR, Song MJ, Schrider DR, Hahn MW, Schaack S. 2017. Genome-Wide Estimates of Transposable Element Insertion and Deletion Rates in Drosophila Melanogaster. *Genome Biol Evol* 9:1329–1340.

Alekseyenko AA, Gorchakov AA, Zee BM, Fuchs SM, Kharchenko PV, Kuroda MI. 2014. Heterochromatin-associated interactions of Drosophila HP1a with dADD1, HIPP1, and repetitive RNAs. *Genes Dev* 28:1445–1460.

Allshire RC, Madhani HD. 2018. Ten principles of heterochromatin formation and function. *Nature Reviews Molecular Cell Biology* 19:229–244.

Andersen PR, Tirian L, Vunjak M, Brennecke J. 2017. A heterochromatin-dependent transcription machinery drives piRNA expression. *Nature* 549:54–59.

Andreev VI, Yu C, Wang J, Schnabl J, Tirian L, Gehre M, Handler D, Duchek P, Novatchkova M, Baumgartner L, et al. 2022. Panoramix SUMOylation on chromatin connects the piRNA pathway to the cellular heterochromatin machinery. *Nat Struct Mol Biol* 29:130–142.

Bannister AJ, Kouzarides T. 2011. Regulation of chromatin by histone modifications. *Cell Research* 21:381–395.

Bao X, Deng H, Johansen J, Girton J, Johansen KM. 2007. Loss-of-function alleles of the JIL-1 histone H3S10 kinase enhance position-effect variegation at pericentric sites in Drosophila heterochromatin. *Genetics* 176:1355–1358.

Barbash DA, Siino DF, Aaron M, Roote J. 2003. A rapidly evolving MYB-related protein causes species isolation in Drosophila. *Proc. Natl. Acad. Sci. U.S.A* 100:5302–5307.

Bell O, Burton A, Dean C, Gasser SM, Torres-Padilla M-E. 2023. Heterochromatin definition and function. *Nat Rev Mol Cell Biol* 24:691–694.

Blomberg SP, Garland Jr T. 2002. Tempo and mode in evolution: phylogenetic inertia, adaptation and comparative methods. *Journal of Evolutionary Biology* 15:899–910.

Blomberg SP, Garland T, Ives AR. 2003. TESTING FOR PHYLOGENETIC SIGNAL IN COMPARATIVE DATA: BEHAVIORAL TRAITS ARE MORE LABILE. *Evolution* 57:717–745.

Blumenstiel JP, Erwin AA, Hemmer LW. 2016. What Drives Positive Selection in the Drosophila piRNA Machinery? The Genomic Autoimmunity Hypothesis. *Yale J Biol Med* 89:499–512.

Brand CL, Levine MT. 2021. Functional Diversification of Chromatin on Rapid Evolutionary Timescales. *Annual Review of Genetics* 55:401–425.

Brand CL, Oliver GT, Farkas IZ, Buszczak M, Levine MT. 2024. Recurrent Duplication and Diversification of a Vital DNA Repair Gene Family Across Drosophila. *Molecular Biology and Evolution* 41:msae113.

Brown CJ, Takayama S, Campen AM, Vise P, Marshall TW, Oldfield CJ, Williams CJ, Dunker AK. 2002. Evolutionary rate heterogeneity in proteins with long disordered regions. *J Mol Evol* 55:104–110.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Charlesworth B, Langley CH. 1989. The Population Genetics of Drosophila Transposable Elements. *Annual Review of Genetics* 23:251–287.

Chen P, Kotov AA, Godneeva BK, Bazylev SS, Olenina LV, Aravin AA. 2021. piRNA-mediated gene regulation and adaptation to sex-specific transposon expression in D. melanogaster male germline. *Genes Dev.* 35:914–935.

Choi JY, Lee YCG. 2020. Double-edged sword: The evolutionary consequences of the epigenetic silencing of transposable elements. *PLOS Genetics* 16:e1008872.

Coronado-Zamora M, González J. 2023. The epigenetics effects of transposable elements are

context dependent and not restricted to gene silencing. :2023.11.27.568862. Available from: https://www.biorxiv.org/content/10.1101/2023.11.27.568862v1

Cosby RL, Chang N-C, Feschotte C. 2019. Host–transposon interactions: conflict, cooperation, and cooption. *Genes Dev.* 33:1098–1116.

Crona F, Dahlberg O, Lundberg LE, Larsson J, Mannervik M. 2013. Gene regulation by the lysine demethylase KDM4A in *Drosophila*. *Developmental Biology* 373:453–463.

Czermin B, Schotta G, Hülsmann BB, Brehm A, Becker PB, Reuter G, Imhof A. 2001. Physical and functional association of SU(VAR)3-9 and HDAC1 in Drosophila. *EMBO Rep* 2:915–919.

Deng H, Bao X, Cai W, Blacketer MJ, Belmont AS, Girton J, Johansen J, Johansen KM. 2008. Ectopic histone H3S10 phosphorylation causes chromatin structure remodeling in Drosophila. *Development* 135:699–705.

Dimova DK, Stevaux O, Frolov MV, Dyson NJ. 2003. Cell cycle-dependent and cell cycle-independent control of transcription by the Drosophila E2F/RB pathway. *Genes Dev.* 17:2308–2320.

Eissenberg JC, Elgin SCR. 2014. HP1a: a structural chromosomal protein regulating transcription. *Trends in Genetics* 30:103–110.

Elgin SCR, Reuter G. 2013. Position-effect variegation, heterochromatin formation, and gene silencing in Drosophila. *Cold Spring Harb Perspect Biol* 5:a017780.

Feller C, Forné I, Imhof A, Becker PB. 2015. Global and Specific Responses of the Histone Acetylome to Systematic Perturbation. *Molecular Cell* 57:559–571.

Feng W, Michaels SD. 2015. Accessing the Inaccessible: The Organization, Transcription, Replication, and Repair of Heterochromatin in Plants. *Annu. Rev. Genet.* 49:439–459.

Flores HA, DuMont VLB, Fatoo A, Hubbard D, Hijji M, Barbash DA, Aquadro CF. 2015. Adaptive evolution of genes involved in the regulation of germline stem cells in Drosophila melanogaster and D. simulans. *G3 (Bethesda)* 5:583–592.

Forman-Kay JD, Mittag T. 2013. From sequence and forces to structure, function, and evolution of intrinsically disordered proteins. *Structure* 21:1492–1499.

Gallach M, Betrán E. 2011. Intralocus sexual conflict resolved through gene duplication. *Trends Ecol Evol* 26:222–228.

Garzino V, Pereira A, Laurenti P, Graba Y, Levis RW, Le Parco Y, Pradel J. 1992. Cell lineage-specific expression of modulo, a dose-dependent modifier of variegation in Drosophila. *EMBO J* 11:4471–4479.

Ghildiyal M, Seitz H, Horwich MD, Li C, Du T, Lee S, Xu J, Kittler ELW, Zapp ML, Weng Z, et al. 2008. Endogenous siRNAs Derived from Transposons and mRNAs in Drosophila Somatic Cells. *Science* 320:1077–1081.

Girton JR, Johansen KM. 2008. Chapter 1 Chromatin Structure and the Regulation of Gene Expression: The Lessons of PEV in Drosophila. In: Genetics B-A in, editor. Vol. 61. Long-Range Control of Gene Expression. Academic Press. p. 1–43. Available from: http://www.sciencedirect.com/science/article/pii/S0065266007000016

Graba Y, Laurenti P, Perrin L, Aragnol D, Pradel J. 1994. The modifier of variegation modulo gene acts downstream of dorsoventral and HOM-C genes and is required for morphogenesis in Drosophila. *Dev Biol* 166:704–715.

Greil F, de Wit E, Bussemaker HJ, van Steensel B. 2007. HP1 controls genomic targeting of four novel heterochromatin proteins in Drosophila. *The EMBO Journal* 26:741–751.

Guthmann M, Burton A, Torres-Padilla M. 2019. Expression and phase separation potential of heterochromatin proteins during early mouse development. *EMBO Rep* 20:e47952.

Hahn M, Dambacher S, Schotta G. 2010. Heterochromatin dysregulation in human diseases. *Journal of Applied Physiology* 109:232–242.

Haig D. 2016. Transposable elements: Self-seekers of the germline, team-players of the soma. *Bioessays* 38:1158–1166.

Hastings P, Lupski JR, Rosenberg SM, Ira G. 2009. Mechanisms of change in gene copy number. *Nat Rev Genet* 10:551–564.

Heitz E. 1928. Das Heterochromatin der Moose. *Jahrb Wiss Botanik* 69:762–818.

Helleu Q, Levine MT. 2018. Recurrent Amplification of the Heterochromatin Protein 1 (HP1) Gene Family across Diptera. *Molecular Biology and Evolution* 35:2375–2389.

Henikoff S, Ahmad K, Malik HS. 2001. The Centromere Paradox: Stable Inheritance with Rapidly Evolving DNA. *Science* 293:1098–1102.

Herz H-M, Morgan M, Gao X, Jackson J, Rickels R, Swanson SK, Florens L, Washburn MP, Eissenberg JC, Shilatifard A. 2014. Histone H3 lysine-to-methionine mutants as a paradigm to study chromatin signaling. *Science* 345:1065–1070.

Hilfiker A, Hilfiker-Kleiner D, Pannuti A, Lucchesi JC. 1997. mof, a putative acetyl transferase gene related to the Tip60 and MOZ human genes and to the SAS genes of yeast, is required for dosage compensation in Drosophila. *EMBO J* 16:2054–2060.

Hoskins RA, Carlson JW, Kennedy C, Acevedo D, Evans-Holm M, Frise E, Wan KH, Park S, Mendez-Lago M, Rossi F, et al. 2007. Sequence Finishing and Mapping of Drosophila melanogaster Heterochromatin. *Science* 316:1625–1628.

Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, Booth BW, Pfeiffer BD, George RA, Svirskas R, et al. 2015. The Release 6 reference sequence of the Drosophila melanogaster genome. *Genome Res.*:gr.185579.114.

Hu G, Katuwawala A, Wang K, Wu Z, Ghadermarzi S, Gao J, Kurgan L. 2021. flDPnn: Accurate intrinsic disorder prediction with putative propensities of disorder functions. *Nat Commun* 12:4438.

Huang Y, Lee YCG. Blessing or curse: how the epigenetic resolution of host-transposable element conflicts shapes their evolutionary dynamics. *Proc Biol Sci* 291:20232775.

Huang Y, Shukla H, Lee YCG. 2022. Species-specific chromatin landscape determines how transposable elements shape genome evolution.Nordborg M, Przeworski M, editors. *eLife* 11:e81567.

Janssen A, Colmenares SU, Karpen GH. 2018. Heterochromatin: Guardian of the Genome. *Annual Review of Cell and Developmental Biology* 34:265–288.

Kamilar JM, Cooper N. 2013. Phylogenetic signal in primate behaviour, ecology and life history. *Philos Trans R Soc Lond B Biol Sci* 368:20120341.

Kasinathan B, Colmenares SU III, McConnell H, Young JM, Karpen GH, Malik HS. 2020. Innovation of heterochromatin functions drives rapid evolution of essential ZAD-ZNF genes in Drosophila.Wittkopp PJ, editor. *eLife* 9:e63368.

Kassis JA, Kennison JA, Tamkun JW. 2017. Polycomb and Trithorax Group Genes in Drosophila. *Genetics* 206:1699–1725.

Kelleher ES. 2021. Protein–Protein Interactions Shape Genomic Autoimmunity in the Adaptively Evolving Rhino-Deadlock-Cutoff Complex.Betran E, editor. *Genome Biology and Evolution* 13:evab132.

Kellner WA, Ramos E, Bortle KV, Takenaka N, Corces VG. 2012. Genome-wide phosphoacetylation of histone H3 at Drosophila enhancers and promoters. *Genome Res.* 22:1081–1088.

Kendek A, Wensveen MR, Janssen A. 2021. The Sound of Silence: How Silenced Chromatin Orchestrates the Repair of Double-Strand Breaks. *Genes* 12:1415.

Khan T, Douglas GM, Patel P, Nguyen Ba AN, Moses AM. 2015. Polymorphism Analysis Reveals Reduced Negative Selection and Elevated Rate of Insertions and Deletions in Intrinsically Disordered Protein Regions. *Genome Biology and Evolution* 7:1815–1826.

Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, Comeault AA, Peede D, D'Agostino ER, Pelaez J, et al. 2021. Highly contiguous assemblies of 101 drosophilid genomes.Coop G, Wittkopp PJ, Sackton TB, editors. *eLife* 10:e66405.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37:907–915.

King TD, Leonard CJ, Cooper JC, Nguyen S, Joyce EF, Phadnis N. 2019. Recurrent Losses and Rapid Evolution of the Condensin II Complex in Insects. *Mol. Biol. Evol.* 36:2195–2204.

Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, Zhang F, Schultz N, Koppetsch BS, Nowosielska A, et al. 2009. The Drosophila HP1 homolog Rhino is required for transposon silencing and piRNA production by dual-strand clusters. *Cell* 138:1137–1149.

Kokot M, Długosz M, Deorowicz S. 2017. KMC 3: counting and manipulating k-mer statistics. *Bioinformatics* 33:2759–2761.

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019. Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome Biology* 20:278.

Krause SA, Overend G, Dow JAT, Leader DP. 2022. FlyAtlas 2 in 2022: enhancements to the Drosophila melanogaster expression atlas. *Nucleic Acids Res* 50:D1010–D1015.

Kuo Y-M, Andrews AJ. 2013. Quantitating the Specificity and Selectivity of Gcn5-Mediated Acetylation of Histone H3. *PLOS ONE* 8:e54896.

Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov EM. 2023. OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. *Nucleic Acids Research* 51:D445–D451.

Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, Langley CH, Pool JE. 2015. The Drosophila Genome Nexus: A Population Genomic Resource of 623 Drosophila melanogaster Genomes, Including 197 from a Single Ancestral Range Population. *Genetics*:genetics.115.174664.

Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA, Redding S, Narlikar GJ. 2017. Liquid droplet formation by HP1α suggests a role for phase separation in heterochromatin. *Nature* 547:236–240.

Lartillot N, Poujol R. 2011. A phylogenetic model for investigating correlated evolution of substitution rates and continuous phenotypic characters. *Mol Biol Evol* 28:729–744.

Lee J-H, Kim EW, Croteau DL, Bohr VA. 2020. Heterochromatin: an epigenetic point of view in aging. *Exp Mol Med* 52:1466–1474.

Lee YCG. 2015. The Role of piRNA-Mediated Epigenetic Silencing in the Population Dynamics of Transposable Elements in Drosophila melanogaster. *PLoS Genet* 11:e1005269.

Lee YCG, Karpen GH. 2017. Pervasive epigenetic effects of Drosophila euchromatic transposable elements impact their evolution. *eLife* [Internet] 6. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5505702/

Lee YCG, Leek C, Levine MT. 2017. Recurrent Innovation at Genes Required for Telomere Integrity in Drosophila. *Molecular Biology and Evolution* 34:467–482.

Lee YCG, Ogiyama Y, Martins NMC, Beliveau BJ, Acevedo D, Wu C -ting, Cavalli G, Karpen GH. 2020. Pericentromeric heterochromatin is hierarchically organized and spatially contacts H3K9me2 islands in euchromatin. *PLOS Genetics* 16:e1008673.

Levine MT, McCoy C, Vermaak D, Lee YCG, Hiatt MA, Matsen FA, Malik HS. 2012. Phylogenomic Analysis Reveals Dynamic Evolutionary History of the Drosophila Heterochromatin Protein 1 (HP1) Gene Family. *PLOS Genetics* 8:e1002729.

Lewis SH, Salmela H, Obbard DJ. 2016. Duplication and Diversification of Dipteran Argonaute Genes, and the Evolutionary Divergence of Piwi and Aubergine. *Genome Biol Evol* 8:507–518.

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27:2987–2993.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.

Li Y, Danzer JR, Alvarez P, Belmont AS, Wallrath LL. 2003. Effects of tethering HP1 to euchromatic regions of the Drosophila genome. *Development* 130:1817–1824.

de Lima LG, Ruiz-Ruano FJ. 2022. In-Depth Satellitome Analyses of 37 Drosophila Species Illuminate Repetitive DNA Evolution in the Drosophila Genus. *Genome Biology and Evolution* 14:evac064.

Lin Y, Currie SL, Rosen MK. 2017. Intrinsically disordered sequences enable modulation of protein phase separation through distributed tyrosine motifs. *J Biol Chem* 292:19110–19120.

Liu J, Ali M, Zhou Q. 2020. Establishment and evolution of heterochromatin. *Ann N Y Acad Sci* 1476:59–77.

Locke J, Kotarski MA, Tartof KD. 1988. Dosage-dependent modifiers of position effect variegation in Drosophila and a mass action model that explains their effect. *Genetics* 120:181–198.

Malik HS, Vermaak D, Henikoff S. 2002. Recurrent evolution of DNA-binding motifs in the Drosophila centromeric histone. *Proc Natl Acad Sci U S A* 99:1449–1454.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the Adh locus in Drosophila. *Nature* 351:652–654.

Meignin C, Davis I. 2008. UAP56 RNA helicase is required for axis specification and cytoplasmic mRNA localization in *Drosophila*. *Developmental Biology* 315:89–98.

Mérel V, Gibert P, Buch I, Rodriguez Rada V, Estoup A, Gautier M, Fablet M, Boulesteix M, Vieira C. 2021. The Worldwide Invasion of Drosophila suzukii Is Accompanied by a Large Increase of Transposable Element Load and a Small Number of Putatively Adaptive Insertions. *Molecular Biology and Evolution* [Internet]. Available from: https://doi.org/10.1093/molbev/msab155

Mohn F, Sienski G, Handler D, Brennecke J. 2014. The Rhino-Deadlock-Cutoff Complex Licenses Noncanonical Transcription of Dual-Strand piRNA Clusters in Drosophila. *Cell* 157:1364–1379.

Muller HJ. 1930. Types of visible variations induced by X-rays inDrosophila. *Journ. of Gen.* 22:299–334.

Necci M, Piovesan D, Tosatto SCE. 2021. Critical assessment of protein intrinsic disorder prediction. *Nat Methods* 18:472–481.

Ninova M, Chen Y-CA, Godneeva B, Rogers AK, Luo Y, Fejes Tóth K, Aravin AA. 2020. Su(var)2-10 and the SUMO Pathway Link piRNA-Guided Target Recognition to Chromatin Silencing. *Molecular Cell* 77:556-570.e6.

Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, Craggs TD, Bazett-Jones DP, Pawson T, Forman-Kay JD, et al. 2015. Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles. *Mol Cell* 57:936–947.

Obbard DJ, Gordon KHJ, Buck AH, Jiggins FM. 2009. The evolution of RNAi as a defence against viruses and transposable elements. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 364:99–115.

Obbard DJ, Jiggins FM, Halligan DL, Little TJ. 2006. Natural selection drives extremely rapid evolution in antiviral RNAi genes. *Curr. Biol.* 16:580–585.

Pak CW, Kosno M, Holehouse AS, Padrick SB, Mittal A, Ali R, Yunus AA, Liu DR, Pappu RV, Rosen MK. 2016. Sequence Determinants of Intracellular Phase Separation by Complex Coacervation of a Disordered Protein. *Mol Cell* 63:72–85.

Parhad SS, Tu S, Weng Z, Theurkauf WE. 2017. Adaptive Evolution Leads to Cross-Species Incompatibility in the piRNA Transposon Silencing Machinery. *Developmental Cell* 43:60-70.e5.

Park JI, Bell GW, Yamashita YM. 2023. Derepression of Y-linked multicopy protamine-like genes interferes with sperm nuclear compaction in D. melanogaster. *Proc Natl Acad Sci U S A* 120:e2220576120.

Peacock WJ, Lohe AR, Gerlach WL, Dunsmuir P, Dennis ES, Appels R. 1978. Fine structure and evolution of DNA in heterochromatin. *Cold Spring Harb Symp Quant Biol* 42 Pt 2:1121–1135.

Regnard C, Straub T, Mitterweger A, Dahlsveen IK, Fabian V, Becker PB. 2011. Global Analysis of the Relationship between JIL-1 Kinase and Transcription. *PLOS Genetics* 7:e1001327.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3:217–223.

Rosin L, Mellone BG. 2016. Co-evolving CENP-A and CAL1 Domains Mediate Centromeric CENP-A Deposition across Drosophila Species. *Dev Cell* 37:136–147.

Ross BD, Rosin L, Thomae AW, Hiatt MA, Vermaak D, Cruz AFA de la, Imhof A, Mellone BG, Malik HS. 2013. Stepwise Evolution of Essential Centromere Function in a Drosophila Neogene. *Science* 340:1211–1214.

Rudolph T, Yonezawa M, Lein S, Heidrich K, Kubicek S, Schäfer C, Phalke S, Walther M, Schmidt A, Jenuwein T, et al. 2007. Heterochromatin Formation in Drosophila Is Initiated through Active Removal of H3K4 Methylation by the LSD1 Homolog SU(VAR)3-3. *Molecular Cell* 26:103–115.

Satyaki PRV, Cuykendall TN, Wei KH-C, Brideau NJ, Kwak H, Aruna S, Ferree PM, Ji S, Barbash DA. 2014. The Hmr and Lhr Hybrid Incompatibility Genes Suppress a Broad Range of Heterochromatic Repeats. *PLOS Genetics* 10:e1004240.

Sawyer IA, Sturgill D, Dundr M. 2019. Membraneless nuclear organelles and the search for phases within phases. *WIREs RNA* 10:e1514.

Schotta G. 2002. Central role of Drosophila SU(VAR)3-9 in histone H3-K9 methylation and heterochromatic gene silencing. *The EMBO Journal* 21:1121–1131.

Schotta G, Lachner M, Sarma K, Ebert A, Sengupta R, Reuter G, Reinberg D, Jenuwein T. 2004. A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev* 18:1251–1262.

Shaffer CD, Cenci G, Thompson B, Stephens GE, Slawson EE, Adu-Wusu K, Gatti M, Elgin SCR. 2006. The Large Isoform of Drosophila melanogaster Heterochromatin Protein 2 Plays a Critical Role in Gene Silencing and Chromosome Structure. *Genetics* 174:1189–1204.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, et al. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7:539.

Smith NGC, Eyre-Walker A. 2002. Adaptive protein evolution in Drosophila. *Nature* 415:1022–1024.

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of transposable element families, sequence models, and genome annotations. *Mob DNA* 12:2.

Strom AR, Emelyanov AV, Mir M, Fyodorov DV, Darzacq X, Karpen GH. 2017. Phase separation drives heterochromatin domain formation. *Nature* 547:241–245.

Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, Price DK, Waddell PJ, Lang M, Courtier-Orgogozo V, et al. 2022. Widespread introgression across a phylogeny of 155 Drosophila genomes. *Current Biology* 32:111-123.e5.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Research* 34:W609–W612.

Swanson WJ, Nielsen R, Yang Q. 2003. Pervasive Adaptive Evolution in Mammalian

Fertilization Proteins. *Molecular Biology and Evolution* 20:18–20.

Swenson JM, Colmenares SU, Strom AR, Costes SV, Karpen GH. 2016. The composition and organization of Drosophila heterochromatin are heterogeneous and dynamic. *eLife* 5:e16096.

Tatavosian R, Kent S, Brown K, Yao T, Duc HN, Huynh TN, Zhen CY, Ma B, Wang H, Ren X. 2019. Nuclear condensates of the Polycomb protein chromobox 2 (CBX2) assemble through phase separation. *J Biol Chem* 294:1451–1463.

The UniProt Consortium. 2023. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Research* 51:D523–D531.

Vermaak D, Henikoff S, Malik HS. 2005. Positive selection drives the evolution of *rhino*, a member of the heterochromatin protein 1 family in *Drosophila*. *PLoS Genet* 1:96–108.

Verschure PJ, van der Kraan I, de Leeuw W, van der Vlag J, Carpenter AE, Belmont AS, van Driel R. 2005. In Vivo HP1 Targeting Causes Large-Scale Chromatin Condensation and Enhanced Histone Lysine Methylation. *Molecular and Cellular Biology* 25:4552–4564.

Villeponteau B. 1997. The heterochromatin loss model of aging. *Exp Gerontol* 32:383–394.

Wang L-G, Lam TT-Y, Xu S, Dai Z, Zhou L, Feng T, Guo P, Dunn CW, Jones BR, Bradley T, et al. 2020. Treeio: An R Package for Phylogenetic Tree Input and Output with Richly Annotated and Associated Data. *Molecular Biology and Evolution* 37:599–603.

Wang Y, McNeil P, Abdulazeez R, Pascual M, Johnston SE, Keightley PD, Obbard D. 2023. Variation in mutation, recombination, and transposition rates in Drosophila melanogaster and Drosophila simulans. *Genome Res.*:gr.277383.122.

Wei KH-C, Grenier JK, Barbash DA, Clark AG. 2014. Correlated variation and population differentiation in satellite DNA abundance among lines of Drosophila melanogaster. *PNAS* 111:18793–18798.

Wei KH-C, Lower SE, Caldas IV, Sless TJS, Barbash DA, Clark AG. 2018. Variable Rates of Simple Satellite Gains across the Drosophila Phylogeny. *Mol. Biol. Evol.* 35:925–941.

Wyman MJ, Cutter AD, Rowe L. 2012. GENE DUPLICATION IN THE EVOLUTION OF SEXUAL DIMORPHISM. *Evolution* 66:1556–1566.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24:1586–1591.

Yang Z, Wong WSW, Nielsen R. 2005. Bayes empirical bayes inference of amino acid sites under positive selection. *Mol Biol Evol* 22:1107–1118.

Zhang Y, Chu J, Cheng H, Li H. 2023. De novo reconstruction of satellite repeat units from sequence data. *Genome Res.* [Internet]. Available from: https://genome.cshlp.org/content/early/2023/11/20/gr.278005.123