1 **Whole-brain neural substrates of behavioral variability in the larval zebrafish**
2

3 Jason Manley[1,2] and Alipasha Vaziri[1,2,*]
4
5 1) Laboratory of Neurotechnology and Biophysics, The Rockefeller University, New York,
6   NY 10065, USA
7 2) The Kavli Neural Systems Institute, The Rockefeller University, New York, NY 10065, USA
8 * Correspondence: vaziri@rockefeller.edu

9
10
11 **ABSTRACT**
12

Animals engaged in naturalistic behavior can exhibit a large degree of behavioral variability
even under sensory invariant conditions. Such behavioral variability can include not only
variations of the same behavior, but also variability across qualitatively different behaviors
driven by divergent cognitive states, such as fight-or-flight decisions. However, the neural
circuit mechanisms that generate such divergent behaviors across trials are not well
understood. To investigate this question, here we studied the visual-evoked responses of larval
zebrafish to moving objects of various sizes, which we found exhibited highly variable and
divergent responses across repetitions of the same stimulus. Given that the neuronal circuits
underlying such behaviors span sensory, motor, and other brain areas, we built a novel Fourier
light field microscope which enables high-resolution, whole-brain imaging of larval zebrafish
during behavior. This enabled us to screen for neural loci which exhibited activity patterns
correlated with behavioral variability. We found that despite the highly variable activity of
single neurons, visual stimuli were robustly encoded at the population level, and the visual-
encoding dimensions of neural activity did not explain behavioral variability. This robustness
despite apparent single neuron variability was due to the multi-dimensional geometry of the
neuronal population dynamics: almost all neural dimensions that were variable across
individual trials, i.e. the "noise" modes, were orthogonal to those encoding for sensory
information. Investigating this neuronal variability further, we identified two sparsely-
distributed, brain-wide neuronal populations whose pre-motor activity predicted whether the
larva would respond to a stimulus and, if so, which direction it would turn on a single-trial level.
These populations predicted single-trial behavior seconds before stimulus onset, indicating they
encoded time-varying internal modulating behavior, perhaps organizing behavior over longer
timescales or enabling flexible behavior routines dependent on the animal's internal state. Our
results provide the first whole-brain confirmation that sensory, motor, and internal variables
are encoded in a highly mixed fashion throughout the brain and demonstrate that de-mixing
each of these components at the neuronal population level is critical to understanding the
mechanisms underlying the brain's remarkable flexibility and robustness.

INTRODUCTION

Animals are not deterministic input-output machines, but instead display highly flexible behavioral responses even under sensory invariant conditions. A holistic understanding of the neuronal mechanisms underlying decision making requires explaining such variability in behavior at the single trial level. However, there are many putative sources of trial-to-trial neuronal variability. In particular, it is unclear whether such variability results from deterministic sources, such as internal states and long-timescale dynamics, or stochasticity. Individual neurons exhibit non-negligible noise from a variety of sources, including electrical noise due to ion channel fluctuations, synaptic noise due to biochemical processes, jitter in the timing of action potentials, and "digitization" noise due to the all-or-nothing nature of the action potential (Faisal et al., 2008).

However, trial-to-trial variability in neuronal firing is generally not independent across neurons, but instead highly correlated over repeated stimulus presentations within certain groups of neurons (Kohn et al., 2015; Shadlen and Newsome, 1998; Zohary et al., 1994), suggesting that variability in neural circuits may not be dominated by the stochasticity of individual neurons. Such structured covariation is referred to as "noise correlations" (Cohen and Kohn, 2011) because they are unrelated to the often experimenter-controlled external variable that is simultaneously encoded within the neuronal population. While certain structures of noise correlations are known to limit the information encoding capacity of the neuronal population (Moreno-Bote et al., 2014; Bartolo et al., 2020; Rumyantsev et al., 2020), there is little consensus about both their origin (Kanitscheider et al., 2015; Zylberberg et al., 2016) and behavioral impact (Zohary et al., 1994; Huang and Lisberger, 2009). In particular, there are opposing reports in different species and under different task conditions as to whether such inter-neuronal correlations can improve (Cafaro and Rieke, 2010; Zylberberg et al., 2016; Valente et al., 2021) or interfere (Cohen and Maunsell, 2009; Ruda et al., 2020; Kafashan et al., 2021) with the neural computations underlying decision making. As such, a general framework capable of reconciling such widely varying reports and explaining the functional impact of noise correlations on neural computation and behavior is currently missing.

Further, few experimental studies have linked noise correlations in sensory regions to pre-motor dynamics and ultimately behavior (Valente et al., 2021). Instead, most reports have mainly focused on the scale and structure of such covariations across sensory neurons in anesthetized animals or without regard to the animal's behavioral state. However, recent studies have shown that many species encode the animal's own movements in a highly distributed fashion, including across early sensory regions (Kauvar et al., 2020; Musall et al., 2019; Stringer et al., 2019). These brain-wide, motor-related fluctuations must then drive at least some of the observed neuronal covariation across stimulus presentations. Thus, noise correlations are not merely stochastic "noise", but instead may encode important variables that are stimulus-independent and perhaps related to the animal's proprioception and context, such as long-timescale behavioral and internal state dynamics.

Such highly overlapping representations of sensory information and additional contextual
variables may underly the brain's remarkable ability to flexibly control behavior and select the
most appropriate action at any given moment based on the animal's goals and internal state. As
such, naturalistic decision making is a function of not just incoming sensory stimuli, but also
internal variables such as the animal's motivations and recent experience or learned behaviors.
For example, foraging has been shown across species to be modulated by hunger (Filosa et al.,
2016), long-timescale exploration versus exploitation states (Flavell et al., 2013; Marques et al.,
2020), and other hidden internal variables (Sternson, 2020; Torigoe et al., 2021; Flavell et al.,
2022). This evidence suggests that answering longstanding questions regarding the neuronal
mechanisms underlying naturalistic decision making will require understanding the intersection
of numerous neural circuits distributed throughout the vertebrate brain.
99
Thus, probing the structure of the neuronal mechanisms underlying behavioral variability
requires high spatiotemporal resolution recording of brain-wide neuronal dynamics and
behavior across many trials. While it is often necessary to average across trials to deal with
both the inherent variable firing of single neurons and experimental measurement noise, such
trial averaging inherently precludes any investigation of the link between neural and behavioral
variability on the trial-by-trial level. Instead, methods that utilize the entire neuronal population
dynamics have shown that pooling information across neurons can allow for successful
decoding of information that is not possible from individual neurons. This is because
simultaneous multi-neuron recordings can leverage statistical power across neurons to capture
the most salient aspects of population dynamics within single trials. Such population-level
approaches have revolutionized the fields of motor planning and decision making, providing
highly robust brain-machine interfaces (Kao et al., 2015; Pandarinath et al., 2018) that combine
information from at least a few recording sites and making it possible to predict decisions (Wei
et al., 2019; Lin et al., 2020), reaction time (Afshar et al., 2011), and additional behavioral states
(Kaufman et al., 2015; Marques et al., 2020) at the single trial level. Recently, optical imaging
techniques (Weisenburger and Vaziri, 2018; Kim and Schnitzer, 2022; Machado et al., 2022)
have dramatically increased the attainable number of simultaneously recorded neurons within
increasingly larger brain volumes at high speed and resolution. Thus, large-scale neural
recording provides a promising approach to both screening for potential sources of variability
across the brain and identifying robust population-level dynamics at the level of single trials.
However, due to the difficulty of obtaining whole-brain optical access in many vertebrates,
previous studies have not been able to quantify brain-wide noise correlations at the single
neuron level and link such noise modes to variable decisions at the single trial level.
123
In this study, we sought to identify the neural loci which drive or encode information related to
behavioral variability across trials. To do so, we performed whole-brain recording at cellular
resolution in larval zebrafish engaged in ethologically relevant visually-evoked behaviors to
identify neuronal populations that were predictive of behavioral variability under sensory
invariant conditions. While it is well-known that larval zebrafish on average exhibit a target-
directed prey capture response to small visual stimuli and a target-avoidance escape response
to large visual stimuli, we demonstrated that these ethological, oppositely valanced, and highly
divergent responses exhibited significant variability across trials of identical stimulus

presentations. Turning first to visually-evoked neural populations, we found that despite trial-
to-trial variability in the responses of single neurons, visual information was reliably encoded at
the population level across trials of each stimulus. As such, the visual-encoding neuronal
activity patterns did not account for the larvae's variable behavioral responses across trials.
136
A key feature of our system, unlike previous studies, was that it allowed investigating the
geometry of trial-to-trial variability of neuronal dynamics on the whole-brain-level, by
simultaneous observation of brain-wide neural variability during a visual decision-making
behavior. Thus, we were able to identify whole-brain modes of neural variability which were
related to the larva's motor actions. In particular, we identified highly distributed neuronal
populations whose pre-motor activity predicted the larvae's responsiveness and turn direction
on a single trial level, indicating that behavioral variability is not represented in particular
anatomical neural loci. We found that this predictability exhibited two dominant timescales: a
longer-timescale and pre-stimulus turn bias that was not time-locked to the stimulus
presentations or motor timing; and a rapid increase in predictability and ramping activity about
one to two seconds before movement initiation. Consistent with result from previous studies,
we speculate that the former, longer-timescale bias is related to a circuitry that has evolved to
optimize an exploration versus exploitation behavior in forging while the latter, short-timescale
ramping activity likely drives the downstream motor circuitry to execute the actual selected
action in each trial. Our data suggest that behavioral variability in response to repetitions of the
same sensory stimulus may not be driven by a single brain region. Rather, it is more likely
generated by a combination of factors encoded by neurons throughout the brain, including a
time-varying and internal turn direction bias, in addition to the well-studied visuomotor
transformations.
156
157
158 **RESULTS**
159
160 **Fourier Light Field Microscopy enables high-speed imaging of neuronal population dynamics**
161
In order to investigate the whole-brain neural correlates of behavioral variability, we designed
and built a Fourier light field microscope (fLFM) that was optimized for high-resolution, whole-
brain recoding of neuronal activity in the larval zebrafish. Conventional light field microscopy
(cLFM) provides an elegant solution to volumetric snapshot imaging by placing a microlens
array (MLA) in the image plane to encode both the position and angle of incidence of the light
rays onto a camera sensor (Levoy et al., 2006). Combined with 3D deconvolution algorithms for
computational reconstruction of the recorded volume (Agard, 1984; Broxton et al., 2013;
Prevedel et al., 2014), LFM enables imaging of an entire volume without requiring any scanning.
Due to its high speed and volumetric capabilities, cLFM has been widely applied to image the
dynamics of *in vivo* biological systems, particularly neuronal dynamics (Prevedel et al., 2014;
Aimon et al., 2019; Lin et al., 2020; Quicke et al., 2020; Yoon et al., 2020). Further, additional
computational strategies have been devised to image deep into scattering tissue, such as the
mouse brain (Nöbauer et al., 2017; Skocek et al., 2018; Nöbauer et al., 2023). However, a
primary limitation of cLFM is a dramatic drop in resolution near the native image plane (NIP)

because its spatio-angular sampling is uneven across depth and highly redundant near the NIP.
This low resolution at the center of the sample can only be partially overcome using techniques
such as wavefront coding (Cohen et al., 2014). In contrast, fLFM has been shown to provide
high-resolution imaging across a two- to three-fold extended depth by processing the light field
information through the Fourier domain (Llavador et al., 2016; Scrofani et al., 2017; Cong et al.,
2017; Guo et al., 2019; Liu et al., 2020); i.e., placing the MLA conjugate to the back focal plane
of the objective. In this configuration, the MLA segments the wavefront by transmitting the
various spatial frequencies (which correspond to angular information) into images on different
regions of the camera, providing a well-aliased sampling of the spatio-angular information
without significant redundancy near the NIP.
186
We simultaneously optimized our design for simplicity and cost-effectiveness by using
exclusively off-the-shelf elements, as opposed to previous implementations of fLFM which
relied on custom optical components. Our design (Figure 1A, see Methods for details) enables
imaging of an approximately $750 \times 750 \times 200$ μm^3 volume, corresponding to roughly the whole
brain of a 7 days post fertilization (dpf) larval zebrafish. 3D information is captured in the raw
images (Figure 1B) because each lenslet in the MLA segments a region of the Fourier plane and
thus forms an image of the sample from a unique perspective. Experimentally, the point spread
function (PSF) full width at half maximum (FWHM) of our system measured 3.3 ± 0.21 μm (95%
confidence interval) laterally and 5.4 ± 0.4 μm axially (Figure 1C), consistent with theoretical
estimates of 3.1 μm and 4.7 μm, respectively. This represents a substantial improvement in
axial resolution compared to our previous cLFM, which had a PSF FWHM of 3.4 μm laterally and
11.3 μm axially (Prevedel et al., 2014). As such, our design enabled cellular-resolution imaging
across the whole brain of the larval zebrafish.
200
Next, we developed a custom pipeline (Figure 1D) for 3D reconstruction and neuronal
segmentation to extract the dynamics of neurons across the field of view. This consisted of
three main steps: denoising, 3D reconstruction, and neuronal segmentation. As one of the key
limitations of LFM is signal-to-noise ratio (SNR) due to the distribution of emitted photons onto
a sensor array, we first denoised the raw sensor images. To do so we trained a
DeepInterpolation (Lecoq et al., 2021) deep neural network model, which has been shown to
increase the SNR of the resulting neuronal timeseries. Next, the full 3D volume was
reconstructed from denoised sensor images using Richardson-Lucy deconvolution (Agard, 1984;
Broxton et al., 2013; Prevedel et al., 2014) and an experimentally measured PSF. In addition to
enabling high-resolution imaging across an extended depth of field, a key advantage of fLFM is
that the reconstruction of the volume images can be performed approximately 100 times faster
than in cLFM, due to fLFM's shift-invariant point spread function. Finally, to identify neuronal
regions of interest (ROIs) and their dynamics within the reconstructed volume, we utilized
CNMF-E (Zhou et al., 2018), a constrained matrix factorization approach to extract *in vivo*
calcium signals from one-photon imaging data. The CNMF-E algorithm was applied to each
plane in parallel, after which the putative ROIs from each plane were collated and duplicate
neurons across planes were merged.
218

To validate our setup, we imaged whole-brain dynamics from head-immobilized larval zebrafish
expressing nuclear-localized GCaMP6s (NL-GCaMP6s) pan-neuronally (Vladimirov et al., 2014)
while monitoring the larva's tail movements using a high-speed camera and presenting visual
stimuli, which as we will discuss below consisted of single dots of 3-5 different sizes moving
from the center toward the left or right visual field. Across our 1-2 hour recordings, we
identified 16,524 $\pm$ 3,942 neurons per animal (mean $\pm$ 95% confidence interval; range: 6,425 to
35,928). Thus, the combination of fLFM and the improvement in our reconstruction pipeline
enabled the detection of significantly higher neuron numbers than the ~5,000 neurons as
reported in our previous cLFM realizations (Prevedel et al., 2014; Lin et al., 2020). The
performance improvement over cLFM could also be visualized by inspecting the maximum
intensity projection (MIP) of a zebrafish recording in each modality. While cLFM (Figure 1E)
exhibits lower resolution and grid-like artifacts near the native image plane (black arrows) due
to redundant sampling near the NIP, fLFM's well-aliased sampling of spatio-angular information
provides cellular-resolution imaging throughout the entire volume (Figure 1F).
233
234   Within these observed large-scale dynamics, we found a diversity of neuronal activity patterns
235   (Figure 1G). For example, we identified neurons tuned to the vigor the larva's tail movements
236   (see Methods for details) and those tuned to the presentation of visual stimuli as described
237   below (Figure 1H), providing a proof of principle that fLFM enables whole-brain imaging of the
238   relevant neuronal population activity encoding sensory inputs and behavior.
239
240   **Larval zebrafish exhibit highly variable motor responses to visual stimuli**
241
Whole-brain, cellular-resolution imaging at high speed enabled us to screen for regions across
the brain which exhibited covariations related to variability in behavior and naturalistic decision
making. We thus set out to investigate the neuronal basis of trial-to-trial variability in action
selection during two different ethologically-relevant behaviors which are known to be
generated by brain-wide sensorimotor circuitries (Chen et al., 2018). In the larval zebrafish,
several visually-evoked behaviors such as prey capture (Borla et al., 2002; Patterson et al.,
2013) and escape response (Temizer et al., 2015) have been widely studied. In particular,
varying a single stimulus parameter, the size of a visual object, can elicit dramatically distinct
average behavioral responses (Barker and Baier, 2015), ranging from target-directed responses
(e.g., prey capture) to small stimuli, to target-avoidance behaviors (e.g., escape response) to
large stimuli. These distinct behaviors can be evoked in the laboratory using simple visual
stimuli and in a head-fixed imaging preparation (Bianco et al., 2011; Semmelhack et al., 2014;
Temizer et al., 2015; Barker and Baier, 2015; Bianco and Engert, 2015; Timothy W. Dunn et al.,
2016; Filosa et al., 2016; Förster et al., 2020; Oldfield et al., 2020). As such, the neural circuitry
underlying each of the individual behaviors is well understood (Portugues and Engert, 2009;
Bollmann, 2019) on the trial-averaged level. However, importantly these behaviors are not
reflexive. They involve the processing of information across multiple brain regions and are
subjected to neuromodulatory effects and longer time scale changes of internal states (Filosa et
al., 2016; Marques et al., 2020). As such, the larvae do not deterministically perform the same
actions over multiple sensory invariant trials. Even under highly optimized and invariant
conditions, prey capture is only observed in up to 30% of trials (Bianco et al., 2011), whereas

escape response is more consistently observed in 80% of trials (Temizer et al., 2015). However,
the neuronal populations and the underlying circuitry driving such variability in responsiveness
and action selection in either of these paradigms have not been identified, yet these
ethologically highly divergent behaviors can be elicited by tuning a single parameter. We thus
hypothesized that modulating the size of the visual object should reveal a regime of behavioral
variability between these highly divergent, naturalistic behaviors, providing a useful means and
context to study questions such as how behavioral strategies may switch across trials or how
neural and behavioral variability changes as a function of the ambiguity of the presented
stimulus.
272
Thus, we first sought to characterize the trial-to-trial variability of the larvae's behavioral
responses to repetitions of simple visual stimuli of various sizes. In a freely swimming context
(Figure 2A), we displayed moving dots of various sizes ranging from 1 to 40 visual degrees (see
Methods for details) on a screen positioned below the larvae. Analyzing these behavioral
recordings for different dot sizes, we indeed found characteristic target-avoidance bouts similar
to escape response (Figure 2B) and target-directed bouts similar to prey capture response
(Figure 2C; see also Video 1). For each bout of behavior in which a larva was close to a visual
object, we quantified whether it approached or avoided the target using the larva's change in
distance from the object as a metric (see Methods for details). Consistent with previous work
(Barker and Baier, 2015), on average the larvae exhibited target-directed behavior to small
stimuli (1-5°), followed by a crossover point at which the larvae were not preferentially target-
directed or target-avoiding for stimuli of 7° in size, and then a target-avoidance behavioral
regime to stimuli at above 10° or larger which elicited characteristic escape responses on
average (Figure 2D). Moreover, across individual trials the larvae's responses exhibited
significant variability, yielding many trials in which the larvae either did not respond or
responded in the opposite direction of the average response, as indicated by the large standard
deviation across bouts for the same stimulus size (Figure 2D).
290
Next, we confirmed that these behaviors were preserved in a head-fixed imaging preparation,
in which the larvae were embedded in agarose with their eyes and tail cut free (Figure 2E), such
that their variable behavioral responses to visual stimuli could be tracked while performing
whole-brain neuronal recordings using our fLFM system. To do so, we presented visual stimuli
as dots of a certain size moving in either the left or right visual field of the animals (see
Methods for details). Visual stimuli were created using a projector that projected onto the edge
of the circular 35 mm petri dish in which the larvae were embedded. Utilizing curvature of the
tail to the left or the right as a metric for target-directed versus -avoidance behavior, we again
identified the stimulus-size dependence of the larvae's average behavioral response (Figure 2F)
that recapitulated what was observed in freely behaving larvae. We also confirmed that the
corresponding opposite tail responses were observed when the stimuli were presented on the
other side of the visual field, demonstrating that even during head-fixation the larvae exhibit on
average directed behaviors that depend on the stimulus size. However, at the single trial level,
we observed behavioral variability to nearly all stimulus sizes (Figure 2G), even in the stimulus
size regime known to optimally drive prey capture and escape response behaviors. The most
widely variable responses were at 4° and 11°, corresponding to stimuli in the target-directed

regime and near the crossover between target-directed and avoidance, respectively. This high
degree of variability among responses to small and intermediate stimuli is consistent with
previous literature estimates that even in optimized setups, the prey capture response is
observed at rates less than 30% during head-fixation (Bianco et al., 2011) and any motor
response at all to small visual stimuli is observed at rates up to 60% of trials in free behavior
(Filosa et al., 2016). However, the neural mechanisms and circuitry underlying such trial-to-trial
behavioral variability are not understood. In particular, it is unclear whether such variability
within these ethologically relevant, yet divergent behaviors is driven by noise and inherent
stochasticity in neuronal circuits, or represents modulation by time-varying internal states, such
as motivation and hunger.
317
318   **Trial-to-trial variability in visually-evoked neurons is largely orthogonal to visual decoding**
319   **dimensions**
320
Previous studies have reported that individual neurons tuned to specific stimuli often exhibit
significant variability in their responses over multiple presentations of the same stimulus
(Montijn et al., 2016; Rumyantsev et al., 2020; Shadlen and Newsome, 1998; Zohary et al.,
1994; Zylberberg et al., 2016). Thus, one potential mechanism underlying our observed
behavioral variability could be trial-to-trial deviations in the neural encoding of visual stimuli.
Given that we had observed behavioral variability across the entire studied range of stimulus
sizes, we proceeded to record whole-brain dynamics with the fLFM while presenting many
repetitions of 3 to 6 different stimulus sizes spanning 1° to 40° (see Methods for details).
Investigating the responses of individual neurons across the whole brain, we indeed found that
even neurons highly tuned to a particular stimulus exhibited variability in their responses across
multiple presentations of the same stimuli (Figure 3A). Given that downstream decision-making
neuronal ensembles likely pool information from many visually tuned neurons, we asked
whether an optimal population decoder could reliably extract information about the visual
stimulus across trials. We proceeded to build various logistic regression classifiers to decode
which visual stimulus was presented from the whole-brain neuronal activity pattern during the
stimulus presentation period of each trial. We found that despite the observed variability on
the single neuron level, the stimuli could be reliably decoded from the visually-tuned neurons
identified within the whole-brain population activity (Figure 3B) at the single trial level with an
accuracy of $94 \pm 2\%$ (mean $\pm$ 95% confidence interval). This robust decodability suggests that
the observed single neuron variability does not severely limit the information encoding capacity
at the population level.
342
Given the wealth of literature debating whether trial-to-trial noise correlations could interfere
with stimulus encoding (Moreno-Bote et al., 2014; Kanitscheider et al., 2015; Zylberberg et al.,
2017; Bartolo et al., 2020; Rumyantsev et al., 2020; Valente et al., 2021; Kafashan et al., 2021)
and potentially drive behavioral variability, we next asked how the observed neuronal
variability was structured to prevent degradation of the visual encoding. We began by
considering the dimensions that maximally explain trial-to-trial neuronal variability within the
high-dimensional space of neuronal population dynamics, which are sometimes referred to as
"noise modes" (Rumyantsev et al., 2020). Historically, the term "noise" is used to describe such

trial-to-trial variability that is not related to the external stimulus; however, such noise could
represent significant internal neuronal dynamics and not just measurement shot noise, for
example. If the primary noise modes (represented by neural activity along the vectors $\vec{e}_\alpha$) are
orthogonal to the optimal stimulus decoding direction ($\vec{w}_{opt}$), then trial-to-trial variability will
not hinder the visual stimulus decoding (Figure 3C(i)). In this case, the neuronal population may
carry a multi-dimensional set of at least two variables and the visual stimulus encoding would
be preserved. Trial-to-trial variability could also not be limited to a particular set of dimensions,
for example if it represented noise that is uniformly distributed in the neural state space (Figure
3C(ii)), in which case pooling or averaging across sufficient neurons may still recover adequate
stimulus information. Finally, trial-to-trial noise modes may be aligned with the stimulus
decoding dimension (Figure 3C(iii)), which would maximally degrade the visual stimulus
decoding.
363
To assess which of these scenarios applied to the geometry of trial-to-trial neural variability in
our data, we first utilized partial least squares (PLS) regression to identify a low-dimensional
subspace of the whole-brain neural activity that optimally preserved information about the
visual stimulus (Figure S3A, see Methods for details), which we refer to as the visually-evoked
neuronal activity patterns. Importantly, this approach ensures that we are identifying trial-to-
trial neural variation that is contained within the visually-evoked neurons, as opposed to
variability that is completely unrelated to visual encoding. Within these visually-evoked
neurons, PLS regression also identifies the optimal stimulus decoding direction $\vec{w}_{opt}$ (Figure
S3B). Additionally, trial-to-trial variability of the visually-evoked neurons is summarized by their
noise covariance matrix, which describes how the activity of neurons covaries across trials
(Moreno-Bote et al., 2014; Kohn et al., 2015). The average noise correlation matrix across all
stimuli (see Methods for details) appeared highly structured and multi-dimensional (Figure 3D),
indicating the presence of multiple neuronal ensembles whose activity is strongly correlated
across trials. We analyzed the structure of this trial-to-trial noise by finding the eigenvectors $\vec{e}_\alpha$
of the average neural noise covariance matrix across all stimuli, following (Rumyantsev et al.,
2020). As such, these noise modes $\vec{e}_\alpha$ represented the dimensions of neuronal activity within
the visually-evoked neurons that maximally covaried across trials. We found that the noise
modes were strongly structured since the trial-to-trial variance was concentrated in relatively
few dimensions. The single largest noise mode captured up to 50% of the variance across larvae
(Figure 3E, y-axis), indicating that such activity is likely not merely "noise" but potentially
physiologically relevant. Thus, the observed neuronal variability was not independent across
neurons, but strongly correlated. To assess the impact these noise modes could have on the
stimulus decoding, we calculated the angles between the noise modes $\vec{e}_\alpha$ and $\vec{w}_{opt}$. We
discovered that the noise modes containing the majority of the variance were orthogonal to
$\vec{w}_{opt}$, while only the smallest noise modes contained any overlap the stimulus decoding
direction $\vec{w}_{opt}$ (Figure 3E). This finding demonstrates that the structure underlying trial-to-trial
neuronal variability in the zebrafish visual system lies in a regime closely resembling the
orthogonal variability as illustrated in Figure 3C(i), as also previously observed in a subregion of
mouse primary visual cortex (Rumyantsev et al., 2020). Thus, while even a small degree of
observed overlap between sensory and noise modes can in principle limit the information

encoding capacity of the visually-evoked neurons (Moreno-Bote et al., 2014), this orthogonal
structure minimizes this limitation and is consistent with our previous finding that the stimuli
presented in this study are highly decodable from the neuronal population dynamics (Figure
3B). The orthogonal relationship between sensory and noise modes can be visualized by
projecting average neural activity within each trial onto the stimulus decoding direction $\vec{w}_{opt}$
and the noise modes (Figure 3F).

We next mapped these neural coefficient vectors defining the stimulus decoding direction and
the noise modes onto their anatomical locations of the larvae's brain. As expected, the visually-
evoked neurons were highly concentrated within the optic tectum, such that both the stimulus
decoding direction (Figure 3G) and noise modes (Figure 3H) highly overlapped. The significance
of each neuron's contribution to $\vec{w}_{opt}$ and $\vec{e}_\alpha$ was assessed by comparing its coefficient to a
distribution of coefficients derived from trial-shuffled data (see Methods for details). Overall, 30
$\pm$ 6% of visually-evoked neurons that significantly contributed to the largest noise mode $\vec{e}_1$ also
exhibited significant contributions to the stimulus decoding direction $\vec{w}_{opt}$ (mean $\pm$ 95% CI
across n=10 larvae). Therefore, while many visually-evoked neurons have response components
along both $\vec{e}_\alpha$ and $\vec{w}_{opt}$, the neuronal population dynamics are highly structured such that
activity patterns encoding external visual stimuli are reliable and largely independent from any
trial-to-trial noise modes.

Given that these noise modes represented highly-structured neuronal variability across trials,
we next asked whether they were at all related to the larvae's behavior. First, we asked
whether any noise modes were correlated with the vigor of the larvae's tail movements,
defined as the absolute tail curvature time series convolved with the GCaMP response kernel
(see Methods for details). We found that the largest noise modes were correlated with the
larvae's instantaneous tail vigor (Figure 3I). While we identified the noise modes within the
visually-evoked neuronal population, these noise modes were also highly correlated with the
largest principal component (PC) of the whole-brain dynamics (Figure 3J). This indicates that
both the observed trial-to-trial neuronal variability in visual regions and elsewhere across the
brain are dominated by behavior-related activity (Figure S3C). In fact, 35 $\pm$ 15% of neurons with
significant coefficients in the largest visual noise mode $\vec{e}_1$ also significantly contributed to the
first brain-wide PC (mean $\pm$ 95% CI across n=10 larvae, see Methods for details). While the
lower variance noise modes did not exhibit clear motor-related activity, we suspected they may
be involved in additional internal computations that may modulate decision making across
trials. Taken together, these results show that visual stimuli are faithfully represented on the
population level, and that the observed behavioral variability is not driven by changes in the
fidelity of sensory encoding across trials. Instead, behavioral variability may be explained by
additional, orthogonal dimensions of neuronal activity either within the visually-evoked
population or elsewhere in the brain that are stimulus-independent but predictive of the
larvae's behavior.

Pre-motor neuronal populations predictive of single-trial behavior
436

We hypothesized that there exists a pre-motor neuronal population that would be predictive of
the larvae's behavioral response in any given trial *before* movement initiation. As such, we
aimed to identify any neural locus predictive of the larvae's turn direction at various timepoints
through each visual stimulus presentation. Thus, we looked beyond only the visually-evoked
population and returned to the whole-brain activity patterns. In order to build predictive
models from the neuronal dynamics despite varying reaction time across trials, we applied
time-warping to the neuronal timeseries within each trial such that the stimulus onset and
movement initiation timepoints were aligned across trials (Figure 4A, see Methods for details).
We selected trials in which the larvae had a reaction time of at least one second from the
stimulus onset (providing sufficient pre-motor neuronal data) and made its first tail movement
with a defined minimum tail curvature to either the left or right, so as to separate clear left and
right turns from simple forward locomotion. Further, we utilized trials from all visual object
sizes to collect enough trials to reliably train models and thus identify populations predictive of
decisions regardless of incoming sensory information. We found that there appeared to be
neurons with differing time-warped activity patterns during left and right turn trials (Figure 4A,
bottom). Additionally, we extracted trials in which the larva did not respond following the
stimulus presentation, which were time-warped according to a randomly selected reaction time
from the distribution in responsive trials. As such, each trial could be categorized according to
the stimulus presented, the larva's responsiveness (response/no response), and its turn
direction (left/right) within responsive trials.
457
We first asked whether the larvae's responsiveness and turn direction could be accurately
classified from the whole-brain neuronal dynamics within single trials, as well as how this
predictability behaved over the pre-motor period. To identify the timepoints throughout the
trial within which these could be accurately classified, we varied the time window of neuronal
dynamics used to build a binary logistic regression classification model (Figure S4A). Neuronal
activity across all neurons was integrated from the window start to the window end separately
for each trial and then a model was trained on 80% of the trials. To assess the prediction
accuracy at various timepoints throughout the trial, we utilized the mean F score across larvae
on held-out trials, which measures the harmonic mean of the precision and recall. As expected,
the multiclass stimulus prediction accuracy peaked shortly after the stimulus onset and
remained high throughout the pre-motor and decision periods (Figure 4B). For both the
responsiveness (Figure 4C) and the turn direction (Figure 4D(i)), the mean F score was
consistent across the pre-stimulus and stimulus periods, before quickly ramping near the
movement initiation timepoint. Interestingly, the classification performance was significantly
higher than shuffled data at all timepoints (Figure 4C-D), including before the start of the
stimulus presentation, with an average accuracy of $77.4 \pm 4.4\%$ (mean $\pm$ 95% CI across n=10
larvae) during the pre-motor period (Figure 4D(i)). These results suggest there are two
dominant timescales which contribute to the larva's behavioral response: a longer-timescale
and relatively weak (but significant) activity pattern that does not appear aligned to stimulus or
behavioral events, which we term the pre-stimulus turn bias; and a fast-timescale dramatic
increase in predictability very near the decision, which we term the movement initiation
dynamics.
480

Given that we observed the largest degree of behavioral variability in the responses to small-to-
intermediate size stimuli (Figure 2G), we hypothesized that the pre-stimulus turn bias would be
most influential (and thus most predictive of behavior) during such stimulus presentations, and
less so during presentations of larger stimuli. Indeed, while the mean F scores for turn direction
prediction were highly variable across larvae, on average the largest stimuli exhibited the
lowest predictability from the pre-stimulus turn bias signal (Figure S4B, blue line). This likely
reflects a more salient stimulus drive during large stimulus trials, which in the natural
environment could reflect predators and a potential life-or-death decision for the larvae.
However, significant predictability was observed across all stimulus sizes during the movement
initiation dynamics (Figure S4B, orange line). We thus argue that the pre-stimulus turn bias is
subsequently combined with incoming stimulus information during action selection. Ultimately,
the larva's decision is then signaled by the dramatic increase in predictability during the
movement initiation dynamics.
494
These single-trial predictions could be visualized by projecting the brain-wide neuronal activity
during each trial onto the neural dimensions that optimally classified turn direction and
responsiveness. Consistent with our classification performance, we found that the trajectories
during single trials (Figure 4E) were highly separated according to the ultimate behavioral
response. On average and across larvae, these trajectories remained localized in distinct
quadrants throughout the entire trial (Figure 4F), including before and during the stimulus
presentation period. This suggests that the observed behavioral variability is due in part to the
initial internal state of the neural dynamics at the time of stimulus presentation and decision.
Indeed, by removing the time-warping and observing the dynamics during single trials in real
time, left and right turn trials appear separated for many seconds prior to the movement
(longer than the 3 second visual trials). Ultimately, the decision to turn during the movement
initiation dynamics appeared driven by a strong ramping activity approximately one second
before the turn (Figure 4G). Thus, the identified neuronal dimensions exhibited reliable
decoding within single trials driven by an initial bias and ultimately resulting in the observed
movement initiation dynamics.
510
Next, we asked which specific neuronal populations across the brain contributed to the
accurate pre-motor classification of behaviors, and thus could be involved in driving the
decision of when and how to act. We considered a neuron to be significantly contributing to the
behavioral prediction if its distribution of absolute coefficients across 6-fold cross-validation
was significantly higher (one-tailed t-test, $p < 0.05$) when fit to real behavioral data as opposed
to a null distribution created by randomly permuting the trial type labels. We found that both
the ensemble of neurons which significantly contributed to the turn direction prediction
(Figures 4H-I) and the responsiveness ensemble (Figures 4J-K) were highly distributed across
the brain. Both of these distinct neuronal ensembles showed no significant difference in their
distribution across the four major brain regions: the telencephalon (Tel), optic tectum (Tec),
cerebellum (Cer), and hindbrain (Hind). Further, while this population included visually-evoked
neurons within the optic tectum, we found that our previously identified noise modes $\vec{e}_\alpha$ were
similarly predictive of single trial turn direction (Figure S4C), whereas the optimal stimulus
decoding direction $\vec{w}_{opt}$ (Figure 3G) was not able to predict single-trial turn direction (Figure

S4D). Thus, our data suggest that the neural dynamics underlying single-trial action selection
are the result of a widely-distributed circuit that contains subpopulations encoding internal
time-varying biases related to both the larva's responsiveness and turn direction, yet distinct
from the sensory encoding circuitry.
529
What could be the origin of these trial-to-trial biases in the larvae's behavior? Given that our
results demonstrate that it is possible to predict a behavioral bias before the visual stimulus is
shown, we hypothesized these results could reflect the influence of the zebrafish neuronal
circuitry for spontaneous turns, which includes the hindbrain oscillator (or anterior
rhombencephalic turning region, ARTR) known to bias sequential turns towards a given
direction (Dunn et al. 2016). While our pre-motor neuronal population is not exclusively
localized to the ARTR, we did identify many neurons in the cerebellum and hindbrain (Figures
4J-K) whose locations were consistent with these known populations. As such, we asked
whether spontaneous turns could be predicted from this same turn direction neuronal
ensemble (Figure 5A). To address this question, we selected spontaneous turns that occurred
either during inter-trial intervals or within a 2-minute period at the beginning of each recording
without any stimulus presentations, utilizing the same criteria to define left and right turns as
previously (see Methods for details). We then asked whether the spontaneous turn direction
during periods without sensory stimulation could be predicted from the same turn direction
neuronal ensemble and coefficients previously identified in Figure 4H-I. We found a significant
correlation between the pre-motor turn direction predictions and the observed spontaneous
tail curvature across larvae (Figure 5B, r=0.41, p<0.05), suggesting this same ensemble is
activated prior to both spontaneous and visually-evoked turns. This ensemble had a pre-motor
turn direction classification accuracy of 70.2 $\pm$ 6.0% (mean $\pm$ 95% CI across n=5 larvae, Figure
5C). Thus, these results demonstrate that a portion of the observed behavioral variability is
overlapping with the larva's motor circuitry responsible for generation of spontaneous
behavior.
552
Our data highlight that the neural mechanisms involved in single-trial decision making are
reflected in a highly distributed ensemble of neurons across the larval zebrafish brain. Further,
behavioral variability, particularly in the larvae's responses to small-to-intermediate size stimuli,
can be partially explained by an internal, task-independent turn bias that is unrelated to the
visual stimulation, while their stimulus-driven responses to larger predator-like stimuli
exhibited a comparatively lower level of behavioral variability. This suggests the importance of
considering how internal biases and circuits governing spontaneous movements interact with
the well-studied sensorimotor circuits to govern how the brain generates behavior in a
naturalistic context.
562
563   **DISCUSSION**
564
In this study, we designed and built an optimized, high-resolution Fourier light field microscope
(fLFM) to perform whole-brain imaging of the larval zebrafish in order to investigate the nature
of trial-to-trial variability in neuronal dynamics and behavior on whole-brain yet cellular level.
To do so, we studied visually-evoked turns within two ethologically relevant behaviors with

opposing valance, prey capture and escape response, each of which are on average driven by
distinct pre-motor neuronal ensembles (Bollmann, 2019) dependent on the size of the given
sensory stimulus. Consistent with previous results (Barker and Baier, 2015), we found that the
larvae's behavioral responses were highly variable across presentations of the same stimulus,
with behavioral variability peaking in response to stimulus of an intermediate size between
those which optimally drive attraction or avoidance behaviors. Given that we lack a mechanistic
understanding of which specific neuronal circuits drive the observed variance across trials, we
utilized fLFM to screen whole-brain dynamics at cellular resolution and identify neuronal
ensembles which contribute to behavioral variability at the single-trial level.
578
We first asked whether the observed behavioral variability could be explained by noise or trial-
to-trial deviations in encoding of the visual stimuli. We found that despite the variable
responses of individual neurons to visual stimuli across trials, the population response faithfully
and reliably encoded the visual stimulus in each trial. Further, we discovered that the visually-
evoked neuronal activity patterns were largely orthogonal to those dimensions that maximally
covaried across stimulus repetitions, i.e. the noise modes, indicating that the source of
behavioral variability was not unfaithful representation of the sensory inputs. Instead, we found
that at least a third of the visually-evoked neurons which contributed to noise modes were
related to motor behavior. We ultimately identified two brain-wide neuronal populations which
could predict the larvae's turn direction and responsiveness at the single-trial level. Surprisingly,
a pre-stimulus bias component of these neuronal population activity could predict the larvae's
trial-by-trial turn direction even before the onset of the stimulus presentation with an average
accuracy of 77%. while after stimulus onset, a sharp ramping of the activity of this neuronal
population approximately one second before movement initiation allowed for an increased
turn direction prediction accuracy of 90%. Taken together, our data show that the larva's trial-
by-trial decisions in response to any given stimulus are partially explained by pre-stimulus turn
biases, and then ultimately driven by movement initiation dynamics which are broadcast within
brain-wide neuronal ensembles.
597
In this context, the design of our behavioral paradigm has allowed us to gain insights into the
nature of trial-to-trail neuronal and behavioral variability within and across two different
ethologically relevant but highly divergent behaviors by tuning only a single parameter. This has
allowed us to show that functionally different neuronal circuits have evolved to exhibit
different levels of variability. Further, the differential interaction of such neuronal variability
with the well-studied sensorimotor circuits may govern how the brain generates optimal
behavior in different naturalistic contexts.
605
In addition, our study represents the first to our knowledge that identifies behavior-related
noise correlations throughout an entire brain-wide sensorimotor decision-making circuit. The
nearly orthogonal relationship between the stimulus decoding direction and trial-to-trial noise
correlations is highly consistent with previous studies which found that noise correlations
within single brain regions are organized so as to prevent additional information encoding
channels within the same population from limiting sensory encoding capacity (Kafashan et al.,
2021; Rumyantsev et al., 2020). However, even a small degree of overlap between noise and

sensory dimensions may decrease the information encoding capacity of the population
(Moreno-Bote et al., 2014). Why then might the brain favor a coding scheme in which
populations mix information encoding with additional noise correlation structure?
616
Our data offer one possibility: that such additional correlations could encode internal or
behavioral states that modulate sensorimotor transformations and behavioral output, enabling
dynamic and flexible behavioral responses. Indeed, we found that a large fraction of the
observed trial-to-trial neuronal variability was related to the larva's behavior: including activity
correlated the instantaneous behavioral state of the animal as well as pre-stimulus turn biases
that correlated with responsiveness and turn direction. We found that the largest trial-to-trial
noise mode within the visually-evoked neurons was a subset of a brain-wide population
correlated with the vigor of the larva's tail movements. This is conceptually similar to recent
evidence in rodents that a large fraction of spontaneous neural activity reflects spontaneous
and uninstructed movements (Musall et al., 2019; Stringer et al., 2019).
627
Taken together, our study represents the first whole-brain confirmation that behaviorally
relevant information is highly mixed throughout neuronal populations involved in processing
sensory and other information, potentially enabling flexible and context-dependent behavior.
Our observation of time-varying neuronal ensembles encoding a turn direction bias is
reminiscent of the larval zebrafish's spontaneous swimming circuitry, which involves a central
pattern generator (the ARTR) that biases the larva to swim in chains of turns in alternating
directions (Timothy W Dunn et al., 2016); however, the neuronal populations we identified
were not localized to the previously identified circuitry, and instead were distributed across the
brain. While such spontaneous turn sequences are thought to underlie efficient exploration of
local environments, our data could reflect a similar sensorimotor circuitry that injects variability
as a learning mechanism to explore the space of possible sensorimotor consequences. Given
the brain-wide distribution of the behavior-related ensembles within the variable visually-
evoked behaviors studied here, we expect that they are not driven solely by a single central
pattern generator as with ARTR and spontaneous turns, but instead involve the interaction of
such pattern generators with the larva's internal state and recent experience. It has been
shown that such hindbrain oscillatory activity integrates recent visual to enable efficient
phototaxis (Wolf et al., 2017), suggesting these pattern generators could act in concert with
additional contextual populations. For example, it has been shown that hunger shifts decisions
from avoidance to approach due to neuromodulation from the hypothalamic and serotonergic
systems (Filosa et al., 2016). Further, our observed dynamics could also reflect the foraging
state of the larva, which is known to be encoded by an oscillatory, neuromodulatory network
distributed across the brain (Marques et al., 2020) and is a likely candidate to modulate a
larva's response to prey-sized stimuli. We envision that utilization of the wide array of
genetically labelled lines that have been associated with these sensorimotor and
neuromodulatory circuits (Semmelhack et al., 2014; Barker and Baier, 2015; Marques et al.,
2020; Okamoto et al., 2021) could disentangle the specific contributions of the sensory, pre-
motor, and internal state circuitry within the brain-wide populations observed here. Of
particular interest will be how the activity of these populations is organized over longer
timescales and modulated by visual input or internal states related to hunger and foraging,

which are critical to balance feeding behaviors and predator avoidance within the developing
larvae.
659
660

661 **ACKNOWLEDGEMENTS**
662
We thank Q. Lin and S. Otero Coronel for discussions and feedback on the manuscript, T.
Nöbauer and Y. Zhang for discussions regarding optical designs, and the members of the Vaziri
Lab for discussions regarding the experiments and data analysis. We also thank A. Kaczynska, S.
Campbell, and J. Hudspeth for housing zebrafish, and the Rockefeller University High
Performance Computing Cluster for compute access. This work was supported in part by the
National Institute of Neurological Disorders and Stroke of the National Institutes of Health
under award numbers 1RF1NS113251 and 1RF1NS110501 and the Kavli Foundation through the
Kavli Neural Systems Institute.

671
672

673 **AUTHOR CONTRIBUTIONS**
674
675 J.M. contributed to the project conceptualization, designed and built the imaging system,
676 performed experiments, analyzed data, and wrote the manuscript. A.V. conceived, led and
677 supervised the project, designed the imaging system and biological experiments, guided data
678 analysis, and wrote the manuscript.

679
680
681 **FIGURE CAPTIONS**
682
683 **Figure 1: Fourier Light Field Microscopy (fLFM) provides a simple and cost-effective method**
684 **for whole-brain imaging of larval zebrafish during behavior**
685
686 **A.** Schematic of the fLFM system. The sample is illuminated with a 470 nm LED through a
687 20×/1.0-NA imaging objective (Obj). The fluorescence is sent to the imaging path via a dichroic
688 mirror (DM). An Olympus-style f=180mm tube lens (TL) and f=180mm Fourier lens (FL) are used
689 to conjugate the back focal plane of the objective onto a microlens array (MLA). An sCMOS
690 sensor is positioned at the focal plane of the MLA to capture the raw fLFM images.
691
692 **B.** An example raw sensor image. Each lenslet in the 8×8 array forms an image of the sample
693 from a slightly different perspective, allowing for reconstruction of the 3D volume.
694
695 **C.** Experimental measurement of the point spread function (PSF). Left: the x-y (top) and x-z
696 (bottom) profiles of a reconstructed image of a 1 μm fluorescent bead. Right: corresponding
697 cross sections of the PSF (points). A Gaussian profile (lines) was fit to these data to measure the
698 full width at half maximum (FWHM), which was 3.3 μm laterally and 5.4 μm axially.
699
700 **D.** Schematic of our fLFM data processing pipeline (see Methods for detailed description).

702 **E.** A maximum intensity projection (MIP) of a conventional LFM (cLFM) larval zebrafish
703 recording resulting in strong grid-like artifacts and low resolution near the native image plane
704 (black arrows). A reconstructed volume of $750 \times 750 \times 200$ μm^3 is shown.
705

706 **F.** An MIP of a fLFM recording. fLFM exhibits higher axial resolution and does not contain
707 artifacts at the native image plane. A reconstructed volume of $750 \times 375 \times 200$ μm^3 is shown.
708

709 **G.** A heatmap of extracted neuronal activity from an example one hour recording of 15,286
710 neurons. Neurons are sorted using the rastermap algorithm (Stringer et al., 2019), such that
711 nearby neurons exhibit similar temporal activity patterns. Multiple distinct activity patterns are
712 seen, included strong activity during two sets of drifting grating-induced optomotor response
713 trials (black arrows).
714

715 **H.** Example neurons tuned to tail movements and visual stimuli. In black are six example neuron
716 traces from the designated region in panel G, which exhibited correlations with the GCaMP-
717 kernel-convolved tail vigor (top trace, see Methods for details). In red are six example neuron
718 traces from the designated region in panel G, which exhibited correlations with the visual
719 stimuli presented in the left visual field (denoted by the red lines at the bottom of the plot).
720

721 **Figure 2: Zebrafish exhibit highly variable motor responses to visual stimuli**
722

723 **A.** Schematic of the freely-behaving experimental setup. A single larva is placed in a 90 mm
724 petri dish with a shallow (approximately 3 mm) amount of water. The dish is placed on a screen
725 which displays visual stimuli consisting of dots of various sizes drifting in a constant direction
726 and speed. The behavior is monitored with a camera from above.
727

728 **B.** An example target-avoidance bout. A composite image taken of four frames from start
729 (white arrow) to near the finish (black arrow) of a bout in which the larvae avoided a large
730 visual stimulus. See also Video 1.
731

732 **C.** An example target-directed bout. A composite image taken of three frames from start (white
733 arrow) to finish (black arrow) of a bout in which the larvae made a directed movement toward
734 a small visual stimulus. See also Video 1.
735

D. The behavioral response of larvae to various size visual stimuli. For each size stimulus, all
bouts of movement in which the larva was within 10 body lengths of a visual stimulus were
considered. For each bout, the change in distance between the larvae and the stimulus during
the movement was monitored. Thus, target-directed bouts exhibited negative values
corresponding to a decrease in the distance to the stimulus, whereas target-avoidance bouts
exhibited positive values. Stimuli less than 7° in size evoked target-directed responses on
average, whereas stimulus greater than 10° evoked target-avoidance responses. Shown is the
mean $\pm$ standard deviation of n=9 larvae.
744

E. Schematic of the head-fixed experimental setup. A larva is embedding in agarose in order to keep its head fixed during whole-brain imaging. The agarose around the tail and eyes are removed, such that the tail can be tracked (indicated by black dots along the tail) and visual stimuli can be presented. Visual stimuli of various sizes are presented moving from the center of visual field to either the left or right.

F. The behavioral response of larvae to various size visual stimuli during head fixation. Visual stimuli are presented in either the left (pink) or right (green) visual field. The directedness of tail movements is monitored by computing the mean tail curvature during a bout of movement, with positive values indicating leftward motions. Similar to freely-behaving experiments, visual stimuli of 7° or less evoked target-directed responses, whereas stimuli larger than 10° evoked target-avoidance. Shown is the mean $\pm$ standard deviation of n=10 larvae. Asterisks indicate stimuli with significant difference between presentations on the left and right visual field ($p < 0.05$, paired t-test).

G. Example behavioral responses to various stimuli. For each of four stimulus sizes on the left and right visual fields, a histogram of the mean tail curvature during bouts is shown for an example larva. While stimulus-evoked behavioral responses are variable in all cases, they appear the least variable in the case of largely target-avoidance bouts to large 44° stimuli (rightmost column).

Figure 3: Trial-to-trial variability in visually-evoked neurons is largely orthogonal to visual decoding dimensions

A. Individual visually tuned neurons exhibit trial-to-trial variability in their responses to stimuli. For each of the six visual stimuli (three object sizes on both the left and right visual fields), the two neurons exhibiting the highest correlation with the stimulus kernel (see Methods for details) are shown for an example larva. Each column represents a neuron, and each line represents its response to a given stimulus during a single trial.

B. Visual stimuli are reliably decodable from whole-brain dynamics on the single trial level. The visual stimuli were decoded using a logistic regression classifier with lasso regularization (see Methods for details). For each larva, a confusion matrix is computed for the test trials during 6-fold cross-validation. Shown is the average confusion matrix across n=8 larvae which were shown six visual stimuli.

C. Schematic of potential geometric relationships between sensory decoding and neural variability dimensions. In each plot, each dot represents the neural response during a single presentation of stimulus A or B. The decision boundary for an optimal classifier is denoted with a dashed line, and the optimal stimulus decoding direction is denoted by the vector $\vec{w}_{opt}$. The direction representing the maximal trial-to-trial variance is denoted by $\vec{e}_1$, and can be calculated by finding the first eigenvector of the noise covariance matrix (see Methods for details). These vectors can be: (i) orthogonal, such that neuronal variability does not limit the

788 stimulus decoding; (ii) show little relationship, for example in the case of uniform variability; or
789 (iii) aligned, such that variability likely limits the information encoding capacity along $\vec{w}_{opt}$.
790
D. The trial-to-trial noise correlation matrix appears multi-dimensional. Shown is the average
noise correlation matrix across all stimulus types presented. The neurons are sorted using
rastermap, which produces a one-dimensional embedding of the neurons, such that neurons
which show similar correlation profiles are placed near to one another. A number of neuronal
populations exhibiting correlations across trials are apparent from the clusters of high
correlations near the diagonal.
797
E. Trial-to-trial variability in the visually-evoked neurons is largely orthogonal to visual decoding
dimensions. The fraction of trial-to-trial variance explained by each noise mode $\vec{e}_\alpha$ is plotting
against the angle between $\vec{e}_\alpha$ and the optimal stimulus decoding direction $\vec{w}_{opt}$. Shown are the
$\vec{e}_\alpha$ for n=8 larvae, colored by their rank order α based on the fraction of variance explained. The
largest noise modes were approximately orthogonal (~90°) to the stimulus decoding direction,
whereas only a few of the smallest noise modes exhibited angles less than 90°.
804
F. Example projections of single trial neural activity along the stimulus decoding and noise
dimensions. Each dot represents the average neural activity within a single trial projected along
$\vec{w}_{opt}$ and $\vec{e}_1$ for an example larva. Each of the six visual stimuli, three object sizes presented on
either the right (green) or left (pink) visual field, are robustly encoded along $\vec{w}_{opt}$ across trials;
however, in all stimuli there is strong orthogonal variability along $\vec{e}_1$, the largest noise mode
representing 39.5% of the trial-to-trial variance.
811
812 **G.** Example neuron coefficients for $\vec{w}_{opt}$. Shown are the 366 neurons with the largest weights
813 over a maximum intensity projection of the recorded volume. The visually-evoked neurons
814 which encode stimulus information are concentrated within the optic tectum.
815
816 **H.** Example neuron coefficients for $\vec{e}_1$. Shown are the 828 neurons with the largest weights over
817 a maximum intensity projection of the recorded volume. The visually-evoked neurons
818 contributing to the largest noise mode are highly overlapping with the neurons contributing to
819 $\vec{w}_{opt}$ in panel G.
820
821 **I.** The neural noise modes are highly correlated with tail movements. Shown are both the
822 GCaMP kernel-convolved tail vigor and the neuronal projection onto the first noise mode $\vec{e}_1$ for
823 a representative larva. Over the full two-hour recording, the tail vigor and noise mode
824 projection exhibit a significant correlation of r=0.58, p<10^{-6}.
825
826 **J.** The largest neural noise modes reflect brain-wide motor encoding. Shown are the
827 correlations between the first noise mode $\vec{e}_1$ and the tail vigor (blue) or the first principal
828 component (PC 1, orange) of whole-brain data (see Figure S3C). Dots show data from individual
829 larvae, whereas the violin plots below show the null distribution for temporally shuffled data, in
830 which the tail vigor or PC 1 are circularly permuted.

Supplementary Figure S3, related to Figure 3, trial-to-trial variability in visually-evoked neurons is largely orthogonal to visual decoding dimensions

A. Schematic of trial-to-trial noise mode identification. First, dimensionality reduction is applied to the whole-brain dynamics (left) to identify the neuronal activity patterns that optimally preserve visual information using partial least squares regression. Within these visually-evoked neurons, each trial's average activity during the visual presentation is computed (middle, each dot represents a single trial corresponding to the pink or green stimulus condition). Finally, the covariance across trials is decomposed to find the noise modes $\vec{e}_\alpha$, which represent orthogonal directions within the visually-evoked neurons and ranked in order of decreasing variance.

B. $\vec{w}_{opt}$ robustly encodes the visual stimulus. Shown are example time traces of the stimulus kernel (blue, where the size of the stimulus is encoded by the magnitude and which visual field it is presented in is encoded by the sign) and the neural projection onto $\vec{w}_{opt}$ (orange), which exhibit a correlation of r=0.88, $p<10^{-6}$.

C. The largest brain-wide principal component (PC) encodes motor behavior. (i) Example time traces of the tail vigor (blue) and PC 1 (orange), which exhibit a correlation of r=0.51, $p<10^{-6}$. (ii) The coefficients of the neurons contributing most to the PC, which are distributed across the brain and are nearly all positively correlated with tail vigor.

Figure 4: Pre-motor neuronal populations predictive of single-trial behavior.

A. Trials are time-warped to align stimulus and decision onsets before classifying the turn direction. Top: Time-warped tail curvatures for trials in which the fish performed leftward or rightward turns, on the left and right, respectively. Bottom: Trial-averaged and time-warped neuronal timeseries for 15,286 neurons during left and right turns. The neurons are sorted using the rastermap algorithm. A time window is swept across the stimulus and decision timepoints to train binary classification models to predict the turn direction from the associated neuronal dynamics.

B. Stimulus classification accuracy peaks after the onset of visual stimulation. The mean F score across n=7 larvae is used to assess the performance of 6-way multiclass classification of the presented visual stimulus as a function of warped time surrounding the stimulus onset (Stim.) and decision timepoint (Dec.). Shown is the mean $\pm$ 95% confidence interval of the F score for the best time window ending at the given timepoint (Data), compared to shuffled data in which the class labels are randomized. The black bar at the bottom indicates timepoints where the data have a significantly higher F score than the shuffled data (p<0.05, paired t-test).

C. Binary classification of responsiveness, whether or not the fish responds in a given trial, is significant throughout all time periods but accuracy peaks near movement initiation. As in panel B, except for binary classification of responsiveness. Nonresponsive trials are time-

warped by randomly selecting a reaction time from the response trials and applying the same
transformation.

D. (i) Turn direction classification accuracy is significantly higher than shuffled data across the
entire time-warped interval, but peaks near movement initiation. As in panel B, except for
binary classification of turn direction. **(ii)** Single trial classification of turn direction across
larvae. The mean confusion matrix across n=7 larvae, which show an accuracy of $77 \pm 4\%$
(mean $\pm$ 95% confidence interval).

E. Single trial trajectories are separated based on responsiveness and turn direction. Shown are
neural activity trajectories during single trials in an example larva projected onto the brain-wide
neural dimensions that optimally separated turn direction and responsiveness.

F. Consistent trial-averaged trajectories across larvae. As in panel F, except for the trial-
averaged responses for n=6 example larvae. For the one bold animal, timepoints across the trial
are indicated by a circle for trial start, diamond for the decision timepoint, and an X for the trial
end.

G. Real-time single-trial dynamics in an example larva. Along the turn direction neural
projection, left and right trials are separated for many seconds before the decision timepoint,
which is longer than the three second length of visual presentations. Activity along this
dimension shows consistent ramping across trials approximately one second before movement.

H. Example turn direction neuronal ensemble. Shown are the coefficients for all neurons which
showed significantly higher (one-tailed t-test, $p < 0.05$) absolute coefficients in the real models
compared to shuffled data in which the turn direction labels are randomly permuted.

I. Highly distributed encoding of turn direction across larvae. The percentage of significant turn
direction neurons located with the four major brain regions (Tel – telencephalon, Tec – optic
tectum, Cer – cerebellum, and Hind – hindbrain) are shown for n=10 larvae. There is no
significant difference between the percentage of neurons across brain regions ($p > 0.05$, paired t-
test).

J. Example responsiveness neuronal ensemble. As in panel I, except for responsiveness. Shown
are the coefficients for all neurons which showed significantly higher (one-tailed t-test, $p < 0.05$)
absolute coefficients in the real models compared to shuffled data in which the turn direction
labels are randomly permuted.

K. Highly distributed encoding of responsiveness across larvae. As in panel J, except for
responsiveness. The percentage of significant turn direction neurons located with the four
major brain regions (Tel – telencephalon, Tec – optic tectum, Cer – cerebellum, and Hind –
hindbrain) are shown for n=10 larvae. There is no significant difference between the percentage
of neurons across brain regions ($p > 0.05$, paired t-test).

Supplementary Figure S4: related to Figure 4, pre-motor neuronal populations predictive of single-trial behavior.

A. Classification of turn direction as a function of time window analyzed. The mean F score across n=7 larvae is used to assess the performance of multi-class classification of the visual stimulus. Top: a time window is swept across various start timepoints (y-axis) and end timepoints (x-axis) to determine the classification performance across various time intervals of the neuronal dynamics. Bottom: Mean $\pm$ 95% confidence interval of the F score for the best time interval ending at the given timepoint (Data), compared to shuffled data in which the stimulus labels are randomized. The classification accuracy is significantly higher than shuffled data ($p<0.05$, paired t-test) only after stimulus onset.

B. Turn direction classification as a function of stimulus size. On average before stimulus onset (blue line), the F score is highest for the smallest size stimuli, whereas the F score is relatively constant at the time of movement initiation (orange line, mean $\pm$ 95% CI across n=5 larvae).

C. As in A, except for classifying left or right turn direction from the projections onto the trial-to-trial noise modes $\vec{e}_\alpha$, as opposed to the whole-brain dynamics as used in Figure 4B.

D. As in A, except for classifying left or right turn direction from the projection onto the optimal stimulus decoding dimension $\vec{w}_{opt}$.

Figure 5: Spontaneous turns are predictable from the same pre-motor neuronal population

A. Schematic of the approach to predict spontaneous turns. Models are fit to predict left or right turn direction during visual trials (top), as in Figure 4. They are then tested on pre-motor periods one second before the spontaneous turn.

B. Spontaneous turns are predicted from visual-evoked model. Shown is the relationship between the spontaneous tail curvature and the predicted pre-motor turn direction using the visual-evoked model. Each dot is a single spontaneous turn and each color represents a different larva. They exhibit a significant correlation of r=0.41, $p<0.05$.

C. Spontaneous turn classification accuracy. The mean cross-validated confusion matrix for spontaneous turn classification over n=5 larvae. Spontaneous turns are predicted with an accuracy of 70.2 $\pm$ 6.0% (mean $\pm$ 95% CI across n=5 larvae).

Video 1: related to Figure 2. Example target-directed and target-avoidance bouts in free behavior.

On the left, an example target-directed bout showing a 7dpf larval zebrafish approaching a stimulus presented on a screen below the dish (see Figure 2A). On the right, a similar target-avoidance bout from the same larva. The videos are slowed down 10x. These videos correspond to the composite images shown in Figure 2B-C.

962
963
964 **METHODS**
965
966 **fLFM setup**
967
The sample is illuminated with an LED excitation light (470 nm, Thorlabs M470L4) via Köhler
illumination through a standard GFP filter set (Thorlabs MDF-GFP) and a 20×/1.0-NA water
immersion objective (Olympus XLUMPLFLN). In order to reduce the amount of excitation light
reaching the eyes of the zebrafish, an aluminum mask is placed in the Köhler illumination path
conjugate to the sample plane. The imaging path consisted of a f=180 mm Olympus-style tube
lens (Thorlabs TTL180-A), a f=180 mm achromatic doublet (Thorlabs AC508-180-A-ML) for the
Fourier lens, and a 13×13 mm microlens array with 1.5 mm pitch and r=13.9 mm radius of
curvature (OKO Tech APO-GB-P1500-R13.9). The microlens array is mounted inside a five-axis
kinematic mount (Thorlabs) to allow precise placement of the array orthogonal to the optical
axis. A 5.5-megapixel sCMOS camera (Andor Zyla) is then positioned at the focus of the
microlens array to capture the fLFM images.
979
The theoretical lateral resolution of an fLFM Is limited by the numerical aperture NA_{ML} of the
microlenses and the diffraction-limited spot size at the sensor. For an emission wavelength λ,
Fourier lens focal length f_{FL}, objective magnification of M, and microlens diameter d_{MLA}, the
Abbe diffraction limit when converted to object space and under the paraxial approximation is
given by $R_{xy} = \frac{\lambda f_{FL}}{d_{MLA}M}$. The theoretical axial resolution for a microlens array with a radius d_{max}
is given by $R_z = \frac{d_{MLA}R_{xy}^2}{\lambda d_{max}}$ (Guo et al., 2019).
986
987 **Zebrafish experiments**
988
For head-fixed zebrafish experiments, n=13 *huc:h2b-gcamp6s* larvae with pan-neuronal and
nuclear-localized GCaMP6s (NL-GCaMP6s) expression were imaged 6-9 days post fertilization
(dpf). The day before experiments were performed, we immobilized larvae by embedding them
in 2.5% agarose approximately 10 mm from the edge of a 35mm petri dish. We then removed
the agarose around their eyes and tail to allow for visual stimulation and tail movement. Larvae
were not fed after immobilization and were thus in a starved state on the day of the
experiment.
996
The petri dish was covered in a rear projection film (Screen Solutions Int.) such that visual
stimuli could be projected directly onto the dish. The projector (BenQ TH671ST) output was
filtered with a 610 nm longpass filter (Schott RG610) such that only red light was displayed,
removing crosstalk between the stimulation and GCaMP fluorescence. The tail was illuminated
with an 850 nm LED and then monitored from below using a near infrared CMOS camera
(Ximea MQ013RG-ON).
1003

Visual stimulation was controlled by Stytra (Štih et al., 2019). To check for intact visual circuitry,
drifting gratings with a period of 40° (visual angle) and speed of 40°/s to the left or right were
shown three times every 30 minutes. Only larvae that displayed a robust optomotor response
were further analyzed. The remaining trials consisted of drifting dots that began in the center of
the visual field and moved to either the left or right at 30°/s. Dots were shown at maximum
contrast against a dark background. Various diameters were shown from 0.3° to 44° visual
angle, with 3 to 6 different sizes used in a recording. Each trial lasted 3 seconds and the inter-
trial interval was randomly distributed with a mean of 9 seconds and a standard deviation of 3
seconds.
1013
1014   **Data acquisition**
1015
1016   The fLFM camera was triggered at 5 or 10 Hz using a microcontroller (Adafruit Grand Central).
1017   The first trigger was used to initiate the tail behavior camera. The tail was monitored in real-
1018   time using Stytra (Štih et al., 2019) at approximately 200 Hz. The recording duration lasted 1 to
1019   2 hours.
1020
1021   **Data processing**
1022
Raw fLFM images were denoised using a custom-trained DeepInterpolation (Lecoq et al., 2021)
model. We trained a "unet_single_1024" model with 5 pre and post frames on a training set of
n=20 example recordings and validation set of n=5 recordings. We ran the training until
convergence of the validation loss. Denoised raw fLFM images were then reconstructed using
the Richardson-Lucy deconvolution algorithm. We performed the deconvolution with an
experimental PSF found by measuring the profile of a 1 μm fluorescent bead through the axial
depth in 2 μm increments. The Richardson-Lucy algorithm took about 1 second per iteration on
a GPU (NVIDIA TITAN V) and we used 12 to 20 iterations per frame. This resulted in a
reconstructed volume of approximately $760 \times 360 \times 280$ μm^3 with voxels of 2.4 μm laterally
and 4 μm axially.
1033
Neuronal ROIs and timeseries were then extracted using the CNMF-E (Zhou et al., 2018) variant
of the CaImAn software package (Giovannucci et al., 2019), which is optimized for processing of
one-photon imaging data. Each plane was analyzed individually and then the results were
merged. To determine neuronal ROIs, the SNR threshold was set at 2, the spatial consistency
(rval) threshold was set at 0.8, and the CNN-based classifier threshold was set at 0.9. Neuronal
timeseries were not deconvolved. Background fluorescence was factored out using the CNMF-E
ring model. The planes were then collated by merging any units with a correlation greater than
0.7, lateral distance less than 10 μm, and axial distance less than 16 μm. Overlapping units were
merged by summing each timeseries weighted by its SNR. Effects of photobleaching were
removed by normalizing by a fit to the mean signal $f(t)$ across all neurons over the recording
with a bi-exponential model $\hat{f}(t) = ae^{bt} + ce^{dt}$. Each neuronal timeseries was then z-scored
before all following analyses.
1046
1047   **Analysis of freely swimming behavior**

1048
For freely swimming experiments, n=9 6-9 dpf *huc:h2b-gcamp6s* larvae were individually placed
in a 90 mm petri dish with a shallow (~3 mm) amount of water. The dish was placed above a
display (Apple iPad) with a 6 mm piece of clear acrylic in between. Black dots of sizes ranging
from 1° to 30° were randomly shown at maximum contrast and moving across the dish in one
direction at 30°/s. A camera (FLIR Grasshopper3 GS3-U3-41C6M-C) with a zoom lens (Navitar
MVL7000) was mounted above to monitor the larvae's behavior at >100 Hz.
1055
Each larva's movements were tracked using DeepLabCut (Mathis et al., 2018), which was
trained to track the position of the eyes, the swim bladder, and four points along the spline of
the tail. Anipose was used to build a pipeline to process large numbers of videos (Karashchuk et
al., 2020). Movement bouts were extracted by applying a threshold to the velocity of the
centroid of the tracked points. Only movement bouts in which the centroid was within
approximately six body lengths of the visual stimulus were further analyzed. For each bout, we
then calculated the mean change in distance to the visual stimulus, with positive values
denoting the larvae moving further away from the visual stimulus.
1064
1065 **Analysis of head-fixed behavior**
1066
1067 During head-fixed visual experiments, 20 points along the spline of the larva's tail were tracked
1068 in real time using Stytra (Štih et al., 2019). The overall tail curvature was then quantified by
1069 summing the angles between each segment of the tail tracking. For comparison with neural
1070 data, this was convolved with the NL-GCaMP6s kernel to define the tail direction kernel, while
1071 the GCaMP-kernel-convolved absolute tail curvature defined the tail vigor kernel.
1072
Bouts of movement were extracted by thresholding the absolute tail curvature with a value
σ_{active} set to one standard deviation of the absolute tail curvature over time. To study the
behavioral response to various stimuli, the mean tail curvature during active bouts was
computed for each stimulus presentation period; trials in which there were no movement
bouts were excluded. For turn direction analyses, trials with a mean tail curvature during bouts
$> \sigma_{active}$ were considered as left turn trials, whereas trials with a mean tail curvature during
bouts $< -\sigma_{active}$ were considered right turn trials.
1080
1081 **Visual stimulus decoding**
1082
For the decoding shown in Figure 3B, the stimulus-evoked neural response was taken as the
average response of each neuron during a single stimulus presentation. For each larva, a logistic
regression model with lasso regularization was trained on all trials using 6-fold cross-validation
with 3 repetitions. The lasso regularization parameter, which induces sparsity, was swept and
the model with the highest average cross-validated accuracy was used. The confusion matrix,
computed on the held-out testing trials, was reported as the average over all rounds of cross-
validation.
1090
1091 **Identification of stimulus decoding and noise dimensions**

1092

The comparison of the neural dimensions encoding visual stimuli versus trial-to-trial noise was
modeled after (Rumyantsev et al., 2020). Partial least squares (PLS) regression was used to find
a low-dimensional space that optimally predicted the visual stimuli, which we refer to as the
visually-evoked neuronal activity patterns. To perform regression, a visual stimulus kernel was
constructed by summing the timeseries of each individual stimulus type, weighted by the
stimulus size and negated for trials on the right visual field, thus providing a single response
variable encoding both the location, size, and timing of all the stimulus presentations. This
stimulus kernel was the convolved with the temporal response kernel of our calcium indicator
(NL-GCaMP6s). The dimensionality d of PLS regression was optimized using 6-fold cross-
validation with 3 repeats and choosing the dimensionalities with the lowest cross-validated
mean squared error. The optimal stimulus decoding direction $\vec{w}_{opt}$ was identified by re-fitting
the optimal PLS regression model with all timepoints.

1105

For each stimulus type, the noise covariance matrix C was computed in the low-dimensional
PLS space, given that direct estimation of the noise covariances across many thousands of
neurons would likely be unreliable. A noise covariance matrix was calculated separately for
each stimulus, and then averaged across all stimuli. The noise modes $\vec{e}_\alpha$ for $\alpha = 1 \dots d$ were
identified by eigendecomposition of the mean noise covariance matrix across all stimuli,
$C = \sum \lambda_\alpha \vec{e}_\alpha \vec{e}_\alpha{}^T$. The angle between the optimal stimulus decoding direction and the noise
modes is thus given by $\cos \theta_\alpha = (\vec{w}_{opt} \cdot \vec{e}_\alpha)/(|\vec{w}_{opt}| \, |\vec{e}_\alpha|)$.

1113

The significance of each neurons' contribution to a given neural population vector was
evaluated by comparison to shuffled datasets. To created shuffled datasets, the ground truth
stimulus presentation labels were randomly permuted. The same analysis above was
performed on n=10 shuffled datasets, to identify a distribution of shuffled coefficients for the
optimal stimulus decoding direction, noise modes, and largest principal component. A neuron
was then considered to be significantly contributing to a given population vector if its absolute
coefficient was greater than three standard deviations above the mean of that of the shuffled
datasets.

1122

1123   **Classification of single trial turn direction, responsiveness, and stimulus**

1124

To predict decisions from neural activity despite the variable reaction times, each trial was
time-warped in order to align the stimulus onset and the movement initiation timepoints,
following (Lin et al., 2020). For no response trials, a random reaction time was selected from
the responsive trials and the corresponding time-warping was applied. Logistic regression
binary classification models with lasso regularization were fit to classify left versus right turn
trials or response versus no response trials from various time windows of the time-warped
neural activity. Thus, time-warped neuronal activity was averaged over windows from various
start and end timepoints to determine the predictability at each time window. The optimal
predictability at any given timepoint was then considered as the best time window that ended
at that timepoint. For each larvae, 6-fold cross-validation with 3 repetitions was utilized to

determine the cross-validated F score, accuracy, and confusion matrix. For stimulus
classification, logistic regression was similarly applied for multi-class classification using the
one-versus-rest scheme. Overall classification performance was quantified using the mean F
score over larvae and was compared to shuffled data, in which the class labels had been
randomly permuted.
1140
1141 Visualizations of single trial trajectories were made by projecting the time-warped trials onto
1142 the weights of an example, randomly chosen logistic regression model.
1143
Significant neurons contributing to turn direction and responsiveness classification were
identified by comparing a neuron's absolute coefficient across the multiple cross-validation
rounds to that from models fit to shuffled class labels. Neurons were deemed to be significantly
contributed using a one-tailed t-test, $p<0.05$. Brain regions (telencephalon, optic tectum,
cerebellum, and hindbrain) were manually annotated to quantify the brain-wide distribution of
significant neurons.
1150
1151
REFERENCES

Afshar A, Santhanam G, Yu BM, Ryu SI, Sahani M, Shenoy KV. 2011. Single-Trial Neural
Correlates of Arm Movement Preparation. *Neuron* 71:555–564.
doi:10.1016/j.neuron.2011.05.047

Agard DA. 1984. Optical Sectioning Microscopy: Cellular Architecture in Three Dimensions.
Annu Rev Bioph Biom 13:191–219. doi:10.1146/annurev.bb.13.060184.001203

Aimon S, Katsuki T, Jia T, Grosenick L, Broxton M, Deisseroth K, Sejnowski TJ, Greenspan RJ.
2019. Fast near-whole–brain imaging in adult Drosophila during responses to stimuli and
behavior. *Plos Biol* 17:e2006732. doi:10.1371/journal.pbio.2006732

Barker AJ, Baier H. 2015. Sensorimotor Decision Making in the Zebrafish Tectum. *Curr Biol*
25:2804–2814. doi:10.1016/j.cub.2015.09.055

Bartolo R, Saunders RC, Mitz AR, Averbeck BB. 2020. Information-Limiting Correlations in Large
Neural Populations. *J Neurosci* 40:1668–1678. doi:10.1523/jneurosci.2072-19.2019

Bianco IH, Engert F. 2015. Visuomotor Transformations Underlying Hunting Behavior in
Zebrafish. *Curr Biol* 25:831–846. doi:10.1016/j.cub.2015.01.042

Bianco IH, Kampff AR, Engert F. 2011. Prey Capture Behavior Evoked by Simple Visual Stimuli in
Larval Zebrafish. *Frontiers Syst Neurosci* 5:101. doi:10.3389/fnsys.2011.00101

Bollmann JH. 2019. The Zebrafish Visual System: From Circuits to Behavior. *Annu Rev Vis Sc*
5:269–293. doi:10.1146/annurev-vision-091718-014723

Borla MA, Palecek B, Budick S, O'Malley DM. 2002. Prey Capture by Larval Zebrafish: Evidence
for Fine Axial Motor Control. *Brain Behav Evol* 60:207–229. doi:10.1159/000066699

Broxton M, Grosenick L, Yang S, Cohen N, Andalman A, Deisseroth K, Levoy M. 2013. Wave
optics theory and 3-D deconvolution for the light field microscope. *Opt Express* 21:25418–
39. doi:10.1364/oe.21.025418

Chen X, Mu Y, Hu Y, Kuan AT, Nikitchenko M, Randlett O, Chen AB, Gavornik JP, Sompolinsky H,
Engert F, Ahrens MB. 2018. Brain-wide Organization of Neuronal Activity and Convergent
Sensorimotor Transformations in Larval Zebrafish. *Neuron* 100:876-890.e5.
doi:10.1016/j.neuron.2018.09.042

Cohen MR, Kohn A. 2011. Measuring and interpreting neuronal correlations. *Nat Neurosci*
14:811–819. doi:10.1038/nn.2842

Cohen MR, Maunsell JHR. 2009. Attention improves performance primarily by reducing
interneuronal correlations. *Nat Neurosci* 12:1594–1600. doi:10.1038/nn.2439

Cohen N, Yang S, Andalman A, Broxton M, Grosenick L, Deisseroth K, Horowitz M, Levoy M.
2014. Enhancing the performance of the light field microscope using wavefront coding. *Opt*
Express 22:24817–39. doi:10.1364/oe.22.024817

Cong L, Wang Z, Chai Y, Hang W, Shang C, Yang W, Bai L, Du J, Wang K, Wen Q. 2017. Rapid
whole brain imaging of neural activity in freely behaving larval zebrafish (Danio rerio). *Elife*
6:e28158. doi:10.7554/elife.28158

Dunn Timothy W., Gebhardt C, Naumann EA, Riegler C, Ahrens MB, Engert F, Del Bene F. 2016.
Neural Circuits Underlying Visually Evoked Escapes in Larval Zebrafish. *Neuron* 89:613–628.
doi:10.1016/j.neuron.2015.12.021

Dunn Timothy W, Mu Y, Narayan S, Randlett O, Naumann EA, Yang C-T, Schier AF, Freeman J,
Engert F, Ahrens MB. 2016. Brain-wide mapping of neural activity controlling zebrafish
exploratory locomotion. *Elife* 5:e12741. doi:10.7554/elife.12741

Faisal AA, Selen LPJ, Wolpert DM. 2008. Noise in the nervous system. *Nat Rev Neurosci* 9:292–
303. doi:10.1038/nrn2258

Filosa A, Barker AJ, Dal Maschio M, Baier H. 2016. Feeding State Modulates Behavioral Choice
and Processing of Prey Stimuli in the Zebrafish Tectum. *Neuron* 90:596–608.
doi:10.1016/j.neuron.2016.03.014

Flavell SW, Gogolla N, Lovett-Barron M, Zelikowsky M. 2022. The emergence and influence of
internal states. *Neuron* 110:2545–2570. doi:10.1016/j.neuron.2022.04.030

Flavell SW, Pokala N, Macosko EZ, Albrecht DR, Larsch J, Bargmann CI. 2013. Serotonin and the
Neuropeptide PDF Initiate and Extend Opposing Behavioral States in C. elegans. *Cell*
154:1023–1035. doi:10.1016/j.cell.2013.08.001

Förster D, Helmbrecht TO, Mearns DS, Jordan L, Mokayes N, Baier H. 2020. Retinotectal
circuitry of larval zebrafish is adapted to detection and pursuit of prey. *Elife* 9:e58596.
doi:10.7554/elife.58596

Giovannucci A, Friedrich J, Gunn P, Kalfon J, Brown BL, Koay SA, Taxidis J, Najafi F, Gauthier JL,
Zhou P, Khakh BS, Tank DW, Chklovskii DB, Pnevmatikakis EA. 2019. CaImAn an open source
tool for scalable calcium imaging data analysis. *Elife* 8:e38173. doi:10.7554/elife.38173

Guo C, Liu W, Hua X, Li H, Jia S. 2019. Fourier light-field microscopy. *Opt Express* 27:25573.
doi:10.1364/oe.27.025573

Huang X, Lisberger SG. 2009. Noise correlations in cortical area MT and their potential impact
on trial-by-trial variation in the direction and speed of smooth-pursuit eye movements. *J*
Neurophysiol 101:3012–30. doi:10.1152/jn.00010.2009

Kafashan M, Jaffe AW, Chettih SN, Nogueira R, Arandia-Romero I, Harvey CD, Moreno-Bote R,
Drugowitsch J. 2021. Scaling of sensory information in large neural populations shows
signatures of information-limiting correlations. *Nat Commun* 12:473. doi:10.1038/s41467-
020-20722-y

Kanitscheider I, Coen-Cagli R, Pouget A. 2015. Origin of information-limiting noise correlations.
Proc National Acad Sci 112:E6973–E6982. doi:10.1073/pnas.1508738112

Kao JC, Nuyujukian P, Ryu SI, Churchland MM, Cunningham JP, Shenoy KV. 2015. Single-trial
dynamics of motor cortex and their applications to brain-machine interfaces. *Nat Commun*
6:7759. doi:10.1038/ncomms8759

Karashchuk P, Rupp KL, Dickinson ES, Walling-Bell S, Sanders E, Azim E, Brunton BW, Tuthill JC.
2020. Anipose: A toolkit for robust markerless 3D pose estimation. *Cell Reports* 36:109730.
doi:10.1016/j.celrep.2021.109730

Kaufman MT, Churchland MM, Ryu SI, Shenoy KV. 2015. Vacillation, indecision and hesitation in
moment-by-moment decoding of monkey motor cortex. *Elife* 4:e04677.
doi:10.7554/elife.04677

Kauvar IV, Machado TA, Yuen E, Kochalka J, Choi M, Allen WE, Wetzstein G, Deisseroth K. 2020.
Cortical Observation by Synchronous Multifocal Optical Sampling Reveals Widespread
Population Encoding of Actions. *Neuron*. doi:10.1016/j.neuron.2020.04.023

Kim TH, Schnitzer MJ. 2022. Fluorescence imaging of large-scale neural ensemble dynamics. *Cell*
185:9–41. doi:10.1016/j.cell.2021.12.007

Kohn A, Coen-Cagli R, Kanitscheider I, Pouget A. 2015. Correlations and Neuronal Population
Information. *Annu Rev Neurosci* 39:1–20. doi:10.1146/annurev-neuro-070815-013851

Lecoq J, Oliver M, Siegle JH, Orlova N, Ledochowitsch P, Koch C. 2021. Removing independent
noise in systems neuroscience data using DeepInterpolation. *Nat Methods* 18:1401–1408.
doi:10.1038/s41592-021-01285-2

Levoy M, Ng R, Adams A, Footer M, Horowitz M. 2006. Light field microscopy. *Acm T Graphic*
25:924. doi:10.1145/1141911.1141976

Lin Q, Manley J, Helmreich M, Schlumm F, Li JM, Robson DN, Engert F, Schier A, Nöbauer T,
Vaziri A. 2020. Cerebellar Neurodynamics Predict Decision Timing and Outcome on the
Single-Trial Level. *Cell* 180:536-551.e17. doi:10.1016/j.cell.2019.12.018

Liu FL, Kuo G, Antipa N, Yanny K, Waller L. 2020. Fourier DiffuserScope: Single-shot 3D Fourier
light field microscopy with a diffuser. *Arxiv*. doi:10.1364/oe.400876

Llavador A, Sola-Pikabea J, Saavedra G, Javidi B, Martínez-Corral M. 2016. Resolution
improvements in integral microscopy with Fourier plane recording. *Opt Express* 24:20792–8.
doi:10.1364/oe.24.020792

Machado TA, Kauvar IV, Deisseroth K. 2022. Multiregion neuronal activity: the forest and the
trees. *Nat Rev Neurosci* 23:683–704. doi:10.1038/s41583-022-00634-0

Marques JC, Li M, Schaak D, Robson DN, Li JM. 2020. Internal state dynamics shape brainwide
activity and foraging behaviour. *Nature* 577:239–243. doi:10.1038/s41586-019-1858-z

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018.
DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat
Neurosci* 21:1281–1289. doi:10.1038/s41593-018-0209-y

Montijn JS, Meijer GT, Lansink CS, Pennartz CMA. 2016. Population-Level Neural Codes Are
Robust to Single-Neuron Variability from a Multidimensional Coding Perspective. *Cell
Reports* 16:2486–2498. doi:10.1016/j.celrep.2016.07.065

Moreno-Bote R, Beck J, Kanitscheider I, Pitkow X, Latham P, Pouget A. 2014. Information-
limiting correlations. *Nat Neurosci* 17:1410–1417. doi:10.1038/nn.3807

Musall S, Kaufman MT, Juavinett AL, Gluf S, Churchland AK. 2019. Single-trial neural dynamics
are dominated by richly varied movements. *Nat Neurosci* 22:1677–1686.
doi:10.1038/s41593-019-0502-4

Nöbauer T, Skocek O, Pernía-Andrade AJ, Weilguny L, Traub FM, Molodtsov MI, Vaziri A. 2017.
Video rate volumetric Ca2+ imaging across cortex using seeded iterative demixing (SID)
microscopy. *Nat Methods* 14:811–818. doi:10.1038/nmeth.4341

Nöbauer T, Zhang Y, Kim H, Vaziri A. 2023. Mesoscale volumetric light-field (MesoLF) imaging of
neuroactivity across cortical areas at 18 Hz. *Nat Methods* **20**:600–609. doi:10.1038/s41592-
023-01789-z

Okamoto H, Cherng B-W, Nakajo H, Chou M-Y, Kinoshita M. 2021. Habenula as the experience-
dependent controlling switchboard of behavior and attention in social conflict and learning.
Curr Opin Neurobiol **68**:36–43. doi:10.1016/j.conb.2020.12.005

Oldfield CS, Grossrubatscher I, Chávez M, Hoagland A, Huth AR, Carroll EC, Prendergast A, Qu T,
Gallant JL, Wyart C, Isacoff EY. 2020. Experience, circuit dynamics and forebrain recruitment
in larval zebrafish prey capture. *Elife* 9:e56619. doi:10.7554/elife.56619

Pandarinath C, Ames KC, Russo AA, Farshchian A, Miller LE, Dyer EL, Kao JC. 2018. Latent
Factors and Dynamics in Motor Cortex and Their Application to Brain–Machine Interfaces. *J
Neurosci* 38:9390–9401. doi:10.1523/jneurosci.1669-18.2018

Patterson BW, Abraham AO, MacIver MA, McLean DL. 2013. Visually guided gradation of prey
capture movements in larval zebrafish. *J Exp Biology* 216:3071–3083.
doi:10.1242/jeb.087742

Portugues R, Engert F. 2009. The neural basis of visual behaviors in the larval zebrafish. *Curr
Opin Neurobiol* 19:644–647. doi:10.1016/j.conb.2009.10.007

Prevedel R, Yoon Y-G, Hoffmann M, Pak N, Wetzstein G, Kato S, Schrödel T, Raskar R, Zimmer
M, Boyden ES, Vaziri A. 2014. Simultaneous whole-animal 3D imaging of neuronal activity
using light-field microscopy. *Nat Methods* 11:727–730. doi:10.1038/nmeth.2964

Quicke P, Howe CL, Song P, Jadan HV, Song C, Knöpfel T, Neil M, Dragotti PL, Schultz SR, Foust
AJ. 2020. Subcellular resolution 3D light field imaging with genetically encoded voltage
indicators. *Biorxiv* 2020.05.22.108191. doi:10.1101/2020.05.22.108191

Rumyantsev OI, Lecoq JA, Hernandez O, Zhang Y, Savall J, Chrapkiewicz R, Li J, Zeng H, Ganguli S,
Schnitzer MJ. 2020. Fundamental bounds on the fidelity of sensory cortical coding. *Nature*
580:100–105. doi:10.1038/s41586-020-2130-2

Scrofani G, Sola-Pikabea J, Llavador A, Sanchez-Ortiga E, Barreiro JC, Saavedra G, Garcia-
Sucerquia J, Martínez-Corral M. 2017. FIMic: design for ultimate 3D-integral microscopy of
in-vivo biological samples. *Biomed Opt Express* 9:335. doi:10.1364/boe.9.000335

Semmelhack JL, Donovan JC, Thiele TR, Kuehn E, Laurell E, Baier H. 2014. A dedicated visual
pathway for prey detection in larval zebrafish. *Elife* 3:e04878. doi:10.7554/elife.04878

Shadlen MN, Newsome WT. 1998. The Variable Discharge of Cortical Neurons: Implications for
Connectivity, Computation, and Information Coding. *J Neurosci* 18:3870–3896.
doi:10.1523/jneurosci.18-10-03870.1998

Skocek O, Nöbauer T, Weilguny L, Traub FM, Xia CN, Molodtsov MI, Grama A, Yamagata M,
Aharoni D, Cox DD, Golshani P, Vaziri A. 2018. High-speed volumetric imaging of neuronal
activity in freely moving rodents. *Nat Methods* 15:429–432. doi:10.1038/s41592-018-0008-0

Sternson SM. 2020. Exploring internal state-coding across the rodent brain. *Curr Opin Neurobiol*
65:20–26. doi:10.1016/j.conb.2020.08.009

Štih V, Petrucco L, Kist AM, Portugues R. 2019. Stytra: An open-source, integrated system for
stimulation, tracking and closed-loop behavioral experiments. *Plos Comput Biol*
15:e1006699. doi:10.1371/journal.pcbi.1006699

Stringer C, Pachitariu M, Steinmetz N, Reddy CB, Carandini M, Harris KD. 2019. Spontaneous
behaviors drive multidimensional, brainwide activity. *Science* 364:eaav7893.
doi:10.1126/science.aav7893

Temizer I, Donovan JC, Baier H, Semmelhack JL. 2015. A Visual Pathway for Looming-Evoked
Escape in Larval Zebrafish. *Curr Biol* 25:1823–1834. doi:10.1016/j.cub.2015.06.002

Torigoe M, Islam T, Kakinuma H, Fung CCA, Isomura T, Shimazaki H, Aoki T, Fukai T, Okamoto H.
2021. Zebrafish capable of generating future state prediction error show improved active
avoidance behavior in virtual reality. *Nat Commun* 12:5712. doi:10.1038/s41467-021-26010-
7

Valente M, Pica G, Bondanelli G, Moroni M, Runyan CA, Morcos AS, Harvey CD, Panzeri S. 2021.
Correlations enhance the behavioral readout of neural population activity in association
cortex. *Nat Neurosci* 24:975–986. doi:10.1038/s41593-021-00845-1

Vladimirov N, Mu Y, Kawashima T, Bennett DV, Yang C-T, Looger LL, Keller PJ, Freeman J, Ahrens
MB. 2014. Light-sheet functional imaging in fictively behaving zebrafish. *Nat Methods*
11:883–884. doi:10.1038/nmeth.3040

Wei Z, Inagaki H, Li N, Svoboda K, Druckmann S. 2019. An orderly single-trial organization of
population dynamics in premotor cortex predicts behavioral variability. *Nat Commun* 10:216.
doi:10.1038/s41467-018-08141-6

Weisenburger S, Vaziri A. 2018. A Guide to Emerging Technologies for Large-Scale and Whole-
Brain Optical Imaging of Neuronal Activity. *Annu Rev Neurosci* 41:431–452.
doi:10.1146/annurev-neuro-072116-031458

Wolf S, Dubreuil AM, Bertoni T, Böhm UL, Bormuth V, Candelier R, Karpenko S, Hildebrand DGC,
Bianco IH, Monasson R, Debrégeas G. 2017. Sensorimotor computation underlying
phototaxis in zebrafish. *Nat Commun* 8:651. doi:10.1038/s41467-017-00310-3

Yoon Y-G, Wang Z, Pak N, Park D, Dai P, Kang JS, Suk H-J, Symvoulidis P, Guner-Ataman B, Wang
K, Boyden ES. 2020. Sparse decomposition light-field microscopy for high speed imaging of
neuronal activity. *Optica* 7:1457. doi:10.1364/optica.392805

Zhou P, Resendez SL, Rodriguez-Romaguera J, Jimenez JC, Neufeld SQ, Giovannucci A, Friedrich
J, Pnevmatikakis EA, Stuber GD, Hen R, Kheirbek MA, Sabatini BL, Kass RE, Paninski L. 2018.
Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data.
Elife 7:e28728. doi:10.7554/elife.28728

Zohary E, Shadlen MN, Newsome WT. 1994. Correlated neuronal discharge rate and its
implications for psychophysical performance. *Nature* 370:140–143. doi:10.1038/370140a0

Zylberberg J, Cafaro J, Turner MH, Shea-Brown E, Rieke F. 2016. Direction-Selective Circuits
Shape Noise to Ensure a Precise Population Code. *Neuron* 89:369–383.
doi:10.1016/j.neuron.2015.11.019

Zylberberg J, Pouget A, Latham PE, Shea-Brown E. 2017. Robust information propagation
through noisy neural circuits. *Plos Comput Biol* 13:e1005497.
doi:10.1371/journal.pcbi.1005497

1351

Figure 1: Fourier Light Field Microscopy (fLFM) provides a simple and cost-effective method for whole-brain imaging of larval zebrafish during behavior

Figure 2: Zebrafish exhibit highly variable motor responses to visual stimuli

A Freely-swimming behavior

B Example target-avoidance bout

C Example target-directed bout

D Simple stimuli of varying size elicit divergent behavioral responses

E Head-fixed behavior

F Behavioral responses are recapitulated during head-fixed imaging

G Head-fixed behavioral responses are highly variable across trials for nearly all stimuli

Figure 3: Trial-to-trial variability in visually-evoked neurons is largely orthogonal to visual decoding dimensions

A Individual tuned neurons exhibit trial-to-trial variability

B Stimuli are robustly decodable at the population level

C Trial-to-trial variability (e_1) could be aligned with or othogonal to decoding dimensions (w_{opt})

(i) Orthogonal variability   **(ii)** Uniform variability   **(iii)** Aligned variability

D Noise correlation matrix appears multi-dimensional

E Trial-to-trial neural variability is largely orthogonal to decoding dimensions

F Example noise versus stimulus activity patterns

G w_{opt} coefficients

H e_1 coefficients

I Neural variability correlates with tail vigor

J Largest noise modes correlate with brain-wide motor activity

Figure 4: Pre-motor neuronal populations predictive of single-trial behavior

Figure 5: Spontaneous turns are predictable from the same pre-motor neuronal population

A Fit models on visually-evoked turns

Tail curvature

Warped time   Warped time

Predict spontaneous turns

Tail curvature

300   320   340
Time (sec)

B Spontaneous turns are predicted from visual-evoked model

Pre-motor turn direction prediction (a.u.)

r=0.41

n=5 larvae

-270   0   270
Tail curvature (deg)

C Spontaneous pre-motor accuracy across larvae

True turn direction

Left: 63.3%   36.7%
Right: 22.9%   77.1%

Left   Right
Predicted turn direction

% of true trials
0