The telencephalon is a neuronal substrate for systemic inflammatory responses in teleosts via polyamine metabolism

Amir Mani[1], Farah Haddad[2], Daniel R. Barreda[2], Irene Salinas[1]*

[1] Center for Evolutionary and Theoretical Immunology, Department of Biology, University of New Mexico, Albuquerque, New Mexico, 87108, USA.

[2] Department of Biological Sciences, University of Alberta, Edmonton, Canada.

*Irene Salinas

Email: isalinas@unm.edu

Author Contributions: Conceptualization: IS, AM; Investigation: AM, FH, DB, IS; Methodology: AM, IS; Formal analysis: AM, FH; Writing-original draft: AM, IS; Writing-reviewing and editing: AM, FH, DB, IS.

Competing Interest Statement: Authors have no competing interests.

Classification: Biological sciences. Immunology and Inflammation.

Keywords: teleosts; telencephalon; hippocampus; polyamines; sickness behavior; behavioral fever, rainbow trout

Abstract

Systemic inflammation elicits sickness behaviors and fever by engaging a complex neuronal circuitry that begins in the preoptic area of the hypothalamus. Ectotherms such as teleost fish display sickness behaviors in response to infection or inflammation, seeking warmer temperatures to enhance survival via behavioral fever responses. To date, the hypothalamus is the only brain region implicated in sickness behaviors and behavioral fever in teleosts. Yet, the complexity of neurobehavioral manifestations underlying sickness responses in teleosts suggests engagement of higher processing areas of the brain. Using two *in vivo* models of systemic inflammation in rainbow trout, we find canonical pyrogenic cytokine responses in the hypothalamus whereas in the telencephalon *il-1b* and *tnfa* expression is decoupled from *il-6* expression. Polyamine metabolism changes, characterized by accumulation of putrescine and decreases in spermine and spermidine, are recorded in the telencephalon but not hippocampus upon systemic injection of bacteria. While systemic inflammation causes canonical behavioral fever in trout, blockade of bacterial polyamine metabolism prior to injection abrogates behavioral fever, polyamine responses and telencephalic but not hypothalamic cytokine responses. Combined, our work identifies the telencephalon as a neuronal substrate for brain responses to systemic inflammation in teleosts and uncovers the role of polyamines as critical chemical mediators in sickness behaviors.

Significance Statement

Systemic inflammation induces neuroimmune responses in the brain resulting in sickness behaviors and fever. In endotherms, sickness behaviors and fever are initiated in the hypothalamus but also engage a complex neuronal circuitry in higher areas of the brain. In ectotherms, only the hypothalamus has been linked to sickness behaviors and behavioral fever. Here we demonstrate that the telencephalon, a critical region of the teleost brain responsible for higher order processing, mounts pyrogenic cytokine responses to systemic inflammation in teleosts that are different from those of the hypothalamus. We identify polyamine metabolism as a

core response of the teleost telencephalon to systemic inflammation and report that bacterial polyamines are triggers of behavioral fever in rainbow trout. Our work uncovers a previously unrecognized role for the telencephalon and polyamine metabolism in sickness behaviors and behavioral fever in teleosts with implications in fish health and fish conservation.

Main Text

Introduction

The bidirectional communication between the periphery and the brain regulates homeostasis and disease states in animals (1–8). Systemic inflammation results in sickness behaviors, such as anorexia, depression, lethargy, sleepiness, social withdrawal, and sometimes fever responses. Sickness behaviors and fever are cytokine orchestrated, complex neurobehavioral phenomena that involve multiple neural substrates (9–13). In mammals, the Median Preoptic Nucleus (MnPO) of the hypothalamus is the first region to detect systemic inflammatory signals (11–14) but extensive experimental evidence indicates that the hypothalamus is not the only brain region responding to systemic inflammation. For instance, systemic inflammation engages neurons in the mouse insular cortex and these neurons can evoke past inflammatory states in the periphery following optogenetic neuronal activation (15, 16). Furthermore, bacterial lipopolysaccharide (LPS) injection, the gold standard model for inducing sickness behaviors in mammals, results in significant neuronal activation not only in the hypothalamus but also in key limbic regions, including the amygdala, hippocampus, nucleus accumbens, and the stria terminalis, indicating a broad neuroanatomical network responding to systemic inflammation (17, 18). Similarly, in rats, LPS injection triggers astrocyte activation within the striatum and hippocampus with significant volumetric expansions occurring in the splenium, retrosplenial cortex, and pericallosal hippocampal cortex at 2 hours post- injection (hpi) (19). Combined, mammalian studies indicate that systemic inflammation disturbs several higher-order cortical areas (13, 15, 20, 21), their connectivity across the entire brain yet to be fully resolved.

Ectotherms, like mammals, also display sickness behaviors in response to infection, stress and inflammation (22–26). For instance, teleosts become lethargic, stop feeding and seek warmer water temperatures in a process known as behavioral fever (27–32). This behaviorally induced fever provides a survival advantage, as it enhances the immune system's efficacy, hinders the proliferation of pathogens and resolves wound healing more effectively (27, 28, 31–33). Similar to mammals, systemic and brain increases in pyrogenic cytokines like IL1β, TNFα, and IL6, alongside PGE2, have been identified as key players in teleosts behavioral fever responses. Furthermore, the preoptic area of the hypothalamus (POA) is thought to be, as in mammals, the main brain region responding to systemic infection and driving behavioral fever responses in ectotherms (30, 33–35). Yet, expression of the prostaglandin receptor EP3 was detected in the telencephalon and optic lobe of Atlantic salmon (29), suggesting that other brain regions detect systemic inflammation in teleosts. Given the complexity of sickness behaviors and fever in teleosts, we hypothesized that the teleost brain must not only sense and produce proinflammatory cytokines in response to systemic inflammation, but also engage higher order processing areas responsible for decision-making and voluntary movement.

The teleost brain, though structurally different from that of mammals, performs similarly complex functions (36). The telencephalon of teleosts, comprising the pallium and subpallium, is increasingly recognized for its role in higher-order processing including spatial cognition, sensation, and voluntary movement, all of which are integral to behavioral fever (37–43). For instance, the teleost telencephalon plays pivotal roles in spatial navigation (40, 41, 43), avoidance behaviors (38), and the coordination of movement towards warmer environments (41, 44, 45). Interestingly, the dorsal, medial, and ventrodorsal telencephalon in teleosts are analogous to the mammalian hippocampus, amygdala, and basal ganglia, respectively (37–43, 46), all of which respond to systemic inflammation in mammals. Based on these premises, along with the EP3 expression in the teleost telencephalon, we hypothesized that this region would be a likely

neuronal substrate candidate in systemic inflammation and behavioral fever responses in teleosts.

Herein we investigate the neuroimmune responses of the rainbow trout brain to systemic inflammation. By comparing the telencephalon to the hippocampus, we identify sharp differences in how these two brain regions respond to inflammation. We also provide experimental evidence that shifts in polyamine metabolism occur specifically in the telencephalon in response to systemic inflammation, and that bacterial derived polyamines regulate the onset of behavioral fever. Our study reveals a novel neuronal substrate beyond the POA that governs responses to systemic inflammation in higher processing areas of the teleost brain.

Results

Systemic microbiota and LPS induce distinct cytokine responses in the trout telencephalon and hippocampus

To investigate brain neuroimmune responses to systemic inflammation we developed two *in vivo* models in juvenile rainbow trout (Fig. 1A). In the first model, we collected bacteria from the gut of healthy rainbow trout and injected it intravenously (i.v) (fig. S1, A and B). Since ectotherms are highly refractory to *E. coli* LPS injection (47–49), we decided to isolate LPS from the native trout gut microbiota instead of using commercial sources. We first measured LPS levels in the serum of trout injected i.v with the isolated bacteria (Fig. 1B). Notably, LPS levels were only elevated in the blood serum of fish injected with gut bacteria at 4 hpi, returning to baseline levels by 1 day (Fig. 1B). Next, we adjusted how much of the commensal LPS we would need to inject to match this value in both models (Fig. 1A and B). In both models, we noted canonical reductions in fish locomotion, but no mortalities occurred during any of the experiments.

Previous studies have established that like mammals, the teleost hippocampus (Hyp) expresses pyrogenic cytokines in response to systemic or systemic infection (28, 32, 35). However, whether other regions of the teleost brain respond to systemic inflammation is currently unknown. We hypothesized, given the role of the teleost telencephalon (Tel) in higher order processing, that this region would be part of the neuronal circuitry responding to systemic insults. Thus, we compared the expression of pyrogenic cytokine genes in the Hyp and Tel in both of our *in vivo* models. As previously reported in the Hyp, *il-1b*, *tnfa*, and *il-6* expression was significantly higher (2-4-fold increase) in commensal-injected and LPS-injected trout 4 hpi (Fig. 1D, F, and H). Expression levels of all the cytokines returned to baseline levels by 1 day for both the commensal and LPS-injected groups (Fig. 1D, F, and H). In the Tel, *il-1b* and *tnfa* expression rose 2-3 folds at 4 hpi for both commensals or LPS injection (Fig. 1C and E). However, *il-6* expression remained unchanged at all tested timepoints and treatments (Fig. 1G). Like the Hyp, Tel cytokine expression levels returned to basal levels by 1 day. These results indicate that the trout Tel responds to systemic inflammation with a unique cytokine signature compared to the hippocampus.

LPS and other systemic inflammation mediators cause breakdown of the blood brain barrier (BBB) in mammals (50, 51). We measured the integrity of the trout BBB in response to the injection of gut-derived bacteria using Evans blue extravasation, a widely used technique to measure BBB permeability (52–54) (Fig. 1I and J). We measured Evans blue extravasation in five different brain regions including olfactory bulb (OB), Tel, Hyp, optic tectum (OT) and cerebellum (Cer). BBB permeability increased by 2-fold compared to vehicle controls in all regions examined by 4 hours (fig. S1, C to E) except for the Tel, where we observed a >3-fold increase in Evans blue extravasation at 4 hpi when compared to baseline levels (Fig. 1J). This increased permeability was transient, with levels decreasing towards the baseline within 1 day post-injection, indicating a temporary disruption of the BBB in the Tel (Fig. 1J). In contrast, other brain regions including the Hyp, showed no significant BBB disruption by 1 day (Fig. 1J and fig. S1, C to E). These data indicate that systemic inflammation preferentially breaks down the Tel BBB compared to other brain regions in rainbow trout.

153
154   Systemic bacteria transiently and selectively penetrate the trout telencephalon
155
Our two *in vivo* models indicated that the Tel and the Hyp respond differently to systemic
inflammation. We have recently reported that salmonids have a resident bacterial community in
their brain at steady state (55). We therefore wondered if upon systemic injection of microbiota,
bacteria would be able to penetrate the brain. We quantified total bacterial loads in the spleen
and five regions of the brain 4 hours and 1 day post-injection using 16s rDNA qPCR. In the Tel, a
substantial increase in bacterial loads was observed at 4 hpi with gut commensals. Bacterial
presence was transient, as levels at 1 day did not differ significantly from baseline measurements
(Fig. 2A). Conversely, the bacterial loads in the Hyp as well as all other brain regions and the
spleen remained unchanged (Fig. 2B and fig. S2, E to J). As predicted, the administration of LPS
did not change bacterial loads in any tissue sampled at either 4 hours or 1 day post-injection (fig.
S2, E to J). We next tested a second model of gut bacterial translocation and systemic
inflammation (fig. S3, A). We employed the recently developed DSS-induced colitis in rainbow
trout (56) and 1 day after the last DSS gavage, we collected the same tissues as before except
the Hyp (fig. S3, A). Similar to the i.v injection model, we noted significant increases in serum
endotoxin levels in DSS treated compared to vehicle treated controls (fig. S3, B), but bacterial
loads were only significantly higher in the Tel of DSS-treated animals compared to controls, with
no other significant changes noted in the other brain regions (fig. S3, C to G). Combined, these
experiments indicate that transient and region-specific bacterial translocation occurs in the trout
brain during systemic inflammation.
175
Given that bacteria penetrate the trout Tel, we next sought to evaluate whether shifts in the Tel
microbial community occur following i.v bacterial injection using 16S rDNA sequencing. Collected
gut bacteria from healthy laboratory rainbow trout were predominantly composed of four bacterial
phyla: Tenericutes (72.2%), Proteobacteria (11.2%), Fusobacteria (7.4%), and Firmicutes (7.2%).
Further analysis at the family level revealed the bacterial community to consist primarily of eight
families: Mycoplasmataceae (72.2%), Fusobacteriaceae (7.2%), Lachnospiraceae (4.9%),
Chitinibacteriaceae (3.0%), and Burkholderiaceae (1.9%) (fig. S1, A and B) and were markedly
different from the tank water microbial community (fig. S4, A and B). The predominant bacterial
phyla identified in the Tel of control animals included Proteobacteria (37.93%), Tenericutes
(27.0%), Actinobacteria (16.5%), and Bacteroidetes (7.9%) mainly composed of
Mycoplasmataceae (27.04%), propionibacteriaceae (11.3%), Chitinibacteraceae (10.7%) and
Shewanellaceae (9.1%) families (Fig. 2C and D). Finally, the bacterial composition in the tank
water was distinctly different from that observed in the Tel and composed mostly of members of
Acidobacteria (29.5%), Bacteroidetes (31.2%), Proteobacteria (17.3%), and Nitrospirae (6.2%)
phyla, such as the families Blastocatellia (uncultured family, 26.6%), Microsillaceae (8.1%),
Nitrososospiraceae (6.2%), and Saprospiraceae (11.4%), indicating a unique microbial community
compared to that of the Tel (fig. S2, A and B). Following i.v injection of bacteria, there was an
increase in the relative abundance of Tenericutes, Firmicutes, and Fusobacteria, whereas
Proteobacteria, Bacteroidetes, and Actinobacteria relative abundance decreased compared to the
vehicle group at the 4-hour and 1-day time points (Fig. 2C). At the family level, the relative
abundance of Mycoplasmataceae, Lachnospiraceae, and Chromobacteriaceae increased at both
time points (Fig. 2D). Conversely, the relative abundance of families such as Enterobacteriaceae,
Moraxellaceae, and Shewanellaceae diminished at the same intervals (Fig. 2D and G). While the
Tel microbial community composition at the 4-hour and 1 day post-injection time points were
largely similar, Ruminococcaceae and Streptococcaceae accounted for 10.4% and 7%,
respectively, of the overall diversity at 4 hours whereas they were only 1.3% and 0.3%,
respectively, at 1 day. At 1 day, Fusobacteriaceae and Chitinibacteraceae relative abundances
were 15.4%, and 8.3%, respectively, a much larger percentage than at 4 hours (0% and 0.1%)
(Fig. 2D). The Weighted UniFrac distance between the Tel bacterial community of vehicle-
injected group and commensal-injected group was significantly different at 1 day (Fig. 2E).
Principal coordinate analysis revealed distinct microbial community compositions between the
vehicle and commensal injected groups, with the Tel microbial communities of the injected

animals remaining closely clustered (Fig. 2F). Alpha diversity metrics (Shannon diversity, Chao1,
and Pielou's evenness) did not show any significant differences in bacteria-injected versus
vehicle-injected animals (fig. S4, C to E).
211
PICRUSt2 analysis revealed shifts in functional pathways of the Tel microbial community
following i.v injection of bacteria. A total of 27 pathways were significantly modulated in the Tel
microbial community at 4 hour and 1 day compared to vehicle controls. Several polyamine
biosynthesis pathways, such as putrescine, spermidine, and spermine biosynthesis were
upregulated at both time points compared to controls (Fig. 2H). Additionally, nucleotide
biosynthesis, including adenine and guanine pathways, L-glutamine and L-lysine biosynthesis,
and degradation pathways like L-histidine degradation and fatty acid salvage were modulated as
a result of the bacterial injection treatment (Fig. 2H). These results indicate that, despite bacterial
loads dropping by 1 day, the systemic inflammation in the Tel microbial community composition
and its predicted functions has longer lasting impacts. Combined, these results show that only
specific suites of gut systemic bacteria penetrate the Tel of rainbow trout altering its community
composition and predict changes in polyamine metabolism in the Tel bacterial community.
224
225 Systemic inflammation modulates host brain polyamines in a region-specific manner
226
Polyamines are polycationic organic molecules with two or more amino groups. Polyamines are
universal molecules in nature, produced by archaea, bacteria and eukaryotes and are involved in
fundamental biological processes such as growth, cell proliferation, differentiation, and death (57–
60). Given that the Tel microbial composition showed predicted changes in bacterial polyamine
synthesis pathways, we next asked if these shifts had an impact on host polyamine synthesis
pathways. Thus, we measure expression levels of four genes encoding for polyamine synthesis
enzymes (*odc1*, *odc2*, *samdc1*, and *samdc2*) in the trout Tel and Hyp (Fig. 3A and B). We
observed a marked increase in the expression of *odc* genes in the Tel, with two-fold and three-
fold elevations at 4 hours and 1 day post-injection of gut commensals, respectively (Fig. 3C and
fig. S5, A). Conversely, a significant decrease in the levels of *samdc* transcripts was noted in the
Tel, with two-fold and three-fold reductions at 4 hours and 1 day post-injection of gut
commensals, respectively (Fig. 3D and fig. S5, C). LPS injection, however, did not cause any
changes in *odc* or *samdc* expression in the Tel. Interestingly, *odc* and *samdc* gene expression did
not change in the Hyp at any given time point or treatment group (Fig. 3E and F and fig. S5, B
and D), indicating the host polyamine responses to systemic bacteria are region-specific.
242
In order to test if host changes in polyamine synthesis gene expression are dependent on the
ability of bacteria to make polyamines, we blocked polyamine metabolism in the microbiota prior
to injecting them into the host. We treated isolated gut commensals with the polyamine pathway
inhibitors DFMA (α-difluoromethylarginine), an irreversible inhibitor of arginine decarboxylase,
and DFMO (α-difluoromethylornithine), an irreversible suicide inhibitor of ornithine decarboxylase
(fig. S2, A to D). Since polyamines can be essential for bacterial growth (61–64) we first
performed *in vitro* experiments to establish the highest dose of DFMA and DFMO that would not
compromise bacterial growth (fig. S2, A to D). Based on these assays we selected 0.25 mM for
each drug for all subsequent *in vivo* experiments (Fig. 3B). Bacteria were washed three times
prior to injection to ensure the removal of DFMA and DFMO (65). We first assessed if treated
bacteria were still capable of translocating into the trout Tel and found that polyamine synthesis
blockade did not impact the ability of bacteria to enter the Tel 4h post-injection (fig. S2, E to J). In
response to injection of DFMA-DFMO treated bacteria, trout *odc* or *samdc* expression levels did
not change in the Tel or Hyp at the time points tested (Fig. 3C and fig. S5, A to D). Combined,
these results indicate that bacterial-derived polyamines selectively modulate host polyamine
synthesis pathways in the Tel but not the Hyp in rainbow trout.
259
260 We next asked if changes in the expression of polyamine synthesis genes in the trout Tel
261 correlated with changes in the Tel polyamine levels. We measured putrescine, spermidine and
262 spermine levels in telencephalic tissue extracts under the different experimental conditions using
263 ELISA. Systemic injection of gut commensals resulted in a 2-fold increase in telencephalic

putrescine levels at both 4 hours and 1 day post-injection (Fig. 3G). Conversely, a sharp decrease in spermidine, and spermine levels was observed in commensal-injected trout, with a 3-5-fold reduction at both time points (Fig. 3H and I). Putrescine, spermidine and spermine levels in the Tel of trout injected with DFMA-DFMO treated bacteria were not different from vehicle controls. (Fig. 3G to I). These results indicate that bacteria-derived polyamines alter host polyamine metabolism in the Tel resulting in putrescine accumulation and downstream reduction of spermidine and spermine.

Bacterial-derived polyamines selectively modulate telencephalic cytokine responses

Polyamines play crucial roles in innate immunity in mammals (66–68). We therefore asked if polyamines from microbiota could drive the expression of pyrogenic cytokines in the trout brain and hypothesized that this may occur in a region-specific manner. We administered DFMA-DFMO treated or untreated bacteria to animals (Fig. 4A) and measured *il-1b*, *tnfa*, and *il-6* transcript levels at 4 hours and 1 day post-injection. Untreated bacteria augmented the expression of *il-1b* and *tnfa* in the Tel 4 hpi. In sharp contrast, DFMA-DFMO treated bacteria failed to induce any significant changes in the expression levels of these cytokines (Fig. 4B and D). Additionally, the expression pattern of *il-6* in the Tel remained consistent across all treatment groups and the vehicle control (Fig. 4F). Conversely, the Hyp, which exhibited no changes in bacterial loads or in the expression of polyamine synthesis enzymes following injection, was agnostic to the ability of bacteria to synthesize polyamines. Bacteria injection, whether treated with DFMA-DFMO or untreated, resulted in an increase in the expression levels of *il-1b*, *tnfa*, and *il-6*, with fold changes ranging from 2 to 3, 4 hpi (Fig. 4C, E and G). Subsequently, the expression levels of these cytokines returned to their normal baseline within 1 day post-injection. These results suggest that polyamines derived from systemic bacteria and/or from the bacteria that penetrate the Tel directly modulate pyrogenic cytokine responses in the Tel but not the Hyp.

Systemic bacteria induce behavioral fever in rainbow trout in a polyamine-dependent manner

Behavioral fever is teleosts has been demonstrated in several models including viral and bacterial infections, killed bacterial challenges, and exposure to inflammatory agents like LPS or zymosan (30, 32, 33). However, whether the presence of microbiota in the systemic compartment induces behavioral fever in teleosts is currently unknown. We utilized a specialized annular thermal preference tank recently developed to document behavioral changes during fish fever (33) to assess the responses of four experimental groups (Fig. 4H). We observed that both i.v injection of gut commensals and LPS isolated from gut bacteria induced behavioral fever. Both groups reached a peak preferred temperature of 16°C (Fig. 4 I to L). However, fish injected i.v with gut commensals exhibited a more rapid onset of warm-seeking behavior compared to the LPS-injected group (Supplementary figure 4A). Two lethargy indicators were also evident in the commensal-injected and LPS-injected groups. Both groups displayed a marked decrease in swimming mobility and velocity (Fig. 4 K and L). Notably, maximum lethargy induced by systemic commensals occurred 2 hours earlier than by LPS (fig. S6, A and B). In contrast, treatment of bacteria with DFMA-DFMO prior to i.v injection prevented the increase in thermal preference and fever-associated lethargy behaviors. These fish showed no significant deviation from those behaviors in the vehicle-injected group (Fig. 4 I to L). Across all measured parameters, the response of animals injected with DFMA-DFMO treated commensals mirrored the behavioral patterns shown in the vehicle-injected group throughout the 24-hour study duration (Fig. 4 I, fig. S6, A and B). Together, these experiments revealed that 1) systemic inflammation induces behavioral fever in trout, and 2) commensal-derived polyamines are necessary to induce behavioral fever in trout under a scenario where systemic inflammation facilitates microbial translocation into the telencephalon.

Discussion

Sickness behaviors and fever are intricate, coordinated neurobehavioral processes that enhance
survival rates in vertebrates (12, 29, 69–72). As such, the vertebrate brain must sense and
respond to peripheral and systemic insults to help the host return to homeostasis. The complex
neuronal circuitry and immune mediators involved in mammalian sickness behaviors are fairly
well characterized, reaching higher order brain regions beyond the Hyp (9–13). However, the
neuroimmunological basis for sickness behaviors in ectotherms remains poorly understood.
325
Teleost fish, like other ectotherms, display sickness behaviors in response to infection
characterized by reduced locomotor activity, social withdrawal, reduced appetite and behavioral
fever (29, 32, 33, 35). Until now, the POV of the Hyp has been the only brain region implicated in
responses during sickness behaviors and fever in teleosts (30–32, 73, 74). We hypothesized that
higher order processing regions of the teleost brain, specifically the Tel, must participate in the
neuroimmune response to systemic inflammation.
332
Here we present two novel acute systemic inflammation models in rainbow trout that mimic the
presence of leaked microbiota or their products in the systemic compartment. Since teleost fish
are highly refractory to *E. coli* LPS injections (47–49), we reasoned that direct injection of
commensals or commensal-derived LPS i.v would be a more biologically relevant model of acute
systemic inflammation than commercial LPS sources. In support, both *in vivo* models triggered
canonical pyrogenic cytokine responses in the Hyp by 4 hours, as previously demonstrated by
others using infection models (35, 40). In contrast, while *il-1b* and *tnfa* expression also increased
in the Tel, *il-6* responses were not induced. In mammals, IL-6 can have pro- and anti-
inflammatory roles in the brain (75–77). When injected systemically or into the brain, IL-6 causes
some sickness behaviors in mammals (78), with unique biological activities compared to IL-1β
(78). For instance, following a high dose of LPS injection, IL-1β knockout mice display reduced
fever compared to controls, however, IL-6 knockout mice have normal fever responses (79).
Additionally, IL-6 does not induce septic shock (80, 81) and injection of anti-IL-6 directly into the
anterior POA suppresses PGE2 (82) and Cox2 (83). In teleosts, we know very little about the
biological functions of IL-6. However, a study found that trout recombinant IL-6 reduces *il-1b* and
tnfa expression in macrophages *in vitro* (84), suggesting that IL-6 responses dampen
inflammation in trout. Assuming a similar function in the brain, *il-6* expression in the hippocampus
would reduce *il-1b* and *tnfa* expression creating a negative feedback loop that limits inflammation,
whereas in the Tel, *il-1b* and *tnfa* effects should be prolonged. Intriguingly, trout have two fish-
specific IL-6 family genes, the ciliary neurotrophic factor (CNTF-like) and M17 and CNTF-like is
highly expressed in the brain (85). Thus, we cannot rule out that one or both molecules play
alternative or complementary functions to canonical IL-6 in the teleost brain.
355
The BBB becomes leaky in response to systemic inflammation such as LPS injection, a
phenomenon well-documented in mammals (50, 51). Loss of BBB integrity in response to LPS
injection varies across different brain regions in mammals (50). Similarly, our studies indicate that
the teleost BBB vulnerability to systemic inflammation varies among brain regions and that the
Tel BBB is particularly susceptible to disruption in rainbow trout. Whether this is due to the unique
cellular and molecular composition of the Tel's BBB remains open for further investigation.
Interestingly, our study surprisingly found that, in rainbow trout, circulatory bacteria transiently
penetrate the Tel but not other brain regions under both acute (intravenous injection) and chronic
(DSS-induced colitis) conditions. Prior research identified the insular cortex as a critical detector
of peripheral inflammation in mice during DSS-induced colitis (16). However, this study did not
explore whether bacterial translocation from the gastrointestinal tract to the bloodstream resulted
in consequent bacterial infiltration into the brain. Nonetheless, the rapid and transient increase in
bacterial loads observed in the trout Tel suggests a controlled mechanism by which the teleost
brain may directly detect or sample changes in blood-borne bacteria. Whereas bacterial presence
in the mammalian brain is thought to be a sign of disease (86–89), regulated bacterial incursions
may not significantly harm the teleost brain. This hypothetical scenario is supported by the fact
that, compared to mammals, the teleost brain harbors a higher number of immune cells under
normal conditions (90) and possesses remarkable regenerative capabilities following injury (91–
95).

375
We have recently reported that the trout brain has a resident bacterial community (55). Here we
show that significant shifts in microbial community composition occur in the trout Tel following an
i.v injection of gut bacteria. Following i.v injection, some bacteria will rapidly be eliminated by the
blood immune system. Yet, we noted changes in the telencephalic microbial community 4 hpi that
persisted by 1 day. We noted expansions of specific taxa such as Ruminococcaceae,
Streptococcaceae and Mycoplasmataceae at 4 hours, that were mostly undetectable by 1 day
indicating a transient colonization. This may be explained by rapid elimination of bacteria by the
trout brain's innate immune response or failure to outcompete other resident microbes. In
contrast, Fusobacteriaceae and Chitinibacteraceae were expanded in the Tel after 1 day,
potentially indicating longer-term dysbiosis in the brain following a single systemic inflammation
event. We currently do not know how systemic bacteria penetrate the trout Tel, although our data
suggest that the greater BBB disruption of the Tel may facilitate bacteria penetration. In support,
we recently reported that mature Chinook salmon, where the gut barrier is disrupted, LPS levels
increase in circulation and bacterial loads increase in the Tel (55). Thus, our findings suggest that
teleosts may experience shifts in brain microbiomes throughout their lifespan, some associated
with physiological and developmental changes, some because of stress, inflammation or
infection.
393
Polyamines are polycationic compounds produced by eukaryotic and prokaryotic cells. In the
vertebrate nervous system, polyamines are involved in neurodevelopment, synaptic plasticity,
cognition, aging, and behavior (96–99). Importantly, polyamine metabolism and accumulation of
putrescine are thought to be a protective biochemical program following brain insults in both
mammals and fish (100–103). Our results indicate that specific regions of the trout brain (the Tel),
but not others (the Hyp) quickly respond to systemic inflammatory cues by shifting polyamine
metabolism. Specifically, we observed an accumulation of putrescine and a decrease in
spermidine and spermine levels in the Tel. In support, putrescine was shown to accumulate in the
goldfish brain following cold exposure suggesting that this protective response helps fish adapt to
environmental changes (100). We also noted drastic decreases in telencephalic spermine levels.
Since spermine inhibits proinflammatory cytokine responses, at least in humans (104),
decreasing spermine levels in the Tel may be an effective mechanism to ensure pyrogen
production, a hypothesis that merits further investigation. It is intriguing that changes in polyamine
metabolism were not evident in the Hyp, reinforcing the notion that compartmentalized responses
to systemic inflammation occur in the teleost brain, with the Tel being the focal point of
polyamine-mediated responses.
410
The role of polyamines in mammalian fever responses remains unclear, with studies concluding
either a lack of function as pyretics (105) or their antipyretic functions (106, 107). However,
polyamines are precursors to gamma-aminobutyric acid (GABA) and GABA is known to act on
the POA regulating body temperature by inhibiting heat production in mammals exposed to hot
temperatures (108, 109). Interestingly, a recent study in human ulcerative colitis (UC) patients
detected significant increases in serum N1-acetylspermidine levels, a bacterial-derived
polyamine, in patients with moderate to severe UC compared to those with inactive disease (110)
suggesting a role for N1-acetylspermidine in the pathology of UC or as a biomarker for disease
severity. Interestingly, 40% of UC patients suffer from low fever of unknown sources (111) and
therefore the possibility that bacterial-derived polyamines mediate this phenomenon is worth
further investigation. A critical question of our study is whether the changes in polyamine
metabolism detected in the Tel as well as subsequent behavioral fever responses require the
incursion of microbiota into the Tel or they occur independently, for instance in response to the
presence of bacteria in the systemic compartment Answering this question would require injecting
bacteria mutants that lack the ability to enter the Tel so that we can disentangle systemic versus
direct Tel responses.
427
428 **Conclusion**
429

In conclusion, the present study suggests that polyamine metabolism ties immune regulation and
behavioral adaptations to inflammation in teleosts, the Tel being the hub for this coordinated
response. Yet, many questions remain to be answered, including the exact neuronal inhibitory
and excitatory circuitry that connects the POV with the Tel during sickness behaviors and fever in
ectotherms. Given that teleosts appear to live in symbiosis with bacteria during healthy states, as
evidenced by their blood and brain microbiomes, we postulate that the Tel acts as a major brain
sensor of systemic inflammatory states, allowing teleosts to rapidly respond to shifts in blood
bacterial levels deviating from homeostasis.
438
439
440   **Materials and Methods**
441
442   Isolation of gut bacteria from juvenile rainbow trout
443
Rainbow trout, initially weighing 0.2 grams, were sourced from Trout Lodge (Washington, USA).
They were maintained at a constant temperature of 16°C and fed daily with 2% of their body
weight using commercial trout diet (Skretting). Feeding was discontinued one day prior to the
collection of gut commensal bacteria. Trout (N=3 per collection group, mean body weight 23±3g)
were euthanized using an overdose of MS-222 in water (Syndel, 200 ppm). Animals were then
perfused by a cardiac injection of sterile PBS-heparin solution (Sigma Aldrich) to clear any blood
from the gut-associated vasculature. Subsequently, the intestines were carefully removed and
dissected lengthwise. Following mucus scraping, bacteria were isolated as previously described.
The bacterial pellet was then washed in triplicate with a 1% PBS-BSA solution (pH 7.2),
resuspended in sterile PBS and all five pellets were combined. The suspension was adjusted to
achieve an optical density (OD) of 0.4 and injected using heparinized syringes.
455
456   Inhibition of polyamine synthesis in isolated trout gut bacteria
457
To irreversibly suppress polyamine synthesis in isolated gut commensals from rainbow trout, we
used the inhibitors DFMA (α-difluoromethylarginine) and DFMO (α-difluoromethylornithine) as
previously described (65, 112, 113). Pilot experiments were performed to assess the growth
kinetics of isolated gut bacteria in the presence of varying concentrations of each
pharmacological agent tested independently (1 mM, 0.5 mM, 0.25 mM, 0.1 mM, 0.05 mM, 0.025
mM, 0.01 mM, and a control vehicle (PBS). Gut bacteria were adjusted to an OD of 0.05 in TSB
and XX ul of the suspension were pipetted onto sterile flat-bottom 96-well plates (N=4/treatment).
Plates were incubated at 30°C under continuous shaking. Optical density was measured every
three hours for 24 hours in an iMark microplate reader (Bio-rad). Next, we evaluated the effects of
growing gut bacteria with equimolar concentrations DFMA and DFMO concurrently using the
same experimental set-up. These experiments identified the maximum concentration of DFMA
and DFMO that could be used without impacting trout gut bacterial growth, which was 0.25 mM
each.

The bacterial cultures were therefore incubated with a solution containing 0.25 mM of each DFMA
and DFMO in phosphate-buffered saline (PBS) at a temperature of 4°C for a duration of 2 hours
as previously described. Following the treatment period, the bacteria were then pelleted through
centrifugation at 7,500 g for 20 minutes at 4°C. Next, the bacterial pellets underwent three
separate washes with PBS. After each wash, the suspended bacterial cells were again
centrifuged at 7,500 g for 20 minutes at 4°C to ensure the complete removal of any residual
DFMA and DFMO. The optical density of the DFMA-DFMO treated bacterial suspension was
adjusted to OD=0.4. This standardized bacterial preparation was then administered to the
animals by i.v injection into the caudal vein.
480
481   Lipopolysaccharide (LPS) Extraction
482
Isolated bacteria from rainbow trout gut were pellet down by centrifugation at 4°C for 25 min at
10000 g. LPS was isolated using the LPS Isolation Kit (MAK339, Millipore Sigma). In brief, the

bacterial pellet underwent lysis via continuous pulse sonication for three 30-second intervals at 10
watts, utilizing a lysing buffer from the kit at a ratio of 1/10 mg/µL pellet to lysing buffer. Next,
samples were incubated on ice for 10 minutes and subsequently centrifuged at 4°C for 10
minutes at 2,500 g. The supernatant was carefully transferred to a new 1.5 mL tube. Proteinase K
was then added to achieve a final concentration of 0.1 mg/mL, and the mixture was incubated at
60°C for 60 minutes. Following incubation, the heated lysate was centrifuged at 4°C for 10
minutes at 2,500 g, and the supernatant was subjected to vacuum drying at 25°C. The dried
sample was then reconstituted in an appropriate vehicle. The purity of the extracted LPS was
evaluated using the total carbohydrate assay kit (MAK104, Millipore Sigma) which determined
that the purity was >99.5%.
495
496 In vivo systemic inflammation models
497
Juvenile trout (23 ±3.2 g body weight) were first sedated by MS-222 anesthesia in water (Syndel,
100 ppm). Animals (N=7/experimental group) were i.v injected with 70 µl of vehicle, isolated
commensal bacteria (OD=0.4), LPS or DFMA-DFMO treated bacteria (OD=o.4) through the
caudal vein. After 4 hours and 1 day, trout were euthanized by an overdose of 200 ppm MS-222
anesthesia (Syndel). Prior to tissue collection, animals were transcardiaclly perfused with sterile
PBS-heparin solution (Sigma Aldrich) prior to tissue collection. The injections comprised 70 µl of
either a vehicle, isolated commensal bacteria with OD=0.4, lipopolysaccharide (LPS), or bacteria
treated with DFMA-DFMO at an OD of 0.4. Trout were euthanized at 4 hours and 1 day post-
injection by an overdose of MS-222 in water until opercular movement stopped (Syndel). Prior to
tissue collection, a sterile PBS-heparin solution (Sigma Aldrich) was used for cardiac perfusion to
ensure the removal of blood and maintain tissue integrity.
509
510 To model chronic dysbiosis in trout, dextran sodium sulfate (DSS, MP Biomedicals) was
511 employed to induce colitis-like symptoms following the protocol by Ding *et al.* (56). This involved
512 administering a 3% DSS solution (by weight/volume in water) by oral gavage to the fish for a
513 consecutive period of 10 days. Throughout the duration of the experiment, fish were fed daily 2%
514 of their total body mass.
515
516 All animal procedures were approved by the University of New Mexico Institutional Animal Care
517 and Use Committee under protocol number 22-201239-MC.
518
519 Quantification of serum LPS levels
520
To assess the serum LPS levels, serum samples were first diluted at a 1:20 ratio using endotoxin
free PBS (Millipore). Diluted serum samples were then subjected to a heat inactivation step,
maintained at 70°C for 15 min. Following this, the endotoxin quantities in the serum were
determined using the Pierce™ chromogenic endotoxin quantification assay (Thermo Fisher
Scientific) according to manufacturer's instructions. To prevent any interference from pyrogenic
substances, all laboratory consumables utilized in this process, such as pipette tips, Eppendorf
tubes, and microtiter plates, were pre-validated to confirm their non-pyrogenicity.
528
529 Quantitative real-time PCR for host gene expression profiling
530
Total RNA was extracted from the dissected tissues by the Trizol method. Tissue lysis was
performed by the addition of two tungsten beads to 1.7 mL Eppendorf tubes containing the tissue
and Trizol. The mixture was then subjected to vigorous shaking at 30 Hz for 5 minutes. Next, the
homogenates/lysates were collected, and RNA extraction was performed following a standard
chloroform/phenol extraction protocol, as delineated in previous studies. Total RNA was then
quantified using a Nanodrop spectrophotometer. For cDNA synthesis, 1 µg of RNA underwent
reverse transcription employing the Superscript III First Strand Synthesis System (Thermo Fisher
Scientific, Catalog No. 18080051). Quantitative PCR (qPCR) was subsequently done using the
SSOadvanced Universal SYBR Green Supermix (Bio-Rad, Catalog No. 1725270), with the

annealing temperature set at 62°C using specific primers (Supplementary table 1) for the
following genes: *il-1b, il-6, tnfa, odc1, odc2, samdc1, samdc2*. Changes in gene expression were
quantified employing the Pfaffl method using *efa1* as the reference gene (114).
543
544 DNA extraction and 16S rDNA library preparation
545
All the aseptic techniques were strictly practiced as previously described (55, 115). To lyse the
specimens a pair of tungsten carbide beads was added to each tube, followed by vigorous lysis at
30 Hz for five minutes using the TissueLyser II (Qiagen). Post-mechanical lysis, we introduced
200 µl of a 1% CTAB (hexadecyltrimethylammonium bromide, Sigma) and 3 µl of proteinase K
solution (100 mg/ml, Sigma) into each tube. The DNA extraction protocol was conducted in line
with procedures previously established by Mitchell & Takacs-Vesbach (116). The purity and
quantity of the extracted DNA were assessed using the Nanodrop ND 1000 spectrophotometer,
with subsequent dilution of the DNA at either a 1:10 or 1:20 ratio. Our library construction for
sequencing included both negative and positive controls, consistent with established norms within
the microbiome research criteria (117, 118). To mitigate the influence of possible contaminants,
we standardized the template DNA concentration for each PCR reaction at a minimum threshold
of 200 ng. Despite the negative controls not meeting this concentration, the extraction elution was
still subjected to PCR amplification. PCR was executed in triplicate for each biological specimen,
with the resultant amplicons being combined before undergoing cleaning and library preparation.
Every sequencing batch incorporated a positive control comprising DNA from seven laboratory-
cultured bacterial strains to monitor sequencing accuracy. Additionally, a previously sequenced
mouse colon DNA sample was used as a control in each sequence run to ensure methodological
consistency.
564
To amplify the V1-V3 regions of the prokaryotic 16S rDNA, three distinct PCR reactions were
performed on each sample using the designated primers 28F and 519R, as previously detailed
(56). The resulting amplification mixtures were then pooled and subjected to a clean-up process
by the AxyPrep Mag PCR Clean-up Kit (Thermo Fisher Scientific). The PCR products were then
tagged using the Nextera XT Index Kit v2 Sets A to B (Illumina), with the amplicon concentrations
normalized to 200 ng/µl via the Qubit high sensitivity dsDNA assay before pooling. An additional
clean-up stage was done using the Axygen PCR clean-up system. The library was then
sequenced on the Illumina NextSeq 2000 platform, by the 600-cycle reagent kit, at the Clinical
and Translational Sciences Center, University of New Mexico Health Sciences Center.
574
575 Bacterial load quantification in trout tissues
576
Bacterial loads within each sample were quantified via quantitative PCR targeting the 16S rDNA
as previously described (55, 119). This involved the amplification of bacterial 16S rDNA
sequences in triplicate, using primers specific to the V1-V3 hypervariable regions as established
in prior research (56). The qPCR mixture was prepared in a 96-well format, containing 2 µl of
standardized DNA/cDNA (10 ng/µl), a 2 µl aliquot of the primer mix, 6 µl of nuclease-free water,
and 10 µl of SsoAdvanced Universal SYBR Green supermix (Bio-Rad). The amplification cycle
commenced with an enzyme activation at 94°C for 1.5 minutes, followed by 33 cycles each
consisting of denaturation at 94°C for 30 seconds, annealing at 52°C for 30 seconds, and
extension at 72°C for 1.5 minutes. This was concluded by a final extension period at 72°C for 7
minutes and a subsequent cooling phase at 4°C. The process was carried out using the Bio-Rad
CFX96 C1000 Touch Thermal Cycler. For quantification, the number of 16S rDNA copies in each
sample was determined by comparison to a standard curve, which was generated using a range
of E. coli 16S rDNA gene copies from 10^9 down to 10 for the V1-V3 variable regions. All runs
included negative controls with water as template.
Microbiome data analyses
592
Microbial sequence data were analyzed using the Quantitative Insights Into Microbial Ecology 2
package (Qiime2, version 2023.7) (120). The Divisive Amplicon Denoising Algorithm 2 (DADA2)
was utilized for partitioning demultiplexed reads into distinct amplicon sequence variants (ASVs)

as described (121). These ASVs were then aligned against the Silva 16S rDNA reference
database (version 138) for taxonomic characterization (122). Before proceeding to in-depth
diversity analyses, samples were normalized to a uniform sampling depth of 4,000 reads per
sample to ensure comprehensive rarefaction. The ensuing core diversity assessment was
conducted while considering both temporal and experimental treatment factors. Using QIIME2,
we computed various alpha diversity indices, including Shannon diversity, Chao1 richness,
Pielou's evenness index, and Faith's phylogenetic diversity (PD). Beta diversity analyses
(Weighted UniFrac distances) were performed in QIIME2 and principal coordinates analysis
(PCoA) plots were visualized by the Qiime2R package in RStudio (version 1.3.959) (123). To
infer the potential functional profiles of the microbial communities, we employed the PICRUSt2
(124) algorithm in Linux and visualized the outputs in RStudio (version 1.3.959) (123).
607
608 Evan's blue injection and extravasation quantification protocol
609
A 1% (w/v) stock solution of Evan's blue dye was prepared in sterile PBS and subsequently
filtered through 0.2 μm sterile filters to ensure purity. For the experimental procedure, 75 μL of the
dye, either in the vehicle or mixed with the gut bacteria suspension, was injected into the caudal
vein of rainbow trout, with each fish weighing approximately 10 g (± 1.35). The fish were then
subjected to sampling at 4 hours and 1 day. Following euthanasia as above, each fish was
perfused with a sterile perfusion solution directly through the heart to ensure the thorough
removal of residual blood and the albumin-Evan's blue complex from the vasculature. Prior to
perfusion, the caudal vein and secondary branchial arch veins were incised to facilitate removal
of any residual blood. Subsequently, the olfactory bulbs, telencephalon, hypothalamus, optic
tectum and cerebellum were dissected. The weight of each dissected brain part was recorded,
and the tissues were immediately immersed in 50% (w/v) trichloroacetic acid (TCA) and
preserved on ice as described (52). Next, the tissues were subjected to a mechanical lysis step
consisting of a 5-minute shaking at a frequency of 30 Hz. The resulting homogenate was then
centrifuged at 10,000 rpm for 20 minutes at 4°C (52). The quantification of extravasated Evans
Blue in each sample was achieved through colorimetry, comparing the absorbance of the
samples to a standard curve generated from a known serial dilution of Evans Blue in 50% TCA at
610 nm (53). The calculated dye concentrations were then normalized to the weight unit of each
sample, allowing for accurate comparative analysis between samples.
628
629 Polyamine quantification by ELISA assay
630
631 Collected specimens were lysed using 1% Triton X-100 (v/v) (Sigma) as described (125). The
632 polyamine profile of each sample including putrescine, spermidine, and spermine levels were
633 quantified employing commercially available ELISA kits (MyBioSource) following the
634 manufacturer's instructions.
635
636 Behavioral assays
637
Rainbow trout (40±7.6 g), were acquired from Smoky Farm Ltd. (Red Deer, Alberta, Canada).
Upon arrival, the fish were accommodated in opaque, continuous flow-through aquaria
maintained at temperatures of 10-12°C. These aquaria were equipped with constant aeration to
ensure appropriate living conditions. A minimum acclimatization period of 14 days post-transport
was held, during which the fish were subject to continuous monitoring. The dietary regimen
consisted of floating pellets, administered once daily (2% of body weight) to provide the
necessary nutrients.
645
Water quality was rigorously controlled, with dissolved oxygen levels maintained between 6.5-7.0
ppm and pH levels sustained within the range of 6.8 to 7.8. Prior to any experimental
manipulation, fish were randomly selected, netted, and anesthetized using a 100 ppm tricaine
methane sulfonate solution (Syndel), adhering to the ethical guidelines sanctioned by the
Canadian Animal Care and Use Committees as well as the Science Animal Support Services.

Cervical dislocation was employed as the method of sacrifice, ensuring adherence to ethical
standards aimed at minimizing animal distress during experimental procedures.
653
All specimens were housed within the Aquatic Facility of the Department of Biological Sciences at
the University of Alberta. The facility replicated natural light conditions, providing a photoperiod of
12 hours of light followed by 12 hours of darkness. Experimental interventions included
intravenous injections of live commensal bacteria (OD = 0.4), isolated LPS (sourced from the
same stock as utilized by the Salinas laboratory), or commensal bacteria treated with
DMFA/DMFO, alongside a vehicle control (N=5). Each fish was then placed in an annular thermal
preference tank (ATPT) for a 24-hour behavior analysis period as described (33).
661
Behavioral assessments were conducted using high-resolution positional tracking, facilitated by
an infrared Panasonic CCTV color camera (model WV-CP620) equipped with a 2X variable focal
lens (model WV-LZ61/2S) and a centrally positioned overhead lighting system. Video recording
was achieved using Bandicam software (Bandicam Company, USA). The analysis of recorded
videos was performed with Ethovision XT, Version 11 (Noldus, Wageningen, Netherlands),
allowing for the quantification of behavioral parameters such as changes in thermal preference,
swimming velocity, and the frequency of transitions between thermal zones as previously
described (33).
670
671 Statistical analyses
672
To assess the levels of endotoxin in blood serum among various groups, we initially conducted
tests for normality using both the F-test and Bartlett's test. Group differences were evaluated
using either unpaired t-tests or one-way Analysis of Variance (ANOVA), with subsequent analysis
through Tukey's post hoc test for specific comparisons. To evaluate bacterial loads, indicated by
16S gene copy numbers in different tissues, data underwent checks for normal distribution as
mentioned. To examine variations across tissues and between groups' bacterial loads, we utilized
both the Brown-Forsythe and Welch's ANOVA tests; the same methodology was utilized to
analyze the behavioral data. To analyze differences in Shannon Diversity indices, we first
performed normality tests as mentioned earlier. Due to the non-parametric nature of these data,
differences among groups were determined using the Kruskal-Wallis test. In instances of
significant disparities, post hoc pairwise analysis was conducted, applying corrections for multiple
comparisons. For visual interpretation of Weighted UniFrac distances, Principal Coordinates
Analysis (PCoA) plots were used. Significance in clustering differences on these plots between
groups was ascertained through multivariate dispersion analysis. All statistical evaluations were
performed using GraphPad Prism (version 10.0.3) and RStudio (version 4.2.3), considering a P-
value of less than 0.05 as statistically significant.
689
690 Data availability
691
692 All microbiome sequencing data is available from NCBI under BioProject number
693 PRJNA1079394.
694
695
696 **Acknowledgments**
697

698 Authors thank Darrel Dinwiddie and Kurt Buchmann for support with the Illumina library
699 sequencing and the University of New Mexico Center for Advanced Research Computing (CARC)
700 for computing resources and data storage. Schematics were made in Biorender.

707
708 **References**
709
710
1. S. S. Chavan, V. A. Pavlov, K. J. Tracey, Mechanisms and therapeutic relevance of
neuro-immune communication. *Immunity* **46**, 927–942 (2017).

2. V. A. Pavlov, K. J. Tracey, Neural regulators of innate immune responses and
inflammation. *Cell. Mol. life Sci. C.* **61**, 2322–2331 (2004).

3. I. J. Elenkov, R. L. Wilder, G. P. Chrousos, E. S. Vizi, The sympathetic nerve—an
integrative interface between two supersystems: the brain and the immune system.
Pharmacol. Rev. **52**, 595–638 (2000).

4. J. Roth, G. E. P. De Souza, Fever induction pathways: evidence from responses to
systemic or local cytokine formation. *Brazilian J. Med. Biol. Res.* **34**, 301–314 (2001).

5. A. M. Vasilache, H. Qian, A. Blomqvist, Immune challenge by intraperitoneal
administration of lipopolysaccharide directs gene expression in distinct blood–brain barrier
cells toward enhanced prostaglandin E2 signaling. *Brain. Behav. Immun.* **48**, 31–41
(2015).

6. Z. Fitzpatrick, *et al.*, Gut-educated IgA plasma cells defend the meningeal venous sinuses.
Nature **587**, 472–476 (2020).

7. A. Kertser, *et al.*, Corticosteroid signaling at the brain-immune interface impedes coping
with severe psychological stress. *Sci. Adv.* **5**, eaav4111 (2019).

8. A. Wang, R. Medzhitov, Counting calories: the cost of inflammation. *Cell* **177**, 223–224
(2019).

9. C. B. Saper, A. A. Romanovsky, T. E. Scammell, Neural circuitry engaged by
prostaglandins during the sickness syndrome. *Nat. Neurosci.* **15**, 1088–1095 (2012).

10. D. Mittli, Inflammatory processes in the prefrontal cortex induced by systemic immune
challenge: focusing on neurons. *Brain, Behav. Immunity-Health* **34**, 100703 (2023).

11. C. Cao, K. Matsumura, K. Yamagata, Y. Watanabe, Endothelial cells of the rat brain
vasculature express cyclooxygenase-2 mRNA in response to systemic interleukin-1β: a
possible site of prostaglandin synthesis responsible for fever. *Brain Res.* **733**, 263–272
(1996).

12. R. Dantzer, J. C. O'connor, G. G. Freund, R. W. Johnson, K. W. Kelley, From
inflammation to sickness and depression: when the immune system subjugates the brain.
Nat. Rev. Neurosci. **9**, 46–56 (2008).

13. J. A. Osterhout, *et al.*, A preoptic neuronal population controls fever and appetite during
sickness. *Nature* **606**, 937–944 (2022).

14. A. Garami, A. A. Steiner, A. A. Romanovsky, Fever and hypothermia in systemic
inflammation. *Handb. Clin. Neurol.* **157**, 565–597 (2018).

15. C. L. Tan, *et al.*, Warm-sensitive neurons that control body temperature. *Cell* **167**, 47–59
(2016).

16. T. Koren, *et al.*, Insular cortex neurons encode and retrieve specific immune responses.
Cell **184**, 5902–5915 (2021).

17. F. Frenois, *et al.*, Lipopolysaccharide induces delayed FosB/DeltaFosB immunostaining
within the mouse extended amygdala, hippocampus and hypothalamus, that parallel the
expression of depressive-like behavior. *Psychoneuroendocrinology* **32**, 516–531 (2007).

18. K. M. Ibrahim, *et al.*, Dorsal hippocampus to nucleus accumbens projections drive
reinforcement via activation of accumbal dynorphin neurons. *Nat. Commun.* **15**, 750
(2024).

19. M. Fritz, *et al.*, Structural and biochemical imaging reveals systemic LPS-induced changes
in the rat brain. *J. Neuroimmunol.* **348**, 577367 (2020).

20. S. N. Kurki, *et al.*, A brain cytokine-independent switch in cortical activity marks the onset
of sickness behavior triggered by acute systemic inflammation. *J. Neuroinflammation* **20**,
176 (2023).

21. M. R. Smith, *et al.*, Integrative analysis of disease signatures shows inflammation disrupts
juvenile experience-dependent cortical plasticity. *eneuro* **3** (2016).

22. M. Mojzesz, *et al.*, Tilapia lake virus-induced neuroinflammation in Zebrafish: Microglia
Activation and Sickness Behavior. *Front. Immunol.* **12** (2021).

23. K. Kirsten, S. M. Soares, G. Koakoski, L. C. Kreutz, L. J. G. Barcellos, Characterization of
sickness behavior in zebrafish. *Brain. Behav. Immun.* **73**, 596–602 (2018).

24. P. C. Lopes, S. S. French, D. C. Woodhams, S. A. Binning, Sickness behaviors across
vertebrate taxa: proximate and ultimate mechanisms. *J. Exp. Biol.* **224**, jeb225847 (2021).

25. D. Llewellyn, G. P. Brown, M. B. Thompson, R. Shine, Behavioral responses to immune-
system activation in an anuran (the cane toad, Bufo marinus): field and laboratory studies.
Physiol. Biochem. Zool. **84**, 77–86 (2011).

26. C. M. Deen, V. H. Hutchison, Effects of lipopolysaccharide and acclimation temperature
on induced behavioral fever in juvenile Iguana iguana. *J. Therm. Biol.* **26**, 55–63 (2001).

27. M. J. Kluger, Fever in ectotherms: evolutionary implications. *Am. Zool.* **19**, 295–304
(1979).

28. M. Cabanac, F. Laberge, Fever in goldfish is induced by pyrogens but not by handling.
Physiol. Behav. **63**, 377–379 (1998).

29. J. B. Covert, W. W. Reynolds, Survival value of fever in fish. *Nature* **267**, 43–45 (1977).

30. S. Boltana, *et al.*, The expression of TRPV channels, prostaglandin E2 and pro-
inflammatory cytokines during behavioural fever in fish. *Brain. Behav. Immun.* **71**, 169–
181 (2018).

31. K. Rakus, *et al.*, Conserved fever pathways across vertebrates: a herpesvirus expressed
decoy TNF-α receptor delays behavioral fever in fish. *Cell Host Microbe* **21**, 244–253
(2017).

32. S. Rey, V. Moiche, S. Boltana, M. Teles, S. MacKenzie, Behavioural fever in zebrafish
larvae. *Dev. Comp. Immunol.* **67**, 287–292 (2017).

33. F. Haddad, *et al.*, Fever integrates antimicrobial defences, inflammation control, and
tissue repair in a cold-blooded vertebrate. *Elife* **12**, e83644 (2023).

34. N. Sanhueza, *et al.*, Behavioural fever promotes an inflammatory reflex circuit in
ectotherms. *Int. J. Mol. Sci.* **22**, 8860 (2021).

35.   K. Rakus, M. Ronsmans, A. Vanderplasschen, Behavioral fever in ectothermic
791   vertebrates. *Dev. Comp. Immunol.* **66**, 84–91 (2017).

792   36.   Y. Ishikawa, N. Yamamoto, H. Hagio, "Functional Neuroanatomy of Teleost Brains" in
793   *Brain Development of Medaka Fish: A New Concept of Brain Morphogenesis in*
794   *Vertebrates*, (Springer, 2022), pp. 231–248.

795   37.   L. S. Demski, The pallium and mind/behavior relationships in teleost fishes. *Brain Behav.*
796   *Evol.* **82**, 31–44 (2013).

797   38.   J. C. Prechtl, *et al.*, Sensory processing in the pallium of a mormyrid fish. *J. Neurosci.* **18**,
798   7381–7393 (1998).

799   39.   M. Portavella, B. Torres, C. Salas, Avoidance response in goldfish: emotional and
800   temporal involvement of medial and lateral telencephalic pallium. *J. Neurosci.* **24**, 2335–
801   2342 (2004).

802   40.   E. Durán, F. M. Ocaña, C. Broglio, F. Rodríguez, C. Salas, Lateral but not medial
803   telencephalic pallium ablation impairs the use of goldfish spatial allocentric strategies in a
804   "hole-board" task. *Behav. Brain Res.* **214**, 480–487 (2010).

805   41.   C. Broglio, F. Rodríguez, A. Gómez, J. L. Arias, C. Salas, Selective involvement of the
806   goldfish lateral pallium in spatial memory. *Behav. Brain Res.* **210**, 191–201 (2010).

807   42.   B. Rodríguez-Expósito, *et al.*, Goldfish hippocampal pallium is essential to associate
808   temporally discontiguous events. *Neurobiol. Learn. Mem.* **139**, 128–134 (2017).

809   43.   F. Rodrıguez, *et al.*, Spatial memory and hippocampal pallium through vertebrate
810   evolution: insights from reptiles and teleost fish. *Brain Res. Bull.* **57**, 499–503 (2002).

811   44.   E. H. Van den Burg, *et al.*, Activation of a sensorimotor pathway in response to a water
812   temperature drop in a teleost fish. *J. Exp. Biol.* **209**, 2015–2024 (2006).

813   45.   S. Rey, *et al.*, Fish can show emotional fever: stress-induced hyperthermia in zebrafish.
814   *Proc. R. Soc. B Biol. Sci.* **282**, 20152266 (2015).

815   46.   B. A. Porter, T. Mueller, The zebrafish amygdaloid complex–functional ground plan,
816   molecular delineation, and everted topology. *Front. Neurosci.* **14**, 608 (2020).

817   47.   A. E. Gauthier, *et al.*, Deep-sea microbes as tools to refine the rules of innate immune
818   pattern recognition. *Sci. Immunol.* **6**, eabe0531 (2021).

819   48.   B. Novoa, T. V Bowman, L. Zon, A. Figueras, LPS response and tolerance in the zebrafish
820   (Danio rerio). *Fish Shellfish Immunol.* **26**, 326–331 (2009).

821   49.   D. B. Iliev, J. C. Roach, S. Mackenzie, J. V Planas, F. W. Goetz, Endotoxin recognition: in
822   fish or not in fish? *FEBS Lett.* **579**, 6519–6528 (2005).

823   50.   W. A. Banks, *et al.*, Lipopolysaccharide-induced blood-brain barrier disruption: roles of
824   cyclooxygenase, oxidative stress, neuroinflammation, and elements of the neurovascular
825   unit. *J. Neuroinflammation* **12**, 1–15 (2015).

826   51.   X. Peng, Z. Luo, S. He, L. Zhang, Y. Li, Blood-brain barrier disruption by
827   lipopolysaccharide and sepsis-associated encephalopathy. *Front. Cell. Infect. Microbiol.*,
828   1083 (2021).

829   52.   O. Y. Waithe, X. Peng, E. W. Childs, B. Tharakan, "Measurement of Blood-Brain Barrier
830   Hyperpermeability Using Evans Blue Extravasation Assay" in *Vascular Hyperpermeability:*

831   *Methods and Protocols*, (Springer, 2023), pp. 177–184.

832   53.   H.-L. Wang, T. W. Lai, Optimization of Evans blue quantitation in limited rat tissue
833   samples. *Sci. Rep.* **4**, 6588 (2014).

834   54.   A. Fleming, H. Diekmann, P. Goldsmith, Functional characterisation of the maturation of
835   the blood-brain barrier in larval zebrafish. *PLoS One* **8**, e77548 (2013).

836   55.   A. Mani, *et al.*, A niche-adapted brain microbiome in salmonids at homeostasis. *bioRxiv*,
837   2012–2023 (2023).

838   56.   Y. Ding, *et al.*, Secretory IgM (sIgM) is an ancient master regulator of microbiota
839   homeostasis and metabolism. *bioRxiv*, 2002–2023 (2023).

840   57.   A. J. Michael, Polyamines in eukaryotes, bacteria, and archaea. *J. Biol. Chem.* **291**,
841   14896–14903 (2016).

842   58.   Y. Y. Lenis, M. A. Elmetwally, J. G. Maldonado-Estrada, F. W. Bazer, Physiological
843   importance of polyamines. *Zygote* **25**, 244–255 (2017).

844   59.   Tt. Thomas*, T. J. Thomas, Polyamines in cell growth and cell death: molecular
845   mechanisms and therapeutic applications. *Cell. Mol. Life Sci. C.* **58**, 244–258 (2001).

846   60.   M. Xuan, *et al.*, Polyamines: their significance for maintaining health and contributing to
847   diseases. *Cell Commun. Signal.* **21**, 348 (2023).

848   61.   A. Ö. Gevrekci, The roles of polyamines in microorganisms. *World J. Microbiol.*
849   *Biotechnol.* **33**, 204 (2017).

850   62.   L. Miller-Fleming, V. Olin-Sandoval, K. Campbell, M. Ralser, Remaining mysteries of
851   molecular biology: the role of polyamines in the cell. *J. Mol. Biol.* **427**, 3389–3406 (2015).

852   63.   A. J. Michael, Biosynthesis of polyamines in eukaryotes, archaea, and bacteria. *Polyam. A*
853   *Univers. Mol. nexus growth, Surviv. Spec. Metab.*, 3–14 (2015).

854   64.   K. Igarashi, K. Kashiwagi, Polyamine modulon in Escherichia coli: genes involved in the
855   stimulation of cell growth by polyamines. *J. Biochem.* **139**, 11–16 (2006).

856   65.   A. J. Bitonti, P. P. Mc Cann, *Inhibition of polyamine biosynthesis in microorganisms*
857   (Academic Press, 1987).

858   66.   D. Soulet, S. Rivest, Polyamines play a critical role in the control of the innate immune
859   response in the mouse central nervous system. *J. Cell Biol.* **162**, 257–268 (2003).

860   67.   Y. L. Latour, A. P. Gobert, K. T. Wilson, The role of polyamines in the regulation of
861   macrophage polarization and function. *Amino Acids* **52**, 151–160 (2020).

862   68.   R. S. Hesterberg, J. L. Cleveland, P. K. Epling-Burnette, Role of polyamines in immune
863   cell functions. *Med. Sci.* **6**, 22 (2018).

864   69.   B. L. Hart, Biological basis of the behavior of sick animals. *Neurosci. Biobehav. Rev.* **12**,
865   123–137 (1988).

866   70.   B. L. Hart, Behavioral adaptations to pathogens and parasites: five strategies. *Neurosci.*
867   *Biobehav. Rev.* **14**, 273–294 (1990).

868   71.   R. Dantzer, K. W. Kelley, Twenty years of research on cytokine-induced sickness
869   behavior. *Brain. Behav. Immun.* **21**, 153–160 (2007).

870   72.   P. A. Rittenhouse, A. D. Levy, Q. Li, C. L. Bethea, L. D. Van de Kar, Neurons in the
871   hypothalamic paraventricular nucleus mediate the serotonergic stimulation of prolactin
872   secretion via 5-HT1c/2 receptors. *Endocrinology* **133**, 661–667 (1993).

873   73.   S. Boltana, *et al.*, Behavioural fever is a synergic signal amplifying the innate immune
874   response. *Proc. R. Soc. B Biol. Sci.* **280**, 20131381 (2013).

875   74.   W. W. Reynolds, M. E. Casterlin, J. B. COVERT, Behavioural fever in teleost fishes.
876   *Nature* **259**, 41–42 (1976).

877   75.   M. Erta, A. Quintana, J. Hidalgo, Interleukin-6, a major cytokine in the central nervous
878   system. *Int. J. Biol. Sci.* **8**, 1254 (2012).

879   76.   J. Scheller, A. Chalaris, D. Schmidt-Arras, S. Rose-John, The pro-and anti-inflammatory
880   properties of the cytokine interleukin-6. *Biochim. Biophys. Acta (BBA)-Molecular Cell Res.*
881   **1813**, 878–888 (2011).

882   77.   J. J. Fuster, K. Walsh, The Good, the Bad, and the Ugly of interleukin-6 signaling. *EMBO*
883   *J.* **33**, 1425–1427 (2014).

884   78.   R.-M. Bluthé, B. Michaud, V. Poli, R. Dantzer, Role of IL-6 in cytokine-induced sickness
885   behavior: a study with IL-6 deficient mice. *Physiol. Behav.* **70**, 367–373 (2000).

886   79.   W. Kozak, *et al.*, IL-6 and IL-1β in Fever: Studies Using Cytokine-Deficient (Knockout)
887   Mice a. *Ann. N. Y. Acad. Sci.* **856**, 33–47 (1998).

888   80.   B. E. Barton, J. V Jackson, Protective role of interleukin 6 in the lipopolysaccharide-
889   galactosamine septic shock model. *Infect. Immun.* **61**, 1496–1499 (1993).

890   81.   C. Libert, *et al.*, Limited involvement of interleukin-6 in the pathogenesis of lethal septic
891   shock as revealed by the effect of monoclonal antibodies against interleukin-6 or its
892   receptor in various murine models. *Eur. J. Immunol.* **22**, 2625–2630 (1992).

893   82.   A. Fernandez-Alonso, K. Benamar, M. Sancibrian, F. J. Lopez-Valpuesta, F. J. Mifiano,
894   Role of interleukin-1β, interleukin-6 and macrophage inflammatory protein-1β in
895   prostaglandin-E2-induced hyperthermia in rats. *Life Sci.* **59**, PL185–PL190 (1996).

896   83.   B. L. Fiebich, *et al.*, Prostaglandin E2 induces interleukin-6 synthesis in human
897   astrocytoma cells. *J. Neurochem.* **68**, 704–709 (1997).

898   84.   M. M. Costa, T. Maehr, P. Diaz-Rosales, C. J. Secombes, T. Wang, Bioactivity studies of
899   rainbow trout (Oncorhynchus mykiss) interleukin-6: effects on macrophage growth and
900   antimicrobial peptide gene expression. *Mol. Immunol.* **48**, 1903–1916 (2011).

901   85.   T. Wang, C. J. Secombes, Identification and expression analysis of two fish-specific IL-6
902   cytokine family members, the ciliary neurotrophic factor (CNTF)-like and M17 genes, in
903   rainbow trout Oncorhynchus mykiss. *Mol. Immunol.* **46**, 2290–2298 (2009).

904   86.   Z. Hu, C.-A. McKenzie, C. Smith, J. G. Haas, R. Lathe, The remarkable complexity of the
905   brain microbiome in health and disease. *BioRxiv*, 2002–2023 (2023).

906   87.   W. G. Branton, *et al.*, Brain microbial populations in HIV/AIDS: α-proteobacteria
907   predominate independent of host immune status. *PLoS One* **8**, e54673 (2013).

908   88.   M. Prudencio, *et al.*, Distinct brain transcriptome profiles in C9orf72-associated and
909   sporadic ALS. *Nat. Neurosci.* **18**, 1175–1182 (2015).

910   89.   R. Alonso, D. Pisa, A. M. Fernández-Fernández, L. Carrasco, Infection of fungi and

911   bacteria in brain tissue from elderly persons and patients with Alzheimer's disease. *Front.*
912   *Aging Neurosci.* **10**, 159 (2018).

913   90.   A. J. Dowding, J. Scholes, Lymphocytes and macrophages outnumber oligodendroglia in
914   normal fish spinal cord. *Proc. Natl. Acad. Sci.* **90**, 10183–10187 (1993).

915   91.   G. K. H. Zupanc, Adult neurogenesis in the central nervous system of teleost fish: From
916   stem cells to function and evolution. *J. Exp. Biol.* **224**, jeb226357 (2021).

917   92.   G. K. H. Zupanc, R. F. Sîrbulescu, Teleost fish as a model system to study successful
918   regeneration of the central nervous system. *New Perspect. Regen.*, 193–233 (2012).

919   93.   G. K. H. Zupanc, "Adult neurogenesis in teleost fish" in *Neurogenesis in the Adult Brain I*,
920   (Springer, 2011), pp. 137–167.

921   94.   M. J. Anderson, S. G. Waxman, 20 Neurogenesis in Adult Vertebrate Spinal Cord In Situ
922   and In Vitro: A New Model System. *Form Funct. Brain Spinal Cord Perspect. a Neurol.*,
923   235 (2003).

924   95.   P. Hitchcock, M. Ochocinska, A. Sieh, D. Otteson, Persistent and injury-induced
925   neurogenesis in the vertebrate retina. *Prog. Retin. Eye Res.* **23**, 183–194 (2004).

926   96.   S. Nakanishi, J. L. Cleveland, Polyamine homeostasis in development and disease. *Med.*
927   *Sci.* **9**, 28 (2021).

928   97.   T. A. Slotkin, J. Bartolome, Role of ornithine decarboxylase and the polyamines in nervous
929   system development: a review. *Brain Res. Bull.* **17**, 307–320 (1986).

930   98.   S. R. Kesler, C. Schwartz, R. E. Stevenson, A. L. Reiss, The impact of spermine synthase
931   (SMS) mutations on brain morphology. *Neurogenetics* **10**, 299–305 (2009).

932   99.   F. N. Bolkenius, N. Seiler, Developmental aspects of polyamine interconversion in rat
933   brain. *Int. J. Dev. Neurosci.* **4**, 217–224 (1986).

934   100.   S. Notari, R. Lucchi, U. Traversa, E. Fabbri, A. Poli, Reversible changes in goldfish brain
935   polyamine concentrations and synthetic enzymes after cold exposure. *Brain Res.* **1006**,
936   241–247 (2004).

937   101.   C. M. HENLEY, C. MUSZYNSKI, L. CHERIAN, C. S. ROBERTSON, Activation of
938   ornithine decarboxylase and accumulation of putrescine after traumatic brain injury. *J.*
939   *Neurotrauma* **13**, 487–496 (1996).

940   102.   D. Dasdelen, N. Cetin, E. Menevse, A. K. Baltaci, R. Mogulkoc, Effects of putrescine on
941   oxidative stress, spermidine/spermine-N (1)-acetyltransferase, inflammation and energy
942   levels in liver and serum in rats with brain ischemia-reperfusion. *Physiol. Int.* **110**, 34–45
943   (2023).

944   103.   T. Masuko, *et al.*, Polyamine transport, accumulation, and release in brain. *J. Neurochem.*
945   **84**, 610–617 (2003).

946   104.   M. Zhang, *et al.*, Spermine inhibits proinflammatory cytokine synthesis in human
947   mononuclear cells: a counterregulatory mechanism that restrains the immune response. *J.*
948   *Exp. Med.* **185**, 1759–1768 (1997).

949   105.   M. Huang, W. Zhang, H. Chen, J. Zeng, Targeting polyamine metabolism for control of
950   human viral diseases. *Infect. Drug Resist.*, 4335–4346 (2020).

106.   M. J. Dascombe, K. D. Haegele, Evidence against putrescine and polyamines as

endogenous mediators of fever. *J. Pharm. Pharmacol.* **37**, 365–366 (1985).

107.   S. Genedani, M. Bernardi, S. Tagliavini, A. Bertolini, Putrescine has hypothermic and
antipyretic activity, in rats. *Life Sci.* **38**, 1293–1298 (1986).

108.   T. A. Wang, *et al.*, Thermoregulation via temperature-dependent PGD2 production in
mouse preoptic area. *Neuron* **103**, 309–322 (2019).

109.   Z. Heli, *et al.*, Recent advances of γ-aminobutyric acid: Physiological and immunity
function, enrichment, and metabolic pathway. *Front. Nutr.* **9**, 1076223 (2022).

110.   M. Schirmer, *et al.*, Linking microbial genes to plasma and stool metabolites uncovers
host-microbial interactions underlying ulcerative colitis disease course. *Cell Host Microbe*
(2024).

111.   B. A. Hendrickson, R. Gokhale, J. H. Cho, Clinical aspects and pathophysiology of
inflammatory bowel disease. *Clin. Microbiol. Rev.* **15**, 79–94 (2002).

112.   A. S. Tyms, J. D. Williamson, C. J. Bacchi, Polyamine inhibitors in antimicrobial
chemotherapy. *J. Antimicrob. Chemother.* **22**, 403–427 (1988).

113.   T. Ponappa, D. R. Lawson, A. Raymond Miller, Inhibition of ornithine decarboxylase
activity reduces polyamine levels and growth of Agrobacterium tumefaciens. *Curr.*
Microbiol. **24**, 269–273 (1992).

114.   M. W. Pfaffl, A new mathematical model for relative quantification in real-time RT–PCR.
Nucleic Acids Res. **29**, e45–e45 (2001).

115.   C. M. Phelps, J. H. Shapira, C. R. Laughlin, M. Meisel, Detection of viable commensal
bacteria in murine melanoma tumors by culturomics. *STAR Protoc.* **4**, 102492 (2023).

116.   K. R. Mitchell, C. D. Takacs-Vesbach, A comparison of methods for total community DNA
preservation and extraction from various thermal environments. *J. Ind. Microbiol.*
Biotechnol. **35**, 1139–1147 (2008).

117.   R. Eisenhofer, *et al.*, Contamination in low microbial biomass microbiome studies: issues
and recommendations. *Trends Microbiol.* **27**, 105–117 (2019).

118.   D. Kim, *et al.*, Optimizing methods and dodging pitfalls in microbiome research.
Microbiome **5**, 1–14 (2017).

119.   O. Zemb, *et al.*, Absolute quantitation of microbes using 16S rRNA gene metabarcoding:
A rapid normalization of relative abundances by quantitative PCR targeting a 16S rRNA
gene spike-in standard. *Microbiologyopen* **9**, e977 (2020).

120.   E. Bolyen, *et al.*, Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

121.   B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat. Methods* **13**, 581–583 (2016).

122.   C. Quast, *et al.*, The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2012).

123.   J. E. Bisanz, qiime2R: Importing QIIME2 artifacts and associated data into R sessions.
Version 0.99 **13** (2018).

124.   G. M. Douglas, *et al.*, PICRUSt2 for prediction of metagenome functions. *Nat. Biotechnol.*

38, 685–688 (2020).

125.   L. Jelsbak, L. E. Thomsen, I. Wallrodt, P. R. Jensen, J. E. Olsen, Polyamines are required
for virulence in Salmonella enterica serovar Typhimurium. *PLoS One* **7**, e36149 (2012).

996
997   **Figures and Tables**

998
Figure 1: Differential responses in the trout telencephalon and the hippocampus to
systemic inflammation
(A) Schematic representation of the *in vivo* acute commensal translocation models used in the
present study. Rainbow trout received an i.v injection of gut commensal bacteria (Comm),
isolated lipopolysaccharides (LPS), or vehicle (control). Animals were sampled 4 hours or 24
hours later and blood, spleen and five regions of the brain were sampled for subsequent analysis.
(B) Changes in serum endotoxin levels post-injection with gut commensals or LPS (N= 6). Mean
fold-change in expression levels of the pro-inflammatory cytokine genes *il-1b*, *tnfa* and *il-6* in the
Tel (C, E, G) and Hyp (D, F, H) after injection of gut commensals or LPS compared to vehicle
controls as measured by RT-qPCR (N= 6). Vehicles show N=12, N=6 for 4 hours and N=6 for 1
day, with differences between each group. (I) Schematic representation of the *in vivo* Evan's blue
extravasation assay to assess the permeability of the blood-brain barrier in the different
experimental groups and brain regions. (J) Analysis of Evan's blue dye extravasation in the Tel

1016
1017
**Figure 2: Systemically injected commensals translocate into the telencephalon but not
other regions of the trout brain**
16S rRNA gene copies in the Tel (A) and Hyp (B) of rainbow trout at 4 hours and 1 day post
bacterial injection (N=6). Vehicles consist of N=12 because they include N=6 vehicle controls for
each time point with no differences among them. Statistical significance is indicated by different
letters based on Welch's ANOVA (P<0.05). Relative abundance at the phylum level (C) and
family level (D) in the trout Tel 4 hours or 1 day post-injection (N= 6). (E) Weighted UniFrac
distances showing significant differences in microbial community structure in the Tel at 4 hours
and 1 day post-injection, as determined by the Kruskal-Wallis test (P<0.05) (N=6). (F) Principal
coordinate analysis (PCoA) of the Weighted Unifrac distances, with ellipses denoting a 95%
confidence interval. (G) Linear discriminant analysis effect size (LEfSe) identifying significantly
different taxa in the Tel at specified post-injection time points. (H) Predicted metabolic pathways
significantly modified in the microbial communities of the telencephalon 4 hours or 1 day post-
injection of commensal bacteria compared to vehicle controls based on PICRUSt2 analysis
(P<0.05).

1033
1034
Figure 3: Systemic inflammation results in telencephalic polyamine responses
(A) Diagram depicting the bacterial polyamine synthesis pathway, with the inhibitory effects of
DFMA and DFMO on bacterial ornithine decarboxylase and arginine decarboxylase, respectively.
(B) Schematic diagram showing the experimental systemic inflammation model of the study
consisting of a single i.v injection of gut commensal bacteria (Comm), gut commensal bacteria
treated with DFMA-DFMO (DFM), isolated LPS (LPS), or vehicle controls. Changes in the
expression levels of polyamine synthesis enzymes *odc1* and *samdc1* in the Tel (C and D) and
Hyp (E and F) post-injection measured by RT-qPCR. The expression levels are reported as the
mean fold change compared to vehicle controls (N=7 fish/group). (G-I) Mean concentration of
putrescine, spermidine and spermine in the trout Tel 4 hours or 1 day post-injection for vehicle,
commensal-injected (Comm) and commensal treated with DFMA-DFMO (DFM) (N=6) fish/group).
The vehicle group includes N=12 because it combines N=6 vehicle controls for each time point
with no differences among them. Different letters indicate statistically significant groups at 95%
interval using one-way ANOVA followed by Tukey's post-hoc test (p<0.05).

Figure 4: Systemic inflammation triggers pyrogenic cytokine expression and behavioral fever in a polyamine-dependent manner.

(A) Schematic representation of the rainbow trout in vivo model used in this study. Trout received a single i.v injection of gut commensal bacteria treated or not with DFMA-DFMO to inhibit polyamine synthesis. (B-G) Changes in pro-inflammatory cytokine gene expression, *il-1b*, *tnfa*, and *il-6* in the Tel (B, D, F) and Hyp (C, E, G) 4 hours and 1 day post-injection of gut bacteria treated or not with polyamine synthesis inhibitors (N=7 fish/group). Expression values from the vehicle and commensal injected animals are the same as in Fig. 1. Statistical significance was tested at 95% interval using one-way ANOVA followed by Tukey's post-hoc test (p<0.05). (H) Behavioral fever paradigm used in this study with a continuous ring-shaped swim chamber allowing rainbow trout to exhibit thermal preference between 11°C and 19°C. (I) Mean preferred temperature over the 24-hour observation period post-injection in vehicle, commensal injected, commensal + DFMA-DFMO (DFM) or LPS injected trout. (J) Mean preferred temperature of

1064   rainbow trout 4 hpi in each experimental group. (K) Mean zone transitions 4 hpi in each
1065   experimental group. (L) Mean swimming velocity 4 hpi in each experimental group. Different
1066   letters indicate statistically different groups by Brown-Forsythe and Welch's one-way ANOVA test
1067   ($p<0.05$) (N=5).
1068

1069

fig. S1. Blood-brain barrier integrity in various brain regions of rainbow trout during
systemic inflammation

(A) 16S rDNA microbial community composition at the phylum (A) and family (B) levels of the
isolated gut commensals prior to injection. (C) Schematic of the Evans blue extravasation assay.
(D) Relative Evans Blue levels in the OB, OT and Cer 4 hours and 1 day post commensal
injection (N= 6) relative to vehicle controls. (E) Images of whole dissected rainbow trout brains
injected with vehicle or gut commensals at 4 hours and 1 day post-injection. Data were analyzed
by one-way ANOVA followed by Tukey's post hoc test ($p<0.05$).

1078

1079

1080

1081

1082

fig. S2. Impact of DFMA-DFMO on commensal bacteria growth and in vivo translocation of bacteria

(A) Optical density (O.D=600) of trout commensal bacterial cultures for 24 hours following incubation with different concentrations of DFMA (N=4). (B) Optical density of trout gut commensal cultures at the 24 hours endpoint following incubation with different concentrations of DFMO (N=4). (C) Growth curves of trout commensal bacteria exposed to combined DFMA-DFMO treatment for a 24 hour period (N=4). (D) Effect of combined DFMA-DFMO treatment on bacterial optical density at the 24 hours endpoint. Different letters indicate statistically significant groups by one-way ANOVA followed by Tukey's post hoc test (N=4). (E-J) Quantitative PCR analysis of 16S rRNA gene copy numbers in the spleen, olfactory bulbs, telencephalon, hypothalamus, optic tectum, and cerebellum after i.v injection of commensal bacteria treated or not with DFMA-DFMO, or LPS at 4 hours and 1-day post-injection (N=6), The vehicle group has N=12 because it combines N=6 vehicle controls for each time point that show no significance differences between them. Statistical significance is denoted by distinct alphabetical markers as determined by ANOVA Welch's test ($p < 0.05$).

1098

1099

1100 **fig. S3:** (A) Schematic illustration of the DSS-induced colitis model used in this study. (B)
1101 Comparative analysis of blood serum endotoxin levels between DSS treated animals and vehicle
1102 group. (C-G) Quantitative polymerase chain reaction (qPCR) assessment of 16S rRNA gene
1103 abundance in various brain regions and the spleen of both vehicle and DSS-induced colitis trout
1104 (N=6). Statistical significance is denoted as determined by unpaired two-tailed t-test (N=6).

1105

1106

fig. S4. Microbial community analysis of tank water and rainbow trout telencephalon
microbiome alpha diversity metrics post microbiota injection

Microbial community composition at the phylum (A) and family (B) levels of the tank water from
the aquarium system where trout were maintained. (C-E) Alpha diversity metrics for the rainbow
trout Tel microbial community at 4 hours and 1 day post gut commensal injection. Shannon
diversity index (C), the Chao1 diversity index (D), and Pielou's evenness index (E) (N=6).
Different letters denote statistically significant groups (P<0.05) by Kruskal-Wallis test.

1114

1115

1116

1117

1118

1119

fig. S5: Changes in expression levels of the polyamine synthesis enzymes *odc2* and *samdc2* in
the telencephalon and hypothalamus post-injection of gut commensals treated or not with DFMA
and DFMO or LPS extracted from gut commensals as measured by RT-qPCR (N=6-7).
Expression levels were normalized to *ef1a*. Data were analyzed used the Pfaffl method. Different
letters indicate statistically significant groups based on one-way ANOVA followed by Tukey's post
hoc test (p<0.05).

1126

1127

1128

fig. S6. Behavioral responses in rainbow trout following microbiota and LPS Injections.

(A) Mean zone transition frequency over a 24-hour period post-injection, reflecting the locomotor
activity of rainbow trout in response to various treatments. The graph illustrates the trout's
exploratory behavior, with notable differences among groups treated with vehicle, gut
commensals (Comm), DFMA-DFMO treated commensals (DFM), and LPS. (B) Mean swimming
velocity over 24 hours post-treatment in the four experimental groups (N=5 fish/treatment).

1135

1136

1137   **Table S1.** Primer sequences employed in this study, adapted from published sources

Gene	Sense	Primer 5'-3'	Amplicon size (bp)	Reference
il-1b	Forward	ACATTGCCAACCTCATCATCG	91bp	(1)
	Reverse	TTGAGCAGGTCCTTGTCCTTG		
il-6	Forward	ACTCCCCTCTGTCACACACC	295	(2)
	Reverse	GGCAGACAGGTCCTCCACTA		
tnfa	Forward	GGGGACAAACTGTGGACTGA	66	(2)
	Reverse	GAAGTTCTTGCCCTGCTCTG		
odc1	Forward	CGTGTGCCAGCTCAGTGTC	179	(3)
	Reverse	CCATGTCAAAGACACAGCGG		
odc2	Forward	TGGTGCCACCCTGAAGGCC	128	(3)
	Reverse	AGATGGCCTGGCTGTAGGTG		
samdc1	Forward	GCAAGGACAAGCTAATTAAG	185	(3)
	Reverse	AACCTTGGGATGGTACGGAG		
samdc2	Forward	AACTCACGATGGAAGCGAAC	121	(3)
	Reverse	AACCTTGGGATGGTACGGAG		

1138

1139   **SI References**

1140

1.   J. Zou, P. S. Grabowski, C. Cunningham, C. J. Secombes, Molecular cloning of interleukin
1β from rainbow trout Oncorhynchus mykiss reveals no evidence of an ice cut site.
Cytokine **11**, 552–560 (1999).

2.   T. R. Jørgensen, M. K. Raida, P. W. Kania, K. Buchmann, Response of rainbow trout
(Oncorhynchus mykiss) in skin and fin tissue during infection with a variant of
Gyrodactylus salaris (Monogenea: Gyrodactylidae). *Folia Parasitol. (Praha).* **56**, 251
(2009).

3.   T. C. Clark, J. Tinsley, D. J. Macqueen, S. A. M. Martin, Rainbow trout (Oncorhynchus
mykiss) urea cycle and polyamine synthesis gene families show dynamic expression
responses to inflammation. *Fish Shellfish Immunol.* **89**, 290–300 (2019).

1151

1152