**The adoption of a westernized gut microbiome in Indian Immigrants and Indo-Canadians is associated with dietary acculturation**

*Leah D. D'Aloisio[1], Mamatha Ballal[5], Sanjoy Ghosh[1,10], Natasha Haskey[1], Nijiati Abulizi[1], Ramin Karimianghadim[2], Chuyi Liu[3], Pacha Sruthi[4], Lakshmipriya Nagarajan[4], Sudha Vasudevan[4], Vignesh Shetty[5,6], Mrudgandha Purandare[5], Ushasi Bhaumik[7], Debaki Ranjan Howlader[8], Sepideh Pakpour[1,9], Jacqueline Barnett[10], Deanna L. Gibson*[1,10,11].*

[1] Biology, University of British Columbia Okanagan Campus, Kelowna, Canada

[2] Applied Science, University of British Columbia Okanagan Campus, Kelowna, Canada

[3] College of Pharmacy and Nutrition, University of Saskatchewan, Saskatchewan, Canada

[4] Food & Nutrition Research, Madras Diabetes Research Foundation, Chennai, India

[5] Enteric Disease Division, Department of Microbiology, Kasturba Medical College, Manipal Academy of Higher Education, Karnataka State, India

[6] Department of Medicine, University of Cambridge, Cambridge, United Kingdom

[7] Division of Bacteriology, ICMR-National Institute for Research in Bacterial Infections

[8] Department of Veterinary Pathobiology, University of Missouri, Missouri, USA

[9] School of Engineering, University of British Columbia Okanagan Campus, Kelowna, Canada

[10] Biochemistry and Molecular Biology Program, University of British Columbia Okanagan Campus, Kelowna, Canada

[11] Department of Medicine, University of British Columbia Okanagan Campus, Kelowna, Canada

* Corresponding author: deanna.gibson@ubc.ca

## ABSTRACT

Indian immigration to westernized countries has recently surged, increasing their risk of Inflammatory Bowel Disease (IBD) post-migration. While crucial for understanding IBD risk, the gut microbiome remains understudied in Indians. This cross-sectional study examines the gut microbiomes of Indians residing in India, Indo-Immigrants, and Indo-Canadians in comparison to Euro-Canadian and Euro-Immigrant controls to understand the impact of westernization on their gut. Stool samples for 16S rRNA and shotgun sequencing assessed microbial taxa and functional profiles, complemented by dietary and demographic data to evaluate lifestyle patterns. Results revealed distinct microbiotas in Indians and Indo-Immigrants compared to control groups, with

high abundances of *Prevotella* spp. and CAZymes reflecting their high complex carbohydrate diet.

Indo-Canadians exhibited a transitional microbiome towards westernization, mirroring increasing

dietary acculturation. Considering 44% of Canadians are first- and second-generation immigrants

and the global adoption of westernized practices, future research should investigate the health

implications of such microbiome transitions in immigrant populations and newly industrialized

nations.

**BACKGROUND**

The growing immigrant population in North America is rapidly westernizing through the

adoption of practices like reduced physical activity, changes in hygiene/medical practices, and

acculturation of the westernized diet, commonly characterized by caloric dense, highly processed

foods with low fiber intake. Amidst this backdrop are health disparities emerging in these

populations. Furthermore, the global spread of industrialization raises concern for the increasing

prevalence of modern diseases.[1] For instance, inflammatory bowel disease (IBD) is increasing in

prevalence globally, and alarming evidence reveals a specifically increased risk in young Indian

immigrants and Indians born in several westernized countries.[2-4] In 2019, India had an incidence

rate of 2.34, whereas South Asians in Canada showed an incidence rate of 14.6 per 100,000

person-years, which was comparable to the general population in Canada.[5,6] A likely contributor

to this elevated risk in westernized countries like Canada may stem from changes in the gut

microbiome as a response to living in an industrialized environment.

The industrialized microbiome has been previously characterized by the decline of

VANISH (volatile and/or associated negatively with industrialized societies of humans) taxa,

frequently found in populations adhering to traditional lifestyles, and the rise of BloSSUM (bloom

or selected in societies of urbanization/modernization) taxa, which appear in modern,

industrialized individuals.[7,8] Furthermore, health consequences following the assimilation into

westernized culture have been previously documented. For example, non-industrialized Irish

Travellers who adopted modern practices and Hmong and Karen migrants from Thailand who

relocated to the United States (US) both experienced gut microbiome shifts that were associated

with an increased risk for metabolic disorders.[9,10] Currently, there is only one study published on

the Indian immigrant gut microbiome, highlighting the gap in non-western/non-European

microbiome research. Indeed, over 71% of all microbiome research comes from westernized

cohorts, skewing our interpretation of a healthy human gut.[11] Here, we aimed to investigate the

distinction between the Indian and westernized gut microbiomes and the differences in the gut of

Indians who migrate to Canada, alongside the adoption of the westernized diet.

**METHODS**

***Study Design***

A cross-sectional study was conducted including three Indian groups (Indian ancestry was

confirmed in each group): (1) Indians residing in India ("Indians"), (2) Indians who migrated to

Canada ("Indo-Immigrants") and (3) Canadians born in Canada from Indian ancestry ("Indo-

Canadians"). Caucasian individuals born in Canada with European ancestry ("Euro-Canadian")

were used as a control, and immigrants with European ancestry from a westernized country

("Euro-Immigrants") as a westernized immigrant control. Healthy participants aged 17-55 years

were recruited. Exclusion criteria included pregnancy, diagnosis of any chronic inflammatory

condition, antibiotics used or travel to India within three months prior to stool collection.

Participants completed a demographic questionnaire to assess baseline characteristics.

Immigrants also completed a lifestyle survey to assess changes in lifestyle and immigration-

related stress.

***Stool Collection & DNA Extraction***

Stool samples were collected from India (Kolkata and Manipal) and Canada (Kelowna)

and processed on site as previously described.[12] DNA was sent to Gut4Health for 16S rRNA gene

sequencing on the Illumina MiSeq platform (V4-V4, 515F/806R primers) (~75,800 reads per

sample) and to the Center for Health Genomics and Informatics for shotgun sequencing on the

Illumina NovaSeq platform (~9.9M reads per sample). Additional details are in Supplementary

Methods.

***Taxonomic Analysis***

Demultiplexed 16S rRNA gene reads were processed in QIIME 2 (Version 2023.9)[13] with

DADA2[14] for quality control and taxonomic classification with GreenGenes2 (10.28.22).[15]

Demultiplexed raw shotgun reads underwent quality assessment with FastQC/MultiQC and used

KneadData (Version 0.10) for trimming and quality control. Taxonomic profiling was conducted

using MetaPhlAn4 for higher taxonomic resolution at the species level,[16] and each sample was

normalized to relative abundances. Diversity metrics included Shannon and Pielou's Evenness

(Kruskal-Wallis), and Bray Curtis and Weighted UniFrac (PERMANOVA). Differentially abundant

bacteria were determined using the Linear Discriminate Analysis (LDA) Effect Size (LEfSe)

algorithm with a 3.5 threshold and alpha 0.05.[17] Using shotgun data, the normalized relative

abundance MetaPhlAn4 output table was imported into MicrobiomeAnalyst 2.0 to apply a Random

Forest algorithm ranked according to their Mean Decrease Accuracy score, indicating the extent

to which each taxon's presence or absence influences the overall accuracy of classifying samples

into their respective cohort.[18] These taxa were then visualized in a heatmap. Using 16S rRNA

gene data, BugBase was used to predict microbiome phenotypes,[19,20] and associations between

beta diversity, dietary patterns and baseline characteristics were analyzed with distance-based

redundancy analysis (dbDRA). Additional details for taxonomic analysis methods can be found in

Supplementary Methods.

*Functional Profiling*

Shotgun reads were analyzed for functional potential using HUMAnN (Version 3.6).[21] Gene family and pathway abundances were annotated using the UniRef90[22] and MetaCyc[23] databases, respectively, then normalized to reads per kilobase to account for gene length. Both tables were further normalized to relative abundance to account for sampling depth. The normalized gene family abundance output was then regrouped to the CAZy and KEGG databases. To determine differentially abundant genes/pathways across cohorts, tables were collapsed to perform LEfSe analysis on the unstratified data using a one-against-all strategy with a threshold of 3.0 on the logarithmic LDA score and an alpha of 0.05 for Kruskal-Wallis test among classes.[17] Differentially abundant features were then plotted as a heatmap using MicrobiomeAnalyst 2.0.[18]

*Nutritional Analysis*

Participants filled out a 3-day food log prior to their stool collection, then researchers retrospectively inputted this data into the ESHA Food Processor® software (Version 11.11.0) to obtain the calculated nutrient intake for each subject. To mitigate bias from using a North American nutritional software, Indian and Indo-Immigrant food logs were also entered into the MDRF EpiNu® Nutritional software, tailored for Indian populations. Additional details on our approach for dietary analysis can be found in Supplementary Methods.

*Statistical Analysis*

Statistical analyses for demographics and dietary data were performed using both R statistical software and GraphPad Prism (Version 10.0.3). We applied the Kruskal-Wallis test coupled with Dunn's post hoc analysis for nonparametric data. For data that passed the Shapiro-Wilk test of normality, an ordinary one-way ANOVA followed by Tukey's post hoc test was performed. Categorical data were analyzed using either a Chi-Square, Fisher's Exact or Fisher's

Exact with Monte Carlo simulation, depending on the data structure. Results were reported as

median values and interquartile range (IQR), unless otherwise specified.

**RESULTS**

***Participant Characteristics***

The study included Indians ($n$ = 61), Indo-Immigrants ($n$ = 32), Indo-Canadians ($n$= 17),

Euro-Canadians ($n$ = 41) and Euro-Immigrants ($n$ = 23) (Figure S1). Baseline demographics were

comparable across groups, though Indians were older and consumed significantly less alcohol

($p_{BONF}$ = 3.29e-09) (Table 1). Indo-Immigrants exhibited more of a change towards westernization

(p = 0.002) and higher immigration-related stress (p = 0.04), with no significant differences in age

at or years since immigration (Table 2).

***Indian Microbiota is Distinctive from Westernized Groups***

To examine the taxa and complexity of the microbial ecosystem in Indians compared to

westerners, alpha diversity revealed the lowest diversity in Indians and highest in Euro-Canadians

(Figure 1A-B). Average percentage of unclassified reads per sample in each cohort can be found

in Table S1. Bray Curtis dissimilarity of shotgun sequence data showed significant differences

across groups, with 26.3% of variation explained on the first 2 axes (Figure 1C). Indians and Indo-

Immigrants' microbiotas differed from westernized cohorts (Table 3), and each other (pseudo-F =

9.132, $p_{BONF}$ = 0.01). Weighted UniFrac also demonstrated distinctions in the Indian and Indo-

Immigrant microbiomes (pseudo-F = 25.98, $p_{BONF}$ = 0.01), with 61.8% of the variation explained

on the first 2 axes (Figure 1D) with similar findings from 16S rRNA gene amplicon sequence data

(Figure S2).

### Table 1. Baseline Characteristics of Participants

| Baseline characteristic | Indian n = 61 | Indo-Immigr n = 32 | Indo-Can n = 17 | Euro-Can n = 41 | Euro-Immigr n = 23 | P value |
|---|---|---|---|---|---|---|
| **Sex** | | | | | | |
| Female | 37 (61) | 18 (60) | 6 (35) | 16 (39) | 10 (43) | 0.10 [a] |
| Male | 24 (39) | 12 (40) | 11 (65) | 25 (61) | 13 (57) | |
| **Age, (years)** | | | | | | |
| Median (IQR) | 31 (12) | 25 (8) | 23 (7) | 24 (5) | 29 (15) | |
| Min-max | 17-52 | 18-53 | 18-38 | 19-50 | 19-53 | |
| **Age classification, n (%)** | | | | | | |
| 17-29 years | 26 (43) | 27 (84) | 14 (82) | 32 (78) | 12 (52) | 0.002 *[e] |
| 30-39 years | 21 (34) | 4 (13) | 3 (18) | 4 (10) | 7 (30) | |
| 40-49 years | 10 (16) | 0 (0) | 0 (0) | 4 (10) | 2 (9) | |
| 50-55 years | 4 (7) | 1 (3) | 0 (0) | 1 (2) | 2 (9) | |
| **Mode of birth, n (%)** | | | | | | |
| Vaginal | 44 (75) | 19 (63) | 14 (88) | 34 (83) | 19 (83) | 0.23 [a] |
| Caesarian | 15 (25) | 11 (37) | 2 (12) | 7 (17) | 4 (17) | |
| **BMI** | | | | | | |
| Mean (SD) | 24 (3) | 24 (4) | 22 (3) | 23 (3) | 22 (2) | |
| Min-max | 17-32 | 17-34 | 17-30 | 17-31 | 18-25 | |
| **BMI classification, n (%)** | | | | | | |
| Underweight (< 18.5) | 3 (5.0) | 1 (4.2) | 2 (12.5) | 3 (8.0) | 1 (5.0) | 0.08 [b] |
| Normal weight (18.5 - < 25) | 40 (66.7) | 14 (58.3) | 12 (75) | 24 (63.0) | 21 (95.0) | |
| Overweight (25 - < 30) | 13 (21.7) | 8 (33.3) | 2 (12.5) | 10 (26.0) | 0 (0) | |
| Obese (> 30) | 4 (6.7) | 1 (4.2) | 0 (0) | 1 (3.0) | 0 (0) | |
| **Smokers, n (%)** | | | | | | |
| Yes | 6 (10) | 0 (0) | 0 (0) | 2 (5) | 2 (9) | 0.2908 [c] |
| No | 56 (90) | 32 (100) | 17 (100) | 39 (95) | 21 (91) | |
| **Alcohol, n (%)** | | | | | | |
| Yes | 15 (24) | 15 (47) | 14 (82) | 33 (80) | 18 (78) | <0.0001†[a] |
| No | 47 (76) | 17 (53) | 3 (18) | 8 (20) | 5 (22) | |
| **Physical activity- walking (mins)** | | | | | | |
| Median (IQR) | NA | 90 (128) | 90 (120) | 60 (75) | 60 (90) | 0.56 [d] |
| Min-max | NA | 0-330 | 30-300 | 12-480 | 12-420 | |
| **Physical activity- exercise (days)** | | | | | | |
| Median (IQR) | NA | 3 (2) | 3 (3) | 4 (3) | 4.5 (3) | 0.18 [e] |
| Min-max | NA | 0-7 | 0-7 | 0.5-6.5 | 1-6 | |
| **Socioeconomic class, n (%)** | | | | | | |
| Low | 15 (25) | 8 (25) | 5 (33) | 10 (25) | 4 (17) | 0.70 [e] |
| Middle | 40 (67) | 21 (66) | 7 (47) | 26 (65) | 14 (61) | |
| Upper | 5 (8) | 3 (9) | 3 (20) | 4 (10) | 5 (22) | |
| **Lifestyle born/raised, n (%)** | | | | | | |
| Rural | NA | 2 (6) | 5 (31) | 5 (12) | 5 (22) | 0.10 [c] |
| Urban | NA | 30 (94) | 11 (69) | 36 (88) | 18 (78) | |

Abbreviations: SD, standard deviation; IQR, interquartile range; BMI, body mass index; [a]Chi-Square, [b]ANOVA, [c]Fisher's Exact, [d]Kruskal-Wallis, [e]Fisher's Exact (Monte Carlo); * Bonferroni correction revealed significant differences between Indian vs. Indo-Immigr ($p_{BONF}$ = 0.006), Indian vs. Euro-Can ($p_{BONF}$ = 0.03); † Bonferroni correct revealed significant differences ($p_{BONF}$ = 3.29e-09).

**Table 2. Characteristics of Immigrant Participants**

| Characteristic | Indo-Immigr n = 32 | Euro-Immigr n = 23 | P value |
|---|---|---|---|
| **△ westernization, n (%)** | | | |
| No westernization | 0 (0) | 4 (18) | 0.002 [a] |
| Little westernization | 12 (38) | 13 (59) | |
| Moderate westernization | 17 (53) | 4 (18) | |
| High westernization | 3 (9) | 1 (4) | |
| **Immigration-related stress, n (%)** | | | |
| Low stress | 22 (69) | 21 (91) | 0.04 [a] |
| Moderate stress | 8 (25) | 2 (9) | |
| High stress | 2 (6) | 0 (0) | |
| **Age at immigration, n (%)** | | | |
| 0-3 years | 0 (0) | 1 (4) | 0.19 [b] |
| 4-17 years | 9 (28) | 3 (13) | |
| 18-49 years | 23 (72) | 19 (83) | |
| **Years since immigration** | | | |
| Median (IQR) | 4 (6) | 3 (9) | 0.65 [c] |
| Min-max | 0-22 | 0-48 | |

Abbreviations: IQR, interquartile range; [a]Chi-Square for Trend, [b]Fisher's Exact, [c]Mann-Whitney

LEfSe identified dominant taxa in each group, with Bacteroidota (formerly Bacteroidetes)

and Pseudomonadota (formerly Proteobacteria) in Indians and Firmicutes in Euro-Canadians

(Figure 1E, Table S1). Indo-Immigrants displayed highest abundances of *Eubacterium* and

Erysipelotrichaceae (a VANISH taxon)[24], both previously found to be enriched in non-

industrialized populations[25], though abundance plots of both taxa reveal low abundances in

Indians (Figure S4). As expected, genera common in the industrialized microbiome were enriched

in the westernized European cohorts such as *Alistipes*, *Bacteroides*, and Clostridia, all previously

noted at high abundances in industrialized groups.[25,26] Nonetheless, average relative abundances

for *Alistipes* and *Bacteroides* were higher in Indians than Indo-Immigrants. Indo-Canadians

represented a transitional gut microbiota, harbouring both taxa found in non-industrialized, such

as *Collinsella*, and industrialized populations, such as *Blautia* and *Anaerostipes* (both noted as

BloSSUM taxa).[25,26] Similar results were found in 16S rRNA gene sequence data (Figure S3,

Table S2).

207  **Figure 1. Indian and Indo-Immigrant Microbiotas are Distinct from Westernized Groups**

208  **(A)** Pielou's Evenness (H = 85.7, p = 1.07e-17). **(B)** Shannon's Diversity (H = 79.8, p = 1.89e-16). Beta diversity was explored with Bray Curtis dissimilarity and Weighted UniFrac, using Pairwise Permutational Multivariate Analysis of Variance (PERMANOVA) to test differences between groups. **(C)** Bray Curtis principal coordinate analysis (PCoA) plot shows 26.3% of variation was captured on the

209  first two axes. **(D)** Weighted UniFrac PCoA plot shows 61.8% of the variation was captured on the first two axes. Pairwise comparisons results can be found in Table S1. **(E)** LEfSe (Linear discriminate analysis Effect Size) cladogram results, depicting differentially abundant bacteria across cohorts, with genus set as lowest taxonomic rank. *** = q ≤ 0.001, ** = q ≤ 0.01, * = q ≤ 0.05. Boxes represent the

210  interquartile range (IQR) between the first and third quartiles, the horizontal line indicates the median, and whiskers are the upper and lower values within 1.5 times the IQR.

211 **Table 3. PERMANOVA Results from Beta Diversity Analysis**

| | Bray Curtis | | Weighted UniFrac | |
|---|---|---|---|---|
| Pair | pseudo-F | *P* value | pseudo-F | *P* value |
| Indo-Immigr vs. Indo-Can | 4.215003 | 0.01 | 4.670629 | 0.01 |
| Indo-Immigr vs. Euro-Can | 8.976951 | 0.01 | 11.932016 | 0.01 |
| Indo-Immigr vs. Euro-Immigr | 5.313757 | 0.01 | 6.903393 | 0.01 |
| Indo-Immigr vs. Indian | 9.132465 | 0.01 | 25.984365 | 0.01 |
| Indo-Can vs. Euro-Can | 3.084292 | 0.01 | 5.409267 | 0.01 |
| Indo-Can vs. Euro-Immigr | 2.166793 | 0.01 | 2.360739 | 0.10 |
| Indo-Can vs. Indian | 8.660676 | 0.01 | 25.206253 | 0.01 |
| Euro-Can vs. Euro-Immigr | 1.163757 | 1.00 | 1.116192 | 1.00 |
| Euro-Can vs. Indian | 18.284343 | 0.01 | 49.611215 | 0.01 |
| Euro-Immigr vs. Indian | 11.182855 | 0.01 | 31.565399 | 0.01 |

213 Adjusted *P* values are displayed, which were obtained using 999 permutations and Bonferroni correction applied. Beta diversity data from shotgun sequence data.

**Figure 2. Significantly Different Gut Bacterial Abundances Detected Across Cohorts**

**(A)** Taxonomic stacked bar plots depicting relative abundances of top 18 most abundant bacteria across all samples from shotgun sequence data (N = 174; Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). **(B)** Heatmap generated in MicrobiomeAnalyst 2.0, displaying taxa identified by Random Forest as key features that contribute to the predictive accuracy of classifying samples into their respective groups (Figure S5). For the heatmap, features were filtered for minimum 4 counts in 20% or more samples, and low variance filter of 10%. Bars on the top represent the cohorts that group the individual sample columns displayed. Taxa are labelled in rows, with taxonomic rank noted before the bacteria name. Colours on heatmap represent the relative abundances of each bacteria in a given sample. **(C)** Heatmap generated in R Studio, displaying the average relative abundances of *P. copri* clades in each cohort. **(D)** LEfSe results of top differentially abundant bacteria. A Kruskal-Wallis test was performed with a significance ($\alpha$) at 0.05 for one-against-all comparisons. Differentially abundant bacteria were detected using a Linear Discriminate Analysis (LDA) score (equal to or greater than 3.5). **(E)** Abundance plot of notable differentially abundant species identified by LEfSe, displaying average relative abundance per cohort.

When assessing differences in bacterial species abundances amongst groups, LEfSe

showed that Indians harboured higher abundances of *Phocaeicola plebeius*, *Megasphaera sp.*,

*Dialister hominis,* and *Escherichia coli* (Figure 2). Among Indo-Immigrants, *Dorea longicatena*,

*Faecalibacterium prausnitizii* and *Lachnospiraceae bacterium* were notably abundant, while Indo-

Canadians had increased abundances of two *Ruminoccocus torques* SGBs, *Blautia wexlerae,*

*Eubacterium rectale, Blautia massiliensis,* and *Anaerostipes hadrus*. Taxa enriched in Euro-

Canadians included *Phocaeicola vulgatus*, *Lachnospiraceae bacterium, Clostridium sp.,* and

*Blautia faecis*, and Euro-Immigrants showed enrichment of *Fusicatenibacter saccharivorans,*

*Clostridia bacterium,* and *Blautia obeum.* A key distinguishing feature from Indians to Indo-

Canadians was the decline in *Prevotella* spp., with *Prevotella copri* (clade A) being over five times

more abundant in Indians (Table 4). *P. copri* Clade A relative abundances were 10.8 for Indians,

5.89 for Indo-Immigrants, and 2.75 for Indo-Canadians (Figure 2C). A similar pattern of

decreasing abundance across Indian cohorts was also observed for Clades B, C and D. All clades

were lowest in abundance in Euro-Canadians. Overall, these findings demonstrate that the gut

microbiome in Indians are distinct from the West, and Indo-Canadians exhibit a transition towards

the industrialized microbiome.

**Indian Gut Microbiome Predicted to be More Pathogen-Tolerant**

To understand potential functionality of the gut microbiota in Indians, we used BugBase

predictions (Figure 3), which estimated that Indians appeared to demonstrate higher stress

tolerance driven by Pseudomonadota (Proteobacteria). The Indian cohorts had the highest

potentially pathogenic bacteria ($p_{FDR}$ = 2.07e-18) mostly dominated by Gram negative bacteria

driven mainly by a combination of Pseudomonadota, Bacteroidota (Bacteroidetes), and

Firmicutes. In addition, BugBase predicted that the Indian gut microbiome had the highest

potential for biofilm formation. Overall, these results

**Table 4. Differentially abundant bacteria from shotgun sequence data**

| Taxon | Log Score | Class | LDA | P value |
|---|---|---|---|---|
| Veillonellaceae GGB4266 (SGB5809) | 4.023 | Indian | 3.680 | 0.003000 |
| *Phocaeicola plebeius (*SGB1903) | 4.658 | Indian | 4.361 | 2.40E-12 |
| *Prevotella copri* clade A (SGB1626) | 5.034 | Indian | 4.735 | 9.25E-13 |
| *Escherichia coli* (SGB10068) | 4.157 | Indian | 3.801 | 4.21E-12 |
| *Megasphaera sp.* (BL7 SGB5858) | 4.456 | Indian | 4.159 | 2.04E-10 |
| *Dialister hominis* (SGB5803) | 4.147 | Indian | 3.838 | 5.42E-06 |
| *Acidaminococcus intestini* (SGB5736) | 3.758 | Indian | 3.544 | 0.001382 |
| *Dorea longicatena* | 4.390 | Indo-Immigr | 4.019 | 2.99E-16 |
| *Faecalibacterium prausnitzii* (SGB15342) | 4.242 | Indo-Immigr | 3.852 | 3.18E-11 |
| *Lachnospiraceae bacterium* (WCA3 601 WT 6H SGB4910) | 4.079 | Indo-Immigr | 3.685 | 0.009908 |
| *Dorea longicatena* (SGB4581) | 4.301 | Indo-Immigr | 3.922 | 2.04E-07 |
| *Faecalibacterium prausnitzii* | 4.930 | Indo-Can | 4.513 | 1.07E-16 |
| *Ruminococcus torques* | 4.410 | Indo-Can | 4.083 | 4.02E-16 |
| *Anaerobutyricum hallii* | 4.033 | Indo-Can | 3.693 | 3.48E-21 |
| *Collinsella aerofaciens* (SGB14546) | 4.478 | Indo-Can | 4.099 | 1.69E-11 |
| *Dialister invisus* (SGB5825) | 4.270 | Indo-Can | 3.963 | 4.37E-08 |
| *Blautia wexlerae* (SGB4837) | 4.657 | Indo-Can | 4.292 | 5.37E-21 |
| *Gemmiger formicilis* (SGB15300) | 4.422 | Indo-Can | 4.001 | 3.60E-06 |
| *Faecalibacterium prausnitzii* (SGB15318) | 4.436 | Indo-Can | 4.108 | 3.78E-14 |
| *Blautia massiliensis* (SGB4826) | 4.232 | Indo-Can | 3.925 | 1.82E-12 |
| *Anaerostipes hadrus* (SGB4540) | 4.189 | Indo-Can | 3.849 | 3.45E-19 |
| *Ruminococcus torques* (SGB4563) | 4.172 | Indo-Can | 3.843 | 1.32E-10 |
| *Faecalibacterium prausnitzii* (SGB15332) | 4.368 | Indo-Can | 4.007 | 2.44E-09 |
| *Anaerobutyricum hallii* (SGB4532) | 3.918 | Indo-Can | 3.583 | 6.69E-20 |
| *Eubacterium rectale* (SGB4933) | 4.656 | Indo-Can | 4.179 | 5.36E-05 |
| *Clostridia bacterium* (SGB14861) | 4.027 | Indo-Can | 3.739 | 7.03E-16 |
| Ruminococcus torques (SGB4608) | 4.035 | Indo-Can | 3.710 | 2.92E-13 |
| *Lachnospiraceae bacterium* | 4.275 | Euro-Can | 3.861 | 3.12E-16 |
| *Blautia faecis* (SGB4820) | 3.909 | Euro-Can | 3.564 | 3.08E-19 |
| *Phocaeicola vulgatus* (SGB1814) | 4.488 | Euro-Can | 4.056 | 5.66E-05 |
| *Clostridium sp* (AF34 10BH SGB4914) | 4.083 | Euro-Can | 3.654 | 0.007673 |
| Ruminococcaceae GGB9635 (SGB15106) | 3.871 | Euro-Immigr | 3.588 | 0.014807 |
| *Clostridia bacterium* | 4.522 | Euro-Immigr | 4.047 | 3.81E-08 |
| *Blautia obeum* | 3.917 | Euro-Immigr | 3.566 | 5.24E-15 |
| *Fusicatenibacter saccharivorans* (SGB4874) | 4.485 | Euro-Immigr | 4.154 | 3.84E-18 |
| *Lacrimispora celerecrescens* (SGB4868) | 3.960 | Euro-Immigr | 3.635 | 2.47E-08 |

**Figure 3. BugBase Predicts Higher Pathogenic Potential and Stress-Tolerant Microbiome in Indians**

A Kruskal-Wallis test was performed in BugBase followed by Mann-Whitney-Wilcoxon tests for pairwise comparisons with adjusted *P* values shown. Relative abundance is presented on the y-axis. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). **(A)** Stress tolerance **(B)** Potentially pathogenic bacteria **(C)** Gram negative bacteria **(D)** Containing mobile elements **(E)** Aerobic bacteria **(F)** Anaerobic bacteria **(G)** Facultatively Anaerobic bacteria **(H)** Biofilm formation. Taxonomic contributions are also displayed for each phenotype prediction. *** = p ≤ 0.001, ** = p ≤ 0.01, * = p ≤ 0.05

suggest the Indian gut microbiome may be more pathogen-tolerant, as healthy individuals were

predicted to harbour microbes with higher pathogenic potential and stress tolerance. However, it

is important to note that results from BugBase are only predictions made using an older

GreenGenes database, therefore these findings should be further substantiated in future studies.

**Gut Functional Potential in Indians is Distinctive from Westernized Groups**

To further understand functional attributes of the Indian gut microbiome, pathway analysis

using MetaCyc was used and revealed pathways enriched in lipopolysaccharide (LPS)

production, peptidoglycan synthesis, microbial growth/metabolism, protein synthesis and DNA

synthesis/repair (Figure 4). In Indo-Canadians, indications of increased glucose storage were

detected. Euro-Canadians had increased pathways related to fatty acid biosynthesis, and Euro-

Immigrants expressed pathways indicating sugar metabolism (Table S3). CAZy gene family

analysis revealed distinct profiles in the Indian gut including glycoside hydrolases (GH)43, GH51,

and GH10, all of which are known for their degradation of plant-based polysaccharides (Table

S4). *P. copri* was the top taxa contributing to these CAZy families (Figure 5A). In addition, KEGG

identities K00561 and K18220 were enriched in Indians and Indo-Immigrants, respectively, both

recognized for antimicrobial resistance genes (Figure 5B). Overall, these results suggest that both

environmental factors, diets and drugs, were influencing the gut microbiome of Indians.

**Westernized Dietary Acculturation Observed in Indo-Canadians**

To understand the role of diet associated with the gut microbiome in our Indian cohort, we

collected and analyzed dietary recalls using both validated software from North America and from

India to account for Indian foods. Indo-Canadians had the highest daily intake of NOVA Group 4

Ultra-Processed Food (UPF), constituting 61% of their caloric intake (95% CI = 467.9 – 780.5

kcal, per 1000 kcal), while Indians consumed significantly less than all other cohorts, reporting

12% of their daily intake (95% CI = 63.66 – 127.1 kcal per 1000 kcal) (Figure 6). Consistent with

high UPF intake, total fiber was also found to be the lowest in Indo-Canadians, opposite in Indians

who consumed more fiber than Indo-Canadians ($p = 0.0028$) and Euro-Canadians ($p = 0.0110$).

The highest proportion of non-meat eaters were Indians at 60% ($n = 36/61$, $p = <0.0001$), with

30% vegetarian and 30% pescetarian. In Indo-Immigrants, 19% ($n = 6/32$), while 12% ($n = 2/17$)

of Indo-Canadians, 10% (*n* = 4/41) Euro-Canadians and 39% (*n* = 9/23) of Euro-Immigrants were

non-meat eaters. Cooking oil usage highlighted sunflower oil, ghee and mustard oil as the top

choices of Indians, whereas Indo-Immigrants preferred olive oil, butter and ghee, with olive oil

and butter being the primary fats used by the three westernized groups (Figure S6, Table 5).

These results highlight the westernized diet adopted by Indo-Canadians, including high levels of

processed food and low levels of fiber, are potential drivers in the westernized gut microbiota.

**Table 5. Percentages of respondents who reported use of each cooking oil**

| Oil Type | Indian *n = 43* | Indo-Immigr *n = 18* | Indo-Can *n = 9* | Euro-Can *n = 35* | Euro-Immigr *n = 18* |
|---|---|---|---|---|---|
| Butter | 5% | 24% | 27% | 24% | 26% |
| Ghee | 12% | 16% | 0% | 0% | 0% |
| Coconut | 9% | 4% | 0% | 5% | 10% |
| Olive | 1% | 28% | 45% | 44% | 39% |
| Sunflower | 30% | 0% | 0% | 2% | 0% |
| Canola | 0% | 12% | 0% | 13% | 6% |
| Vegetable | 5% | 4% | 18% | 2% | 10% |
| Mustard | 11% | 8% | 0% | 0% | 0% |
| Avocado | 0% | 0% | 0% | 5% | 6% |
| Groundnut | 8% | 0% | 0% | 0% | 0% |
| Rice | 8% | 0% | 0% | 0% | 0% |
| Other | 9% | 4% | 9% | 4% | 3% |

**Figure 4. Microbial Metabolic Pathways Show Distinction in Functional Potential of Indian Gut Microbiome**

Pathway abundances were annotated using MetaCyc, then normalized to account for gene length and sampling depth. LEfSe (Linear discriminate analysis Effect Size) was used to test for differentially abundant microbial metabolic pathways using unstratified data. A Kruskal-Wallis test was performed with a significance (a) at 0.05 for one-against-all comparisons. Discriminate features were identified using a Linear Discriminate Analysis (LDA) score, with an LDA score equal to or greater than 3.0. Using MicrobiomeAnalyst 2.0, a heatmap was generated to display the differentially abundant pathways identified from LEfSe. Each column is a sample, and the coloured bars on top represent the cohorts. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). Metabolic pathways are labelled in rows and annotated with a colour that describes their general function. Colours on the heatmap represent the relative abundances of each pathway. * *L-orthinine biosynthesis I* is indirectly linked with protein synthesis and also involved with urea cycle; * *CMP-3-deoxy-D-manno-octulosonate biosynthesis* is directly involved in lipopolysaccharide (LPS) production.

**Figure 5. *Prevotella copri* is the Top Taxa Contributing to CAZy Families Abundant in Indians**

Normalized gene family abundance table was regrouped to CAZy database using *human_regroup_table*. Barplots for stratified taxa contributions for each CAZy gene family was generated using *human_barplot*. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). *Prevotella copri* is the top contributor to CAZyme families **(A)** GH10, GH43, GH51, all which were enriched in Indians. **(B)** Top taxa contributions to antimicrobial resistance-related KEGG orthologues abundant in Indians and Indo-Immigrants. Normalized gene family abundance table was regrouped to KEGG database using *human_regroup_table*. Barplots for stratified taxa contributions for each KEGG ortholog was generated using *human_barplot*. *Prevotella copri AM42 24* is the top contributor to K00561, enriched in Indians and *Coprococcus comes* is the top contributing taxa to K18220, enriched in Indo-Immigrants.

To understand what dietary factors could be driving microbial taxa, the dbRDA model,

accounting for demographics and dietary patterns elucidated 16.62% of the microbial variance on

the first 2 axes (PERMANOVA F = 5.9923, p = 0.001) (Figure 6C). After false discovery rate (FDR)

adjustment, UPF and alcohol intake were principal determinants in differentiating the microbiota

in westernized cohorts ($p_{FDR}$ = 0.003). The pescetarian diet ($p_{FDR}$ = 0.070) and fiber intake ($p_{FDR}$ =

0.2268) appeared to influence clustering of Indian samples, albeit not significant. Advancing age

($p_{FDR}$ = 0.044) influenced another clustering, whereas vegetarianism showed no significant effect

($p_{FDR}$ = 0.371). Overall, these results suggest that UPF and alcohol intake were driving

westernization of the gut microbiome in Indian migrants with this effect most prominent in Indo-

Canadians. Furthermore, the higher fiber intake observed in Indians correlated with high

*Prevotella* spp. and CAZy gene families tailored to the more traditional Indian diet.

Given the above data indicated that the Indo-Canadian group had adopted a westernized

diet, further analysis of diet was examined including absolute macronutrient and micronutrient

intakes (Table 6-7; Table S5-S6). Total energy intake was significantly higher in Euro-Canadian

males vs. Indo-Immigrant males (p = 0.0144, 95% CI = -1461 – -110.0 kcal). Median percentage

of protein intake was highest in Euro-Canadians (16.1%, 95% CI = 13.2 – 17.5%) and lowest

protein intake was in Indians (12.5%, 95% CI = 11.7 – 13.3). Indians consumed the highest

percentage of carbohydrates in their diet (56.5%, 95% CI = 54.7% - 59.0%) and lowest

carbohydrate diet intake was in Indo-Canadians (44.0%, 95% CI = 40.6% – 50.6%). The highest

percentage of fat consumption was in Indo-Canadians (39.7%, CI 95% 33.8% – 42.2%) and the

lowest was in Indians (31.1%, 95% CI = 28.5% – 34.4%). Software comparisons between North

American ESHA and Indian EpiNu revealed no macronutrient composition differences however,

EpiNu calculated a significantly higher intake of PUFAs and MUFAs and lower intake of SFA

amongst Indians, highlighting discrepancy in dietary fat types between dietary software, which

could skew dietary analysis between the countries (Fig 7A-B). When the type of oils consumed

were examined, it was revealed that Indo-Canadians have the least diversity with no traditional

fats added to their diet like ghee (Fig 7C).

410  **Figure 6. Significant Intake of Ultra-Processed Food and Alcohol are Drivers in the Westernized Gut Microbiota**

411  Middle bands are the median values, top and bottom boxes display the first and third quartiles, and whiskers are the min and max values. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). **(A)** Calories from Group 4 Ultra-Processed Food (UPF) was counted for each subject from the ESHA dietary reports then compared to total average daily caloric intake. A Kruskal-Wallis test was performed, followed by Dunn's multiple comparisons, with adjusted p values displayed. **(B)** Fiber intake was normalized per 1000 kcal for each subject. An ordinary one-way ANOVA was performed, followed by Tukey's multiple comparisons, with adjusted p values displayed. **(C)** Distance-based redundancy analysis (dbRDA) plot was generated using Weighted UniFrac distance matrix from 16S rRNA gene amplicon data. This dbRDA plot depicts the distribution of samples and their associations with lifestyle factors. Each dot is a sample, colour coded by cohort. Variance inflation factor values ranged between 1.14 (vegetarian) − 1.38 (pescetarian), indicating each variable uniquely contributed to the model.

**Table 6. Male participant absolute macronutrient intake**

| Macronutrient | RDA/AMDR | Indian n = 37 | Indo-Immigr n = 18 | Indo-Can n = 6 | Euro-Can n = 16 | Euro-Immigr n = 10 | P value |
|---|---|---|---|---|---|---|---|
| **Energy, kcal (IQR)** | - | 2329 (1187) | 2086 (933) | 2274 (532) | 2911 (1342) | 2668 (906) | 0.0343[a] |
| **Protein** | | | | | | | |
| g (IQR) | 56g | 82.9 (41.3) | 78.8 (39.1) | 138 (37.2) | 119 (62.6) | 103 (42.5) | <0.0001[a] |
| % of total energy | 10-35% | 13.1% | 14.3% | 23.1% | 17.8% | 16.4% | |
| g per 1000 kcal | - | 31.3 | 35.1 | 37.5 | 40.4 | 37.6 | |
| **Fat** | | | | | | | |
| g (IQR) | - | 80.2 (43.1) | 76.5 (40.8) | 109 (39.9) | 113 (63.9) | 98.5 (45.2) | 0.0109[a] |
| % of total energy | 20-35% | 29.2% | 34.6% | 39.2% | 38.2% | 37.1% | |
| g per 1000 kcal | - | 34.6 | 38.2 | 44.1 | 42.9 | 40.7 | |
| **PUFA, g (IQR)** | - | 13.1 (13.7) | 10.6 (6.33) | 13.7 (6.32) | 22.2 (13.3) | 11.7 (9.35) | 0.0254[b] |
| **Omega-3, g (IQR)** | 1.6 | 1.44 (1.77) | 0.93 (0.53) | 1.24 (1.01) | 2.31 (3.44) | 0.85 (1.63) | 0.0080[b] |
| **Omgea-6, g (IQR)** | 17 | 11.5 (10.9) | 8.77 (4.64) | 10.5 (5.93) | 16.7 (10.8) | 9.20 (7.15) | 0.0127[b] |
| **MUFA, g (IQR)** | - | 17.8 (14.3) | 17.7 (11.8) | 17.2 (22.3) | 33.0 (24.0) | 24.6 (13.7) | 0.0100[b] |
| **SFA, g (IQR)** | - | 26.4 (20.0) | 23.0 (12.2) | 34.5 (15.0) | 37.3 (22.3) | 30.1 (19.0) | 0.0055[a] |
| **Carbohydrate** | | | | | | | |
| g (IQR) | 130g | 336 (58.6) | 267 (51.8) | 207 (39.7) | 268 (41.1) | 284 (43.0) | <0.0001[a] |
| % of total energy | 45-65% | 58.6% | 51.8% | 39.8% | 41.1% | 43.0% | |
| g per 1000 kcal | - | 141 | 131 | 110 | 113 | 112 | |
| **Fiber, g (IQR)** | 38g | 34.9 (17.4) | 28.3 (20.1) | 13.1 (9.70) | 23.9 (15.5) | 22.4 (22.6) | 0.0016[b] |
| Soluble fiber | | 0.37 | 0.59 | 0.03 | 0.66 | 0.56 | 0.1223[b] |

Abbreviations: [a]ANOVA, [b] Kruskal-Wallis. Median and IQR reported.

416 **Table 7. Female participant absolute macronutrient intake**

| Macronutrient | RDA/AMDR | Indian n = 24 | Indo-Immigr n = 14 | Indo-Can n = 11 | Euro-Can n = 25 | Euro-Immigr n = 13 | P value |
|---|---|---|---|---|---|---|---|
| **Energy, kcal (IQR)** | - | 1747 (630) | 1976 (1136) | 1812 (527) | 2265 (910) | 2227 (1060) | 0.0332* |
| **Protein** | | | | | | | |
| g (IQR) | 46g | 51.5 (21.8) | 65.3 (47.9) | 63.1 (20.5) | 81.0 (20.8) | 66.6 (50.7) | 0.0004 [b] |
| % of total energy | 10-35% | 12.3% | 13.8% | 12.9% | 14.1% | 14.0% | |
| g per 1000 kcal | - | 31.3 | 35.1 | 37.5 | 40.4 | 37.6 | |
| **Fat** | | | | | | | |
| g (IQR) | - | 62.7 (32.3) | 69.8 (38.0) | 73.1 (27.5) | 89.2 (40.4) | 103 (57.0) | 0.0018 [b] |
| % of total energy | 20-35% | 33.4% | 34.4% | 39.7% | 38.7% | 36.7% | |
| g per 1000 kcal | - | 34.6 | 38.2 | 44.1 | 42.9 | 40.7 | |
| **PUFA, g (IQR)** | - | 12.1 (9.49) | 12.9 (11.0) | 8.06 (4.33) | 13.3 (8.90) | 9.87 (13.0) | ns |
| **Omega-3, g (IQR)** | 1.1 | 1.24 (0.85) | 1.46 (1.02) | 0.84 (0.44) | 1.42 (1.25) | 1.06 (1.16) | ns |
| **Omgea-6, g (IQR)** | 12 | 10.5 (8.72) | 8.80 (8.25) | 7.00 (4.04) | 10.0 (8.92) | 8.73 (7.67) | ns |
| **MUFA, g (IQR)** | - | 16.5 (10.5) | 15.6 (13.4) | 15.3 (9.77) | 23.8 (13.7) | 19.0 (23.6) | ns |
| **SFA, g (IQR)** | - | 18.3 (14.9) | 23.4 (10.6) | 22.9 (5.66) | 30.6 (14.3) | 41.7 (35.1) | ns |
| **Carbohydrate** | | | | | | | |
| g (IQR) | 130g | 258 (86.4) | 256 (116) | 218 (99.5) | 272 (119) | 248 (141) | ns |
| % of total energy | 45-65% | 55.7% | 53.6% | 38.4% | 47.6% | 49.4% | |
| g per 1000 kcal | - | 141 | 131 | 110 | 113 | 112 | |
| **Fiber, g (IQR)** | 25g | 24.6 (17.8) | 23.1 (18.7) | 14.7 (6.30) | 27.0 (15.2) | 28.8 (28.4) | 0.0163 [a] |
| Soluble fiber | | 0.21 | 0.69 | 0.05 | 0.75 | 0.45 | 0.0003 [b] |

Abbreviations: [a]ANOVA, [b] Kruskal-Wallis. Median and IQR reported.
*Tukey's post hoc analysis revealed no significant differences between groups

**Figure 7. Dietary Fat Types are Significantly Different Between EpiNu vs. ESHA**

**(A)** Percentages of fat, protein and carbohydrates were calculated in ESHA and EpiNu for each participant, then mean values were calculated for each cohort and an ordinary one-way Analysis of Variance (ANOVA) was used. **(B)** Significant differences in types of fat consumed by Indians (n = 61) and Indo-Immigrants (n = 32) were calculated between EpiNu vs. ESHA. Dietary fats displayed in grams per 1000 Calories (kcal). Kruskal-Wallis tests were conducted to determine differences between groups. The middle band inside of the boxes is the median, the bottom and top boxes are the first and third quartiles, and the whiskers are the min to max values.

## DISCUSSION

**Westernization is Linked to Both Taxonomic and Functional Transitions in the Gut**

**Microbiomes of Indian Migrants**

Earlier studies identified that the Indian gut microbiota is distinct, being enriched in

*Prevotella* spp*., Dialister* spp*.,* and *Megasphaera* spp.,[27-33] unlike westernized groups dominated

by *Bacteroides* spp.[34] and *Phocaeicola vulgatus*[35]. The taxonomic heterogeneity within Indians

also highlights how generalizing the "Indian microbiome" may inaccurately represent this

population, given the cultural variations that exist within the country. Contrary to associations

between low alpha diversity and disease in western populations, healthy Indians showed lower

alpha diversity than westernized cohorts, suggesting this diversity measure alone may not always

be a reliable marker of westernization or health status.[36] However, the dominance of *Prevotella*

*spp*. in the Indian gut may have masked the detection of rarer species, potentially underestimating

richness estimates in Indians.

Taxonomic data revealed Indians and Indo-Immigrants harbour higher abundances of

bacteria characteristic of pre-industrialized societies, including some previously characterized as

VANISH taxa. In contrast, Caucasian cohorts possess microbes typical of industrialized

populations, including BloSSUM taxa noted in previous studies. However, Indo-Canadians

exhibited a transitional microbiota that embodied a blend of both lifestyles, a trend previously

detected in infants living in transitional societies (intermediate between non-industrialized and

industrialized lifestyles).[37] A key genus that is distinctive in this transition in immigrants is the loss

of *Prevotella* spp.[10,38] In fact, this contrast in *Prevotella* spp. abundance in non-westernized versus

westernized groups has sparked interest both in its relation to lifestyle and disease, previously

being linked to IBD,[39,40] rheumatoid arthritis,[41] and obesity[42]. Yet, contradictory studies also

showed its correlation with reduced risks of cardiovascular disease (CVD) [43] diabetes [44], and

obesity,[45] illustrating the current contradictory understanding around *Prevotella* spp.-dominance

in the gut and its role in diseases.

Globally, Indians have some of the highest abundances of *Prevotella* spp. in their gut,[46]

as reflected in our data, emphasizing the importance of context when investigating the role of

microbes in our gut, as host-microbe interactions may divergently evolve depending on the host's

genetics, geography, and lifestyle. In fact, these factors may have already driven evolutionary

changes in *Prevotella* spp. strains: isolated *P. copri* strains from Indian/non-western populations

showed traits that aid in digestion of plant-based carbohydrates using carbohydrate-active

enzymes (CAZymes), whereas *P. copri* strains from westernized populations were enriched with

virulence factors and resistance genes.[46] In our study, Indians had a significantly higher average

abundance of *P. copri* Clade A, along with notable abundances of Clades B and C, which

subsequently declined in Indo-Immigrants and Indo-Canadians. All *P. copri* clades have

previously been documented at higher abundances in non-westernized individuals than

westernized,[47] and our results highlight this effect of westernization on clade abundances using

groups of the same ethnicity. These findings demonstrate how specific strains can evolve with its

host, cautioning against broad associations between bacterial species and diseases.

Indians, who traditionally consume a high-complex-carbohydrate diet, have been shown

to exhibit increased expression of carbohydrate metabolism genes for complex polysaccharides

in the gut.[27,31,33] Our findings reflected this relationship, as we detected enrichment of CAZy

families that aid in the digestion of xylan and xyloglucan, with *P. copri* as the top contributing taxa.

A similar pattern was in fact detected in another study on immigrants who migrated from Thailand

to the US, where a decrease in *Prevotella* spp. abundance corresponded with a loss of CAZymes

that break down plant fiber.[10] The subsequent loss of *P.copri* in our Indo-Immigrant and Indo-

Canadian cohorts aligned with their reduction of a high carbohydrate, high fiber diet, which was a

trend previously explored among Hadza and Nepali groups.[48] Plant-derived microbiota-accessible

carbohydrates (MACs) are required to maintain abundances of *P. copri*, whereas other species

such as *Bacteroides thetaiotaomicron* can utilize both plant and animal-derived MACs, allowing

for persistence in the gut.[48] This relationship suggests that *Prevotella* spp. abundance in the

Indian gut is highly linked to their specific dietary patterns. Furthermore, the generational loss of

*P. copri* in Indian immigrants prompts questions about gut health implications, meriting further

research on strain-specific *P. copri*-host interactions within this demographic.

A previous study found *Dialister succinatiphilus* and *Megasphaera* more abundant in Indo-

Immigrants versus Indo-Canadians, who had significantly higher *D. invisus.* Our data align with

this trend, with *D. invisus* enriched in Indo-Immigrants, *D. succinatiphilus* in Indo-Canadians, and

*Megasphaera* in our Indian cohort. While little is known about these taxa, a relationship between

a carbohydrate-rich diet and succinate-utilizing bacteria has been previously discussed,[49] with *D.*

*succinatiphilus* being a well-known species in this category. Additionally, *Ruminococcus torques,*

a known mucin degrader,[50] previously linked to IBD risk [51] and identified as a significant

contribution in a Crohn's Disease microbiome-risk score,[52] was notably enriched in Indo-

Canadians. Deeper investigations should explore the role of specific *R. torques* strains in the gut

and its implications in IBD should be explored further.

**India is Currently Industrializing**

India's industrialization is reshaping dietary habits, as reflected in our data showing

increased use of sunflower oil over traditional fats like ghee. This trend is likely influenced by

globalization[53] and dietary guidelines recommending a reduction in saturated fat intake and

increase in PUFA-rich vegetable oils. Overtime, these vegetable oils became inexpensive and

widely accessible in India, contributing to a shift away from traditional fats.[54] Coinciding with

increased PUFA consumption is the rising epidemic of several modern diseases in India such as

IBD[55], type-2 diabetes, and metabolic syndrome.[56] India is now in an accelerating incidence phase

of IBD[1], doubling the number of patients from 1990 to 2019.[57] While dietary studies in Indian IBD

patients are limited, previous research in other populations has shown that increased intake of

linoleic (n-6 PUFA) was associated with an increased risk for UC,[58] whereas increased intake of

$\alpha$-linolenic acid (n-3 PUFA) was associated with reduced UC-risk.[59] These patterns highlight the

importance of further investigating the potential role of vegetable oil consumption in relation to the

rising incidence of IBD in India.

In contrast, the westernized cohorts in our study reported cooking mainly with olive oil and

butter, and few reported to cook with vegetable oils rich in omega-6 PUFAs. While these self-

reports are subject to limitations, it is worth highlighting that while India has adopted the use of

white oils like sunflower oil and canola oil, North America may currently be shifting away from

them. Historically, the westernized diet has been characterized by a much higher intake of omega-

6 PUFA, leading to omega-6:omega-3 ratios of 20:1, which is far from the estimated ratio of 1:1

of our ancestors.[60] Excessive omega-6 PUFAs consumption has been associated with increased

risks of CVD,[61] cancer,[62] and autoimmune diseases.[63] Furthermore, with large-scale studies and

meta-analyses refuting the link between saturated fatty acids and CVD[64,65] and the growing

popularity of the Mediterranean Diet,[66] a cultural shift in the North America may be occurring, once

again preferring olive oil and butter,[67] which was an observation also recently reported from an

Australian cohort.[68]

Another factor of industrialization that is common in India is the overuse of antibiotics.

India has one of the highest threats for antimicrobial resistance,[69] as was reflective in our cohorts

with enriched KEGG identities in Indians and Indo-Immigrants that contain signatures for antibiotic

resistance genes, a pattern previously documented in the Indian gut.[27,70] The widespread use of

antibiotics in India not only increases the prevalence of antibiotic resistance, but also accelerates

microbiome turnover, which was indicative in the pathways enriched in our Indian cohort including

peptidoglycan synthesis, microbial growth/metabolism, and DNA synthesis/repair. While common

usage of antibiotics is a likely catalyst for cell turnover, the higher risk for infectious disease via

water contamination and inadequate sanitation in India may also serve as environmental triggers.

Our findings highlight India's unique transition phase, balancing western influences with traditional

lifestyle practices, all of which are shaping their gut microbiome. Future research should delve

deeper into how westernization is currently influencing the gut microbiome in Indians, as this will

likely pose future health challenges.

**Acculturation of Westernized Diet Drives Functional Shifts in Microbiome**

Coinciding with previous findings, we observed dietary acculturation from the traditional high-complex-carbohydrate Indian diet to an increased intake of NOVA Group 4 UPFs,[38,71,72] which are characterized by industrially made ingredients with non-nutritive additives like emulsifiers, flavouring and sweeteners.[73] This dietary shift might explain the enrichment of pathways tapping into glycogen storage in Indo-Canadians, suggesting microbes had inadequate nutrient availability from the host diet.[74] With a 49% rise in Group 4 UPF intake and significant fiber reduction in Indo-Canadians, concerns emerge regarding their gut health and chronic disease risks.

High UPF consumption has been linked to increased risks of CVD, metabolic disorders, micronutrient deficiencies, and anxiety and depression,[73] which may partly be due to imbalances in the gut microbiome that are a triggered from UPF. While there are several potential culprits involved in the ultra-processing of foods that may have mechanistic interactions with the gut, it is likely an interplay of several factors surrounding the consumption of UPFs. First, despite being affordable and accessible, UPFs are often high in calories and palatability, but low in essential nutrients, leading to overconsumption and displacement of nutritious meals.[75] Additionally, emulsifiers, used to mix components in UPF, have been extensively studied in murine models, showing their ability to modulate the gut microbiota, reduce mucus thickness and increase intestinal permeability, bacterial translocation and inflammation.[76] While studies translating these same mechanistic effects of emulsifiers in humans are scarce, a short-term prospective study (*N* = 588) found a positive association between dietary emulsifiers and the inflammatory biomarker glycoprotein acetyls.[77] Moreover, a randomized double-blind controlled-feeding study (*N* = 16) with the emulsifier carboxymethylcellulose (CMC) found varied responses in the treatment group, with two subjects exhibiting significant alterations in gut composition and microbial encroachment, whereas other participants seemed non-sensitive to CMC.[78,79] Although constrained by a small

sample size, these findings prompt the need to further investigate in large-scale cohorts, as this

may in part explain the inconsistent outcomes observed when transitioning from murine to human

studies. Overall, these findings underscore that emulsifiers may be one of many components in

UPF that influence the gut microbiome, but further human studies are necessary to fully

understand the complex role of UPF on the gut microbiome and inflammation. Future research

should also consider examining the synergistic effects of multiple food additives to mirror the

composition of UPFs typically consumed by humans.

**LIMITATIONS**

In addition to inherent limitations of self-reported food diaries (e.g. accuracy of participant

reporting), our study also faced limitations in representing cultural diets, primarily due to

inadequacies in the existing North American databases for Indian cuisine. This led to the use of

the EpiNu Nutritional software, an Indian dietary tool that is tailored for Indian dishes. Another

limitation was the shallow depth of shotgun sequencing and taxonomic classification methods

used restricted our ability to identify strain-level differences in bacteria like *P. copri*. Uneven

sample sizing across groups also presented a limitation; specifically, recruiting Indo-Canadians

was considerably more difficult, which may reflect a combination of cultural, social and

geographical barriers.

As this project focused on the relative abundance of *Prevotella* spp., quantitative PCR

could also help to determine total biomass and absolute abundances of specific taxa present in

the gut, as this may also be a key factor influencing disease risk.[80] Additionally, while

metagenomics can provide important insights into the genetic content present within the gut,

these data only represent the functional potential of the gut and does not provide information

regarding genetic expression and actual function of microbiome communities. Furthermore,

BugBase analyzes 16S data against an older version of the Greengenes database, which may

influence the accuracy of the predictions, especially in relation to the newer Greengenes2 and

MetaPhlAn databases used. Future work should also aim to resolve strain-level analysis as well

as parasite communities in the microbiomes of these populations, as this may differ in those living

in India versus Canada. Lastly, since this was a cross-sectional study, these data provide

associations between westernization and gut microbiome changes, and further investigation is

needed to understand the mechanistic interactions of specific factors of westernization on the gut

and IBD risk.

**Figure 8. Adoption of Westernized Diet is Associated with a Transition Away from Indian Gut Microbiome**

Graphical representation highlighting the main findings from our study. As Indo-Immigrants and Indo-Canadians subsequently increased ultra-processed food intake, their fiber consumption dramatically reduced. This change in dietary pattern was observed alongside a transition away from high *Prevotella* spp., which were found abundant in the Indian gut. Instead, Indo-Canadians adopted BloSSUM (Bloom or Selected in Societies of Urbanization/Modernization) taxa, which are commonly found in the westernized microbiome. High *Prevotella copri* in the Indian gut contributed to enriched carbohydrate-active enzymes (CAZymes) that are specialized to degrade complex carbohydrates common in their diet, which decline in Indian migrant cohorts. Additionally, the Indian microbiome displayed characteristics of a more robust gut, with predictive functions of higher stress tolerance and increased microbial cell turnover.

## CONCLUSIONS`

Consistent with previous studies, our research reveals that the Indian gut microbiome, enriched in *Prevotella* spp., is adapted to their high carbohydrate, high fiber diet. We discovered that the Indian microbiome exhibits characteristics of higher bacterial cell turnover, pathogenic potential and stress tolerance, which may reflect a chronic alteration in the response to a higher load of pathogen exposure. However, since our subjects were healthy and not displaying GI-related ailments, this may suggest that Indians have a higher resilience against stressors posed on the gut (Figure 8). These findings highlight the importance of studying diverse gut microbiomes, particularly in non-westernized and transitional populations, to better predict and manage diseases like IBD. Our study provides a snapshot on the gut microbiome changes in a population undergoing westernization, revealing a drastic transition within the gut microbiomes of Indo-Immigrants and Indo-Canadians, extending our knowledge of the strong effect of immigration and westernization on the gut microbiome. This transition likely reflects microbial adaptation to the dietary acculturation observed, providing insight into the dynamic nature within our gut. Future investigations should examine the health effects of microbiome changes amidst global westernization, as this transition will likely redefine health outcomes for both migrating populations and those living in rapidly modernizing nations.

**Acknowledgements**

We thank all the individuals who took part in this study. We thank the entire Ghosh family for

support. For dietary analysis, we thank Nadia Anvari. For lab support, we thank Mekenna Smith,

Hephzibah Bomide, Carson McComb, Ayva Lewis, Erik Kaila, Andrea Verdugo Meza and Jessica

Josephson.

**Funding**

This work was supported by the Natural Sciences and Engineering Research Council of Canada

(NSERC) and a UBC Killam Research Award to support D.L.G. on her sabbatical in India.

**Competing Interests**

DLG is on the Board of Directors for Crohn's and Colitis Canada and is also the co-founder and

CSO of Melius MicroBiomics Inc. where LDD is the Medical Science Writer.

**Patient and Public Involvement**

Not applicable.

**Ethics Approval**

This study was approved by the University of British Columbia Clinical Research Ethics Board

(UBC CREB) (H21-01555). Approval for sample collection in India was approved under IEC No

411/2018 and by UBC CREB (H17-01324). All participants signed a consent form prior to

participation. A Southeast Asian (SEA) Microbiome Biobank was created.

**Data Availability Statement**

Data supporting this study's findings are available within the paper and its Supplementary Information. The 16S rRNA gene sequences files and metagenomic sequence files of human fecal samples are publicly available on NCBI under accession number PRJNA1082632.

**Code Availability Statement**

The underlying code for this study is available on GitHub and can be accessed via this link:

https://github.com/leahdaloisio/India-Microbiome-Project

**Supplementary Material**

Please find Supplementary Methods and Supplementary Results attached.

**REFERENCES**

1.  Kaplan GG, Windsor JW. The four epidemiological stages in the global evolution of inflammatory bowel disease. *Nature Reviews Gastroenterology & Hepatology.* 2021;18(1):56-66.
2.  Benchimol E, Mack D, Guttmann A, et al. Inflammatory bowel disease in immigrants to Canada and their children: a population-based cohort study. *The american journal of gastroenterology.* 2015;110(4):553-563.
3.  Carroll MW, Hamilton Z, Gill H, et al. Pediatric inflammatory bowel disease among South Asians living in British Columbia, Canada: a distinct clinical phenotype. *Inflamm Bowel Dis.* 2016;22(2):387-396.
4.  Pinsk V, Lemberg DA, Grewal K, Barker CC, Schreiber RA, Jacobson K. Inflammatory bowel disease in the South Asian pediatric population of British Columbia. *Am J Gastroenterol.* 2007;102(5):1077-1083.
5.  Dhaliwal J, Tuna M, Shah BR, et al. Incidence of Inflammatory Bowel Disease in South Asian and Chinese People: A Population-Based Cohort Study from Ontario, Canada. *Clin Epidemiol.* 2021;13:1109.
6.  Giri S, Jena A, Kumar-M P, Muniswamy JR, Nath P, Sharma V. Burden of inflammatory bowel disease in India: analysis of the Global Burden of Disease study from 1990 to 2019. *Intestinal Research.* 2025.
7.  Fragiadakis GK, Smits SA, Sonnenburg ED, et al. Links between environment, diet, and the hunter-gatherer microbiome. *Gut Microbes.* 2019;10(2):216-227.

706    8.      Sonnenburg ED, Sonnenburg JL. The ancestral and industrialized gut microbiota and
707           implications for human health. *Nature Reviews Microbiology.* 2019;17(6):383-390.

708    9.      Parizadeh M, Arrieta M-C. The global human gut microbiome: genes, lifestyles, and diet.
709           *Trends Mol Med.* 2023;29(10):789-801.

710    10.    Vangay P, Johnson AJ, Ward TL, et al. US immigration westernizes the human gut
711           microbiome. *Cell.* 2018;175(4):962-972. e910.

712    11.    Abdill RJ, Adamowicz EM, Blekhman R. Public human microbiome data are dominated
713           by highly developed countries. *PLoS Biol.* 2022;20(2):e3001536.

714    12.    Gorzelak MA, Gill SK, Tasnim N, Ahmadi-Vand Z, Jay M, Gibson DL. Methods for
715           improving human gut microbiome data by reducing variability through sample
716           processing and storage of stool. *PLoS One.* 2015;10(8):e0134802.

717    13.    Bolyen E, Rideout JR, Dillon MR, et al. Reproducible, interactive, scalable and extensible
718           microbiome data science using QIIME 2. *Nat Biotechnol.* 2019;37(8):852-857.

719    14.    Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
720           resolution sample inference from Illumina amplicon data. *Nature Methods.*
721           2016;13(7):581-583.

722    15.    McDonald D, Jiang Y, Balaban M, et al. Greengenes2 unifies microbial data in a single
723           reference tree. *Nat Biotechnol.* 2024;42(5):715-718.

724    16.    Blanco-Míguez A, Beghini F, Cumbo F, et al. Extending and improving metagenomic
725           taxonomic profiling with uncharacterized species using MetaPhlAn 4. *Nat Biotechnol.*
726           2023:1-12.

727    17.    Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation.
728           *Genome Biol.* 2011;12:1-18.

729    18.    Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J. MicrobiomeAnalyst 2.0: comprehensive
730           statistical, functional and integrative analysis of microbiome data. *Nucleic Acids Res.*
731           2023;51(W1):W310–W318.

732    19.    Ward T, Larson J, Meulemans J, et al. BugBase predicts organism-level microbiome
733           phenotypes. *BioRxiv.* 2017:133462.

734    20.    Langille MG, Zaneveld J, Caporaso JG, et al. Predictive functional profiling of microbial
735           communities using 16S rRNA marker gene sequences. *Nat Biotechnol.* 2013;31(9):814-
736           821.

737    21.    Beghini F, McIver LJ, Blanco-Míguez A, et al. Integrating taxonomic, functional, and
738           strain-level profiling of diverse microbial communities with bioBakery 3. *Elife.*
739           2021;10:e65088.

740    22.    Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef: comprehensive and non-
741           redundant UniProt reference clusters. *Bioinformatics.* 2007;23(10):1282-1288.

742    23.    Caspi R, Altman T, Billington R, et al. The MetaCyc database of metabolic pathways and
743           enzymes and the BioCyc collection of Pathway/Genome Databases. *Nucleic Acids Res.*
744           2014;42(D1):D459-D471.

745    24.    Carter MM, Olm MR, Merrill BD, et al. Ultra-deep sequencing of Hadza hunter-gatherers
746           recovers vanishing gut microbes. *Cell.* 2023;186(14):3111-3124.

747    25.    Sonnenburg JL, Sonnenburg ED. Vulnerability of the industrialized microbiota. *Science.*
748           2019;366(6464).

26.  Jha AR, Davenport ER, Gautam Y, et al. Gut microbiome transition across a lifestyle
gradient in Himalaya. *PLoS Biol.* 2018;16(11):e2005396.
27.  Bhute S, Pande P, Shetty SA, et al. Molecular Characterization and Meta-Analysis of Gut
Microbial Communities Illustrate Enrichment of Prevotella and Megasphaera in Indian
Subjects. *Front Microbiol.* 2016;7:660.
28.  Kumbhare SV, Kumar H, Chowdhury SP, et al. A cross-sectional comparative study of gut
bacterial community of Indian and Finnish children. *Sci Rep.* 2017;7(1):10555.
29.  Jain A, Li XH, Chen WN. Similarities and differences in gut microbiome composition
correlate with dietary patterns of Indian and Chinese adults. *AMB Express.*
2018;8(1):104.
30.  Tandon D, Haque MM, R S, et al. A snapshot of gut microbiota of an adult urban
population from Western region of India. *PLoS One.* 2018;13(4):e0195643.
31.  Dhakan DB, Maji A, Sharma AK, et al. The unique composition of Indian gut microbiome,
gene catalogue, and associated fecal metabolome deciphered using multi-omics
approaches. *GigaScience.* 2019;8(3):01.
32.  Arumugam M, Raes J, Pelletier E, et al. Enterotypes of the human gut microbiome.
*Nature.* 2011;473(7346):174-180.
33.  Kulkarni A, Kumbhare SV, Dhotre DP, Shouche YS. Mining the Core Gut Microbiome
from a Sample Indian Population. *Indian J Microbiol.* 2019;59(1):90-95.
34.  Yatsunenko T, Rey FE, Manary MJ, et al. Human gut microbiome viewed across age and
geography. *Nature.* 2012;486(7402):222-227.
35.  Bodykevich G, Ibtisam A, Mao L. The gut microbiome belonging to Colombian adults of
younger generations has shifted to a more westernized composition compared to older
generations. *Undergraduate Journal of Experimental Microbiology and Immunology.*
2023;9.
36.  Fanfan D, Mulligan CJ, Groer M, et al. The intersection of social determinants of health,
the microbiome, and health outcomes in immigrants: A scoping review. *American
Journal of Biological Anthropology.* 2023;183(1):3-19.
37.  Olm MR, Dahan D, Carter MM, et al. Robust variation in infant gut microbiome assembly
across a spectrum of lifestyles. *Science.* 2022;376(6598):1220-1223.
38.  Copeland JK, Chao G, Vanderhout S, et al. The Impact of Migration on the Gut
Metagenome of South Asian Canadians. *Gut microbes.* 2021;13(1):1-29.
39.  Hertz S, Durack J, Kirk KF, et al. Microscopic colitis patients possess a perturbed and
inflammatory gut microbiota. *Dig Dis Sci.* 2022;67(6):2433-2443.
40.  Iljazovic A, Roy U, Gálvez EJ, et al. Perturbation of the gut microbiome by Prevotella spp.
enhances host susceptibility to mucosal inflammation. *Mucosal Immunol.* 2020:1-12.
41.  Scher JU, Sczesnak A, Longman RS, et al. Expansion of intestinal Prevotella copri
correlates with enhanced susceptibility to arthritis. *elife.* 2013;2:e01202.
42.  Moreno-Indias I, Sánchez-Alcoholado L, García-Fuentes E, Cardona F, Queipo-Ortuño MI,
Tinahones FJ. Insulin resistance is associated with specific gut microbiota in appendix
samples from morbidly obese patients. *American Journal of Translational Research.*
2016;8(12):5672.

43.   Wang Y, Ames NP, Tun HM, Tosh SM, Jones PJ, Khafipour E. High molecular weight
barley β-glucan alters gut microbiota toward reduced cardiovascular disease risk. *Front*
*Microbiol.* 2016;7:129.
44.   Bibbò S, Dore MP, Pes GM, Delitala G, Delitala AP. Is there a role for gut microbiota in
type 1 diabetes pathogenesis? *Ann Med.* 2017;49(1):11-22.
45.   Borgo F, Verduci E, Riva A, et al. Relative abundance in bacterial and fungal gut microbes
in obese children: a case control study. *Childhood Obesity.* 2017;13(1):78-84.
46.   Prasoodanan PK V, Sharma AK, Mahajan S, et al. Western and non-western gut
microbiomes reveal new roles of Prevotella in carbohydrate metabolism and mouth–gut
axis. *NPJ Biofilms and Microbiomes.* 2021;7(1):1-17.
47.   Tett A, Huang KD, Asnicar F, et al. The Prevotella copri complex comprises four distinct
clades underrepresented in westernized populations. *Cell Host Microbe.* 2019;26(5):666-
679. e667.
48.   Gellman RH, Olm MR, Terrapon N, et al. Hadza Prevotella require diet-derived
microbiota-accessible carbohydrates to persist in mice. *Cell Rep.* 2023;42(11).
49.   Nakayama J, Yamamoto A, Palermo-Conde LA, et al. Impact of westernized diet on gut
microbiota in children on Leyte island. *Front Microbiol.* 2017;8:197.
50.   Png CW, Lindén SK, Gilshenan KS, et al. Mucolytic bacteria with increased prevalence in
IBD mucosa augmentin vitroutilization of mucin by other bacteria. *Official Journal of the*
*American College of Gastroenterology.* 2010;105(11):2420-2428.
51.   Ali F, Reddy V, Raza M, et al. S3194 Metformin: A Therapeutic Adjunct for Patients With
Inflammatory Bowel Disease? *Official journal of the American College of*
*Gastroenterology.* 2020;115:S1676.
52.   Garay JAR, Turpin W, Lee S-H, et al. Gut microbiome composition is associated with
future onset of Crohn's Disease in healthy first-degree relatives. *Gastroenterology.*
2023;165(3):670-681.
53.   USDA. The Role of Policy and Industry Structure in India's Oilseed Markets. *USDA*
*Economic Research Service.* 2007.
54.   Mani I, Kurpad AV. Fats & fatty acids in Indian diets: Time for serious introspection. *The*
*Indian Journal of Medical Research.* 2016;144(4):507.
55.   Snell A, Segal J, Limdi J, Banerjee R. Inflammatory bowel disease in India: challenges and
opportunities. *Frontline Gastroenterol.* 2020;12(5):390-396.
56.   Lakshmipriya N, Gayathri R, Praseena K, et al. Type of vegetable oils used in cooking and
risk of metabolic syndrome among Asian Indians. *Int J Food Sci Nutr.* 2013;64(2):131-
139.
57.   Dharni K, Singh A, Sharma S, et al. Trends of inflammatory bowel disease from the
Global Burden of Disease Study (1990-2019). *Indian J Gastroenterol.* 2023:1-11.
58.   Investigators IiES. Linoleic acid, a dietary n-6 polyunsaturated fatty acid, and the
aetiology of ulcerative colitis: a nested case–control study within a European
prospective cohort study. *Gut.* 2009;58(12):1606-1611.
59.   Ananthakrishnan AN, Khalili H, Konijeti GG, et al. Long-term intake of dietary fat and risk
of ulcerative colitis and Crohn's disease. *Gut.* 2014;63(5):776-784.
60.   Simopoulos AP. An increase in the omega-6/omega-3 fatty acid ratio increases the risk
for obesity. *Nutrients.* 2016;8(3):128.

61.  Simopoulos AP. Omega-6/omega-3 essential fatty acid ratio and chronic diseases. *Food*
*reviews international.* 2004;20(1):77-90.
62.  De Lorgeril M, Salen P. New insights into the health effects of dietary saturated and
omega-6 and omega-3 polyunsaturated fatty acids. *BMC Med.* 2012;10(1):1-5.
63.  Simopoulos AP. Evolutionary aspects of diet, the omega-6/omega-3 ratio and genetic
variation: nutritional implications for chronic diseases. *Biomed Pharmacother.*
2006;60(9):502-507.
64.  Dehghan M, Mente A, Zhang X, et al. Associations of fats and carbohydrate intake with
cardiovascular disease and mortality in 18 countries from five continents (PURE): a
prospective cohort study. *The Lancet.* 2017;390(10107):2050-2062.
65.  Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Meta-analysis of prospective cohort studies
evaluating the association of saturated fat with cardiovascular disease. *The American*
*Journal of Clinical Nutrition.* 2010;91(3):535-546.
66.  Vetrani C, Piscitelli P, Muscogiuri G, et al. "Planeterranea": An attempt to broaden the
beneficial effects of the Mediterranean diet worldwide. *Frontiers in Nutrition.*
2022;9:973757.
67.  Fernandes J, Fialho M, Santos R, et al. Is olive oil good for you? A systematic review and
meta-analysis on anti-inflammatory benefits from regular dietary intake. *Nutrition.*
2020;69:110559.
68.  Wilson NA, Villani A, Mantzioris E. Preferences for dietary oils and fats in cooking and
food preparation methods: a cross-sectional analysis of Australian adults. *Proc Nutr Soc.*
2023;82(OCE2):E72.
69.  Gandra S, Joshi J, Trett A, Lamkang AS, Laxminarayan R. Scoping report on antimicrobial
resistance in India. *Center for Disease Dynamics, Economics & Policy: Washington, DC,*
*USA.* 2017;2017:1-146.
70.  Gupta M, Didwal G, Bansal S, et al. Antibiotic-resistant Enterobacteriaceae in healthy gut
flora: A report from north Indian semiurban community. *Indian J Med Res.*
2019;149(2):276-280.
71.  Lesser IA, Gasevic D, Lear SA. The association between acculturation and dietary
patterns of South Asian immigrants. *PLoS One.* 2014;9(2):e88495.
72.  Noor S, Dehghan M, Lear SA, et al. Relationship between diet and acculturation among
South Asian children living in Canada. *Appetite.* 2020;147:104524.
73.  Tristan Asensi M, Napoletano A, Sofi F, Dinu M. Low-Grade Inflammation and Ultra-
Processed Foods Consumption: A Review. *Nutrients.* 2023;15(6):1546.
74.  Esteban-Torres M, Ruiz L, Rossini V, Nally K, van Sinderen D. Intracellular glycogen
accumulation by human gut commensals as a niche adaptation trait. *Gut Microbes.*
2023;15(1):2235067.
75.  Dinu M, Tristan Asensi M, Pagliai G, et al. Consumption of ultra-processed foods is
inversely associated with adherence to the Mediterranean diet: a cross-sectional study.
*Nutrients.* 2022;14(10):2073.
76.  Bancil AS, Sandall AM, Rossi M, Chassaing B, Lindsay JO, Whelan K. Food additive
emulsifiers and their impact on gut microbiome, permeability, and inflammation:
mechanistic insights in inflammatory bowel disease. *Journal of Crohn's and Colitis.*
2021;15(6):1068-1079.

77.    Um CY, Hodge RA, Tran HQ, Campbell PT, Gewirtz AT, McCullough ML. Association of
emulsifier and highly processed food intake with circulating markers of intestinal
permeability and inflammation in the cancer prevention study-3 diet assessment sub-
study. *Nutr Cancer.* 2022;74(5):1701-1711.
78.    Chassaing B, Compher C, Bonhomme B, et al. Randomized controlled-feeding study of
dietary emulsifier carboxymethylcellulose reveals detrimental impacts on the gut
microbiota and metabolome. *Gastroenterology.* 2022;162(3):743-756.
79.    Daniel N, Wu GD, Walters W, et al. Human intestinal microbiome determines
individualized inflammatory response to dietary emulsifier carboxymethylcellulose
consumption. *Cell Mol Gastroenterol Hepatol.* 2024;17(2):315-318.
80.    Vandeputte D, Kathagen G, D'hoe K, et al. Quantitative microbiome profiling links gut
community variation to microbial load. *Nature.* 2017;551(7681):507-511.
891