The transition from a non-westernized to westernized gut microbiome in Indian-Immigrants

and Indo-Canadians is associated with dietary acculturation

Leah D. D'Aloisio[1], Mamatha Ballal[5], Sanjoy Ghosh[1,10], Natasha Haskey[1], Nijiati Abulizi[1], Ramin

Karimianghadim[2], Chuyi Liu[3], Pacha Sruthi[4], Lakshmipriya Nagarajan[4], Sudha Vasudevan[4],

Vignesh Shetty[5,6], Mrudgandha Purandare[5], Ushasi Bhaumik[7], Debaki Ranjan Howlader[8],

Sepideh Pakpour[1,9], Jacqueline Barnett[10], Deanna L. Gibson* [1,10,11].

8
9   [1] Biology, University of British Columbia Okanagan Campus, Kelowna, Canada
10
11 [2] Applied Science, University of British Columbia Okanagan Campus, Kelowna, Canada
12
13 [3] College of Pharmacy and Nutrition, University of Saskatchewan, Saskatchewan, Canada
14
15 [4] Food & Nutrition Research, Madras Diabetes Research Foundation, Chennai, India
16
17 [5] Enteric Disease Division, Department of Microbiology, Kasturba Medical College, Manipal Academy of Higher
18 Education, Manipal, India
19
20 [6] Department of Medicine, University of Cambridge, Cambridge, United Kingdom
21
22 [7] School of Medicine, University of Maryland, Maryland, United States
23
24 [8] Department of Veterinary Pathobiology, University of Missouri, Missouri, USA
25
26 [9] School of Engineering, University of British Columbia Okanagan Campus, Kelowna, Canada
27
28 [10] Biochemistry and Molecular Biology Program, University of British Columbia Okanagan Campus, Kelowna, Canada
29
30 [11] Department of Medicine, University of British Columbia Okanagan Campus, Kelowna, Canada
31
32
33 * Corresponding author: deanna.gibson@ubc.ca
34

Background: Indian immigration to westernized countries has surged in recent years, with this

demographic facing a higher risk of Inflammatory Bowel Disease (IBD) after migration. Studying

the gut microbiome is crucial to understanding the factors contributing to IBD, yet Indians remain

underrepresented in microbiome research,

Objective: This study investigates the gut microbiome of Indians residing in India, Indian

immigrants in Canada, and Canadians of Indian ancestry to understand the impact of

westernization on their gut. Given the higher risk of IBD among Indo-Canadians, these results

provide insight into how westernization of the gut microbiome may influence incidence of IBD.

Stool samples were collected from healthy Indians in India, Indian Immigrants in Canada, Indo-

Canadians, plus Canadian-borne individuals with European descent and Euro-Immigrants as

non-Indian/westernized controls (N = 174). 16S rRNA gene and shotgun sequencing compared

microbial taxa and functional profiles across groups. Dietary and demographic data were collected

to assess lifestyle patterns.

Results: Indians and Indo-Immigrants had distinct gut microbiotas from westernized groups, with

high abundances of *Prevotella* spp. and a CAZyme profile reflecting their traditional high complex

carbohydrate diet. The Indian microbiome also showed characteristics of high bacterial cell

turnover, pathogenic potential, and stress tolerance, indicating their gut may be better primed to

handle stressors on the gut. Indo-Canadians displayed a transitional microbiome towards

westernization, which followed a pattern of increasing dietary acculturation.

Conclusions: The shift from a non-westernized to westernized microbiome in Indo-Canadians

highlights the significant influence that westernization and dietary acculturation has on the gut

microbiome. Given that 44% of the Canadian population is made up of first- and second-

generation immigrants and westernized practices are being adopted worldwide, future research

should investigate the health implications that such microbiome transitions have on immigrant

populations and newly industrialized nations.

BACKGROUND

The growing immigrant population in North America is rapidly westernizing through the

adoption of practices like reduced physical activity, changes in hygiene/medical practices, and

acculturation of the westernized diet, commonly characterized by caloric dense, highly processed

foods with low fiber intake. Amidst this backdrop are health disparities emerging in these

populations. Furthermore, the global spread of industrialization raises concern for the increasing

prevalence of modern diseases.[1] For instance, inflammatory bowel disease (IBD) is increasing in

prevalence globally, and alarming evidence reveals a specifically increased risk in young Indian

immigrants and Indians born in Westernized countries.[2-4] A likely contributor to this elevated risk

stems from changes in the gut microbiome as a response to living in an industrialized

environment.

The industrialized microbiome has been previously characterized by the decline of

VANISH (volatile and/or associated negatively with industrialized societies of humans) taxa,

frequently found in populations adhering to traditional lifestyles, and the rise of BloSSUM (bloom

or selected in societies of urbanization/modernization) taxa, which appear in modern,

industrialized populations.[5,6] Furthermore, health consequences following the assimilation into

Westernized culture have been previously documented. For example, non-industrialized Irish

Travellers who adopted modern practices and Hmong and Karen migrants from Thailand who

relocated to the United States (US) both experienced gut microbiome shifts that were associated

with an increased risk for metabolic disorders.[7,8] Currently, there is only one study published on

the Indian immigrant gut microbiome, highlighting the gap in non-western/non-European

microbiome research. Indeed, over 71% of all microbiome research comes from Westernized

cohorts, skewing our interpretation of a healthy human gut.[9] Here, we aimed to investigate the

distinction between the Indian and Westernized gut microbiome and what changes occur in the

gut when Indians migrate to Canada, alongside the adoption of the westernized diet.

METHODS

Study Design

A prospective case-control study was conducted including three Indian groups: (1) Indians

residing in India ("Indians"), (2) Indians who migrated to Canada ("Indo-Immigrants") and (3)

Canadians born in Canada from Indian ancestry ("Indo-Canadians"). Individuals born in Canada

with European ancestry ("Euro-Canadian") were used as a control, and immigrants with European

ancestry from a Westernized country ("Euro-Immigrants") as a westernized immigrant control.

Healthy participants aged 17-55 years were recruited. Exclusion criteria included pregnancy,

diagnosis of any chronic inflammatory condition, antibiotics used or travel to India within three

months prior to stool collection. Participants completed a demographic questionnaire to assess

baseline characteristics. Immigrants also completed a lifestyle survey to assess changes in

lifestyle and immigration-related stress.

Stool Collection & DNA Extraction

Stool samples were collected from India (Kolkata and Manipal) and Canada (Kelowna)

and processed on site as previously described.[10] DNA was sent to Gut4Health for 16S rRNA gene

sequencing on the Illumina MiSeq platform (V4-V4, 515F/806R primers) (~75,800 reads per

sample) and to the Center for Health Genomics and Informatics for shotgun sequencing on the

Illumina NovaSeq platform (~9.9M reads per sample). Additional details are in Supplementary

Methods.

Taxonomic Analysis

Demultiplexed 16S rRNA gene reads were processed in QIIME 2 (Version 2022.2)[11] with

DADA2[12] for quality control and taxonomic classification with GreenGenes2 (10.28.22).[13]

Demultiplexed raw shotgun reads underwent quality assessment with FastQC/MultiQC and used

KneadData (Version 0.10) for trimming and quality control. Taxonomic profiling was conducted

using MetaPhlAn4,[14] then each sample was normalized to relative abundances. Diversity metrics

included Shannon and Pielou's Evenness (Kruskal-Wallis), and Bray Curtis and Weighted

UniFrac (PERMANOVA). Differentially abundant bacteria were determined using the Linear

Discriminate Analysis (LDA) Effect Size (LEfSe) algorithm with a 3.5 threshold and alpha 0.05.[15]

Using shotgun data, the normalized relative abundance MetaPhlAn4 output table was imported

into MicrobiomeAnalyst 2.0 to apply a Random Forest algorithm ranked according to their Mean

Decrease Accuracy score, indicating the extent to which each taxon's presence or absence

influences the overall accuracy of classifying samples into their respective cohort[16].[16] These taxa

were then visualized in a heatmap. Using 16S rRNA gene data, BugBase was used to predict

microbiome phenotypes,[17,18] and associations between beta diversity, dietary patterns and

baseline characteristics were analyzed with distance-based redundancy analysis (dbDRA).

Additional details for taxonomic analysis methods can be found in Supplementary Methods.

Functional Profiling

Shotgun reads were analyzed for functional potential using HUMAnN (Version 3.6).[19]

Gene family and pathway abundances were annotated using the UniRef90[20] and MetaCyc[21]

databases, respectively, then normalized to reads per kilobase to account for gene length. Both

tables were further normalized to relative abundance to account for sampling depth. The

normalized gene family abundance output was then regrouped to the CAZy and KEGG

databases. To determine differentially abundant genes/pathways across cohorts, tables were

collapsed to perform LEfSe analysis on the unstratified data using a one-against-all strategy with

a threshold of 3.0 on the logarithmic LDA score and an alpha of 0.05 for Kruskal-Wallis test among

classes.[15] Differentially abundant features were then plotted as a heatmap using

MicrobiomeAnalyst 2.0.[16]

Nutritional Analysis

Participants filled out a 3-day food log prior to their stool collection, then researchers

retrospectively inputted this data into the ESHA Food Processor® software (Version 11.11.0) to

obtain the calculated nutrient intake for each subject. To mitigate bias from using a North

American nutritional software, Indian and Indo-Immigrant food logs were also entered into the

MDRF EpiNu® Nutritional software, tailored for Indian population. Additional details on our

approach for dietary analysis can be found in Supplementary Methods.

Statistical Analysis

Statistical analyses for demographics and dietary data were performed using both R

statistical software and GraphPad Prism (Version 10.0.3). We applied the Kruskal-Wallis test

coupled with Dunn's post hoc analysis for nonparametric data. For data that passed normality, an

ordinary one-way ANOVA with post hoc Tukey's multiple comparisons test was performed.

Categorical data were analyzed using either a Chi-Square, Fisher's Exact or Fisher's Exact with

Monte Carlo simulation, depending on the data structure. Results were reported as median values

and interquartile range (IQR), unless otherwise specified.

RESULTS

Participant Characteristics

The study included Indians ($n = 61$), Indo-Immigrants ($n = 32$), Indo-Canadians ($n= 17$),

Euro-Canadians ($n = 41$) and Euro-Immigrants ($n = 23$) (Figure S1). Baseline demographics were

comparable across groups, though Indians were older and consumed significantly less alcohol

($p_{BONF} = 3.29e-09$) (Table 1). Indo-Immigrants exhibited more of a change towards westernization

($p = 0.002$) and higher immigration-related stress ($p = 0.04$), with no significant differences in age

at or years since immigration (Table 2).

Indian Microbiota is Distinctive from Westernized Groups

To examine the taxa and complexity of the microbial ecosystem in Indians compared to

Westerners, alpha diversity revealed the lowest diversity in Indians and highest in Euro-

Canadians (Figure 1A-B). Bray Curtis dissimilarity of shotgun sequence data showed significant

differences across groups, with 45.3% of variation explained on the first 2 axes (Figure 1C).

Indians and Indo-Immigrants' microbiotas differed from westernized cohorts (Table 3), and each

other (pseudo-F = 9.132, $p_{BONF} = 0.01$). Weighted UniFrac also demonstrated distinctions in the

Indian and Indo-Immigrant microbiomes (pseudo-F = 25.98, $p_{BONF} = 0.01$), with 61.8% of the

variation explained on the first 2 axes (Figure 1D) with similar findings from 16S rRNA gene

amplicon sequence data (Figure S2).

Figure 1. Indian and Indo-Immigrant Microbiotas are Distinct from Westernized Groups

Alpha Diversity - Boxes represent the interquartile range (IQR) between the first and third quartiles, the horizontal line indicates the median, and whiskers are the upper and lower values within 1.5 times the IQR. Alpha diversity was compared across groups using the Kruskal-Wallis test. **(A)** Pielou's Evenness showed significant differences (H = 85.7, p = 1.07e-17). Pairwise comparisons reveal Indians

(n = 61) had significantly lower evennnes than Euro-Canadians (n = 41) (H = 56.2, q = 6.51e-13), Euro-Immigrants (n = 23) (H = 17.0, q = 7.32e-05) and Indo-Canadians (n= 17) (H = 10.8, q =1.69e-03). Euro-Canadian controls also score significantly higher evenness than Indo-Immigrants (n = 32) (H = 41.4, q = 6.09e-10) and Indo-Canadians (H = 21.7, q = 8.10e-06). **(B)** Similar results were also found with

Shannon's Diversity, showing significant differences (H = 79.8, p = 1.89e-16). *** = q ≤ 0.001, ** = q ≤ 0.01, * = q ≤ 0.05

Beta Diversity - Beta diversity was explored with Bray Curtis dissimilarity and Weighted UniFrac, using Pairwise Permutational Multivariate Analysis of Variance (PERMANOVA) to test differences between groups. **(C)** Bray Curtis principal coordinate analysis (PCoA) plot shows 45.3% of variation was captured on the first two axes. Pairwise comparisons with Bonferroni correction revealed

significant differences between most groups, with distinct clustering of Indians and Indo-Immigrants from westernized cohorts. **(D)** Weighted UniFrac PCoA plot shows 61.8% of the variation was captured on the first two axes, with significant differences also detected. Pairwise comparisons results can be found in Table S1.

(E) LEfSe (Linear discriminate analysis Effect Size) cladogram results, depicting differentially abundant bacteria across cohorts, with genus set as lowest taxonomic rank. Indian gut displayed a dominance of Bacteroidetes and Proteobacteria, Euro-Canadians showed high abundance of Firmicutes. Indo-Immigrants had high abundances of VANISH taxa including *Eubacterium* and Erysipelotrichaceae. Indo-Canadians harboured high abundances of both VANISH taxa like *Collinsella*, and BloSSUM taxa such as *Anaerostipes* and *Blautia*.

LEfSe identified dominant taxa in each group, with Bacteroidota (formerly Bacteroidetes)

and Pseudomonadota (formerly Proteobacteria) in Indians and Firmicutes in Euro-Canadians

(Figure 1E, Table S1). Indo-Immigrants displayed high abundances of *Eubacterium* and

Erysipelotrichaceae, previously categorized as VANISH taxa.[22] As expected, genera common in

the industrialized microbiome were enriched in the westernized European cohorts such as

Alistipes, *Bacteroides*, and Clostridia, previously categorized as BloSSUM taxa.[22] Indo-

Canadians represented a transitional gut microbiota, harbouring both VANISH and BloSSUM

taxa.[23] Similar results were found in 16S rRNA gene sequence data (Figure S3, Table S1).

When assessing differences in bacterial species abundances amongst groups, LEfSe

showed that Indians harboured higher abundances of *Phocaeicola plebeius*, *Megasphaera sp.*,

Dialister hominis, and *Escherichia coli* (Figure 2). Among Indo-Immigrants, *Dorea longicatena*,

Faecalibacterium prausnitizii and *Lachnospiraceae bacterium* were notably abundant, while Indo-

Canadians had increased abundances of two *Ruminoccocus torques* SGBs, *Blautia wexlerae,*

Eubacterium rectale, Blautia massiliensis, and *Anaerostipes hadrus*. Taxa enriched in Euro-

Canadians included *Phocaeicola vulgatus*, *Lachnospiraceae bacterium, Clostridium sp.,* and

Blautia faecis, and Euro-Immigrants showed enrichment of *Fusicatenibacter saccharivorans,*

Clostridia bacterium, and *Blautia obeum.* A key distinguishing feature from Indians to Indo-

Table 1. Baseline Characteristics of Participants

Baseline characteristic	Indian *n = 61*	Indo-Immigr *n = 32*	Indo-Can *n = 17*	Euro-Can *n = 41*	Euro-Immigr *n = 23*	*P* value
Sex						
Female	37 (61)	18 (60)	6 (35)	16 (39)	10 (43)	0.10 [a]
Male	24 (39)	12 (40)	11 (65)	25 (61)	13 (57)	
Age, (years)						
Median (IQR)	31 (12)	25 (8)	23 (7)	24 (5)	29 (15)	
Min-max	17-52	18-53	18-38	19-50	19-53	
Age classification, n (%)						
17-29 years	26 (43)	27 (84)	14 (82)	32 (78)	12 (52)	0.002 *[e]
30-39 years	21 (34)	4 (13)	3 (18)	4 (10)	7 (30)	
40-49 years	10 (16)	0 (0)	0 (0)	4 (10)	2 (9)	
50-55 years	4 (7)	1 (3)	0 (0)	1 (2)	2 (9)	
Mode of birth, n (%)						
Vaginal	44 (75)	19 (63)	14 (88)	34 (83)	19 (83)	0.23 [a]
Caesarian	15 (25)	11 (37)	2 (12)	7 (17)	4 (17)	
BMI						
Mean (SD)	24 (3)	24 (4)	22 (3)	23 (3)	22 (2)	
Min-max	17-32	17-34	17-30	17-31	18-25	
BMI classification, n (%)						
Underweight (< 18.5)	3 (5.0)	1 (4.2)	2 (12.5)	3 (8.0)	1 (5.0)	0.08 [b]
Normal weight (18.5 - < 25)	40 (66.7)	14 (58.3)	12 (75)	24 (63.0)	21 (95.0)	
Overweight (25 - < 30)	13 (21.7)	8 (33.3)	2 (12.5)	10 (26.0)	0 (0)	
Obese (> 30)	4 (6.7)	1 (4.2)	0 (0)	1 (3.0)	0 (0)	
Smokers, n (%)						
Yes	6 (10)	0 (0)	0 (0)	2 (5)	2 (9)	0.2908 [c]
No	56 (90)	32 (100)	17 (100)	39 (95)	21 (91)	
Alcohol, n (%)						
Yes	15 (24)	15 (47)	14 (82)	33 (80)	18 (78)	<0.0001†[a]
No	47 (76)	17 (53)	3 (18)	8 (20)	5 (22)	
Physical activity- walking (mins)						
Median (IQR)	NA	90 (128)	90 (120)	60 (75)	60 (90)	0.56 [d]
Min-max	NA	0-330	30-300	12-480	12-420	
Physical activity- exercise (days)						
Median (IQR)	NA	3 (2)	3 (3)	4 (3)	4.5 (3)	0.18 [e]
Min-max	NA	0-7	0-7	0.5-6.5	1-6	
Socioeconomic class, n (%)						
Low	15 (25)	8 (25)	5 (33)	10 (25)	4 (17)	0.70 [e]

Middle	40 (67)	21 (66)	7 (47)	26 (65)	14 (61)	
Upper	5 (8)	3 (9)	3 (20)	4 (10)	5 (22)	
Lifestyle born/raised, n (%)						
Rural	NA	2 (6)	5 (31)	5 (12)	5 (22)	0.10 [c]
Urban	NA	30 (94)	11 (69)	36 (88)	18 (78)	

Abbreviations: SD, standard deviation; IQR, interquartile range; BMI, body mass index; [a]Chi-Square, [b]ANOVA, [c]Fisher's Exact, [d]Kruskal-Wallis, [e]Fisher's Exact (Monte Carlo); * Bonferroni correction revealed significant differences between Indian vs. Indo-Immigr (p_{BONF} = 0.006), Indian vs. Euro-Can (p_{BONF} = 0.03); † Bonferroni correct revealed significant differences (p_{BONF} = 3.29e-09).

Table 2. Characteristics of Immigrant Participants

Characteristic	Indo-Immigr $n = 32$	Euro-Immigr $n = 23$	P value
△ westernization, n (%)			
No westernization	0 (0)	4 (18)	0.002 [a]
Little westernization	12 (38)	13 (59)	
Moderate westernization	17 (53)	4 (18)	
High westernization	3 (9)	1 (4)	
Immigration-related stress, n (%)			
Low stress	22 (69)	21 (91)	0.04 [a]
Moderate stress	8 (25)	2 (9)	
High stress	2 (6)	0 (0)	
Age at immigration, n (%)			
0-3 years	0 (0)	1 (4)	0.19 [b]
4-17 years	9 (28)	3 (13)	
18-49 years	23 (72)	19 (83)	
Years since immigration			
Median (IQR)	4 (6)	3 (9)	0.65 [c]
Min-max	0-22	0-48	

Abbreviations: IQR, interquartile range; [a]Chi-Square for Trend, [b]Fisher's Exact, [c]Mann-Whitney

Table 3. PERMANOVA Results from Beta Diversity Analysis

	Bray Curtis		Weighted UniFrac	
Pair	pseudo-F	P value	pseudo-F	P value
Indo-Immigr vs. Indo-Can	4.215003	0.01	4.670629	0.01
Indo-Immigr vs. Euro-Can	8.976951	0.01	11.932016	0.01
Indo-Immigr vs. Euro-Immigr	5.313757	0.01	6.903393	0.01

Indo-Immigr vs. Indian	9.132465	0.01	25.984365	0.01
Indo-Can vs. Euro-Can	3.084292	0.01	5.409267	0.01
Indo-Can vs. Euro-Immigr	2.166793	0.01	2.360739	0.10
Indo-Can vs. Indian	8.660676	0.01	25.206253	0.01
Euro-Can vs. Euro-Immigr	1.163757	1.00	1.116192	1.00
Euro-Can vs. Indian	18.284343	0.01	49.611215	0.01
Euro-Immigr vs. Indian	11.182855	0.01	31.565399	0.01

Adjusted *P* values are displayed, which were obtained using 999 permutations and Bonferroni correction applied. Beta diversity data from shotgun sequence data.

Canadians was the decline in *Prevotella* spp., with *Prevotella copri* (clade A) being over five times

more abundant in Indians (Table 4). Overall, these findings demonstrate that the gut microbiome

in Indians are distinct from the West, and Indo-Canadians exhibit a transition towards the

industrialized microbiome.

Indian Gut Microbiome Predicted to be More Robust

To understand potential functionality of the gut microbiota from Indians, we used BugBase

predictions (Figure 3), finding Indians appeared to demonstrate higher stress tolerance driven by

Pseudomonadota (Proteobacteria). The Indian cohorts had the highest potentially pathogenic

bacteria (p_{FDR} = 2.07e-18) mostly dominated by Gram negative bacteria driven mainly by a

combination of Pseudomonadota, Bacteroidota (Bacteroidetes), and Firmicutes. In addition, the

Indian gut microbiome had the highest potential for biofilm formation. Overall, these results

suggest the India gut microbiome may be more robust, as healthy individuals were predicted to

harbour microbes with higher pathogenic potential and stress tolerance.

Gut Functional Potential in Indians is Distinctive from Westernized Groups

To further understand functional attributes of the Indian gut microbiome, pathway analysis using MetaCyc was used and revealed pathways enriched in lipopolysaccharide (LPS) production, peptidoglycan synthesis, microbial growth/metabolism, protein synthesis and DNA synthesis/repair (Figure 4). In Indo-Canadians, indications of increased glucose storage were detected. Euro-Canadians had increased pathways related to fatty acid biosynthesis, and Euro-Immigrants expressed pathways indicating sugar metabolism (Table S2). CAZy gene family analysis revealed distinct profiles in the Indian gut including glycoside hydrolases (GH)43, GH51, and GH10, all of which are known for their degradation of plant-based polysaccharides (Table S3). *P. copri* was the top taxa contributing to these CAZy families (Figure 5A). In addition, KEGG identities K00561 and K18220 were enriched in Indians and Indo-Immigrants, respectively, both recognized for antimicrobial resistance genes (Figure 5B). Overall, these results suggest that both environmental factors, diets and drugs, were influencing the gut microbiome of Indians.

Westernized Dietary Acculturation Observed in Indo-Canadians

To understand the role of diet associated with the gut microbiome in our Indian cohort, we collected and analyzed dietary recalls using both validated software from North America and from India to account for Indian foods. Indo-Canadians had the highest daily intake of NOVA Group 4 Ultra-Processed Food (UPF), constituting 61% of their caloric intake (95% CI = 467.9 – 780.5 kcal, per 1000 kcal), while Indians consumed significantly less than all other cohorts, reporting 12% of their daily intake (95% CI = 63.66 – 127.1 kcal per 1000 kcal) (Figure 6). Consistent with high UPF intake, total fiber was also found to be the lowest in Indo-Canadians, opposite in Indians who consumed more fiber than Indo-Canadians ($p = 0.0028$) and Euro-Canadians ($p = 0.0110$).

Figure 2. Significantly Different Gut Bacterial Abundances Detected Across Cohorts

Taxonomic stacked bar plots depicting relative abundances of top 18 most abundant bacteria across all samples (N = 174; Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). Both **(A)** 16S rRNA and **(B)** shotgun sequence data demonstrate a high abundance of *Prevotella* spp. in Indians that declines in Indian migrant cohorts. Overall gut bacterial composition is both distinct and more heterogenous in Indians compared to all other cohorts. **(C)** LEfSe results of top differentially abundant bacteria reveals distinct patterns in bacterial abundances in those living in Canada versus India. A Kruskal-Wallis test was performed with a significance (α) at 0.05 for one-against-all comparisons. Differentially abundant bacteria were detected using a Linear Disciminate Analysis (LDA) score (equal to or greater than 3.5). In Indians, *Prevotella copri* (clade A SGB1626) was over 5 times more abundant (LDA score = 4.73, p = 9.25e-13). **(D)** Heatmap generated in MicrobiomeAnalyst 2.0, displaying taxa identified by Random Forest as key features that contribute to the predictive accuracy of classifying samples into their respective groups (Figure S4). Features were filtered for minimum 4 counts in 20% or more samples, and low variance filter of 10%. Bars on the top represent the cohorts that group the individual sample columns displayed. Taxa are labelled in rows, with taxonomic rank noted before the bacteria name. Colours on heatmap represent the relative abundances of each bacteria in a given sample. **(E)** Heatmap generated in R Studio, displaying the average relative abundances of *P. copri* clades in each cohort. Clade A relative abundances were 10.8 for Indians, 5.89 for Indo-Immigrants, and 2.75 for Indo-Canadians. A similar pattern of decreasing abundance across Indian cohorts was also observed for Clades B, C and D. All clades were lowest in abundance in Euro-Canadians.

Table 4. Differentially abundant bacteria from shotgun sequence data

Taxon	Log Score	Class	LDA	P value
Veillonellaceae GGB4266 (SGB5809)	4.023	Indian	3.680	0.003000
*Phocaeicola plebeius (*SGB1903)	4.658	Indian	4.361	2.40E-12
Prevotella copri clade A (SGB1626)	5.034	Indian	4.735	9.25E-13
Escherichia coli (SGB10068)	4.157	Indian	3.801	4.21E-12
Megasphaera sp. (BL7 SGB5858)	4.456	Indian	4.159	2.04E-10
Dialister hominis (SGB5803)	4.147	Indian	3.838	5.42E-06
Acidaminococcus intestini (SGB5736)	3.758	Indian	3.544	0.001382
Dorea longicatena	4.390	Indo-Immigr	4.019	2.99E-16
Faecalibacterium prausnitzii (SGB15342)	4.242	Indo-Immigr	3.852	3.18E-11
Lachnospiraceae bacterium (WCA3 601 WT 6H SGB4910)	4.079	Indo-Immigr	3.685	0.009908
Dorea longicatena (SGB4581)	4.301	Indo-Immigr	3.922	2.04E-07
Faecalibacterium prausnitzii	4.930	Indo-Can	4.513	1.07E-16
Ruminococcus torques	4.410	Indo-Can	4.083	4.02E-16
Anaerobutyricum hallii	4.033	Indo-Can	3.693	3.48E-21
Collinsella aerofaciens (SGB14546)	4.478	Indo-Can	4.099	1.69E-11
Dialister invisus (SGB5825)	4.270	Indo-Can	3.963	4.37E-08
Blautia wexlerae (SGB4837)	4.657	Indo-Can	4.292	5.37E-21
Gemmiger formicilis (SGB15300)	4.422	Indo-Can	4.001	3.60E-06
Faecalibacterium prausnitzii (SGB15318)	4.436	Indo-Can	4.108	3.78E-14
Blautia massiliensis (SGB4826)	4.232	Indo-Can	3.925	1.82E-12
Anaerostipes hadrus (SGB4540)	4.189	Indo-Can	3.849	3.45E-19
Ruminococcus torques (SGB4563)	4.172	Indo-Can	3.843	1.32E-10
Faecalibacterium prausnitzii (SGB15332)	4.368	Indo-Can	4.007	2.44E-09
Anaerobutyricum hallii (SGB4532)	3.918	Indo-Can	3.583	6.69E-20
Eubacterium rectale (SGB4933)	4.656	Indo-Can	4.179	5.36E-05

Clostridia bacterium (SGB14861)		4.027	Indo-Can	3.739	7.03E-10
Ruminococcus torques (SGB4608)		4.035	Indo-Can	3.710	2.92E-13
Lachnospiraceae bacterium		4.275	Euro-Can	3.861	3.12E-16
Blautia faecis (SGB4820)		3.909	Euro-Can	3.564	3.08E-19
Phocaeicola vulgatus (SGB1814)		4.488	Euro-Can	4.056	5.66E-05
Clostridium sp (AF34 10BH SGB4914)		4.083	Euro-Can	3.654	0.007673
Ruminococcaceae GGB9635 (SGB15106)		3.871	Euro-Immigr	3.588	0.014807
Clostridia bacterium		4.522	Euro-Immigr	4.047	3.81E-08
Blautia obeum		3.917	Euro-Immigr	3.566	5.24E-15
Fusicatenibacter saccharivorans (SGB4874)		4.485	Euro-Immigr	4.154	3.84E-18
Lacrimispora celerecrescens (SGB4868)		3.960	Euro-Immigr	3.635	2.47E-08

Figure 3. BugBase Predicts Higher Pathogenic Potential and Stress-Tolerant Microbiome in Indians

A Kruskal-Wallis test was performed in BugBase followed by Mann-Whitney-Wilcoxon tests for pairwise comparisons with adjusted *P*
values shown. Significant phenotypic differences were predicted from the 16S rRNA gene sequence data. Relative abundance is
presented on the y-axis. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). **(A)** Stress
tolerance was significantly higher in Indians (p_{FDR} = 2.40e-12) compared to all cohorts. **(B)** Potentially pathogenic bacteria was
significantly higher in Indians (p_{FDR} = 2.07e-18) compared to all cohorts. Indo-Immigrants also showed significantly higher pathogenic
potential than Indo-Canadians (p_{FDR} = 5.78e-04), Euro-Canadians (p_{FDR} = 1.41e-04) and Euro-Immigrants (p_{FDR} = 2.34e-04). **(C)** Gram
negative bacteria were estimated to be significantly higher in Indians than all other cohorts (p_{FDR} = 2.01e-23). Indo-Immigrants were
predicted to have a significantly higher Gram-negative bacteria abundance than Indo-Canadians (p_{FDR} = 3.32e-06), Euro-Canadians
340 (p_{FDR} = 7.96e-10), and Euro-Immigrants (p_{FDR} = 1.81e-05). **(D)** Predictions of containing mobile elements was significantly lower in
341 Indians than all other cohorts (p_{FDR} = 1.76e-17). Mobile elements were also predicted to be significantly lower in Indians than Indo-
342 Canadians (p_{FDR} = 3.34e-05), Euro-Canadians (p_{FDR} = 1.53e-11), and Euro-Immigrants (p_{FDR} = 6.42e-06). **(E)** Aerobic bacteria was
343 significantly higher in Indians than Indo-Immigrants (p_{FDR} = 0.0149) and Indo-Canadians Immigrants (p_{FDR} = 0.0090), and Indo-
344 Immigrants were significantly lower than Euro-Canadians (p_{FDR} = 0.0023) and Euro-Immigrants (p_{FDR} = 0.0149), and Indo-Canadians
were significantly lower than Euro-Canadians (p_{FDR} = 0.0019) and Euro-Immigrants (p_{FDR} = 0.0090). **(F)** Anaerobic bacteria abundance
345 was significantly lower in Indians (p_{FDR} = 7.55e-07) compared to all cohorts. Indo-Canadians also showed significantly higher anaerobic
346 bacteria than Euro-Canadians (p_{FDR} = 0.0060). **(G)** Facultatively Anaerobic bacteria abundance was significantly more abundant in
347 Indians than all other cohorts (p_{FDR} = 1.57e-07). **(H)** No significant differences were detected in biofilm formation predictions amongst
groups. Taxonomic contributions are also displayed for each phenotype prediction. *** = p ≤ 0.001, ** = p ≤ 0.01, * = p ≤ 0.05

The highest proportion of non-meat eaters were Indians at 60% (*n* = 36/61, p = <0.0001), with

30% vegetarian and 30% pescetarian. In Indo-Immigrants, 19% (*n* = 6/32), while 12% (*n* = 2/17)

of Indo-Canadians, 10% (*n* = 4/41) Euro-Canadians and 39% (*n* = 9/23) of Euro-Immigrants were

non-meat eaters. Cooking oil usage highlighted sunflower oil, ghee and mustard oil as the top

choices of Indians, whereas Indo-Immigrants preferred olive oil, butter and ghee, with olive oil

and butter being the primary fats used by the three westernized groups (Figure S5). These results

highlight the westernized diet adopted by Indo-Canadians, including high levels of processed food

and low levels of fiber, are potential drivers in the westernized gut microbiota.

Figure 4. Microbial Metabolic Pathways Show Distinction in Functional Potential of Indian Gut Microbiome

Pathway abundances were annotated using MetaCyc, then normalized to account for gene length and sampling depth. LEfSe (Linear discriminate analysis Effect Size) was used to test for differentially abundant microbial metabolic pathways using unstratified data. A Kruskal-Wallis test was performed with a significance (a) at 0.05 for one-against-all comparisons. Discriminate features were identified using a Linear Discriminate Analysis (LDA) score, with an LDA score equal to or greater than 3.0. Using MicrobiomeAnalyst 2.0, a heatmap was generated to display the differentially abundant pathways identified from LEfSe. Each column is a sample, and the coloured bars on top represent the cohorts. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). Metabolic pathways are labelled in rows and annotated with a colour that describes their general function. Colours on the heatmap represent the relative abundances of each pathway. This heatmap displays a clear distinction in microbial metabolic pathway abundances in Indians versus cohorts living in Canada. * *L-orthinine biosynthesis I* is indirectly linked with protein synthesis and also involved with urea cycle; * *CMP-3-deoxy-D-manno-octulosonate biosynthesis* is directly involved in lipopolysaccharide (LPS) production.

Figure 5. *Prevotella copri* is the Top Taxa Contributing to CAZy Families Abundant in Indians

Normalized gene family abundance table was regrouped to CAZy database using *human_regroup_table*. Barplots for stratified taxa contributions for each CAZy gene family was generated using *human_barplot*. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). *Prevotella copri* is the top contributor to CAZyme families **(A)** GH10, GH43, GH51, all which were enriched in Indians. Top taxa contributions to antimicrobial resistance-related KEGG orthologues abundant in Indians and Indo-Immigrants. Normalized gene family abundance table was regrouped to KEGG database using *human_regroup_table*. Barplots for stratified taxa contributions for each KEGG ortholog was generated using *human_barplot*. **(B)** *Prevotella copri AM42 24* is the top contributor to K00561, enriched in Indians and *Coprococcus comes* is the top contributing taxa to K18220, enriched in Indo-Immigrants.

Table 5. Percentages of respondents who reported use of each cooking oil

Oil Type	Indian *n = 43*	Indo-Immigr *n = 18*	Indo-Can *n = 9*	Euro-Can *n = 35*	Euro-Immigr *n = 18*
Butter	5%	24%	27%	24%	26%
Ghee	12%	16%	0%	0%	0%
Coconut	9%	4%	0%	5%	10%
Olive	1%	28%	45%	44%	39%
Sunflower	30%	0%	0%	2%	0%
Canola	0%	12%	0%	13%	6%
Vegetable	5%	4%	18%	2%	10%
Mustard	11%	8%	0%	0%	0%
Avocado	0%	0%	0%	5%	6%
Groundnut	8%	0%	0%	0%	0%
Rice	8%	0%	0%	0%	0%
Other	9%	4%	9%	4%	3%

To understand what dietary factors could be driving microbial taxa, the dbRDA model,

accounting for demographics and dietary patterns elucidated 16.62% of the microbial variance on

the first 2 axes (PERMANOVA F = 5.9923, p = 0.001) (Figure 6C). After false discovery rate (FDR)

adjustment, UPF and alcohol intake were principal determinants in differentiating the microbiota

in westernized cohorts (p_{FDR} = 0.003). The pescetarian diet (p_{FDR} = 0.070) and fiber intake (p_{FDR} =

0.2268) appeared to influence clustering of Indian samples, albeit not significant. Advancing age

(p_{FDR} = 0.044) influenced another clustering, whereas vegetarianism showed no significant effect

(p_{FDR} = 0.371). Overall, these results suggest that UPF and alcohol intake were driving

westernization of the gut microbiome in Indian migrants with this effect most prominent in Indo-

Canadians. Furthermore, the higher fiber intake observed in Indians coincided with high

Prevotella spp. and CAZy gene families tailored to the more traditional Indian diet.

Given the above data indicated that the Indo-Canadian group had adopted a westernized

diet, further analysis of diet was examined including absolute macronutrient and micronutrient

intakes (Table 6-7; Table S4-S5). Total energy intake was significantly higher in Euro-Canadian

Figure 6. Significant Intake of Ultra-Processed Food and Alcohol are Drivers in the Westernized Gut Microbiota

Middle bands are the median values, top and bottom boxes display the first and third quartiles, and whiskers are the min and max values. Indian (n = 61), Indo-Immigr (n = 32), Indo-Can (n = 23), Euro-Can (n = 41), Euro-Immigr (n = 23). **(A)** Calories from Group 4 Ultra-Processed Food (UPF) was counted for each subject from the ESHA dietary reports then compared to total average daily caloric intake. A Kruskal-Wallis test was performed, followed by Dunn's multiple comparisons, with adjusted p values displayed. Percentage of daily caloric intake from Group 4 UPF was highest in Indo-Canadians and Indians consumed a significantly lower amount than all other cohorts (p = <0.0001). **(B)** Fiber intake was normalized per 1000 kcal for each subject. An ordinary one-way ANOVA was performed, followed by Tukey's multiple comparisons, with adjusted p values displayed. Indians consumed significantly more fiber than Indo-Canadians (p = <0.0001) and Euro-Canadians (p = 0.0213). Indo Canadians also consumed significantly less fiber than Indo-Immigrants (p = 0.0028) and Euro-Canadians (p = 0.0110). **(C)** Distance-based redundancy analysis (dbRDA) plot was generated using Weighted UniFrac distance matrix from 16S rRNA gene amplicon data. This dbRDA plot depicts the distribution of samples and their associations with lifestyle factors. Each dot is a sample, colour coded by cohort. Variance inflation factor values ranged between 1.14 (vegetarian) – 1.38 (pescetarian), indicating each variable uniquely contributed to the model. The explanatory variables in this model elucidated 16.62% of the microbial variation on the first two axes (F = 6.8528, p = 0.001). Permutational multivariation ANOVA (PERMANOVA) test reveals these explanatory variables collectively were significant (F = 5.9923, p = 0.001), with Group 4 UPF (pFDR = 0.003) and alcohol (pFDR = 0.003) driving distinctions in the microbiota of westernized cohorts. Distinct clustering is shown in Indians, with advancing age being a significant contribution (pFDR = 0.044).

429 **Table 6. Male participant absolute macronutrient intake**

Macronutrient	RDA/AMDR	Indian n = 37	Indo-Immigr n = 18	Indo-Can n = 6	Euro-Can n = 16	Euro-Immigr n = 10	P value
Energy, kcal (IQR)	-	2329 (1187)	2086 (933)	2274 (532)	2911 (1342)	2668 (906)	0.0343 [a]
Protein							
g (IQR)	56g	82.9 (41.3)	78.8 (39.1)	138 (37.2)	119 (62.6)	103 (42.5)	<0.0001
% of total energy	10-35%	13.1%	14.3%	23.1%	17.8%	16.4%	
g per 1000 kcal	-	31.3	35.1	37.5	40.4	37.6	
Fat							
g (IQR)	-	80.2 (43.1)	76.5 (40.8)	109 (39.9)	113 (63.9)	98.5 (45.2)	0.0109 [a]
% of total energy	20-35%	29.2%	34.6%	39.2%	38.2%	37.1%	
g per 1000 kcal	-	34.6	38.2	44.1	42.9	40.7	
PUFA, g (IQR)	-	13.1 (13.7)	10.6 (6.33)	13.7 (6.32)	22.2 (13.3)	11.7 (9.35)	0.0254 [b]
Omega-3, g (IQR)	1.6	1.44 (1.77)	0.93 (0.53)	1.24 (1.01)	2.31 (3.44)	0.85 (1.63)	0.0080
Omgea-6, g (IQR)	17	11.5 (10.9)	8.77 (4.64)	10.5 (5.93)	16.7 (10.8)	9.20 (7.15)	0.0127 [b]
MUFA, g (IQR)	-	17.8 (14.3)	17.7 (11.8)	17.2 (22.3)	33.0 (24.0)	24.6 (13.7)	0.0100 [b]
SFA, g (IQR)	-	26.4 (20.0)	23.0 (12.2)	34.5 (15.0)	37.3 (22.3)	30.1 (19.0)	0.0055 [a]
Carbohydrate							
g (IQR)	130g	336 (58.6)	267 (51.8)	207 (39.7)	268 (41.1)	284 (43.0)	<0.0001 [a]
% of total energy	45-65%	58.6%	51.8%	39.8%	41.1%	43.0%	
g per 1000 kcal	-	141	131	110	113	112	
Fiber, g (IQR)	38g	34.9 (17.4)	28.3 (20.1)	13.1 (9.70)	23.9 (15.5)	22.4 (22.6)	0.0016 [b]
Soluble fiber		0.37	0.59	0.03	0.66	0.56	0.1223

Abbreviations: [a] ANOVA, [b] Kruskal-Wallis. Median and IQR reported.

Table 7. Female participant absolute macronutrient intake

Macronutrient	RDA/AMDR	Indian *n = 24*	Indo-Immigr *n = 14*	Indo-Can *n = 11*	Euro-Can *n = 25*	Euro-Immigr *n = 13*	*P* value
Energy, kcal (IQR)	-	1747 (630)	1976 (1136)	1812 (527)	2265 (910)	2227 (1060)	0.0332
Protein							
g (IQR)	46g	51.5 (21.8)	65.3 (47.9)	63.1 (20.5)	81.0 (20.8)	66.6 (50.7)	0.0004 [b]
% of total energy	10-35%	12.3%	13.8%	12.9%	14.1%	14.0%	
g per 1000 kcal	-	31.3	35.1	37.5	40.4	37.6	
Fat							
g (IQR)	-	62.7 (32.3)	69.8 (38.0)	73.1 (27.5)	89.2 (40.4)	103 (57.0)	0.0018 [b]
% of total energy	20-35%	33.4%	34.4%	39.7%	38.7%	36.7%	
g per 1000 kcal	-	34.6	38.2	44.1	42.9	40.7	
PUFA, g (IQR)	-	12.1 (9.49)	12.9 (11.0)	8.06 (4.33)	13.3 (8.90)	9.87 (13.0)	ns
Omega-3, g (IQR)	1.1	1.24 (0.85)	1.46 (1.02)	0.84 (0.44)	1.42 (1.25)	1.06 (1.16)	ns
Omgea-6, g (IQR)	12	10.5 (8.72)	8.80 (8.25)	7.00 (4.04)	10.0 (8.92)	8.73 (7.67)	ns
MUFA, g (IQR)	-	16.5 (10.5)	15.6 (13.4)	15.3 (9.77)	23.8 (13.7)	19.0 (23.6)	ns
SFA, g (IQR)	-	18.3 (14.9)	23.4 (10.6)	22.9 (5.66)	30.6 (14.3)	41.7 (35.1)	ns
Carbohydrate							
g (IQR)	130g	258 (86.4)	256 (116)	218 (99.5)	272 (119)	248 (141)	ns
% of total energy	45-65%	55.7%	53.6%	38.4%	47.6%	49.4%	
g per 1000 kcal	-	141	131	110	113	112	
Fiber, g (IQR)	25g	24.6 (17.8)	23.1 (18.7)	14.7 (6.30)	27.0 (15.2)	28.8 (28.4)	0.0163 [a]
Soluble fiber		0.21	0.69	0.05	0.75	0.45	0.0003 [b]

Abbreviations: [a]ANOVA, [b] Kruskal-Wallis. Median and IQR reported.
*Tukey's post hoc analysis revealed no significant differences between groups

males vs. Indo-Immigrant males (p = 0.0144, 95% CI = -1461 – -110.0 kcal). Median percentage

of protein intake was highest in Euro-Canadians (16.1%, 95% CI = 13.2 – 17.5%) and lowest

protein intake was in Indians (12.5%, 95% CI = 11.7 – 13.3). Indians consumed the highest

percentage of carbohydrates in their diet (56.5%, 95% CI = 54.7% - 59.0%) and lowest

carbohydrate diet intake was in Indo-Canadians (44.0%, 95% CI = 40.6% – 50.6%). The highest

percentage of fat consumption was in Indo-Canadians (39.7%, CI 95% 33.8% – 42.2%) and the

lowest was in Indians (31.1%, 95% CI = 28.5% – 34.4%). Software comparisons between North

American ESHA and Indian EpiNu revealed no macronutrient composition differences however,

EpiNu calculated a significantly higher intake of PUFAs and MUFAs and lower intake of SFA

amongst Indians, highlighting discrepancy in dietary fat types between dietary software, which

could skew dietary analysis between the countries (Fig 7A-B). When the type of oils consumed

were examined, it was revealed that Indo-Canadians have the least diversity with no traditional

fats added to their diet like ghee (Fig 7C).

Figure 7. Dietary Fat Types are Significantly Different Between EpiNu vs. ESHA

(A) Percentages of fat, protein and carbohydrates were calculated in ESHA and EpiNu for each participant, then mean values were calculated for each cohort. An ordinary one-way Analysis of Variance (ANOVA) was used, and no significant differences in macronutrient intake were

found between ESHA and EpiNu. **(B)** Significant differences in types of fat consumed by Indians (n = 61) and Indo-Immigrants (n = 32) were calculated between EpiNu vs. ESHA. Dietary fats displayed in grams per 1000 Calories (kcal). Kruskal-Wallis tests were conducted to determine differences between groups. The middle band inside of the boxes is the median, the bottom and top boxes are the first and third

quartiles, and the whiskers are the min to max values. When compared to EpiNu, ESHA calculated a significantly higher saturated fatty acid (SFA) intake in Indians (p = <0.0001). EpiNu calculated significantly higher monounsaturated fatty acid (MUFA) intake in both Indians and Indo-Immigrants (p = <0.0001). Compared to ESHA, EpiNu calculated a significantly higher intake of polyunsaturated fatty acid (PUFA) in

Indians.

DISCUSSION

Westernization is Linked to Both Taxonomic and Functional Transitions in the Gut

Microbiomes of Indian Migrants

Earlier studies identified that the Indian gut microbiota is distinct, as it is rich in *Prevotella*

spp*., Dialister* spp*.,* and *Megasphaera* spp.,[24-30] unlike westernized groups dominated by

Bacteroides spp.[31] and *Phocaeicola vulgatus*[32]. Furthermore, the high taxonomic variation in

Indians may in part reflect their diverse dietary practices in India alongside the current western

influence. This heterogeneity also highlights how attempting to generalize the "Indian microbiome"

may inaccurately represent this population due to the great cultural variations that exist within the

country. Contrary to associations of low alpha diversity in westernization and disease, healthy

Indians showed lower alpha diversity than westernized cohorts, suggesting this diversity measure

may not always be a reliable marker of westernization or health status.[33]

Taxonomic data revealed high abundances of VANISH taxa in Indians and Indo-

Immigrants, and BloSSUM taxa enriched in European westernized controls. Indo-Canadians

exhibited a transitional microbiota containing both VANISH and BloSSUM taxa, a trend previously

detected in infants living in transitional societies (intermediate between non-industrialized and

industrialized lifestyles).[34] A key genus that is distinctive in this transition in immigrants is the loss

of *Prevotella* spp.[8,35] In fact, this contrast in *Prevotella* spp. abundance in non-westernized versus

westernized groups has sparked interest both in its relation to lifestyle and disease, previously

being linked to IBD,[36] [37] rheumatoid arthritis,[38] and obesity[39]. Yet, contradictory studies also

showed its correlation with reduced risks of cardiovascular disease (CVD) [40] diabetes [41], and

obesity,[42] illustrating the current contradictory understanding around *Prevotella* spp.-dominance

in the gut and its role in diseases.

Globally, Indians have some of the highest abundances of *Prevotella* spp. in their gut,[43]

as was reflected in our data, which highlights the importance of context when investigating the

role of microbes in our gut, as host-microbe interactions may divergently evolve depending on the

host's genetics, geography, and lifestyle. In fact, these factors may have already driven

evolutionary changes in *Prevotella* spp. strains: isolated *P. copri* strains from Indian/non-western

populations showed traits that aid in digestion of plant-based carbohydrates using carbohydrate-

active enzymes (CAZymes), whereas *P. copri* strains from westernized populations were enriched

with virulence factors and resistance genes.[43] In our study, Indians had a significantly higher

average abundance of *P. copri* Clade A, along with notable abundances of Clades B and C, which

subsequently declined in Indo-Immigrants and Indo-Canadians (Figure 2D). All *P. copri* clades

have previously been documented at higher abundances in non-westernized individuals than

westernized,[44] and our results highlight this effect of westernization on clade abundances using

groups of the same ethnicity. These findings demonstrate how specific strains can evolve with its

host, cautioning against broad associations between bacterial species and diseases.

Indians, known for their high-complex-carbohydrate diet, have been previously indicated

to exhibit an increased expression of carbohydrate metabolism genes for complex

polysaccharides in their gut.[24,28,30] Our findings reflected this relationship, as we detected

enrichment of CAZy families that aid in the digestion of xylan and xyloglucan, with *P. copri* as the

top contributing taxa. A similar pattern was in fact detected in another study on immigrants who

migrated from Thailand to the US, where a decrease in *Prevotella* spp. abundance corresponded

with a loss of CAZymes that break down plant fiber.[8] The subsequent loss of *P.copri* in our Indo-

Immigrant and Indo-Canadian cohorts aligned with their reduction of a high carbohydrate, high

fiber diet, which was a trend previously explored among Hadza and Nepali groups.[45] Plant-derived

microbiota-accessible carbohydrates (MACs) are required to maintain abundances of *P. copri*,

whereas other species such as *Bacteroides thetaiotaomicron* can utilize both plant and animal-

derived MACs, allowing for persistence in the gut.[45] This relationship therefore suggests that the

Prevotella spp. abundance in the Indian gut is highly linked to their specific dietary patterns.

Furthermore, the generational loss of *P. copri* in Indian immigrants prompts questions about gut

health implications, meriting further research on strain-specific *P. copri*-host interactions within

this demographic.

A previous study found *Dialister succinatiphilus* and *Megasphaera* more abundant in Indo-

Immigrants versus Indo-Canadians, who had significantly higher *Dialister invisus.* Our data align

with this trend, with *D. invisus* enriched in Indo-Immigrants, *D. succinatiphilus* in Indo-Canadians,

and *Megasphaera* in our Indian cohort. While little is known about these taxa, a relationship

between a carbohydrate-rich diet and succinate-utilizing bacteria has been previously

discussed,[46] with *D. succinatiphilus* being a well-known species in this category. Additionally,

Ruminococcus torques, a known mucin degrader,[47] previously linked to IBD risk [48] and identified

as a significant contribution in a Crohn's Disease microbiome-risk score,[49] was notably enriched

in Indo-Canadians. Deeper investigations should explore the role of specific *R. torques* strains in

the gut and its implications in IBD should be explored further.

India is Currently Industrializing

India's industrialization is reshaping dietary habits, as seen in our data where sunflower

oil, contradictory to traditional fats like ghee, has become predominately used due to

globalization[50] and dietary guidelines recommending a reduction in saturated fatty acids and

increase in PUFA-rich vegetable oils. Overtime, these vegetable oils became inexpensive and

more readily available in India, therefore there has been a transition away from the use of their

traditional fats.[51] Coinciding with this transition of increased PUFA consumption is the rising

epidemic of several modern diseases such as IBD[52], type-2 diabetes, and metabolic syndrome

(MS).[53] India is now in an accelerating incidence phase of IBD[1], doubling the number of patients

from 1990 to 2019,[54] and the increased intake of linolenic (omega-6 PUFA) in Indians has been

previously associated with an increased risk for UC.[55,56] These findings highlight the need to

investigate the role of vegetable oil consumption in India and its potential contribution to the rising

incidence of IBD.

Surprisingly, the westernized cohorts in our study reported cooking mainly with olive oil and butter,

and few reported to cook with vegetable oils unlike their expected intake of oils rich in omega-6

PUFAs.[57] Though these reports are limiting it is worth highlighting that while India has adopted

the use of white oils like sunflower oil and canola oil, North America may currently be shifting

away from them. For decades, the westernized diet has been characterized by a much higher

intake of omega-6 PUFA, leading to omega-6:omega-3 ratios of 20:1, which is far from the

estimated ratio of 1:1 of our ancestors.[58] This excessive intake of omega-6 PUFAs has been

associated with CVD,[59] cancer,[60] and autoimmune diseases.[61] Furthermore, with large-scale

studies and meta-analyses refuting the link between saturated fatty acids and CVD[62,63] and the

increasing popularity of the Mediterranean Diet,[64] a cultural shift in the North America may be

occurring, once again preferring olive oil and butter,[65] which was an observation also recently

reported from an Australian cohort.[66]

Another factor of industrialization that is common in India is the overuse of antibiotics.

India has one of the highest threats for antimicrobial resistance,[67] as was reflective in our cohorts

with enriched KEGG identities in Indians and Indo-Immigrants that contain signatures for antibiotic

resistance genes, a pattern previously documented in the Indian gut[24 68] The widespread use of

antibiotics in India not only increases the prevalence of antibiotic resistance, but also accelerates

microbiome turnover, which was indicative in the pathways enriched in our Indian cohort including

peptidoglycan synthesis, microbial growth/metabolism, and DNA synthesis/repair. While common

usage of antibiotics is a likely catalyst for cell turnover, the higher risk for infectious disease via

water contamination and inadequate sanitation in India may also serve as environmental triggers.

Our findings highlight India's unique transition phase, balancing Western influences with

traditional lifestyle practices, all of which are shaping their gut microbiome. Future research should

delve deeper into how westernization is currently influencing the gut microbiome in Indians, as

this will likely pose future health challenges.

Acculturation of Westernized Diet Drives Functional Shifts in Microbiome

590   Coinciding with previous findings, we observed dietary acculturation from the traditional

high-complex-carbohydrate Indian diet to an increased intake of NOVA Group 4 UPFs,[35,69],[70]

which are characterized by industrially made ingredients with non-nutritive additives like

emulsifiers, flavouring and sweeteners.[71] This dietary shift might explain the enrichment of

pathways tapping into glycogen storage in Indo-Canadians, suggesting microbes had inadequate

nutrient availability from the host diet.[72] With a 49% rise in Group 4 UPF intake and significant

fiber reduction in Indo-Canadians, concerns emerge regarding their gut health and chronic

disease risks.

599   Increasing UPF consumption has been linked to higher risks for CVD, metabolic disorders,

micronutrient deficiencies, and anxiety and depression,[71] which may partly be due to imbalances

in the gut microbiome that are a triggered from UPF. While there are several potential culprits

involved in the ultra-processing of foods that may have mechanistic interactions with the gut, it is

likely an interplay of several factors surrounding the consumption of UPFs. First, despite being

affordable and accessible, UPFs are often high in calories and palatability, but low in essential

nutrients, leading to overconsumption and displacement of nutritious meals.[73] Additionally,

emulsifiers, used to mix components in UPF, have been extensively studied in murine models,

showing their ability to modulate the gut microbiota, reduce mucus thickness and increase

intestinal permeability, bacterial translocation and inflammation.[74] While studies translating these

same mechanistic effects of emulsifiers in humans are scarce, a short-term prospective study (N

= 588) found a positive association between dietary emulsifiers and the inflammatory biomarker

glycoprotein acetyls.[75] Moreover, a randomized double-blind controlled-feeding study (N = 16)

with the emulsifier carboxymethylcellulose (CMC) found varied responses in the treatment group,

with two subjects exhibiting significant alterations in gut composition and microbial encroachment,

whereas other participants seemed non-sensitive to CMC.[76,77] Although constrained by a small

sample size, these findings prompt the need to further investigate in large-scale cohorts, as this

may in part explain the inconsistent outcomes observed when transitioning from murine to human

studies. Overall, these findings underscore that emulsifiers may be one of many components in

UPF that may affect the gut microbiome, however more human studies are required to fully

understand the complex role of UPF and its effects on the gut microbiome and inflammation.

Future studies should also consider examining the synergistic effects of multiple food additives to

mirror the composition of UPFs typically consumed by humans.

LIMITATIONS

In addition to inherit limitations of self-reported food diaries (e.g. accuracy of the subject's

record), our study also faced limitations in accurately representing specific cultural diets, primarily

due to inadequacies in existing databases for Indian cuisine. This led to the use of EpiNu

Nutritional software, better tailored for Indian dietary data. Additionally, the shallow depth of

shotgun sequencing and taxonomic classification methods used restricted our ability to identify

strain-level differences in bacteria like *P. copri*. Another limitation worth mentioning is the uneven

sample sizing across groups. It was considerably more difficult to recruit Indo-Canadians for our

study, which we believe may have been due to a combination of cultural, social and geographical

barriers.

As this project focused on the relative abundance of *Prevotella* spp., quantitative PCR

could also help to determine absolute abundances present in the gut, as this may also be a key

factor influencing disease risk [78]. Additionally, while metagenomics can provide important insights

into the genetic content present within the gut, these data only represent the functional potential

of the gut and does not provide us information regarding genetic expression and actual function

of microbiome communities. Overall, future work should aim to resolve strain-level analysis and

parasite communities in the microbiomes of this population along with deeper exploration into

absolute abundances of taxa and the functional activity present to better understand their

increased risk to IBD. Lastly, this study merely provides associative data between westernization

and gut microbiome changes; further investigation is needed to understand the mechanistic

interactions of specific factors of westernization on the gut and IBD risk.

Figure 8. Adoption of Westernized Diet is Associated with a Transition Away from Indian Gut Microbiome

Graphical representation highlighting the main findings from our study. As Indo-Immigrants and Indo-Canadians subsequently increased ultra-processed food intake, their fiber consumption dramatically reduced. This change in dietary pattern was observed alongside a transition away from high *Prevotella* spp. and VANISH (Volatile and/or Associated Negatively with Industrialized Societies of Humans) taxa, which were found abundant in the Indian gut. Instead, Indo-Canadians adopted BloSSUM (Bloom or Selected in Societies of Urbanization/Modernization) taxa, which are commonly found in the westernized microbiome. High *Prevotella copri* in the Indian gut contributed to enriched carbohydrate-active enzymes (CAZymes) that are specialized to degrade complex carbohydrates common in their diet, which decline in Indian migrant cohorts. Additionally, the Indian microbiome displayed characteristics of a more robust gut, with predictive functions of higher stress tolerance and increased microbial cell turnover.

CONCLUSIONS

Consistent with previous studies, our research reveals that the Indian gut microbiome, rich in *Prevotella* spp., is adapted to their high carbohydrate, high fiber diet. We discovered that the Indian microbiome exhibits characteristics of higher bacterial cell turnover, pathogenic potential and stress tolerance, indicating its robustness against stressors posed on the gut (Figure 8). These findings highlight the importance in understanding diverse gut microbiomes, particularly in non-westernized and transitional populations, for betting predicting and managing diseases like IBD. Our study sheds light on the gut microbiome changes in a population undergoing westernization, revealing a drastic transition within the gut microbiomes of Indo-Immigrants and Indo-Canadians, extending our knowledge of the strong effect of immigration and westernization on the gut microbiome. This transition likely reflects microbial adaptation to the dietary acculturation observed, providing insight into the dynamic nature within our gut microbiome. Future investigations should examine the health effects of microbiome changes amidst global westernization, as this transition will likely redefine health outcomes for both migrating populations and those living in rapidly modernizing nations.

Acknowledgements

We thank all the individuals who took part in this study. We thank the entire Ghosh family for support. For dietary analysis, we thank Nadia Anvari. For lab support, we thank Mekenna Smith, Hephzibah Bomide, Carson McComb, Ayva Lewis, Erik Kaila, Andrea Verdugo Meza and Jessica Josephson.

Funding

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and a UBC Killam Research Award to support D.L.G. on her sabbatical in India.

Competing Interests

DLG is the co-founder and CSO of Melius MicroBiomics Inc. where LDD is the Science

Communications Specialist.

Patient and Public Involvement

Not applicable.

Ethics Approval

This study was approved by the University of British Columbia Clinical Research Ethics Board

(UBC CREB) (H21-01555). Approval for sample collection in India was approved under IEC No

411/2018 and by UBC CREB (H17-01324). All participants signed a consent form prior to

participation. A Southeast Asian (SEA) Microbiome Biobank was created.

Data Availability Statement

Data supporting this study's findings are available within the paper and its Supplementary

Information. The 16S rRNA gene sequences files and metagenomic sequence files of human

fecal samples are publicly available on NCBI under accession number PRJNA1082632. Scripts

used for the analysis of this study will be made available upon publication (upload pending).

Supplementary Material

Please find Supplementary Methods and Supplementary Results attached.

REFERENCES

1. Kaplan GG, Windsor JW. The four epidemiological stages in the global evolution of
inflammatory bowel disease. *Nature Reviews Gastroenterology & Hepatology.*
2021;18(1):56-66.
2. Benchimol E, Mack D, Guttmann A, et al. Inflammatory bowel disease in immigrants to
Canada and their children: a population-based cohort study. *The american journal of*
gastroenterology. 2015;110(4):553-563.

3.   Carroll MW, Hamilton Z, Gill H, et al. Pediatric inflammatory bowel disease among South
Asians living in British Columbia, Canada: a distinct clinical phenotype. *Inflamm Bowel*
Dis. 2016;22(2):387-396.
4.   Pinsk V, Lemberg DA, Grewal K, Barker CC, Schreiber RA, Jacobson K. Inflammatory
bowel disease in the South Asian pediatric population of British Columbia. *Am J*
Gastroenterol. 2007;102(5):1077-1083.
5.   Fragiadakis GK, Smits SA, Sonnenburg ED, et al. Links between environment, diet, and
the hunter-gatherer microbiome. *Gut Microbes.* 2019;10(2):216-227.
6.   Sonnenburg ED, Sonnenburg JL. The ancestral and industrialized gut microbiota and
implications for human health. *Nature Reviews Microbiology.* 2019;17(6):383-390.
7.   Parizadeh M, Arrieta M-C. The global human gut microbiome: genes, lifestyles, and diet.
Trends Mol Med. 2023;29(10):789-801.
8.   Vangay P, Johnson AJ, Ward TL, et al. US immigration westernizes the human gut
microbiome. *Cell.* 2018;175(4):962-972. e910.
9.   Abdill RJ, Adamowicz EM, Blekhman R. Public human microbiome data are dominated
by highly developed countries. *PLoS Biol.* 2022;20(2):e3001536.
10.   Gorzelak MA, Gill SK, Tasnim N, Ahmadi-Vand Z, Jay M, Gibson DL. Methods for
improving human gut microbiome data by reducing variability through sample
processing and storage of stool. *PLoS One.* 2015;10(8):e0134802.
11.   Bolyen E, Rideout JR, Dillon MR, et al. Reproducible, interactive, scalable and extensible
microbiome data science using QIIME 2. *Nat Biotechnol.* 2019;37(8):852-857.
12.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nature Methods.*
2016;13(7):581-583.
13.   McDonald D, Jiang Y, Balaban M, et al. Greengenes2 enables a shared data universe for
microbiome studies. *bioRxiv.* 2022:2022.2012. 2019.520774.
14.   Blanco-Míguez A, Beghini F, Cumbo F, et al. Extending and improving metagenomic
taxonomic profiling with uncharacterized species using MetaPhlAn 4. *Nat Biotechnol.*
2023:1-12.
15.   Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation.
Genome Biol. 2011;12:1-18.
16.   Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J. MicrobiomeAnalyst 2.0: comprehensive
statistical, functional and integrative analysis of microbiome data. *Nucleic Acids Res.*
2023;51(W1):W310–W318.
17.   Ward T, Larson J, Meulemans J, et al. BugBase predicts organism-level microbiome
phenotypes. *BioRxiv.* 2017:133462.
18.   Langille MG, Zaneveld J, Caporaso JG, et al. Predictive functional profiling of microbial
communities using 16S rRNA marker gene sequences. *Nat Biotechnol.* 2013;31(9):814-
821.
19.   Beghini F, McIver LJ, Blanco-Míguez A, et al. Integrating taxonomic, functional, and
strain-level profiling of diverse microbial communities with bioBakery 3. *Elife.*
2021;10:e65088.
20.   Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef: comprehensive and non-
redundant UniProt reference clusters. *Bioinformatics.* 2007;23(10):1282-1288.

21.   Caspi R, Altman T, Billington R, et al. The MetaCyc database of metabolic pathways and
779   enzymes and the BioCyc collection of Pathway/Genome Databases. *Nucleic Acids Res.*
780   2014;42(D1):D459-D471.
781   22.   Sonnenburg JL, Sonnenburg ED. Vulnerability of the industrialized microbiota. *Science.*
782   2019;366(6464).
783   23.   Jha AR, Davenport ER, Gautam Y, et al. Gut microbiome transition across a lifestyle
784   gradient in Himalaya. *PLoS Biol.* 2018;16(11):e2005396.
785   24.   Bhute S, Pande P, Shetty SA, et al. Molecular Characterization and Meta-Analysis of Gut
786   Microbial Communities Illustrate Enrichment of Prevotella and Megasphaera in Indian
787   Subjects. *Front Microbiol.* 2016;7:660.
788   25.   Kumbhare SV, Kumar H, Chowdhury SP, et al. A cross-sectional comparative study of gut
789   bacterial community of Indian and Finnish children. *Sci Rep.* 2017;7(1):10555.
790   26.   Jain A, Li XH, Chen WN. Similarities and differences in gut microbiome composition
791   correlate with dietary patterns of Indian and Chinese adults. *AMB Express.*
792   2018;8(1):104.
793   27.   Tandon D, Haque MM, R S, et al. A snapshot of gut microbiota of an adult urban
794   population from Western region of India. *PLoS One.* 2018;13(4):e0195643.
795   28.   Dhakan DB, Maji A, Sharma AK, et al. The unique composition of Indian gut microbiome,
796   gene catalogue, and associated fecal metabolome deciphered using multi-omics
797   approaches. *GigaScience.* 2019;8(3):01.
798   29.   Arumugam M, Raes J, Pelletier E, et al. Enterotypes of the human gut microbiome.
799   *Nature.* 2011;473(7346):174-180.
800   30.   Kulkarni A, Kumbhare SV, Dhotre DP, Shouche YS. Mining the Core Gut Microbiome
801   from a Sample Indian Population. *Indian J Microbiol.* 2019;59(1):90-95.
802   31.   Yatsunenko T, Rey FE, Manary MJ, et al. Human gut microbiome viewed across age and
803   geography. *Nature.* 2012;486(7402):222-227.
804   32.   Bodykevich G, Ibtisam A, Mao L. The gut microbiome belonging to Colombian adults of
805   younger generations has shifted to a more westernized composition compared to older
806   generations. *Undergraduate Journal of Experimental Microbiology and Immunology.*
807   2023;9.
808   33.   Fanfan D, Mulligan CJ, Groer M, et al. The intersection of social determinants of health,
809   the microbiome, and health outcomes in immigrants: A scoping review. *American*
810   *Journal of Biological Anthropology.* 2023;183(1):3-19.
811   34.   Olm MR, Dahan D, Carter MM, et al. Robust variation in infant gut microbiome assembly
812   across a spectrum of lifestyles. *Science.* 2022;376(6598):1220-1223.
813   35.   Copeland JK, Chao G, Vanderhout S, et al. The Impact of Migration on the Gut
814   Metagenome of South Asian Canadians. *Gut microbes.* 2021;13(1):1-29.
815   36.   Hertz S, Durack J, Kirk KF, et al. Microscopic colitis patients possess a perturbed and
816   inflammatory gut microbiota. *Dig Dis Sci.* 2022;67(6):2433-2443.
817   37.   Iljazovic A, Roy U, Gálvez EJ, et al. Perturbation of the gut microbiome by Prevotella spp.
818   enhances host susceptibility to mucosal inflammation. *Mucosal Immunol.* 2020:1-12.
819   38.   Scher JU, Sczesnak A, Longman RS, et al. Expansion of intestinal Prevotella copri
820   correlates with enhanced susceptibility to arthritis. *elife.* 2013;2:e01202.

821   39.   Moreno-Indias I, Sánchez-Alcoholado L, García-Fuentes E, Cardona F, Queipo-Ortuño MI,
822   Tinahones FJ. Insulin resistance is associated with specific gut microbiota in appendix
823   samples from morbidly obese patients. *American Journal of Translational Research.*
824   2016;8(12):5672.

825   40.   Wang Y, Ames NP, Tun HM, Tosh SM, Jones PJ, Khafipour E. High molecular weight
826   barley β-glucan alters gut microbiota toward reduced cardiovascular disease risk. *Front*
827   *Microbiol.* 2016;7:129.

828   41.   Bibbò S, Dore MP, Pes GM, Delitala G, Delitala AP. Is there a role for gut microbiota in
829   type 1 diabetes pathogenesis? *Ann Med.* 2017;49(1):11-22.

830   42.   Borgo F, Verduci E, Riva A, et al. Relative abundance in bacterial and fungal gut microbes
831   in obese children: a case control study. *Childhood Obesity.* 2017;13(1):78-84.

832   43.   Prasoodanan PK V, Sharma AK, Mahajan S, et al. Western and non-western gut
833   microbiomes reveal new roles of Prevotella in carbohydrate metabolism and mouth–gut
834   axis. *NPJ Biofilms and Microbiomes.* 2021;7(1):1-17.

835   44.   Tett A, Huang KD, Asnicar F, et al. The Prevotella copri complex comprises four distinct
836   clades underrepresented in westernized populations. *Cell Host Microbe.* 2019;26(5):666-
837   679. e667.

838   45.   Gellman RH, Olm MR, Terrapon N, et al. Hadza Prevotella Require Diet-derived
839   Microbiota Accessible Carbohydrates to Persist in Mice. *bioRxiv.* 2023:2023.2003.
840   2008.531063.

841   46.   Nakayama J, Yamamoto A, Palermo-Conde LA, et al. Impact of westernized diet on gut
842   microbiota in children on Leyte island. *Front Microbiol.* 2017;8:197.

843   47.   Png CW, Lindén SK, Gilshenan KS, et al. Mucolytic bacteria with increased prevalence in
844   IBD mucosa augmentin vitroutilization of mucin by other bacteria. *Official Journal of the*
845   *American College of Gastroenterology.* 2010;105(11):2420-2428.

846   48.   Ali F, Reddy V, Raza M, et al. S3194 Metformin: A Therapeutic Adjunct for Patients With
847   Inflammatory Bowel Disease? *Official journal of the American College of*
848   *Gastroenterology.* 2020;115:S1676.

849   49.   Garay JAR, Turpin W, Lee S-H, et al. Gut microbiome composition is associated with
850   future onset of Crohn's Disease in healthy first-degree relatives. *Gastroenterology.*
851   2023;165(3):670-681.

852   50.   USDA. The Role of Policy and Industry Structure in India's Oilseed Markets. *USDA*
853   *Economic Research Service.* 2007.

854   51.   Mani I, Kurpad AV. Fats & fatty acids in Indian diets: Time for serious introspection. *The*
855   *Indian Journal of Medical Research.* 2016;144(4):507.

856   52.   Snell A, Segal J, Limdi J, Banerjee R. Inflammatory bowel disease in India: challenges and
857   opportunities. *Frontline Gastroenterol.* 2020;12(5):390-396.

858   53.   Lakshmipriya N, Gayathri R, Praseena K, et al. Type of vegetable oils used in cooking and
859   risk of metabolic syndrome among Asian Indians. *Int J Food Sci Nutr.* 2013;64(2):131-
860   139.

861   54.   Dharni K, Singh A, Sharma S, et al. Trends of inflammatory bowel disease from the
862   Global Burden of Disease Study (1990-2019). *Indian J Gastroenterol.* 2023:1-11.

55. Investigators IiES. Linoleic acid, a dietary n-6 polyunsaturated fatty acid, and the
aetiology of ulcerative colitis: a nested case–control study within a European
prospective cohort study. *Gut.* 2009;58(12):1606-1611.
56. Ananthakrishnan AN, Khalili H, Konijeti GG, et al. Long-term intake of dietary fat and risk
of ulcerative colitis and Crohn's disease. *Gut.* 2014;63(5):776-784.
57. Mariamenatu AH, Abdu EM. Overconsumption of omega-6 polyunsaturated fatty acids
(PUFAs) versus deficiency of omega-3 PUFAs in modern-day diets: the disturbing factor
for their "balanced antagonistic metabolic functions" in the human body. *Journal of*
lipids. 2021;2021:1-15.
58. Simopoulos AP. An increase in the omega-6/omega-3 fatty acid ratio increases the risk
for obesity. *Nutrients.* 2016;8(3):128.
59. Simopoulos AP. Omega-6/omega-3 essential fatty acid ratio and chronic diseases. *Food*
reviews international. 2004;20(1):77-90.
60. De Lorgeril M, Salen P. New insights into the health effects of dietary saturated and
omega-6 and omega-3 polyunsaturated fatty acids. *BMC Med.* 2012;10(1):1-5.
61. Simopoulos AP. Evolutionary aspects of diet, the omega-6/omega-3 ratio and genetic
variation: nutritional implications for chronic diseases. *Biomed Pharmacother.*
2006;60(9):502-507.
62. Dehghan M, Mente A, Zhang X, et al. Associations of fats and carbohydrate intake with
cardiovascular disease and mortality in 18 countries from five continents (PURE): a
prospective cohort study. *The Lancet.* 2017;390(10107):2050-2062.
63. Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Meta-analysis of prospective cohort studies
evaluating the association of saturated fat with cardiovascular disease. *The American*
Journal of Clinical Nutrition. 2010;91(3):535-546.
64. Vetrani C, Piscitelli P, Muscogiuri G, et al. "Planeterranea": An attempt to broaden the
beneficial effects of the Mediterranean diet worldwide. *Frontiers in Nutrition.*
2022;9:973757.
65. Fernandes J, Fialho M, Santos R, et al. Is olive oil good for you? A systematic review and
meta-analysis on anti-inflammatory benefits from regular dietary intake. *Nutrition.*
2020;69:110559.
66. Wilson NA, Villani A, Mantzioris E. Preferences for dietary oils and fats in cooking and
food preparation methods: a cross-sectional analysis of Australian adults. *Proc Nutr Soc.*
2023;82(OCE2):E72.
67. Gandra S, Joshi J, Trett A, Lamkang AS, Laxminarayan R. Scoping report on antimicrobial
resistance in India. *Center for Disease Dynamics, Economics & Policy: Washington, DC,*
USA. 2017;2017:1-146.
68. Gupta M, Didwal G, Bansal S, et al. Antibiotic-resistant Enterobacteriaceae in healthy gut
flora: A report from north Indian semiurban community. *Indian J Med Res.*
2019;149(2):276-280.
69. Lesser IA, Gasevic D, Lear SA. The association between acculturation and dietary
patterns of South Asian immigrants. *PLoS One.* 2014;9(2):e88495.
70. Noor S, Dehghan M, Lear SA, et al. Relationship between diet and acculturation among
South Asian children living in Canada. *Appetite.* 2020;147:104524.

71. Tristan Asensi M, Napoletano A, Sofi F, Dinu M. Low-Grade Inflammation and Ultra-
Processed Foods Consumption: A Review. *Nutrients.* 2023;15(6):1546.
72. Esteban-Torres M, Ruiz L, Rossini V, Nally K, van Sinderen D. Intracellular glycogen
accumulation by human gut commensals as a niche adaptation trait. *Gut Microbes.*
2023;15(1):2235067.
73. Dinu M, Tristan Asensi M, Pagliai G, et al. Consumption of ultra-processed foods is
inversely associated with adherence to the Mediterranean diet: a cross-sectional study.
Nutrients. 2022;14(10):2073.
74. Bancil AS, Sandall AM, Rossi M, Chassaing B, Lindsay JO, Whelan K. Food additive
emulsifiers and their impact on gut microbiome, permeability, and inflammation:
mechanistic insights in inflammatory bowel disease. *Journal of Crohn's and Colitis.*
2021;15(6):1068-1079.
75. Um CY, Hodge RA, Tran HQ, Campbell PT, Gewirtz AT, McCullough ML. Association of
emulsifier and highly processed food intake with circulating markers of intestinal
permeability and inflammation in the cancer prevention study-3 diet assessment sub-
study. *Nutr Cancer.* 2022;74(5):1701-1711.
76. Chassaing B, Compher C, Bonhomme B, et al. Randomized controlled-feeding study of
dietary emulsifier carboxymethylcellulose reveals detrimental impacts on the gut
microbiota and metabolome. *Gastroenterology.* 2022;162(3):743-756.
77. Daniel N, Wu GD, Walters W, et al. Human intestinal microbiome determines
individualized inflammatory response to dietary emulsifier carboxymethylcellulose
consumption. *Cell Mol Gastroenterol Hepatol.* 2024;17(2):315-318.
78. Vandeputte D, Kathagen G, D'hoe K, et al. Quantitative microbiome profiling links gut
community variation to microbial load. *Nature.* 2017;551(7681):507-511.
930