1 Enhanced Ca^{2+}-Driven Arrhythmias in Female Patients with Atrial Fibrillation: Insights from

2 Computational Modeling

Xianwei Zhang[1*], Yixuan Wu[1*], Charlotte Smith[1], William E. Louch[2], Stefano Morotti[1], Dobromir

Dobrev[3,4,5], Eleonora Grandi[1#], Haibo Ni[1#]

1. Department of Pharmacology, University of California Davis, Davis, CA, USA

2. Institute for Experimental Medical Research, Oslo University Hospital and University of Oslo,

Norway

3. Institute of Pharmacology, West German Heart and Vascular Center, University Duisburg-

Essen, Essen, Germany.

4. Montréal Heart Institute, Université de Montréal, Montréal, Québec, Canada.

5. Department of Integrative Physiology, Baylor College of Medicine, Houston, TX, USA.

*Contributed Equally to the work

#Shared senior authorship

Running title: Modeling sex differences in AF mechanisms and therapy

Correspondence:

Eleonora Grandi, Department of Pharmacology, University of California Davis
451 Health Science Drive, GBSF 3503, Davis, CA 95616, USA
Email: ele.grandi@gmail.com
Phone: 530-752-4780
25
26 Haibo Ni, Department of Pharmacology, University of California Davis
27 451 Health Science Drive, GBSF 3503, Davis, CA 95616, USA
28 Email: hbni@ucdavis.edu
29 Phone: 530-752-4780

Abstract

Background and Aims. Substantial sex-based differences have been reported in atrial fibrillation (AF), with female patients experiencing worse symptoms, increased complications from drug side effects or ablation, and elevated risk of AF-related stroke and mortality. Recent studies revealed sex-specific alterations in AF-associated Ca^{2+} dysregulation, whereby female cardiomyocytes more frequently exhibit potentially proarrhythmic Ca^{2+}-driven instabilities compared to male cardiomyocytes. In this study, we aim to gain a mechanistic understanding of the Ca^{2+}-handling disturbances and Ca^{2+}-driven arrhythmogenic events in males vs females and establish their responses to Ca^{2+}-targeted interventions.

Methods and Results. We incorporated known sex differences and AF-associated changes in the expression and phosphorylation of key Ca^{2+}-handling proteins and in ultrastructural properties and dimensions of atrial cardiomyocytes into our recently developed 3D atrial cardiomyocyte model that couples electrophysiology with spatially detailed Ca^{2+}-handling processes. Our simulations of quiescent cardiomyocytes show increased incidence of Ca^{2+} sparks in female vs male myocytes in AF, in agreement with previous experimental reports. Additionally, our female model exhibited elevated propensity to develop pacing-induced spontaneous Ca^{2+} releases (SCRs) and augmented beat-to-beat variability in action potential (AP)-elicited Ca^{2+} transients compared with the male model. Parameter sensitivity analysis uncovered precise arrhythmogenic contributions of each component that was implicated in sex and/or AF alterations. Specifically, increased ryanodine receptor phosphorylation in female AF cardiomyocytes emerged as the major SCR contributor, while reduced L-type Ca^{2+} current was protective against SCRs for male AF cardiomyocytes. Furthermore, simulations of tentative Ca^{2+}-targeted interventions identified potential strategies to attenuate Ca^{2+}-driven arrhythmogenic events in female atria (e.g., t-tubule restoration, and inhibition of ryanodine receptor and sarcoplasmic/endoplasmic reticulum Ca^{2+}-ATPase), and revealed enhanced efficacy when applied in combination.

Conclusions. Our sex-specific computational models of human atrial cardiomyocytes uncover increased propensity to Ca^{2+}-driven arrhythmogenic events in female compared to male atrial cardiomyocytes in AF, and point to combined Ca^{2+}-targeted interventions as promising approaches to treat AF in female patients. Our study establishes that AF treatment may benefit from sex-dependent strategies informed by sex-specific mechanisms.

Translational perspective. Accumulating evidence demonstrates substantial sex-related
differences in atrial fibrillation (AF), which is the most common arrhythmia, with female patients
faring worse with the condition. By integrating known sex-differential components into our
computational atrial cardiomyocyte model we found that female atrial cardiomyocytes in AF
exhibit greater propensity to develop Ca^{2+}-driven arrhythmia than male cardiomyocytes. Model
analyses provided novel mechanistic insights and suggested strategies such as t-tubule
restoration, correction of Ca^{2+}-handling disturbances, and the combination of both, as promising
approaches to treat AF in female patients. Our study uncovers and validate sex-specific AF
mechanisms and inform the development of targeted anti-AF strategies.

Keywords: atrial fibrillation; sex differences; computational modeling; Ca^{2+}-driven arrhythmias;
therapy

Graphical abstract: Sex-specific 3D spatiotemporal models of human atrial cardiomyocyte
Ca^{2+} signaling reveal a greater propensity to develop Ca^{2+}-driven arrhythmic events in female vs
male atrial cardiomyocytes in AF. Model analysis links sex-specific AF remodeling to
arrhythmogenic mechanisms. AF, atrial fibrillation; SCR, spontaneous Ca^{2+} release; CaT,
cytosolic Ca^{2+} transient; RyR2-P, phosphorylated ryanodine receptor type 2 (RyR2); CSQ,
calsequestrin; LTCC, L-type Ca^{2+} channel; PLB, phospholamban; SERCA, sarcoendoplasmic
reticulum Ca^{2+} ATPase.

Introduction

Atrial fibrillation (AF), the world's most common cardiac arrhythmia, is a global epidemic with a prevalence projected to increase 3-fold in the next 50 years.[1,2] AF causes significant morbidity and mortality, and places a substantial socioeconomical burden on healthcare systems. Despite decades of advances in both fundamental research and clinical management, current AF therapies (e.g., antiarrhythmic drugs and catheter ablation) exhibit inadequate efficacy[3] and pose significant risks of adverse effects.[4] The challenge in developing effective AF therapeutic strategies is exacerbated by the marked heterogeneity in disease phenotypes across patient groups[3] coupled with a limited understanding of the underlying pathophysiological mechanisms.

Sex-related differences have been reported in the prevalence, clinical manifestation, management strategies, and treatment outcomes of AF.[5–10] Men develop AF years earlier than women;[5,9,10] however, women experience worse symptoms, more complications arising from drug side effects or ablation, diminished quality of life, and greater risk of AF-related stroke and mortality.[5,9,10] Although these differences may be attributed to inherent differences in cardiac function between male and female patients, the insufficient inclusion of female sex in both fundamental research and clinical cohorts[11] has led to an incomplete understanding of the mechanisms that contribute to these sex differences. Because of this, females often obtain less effective but potentially harmful treatment approaches that lead to worse prognoses. While there is a growing interest in exploring sex differences in ventricular electrophysiology,[12–17] far less attention has been directed to the mechanistic understanding of the differences between the atria of male and female patients. As such, a systematic investigation of these sex differences is urgently required to discover the precise mechanisms of sex-dependent disparities observed in AF to lay the foundation for defining sex-specific therapeutic strategies.

While AF comprises electrophysiological and structural remodeling, dysregulated Ca^{2+} handling has emerged as a key factor in its pathophysiology.[18–23] As in the ventricles, Ca^{2+} crucially mediates cardiac excitation-contraction coupling in the atria, whereby electrical excitation of the cardiomyocytes leads to changes in the intracellular Ca^{2+} that mediate mechanical contraction and relaxation by acting on the myofilament machinery.[24] As a consequence of the coupling between Ca^{2+} and voltage, intracellular Ca^{2+}-handling abnormalities in atrial cardiomyocytes from patients with AF[18,19] not only directly impair contractility,[25] but could also promote ectopic (triggered) activity that significantly contributes to AF initiation and maintenance.[18,19] This is driven by spontaneous Ca^{2+} releases (SCRs) from the sarcoplasmic reticulum (SR) during diastole,

which cause increased Ca^{2+} extrusion by the electrogenic Na^+/Ca^{2+} exchanger (NCX) that triggers

transmembrane potential afterdepolarizations, leading to ectopic activity in the atria.[18,19,23,26,27]

Additionally, Ca^{2+} dysregulation further promotes electrical and structural remodeling in the atria,

thereby increasing the likelihood of AF maintenance.[19] Therefore, the correction of Ca^{2+}-handling

abnormalities has emerged as a promising anti-AF strategy.[19,23,28,29]

Previous experimental work has reported sex differences in the expression and phosphorylation

of Ca^{2+}-handling proteins,[30] t-tubular system density,[31] and cell dimensions[31–33] of atrial

cardiomyocytes. For instance, the well-documented AF-typical reduction in L-type Ca^{2+} current

(I_{CaL})[34,35] was observed in atrial cardiomyocytes from male, but not female patients.[30] Conversely,

atrial cardiomyocytes from females with AF displayed a greater incidence of Ca^{2+} sparks than

males.[30] While these findings are indicative of AF-related sex differences in ionic, structural, and

functional aspects, their precise arrhythmogenic consequences and sex-specific implications are

not yet mechanistically dissected and quantitatively determined. Importantly, there is a scarcity of

studies investigating differences in atrial rate dependency of Ca^{2+}-handling processes between

male and female cardiomyocytes, which are crucial in understanding sex-specific mechanisms of

arrhythmogenesis in AF. While experimentally delineating the precise contribution of these

aspects to sex differences in AF presents significant challenges, computational models have

emerged as a powerful tool for unraveling the subcellular mechanisms of Ca^{2+} dysregulation.[36–39]

In this study, we employed computational modeling to systematically assess the mechanisms of

Ca^{2+}-dependent arrhythmia in male and female atrial cardiomyocytes. We also compared male

vs female responses to various putative therapeutic interventions targeting the subcellular

determinants of intracellular Ca^{2+} handling. To achieve this, we utilized our recently developed

three-dimensional (3D) human atrial cardiomyocyte model that couples electrophysiology with

spatially-detailed subcellular Ca^{2+} signalling.[36,37] We built a suite of male- and female-specific

spatially-detailed models of intracellular Ca^{2+} dynamics in normal sinus rhythm (nSR) and AF

conditions by incorporating all known sex- and AF-dependent alterations in Ca^{2+}-handling,[30] t-

tubular system density,[31,40] and cell dimensions.[31–33,41] We found that female AF atrial

cardiomyocytes displayed a higher incidence and amplitude of SCRs, and a greater beat-to-beat

instability of the paced Ca^{2+} transients *vs* male, owing mainly to female-specific AF remodeling in

ryanodine receptor type-2 (RyR2) phosphorylation. Furthermore, our simulations identified

potential intervention strategies targeting the Ca^{2+} handling system that may yield favorable

outcomes in female atrial cardiomyocytes. Overall, our study underscores and validates the role

of sex-specific differences in AF that warrants future fundamental and clinical studies to develop

sex-specific anti-AF approaches.

Methods

Baseline model and overall study design

Our recently developed three-dimensional model of the human atrial cardiomyocytes that couples

electrophysiology, subcellular Ca^{2+} dynamics, and t-tubular structures[36,37] was extended to

incorporate descriptions of reported subcellular gradients of RyR2 and calsequestrin (CSQ)

distribution,[30] and deemed to represent the male atrial cardiomyocyte in normal sinus rhythm

(nSR). This model served as the reference for adjusting parameters to generate additional

subcellular Ca^{2+} handling models of the male cardiomyocyte in AF conditions, and of female

cardiomyocytes in nSR and AF conditions (**Table 1**, detailed descriptions of parameters are listed

in **Supplementary Table S1**). Our sex-specific models were employed to assess the propensity

of atrial cardiomyocytes to developing Ca^{2+} sparks in quiescent cells and SCR events following

burst-pacing, and to mechanistically investigate the impact of AF- and sex-dependent factors on

Ca^{2+}-driven arrhythmogenesis. Furthermore, we systematically evaluated the responses of atrial

cardiomyocytes to several candidate Ca^{2+}-targeted intervention strategies.

Updated structural characteristics and subcellular distribution of RyR2s and CSQ in sex-specific atrial cardiomyocyte models

To account for the previously reported heterogeneous subcellular localization of RyR2s and

CSQ,[30] we incorporated the density gradient of RyR2s and CSQ in surface Ca^{2+} release units

(CRUs, i.e., the 2 outer CRU layers near cell surface) vs inner (i.e., non-surface) CRUs while

keeping the whole-cell total expression unchanged. Specifically, the density ratio of surface-to-

inner CRUs was assigned to 1.5 for RyR2s and 3 for CSQ, based on reported subcellular

gradients.[30] We incorporated descriptions of cell dimensions based on experimental findings of

cell size differences between sexes and with AF-remodeling, with larger cell dimensions in male

vs female cardiomyocytes[31–33] and in cAF vs nSR conditions[40,41] (**Table 1**). Additionally, to account

for detected differences in t-tubular density between sexes[31] and AF-remodeling,[40] the t-tubular

density was lower in female vs male cardiomyocyte models, or for cAF vs nSR models (**Table 1**),

while maintaining the average branch length of the t-tubular structures. The electrical capacitance

of the new sex-specific models of nSR and cAF cardiomyocytes was computed according to the
modifications of cell dimension and t-tubular density, as done previously.[36,37]

Parameterization of Ca²⁺ handling to construct sex-specific atrial cardiomyocyte models

We updated the Ca^{2+}-handling parameters in male/female and nSR/AF models to describe the

reported sex difference and AF-remodeling of Ca^{2+}-handling proteins expression and

phosphorylation.[30] The modifications are summarized in **Table 1** and include the maximum unitary

flux and inactivation of L-type Ca^{2+}-channel (LTCC, with parameters tuned based on simulated

voltage clamp to recapitulate the LTCC changes by AF in male cardiomyocytes[30]), the expression

of CSQ, phospholamban (PLB), and RyR2 phosphorylation (**Table 1**). Corresponding parameter

values are given in **Supplementary Table S1**.

Table 1. Sex- and AF-dependent changes in the substructure and Ca²⁺-handling processes.

Changed parameters		nSR	Male	Female	Ca²⁺-directed Interventions in cAF
		Female vs Male	cAF vs nSR		
Cell Size	Length	↓ 9.1%[31–33]	↑ 10.9%[40,41]	↑ 10.0%[40,41]	—
	Width	↓ 23.5%[31–33]	↑ 17.6%[40,41]	↑ 23.1%[40,41]	—
Tubular Density		↓ 49.4%[31]	↓ 97.6%[40]	↓ 97.1%[40]	Up-regulation
I_{CaL}	Maximum Unitary Flux	—	↓ 50.0%[30]	—	—
	Ca²⁺-dependent Inactivation Tau	—	↑ 71.5%[30]	—	—
CSQ Expression		↓ 24.0%[30]	↓ 45.0%[30]	↓ 27.6%[30]	Up-regulation
PLB Expression *		—	↓ [30]	—	Up-regulation
RyR2 Phosphorylation		—	—	↑ [30]	Down-regulation
Max. SERCA Activity #		—	—	—	Down-regulation

*: Herraiz-Martinez et al.[30] reported a lower threonine-phosphorylated PLB but greater SERCA
expression in female-vs-male atria in nSR. Since these two sex-dependent differences have opposing
effects on the SERCA activity, we did not incorporate these changes in our models.
#: Given the aforementioned model parameterization of PLB/SERCA (*) and that there was no sex
difference in SERCA expression in cAF, we did not incorporate into our model the AF-induced
reduction of SERCA expression in female atria reported by Herraiz-Martinez et al.[30].

Simulation protocol and analysis of Ca²⁺ sparks and SCRs

To investigate the rate dependence of Ca^{2+} dynamics and SCR events following burst pacing, we

employed an AP clamp protocol. The protocol consisted of a 4-s unstimulated period, followed by

a 28-s AP-clamp period, during which the cardiomyocyte was paced at 0.5, 1, 2, 3, 4, or 5 Hz

using AP-clamp protocols. The protocol concluded with a 5-s non-stimulation pause period to

allow for evaluation of SCR events. The AP traces used in the protocol at each pacing rate were

obtained from current-clamp simulations using the Zhang et. al.[36] model. The holding potential

during the non-stimulated period was set to the resting membrane potential of the AP clamp trace.

To identify SCRs, we analyzed the time course of the averaged $[Ca^{2+}]_{Cyto}$ of each line scan, from

the beginning of the last beat to the end of the 5[th] second no-stimulation period, then excluded

the first $[Ca^{2+}]_{Cyto}$ peak (i.e., AP-induced Ca^{2+} release). The incidence of Ca^{2+} sparks was

determined for each CRU on selected line scans by analyzing the time course of CRU-specific

$[Ca^{2+}]_{Cyto}$ during the 15-s no-stimulation period. In both cases, the respective $[Ca^{2+}]_{Cyto}$ time

courses, sampled at 1 kHz, were denoised using a moving-averaging filter (window size = 5). To

determine the incidence and amplitude of SCRs, and latency of the first SCR, as well as the

incidence of Ca^{2+} sparks, we utilized and the 'findpeak' function in MATLAB R2021(MathWorks,

Natick, MA, USA), with 'MinPeakProminence' of 0.05 μM and 'MinPeakDistance' of 50 ms for the

findpeak function to define SCRs for the averaged line scan $[Ca^{2+}]_{Cyto}$, and Ca^{2+} sparks for each

CRU $[Ca^{2+}]_{Cyto}$. The amplitudes of SCR events were calculated as the difference between the

peak and the minimum $[Ca^{2+}]_{Cyto}$ values during the no-stimulation period. To minimize the

dependence of detected SCR events on the location of line scan, we monitored the $[Ca^{2+}]_{Cyto}$

along 13 transversal lines selected from the middle z-stack of simulated cardiomyocytes, with a

longitudinal spacing of 2.7 μm (i.e., 3 CRUs) between the line scans.

Simulation of varying individual Ca²⁺-handling determinants and candidate therapeutic

interventions

We investigated the contribution of each sex- and AF-associated change to atrial arrhythmogenic

events through sensitivity analysis by varying one factor at-a-time. The targets included cell width,

t-tubular density, the relative change of PLB expression, RyR2 phosphorylation, and LTCC activity.

In addition, we simulated several interventions targeting atrial Ca^{2+} handling to demonstrate

potential sex differences in anti-arrhythmia intervention outcomes. These interventions included

t-tubular system restoration, PLB and CSQ overexpression, sarcoplasmic/endoplasmic reticulum

Ca^{2+}-ATPase (SERCA) inhibition, and RyR2 dephosphorylation. A detailed descriptions of the

parameters for these interventions are listed in **Table 1** and values given in **Supplementary**
Table S2.

Numerical methods, statistical analysis, and data availability

Our sex-specific atrial cardiomyocyte models were implemented in C++ and parallelized using

OpenMP 5.1 (https://www.openmp.org/), and with a combination of deterministic and stochastic

modeling as done previously.[36,37] We improved the accuracy of stochastic simulation of the

Markov model of RyR2s by replacing the binomial-distribution-based random generator (which

was applied to simulate the stochastic transition of RyR2s among the 4 RyR2 Markov states) with

a multi-nominal stochastic generator, as implemented in Numpy.[42] The source code of our new

3D   human   atrial   cell   model   can   be   accessed   from

http://elegrandi.wixsite.com/grandilab/downloads and https://github.com/drgrandilab. Statistical

analysis was performed by One-way ANOVA with planned comparison and Bonferroni correction

using MATLAB (Version 2022b, The MathWorks, Natick, MA, USA) with a threshold significance

level of 0.05.

Results

Simulated female vs male AF cardiomyocytes exhibit a greater incidence of Ca^{2+} sparks at

rest, similar to experimental observation

We constructed male- and female-specific spatially detailed stochastic models of Ca^{2+} handling

for both nSR and chronic AF (cAF) conditions within a human atrial cardiomyocyte model. To test

the robustness of our sex- and AF-specific parameterization, we first compared the incidence of

subcellar Ca^{2+} sparks in the male and female atrial cardiomyocyte models at rest with

experimental data from human samples. The membrane voltage of the model cardiomyocytes

were clamped to -72 mV, which corresponds to the resting membrane potential of the baseline

model[36] at 0.5 Hz of steady-state pacing. Transversal line scan analyses (**Fig. 1Ai,ii**) of simulated

$[Ca^{2+}]_{Cyto}$ revealed increased Ca^{2+} spark incidence in AF vs nSR conditions (**Fig. 1B**) in both male

and female cardiomyocytes. There was no sex difference in the Ca^{2+} spark frequency in nSR

conditions. However, in AF-remodeled cardiomyocytes the female model exhibited substantially

greater Ca^{2+} spark abundance compared with the male model. These results are confirmed in

additional simulations wherein the cardiomyocyte resting membrane potential was clamped at

various levels (**Supplementary Fig. S1**). Importantly, the AF- and sex-dependent differences in

Ca^{2+} spark incidence observed in our simulations qualitatively align with a previous experimental

study comparing Ca^{2+} sparks in resting atrial cardiomyocytes from male and female patients with

and without AF (**Fig. 1C**).[30] These simulations suggest that the sex- and AF-related differences

incorporated in our models recapitulate the sex differences observed in atrial Ca^{2+} spark

properties.

Fig. 1. **Simulated effects of sex and AF on Ca^{2+} sparks and comparison to experiments.** (**A**)
Simulated transversal line scan of intracellular Ca^{2+} from (**i, iii**) male and (**ii, iv**) female atrial
cardiomyocytes in (**i, ii**) nSR and (**iii, iv**) cAF conditions. (**B-C**) Quantification of Ca^{2+} spark (**B**)
frequency in simulations, and (**C**) comparison of experimental Ca^{2+} spark (i) sites and (ii) density
reported by Herraiz-Martinez et al.[30].

Fast atrial rate-induced SCR events are more frequent in female vs male cardiomyocytes

in AF

The elevated Ca^{2+}-spark incidence in female AF may be indicative of arrhythmogenic Ca^{2+}-driven

instabilities in female vs male atria. However, although sex difference in Ca^{2+} sparks has been

reported from quiescent human atrial cardiomyocytes,[30] SCRs and their rate-dependence that is

critically relevant to AF, have not yet been investigated. As such, we next sought to determine the

arrhythmogenic propensity of atrial cardiomyocytes using a pacing-pause protocol. Because

experimental characterization of sex differences in atrial electrophysiology is lacking, and a recent

study reported no difference between male and female AP characteristics in AF at 1 Hz,[43] we

applied an AP-clamp protocol to provoke SCRs using the same AP configuration for both sexes.

Fig. 2. Simulated effects of sex and AF on spontaneous Ca²⁺ release (SCR) events in human atrial cardiomyocytes. (A) **(i)** AP-clamp traces at 3 Hz and subsequent resting period, **(ii)** whole-cell averaged cytosolic Ca²⁺, **(iii-v)** transverse line scan of cytosolic Ca²⁺ in **(iii)** nSR vs **(iv)** cAF, and **(v)** the corresponding averaged line-scan Ca²⁺ traces for male (*left*) vs female (*right*) cardiomyocytes. White arrow heads indicate SCRs and black arrows mark the start of an AP. (**B-C**) SCR incidence was significantly higher (***B***) and SCR amplitude was significantly greater (***C***) in female vs male cells in AF conditions, but there was no sex-dependent difference in SCR incidence and amplitude in nSR. (**D**) Latency to the first SCR. Columns (i-iv) of (**B-D**) depict data from AP clamps of 2 Hz to 5 Hz, respectively. The bottom panels represent the amplitude of the first SCR for each line scan. ***: p < 0.001; **: p < 0.01; *: p < 0.05; ns: not significant.

This allowed studying the effects of sex differences in Ca^{2+} handling independent of possible sex
differences in atrial electrophysiology. To achieve this, we generated trains of APs by pacing our
baseline cardiomyocyte model[36,37] at various rates and applied these APs to all cardiomyocyte
models prior to holding at the resting membrane potential to record SCRs (**Fig. 2Ai**). Transversal
line scans of $[Ca^{2+}]_{Cyto}$ were obtained (**Fig. 2Aiii-iv**) to assess local SCRs, which were identified
as post-repolarization instability in the averaged line-scan $[Ca^{2+}]_{Cyto}$ (**Fig. 2Av**). We found that the
stimulation protocol led to SCRs in both male and female atrial cardiomyocytes (**Fig. 2Aiii,iv**).
There was no sex-dependent difference in SCR incidence in nSR conditions except for the 5-Hz
AP clamp (**Fig. 2B**). However, in AF conditions female cardiomyocytes exhibited a greater
incidence of SCRs compared to male cardiomyocytes (**Fig. 2Aiv,v** and **2Bi-iv**). Of note, the sex-
dependent difference in SCR incidence was not attributed to differences in SR load, which was
similar between male and female cardiomyocytes in nSR, and slightly lower in the female vs male
cardiomyocytes in AF (**Supplementary Fig. S4Biv**). Interestingly, AF caused a reduction (vs nSR)
in the maximum amplitude of SCRs in male cardiomyocytes (**Fig. 2Ci-iv**), owing to enhanced
SERCA activity (due to PLB reduction) and whole-cell Ca^{2+} unloading due to LTCC reduction (see
Graphical abstract). In contrast, the maximum SCR amplitude was elevated in AF-vs-nSR
female cardiomyocytes (**Fig. 2Ci-iv**), due to hyperactive RyR2 (see **Graphical abstract**). Overall,
the sex-dependent differences in AF were amplified with faster AP rates, displaying a positive rate
dependence (**Fig. 2Bi-iv,Ci-iv**). We also found that in male AF the latency of SCRs was delayed
compared to female cardiomyocytes and nSR conditions (**Fig. 2Di-iv**). Collectively, these
simulations showed sex differences in the incidence of potentially proarrhythmic SCRs in human
AF and pointing to a higher susceptibility to Ca^{2+}-dependent arrhythmias in female vs male
cardiomyocytes.

Sensitivity analysis unmasks the individual contribution of sex- and AF-dependent
differences to SCR

Our simulations revealed sex-dependent differences in propensity to SCRs due to the combined
effects of sex-related and AF-induced differences in subcellular Ca^{2+}-handling. To dissect the
contribution of individual alterations, we performed parameter sensitivity analysis by varying one
parameter at-a-time in the nSR male-specific model (**Fig. 3**) under AP trains (3-Hz pacing) applied
to provoke SCRs. Simulating varying cell width did not affect SCR incidence, with a trend of
augmented SCR amplitude in wider myocytes (**Fig. 3Ai-ii**). This suggests that the increased
cardiomyocyte size in AF vs nSR conditions or male vs female atria may have a limited
contribution to the simulated disease- and sex-dependent phenotypes. Conversely, decreasing t-

tubular density substantially increased both incidence and amplitude of SCRs (**Fig. 3Bi-ii)**,
indicating a potential proarrhythmic role of the t-tubular loss noted in AF. Indeed, our previous
work demonstrated that loss of tubules promoted SCRs in paced cardiomyocytes through
reduction of NCX-dependent Ca^{2+} export, causing diastolic Ca^{2+} accumulation and increased
RyR2 open probability.[36] These effects were stronger for the RyR2s located at the cell interior
than those at the periphery and enhanced by faster AP rates.[36] Taken together, these results
suggest that the loss of t-tubules promotes arrhythmic events in both sexes.

Fig. 3. Individual contribution of sex- and AF-dependent alterations to SCR event formation
in human atrial cardiomyocytes determined by changing one parameter at-a-time. (i) SCR
incidence, **(ii)** SCR amplitude, and **(iii)** SCR latency simulated using male cardiomyocyte model
in nSR conditions, by varying **(A)** cell width, **(B)** t-tubule density, **(C)** relative change of PLB
expression, **(D)** relative change of RyR2 phosphorylation, and **(E)** relative change of LTCC activity.
***: $p < 0.001$; **: $p < 0.01$; *: $p < 0.05$; ns: not significant.

Similarly, reducing PLB expression (e.g., as seen in male AF cardiomyocytes) caused moderate
elevation of SCR incidence, while slightly decreasing SCR amplitude (by promoting SERCA
activity) and latency (**Fig. 3Ci-iii**). Our simulations showed that increased RyR2 phosphorylation
(as noted in the female atria[30]) alone moderately increased SCR incidence and shortened SCR
latency, although with slightly lower amplitude (**Fig. 3Di-iii**). Furthermore, reducing L-type Ca^{2+}

channel activity drastically diminished SCR incidence and amplitude and delayed SCR latency
(**Fig. 3Ei-iii**), due to unloading of the cytosolic and SR Ca^{2+} (see **Graphical abstract**). Finally,
our previous modeling work demonstrated a SCR-promoting effect of CSQ downregulation.[37]
Therefore, the AF-associated reduction in PLB expression in male atria and elevated RyR2
phosphorylation in the female atria augmented SCRs in male and female cardiomyocytes,
respectively, whereas the AF-induced reduction of L-type Ca^{2+} channel activity in the male atria
could ameliorate the propensity to SCRs, thus contributing substantially to the female-vs-male
increase in Ca^{2+}-dependent arrhythmias in AF.

Interventions targeting individual Ca^{2+}-handling determinants alleviates SCRs

Correcting dysfunctional intracellular Ca^{2+} signaling in AF is a promising anti-AF strategy.[36–39] As
our simulations predicted the increase in AF-associated arrhythmogenic SCRs in female vs male
atrial cardiomyocytes, we next sought to evaluate their responses to putative Ca^{2+}-targeted
interventions. We simulated the upregulation of t-tubular structure (Tubule up), CSQ (CSQ up)
and PLB expression (PLB up), and downregulation of RyR2 (RyR2 down, i.e., dephosphorylation)
and SERCA (SERCA down, i.e., inhibition) (**Fig. 4**). As above, we applied a 3 Hz AP-clamp
protocol (**Fig. 4A,B**) to compare the SCR characteristics in AF cardiomyocytes with the proposed
interventions. We found that all putative interventions could suppress SCR incidence (**Fig. 4C**)
and delay SCR latency (**Fig. 4E**), with t-tubule recovery and CSQ upregulation also decreasing
SCR amplitude in female cardiomyocytes (**Fig. 4Di,iv**). We define the effectiveness of an
intervention as a reduction of SCR incidence post-intervention reaching a level comparable to or
below the baseline observed in male cardiomyocytes under nSR conditions (shown with dashed
lines in **Fig. 4C-E**). T-tubule recovery (**Fig. 4Bi, Ci**), RyR2 dephosphorylation (**Fig. 4Cii,**
Supplementary Fig. S2B), and SERCA inhibition (**Fig. 4Ciii, *Supplementary Fig. S2C***) alone
were effective in preventing SCRs in both sexes, with greater SCR-suppressing responses in
female vs male cardiomyocytes. In contrast, PLB overexpression was effective in male but not
female cardiomyocytes (**Fig. 4Bii, Cv**). CSQ overexpression alone was not effective against SCR
in either sex (***Supplementary Fig. S2D,* Fig. 4Civ**).

Fig. 4. Simulated effects of targeted interventions of Ca²⁺-handling determinants in male and female atrial cardiomyocytes from AF patients. **(A) (i)** AP-clamp traces and subsequent

resting period, **(ii-iii)** simulated transversal line scans of $[Ca^{2+}]_{Cyto}$ for male and female
cardiomyocytes in **(ii)** nSR and **(iii)** AF conditions as in **Fig. 2A** are illustrated for comparison. **(B)**
Simulated transversal line scans of $[Ca^{2+}]_{Cyto}$ with interventions upregulating (**i**) t-tubular structure
and **(ii)** PLB in human atrial cardiomyocytes with AF. White arrowheads indicate SCRs and black
arrows mark the start of AP. **(C-E)** Summary of intervention effects on (**C**) SCR incidence, (**D**)
SCR amplitude, and (**E**) SCR latency; columns (**i-v**) illustrate effects of proposed interventions
with (**i**) Tubule up, (**ii**) RyR2 down, (**iii**) SERCA down, (**iv**) CSQ up, and (**v**) PLB up. In Panel (**C**),
the numbers indicate the mean reduction in SCR incidence due to intervention. ***: $p < 0.001$; **:
$p < 0.01$; *: $p < 0.05$; ns: not significant.

Next, we performed additional simulations to test the effects of multi-target interventions with

pairwise combination of the proposed maneuvers (**Fig. 5A-E** and **Supplementary Fig. S3A-E**).

In general, compared with the single-hit interventions (**Fig. 4**), combined interventions caused a

stronger SCR suppression (**Fig. 5C**), led to greater reductions of SCR amplitude (to values lower

than nSR baseline, **Fig. 5D**), and stronger prolongation of SCR latency (**Fig. 5E**). Overall, the

extent of SCR suppression was more pronounced in female vs male cardiomyocytes. Interestingly,

CSQ overexpression presented an exception, whereby the combined interventions of CSQ

overexpression with another intervention resulted in less effective outcomes than with the

standalone application of the individual intervention (**Supplementary Fig. S3Cii-v**). Collectively,

our simulations suggest that the combined interventions that correct t-tubular structure, RyR2

phosphorylation, SERCA activity and PLB expression, are more effective in suppressing SCRs in

compared to a single-hit approach, with female cardiomyocytes showing additive responses to

these combined interventions.

Beat-to-beat Ca^{2+}-transient variability is greater in female vs male cardiomyocytes in AF

Our simulations uncovered an increased propensity to SCRs in female vs male cardiomyocytes

in AF and demonstrated SCR-suppressing effects of interventions targeting subcellular Ca^{2+}-

handling determinants. However, how sex modifies the characteristics of the Ca^{2+} transient (CaT)

elicited during APs, especially in AF, is unknown. Therefore, we compared the AP-evoked CaT

properties in our male and female atrial cardiomyocyte models in both nSR and AF conditions, by

clamping the membrane with AP trains at frequencies ranging from 0.5 to 5 Hz (**Supplementary**

Fig. S4A). Analyses of whole-cell CaT demonstrated rate-dependent increases of systolic and

diastolic Ca^{2+} levels as well as acceleration of relaxation for the cardiomyocyte models from both

sexes (**Supplementary Fig. S4B**). In both male and female, simulations recapitulated the

experimentally observed reduction of CaT amplitude in atrial cardiomyocytes of AF patients.[22,23]

Fig. 5. Simulated effects of pair-wise combined interventions on male and female atrial cardiomyocytes from AF patients. **(A) (i)** AP-clamp traces, **(ii-iii)** simulated transversal line-scan $[Ca^{2+}]_{Cyto}$ for male and female cardiomyocytes in **(ii)** nSR and **(iii)** AF conditions as in **Fig. 2A** are illustrated for comparison. **(B)** Simulated transversal line scans of $[Ca^{2+}]_{Cyto}$ with combined interventions of (**i**) RyR2 down + Tubule up, and (**ii**) RyR2 down + SERCA down. White arrow heads indicate SCRs and black arrows mark the start of AP. **(C-E)** Summary of intervention effects on **(C)** SCR incidence, **(D)** SCR amplitude, and **(E)** SCR latency; columns **(i-v)** depict effects of combined intervention with (**i**) RyR2 down + Tubule up, (**ii**) SERCA down + Tubule up, (**iii**) PLB

441 up + Tubule up, (**iv**) RyR2 down + SERCA down, and (**v**) RyR2 down + PLB up. In Panel (**C**), the
442 numbers indicate the mean reduction of SCR incidence due to intervention. ***: $p < 0.001$; **: $p <$
443 0.01; *: $p < 0.05$; ns: not significant.
444

However, the reduction of CaT amplitude in male cardiomyocytes was the consequence of the

lower systolic $[Ca^{2+}]_{Cyto}$, whereas in the female cardiomyocytes this was due to the higher diastolic

$[Ca^{2+}]_{Cyto}$ (**Supplementary Fig. S4Bi-ii**). Male cardiomyocytes also displayed a slightly greater

systolic $[Ca^{2+}]_{Cyto}$ than the female cardiomyocytes in nSR conditions, but this pattern was reversed

in AF conditions (**Supplementary Fig. S4A-B**). Finally, the simulated SR Ca^{2+} load was not

different between male and female cardiomyocytes in nSR conditions and slightly lower in the

female vs male cardiomyocytes in AF conditions. These observations in the SR load are

consistent with previous experimental measurements using the time-integral of the caffeine-

induced (NCX) current.[30]

Beat-to-beat variability of the CaT during pacing cycles may induce AP instabilities that could

promote arrhythmogenesis. We sought to determine if the more frequent diastolic SCRs do

increase beat-to-beat variability of the atrial CaT. The simulated transversal line scans of

subcellular CaT measured at the center of the cell revealed a greater asynchrony (i.e., U-shaped

pattern) in centripetal CaT propagation for male vs female cardiomyocytes, a phenomenon that

was promoted by AF in both sexes (**Fig. 6Aii-iii**). Like the SCR elicited following pacing-pause

AP-clamp, these transversal line scans of cytosolic Ca^{2+} revealed diastolic SCRs between the

AP-induced CaTs (**Fig. 6Ai-iv**). The incidence of diastolic SCRs was higher for AF vs nSR

conditions in the female but not male cardiomyocytes (**Fig. 6Aii-iv**), consistent with the greater

incidence of SCRs after pacing in the female cardiomyocytes in AF (**Fig. 2B**).

We found that the diastolic SCRs show a beat-to-beat variation in incidence and amplitude,

especially in AF (**Fig. 6Aii-iii**). This variation may impact SR load and RyR2 refractoriness and

could increase the beat-to-beat variability in AP-coupled CaT amplitude. We computed the beat-

to-beat variation of whole-cell CaT amplitude evoked using AP clamp at 0.5 to 5 Hz

(**Supplementary Fig. S5A-B**). While the beat-to-beat variation in CaT amplitude was similar

between the two sexes in nSR conditions, AF simulations unmasked sex-dependent changes.

Elevated beat-to-beat CaT variation was observed at fast rates in AF vs nSR in female

cardiomyocytes (3 to 5 Hz), whereas beat-to-beat CaT variation was reduced in male

cardiomyocytes (4 and 5 Hz). Thus, at fast rates beat-to-beat CaT variability is greater in female

vs male AF cardiomyocytes, pointing to a greater Ca²⁺-handling instability in the female atrial

cardiomyocytes in AF.

Fig. 6. **Simulated transversal line scans of AP-clamp evoked Ca²⁺ transients in male and**

female atrial cardiomyocytes and responses to interventions in AF. (A) (i) AP-clamp traces,

(ii-iv) simulated transversal line scans of AP clamp-elicited Ca²⁺ transients for male and female

cardiomyocytes in **(ii)** nSR and **(iii)** AF conditions, and **(iv)** whole-cell averaged traces of the
corresponding CaTs. **(B)** Simulated transversal line scans of AP clamp-elicited Ca^{2+} transients
from cardiomyocytes in AF following interventions causing **(i)** Tubule up, **(ii)** RyR2 down, and **(iii)**
comparison of whole-cell averaged traces of the corresponding CaTs with and without
intervention. **(C)** Simulated transversal line scans of AP clamp-elicited Ca^{2+} transients from
cardiomyocytes in AF following combined interventions of **(i)** RyR2 down + Tubule up, **(ii)** RyR2
down + SERCA down, and **(iii)** comparison of whole-cell averaged traces of the corresponding
CaTs with and without intervention. Arrows mark the start of AP, and the diastolic SCRs are
indicated with arrow heads.
490

Sensitivity analysis revealed the impact of individual sex-dependent differences on the beat-to-
beat CaT variability (**Supplementary Fig. S5C**). We found that female-to-male differences, *i.e.,*
lower cell width and t-tubular density, but greater LTCC activity and RyR2 phosphorylation, could
all contribute to greater beat-to-beat CaT variability in female vs male cardiomyocytes, whereas
the PLB loss alone (as noted in male cardiomyocytes) could elevate the beat-to-beat CaT
variability for the male cardiomyocytes (**Supplementary Fig. S5Ciii**).

We also evaluated how the Ca^{2+}-targeted interventions impacted the beat-to-beat CaT variability
in AF (**Fig. 6B,C, Supplementary Fig. S6**). Simulated line-scan $[Ca^{2+}]_{Cyto}$ showed that correcting
RyR2 (**Fig. 6Bii**), SERCA, PLB, or CSQ alone (**Supplementary Fig. S6Aiii-v**) did not affect the
asynchronized, U-shaped subcellular CaT propagation. In contrast, t-tubule recovery (either
increasing t-tubular density alone or combined with an additional intervention) improved the
synchronization of AP-evoked subcellular CaTs (**Fig. 6Bi**). In general, the simulated interventions
reduced the incidence of diastolic SCRs in both sexes (**Fig. 6B,C, Supplementary Fig. S6Aiv-**
viii); however, upregulation of CSQ instead promoted diastolic SCRs in female cardiomyocytes
(**Supplementary Fig. S6Aiii**). The efficacy to suppress diastolic SCRs was enhanced with
combined compared to single-hit interventions, especially for interventions that involved t-tubular
recovery in the male cardiomyocytes (**Fig. 6C, Supplementary Fig. S6Avi-viii**).

Fig. 7 and **Supplementary Fig. S7** summarize the effects of these interventions on the AP-
induced CaT amplitude, CaT beat-to-beat variability, and the post-AP SCR incidence, in female
and male atrial cardiomyocytes with AF, respectively. Indeed, these interventions generally could
suppress SCR incidence (**Fig. 7A**), and lower SCR amplitude (**Fig. 7B**), increase (**Fig. 7C**) the
CaT amplitude, and reduce the beat-to-beat CaT variability (**Fig. 7D**) in female cardiomyocytes.
Similar effects were observed in male cardiomyocytes (**Supplementary Fig. S7A-D**). To compare
the effects of the tested interventions for female atrial cardiomyocytes in AF, we performed a
multi-objective   assessment   of   combined   intervention   regimens.   Given   that   the   male

cardiomyocytes displayed a lower incidence of SCRs, we used the phenotype of male atrial

cardiomyocytes in nSR as reference to assess the intervention outcomes of the female atrial

cardiomyocytes. When solely considering the SCR incidence and amplitude, 8 intervention

regimens (out of 15) were deemed effective for female cardiomyocytes (**Fig. 7 A,B**). This number

dropped to 5 for female cardiomyocytes when a reduction of beat-to-beat CaT variability was

included in the objectives (**Fig. 7D**). The top 6 performing interventions for female atrial

cardiomyocytes in AF are 'double-hit' interventions involving combinations of t-tubular system

improvement with either downregulation of RyR2 or SERCA, or upregulation of PLB (Group 1,

Fig. 7). Overall, these combined intervention regimens outperformed the respective single-hit

interventions. Interestingly, combinations that added CSQ worsened the outcome compared to

single-hit interventions without CSQ (Group 2, **Fig. 7**), while sole PLB and CSQ modulations were

not effective in female cardiomyocytes (Group 3, **Fig. 7**), and CSQ modulation alone failed also

to affect male cardiomyocytes (**Supplementary Fig. S7**). Of note, while all interventions could

increase the CaT amplitude (and thus improve the related atrial contractility), interventions that

incorporated t-tubular system restoration were generally most effective at elevating CaT

amplitude (and thus contractility). Taken together, these results suggest that combining t-tubular

structure restoration with correction of Ca^{2+}-handling protein abnormalities may be a viable

approach to target Ca^{2+}-driven arrhythmias in females with AF.

Fig 7. Summary data comparing the intervention effects on SCR (A) incidence and (B) amplitude, (C) whole-cell CaT amplitude, and (D) CaT beat-to-beat variability in female atrial cardiomyocytes. Dashed lines indicate the referencing values obtained from male cardiomyocytes in nSR conditions. Groups 1 – 3 highlight (1) combined interventions targeting t-tubules, RyR2s, SERCA, and PLB, (2) combinations of CSQ intervention with interventions targeting t-tubules, RyR2s, SERCA, or PLB, and (3) interventions with PLB and CSQ overexpression.

Discussion

A deeper mechanistic understanding of the sex-dependent differences in AF is urgently required to advance the development of tailored, sex-specific approaches to treat this prevalent condition. In this study, we have built a suite of sex-specific models coupling the spatiotemporal dynamics between electrophysiology and subcellar Ca^{2+} signaling. We leveraged our recently developed 3D human atrial cardiomyocyte model to incorporate known sex- and AF-related differences in Ca^{2+} handling and cellular ultrastructure. Our simulations revealed that, in AF conditions, female vs male atrial cardiomyocytes exhibited greater frequency of Ca^{2+} sparks and a higher susceptibility to pacing-induced SCRs. The beat-to-beat variability in the AP-evoked CaT was also greater in female vs male cardiomyocytes. Combined, our simulation data demonstrated a higher propensity to Ca^{2+}-driven arrhythmogenic events in female compared to male atrial cardiomyocytes in AF conditions, which was attributed to sex-dependent differences in AF-remodeled components,

primarily increased RyR2 phosphorylation in female and reduced LTCC activity in male atrial

cardiomyocytes. Furthermore, our modeling identified Ca^{2+}-targeted interventions that could

attenuate the Ca^{2+}-driven arrhythmic events and informed about combined interventions that

correct Ca^{2+}-handling abnormalities (e.g., RyR2 dephosphorylation, SERCA inhibition, and PLB

overexpression) and restore t-tubule system as potentially effective strategies for suppressing

Ca^{2+}-driven arrhythmias in female patients with AF. Collectively, our study establishes that AF

treatment may benefit from sex-dependent therapeutic approaches informed by sex-specific

mechanisms.

Female atrial cardiomyocytes reveal greater propensity to proarrhythmic SCRs

Despite substantial sex-dependent differences in the clinical symptoms, prescribed treatment

options, and outcomes of AF, mechanistic studies to uncover the sex-specific underpinnings of

AF pathophysiology have been insufficient. Our sex-specific models show greater incidence of

Ca^{2+} sparks in female vs male atrial cardiomyocytes with AF at rest (**Fig. 1**), in agreement with

previous experimental reports.[30] Using a pacing protocol that is relevant to studying arrhythmia,

our simulations demonstrated that female vs male cardiomyocytes exhibited much greater

incidence and amplitude of pacing-induced SCRs in AF conditions, along with a shorter SCR

latency (**Fig. 2**), all reflecting more aberrant RyR2 function and pointing to greater propensity to

developing Ca^{2+}-dependent triggered activities in females. Interestingly, previous clinical

observations documented a greater incidence of non-pulmonary vein triggers in female vs male

atria with AF.[7] This may be, at least in part, due to the greater propensity to cellular triggered

activity in female vs males that we identified here. Thus, our simulations offer plausible

mechanistic insights into this sex-dependent difference in clinical AF presentation. Additionally,

our data show that the beat-to-beat variability of AP-elicited CaTs was much greater in female vs

male cardiomyocytes in AF, despite being comparable in nSR conditions (**Fig. 6** and

Supplementary Fig. S5). The elevated beat-to-beat variability in female cardiomyocytes was

paralleled by a greater incidence of diastolic SCRs (**Fig. 6**). Through Ca^{2+}/V_m bidirectional

coupling, the instability in CaT could translate into beat-to-beat variability in APs, and subsequent

increase in spatial AP heterogeneity in tissue that promotes vulnerability to AF-maintaining reentry.

Combined, our simulation data point to elevated propensity to developing both ectopic activity

(through a Ca^{2+}-dependent trigger) and reentry (due to beat-to-beat variability in CaT) in female

vs male atria in AF. On the other hand, our simulations suggest that the triggered activity

mechanism may be less prominent in male atria with AF and imply that reentry (which is caused

by LTCC reduction-mediated shortening of AP and refractoriness) may be the dominant AF-
promoting mechanism in male patients.

Sex-specific molecular and ultrastructural signatures cause SCRs in female atria with AF
Sensitivity analyses of the sex-differential components revealed the individual contribution of each
element to the greater propensity to Ca^{2+}-dependent arrhythmia in female vs male
cardiomyocytes. We found that the greater RyR2 phosphorylation in female and reduced LTCC
function in male cardiomyocytes are major determinants of the observed sex-dependent
differences (**Fig. 3**), positioning aberrant RyR2s as potential targets for AF treatment in female
patients. These results are consistent with our previous study suggesting that reported LTCC
reduction by AF is an adaptive mechanism of the atria to limit triggered activity, whereas enhanced
RyR2 phosphorylation constitutes a maladaptation that drives the arrhythmia.[28] In this regard, our
suite of sex-specific human atrial cardiomyocyte models is unique and instrumental to gain
mechanistic insights into the sex-dependent differences in Ca^{2+}-driven arrhythmogenesis in
human atria, providing a basis to inform putative treatment strategies to correct the deranged Ca^{2+}
signals, especially for the female atria.

Interventions targeting t-tubules and Ca^{2+} handling effectively suppress SCRs in females
Because Ca^{2+}-dependent arrhythmogenesis may be the dominant AF-promoting mechanism in
female atria, suppression of Ca^{2+}-dependent arrhythmia should be a prioritized objective for
therapeutic selection and drug development for female patients. As the Ca^{2+}-dependent trigger is
not a primary target of current AF therapies, the elevated propensity to this trigger may be a
reason why female patients respond more poorly to existing treatment options. For example,
clinical evidence suggests that AF recurrence rates after ablation were greater in female vs male
patients,[44] which could be potentially explained by the greater incidence of triggered activity in the
female atria. Here, our simulated interventions targeting the t-tubular system and the Ca^{2+}-
handling machinery (**Figs. 5 – 7**) show that suppression of SCR incidence, reduction of beat-to-
beat CaT variability, and promotion of subcellular CaT synchronization could constitute promising
anti-AF approaches in female patients. The first two would attenuate Ca^{2+} dependent arrhythmia
to achieve rhythm control, while the latter could help rescue atrial contractile dysfunction, which
is substantially diminished in AF and favors thrombosis. Interestingly, simulated interventions
solely targeting PLB, CSQ, or SERCA did not fully correct SCRs, whereas correcting RyR2s or
the t-tubule system alone did not rescue the reduced CaT amplitude (**Fig. 7**). Overall, our
simulations suggest that combined interventions of multiple therapeutic targets are more effective

than single-hit strategies. Interventions combining t-tubular structure recovery, PLB
overexpression, and a reduction of SERCA and RyR2 activities, as highlighted in Group 1 in **Fig.**
7, are the most effective strategies to suppress proarrhythmic SCRs, while also increasing CaT
amplitude and thus contractility in female atria. Of note, combinations involving CSQ emerged as
exceptions, underscoring the complexity of SCR regulation by CSQ due to the dual effects of
CSQ on RyR2 gating and luminal Ca^{2+} buffering in the SR.[37]

The Ca^{2+}-targeted intervention strategies identified by our modeling work warrant preclinical
experimental assessments. For instance, RyR2 inhibition can be achieved by repurposing
dantrolene, a clinical drug used as a RyR1 inhibitor, or using *ent-verticilide*, which is a selective
RyR2 inhibitor that suppressed AF susceptibility caused by hyperactive RyR2s.[45] Indeed,
dantrolene has been shown to reduce SCRs in human atria and triggered activity in human
ventricles.[46–48] Also, propafenone[49] and flecainide are found to act through inhibition of RyR2s
when treating catecholaminergic polymorphic ventricular tachycardia.[49,50] Interestingly,
propafenone has greater efficacy of pharmacological cardioversion in female patients,[51] while the
efficacy of preventing AF recurrence by class Ic antiarrhythmic drugs (which includes propafenone
and flecainide) was greater in female vs male patients after receiving catheter ablation.[52] In
addition, SERCA inhibition may leverage investigational compounds, such as thapsigargin and
cyclopiazonic acid.[53] Although SERCA upregulation has been pursued for treatment of other heart
conditions, e.g., heart failure,[54] here we found SERCA inhibition instead could reduce SR content
to attenuate RyR2 activity (resulting in reduced SCRs, **Fig. 4**), and augmented CaT amplitude
(**Fig. 7**). Moreover, restoration of t-tubular structure may take advantage of upregulating several
proteins that have been implicated in the regulation of t-tubules, namely junctophilin 2 (JPH2),[55]
amphiphysin II (AMPII, also known as bridging integrator 1/BIN-1),[56,57] and myotubularin
(MTM1),[58] among others. Gene therapies to increase the protein expression have shown
improved outcomes in heart failure by augmenting JPH2,[55] and AAV9 transduction of BIN1
restores t-tubule microdomains to improve cardiac function in stressed[57] and db/db diabetic
mice.[59] Interestingly, treatment using tadalafil, a PDE5 inhibitor, restored t-tubules by upregulating
BIN1 expression.[60] While these studies primarily focused on the ventricles, similar treatments and
effects may be translated into the atria. Lastly, although reports have been limited, gene therapy
may also be applied to augment PLB expression. As such, the therapeutic strategies identified by
our modeling work are all experimentally tractable in the future.

Inclusion of female subjects is key to define sex-specific disease mechanisms

Accumulating evidence show a much lower representation of female sex in both basic and clinical

research of cardiovascular diseases,[61,62] which naturally leads to an inappropriate extrapolation

of study outcomes from men to women.[62] Therefore, a more inclusive representation of female

subjects in research is imperative to identify sex-specific disease mechanisms and facilitate the

development of targeted treatment strategies considering the unique physiological and

pathological aspects of cardiovascular health in female patients. Aligned with this theme, our

study reveals increased Ca^{2+}-driven arrhythmia mechanisms in female patients with AF, and

suggests specific correction of aberrant Ca^{2+}-dependent processes as a potential novel

therapeutic intervention. Moreover, our study provides novel sex-specific computational models

as powerful integrative tools to study emerging biological variables and (patho)physiological

mechanisms that are dependent on sex.

Study limitations

Our modeling approach has several limitations that warrant future investigations. First, our

assessment utilized sex-independent AP clamp to reveal Ca^{2+}-driven arrhythmic events in the

absence of well-established knowledge of sex-dependent differences in atrial APs. Interestingly,

recent characterization of atrial APs from male and female patients revealed some sex-dependent

differences in AP duration and resting membrane potential; however, these differences were only

present in nSR conditions and were diminished in AF.[43] Additionally, previous expression profiling

of human hearts identified several ion channels that were sex-differentially expressed or

modified.[30,63] Therefore, future studies should integrate additional sex-dependent differences in

electrophysiology to extend our sex-specific models to investigate the sex-specific interactive role

of AP and Ca^{2+} processes in promoting AF and as therapeutic targets. Second, our current sex-

specific modeling framework incorporates the direct effectors of electrophysiology and Ca^{2+}

signaling. As previous studies involving animals uncovered sex-dependent differences in

upstream signals including cAMP/phosphodiesterase in the ventricles,[64,65] similar differences, if

confirmed in the atria, can be readily incorporated into our framework, as established in our atrial

signaling model.[28] Third, our current work solely focused on isolated cardiomyocytes. Previous

studies associated female atria with greater fibrosis[44,66] as compared to male atria. Thus, the

increased incidence of Ca^{2+}-dependent triggers in female atria as revealed by our study may be

exacerbated in tissue through mismatched source-sink relation to trigger spontaneous APs and/or

subthreshold delayed afterdepolarizations. The latter could promote tissue dispersion of cell

excitability due to heterogeneous availability in fast Na^+ current due to heterogeneous voltage

and Ca^{2+}-dependent signaling,[28] thereby causing functional conduction block that promotes

reentry.[67] The implications for spontaneous APs and/or subthreshold delayed afterdepolarizations

warrant future investigations using sex-specific tissue models of human atria.

Conclusions

In summary, we have built a family of male- and female-specific 3D models of human atrial

cardiomyocytes in both nSR and AF conditions. Our analyses demonstrated increased propensity

to developing Ca^{2+}-driven arrhythmias in female vs male atrial cardiomyocytes with AF, and

suggested Ca^{2+}-targeted interventions as promising strategies for female AF patients. Our study

advocates for more inclusion of female subjects and provides new sex-specific computational

models as useful means to define sex-specific disease mechanisms and inform the development

of sex-specific therapeutics, ultimately improving effectiveness and equity of AF management.

706 Funding and Acknowledgements

Funding: This work was supported by American Heart Association Postdoctoral Fellowships

20POST35120462 (H.N.), 24POST1195229 (Y.W.), and Predoctoral Fellowship

20PRE35120465 (X.Z.); National Institutes of Health/NHLBI Grants R01HL131517 (E.G. and

D.D.), R01HL170521 (E.G., H.N.), R01HL141214 and P01HL141084 (E.G.), R01HL136389

(D.D.), R01HL089598 (D.D.), R01HL163277 (D.D.), R01HL160992 (D.D.), R01HL165704

(D.D.), R00HL138160 (S.M.); NIH Stimulating Peripheral Activity to Relieve Conditions Grant

1OT2OD026580-01 (E.G.); UC Davis School of Medicine Dean's Fellow Award (E.G.); the

European Union (large-scale integrative project MAESTRIA, No. 965286, D.D.); Burroughs

Wellcome Fund - Doris Duke Charitable Foundation "COVID-19 Fund to Retain Clinical

Scientists" Award (S.M.)

Conflict of Interest: None declared.

References

1. Kornej J, Börschel CS, Benjamin EJ, Schnabel RB. Epidemiology of Atrial Fibrillation in the 21st Century: Novel Methods and New Insights. *Circ Res* 2020;**127**:4–20.

2. Alonso A, Bengtson LGS. A rising tide: the global epidemic of atrial fibrillation. *Circulation* 2014;**129**:829–830.

3. Nattel S, Sager PT, Hüser J, Heijman J, Dobrev D. Why translation from basic discoveries to clinical applications is so difficult for atrial fibrillation and possible approaches to improving it. *Cardiovasc Res* 2021;**117**:1616–1631.

4. Turagam MK, Musikantow D, Whang W, Koruth JS, Miller MA, Langan M-N, Sofi A, Choudry S, Dukkipati SR, Reddy VY. Assessment of Catheter Ablation or Antiarrhythmic Drugs for First-line Therapy of Atrial Fibrillation. *JAMA Cardiol* 2021;**6**:1–9.

5. Odening KE, Deiß S, Dilling-Boer D, Didenko M, Eriksson U, Nedios S, Ng FS, Roca Luque I, Sanchez Borque P, Vernooy K, Wijnmaalen AP, Yorgun H. Mechanisms of sex differences in atrial fibrillation: role of hormones and differences in electrophysiology, structure, function, and remodelling. *EP Eur* 2019;**21**:366–376.

6. Tadros R, Ton A-T, Fiset C, Nattel S. Sex Differences in Cardiac Electrophysiology and Clinical Arrhythmias: Epidemiology, Therapeutics, and Mechanisms. *Can J Cardiol* 2014;**30**:783–792.

7. Andrade JG, Deyell MW, Lee AYK, Macle L. Sex Differences in Atrial Fibrillation. *Can J Cardiol* 2018;**34**:429–436.

8. Magnussen Christina, Niiranen Teemu J., Ojeda Francisco M., Gianfagna Francesco, Blankenberg Stefan, Njølstad Inger, Vartiainen Erkki, Sans Susana, Pasterkamp Gerard, Hughes Maria, Costanzo Simona, Donati Maria Benedetta, Jousilahti Pekka, Linneberg Allan, Palosaari Tarja, de Gaetano Giovanni, Bobak Martin, den Ruijter Hester M., Mathiesen Ellisiv, Jørgensen Torben, Söderberg Stefan, Kuulasmaa Kari, Zeller Tanja, Iacoviello Licia, Salomaa Veikko, Schnabel Renate B. Sex Differences and Similarities in Atrial Fibrillation Epidemiology, Risk Factors, and Mortality in Community Cohorts. *Circulation* 2017;**136**:1588–1597.

9. Gillis AM. Atrial Fibrillation and Ventricular Arrhythmias: Sex Differences in Electrophysiology, Epidemiology, Clinical Presentation, and Clinical Outcomes. *Circulation* 2017;**135**:593–608.

10. Ko D, Rahman F, Schnabel RB, Yin X, Benjamin EJ, Christophersen IE. Atrial fibrillation in women: epidemiology, pathophysiology, presentation, and prognosis. *Nat Rev Cardiol* 2016;**13**:321–332.

11. Vitale C, Fini M, Spoletini I, Lainscak M, Seferovic P, Rosano GM. Under-representation of elderly and women in clinical trials. *Int J Cardiol* 2017;**232**:216–221.

756   12. Fülöp L, Bányász T, Szabó G, Tóth IB, Bíró T, Lôrincz I, Balogh Á, Petô K, Mikó I, Nánási
757   PP. Effects of sex hormones on ECG parameters and expression of cardiac ion channels
758   in dogs. *Acta Physiol* 2006;**188**:163–171.

759   13. Xiao L, Zhang L, Han W, Wang Z, Nattel S. Sex-based transmural differences in cardiac
760   repolarization and ionic-current properties in canine left ventricles. *Am J Physiol-Heart Circ*
761   *Physiol* 2006;**291**:H570–H580.

762   14. James AF, Arberry LA, Hancox JC. Gender-related differences inventricular myocyte
763   repolarization in the guineapig. *Basic Res Cardiol* 2004;**99**:183–192.

764   15. Trépanier-Boulay Véronique, St-Michel Chantale, Tremblay Annie, Fiset Céline. Gender-
765   Based Differences in Cardiac Repolarization in Mouse Ventricle. *Circ Res* 2001;**89**:437–
766   444.

767   16. Saito Tomoaki, Ciobotaru Andrea, Bopassa Jean Chrisostome, Toro Ligia, Stefani Enrico,
768   Eghbali Mansoureh. Estrogen Contributes to Gender Differences in Mouse Ventricular
769   Repolarization. *Circ Res* 2009;**105**:343–352.

770   17. Fogli Iseppe A, Ni H, Zhu S, Zhang X, Coppini R, Yang P-C, Srivatsa U, Clancy CE,
771   Edwards AG, Morotti S, Grandi E. Sex-Specific Classification of Drug-Induced Torsade de
772   Pointes Susceptibility Using Cardiac Simulations and Machine Learning. *Clin Pharmacol*
773   *Ther* 2021;**110**:380–391.

774   18. Heijman J, Voigt N, Nattel S, Dobrev D. Cellular and Molecular Electrophysiology of Atrial
775   Fibrillation Initiation, Maintenance, and Progression. *Circ Res* 2014;**114**:1483–1499.

776   19. Nattel S, Dobrev D. The multidimensional role of calcium in atrial fibrillation
777   pathophysiology: mechanistic insights and therapeutic opportunities. *Eur Heart J*
778   2012;**33**:1870–1877.

779   20. Heijman J, Muna AP, Veleva T, Molina CE, Sutanto H, Tekook M, Wang Q, Abu-Taha IH,
780   Gorka M, Künzel S, El-Armouche A, Reichenspurner H, Kamler M, Nikolaev V, Ravens U,
781   Li N, Nattel S, Wehrens XHT, Dobrev D. Atrial Myocyte NLRP3/CaMKII Nexus Forms a
782   Substrate for Postoperative Atrial Fibrillation. *Circ Res* 2020;**127**:1036–1055.

783   21. Molina CE, Abu-Taha IH, Wang Q, Roselló-Díez E, Kamler M, Nattel S, Ravens U,
784   Wehrens XHT, Hove-Madsen L, Heijman J, Dobrev D. Profibrotic, Electrical, and Calcium-
785   Handling Remodeling of the Atria in Heart Failure Patients With and Without Atrial
786   Fibrillation. *Front Physiol* 2018;**9**:1383.

787   22. Nattel S, Heijman J, Zhou L, Dobrev D. Molecular Basis of Atrial Fibrillation
788   Pathophysiology and Therapy: A Translational Perspective. *Circ Res* 2020;**127**:51–72.

789   23. Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, Sun Q, Wieland T, Ravens U,
790   Nattel S, Wehrens XHT, Dobrev D. Enhanced sarcoplasmic reticulum Ca2+ leak and
791   increased Na+-Ca2+ exchanger function underlie delayed afterdepolarizations in patients
792   with chronic atrial fibrillation. *Circulation* 2012;**125**:2059–2070.

24. Bers DM. Cardiac excitation–contraction coupling. *Nature* 2002;**415**:198–205.

25. Schotten U, Haan S de, Verheule S, Harks EGA, Frechen D, Bodewig E, Greiser M, Ram
R, Maessen J, Kelm M, Allessie M, Wagoner DRV. Blockade of atrial-specific K+-currents
increases atrial but not ventricular contractility by enhancing reverse mode Na+/Ca2+-
exchange. *Cardiovasc Res* 2007;**73**:37–47.

26. Voigt N, Heijman J, Wang Q, Chiang DY, Li N, Karck M, Wehrens XHT, Nattel S, Dobrev
D. Cellular and molecular mechanisms of atrial arrhythmogenesis in patients with
paroxysmal atrial fibrillation. *Circulation* 2014;**129**:145–156.

27. Beavers DL, Wang W, Ather S, Voigt N, Garbino A, Dixit SS, Landstrom AP, Li N, Wang Q,
Olivotto I, Dobrev D, Ackerman MJ, Wehrens XHT. Mutation E169K in junctophilin-2
causes atrial fibrillation due to impaired RyR2 stabilization. *J Am Coll Cardiol*
2013;**62**:2010–2019.

28. Ni H, Morotti S, Zhang X, Dobrev D, Grandi E. Integrative human atrial modelling unravels
interactive protein kinase A and Ca2+/calmodulin-dependent protein kinase II signalling as
key determinants of atrial arrhythmogenesis. *Cardiovasc Res* 2023;**119**:2294–2311.

29. Grammatika Pavlidou N, Dobrev S, Beneke K, Reinhardt F, Pecha S, Jacquet E, Abu-Taha
IH, Schmidt C, Voigt N, Kamler M, Schnabel RB, Baczkó I, Garnier A, Reichenspurner H,
Nikolaev VO, Dobrev D, Molina CE. Phosphodiesterase 8 governs cAMP/PKA-dependent
reduction of L-type calcium current in human atrial fibrillation: a novel arrhythmogenic
mechanism. *Eur Heart J* 2023;**44**:2483–2494.

30. Herraiz-Martínez A, Tarifa C, Jiménez-Sábado V, Llach A, Godoy-Marín H, Colino-Lage H,
Nolla-Colomer C, Casabella-Ramon S, Izquierdo-Castro P, Benítez I, Benítez R, Roselló-
Díez E, Rodríguez-Font E, Viñolas X, Ciruela F, Cinca J, Hove-Madsen L. Influence of sex
on intracellular calcium homoeostasis in patients with atrial fibrillation. *Cardiovasc Res*
2022;**118**:1033–1045.

31. Yue X, Zhang R, Kim B, Ma A, Philipson KD, Goldhaber JI. Heterogeneity of transverse-
axial tubule system in mouse atria: Remodeling in atrial-specific Na+–Ca2+ exchanger
knockout mice. *J Mol Cell Cardiol* 2017;**108**:50–60.

32. Winters J, Isaacs A, Zeemering S, Kawczynski M, Maesen B, Maessen J, Bidar E,
Boukens B, Hermans B, Hunnik A van, Casadei B, Fabritz L, Chua W, Sommerfeld L,
Guasch E, Mont L, Batlle M, Hatem S, Kirchhof P, Wakili R, Sinner M, Stoll M, Goette A,
Verheule S, Schotten U. Heart Failure, Female Sex, and Atrial Fibrillation Are the Main
Drivers of Human Atrial Cardiomyopathy: Results From the CATCH ME Consortium. *J Am*
Heart Assoc 2023;**12**:e031220.

33. Thibault S, Ton A-T, Huynh F, Fiset C. Connexin Lateralization Contributes to Male
Susceptibility to Atrial Fibrillation. *Int J Mol Sci* 2022;**23**:10696.

34. Christ T, Boknik P, Wöhrl S, Wettwer E, Graf EM, Bosch RF, Knaut M, Schmitz W, Ravens
U, Dobrev D. L-Type Ca2+ Current Downregulation in Chronic Human Atrial Fibrillation Is
Associated With Increased Activity of Protein Phosphatases. *Circulation* 2004;**110**:2651–
2657.

35. Bosch RF, Zeng X, Grammer JB, Popovic K, Mewis C, Kühlkamp V. Ionic mechanisms of
electrical remodeling in human atrial fibrillation. *Cardiovasc Res* 1999;**44**:121–131.

36. Zhang X, Ni H, Morotti S, Smith CER, Sato D, Louch WE, Edwards AG, Grandi E.
Mechanisms of spontaneous Ca2+ release-mediated arrhythmia in a novel 3D human
atrial myocyte model: I. Transverse-axial tubule variation. *J Physiol* 2023;**601**:2655–2683.

37. Zhang X, Smith CER, Morotti S, Edwards AG, Sato D, Louch WE, Ni H, Grandi E.
Mechanisms of spontaneous Ca2+ release-mediated arrhythmia in a novel 3D human
atrial myocyte model: II. Ca2+-handling protein variation. *J Physiol* 2023;**601**:2685–2710.

38. Colman MA, Alvarez-Lacalle E, Echebarria B, Sato D, Sutanto H, Heijman J. Multi-Scale
Computational Modeling of Spatial Calcium Handling From Nanodomain to Whole-Heart:
Overview and Perspectives. *Front Physiol* 2022;**13**:836622.

39. Song Z, Ko CY, Nivala M, Weiss JN, Qu Z. Calcium-Voltage Coupling in the Genesis of
Early and Delayed Afterdepolarizations in Cardiac Myocytes. *Biophys J* 2015;**108**:1908–
1921.

40. Lenaerts I, Bito V, Heinzel FR, Driesen RB, Holemans P, D'hooge J, Heidbüchel H, Sipido
KR, Willems R. Ultrastructural and Functional Remodeling of the Coupling Between Ca2+
Influx and Sarcoplasmic Reticulum Ca2+ Release in Right Atrial Myocytes From
Experimental Persistent Atrial Fibrillation. *Circ Res* 2009;**105**:876–885.

41. Neef S, Dybkova N, Sossalla S, Ort KR, Fluschnik N, Neumann K, Seipelt R, Schöndube
FA, Hasenfuss G, Maier LS. CaMKII-Dependent Diastolic SR Ca2+ Leak and Elevated
Diastolic Ca2+ Levels in Right Atrial Myocardium of Patients With Atrial Fibrillation. *Circ*
Res 2010;**106**:1134–1144.

42. Harris CR, Millman KJ, Walt SJ van der, Gommers R, Virtanen P, Cournapeau D, Wieser
E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, Kerkwijk MH van, Brett M,
Haldane A, Río JF del, Wiebe M, Peterson P, Gérard-Marchant P, Sheppard K, Reddy T,
Weckesser W, Abbasi H, Gohlke C, Oliphant TE. Array programming with NumPy. *Nature*
2020;**585**:357–362.

43. Pecha S, Ismaili D, Geelhoed B, Knaut M, Reichenspurner H, Eschenhagen T, Schnabel
RB, Christ T, Ravens U. Resting membrane potential is less negative in trabeculae from
right atrial appendages of women, but action potential duration does not shorten with age.
J Mol Cell Cardiol 2023;**176**:1–10.

44. Wong GR, Nalliah CJ, Lee G, Voskoboinik A, Chieng D, Prabhu S, Parameswaran R,
Sugumar H, Al-Kaisey A, McLellan A, Ling L-H, Sanders P, Kistler PM, Kalman JM. Sex-
Related Differences in Atrial Remodeling in Patients With Atrial Fibrillation: Relationship to
Ablation Outcomes. *Circ Arrhythm Electrophysiol* 2022;**15**:e009925.

45. Kim K, Blackwell DJ, Yuen SL, Thorpe MP, Johnston JN, Cornea RL, Knollmann BC. The
selective RyR2 inhibitor ent-verticilide suppresses atrial fibrillation susceptibility caused by
Pitx2 deficiency. *J Mol Cell Cardiol* 2023;**180**:1–9.

46. Pabel S, Mustroph J, Stehle T, Lebek S, Dybkova N, Keyser A, Rupprecht L, Wagner S,
Neef S, Maier LS, Sossalla S. Dantrolene reduces CaMKIIδC-mediated atrial arrhythmias.
EP Eur 2020;**22**:1111–1118.

47. George SA, Brennan-McLean JA, Trampel KA, Rytkin E, Faye NR, Knollmann BC, Efimov
IR. Ryanodine receptor inhibition with acute dantrolene treatment reduces arrhythmia
susceptibility in human hearts. *Am J Physiol-Heart Circ Physiol* 2023.

48. Hartmann N, Pabel S, Herting J, Schatter F, Renner A, Gummert J, Schotola H, Danner
BC, Maier LS, Frey N, Hasenfuss G, Fischer TH, Sossalla S. Antiarrhythmic effects of
dantrolene in human diseased cardiomyocytes. *Heart Rhythm* 2017;**14**:412–419.

49. Hwang HS, Hasdemir C, Laver D, Mehra D, Turhan K, Faggioni M, Yin H, Knollmann BC.
Inhibition of cardiac Ca2+ release channels (RyR2) determines efficacy of class I
antiarrhythmic drugs in catecholaminergic polymorphic ventricular tachycardia. *Circ*
Arrhythm Electrophysiol 2011;**4**:128–135.

50. Kryshtal Dmytro O., Blackwell Daniel J., Egly Christian L., Smith Abigail N., Batiste
Suzanne M., Johnston Jeffrey N., Laver Derek R., Knollmann Bjorn C. RYR2 Channel
Inhibition Is the Principal Mechanism of Flecainide Action in CPVT. *Circ Res*
2021;**128**:321–331.

51. Wybraniec MT, Maciąg A, Miśkowiec D, Ceynowa-Sielawko B, Balsam P, Wójcik M,
Wróbel W, Farkowski M, Ćwiek-Rębowska E, Szołkiewicz M, Ozierański K, Błaszczyk R,
Bula K, Dembowski T, Peller M, Krzowski B, Wyganowska-Kapryan A, Wańha W, Koziński
M, Kasprzak JD, Szwed H, Mizia-Stec K. Antiarrhythmic drugs for pharmacological
cardioversion of atrial fibrillation and sex differences: Insights from the CANT II Study. *Pol*
Heart J Kardiologia Pol 2023;**81**:1089–1095.

52. Park YJ, Park J-W, Yu HT, Kim T-H, Uhm J-S, Joung B, Lee M-H, Pak H-N. Sex difference
in atrial fibrillation recurrence after catheter ablation and antiarrhythmic drugs. *Heart*
2023;**109**:519–526.

53. Tadini-Buoninsegni F, Smeazzetto S, Gualdani R, Moncelli MR. Drug Interactions With the
Ca2+-ATPase From Sarco(Endo)Plasmic Reticulum (SERCA). *Front Mol Biosci* 2018;**5**:36.

54. Eisner D, Caldwell J, Trafford A. Sarcoplasmic Reticulum Ca-ATPase and Heart Failure 20
Years Later. *Circ Res* 2013;**113**:958–961.

55. Reynolds JO, Chiang DY, Wang W, Beavers DL, Dixit SS, Skapura DG, Landstrom AP,
Song L-S, Ackerman MJ, Wehrens XHT. Junctophilin-2 is necessary for T-tubule
maturation during mouse heart development. *Cardiovasc Res* 2013;**100**:44–53.

56. Li J, Agvanian S, Zhou K, Shaw RM, Hong T. Exogenous Cardiac Bridging Integrator 1
Benefits Mouse Hearts With Pre-existing Pressure Overload-Induced Heart Failure. *Front*
Physiol 2020;**11**.

57. Liu Y, Zhou K, Li J, Agvanian S, Caldaruse A-M, Shaw S, Hitzeman TC, Shaw RM, Hong
T. In Mice Subjected to Chronic Stress, Exogenous cBIN1 Preserves Calcium-Handling
Machinery and Cardiac Function. *JACC Basic Transl Sci* 2020;**5**:561–578.

58. Perdreau-Dahl H, Lipsett DB, Frisk M, Kermani F, Carlson CR, Brech A, Shen X, Bergan-
Dahl A, Hou Y, Tuomainen T, Tavi P, Jones PP, Lunde M, Wasserstrom JA, Laporte J,
Ullrich ND, Christensen G, Morth JP, Louch WE. BIN1, Myotubularin, and Dynamin-2
Coordinate T-Tubule Growth in Cardiomyocytes. *Circ Res* 2023;**132**:e188–e205.

59.　Li J, Richmond B, Cluntun AA, Bia R, Walsh MA, Shaw K, Symons JD, Franklin S, Rutter
J, Funai K, Shaw RM, Hong T. Cardiac gene therapy treats diabetic cardiomyopathy and
lowers blood glucose. *JCI Insight* 2023;**8**.

60.　Lawless M, Caldwell JL, Radcliffe EJ, Smith CER, Madders GWP, Hutchings DC, Woods
LS, Church SJ, Unwin RD, Kirkwood GJ, Becker LK, Pearman CM, Taylor RF, Eisner DA,
Dibb KM, Trafford AW. Phosphodiesterase 5 inhibition improves contractile function and
restores transverse tubule loss and catecholamine responsiveness in heart failure. *Sci Rep*
2019;**9**:6801.

61.　Flórez-Vargas O, Brass A, Karystianis G, Bramhall M, Stevens R, Cruickshank S, Nenadic
G. Bias in the reporting of sex and age in biomedical research on mouse models. Mungall
C, ed. *eLife* 2016;**5**:e13615.

62.　Tobb K, Kocher M, Bullock-Palmer RP. Underrepresentation of women in cardiovascular
trials- it is time to shatter this glass ceiling. *Am Heart J Plus Cardiol Res Pract*
2022;**13**:100109.

63.　Ambrosi CM, Yamada KA, Nerbonne JM, Efimov IR. Gender Differences in
Electrophysiological Gene Expression in Failing and Non-Failing Human Hearts. *PLOS*
ONE 2013;**8**:e54635.

64.　Parks RJ, Ray G, Bienvenu LA, Rose RA, Howlett SE. Sex differences in SR Ca2+ release
in murine ventricular myocytes are regulated by the cAMP/PKA pathway. *J Mol Cell*
Cardiol 2014;**75**:162–173.

65.　Caldwell JL, Lee I-J, Ngo L, Wang L, Bahriz S, Xu B, Bers DM, Navedo MF, Bossuyt J,
Xiang YK, Ripplinger CM. Whole-heart multiparametric optical imaging reveals sex-
dependent heterogeneity in cAMP signaling and repolarization kinetics. *Sci Adv*
2023;**9**:eadd5799.

66.　Akoum N, Mahnkopf C, Kholmovski EG, Brachmann J, Marrouche NF. Age and sex
differences in atrial fibrosis among patients with atrial fibrillation. *EP Eur* 2018;**20**:1086–
1092.

67.　Landstrom AP, Dobrev D, Wehrens XHT. Calcium Signaling and Cardiac Arrhythmias. *Circ*
Res 2017;**120**:1969–1993.