Alanine and glutamate catabolism ensure proper sporulation by preventing premature germination and providing energy respectively

Fengzhi Lyu, Tianyu Zhang, Dong Yang, Lei Rao*, Xiaojun Liao*

College of Food Science and Nutritional Engineering, National Engineering Research Center for Fruit and Vegetable Processing, Key Laboratory of Fruit and Vegetable Processing of Ministry of Agriculture and Rural Affairs, Beijing Key Laboratory for Food Non-Thermal Processing, China Agricultural University, Beijing, China

*Correspondence

Lei Rao, College of Food Science and Nutritional Engineering, China Agricultural University, 100083, Beijing,China.

Email: rao.lei@cau.edu.cn

Xiaojun Liao, College of Food Science and Nutritional Engineering, China Agricultural University, 100083, Beijing,China.

Email: liaoxjun@cau.edu.cn

Abstract

Sporulation as a typical bacterial differentiation process has been studied for decades. However, two crucial aspects of sporulation, (i) the energy sources supporting the process, and (ii) the maintenance of spore dormancy throughout sporulation, are scarcely explored. Here, we reported the crucial role of RocG-mediated glutamate catabolism in regulating mother cell lysis, a critical step for successful sporulation, likely by providing energy metabolite ATP. Notably, *rocG* overexpression resulted in an excessive ATP accumulation in sporulating cells, leading to adverse effects on future spore properties, e.g. increased germination efficiency, reduced DPA content, and lowered heat resistance. Additionally, we revealed that Ald-mediated alanine metabolism decreased the typical germinant L-alanine concentration in sporulating environment, thereby preventing premature germination and maintaining spore dormancy. Our data inferred that sporulation was a highly orchestrated biological process requiring a delicate balance in diverse metabolic pathways, hence ensuring both the completion of sporulation and production of high-quality spores.

Introduction

Spores are generated by spore-forming bacteria such as the orders *Bacillales* and *Clostridiales* in response to unfavorable environmental conditions, such as nutrient limitation (1, 2). Spores are metabolically dormant and considered the most resilient living organisms due to their extreme resistance to harsh environments, and they are capable of surviving for millions of years (3-5). The process of forming spores from bacterial vegetative cells is termed sporulation. Taking the model bacterium *Bacillus subtilis* as an example, the morphological process of sporulation can be divided into several stages: asymmetric division, engulfment, spore maturation, mother cell lysis, and spore release (6, 7). Initially, as vegetative cells commit to sporulation, the earliest visible event is asymmetric division, which produces the septum, dividing the vegetative cell into a larger mother cell and a smaller forespore. Subsequently, the mother cell membrane migrates around the forespore until it is completely enclosed. This phagocytosis-like process is identified as engulfment. Concurrently, the double-membrane structure of the forespore forms, followed by cortex synthesis and spore coat assembly. Then, the forespore chromosomes become saturated with small acid-soluble proteins (SASPs), and the water within the forespore is replaced by dipicolinic acid (DPA) synthesized in the mother cell, resulting in forespore dehydration. These events culminate in the appearance of phase-bright spores. Next, mother cell lysis occurs after spore maturation, allowing the spores to be released into the environment.

Research on the morphological events of the sporulation process, as well as the underlying gene expression and molecular mechanisms, has been continued for decades (6, 8-11). However, relative few studies focus on guaranteeing proper sporulation, especially concerning the quantity and quality of spores. Proper sporulation encompasses two essential aspects. The

first is the normal progression of sporulation. Sporulation is recognized as an energy-intensive biological process (12), implying the importance of energy supply in promoting its progress. Studies have suggested that amino acid metabolism, such as glutamate and alanine catabolism, may serve as potential energy sources driving sporulation (13-15). However, the mechanisms regulating these processes during sporulation remain unclear. Secondly, proper sporulation needs the maintenance of spore dormancy throughout the process. Previous studies report that the deletion of *ylbJ*, *pdaB*, or genes encoding SpoVA protein results in premature germination, indicating the loss of dormancy maintenance ability of generated spores during sporulation (16, 17). The reasons for these types of premature germination are varied, including inappropriate activation of germination receptors, incorrect assembly of spore outer structures, and deficiencies in the SpoVA channel (16, 17). Except for these intrinsic factors, quantities of external factors such as nutrient germinants that induce germination may also exist around the generated spores. How spores maintain dormant in such a tempting environment remains a mystery to be explored. Therefore, defects in either energy supply or dormancy maintenance can lead to abnormal sporulation, adversely affecting the quantity or quality of spores produced.

Here, we reported that the RocG-mediated glutamate catabolism played a crucial role in ensuring proper sporulation, particularly by promoting mother cell lysis through providing energy support. Our research further demonstrated that overexpression of *rocG* resulted in excessively high ATP contents in sporulating cells, which adversely affected the properties of the resulting spores, e.g. elevated germination efficiency, reduced DPA content, and lowered heat resistance. Moreover, we revealed that Ald-mediated alanine catabolism decreased the concentration of typical germinant L-alanine in the sporulating environment to a certain level. This regulation effectively prevented premature germination and contributed to maintain spore dormancy

throughout the sporulation process.

Results

Proteins involved in alanine, aspartate and glutamate metabolism show

enrichment according to proteomics analysis during sporulation of *B.*

subtilis

In order to explore the essential metabolic pathways in *B. subtilis* sporulation,

Tandem Mass Tag-based (TMT) quantitative proteomics analysis was

conducted comparing dormant spores (DS) and sporulating vegetative cells

(VC) at t_0 (Figure 1A). Hierarchical clustering analysis (HCA) was employed to

illustrate the overall differences in proteins between the DS and VC groups

(Figure 1B). The HCA heatmap displayed greater differences between groups

than within groups, indicating significant differences in proteins between DS

and VC. Differentially-expressed proteins were identified based on the criteria

of $p < 0.05$ and fold change > 1.2 (the expression level increased by more

than 1.2-fold or decreased by less than 0.83-fold). 1,259 proteins with

decreased expression as well as 1,248 proteins with increased expression

were screened out in the DS group (Figure 1C). KEGG pathway enrichment

analysis of these differentially expressed proteins revealed the significant

alterations in several metabolic pathways between the DS and VC groups,

including alanine, aspartate and glutamate metabolism, ribosome, flagellar

assembly, glyoxylate and dicarboxylate metabolism, and methane metabolism

(Figure 1D). Of these pathways, the most significant changes were observed

in alanine, aspartate and glutamate metabolism, implying their close

association with the sporulation of *B. subtilis*. Previous studies have indicated

that *ald*, encoding alanine dehydrogenase Ald, and *rocG*, encoding glutamate

dehydrogenase RocG, are crucial regulators of alanine and glutamate

metabolism, respectively (14, 18). In addition, Δ*ald* and Δ*rocG* mutants have

shown notable defects in sporulation (14, 15). However, the deletion of *ansB*

gene, encoding L-aspartase important for aspartate metabolism, has no

significant effect on sporulation (19). Consequently, further exploration was

conducted in the following work to elucidate the effects of alanine and

glutamate metabolism on sporulation.

Alanine and glutamate metabolism collaboratively regulate sporulation

with an additive effect

Given that alanine and glutamate metabolism have been reported separately

to be involved in sporulation (14, 15, 20), we wonder if there are any potential

joint impacts of these two pathways on sporulation. We constructed the Δ*ald*

Δ*rocG* mutant and observed a significantly lower number of phase-bright

spores compared to the Δ*ald* or Δ*rocG* mutants (Figure 2A). This indicated

that the sporulation defect in the Δ*ald* Δ*rocG* mutant was more pronounced

than in either the Δ*ald* or Δ*rocG* mutants. This was further demonstrated by

examining the heat-resistant spores produced in sporulation, as the

percentage of the Δ*ald* and Δ*rocG* spores was 10.9% and 29.8% respectively,

while the Δ*ald* Δ*rocG* mutant was only 0.3% (Figure 2B). The severe

sporulation defect of the double mutant suggested that Ald and RocG jointly

regulated sporulation in an additive manner. Additionally, we observed an

accumulation of phase-dark forespores in the Δ*ald* Δ*rocG* mutant (Figure 2A),

which could be attributed to two factors: (i) premature germination due to

abnormal spore structure assembly or an inappropriate in-situ sporulating

environment (16) , or (ii) failure of spore maturation due to limited energy

supporting sporulation progression. We first deleted *gerAA* to investigate if

premature germination occurred in the Δ*ald* Δ*rocG* mutant. As shown in

Figure 3A, the sporulation defect of the double mutant was partially rescued,

with 29.7% of phase-bright spores formed in the Δ*ald* Δ*rocG* Δ*gerAA* (Δ3)

mutant. This result indicated that premature germination indeed existed in the

Δ*ald* Δ*rocG* mutant. However, the partial rescue effect suggested that
premature germination was not the exclusive reason for sporulation defect in
the Δ*ald* Δ*rocG* mutant. Therefore, the limitation of energy support remained a
substantial explanation for the sporulation defect phenotype observed in the
double mutant strain. Nonetheless, we further explored these two possibilities
in the following work.

RocG-mediated glutamate metabolism, rather than Ald-mediated alanine
metabolism, is essential for ensuring both the sporulation efficiency and
the spore quality, likely through energy supply

As the Ald and RocG mediated alanine and glutamate metabolisms were
proposed as potential energy sources for sporulation (14, 15), we
hypothesized that these two metabolic pathways contribute to energy support
independently. If this was true, excessive complementation of either metabolic
pathway should be capable of rescuing the sporulation defect observed in the
Δ*ald* Δ*rocG* mutant. Here, we used the Δ*ald* Δ*rocG* Δ*gerAA* (Δ3) strain to
exclude the premature germination effect and independently explored the
energy supply mechanism (Figure 3A). Based on this, *ald* and *rocG* were
artificially expressed separately and jointly in Δ3 under an IPTG-inducible
promoter. Results indicated that increasing the expression level of *ald* by
raising the concentration of IPTG up to 5 mM had no significant effect on the
quantity of phase-bright spores in the Δ3 mutant, with the sporulation
percentage ranging between 20% and 30% (Figure S1, Figure S2A).
Accordingly, these spores exhibited significant germination deficiency under
AGFK induction (Figure S2B, Figure 3C). However, elevating the *rocG*
expression level with the addition of at least 10 mM IPTG restored the
sporulation of the Δ3 mutant to 53.4%, similar to that of the wild-type (56.8%)
(Figure 3B). Furthermore, when more than 20 mM IPTG was added, the
germination deficiency of the Δ3 mutant spores was also recovered to the

level of the wild-type (Figure S2B, Figure 3C). Notably, the Δ3 mutant with

IPTG-induced co-expression of *ald* and *rocG* exhibited the same sporulation

and germination phenotypes as those with IPTG-induced sole expression of

rocG (Figure S2B, Figure 3C). Hence, in the Δ3 mutant, sole complementation

of RocG succeeded in rescuing the sporulation defect, unlike in the case of

Ald. This indicated that RocG-mediated glutamate metabolism appeared to

regulate sporulation by providing energy sources, whereas Ald-mediated

alanine metabolism may not play the same role.

To further investigate whether these two catabolic pathways are involved in

controlling the spore quality, mutants of transcription factors of Ald and RocG,

Δ*adeR* and Δ*ahrC* Δ*rocR* (15, 21), were respectively constructed to examine

the germination phenotype of the spores. *gerAA* was also knocked out in

these mutants to ensure the comparability of experimental results. The

deletion of transcription factors was demonstrated to reduce, rather than

completely eliminate, the expression of regulated genes (Figure 4B), and this

rescued the sporulation deficiency (Figure 4A). Interestingly, no significant

germination defect was observed in spores of the Δ*adeR* Δ*gerAA* mutant with

decreased expression of *ald* (Figure 4C). However, spores of the Δ*ahrC*

Δ*rocR* Δ*gerAA* mutant with low expression of *rocG* showed a remarkable

germination deficiency (Figure 4C). Moreover, spores of the Δ*adeR* Δ*ahrC*

Δ*rocR* Δ*gerAA* mutant exhibited a similar germination deficiency phenotype to

that of Δ*ahrC* Δ*rocR* Δ*gerAA* mutant spores (Figure 4C), indicating that the

expression of *rocG*, rather than *ald*, was essential for ensuring spore quality

with normal germination capability. Taken together, these results strongly

implied that RocG-mediated glutamate metabolism regulated both sporulation

efficiency and spore quality, probably through energy supply. As for Ald-

mediated alanine metabolism, its effect on sporulation was unlikely to be

executed by providing energy sources. Instead, it is more reasonably

associated with premature germination, as quantities of phase-dark spores

were observed in the sporulating cells of the Δ*ald* mutant (Figure 2A).

Ald inhibits premature germination during sporulation by reducing L-
alanine content in the external environment of generating spores

As mentioned above, the presence of phase-dark spores in the Δ*ald* mutant
prompted us to investigate the impact of Ald-mediated alanine metabolism on
premature germination. As illustrated in Figure 5A-5B, the Δ*ald* mutant
produced significant quantities of phase-dark spores, and the percentage of
phase-bright spores was only 11.5%. However, upon the deletion of *gerAA* in
the Δ*ald* mutant, the percentage of phase-bright spores significantly raised to
61.5%, approaching levels observed in the wild-type (76.5%). In addition, no
notable deficiency in germination was observed in Δ*ald* Δ*gerAA* spores
compared to the wild-type (Figure 5C). Thus, it can be concluded that the
absence of Ald led to sporulation defect by inducing premature germination
during sporulation. To identify the timing of premature germination occurrence,
the sporulation process of the Δ*ald* mutant was examined by time-lapse
microscopy. Interestingly, two models were observed: (i) forespores
prematurely germinated during mother cell lysis, and then were released; (ii)
dormant spores were released and subsequently induced to premature
germination (Figure 5D).

The connection between Ald and premature germination raised up an
intriguing speculation that the interruption of alanine metabolism may lead to
an over-accumulation of L-alanine, which triggers premature germination. To
investigate this, *ald* was artificially expressed in the Δ*ald* mutant under an
IPTG-inducible promoter, and the sporulation phenotype as well as the
environmental L-alanine content at the late sporulation phase (t_{19}) of these
mutants were examined. The results showed a gradual increase in the
percentage of phase-bright spores with elevating the expression levels of *ald*

(Figure 6A-6C). Notably, the expression of *ald* decreased as the concentration

of added IPTG increased to 1 mM, which was possibly due to the toxicity of

IPTG to cells (Figure 6C). In contrast, an opposite trend was observed in the

environmental L-alanine concentration of the Δ*ald* mutant, which reached

3,397.4 μM without IPTG induction, while decreased to wild-type levels of

145.9 μM when > 200 μM IPTG was added to elevate the *ald* expression

(Figure 6B). Consequently, Ald-mediated alanine metabolism was responsible

for reducing the L-alanine content in the external environment of spores to

prevent premature germination, and thus ensuring proper sporulation.

RocG plays a crucial role in regulating both σ^K-dependent spore release

and spore properties likely by providing energy sources

As indicated above, RocG-mediated glutamate metabolism regulated both

sporulation efficiency and spore quality likely through energy support. To

further explore the underlying mechanism, we focused on identifying the

specific sporulation stage interrupted by *rocG* deletion. We constructed strains

capable of reporting stage-specific sigma factors involved in sporulation,

namely σ^F, σ^E, σ^G, and σ^K, which respectively regulated polar division,

engulfment, spore maturation, and spore release (6, 11, 22, 23). P$_{spoIIQ}$, P$_{spoIID}$,

P$_{sspB}$, and P$_{gerE}$ promotors that was respectively recognized by σ^F, σ^E, σ^G, and

σ^K were fused to *gfp*. In general, the activation of a particular σ factor

correlates with σ-dependent GFP fluorescence, thereby visually displaying the

impaired sporulation stage (22). Our results demonstrated that the activation

patterns of σ^F and σ^E in the Δ*rocG* mutant were similar to those of the wild-

type (Figure S3). The activation of σ^G could be achieved in the Δ*rocG* mutant,

though there was a delay in the activation timepoint (Figure 7A). Remarkably,

σ^K-dependent GFP fluorescence in the Δ*rocG* mutant initially appeared at t$_7$

but abnormally persisted until t$_{25}$ (Figure 7B). Since σ^K regulates cell wall lytic

enzymes, leading to mother cell lysis (24, 25), its fluorescence is supposed to

disappear with spore release. The persistence of σ^K-dependent GFP

fluorescence at the end of sporulation in the Δ*rocG* mutant indicated the

failure of mother cell lysis, a process mainly executed by the sporulation-

specific cell wall lytic enzymes CwlC and CwlH (24, 26). As expected, the

expression level of *cwlC* and *cwlH* were remarkably lower in the Δ*rocG* mutant

compared to those in the wild-type (Figure 7C). Taken together, the deletion of

rocG had no notable effect on the activation of sporulation-specific σ factors

but hindered mother cell lysis by impacting the expression of cell wall lytic

enzymes, resulting in an impaired spore release process (Figure 7D).

To further understand the regulating role of RocG in sporulation, *rocG* was

artificially expressed in the Δ*rocG* mutant under an IPTG-inducible promoter.

The results showed a significant increase in the percentage of released

spores with elevating expression levels of *rocG* (Figure 8A-8C). Moreover, the

addition of more than 10 μM IPTG remarkably improved the percentage of

spore release to the wild-type level (Figure 8A, 8C). As hypothesized

previously, RocG-mediated glutamate metabolism could provide energy

sources to drive sporulation proceeding, we then examined the ATP levels in

sporulating cells to verify this. Since previous research has demonstrated that

the ATP content of mother cells during sporulation peaked at t_1 (15, 27), we

examined ATP content at t_1 in the Δ*rocG* mutant with different *rocG* expression

levels. Accordingly, the level of ATP in the Δ*rocG* mutant increased with the

elevation of *rocG* expression (Figure 8D). The positive correlation between

the ATP level in sporulating cells (Figure 8D) and future spore release (Figure

8C) suggested the crucial role of energy supply in the late sporulation process,

particularly in mother cell lysis. Notably, the ATP content of the Δ*rocG* mutant

with 50 μM IPTG induction was almost double that of the wild-type (Figure 8D).

We then wondered if such a high level of ATP in sporulating cells could affect

the properties of the future spores. To test this, the spores generated under

different concentrations of IPTG induction were purified and examined for

germination phenotypes, as well as DPA content and heat resistance.

Interestingly, Δ*rocG* spores with 50 μM IPTG induction showed higher

germination efficiency (Figure 8G-8I) but significantly lower DPA content

(Figure 8E) as well as decreased heat resistance (Figure 8F) compared to the

wild-type spores. Taken together, the expression of *rocG* can indeed provide

energy support for sporulation, especially for mother cell lysis regulated by σ^K,

thus contributing to the proper sporulation process. However, overexpression

of *rocG* can accumulate excessive ATP in sporulating cells, which might

adversely affect the spore properties.

Discussion

Sporulation as a typical bacterial differentiation process has been extensively

studied for decades, and the morphological events along with the signal

transduction for this process are relatively well elucidated (6, 8-11). However,

as an energy-consuming process, the sources of energy supply and the

underlying regulating mechanism lack research. In addition, how the

generated spores maintain in dormant state during sporulation remains

mysterious. Here, we demonstrated that Ald-mediated alanine metabolism

decreased the concentration of the typical germinant L-alanine in the

sporulating environment to a certain level, thus avoiding premature

germination and maintaining spore dormancy. Moreover, we also provided

evidences supporting that RocG-mediated glutamate metabolism ensured

proper sporulation, especially mother cell lysis, by regulating ATP levels

during sporulation. Additionally, excessively high ATP levels during the

sporulation process was supposed to adversely affect the properties of the

produced spores, including faster germination efficiency, lower DPA content,

along with decreased heat resistance. Our data revealed that sporulation was

a highly orchestrated and exquisite biological process requiring the balance of

diverse metabolic pathways, e.g. alanine catabolism to eliminate surrounding

germinants, and glutamate metabolism providing an appropriate level of energy to ensure both sporulation completion and the high quality of generated spores (Figure 9).

Our finding of alanine catabolism eliminating the germinant L-alanine raises another open question that which catabolic pathways or biological reactions are responsible for regulating the balance of other potential germinants during sporulation. Indeed, sporulation involves protein turnover, in which new proteins are continuously synthesized using the amino acids derived from the breakdown of pre-existing cellular protein (28, 29). Hence, substantial free amino acids exist during sporulation, which could serve as potential germinants. Moreover, non-sporulating cells can produce meso-diaminopimelic acid (m-DAP) type muropeptides, also identified as possible germinants (30). However, the mechanisms by which spores eliminate these potential germinants remain unclear. Our research revealed that alanine catabolism is one of the strategies employed to achieve this. While we also observed that L-alanine was not completely eliminated, as 145.9 µM L-alanine was still detected in the sporulating medium of the wild-type (Figure 6B). Why did the presence of residual L-alanine in the environment not trigger germination? One plausible explanation is that alanine racemases present in the spore coat can convert the germinant L-alanine into the germination inhibitor D-alanine, thereby allowing the spores to persist in dormancy (31). However, whether there are alternative mechanisms preventing the residual L-alanine and other germinants from triggering germination is worth exploring.

Another new finding in our study revealed that RocG-mediated glutamate metabolism plays a crucial role as an energy source for sporulation. Indeed, the catabolic product of glutamate, 2-oxoglutarate (2-OG), directly participates in the tricarboxylic acid (TCA) cycle, showing its superior efficiency in providing energy during nutrient-limited sporulation (32, 33). Concurrently,

various amino acids such as proline, ornithine, citrulline, and arginine can be

converted into glutamate through the arginine degradation pathway (34),

indicating its high availability in sporulating cells. These aspects give

glutamate an advantage in supporting energy during sporulation compared to

other amino acids. Additionally, glutamate stands out among the limited free

amino acids found within diverse spores (35), further suggesting its crucial

role during sporulation. In our study, we showed that ATP produced by

glutamate catabolism is highly corelated with mother cell lysis regulated by

the cell wall lytic enzymes CwlC and CwlH, and the interruption of this

catabolism remarkably reduced the expression of these two enzymes. This

finding implies that the energy derived from glutamate catabolism is crucial for

the expression of genes regulating the final stage of sporulation. Moreover,

we also observed that the overaccumulation of ATP in sporulating cells

through glutamate catabolism adversely affected the properties of future

spores. These effects may be attributed to abnormal changes in the structure

assembly or molecular reservoir in the generated spores (28, 36). Actually,

spores can inherit molecules from sporulating cells, such as alanine

dehydrogenase and ATP, to modulate their revival capability (36). Moreover,

glutamate, as a universal amino group donor in all living organisms, can serve

as a carbon or nitrogen source for synthesizing other amino acids and DPA

during sporulation (34, 35, 37). Additionally, the glutamate catabolic

intermediate 2-OG can contribute to the synthesis of amino acids, nucleotides,

and NADH (32), potentially affecting the structure construction or molecular

modulation of spores. All these evidences support that glutamate is an ideal

substrate for energy supply and the synthesis of new substances during

sporulation, ensuring both the proper sporulation process and the quality of

the spores.

Methods

Strains and plasmids

B. subtilis strains used in this study are listed Table S1. Plasmids construction is listed in Table S2, and primers are described in Table S3. For gene replacement strategy, primer pairs were used to amplify the flanking genomic regions of the corresponding gene. PCR products and the respective antibiotic resistance gene were used for Gibson assembly (NEB, USA) (38). The product was used to transform *B. subtilis* PY79 to obtain the mutant allele.

General methods

All general methods for *B. subtilis* were carried out as described previously with some modifications (15). Cultures of wild-type and mutant strains were cultivated in LB medium (Difco) at 37 °C. Sporulation was carried out at 37°C by suspending overnight cells (OD_{600} = 0.05) in Schaeffer's liquid medium (Difco Sporulation Medium, DSM) (39). Sporulation t_0 was identified as the third hour after spores suspending in DSM. The percentage of sporulation was evaluated by calculating the ratio of total number of colonies forming units (CFU) before and after heat treatment (80°C, 20 min) (40). The percentage of phase-bright or released spores were counted based on the according phase-contrast images. To ensure confidence of the data, > 800 cells were counted for each experiment. Spore germination with different germinants was examined as described previously with some modifications (15, 41). Briefly, purified spores were heat activated at 75°C for 30 min, and then induced by L-Alanine (10 mM) or AGFK (2.5 mM L-Asparagine, 5 mg/mL D-glucose, 5 mg/mL D-fructose, and 50 mM KCl) at 37°C, or DDA (1 mM in 10 mM Tris-HCl, pH 7.4) at 42°C. The germination was tested by determining the DPA release as descried in the following text.

Spore purification

Matured spores were purified as described previously (15). Briefly, 22 hrs DSM culture was centrifuged and washed 3 times by DDW and then kept in 4°C with constant agitation. The suspension was washed once a day and resuspended in DDW. After 7 days, the suspension was centrifuged to collect the pellet. 20% histodenz solution was used to resuspend the pellet at a ratio of 400 μL per 10 mL of DSM for 30 min on ice. Aliquots (200 μL) of resuspension mixture were then added on top of 900 μL 50% histodenz solution, and gradient fractionation was carried out by centrifugation at 15,000 rpm at 4°C for 30 min. The pellet was collected and washed at least 5 times by DDW. Phase contrast microscopy was then used to evaluate the purity of pellet spores. Spores with >99% purity can be used for following experiments, otherwise the purification steps should be carried out more than once.

Tandem Mass Tag-based (TMT) quantitative proteomics analysis

TMT quantitative proteomics analysis was carried out between pure dormant spores (DS) and vegetative cells (VC) at sporulation t_0 by Shanghai Applied Protein Technology Co., Ltd (Shanghai, China). DS and VC samples were collected by centrifugation and then freeze-dried and bead-grinded using FastPrep-24 (M. P. Biomedicals, LLC, USA). Samples were then extracted for proteins and labeled using TMT reagent. Proteomics analysis was then carried out by LC-MS/MS system with on a Q Exactive mass spectrometer (Thermo Scientific) that was coupled to Easy nLC (Proxeon Biosystems, now Thermo Fisher Scientific). The raw data for each sample were searched using the MASCOT engine (Matrix Science, London, UK; version 2.2) embedded into Proteome Discoverer 1.4 software for identification and quantitation analysis. Hierarchical clustering analysis was performed using Cluster 3.0 (http://bonsai.hgc.jp/~mdehoon/software/cluster/software.htm) and Java

Treeview software (http://jtreeview.sourceforge.net). Enrichment analysis was

performed based on KEGG database (http://geneontology.org/). A statistical

analysis was performed using a t-test to determine the significance (p-value)

of differentially-expressed proteins. The expression level of proteins with $p <$

0.05 and fold change > 1.2 (the expression level increased by more than 1.2-

fold or decreased by less than 0.83-fold) were considered as significant

difference.

DPA measurements

DPA release was detected as described previously with some modifications

(42). Briefly, spore germination was induced by L-Ala, AGFK or DDA at 37°C

or 42°C in a 96-well plate. Spores at OD_{600} of 20, 10 mM germinants, 25 mM

K-Hepes buffer (pH 7.4) as well as 50 mM $TbCl_3$ were mixed in 200 μL and

Tb^{3+}-DPA fluorescence intensity was monitored at Ex/Em = 270/545 nm by a

TECAN Spark 10M microplate reader (TECAN, Switzerland). Total DPA

content of spores were evaluated by boiling the spores (OD_{600} of 1) for 20 min

and mixing the spores and 50 mM $TbCl_3$ to 200 μL in a 96-well plate. The DPA

standard solution was serially diluted and detected together to obtain a

standard curve. The detection parameters for DPA release were the same as

above, and the total DPA content of spores was calculated based on the

standard curve.

Phase-contrast and fluorescence microscopy

Phase-contrast and fluorescence microscopy were performed using a Nikon

DS-Qi2 microscope equipped with a Nikon Ph3 DL 100x/1.25 Oil phase

contrast objective. Both bacterial cells (500 μL) and spores (50 μL) were

centrifuged, and the pellets were resuspended with 5 ~ 10 μL PBSx1 and then

imaged. For time-lapse imaging of sporulation, Imaging System Cell Chamber

(AttofluorTM Cell Chamber) was used. Sporulating cells were collected by

centrifugation. The supernatant DSM was collected to make a gel-pad with

1% agarose. The collected sporulating cells were incubated on the DSM gel-

pad in a chamber at 37 °C. Image analysis and processing were performed by

ImageJ2.

Real-Time Quantitative PCR (RT-qPCR)

Real-time quantitative PCR (RT-qPCR) was carried out followed the protocol

described previously (15). RNA samples of sporulating cells (500 µL) were

collected from DSM by centrifugation, and then extracted by FastPure

Cell/Tissue Total RNA Isolation Kit V2 (Vazyme Biotech Co..Ltd). HiScript III

All in-one RT SuperMix for qPCR (Vazyme Biotech Co..Ltd) was used to

reverse transcribed RNA samples. RT-qPCR reactions were conducted with

PerfectStart Green qPCR SuperMix (TransGen Biotech Co., Ltd). CFX

Connect RealTime PCR Dection System (Bio-Rad Laboratories (Shanghai)

Co., Ltd.) was used to detect the fluorescence and *scr* gene was selected to

normalize sample data (zhou23). Each experiment was performed triplicate.

Environmental L-alanine content assay

Measurement of environmental L-Alanine level was performed as the

instruction of Amplite Fluorimetric L-Alanine Assay Kit (AAT Bioquest, Inc.).

DSM media of sporulating cells at sporulation t_{19} was collected from the

supernatant after centrifugation. The fluorescence intensity of L-Alanine in

DSM media was monitored by TECAN Spark 10M microplate reader (TECAN,

Switzerland) at Ex/Em = 540/590 nm. The standard solution of L-alanine was

serially diluted and detected together to obtain a standard curve, and the L-

Alanine content in the environment was calculated based on the standard

curve.

ATP content assay

Measurement of ATP content in mother cell was performed using the BacTiter-Glo Microbial Cell Viability Assay (Promega). As guided by the instructions, sporulating cells in DSM at sporulation t_1 were collected and detected luminescence using a TECAN Spark 10M microplate reader (TECAN, Switzerland). The standard solution of ATP was serially diluted and detected simultaneously to obtain a standard curve, and the ATP level was calculated based on the standard curve.

Data processing

Unless stated otherwise, each experiment was carried out at least triplicate. GraphPad Prism 8 software was used for all statistical analysis, data processing, and graph drawing. One-way ANOVA was performed to analyze the variance and $p < 0.05$ was regarded as significance for all data statistics.

Data availability

The data that support the findings of this study are available from the authors on reasonable request.

Acknowledgements

We are grateful to Dr. Bing Zhou (The Hebrew University of Jerusalem) for valuable discussions and comments. This work was supported by National Natural Science Foundation of China (NSFC) (grant No. 32372470), NSFC (grant No. 32001658), Agricultural Research Outstanding Talents of China (grant No. 13210317) awarded to Lei Rao, and 2115 Talent Development Program of China Agricultural University.

Author contributions

F. L. and L. R. conceived the idea of the experiment. F. L. contributed to the acquisition and analysis of data as well as writing the draft. T. Z. contributed to the construction of mutant strains. Y. D., L. R., and X. L. contributed to revising the paper.

Competing interests

The authors declare no competing interests.

References

1. Driks A. 2002. Maximum shields: the assembly and function of the bacterial spore coat. Trends Microbiol 10:251-254.

2. Stragier P, Losick R. 1996. Molecular genetics of sporulation in *Bacillus subtilis*. Annu Rev Genet 30:297-41.

3. Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. 2000. Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. Microbiol Mol Biol Rev 64:548-572.

4. Setlow P. 2006. Spores of *Bacillus subtilis*: their resistance to and killing by radiation, heat and chemicals. J Appl Microbiol 101:514-525.

5. Vreeland RH, Rosenzweig WD, Powers DW. 2000. Isolation of a 250 million-year-old halotolerant bacterium from a primary salt crystal. Nature 407:897-900.

6. Riley EP, Schwarz C, Derman AI, Lopez-Garrido J. 2020. Milestones in *Bacillus subtilis* sporulation research. Microb Cell 8:1-16.

7. Higgins D, Dworkin J. 2012. Recent progress in *Bacillus subtilis* sporulation. FEMS Microbiol Rev 36:131-148.

8. Tokuyasu K, Yamada E. 1959. Fine Structure of *Bacillus subtilis*: II. Sporulation Progress. The Journal of Cell Biology 5:129-133.

9. Kawata T, Inoue T, Takagi A. 1963. Electron microscopy of spore formation and germination in *Bacillus subtilis*. Japanese Journal of Microbiology 7:23-41.

10. Piggot P, Coote J. 1976. Genetic aspects of bacterial endospore formation. Bacteriological reviews 40:908-962.

11. Errington J. 2003. Regulation of endospore formation in *Bacillus subtilis*. Nat Rev Microbiol 1:117-126.

12. Phillips Z, Strauch M. 2002. *Bacillus subtilis* sporulation and stationary phase gene expression. Cell Mol Life Sci 59:392-402.

13. Charba J, Nakata H. 1977. Role of glutamate in the sporogenesis of

Bacillus cereus. J Bacteriol 130:242-248.

14.   Siranosian KJ, Ireton K, Grossman AD. 1993. Alanine dehydrogenase
(ald) is required for normal sporulation in *Bacillus subtilis*. J Bacteriol
175:6789-6796.

15.   Rao L, Zhou B, Serruya R, Moussaieff A, Sinai L, Ben-Yehuda S. 2022.
Glutamate catabolism during sporulation determines the success of the
future spore germination. iScience 25:105242.

16.   Ramírez‐Guadiana FH, Meeske AJ, Wang X, Rodrigues CD, Rudner
DZ. 2017. The *Bacillus subtilis* germinant receptor GerA triggers
premature germination in response to morphological defects during
sporulation. Mol Microbiol 105:689-704.

17.   Gao Y, Barajas-Ornelas RDC, Amon JD, Ramírez-Guadiana FH, Alon A,
Brock KP, Marks DS, Kruse AC, Rudner DZ. 2022. The SpoVA
membrane complex is required for dipicolinic acid import during
sporulation and export during germination. Genes Dev 36:634-646.

18.   Belitsky BR, Sonenshein AL. 1998. Role and regulation of *Bacillus*
subtilis glutamate dehydrogenase genes. J Bacteriol 180:6298-6305.

19.   Yoshida K, Fujita Y, Ehrlich SD. 1999. Three asparagine synthetase
genes of *Bacillus subtilis*. J Bacteriol 181:6081-91.

20.   de Vries YP, Atmadja RD, Hornstra LM, de Vos WM, Abee T. 2005.
Influence of glutamate on growth, sporulation, and spore properties of
Bacillus cereus ATCC 14579 in defined medium. Appl Environ Microbiol
71:3248-3254.

21.   Lin TH, Wei GT, Su CC, Shaw GC. 2012. AdeR, a PucR-type
transcription factor, activates expression of L-alanine dehydrogenase
and is required for sporulation of *Bacillus subtilis*. J Bacteriol 194:4995-
5001.

22.   Meeske AJ, Rodrigues CD, Brady J, Lim HC, Bernhardt TG, Rudner DZ.
2016. High-throughput genetic screens identify a large and diverse
collection of new sporulation genes in *Bacillus subtilis*. PLoS Biol

14:e1002341.

23.   Hilbert DW, Piggot PJ. 2004. Compartmentalization of gene expression during *Bacillus subtilis* spore formation. Microbiol Mol Biol Rev 68:234-262.

24.   Nugroho FA, Yamamoto H, Kobayashi Y, Sekiguchi J. 1999. Characterization of a New Sigma-K-Dependent Peptidoglycan Hydrolase Gene That Plays a Role in *Bacillus subtilis* Mother Cell Lysis. J Bacteriol 181:6230-6237.

25.   Smith TJ, Foster SJ. 1995. Characterization of the involvement of two compensatory autolysins in mother cell lysis during sporulation of *Bacillus subtilis* 168. J Bacteriol 177:3855-3862.

26.   Kuroda A, Asami Y, Sekiguchi J. 1993. Molecular cloning of a sporulation-specific cell wall hydrolase gene of *Bacillus subtilis*. J Bacteriol 175:6260-6268.

27.   Updegrove TB, Harke J, Anantharaman V, Yang J, Gopalan N, Wu D, Piszczek G, Stevenson DM, Amador-Noguez D, Wang JD. 2021. Reformulation of an extant ATPase active site to mimic ancestral GTPase activity reveals a nucleotide base requirement for function. elife 10:e65845.

28.   Kornberg A, Spudich JA, Nelson DL, Deutscher MP. 1968. Origin of proteins in sporulation. Annu Rev Biochem 37:51-78.

29.   Sekar V, Hageman JH. 1987. Protein turnover and proteolysis during sporulation of *Bacillus subtilis*. Folia Microbiol 32:465-480.

30.   Dworkin J, Shah IM. 2010. Exit from dormancy in microbial organisms. Nature Reviews Microbiology 8:890-896.

31.   Yasuda Y, Kanda K, Nishioka S, Tanimoto Y, Kato C, Saito A, Fukuchi S, Nakanishi Y, Tochikubo K. 1993. Regulation of L-alanine-initiated germination of *Bacillus subtilis* spores by alanine racemase. Amino Acids 4:89-99.

32.   Huergo LF, Dixon R. 2015. The emergence of 2-oxoglutarate as a

master regulator metabolite. Microbiol Mol Biol Rev 79:419-435.

33.   Gunka K, Commichau FM. 2012. Control of glutamate homeostasis in
Bacillus subtilis: a complex interplay between ammonium assimilation,
glutamate biosynthesis and degradation. Mol Microbiol 85:213-224.

34.   Manabe K, Kageyama Y, Morimoto T, Ozawa T, Sawada K, Endo K,
Tohata M, Ara K, Ozaki K, Ogasawara N. 2011. Combined Effect of
Improved Cell Yield and Increased Specific Productivity Enhances
Recombinant Enzyme Production in Genome-Reduced *Bacillus subtilis*
Strain MGB874. Appl Environ Microbiol 77:8370-8381.

35.   Nelson DL, Kornberg A. 1970. Biochemical Studies of Bacterial
Sporulation and Germination: XVIII. FREE AMINO ACIDS IN SPORES.
J Biol Chem 245:1128-1136.

36.   Mutlu A, Trauth S, Ziesack M, Nagler K, Bergeest J-P, Rohr K, Becker
N, Höfer T, Bischofs IB. 2018. Phenotypic memory in *Bacillus subtilis*
links dormancy entry and exit by a spore quantity-quality tradeoff.
Nature Communications 9:69.

37.   Gundlach J, Commichau FM, Stülke J. 2018. Perspective of ions and
messengers: an intricate link between potassium, glutamate, and cyclic
di-AMP. Curr Genet 64:191-195.

38.   Guérout-Fleury A-M, Frandsen N, Stragier P. 1996. Plasmids for
ectopic integration in *Bacillus subtilis*. Gene 180:57-61.

39.   Harwood CR, Cutting SM. 1990. Molecular biological methods for
Bacillus. Chichester ; New York : Wiley.

40.   Zhou B, Semanjski M, Orlovetskie N, Bhattacharya S, Alon S, Argaman
L, Jarrous N, Zhang Y, Macek B, Sinai L, Ben-Yehuda S. 2019.
Arginine dephosphorylation propels spore germination in bacteria. Proc
Natl Acad Sci USA 116:14228-14237.

41.   Vepachedu VR, Setlow P. 2007. Role of SpoVA proteins in release of
dipicolinic acid during germination of *Bacillus subtilis* spores triggered
by dodecylamine or lysozyme. J Bacteriol 189:1565-1572.

42.   Yi X, Setlow P. 2010. Studies of the commitment step in the
germination of spores of *Bacillus* species. J Bacteriol 192:3424-3433.
687
688

Figure legends

Figure 1. TMT quantitative proteomic analysis. (A) Proteins were compared between dormant spores (DS) and vegetative cells (VC) of at the onset of sporulation (t_0); (B) Heatmap of differential expressed proteins in DS samples were grouped using Hierarchical Cluster Analysis. Each line represented a protein, with fold change (FC) > 1.2 and $p < 0.05$ (T-test) as the screening criteria. The proteins with significantly decreased expression were marked in blue, the proteins with significantly increased expression were in red, the proteins without quantitative information were in gray; (C) The volcano map of proteins in DS group was drawn based on FC and the p-value of T test. The proteins with significantly decreased (FC < 0.83, $p < 0.05$) and increased (FC > 1.2, $p < 0.05$) expression were marked in blue and red, respectively, and the non-differentiated proteins were in gray; (D) The enrichment map (Top 20) of KEGG pathway enrichment analysis of differentially expressed proteins in the DS group by Fisher's exact test. The color of the bubble represents the significance of the enriched KEGG pathway, and the color gradient represents the size of the p-value (-log10), and the closer to red, the smaller the p-value. The size of the bubble represents the amount of differential protein.

Figure 2. Alanine and glutamate metabolism collaboratively regulate sporulation with an additive effect. (A) Representative phase-contrast images of sporulating cells at the late sporulation stage t_{19}. *B. subtilis* PY79 (wt), YZ11 (Δ*ald*), YZ19 (Δ*rocG*), YZ12 (Δ*ald* Δ*rocG*), and YZ13 (Δ*ald* Δ*rocG*, *amyE::ald-rocG*) strains were induced to sporulate in DSM at 37°C for 22 hrs and followed by microscopy. Red arrowheads point to premature germinated spores. Scale bar, 2 µm; (B) The percentage of sporulation of the strains described in (A). Data are presented as the percentage of total number of colonies forming units (CFU) before and after heat treatment (80°C, 20 min). Shown is a representative experiment out of three independent biological

repeats.

Figure 3. RocG-mediated glutamate metabolism is essential for ensuring the

sporulation efficiency. (A) Representative phase-contrast images of

sporulating cells at the late sporulation stage t_{19}. *B. subtilis* PY79 (wt), YZ22

(Δald $\Delta rocG$ $\Delta gerAA$), YZ24 (Δald $\Delta rocG$ $\Delta gerAA$, $amyE::P_{IPTG}$-*rocG*), YZ25

(Δald $\Delta rocG$ $\Delta gerAA$, $amyE::P_{IPTG}$-*ald*), and YZ26 (Δald $\Delta rocG$ $\Delta gerAA$,

$amyE::P_{IPTG}$-*rocG-ald*) strains were induced to sporulate in DSM at 37°C for

22 hrs and followed by microscopy. 50 µM IPTG was added to the YZ24,

YZ25 and YZ26 cultures at the sporulation t_0 to induce corresponding gene

expression. Scale bar, 2 µm; (B) Quantification of the experiment described in

(A). Data are presented as percentages of the number of the phase-bright

spores and all sporulating cells in the same image (n ≥ 800 for each strain);

(C) AGFK-induced germination of spores collected in (A). Spores of wt, as

well as YZ24, YZ25, and YZ26 strains with 50 µM IPTG induction, were

purified and incubated with AGFK (10 mM) to trigger germination. DPA

release was measured by detecting the relative fluorescence units (RFU) of

Tb^{3+}-DPA. Shown is a representative experiment out of three independent

biological repeats.

Figure 4. The expression of *rocG* during sporulation is crucial to ensure the

quality of correspondingly generated spores. (A) Representative phase-

contrast images of sporulating cells at the late sporulation stage t_{19}. *B. subtilis*

PY79 (wt), YZ81 ($\Delta adeR$ $\Delta gerAA$), YZ23 ($\Delta ahrC$ $\Delta rocR$ $\Delta gerAA$), and YZ90

($\Delta adeR$ $\Delta ahrC$ $\Delta rocR$ $\Delta gerAA$) strains were induced to sporulate in DSM at

37°C for 22 hrs and followed by microscopy. Scale bar, 2 µm; (B) Expression

of the *ald* gene in wt, YZ11 (Δald) and YZ81 strains, and *rocG* gene in wt,

YZ19 ($\Delta rocG$), and YZ23 strains. Sporulating cells were collected at t_0 and

detected as described in Methods; (C) AGFK-induced germination of spores

collected in (A). Spores of wt, YZ81, YZ23, and YZ90 strains were purified and incubated with AGFK (10 mM) to trigger germination. DPA release was measured by detecting the RFU of Tb^{3+}-DPA. Shown is a representative experiment out of three independent biological repeats.

Figure 5. The absence of Ald induces premature germination during sporulation. (A) Representative phase-contrast images of sporulating cells at the late sporulation stage t_{19}. *B. subtilis* PY79 (wt), YZ11 (Δ*ald*), and YZ21 (Δ*ald* Δ*gerAA*) strains were induced to sporulate in DSM at 37°C for 22 hrs and followed by microscopy. Red arrowheads point to premature germinated spores. Scale bar, 2 μm; (B) Quantification of the experiment described in (A). Data are presented as percentages of the number of the phase-bright spores and all sporulating cells in the same image (n ≥ 800 for each strain); (C) AGFK-induced germination of spores collected in (A). Spores of wt, YZ11, and YZ21 strains were purified and incubated with AGFK (10 mM) to trigger germination. DPA release was measured by detecting the RFU of Tb^{3+}-DPA. Shown is a representative experiment out of three independent biological repeats; (D) Models of the premature germination in the Δ*ald* mutant. YZ11 strain was induced to sporulate in DSM at 37°C. After 14 hrs of incubation, sporulating cells were collected on an exhausted DSM gel-pad as described in Methods, and followed by time-lapse microscopy at a 10 min interval.

Figure 6. Ald-mediated alanine metabolism regulates L-alanine content in the external environment of generated spores. (A) Representative phase-contrast images of sporulating cells at the late sporulation stage t_{19}. YZ31 (Δ*ald*, *amyE::P$_{IPTG}$-ald*) strains were induced to sporulate in DSM at 37°C for 22 hrs and followed by microscopy. 0-1000 μM IPTG was added at the sporulation t_0 to induce *ald* expression. Scale bar, 2 μm; (B) The percentage of phase-bright spores as well as the environmental L-alanine content of the wt and YZ31 strains at the late sporulation stage t_{19}. The percentage of phase-bright spores

are presented as ratio of the number of the phase-bright spores and all sporulating cells in the same image (n ≥ 800 for each strain); The environmental L-alanine content was detected as described in Methods; (C) Expression of the *ald* gene in YZ31 strains with different IPTG induction. Sporulating cells were collected 10 min after IPTG addition and detected as described in Methods. Shown is a representative experiment out of three independent biological repeats.

Figure 7. Cytological sporulation assay reveals the impaired sporulation stage of Δ*rocG* mutants. (A-B) Representative phase contrast and the indicated fluorescent images of wt and YZ19 (Δ*rocG*) cells harboring two transcriptional fusions, σ^G and σ^K, at sporulation t_4, t_7, and t_{25}. Scale bar, 2 μm; (C) Expression of the *cwlC* and *cwlH* gene in wt and YZ19 strains. Sporulating cells were collected at t_7 and detected as described in Methods. Shown is a representative experiment out of three independent biological repeats. (D) Models of the sporulation defect in the Δ*rocG* mutants.

Figure 8. RocG regulates both mother cell lysis and spore properties. (A) Representative phase-contrast images of sporulating cells at the late sporulation stage t_{19}. *B. subtilis* PY79 (wt), YZ19 (Δ*rocG*) and YZ32 (Δ*rocG, amyE::P_{IPTG}-rocG*) strains were induced to sporulate in DSM at 37°C for 22 hrs and followed by microscopy. 0-50 μM IPTG was added to YZ32 at the sporulation t_0 to induce *rocG* expression. Scale bar, 2 μm; (B) Expression of the *rocG* gene in strains indicated in (A). Sporulating cells were collected 30 min after IPTG addition and detected as described in Methods; (C) Quantification of released spores produced by strains indicated in (A). Data are presented as percentages of the number of the released spores and all sporulating cells in the same image (n ≥ 800 for each strain); (D) ATP levels in strains indicated in (A). Sporulating culture was collected at t_1 and analyzed for ATP level as described in Methods; (E) DPA content in spores of wt and

YZ32 with different IPTG induction. Spores were purified and boiling for 20 min. DPA content was measured by detecting the RFU of Tb^{3+}-DPA; (F) Heat resistance of spores collected in (E). Data are presented as the percentage of total number of CFU before and after heat treatment (90°C, 10 min); (G-I) Germination phenotypes of spores collected in (E). Spores were purified and incubated with (G) L-alanine (10 mM), (H) AGFK (10 mM), and (I) DDA (10 mM) to trigger germination. DPA release was measured by detecting the RFU of Tb^{3+}-DPA. Shown is a representative experiment out of three independent biological repeats.

Figure 9. The crucial roles of alanine and glutamate catabolism in ensuring proper sporulation.

Figure 1

Proteomic analysis

B

VC DS

TMT quantitative proteomics

1
0.5
0
-0.5
-1

C

-log10(p-value)

7.5

5.0

2.5

0.0

-4   0   4
log2(DS/VC)

Regulated Type
- unchange
- down
- up

D

KEGG Enrichment

Biofilm formation - Escherichia coli
C5-Branched dibasic acid metabolism
Streptomycin biosynthesis
Inositol phosphate metabolism
Alanine, aspartate and glutamate metabolism
Methane metabolism
Ribosome
One carbon pool by folate
Flagellar assembly
Bacterial chemotaxis
Arginine biosynthesis
Glyoxylate and dicarboxylate metabolism
Thiamine metabolism
Nitrogen metabolism
Nonribosomal peptide structures
Histidine metabolism
Tryptophan metabolism
Glycine, serine and threonine metabolism
Purine metabolism
Glycolysis / Gluconeogenesis

KEGG Pathways(Top 20)

0.8   0.9   1.0
Rich factor

$-\log_{10}$(p.value)
3.0
2.5
2.0
1.5
1.0

Protein number
- 10
- 20
- 30
- 40
- 50

Figure 2

A

B

Figure 3

Figure 4

A

B

C

Figure 5

Figure 6

A

IPTG/μM

B

C

Figure 7

Figure 8

Figure 9