RGS10 deficiency facilitates distant metastasis by inducing epithelial–mesenchymal transition in breast cancer

Yang Liu[1#], Yi Jiang[1#], Peng Qiu[2#], Tie Ma[3#], Jiawen Bu[1], Yueting Hu[1],Ming Jin[1], Tong Zhu[4], Xi Gu[1*]

1 Department of Oncology, Shengjing Hospital of China Medical University, Shenyang, China

2 Department of Anesthsiology, Shengjing Hospital of China Medical University, Shenyang, China

3 Department of Pathology, Shengjing Hospital of China Medical University, Shenyang, China

4 Breast Surgery of Panjin Central Hospital, Panjin, China

# These authors contributed equally to this work.

*For correspondence:

jadegx@163.com (XG);

Abstract

Distant metastasis is the major cause of death in patients with breast cancer.

Epithelial–mesenchymal transition (EMT) contributes to breast cancer metastasis.

Regulator of G protein-signaling (RGS) proteins modulate metastasis in various

cancers. This study identified a novel role for RGS10 in EMT and metastasis in breast

cancer. RGS10 protein levels were significantly lower in breast cancer tissues

compared to normal breast tissues, and deficiency in RGS10 protein predicted a worse

prognosis in patients with breast cancer. RGS10 protein levels were lower in the highly

aggressive cell line MDA-MB-231 than in the poorly aggressive, less invasive cell lines

MCF7 and SKBR3. Silencing RGS10 in SKBR3 cells enhanced EMT and caused

SKBR3 cell migration and invasion. The ability of RGS10 to suppress EMT and

metastasis in breast cancer was dependent on lipocalin-2 and miR-539-5p. These

findings identify RGS10 as a tumor suppressor, prognostic biomarker, and potential

therapeutic target for breast cancer.

Key words: Breast cancer, miRNA, metastasis, RGS10, EMT.

Introduction

Breast cancer is the most common cancer among women worldwide. Globally, in 2020, an estimated 2.3 million new cases of breast cancer were diagnosed, and there were approximately 685,000 deaths from the disease (Sung et al., 2021). The majority of breast cancer mortality is due to distant metastasis, with 5-year survival estimated at 30% (American Cancer Society, 2023). There is a critical need to identify early breast cancer metastasis using prognostic biomarkers to ensure patients receive effective anticancer therapies in a timely manner (Miglietta et al., 2022).

Epithelial-to-mesenchymal transition (EMT) plays a critical role in tumor progression and metastatic invasion in breast cancer. EMT describes the process by which epithelial cells lose their epithelial characteristics and cell–cell contact and increase their invasive potential (Singh and Chakrabarti, 2019). EMT is characterized by a loss of epithelial cell markers, such as cytokeratins and E-cadherin, followed by an upregulation in the expression of mesenchymal cell markers, such as N-cadherin and vimentin. EMT is regulated at different levels by factors involved in cell signaling, transcriptional control, and epigenetic and post-translational modifications (Lai et al., 2020).

Biomarkers of EMT may predict early breast cancer metastasis and facilitate clinical decision-making (Liu et al., 2016). The regulator of G protein signaling 10 (RGS10) belongs to the superfamily of RGS proteins that bind and deactivate heterotrimeric G proteins (Almutairi et al., 2020). RGS proteins are important mediators of essential cellular processes and may be tumor initiators or tumor suppressors (Li et al., 2023). The canonical function of RGS proteins is to act as GTPase-activating proteins (GAPs), accelerate GTP hydrolysis on G-protein alpha subunits, and terminate signaling pathways downstream of G protein-coupled receptors (Hunt et al., 1996). RGS proteins can have noncanonical GAP-independent functions, including the

suppression of transforming growth factor beta (TGF-β)-induced EMT in non-small cell lung

cancer by RGS6 (Wang et al., 2022).

RGS10 is a critical regulator of cell survival, polarization, adhesion, chemotaxis, and

differentiation that exhibits tumor-suppressing effects in ovarian and colorectal cancer. In

ovarian cancer cells, RGS10 suppression increases proliferation by phosphorylation of mTOR,

4E-BP1, p70S6K and rProtein-S6, including in the presence of chemotherapy (Altman et al.,

2015), loss of RGS10 expression contributes to the development of chemoresistance (Cacan et

al., 2014), and modulating RGS10 expression can alter sensitivity to paclitaxel, cisplatin, and

vincristine (Cacan et al., 2014; Hooks et al., 2010; Hooks and Murph, 2015). In ovarian tumors,

transcription of RGS10 is regulated by DNA methylation and histone deacetylation (Nguyen et

al., 2014). In colorectal cancer, RGS10 expression is suppressed, and inhibition of DNA

methylation may contribute to improved prognosis (Caldiran and Cacan, 2022).

In breast cancer, RGS10 is downregulated in heavily metastatic human breast cancer cell

populations compared to weakly metastatic human breast cancer cell populations (Montel et

al., 2005; Steeg, 2005). The mechanisms underlying the metastasis-suppressing function of

RGS10 in breast cancer remain to be elucidated. In this study, we investigated the function of

RGS10 in breast cancer, specifically in breast cancer metastasis. The objectives were to 1)

characterize the expression of RGS10 in freshly resected breast cancer and adjacent normal

breast tissues; 2) determine the prognostic significance of RGS10 expression in patients with

breast cancer; and 3) explore the role of RGS10 and upstream effectors in tumor progression

and metastasis in breast cancer cells *in vitro* and *in vivo*.

Methods

Key resources table

Reagent type (species) or resource	Designation	Source Or reference	Identifiers	Additional information
Cell line (SKBR3)	HER2+ breast cancer cell line	ATCC		Cell line cultured in McCoy's 5A Culture medium supplemented with 10% FBS
Cell line (MCF-7)	ER+/PR+ breast cancer cell line	ATCC		Cell line cultured in DMEM high glucose Culture medium supplemented with 10% FBS
Cell line (MDA-MB-231)	Triple-negative breast cancer cell line	ATCC		Cell line cultured in L-15 Culture medium supplemented with 10% FBS
Transfected construct (human)	RGS10 shRNA#161,321,506	Syngentech Biotechnology		Lentiviral construct to transfect and express the shRNA
Transfected construct (human)	miRNA-539-5p mimics and inhibitor	GenePharma Biotechnology		
Antibody	Anti-RGS10 (rabbit monoclonal)	Abcam	Cat# ab154172	WB (1:250) IHC (1:50)
Antibody	Anti-lipocalin-2 (rabbit monoclonal)	ABclonal	Cat# A2092	WB(1:1000) IHC (1:500)
Antibody	Anti-E-cadherin	Proteintech	Cat# 20874-1-AP	WB (1:1000) IHC (1:500)

				IF (1:100)
Antibody	Anti-vimentin (mouse monoclonal)	HUABIO	Cat# EM0401	WB (1:1000) IHC (1:200) IF (1:100)
Antibody	Anti-snail (rabbit monoclonal)	ABclonal	Cat# A5243	WB (1:1000) IHC (1:00) IF (1:100)
Antibody	Anti-GAPDH (rabbit monoclonal)	CST	Cat# 5174	WB (1:1000)
Antibody	Anti-β-tubulin (rabbit monoclonal)	ABclonal	Cat# AC008	WB (1:5000)
Sequence-based reagent	RGS10_F	This paper	PCR primers	CACGACAGCGAT GGCAGTTCC
Sequence-based reagent	RGS10_R	This paper	PCR primers	CTTTTCACGCCTT CTGGGTCTTCC
Sequence-based reagent	MiR-539-5p_F	This paper	PCR primers	CGCTGCATGGAG AAATTATCCT
Sequence-based reagent	MiR-539-5p_R	This paper	PCR primers	CAGTGCAGGGTC CGAGGT
Software, algorithm	SPSS	SPSS	SPSS, version 25.0 Armonk, NY, USA	

Bioinformatics

The Genotype-*Tissue Expression* database (www.gtexportal.org) was used to analyze

RGS10 mRNA levels in 31 normal human tissues. The Cancer Cell Line Encyclopedia

database (https://sites.broadinstitute.org/ccle) was used to analyze RGS10 mRNA levels in a

series of cell lines (Rahman et al., 2019). The Kaplan–Meier plotter (http://kmplot.com) was

used to assess the relevance of RGS10 mRNA expression for disease-free survival (DFS) and

overall survival (OS) in patients with breast cancer (Győrffy, 2021).

Tissues and patients

Twenty paired breast cancer and normal adjacent tissues were obtained from patients

undergoing surgical resection at Shengjing Hospital of China Medical University.

This study included an additional 133 patients with histologically confirmed invasive

ductal breast carcinoma who received treatment at the Shengjing Hospital of China Medical

University from April 2006 to April 2008. Patients had: 1) undergone surgery with no distant

metastases at the time of the operation; 2) received standard adjuvant therapy after surgery; 3)

≥10 axillary lymph nodes dissected and examined pathologically after surgery (Coates et al.,

2015; Gradishar et al., 2021; Wolff et al., 2013); and 4) ≥10 years of follow-up as outpatients or

by telephone interviews.

This study was approved by the Institutional Review Board of Shengjing Hospital of

China Medical University.

Breast cancer cells and culture

Human breast cancer cell lines SKBR3, MCF-7 and MDA-MB-231 were obtained from

American Type Culture Collection (ATCC, Manassas, VA). The identity of cell lines had been

authenticated by using STR profiling. SKBR3 cells were cultured in McCoy's 5A (modified)

medium supplemented with 10% fetal bovine serum at 37°C and 5% CO_2. MCF7 cells were

cultured in DMEM high glucose (Invitrogen) medium supplemented with 10% fetal bovine

serum at 37°C and 5% CO_2. MDA-MB-231 cells were cultured in Leibovitz's L-15 medium

supplemented with 10% fetal bovine serum at 37°C without CO_2.

Transfection

SKBR3 cells were transfected with short hairpin (sh)RNAs specifically targeting RGS10

(Syngentech Biotechnology, Beijing, China). shRNA sequences were: shRNA-RGS10-161,

5′-GCCTCAAGAGCACAGCCAAAT-3′;   shRNA-RGS10-321, 5′-

GGAGATCTACATGACCTTTCT-3′;   a n d   shRNA-RGS10-506,

5′-GCACCCTCTGATGTTCCAGAA-3′. The shRNA-negative control (NC) sequence was

5′-AAACGTGACACGTTCGGAGAA-3′. As transfection efficiencies of shRNA-RGS10-161

and shRNA-RGS10-506 were better than shRNA-RGS10-321, shRNA-RGS10-161 and

shRNA-RGS10-506 were used for subsequent experiments.

A miR-539-5p mimic, miR-539-5p inhibitor, and corresponding NC were designed and

synthesized by GenePharma Biotechnology (Shanghai, China). The sequences of miRNA

were: miR-539-5p mimic, 5′-GGAGAAAUUAUCCUUGGUGUGU-3′; miR-539-5p mimic

NC,5′-UUCUCCGAACGUGUCACGUTT-3′;   miR-539-5p

inhibitor,5′-ACACACCAAGGAUAAUUUCUCC-3′;   miR-539-5p   inhibitor   NC

5′-CAGUACUUUUGUGUAGUACAA′. SKBR3 cells were transfected with the miR-539-5p

mimic or NC. MDA-MB-231 cells were transfected with the miR-539-5p inhibitor or NC.

Lipofectamine 3000 (L3000015, Thermo Fisher Scientific, Invitrogen, USA) was used for cell

transfection.

Reverse transcription–quantitative polymerase chain reaction (RT-qPCR)

Total RNA was isolated from breast cancer tissues and breast cancer cell lines using

Triquick reagent (Solarbio Life Science, Beijing, China) and reverse-transcribed using a cDNA

synthesis kit (TaKaRa Bio, Kusatsu, Japan). RT-qPCR was conducted using SYBR Green

PCR Master Mix (TaKaRa) on a Fast-Real-Time PCR System. Primers were: RGS10 forward,

5′-CACGACAGCGATGGCAGTTCC-3′; RGS10 reverse,

5′-CTTTTCACGCCTTCTGGGTCTTCC-3′; GAPDH forward,

5′-CCTTCCGTGTCCCCACT-3′; and GAPDH reverse, 5′-GCCTGCTTCACCACCTTC-3′.

Thermocycling conditions were 95°C for 30 s, 40 cycles of 95°C for 3 s, and 60°C for 30 s.

Relative mRNA levels were calculated using the comparative threshold (Cq) cycle method

($2^{-\Delta\Delta Ct}$) (Adnan et al., 2011).

cDNA synthesis in SKBR3 cells transfected with the miR-539-5p mimic or NC and

MDA-MB-231 cells transfected with the miR-539-5p inhibitor or NC was conducted using a

miRNA First Strand cDNA Synthesis kit that adopted A-tailing (B532461, Sangon Biotech).

qPCR was performed using a miRNA qPCR kit (B532461, Sangon Biotech). Relative

miR-539-5p level was normalized to U6 as the internal control. Primers were as follows:

miR-539-5p forward, 5′-CGCTGCATGGAGAAATTATCCT-3′; miR-539-5p reverse,

5′-CAGTGCAGGGTCCGAGGT-3′; U6 forward,

5′-GCTCGCTTCGGCAGCACATATAC-3′, and U6 reverse

5′-CGAATTTGCGTGTCATCCTTGCG-3′. Thermocycling conditions were 95°C for 30 s, 40

cycles of 95°C for 5 s, and 60°C for 30 s. Relative mRNA levels were calculated using the

$2^{-\Delta\Delta Ct}$ method.

Cell Counting Kit-8 (CCK-8) analysis

158   Cells were seeded into 96-well plates at a density of 5000 cells per well. After 0, 24, 48,

159 and 72 h, CCK-8 (Cofitt Life Science Biotechnology) solution was added to each well, and

160 cells were incubated for 3 h. Optical density was measured at 450 nm using a microplate

161 spectrophotometer (Bio-Rad). Experiments were performed in triplicate.

Colony formation assay

SKBR3 and MDA-MB-231 cells were trypsinized and plated on 6-well plates at a density of 1000 cells per well. When visible colonies were formed, the cells were fixed with methyl alcohol and stained with crystal violet. The plates were photographed, and the number of colonies was counted using ImageJ software.

Cell invasion and migration assay

For cell invasion, 20,000 cells in 0.2 mL serum-free culture medium were placed in the upper chamber of a Transwell chamber that had been precoated with Matrigel matrix. The lower chamber was filled with 1 mL culture medium containing 10% fetal bovine serum as the chemoattractant. After culturing at $37^{o}C$ for 24 h, cells that had migrated to the lower surface through the membrane were fixed with 4% paraformaldehyde for 10 min. Fixed cells were stained with crystal violet.

Detection of cell migration was similar to cell invasion, except the upper chamber of the Transwell chamber was not precoated with Matrigel.

Immunohistochemistry

Surgically resected tissues were fixed in 4% formaldehyde, embedded in paraffin, and sectioned into 6-μm-thick slices. Sections were deparaffinized with xylene, rehydrated in a series of graded ethanols, and rinsed in Tris-buffered saline. Sections were incubated with a primary antibody anti-RGS10 (1:50; Abcam, Cambridge, MA; ab154172), anti-lipocalin-2 (LCN2) (1:500, ABclonal, #A2092), anti-E-cadherin (1:500, Proteintech, #20874-1-AP), anti-vimentin (1:200, HUABIO, #EM0401), anti-snail (1:100, ABclonal, # A5243), at 4°C overnight, incubated with a secondary antibody (Zhong Shan Jin Qiao Biotechnology Co.,

Beijing, China) at room temperature for approximately 45 min, and protein was visualized with

a diaminobenzidine staining kit (Zhong Shan Jin Qiao Biotechnology Co.).

Immunofluorescence staining

After transfection with the miR-539-5p mimic or miR-539-5p inhibitor for 24 h, SKBR3

and MDA-MB-231 cells were smeared on glass slides, cultured for 36 h, and fixed with 4%

paraformaldehyde for 20 min. Cells were washed three times with phosphate-buffered saline

(PBS) and blocked with 5% bovine serum albumin in PBS (pH 7.4) at room temperature for 1

h. Cells were incubated with anti-E-cadherin (1:100, Proteintech, #20874-1-AP),

anti-vimentin (1:100, HUABIO, #EM0401), or anti-snail (1:100, ABclonal, #A5243) antibody

at 4°C overnight. Cells were washed three times with PBS, incubated with Alexa-Fluor 594

goat anti-mouse IgG H&L or Alexa-Fluor 488 goat anti-rabbit IgG H&L (Proteintech, 1:400

dilution) at room temperature for 1 h in the dark, washed three times, and incubated with

4′,6-diamidino-2-phenylindole (Solarbio, #C0065) for 10 min. After staining, cells were sealed

with mounting media. Pictures were taken under a fluorescence microscope (Nikon, Japan).

Western blotting

Cells were harvested by trypsinization and lysed in radioimmunoprecipitation assay lysis

buffer. Cell lysate was centrifuged to a pellet (insoluble material). Proteins in the supernatant

(30 μg of protein/lane) were separated by sodium dodecyl sulfate–polyacrylamide gel

electrophoresis and transferred to polyvinylidene fluoride membranes. After blocking with 5%

w/v nonfat dry milk for 1 h, membranes were incubated with anti-RGS10 (1:250; Abcam;

ab154172), anti-GAPDH (1:1000, CST, 5174S), anti-lipocalin-2 (LCN2) (1:1000, ABclonal,

#A2092), anti-E-cadherin (1:1000, Proteintech, #20874-1-AP), anti-vimentin (1:1000,

HUABIO, #EM0401), anti-snail (1:1000, ABclonal, # A5243), or anti-β-tubulin (loading

control) (1:5000, ABclonal, AC008) antibody at 4°C overnight. Membranes were incubated

with a secondary antibody (Zhong Shan Jin Qiao Biotechnology Co.) at 37°C for 1 h. Bands

were visualized using an enhanced chemiluminescent reagent. The relative protein levels of

RGS10, LCN2, E-cadherin, vimentin, and snail were determined by densitometric analysis

using Image Lab™ 6.0.1 software.

Dual-luciferase reporter assay

The StarBase database was used to predict potential binding sites between RGS10 mRNA

and hsa-miR-539-5p. Sequences containing the 3′-untranslated region fragments of RGS10

and the mutated binding site of hsa-miR-539-5p or wild type were designed and constructed by

GeneChem Biotechnology (Shanghai, China). The miR-539-5p mimic or NC was

cotransfected with the wild-type or mutant plasmid using lipofectamine 3000 (Invitrogen).

Luciferase intensity was recorded 48 h after transfection using the dual-luciferase reporter

assay system (Promega, Madison, WI).

Molecular interaction networks

The Search Tool for the Retrieval of Interacting *Genes*/Proteins database

(https://string-db.org/) was used to identify proteins that interact with RGS10, and conduct a

protein–protein network interaction analysis (Szklarczyk et al., 2015). Subsequently, the

Database for Annotation, Visualization, and Integrated Discovery (https://david.ncifc rf.gov)

was used to perform Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes

(KEGG) pathway-enrichment analysis.

In-vivo **animal experiments**

Nine BALB/C female nude mice (6 weeks old; 19–22 g) (Beijing Huafukang Biotechnology

Company) were randomly assigned into three groups (n = 3 each). Mice were maintained in a

specific-pathogen-free environment at 28°C and 50% humidity. The experimental design was

approved by the Animal Experiment Ethics Committee of Shengjing Hospital. A total of $1 \times$

10^6 SKBR3 cells transfected with shRNA-RGS10 or shRNA-NC was injected subcutaneously.

Tumor volume was measured at 25 days after inoculation.

Statistical analysis

Statistical analyses were conducted using SPSS 25.0 software. *In-vitro* experiments were

performed in triplicate. Associations between RGS10 gene and protein expression and

clinicopathological characteristics were evaluated with the chi-squared test or independent

samples *t*-test, as appropriate. Survival curves were generated using the Kaplan–Meier method.

Univariate and multivariate Cox regression analyses were used to identify prognostic

predictors related to DFS and OS. Hazard ratios and corresponding 95% confidence intervals

were calculated. All *P* values were two-sided, with $P < 0.05$ considered statistically significant.

Results

Expression and prognostic associations of *RGS10* in breast cancer

To determine the role of RGS10 in breast cancer, we analyzed RGS10 mRNA levels in

normal tissues (n = 31) from the Genotype-Tissue Expression dataset, which showed RGS10

mRNA levels were high in normal breast, blood, colon, and small intestine tissues and low in

normal heart, liver, and pancreas tissues (Figure 1A). Next, we determined RGS10 mRNA

levels in cell lines representing 21 human cancers from the Cancer Cell Line Encyclopedia

database (Figure 1B). Finally, we applied RT-qPCR in freshly resected breast cancer tissues (n

= 20) and matched adjacent normal breast tissues and showed RGS10 mRNA levels were

lower in breast cancer tissues compared to normal breast tissues ($P = 0.003$; Figure 1C).

Clinicopathological characteristics of these 20 patients are shown in Table 1. This pattern

implied a downregulation of RGS10 expression in breast cancer tissues.

To investigate the biological role and clinical and prognostic significance of RGS10 in

breast cancer tissues, we used survival analyses. In breast cancer samples from the

Kaplan–Meier plotter database, high RGS10 mRNA level was associated with significantly

improved DFS ($P = 0.0066$; Figure 1D) and OS ($P = 0.027$; Figure 1E). In surgically resected

breast cancer tissues (n=133), RGS10 protein expression level detected by

immunohistochemistry (representative images shown in Figure 1F) was positively correlated

with breast cancer subtype ($P = 0.043$), distant metastasis ($P = 0.008$), and survival status ($P =$

0.024). There were no correlations with age, comorbid disease, histological grade, tumor size,

number of positive axillary lymph nodes, number of pregnancies, number of births, age at first

pregnancy, menopausal status, estrogen receptor status, progesterone receptor status, human

epidermal growth factor receptor 2 (HER2) status, or Ki67 status (Table 2). In these patients, a

high RGS10 protein expression level was associated with a longer DFS ($P = 0.003$, Figure 1G)

and OS ($P = 0.022$, Figure 1H). Clinicopathological characteristics associated with DFS and

OS were identified with Cox regression analyses. On multivariate regression analysis,

histological grade and RGS10 protein expression were independent predictors of DFS (Table

3). On univariate regression analysis, age, histological grade, and RGS10 protein expression

were independent predictors of OS (Table 4). This suggests low RGS10 expression is

associated with poor survival in patients with breast cancer.

Taken together, these findings imply that RGS10 has a role in suppressing breast cancer

and RGS10 may represent a potential prognostic biomarker in breast cancer.

RGS10 silencing increases the proliferation and migration of breast cancer cells *in vitro*

To characterize RGS10 protein expression in the breast cancer cell lines MDA-MB-231,

MCF7, and SKBR3, we conducted Western blotti*ng.* MDA-MB-231 is a highly aggressive,

invasive, and poorly differentiated triple-negative breast cancer cell line (Bianchini et al.,

2016). MCF7 is an adenocarcinoma cell line with estrogen, progesterone, and glucocorticoid receptors (Lee et al., 2015). The SKBR3 cell line overexpresses the HER2/c-erb-2 gene product (Merlin et al., 2002). RGS10 protein levels appeared lower in the highly aggressive cell line MDA-MB-231 compared to the poorly aggressive and less invasive cell lines MCF7 and SKBR3 (Figure 2A). This finding validates our previous observations suggesting that RGS10 acts as a tumor suppressor in breast cancer.

To investigate the role of RGS10 in breast cancer cells, we silenced RGS10 in SKBR3 cells using two independent shRNA constructs, shRNA-RGS10-161 and shRNA-RGS10-506, which had significantly improved transfection efficiency compared to shRNA-RGS10-321 (Figure 2B). We evaluated the effects of RGS10 on breast cancer cell colony formation, migration, and invasion in RGS10-depleted (shRNA-RGS10-506, shRNA-RGS10-161) SKBR3 cells and SKBR3 cells transfected with shRNA-negative control (NC). Colony formation and Transwell migration/invasion assays showed significantly increased clonogenic ability, migration ability, and invasive capacity in RGS10-depleted SKBR3 cells compared to NC (P < 0.001) (Figure 2C–F).

These findings show invasion and metastasis were enhanced in breast cancer cells lacking RGS10, suggesting an inhibitory effect of RGS10 in breast cancer metastasis.

RGS10 silencing promotes lipocalin-2 (LCN2) expression and EMT in breast cancer cells

To explore the mechanisms by which RGS10 suppresses breast cancer invasion and metastasis, we analyzed potential downstream tumor metastasis-related genes by comparing the transcriptomes in RGS10-depleted (shRNA-RGS10-506, shRNA-RGS10-161) SKBR3 cells and SKBR3 cells transfected with shRNA-NC. Differential gene expression was visualized using a volcano plot, which revealed 70 genes were significantly upregulated in RGS10-depleted SKBR3 cells (Figure 3A). KEGG pathway analysis and GO enrichment

analysis of differentially expressed genes identified upregulated KEGG pathways, biological

processes, cellular components, and molecular functions (Figure 4B–E). Upregulated KEGG

pathways were associated with cytokine-cytokine receptor interactions and extracellular

matrix-receptor interactions. Gene sets associated with high- and low-RGS10 mRNA

expression were identified by gene set enrichment analysis using the Molecular Signatures

Database (Liberzon et al., 2011). Gene sets associated with high-RGS10 mRNA expression

included allograft rejection, apoptosis, interleukin (IL) 6/Janus kinase/signal transducer and

activator of transcription pathway (STAT) 3, IL2/STAT5 pathway, and inflammatory response

(Table 5). Western blotting showed changes in biomarkers of EMT in RGS10-depleted

SKBR3 cells compared to NC. Neutrophil-derived cytokine LCN2 and vimentin protein levels

were higher and E-cadherin protein levels were lower in RGS10-depleted SKBR3 cells

compared to NC (Figure 3F–G).

These findings show that RGS10 deficiency promotes invasion and metastasis by

activating the LCN2 pathway to induce EMT in breast cancer cells, supporting the potential of

RGS10 as a prognostic biomarker in breast cancer.

miR-539-5p regulates the migration, invasion, proliferation, and EMT of breast cancer

cells

To study the upstream regulatory mechanism of RGS10 in breast cancer, we used the

StarBase database to predict miRNAs that could potentially bind to RGS10. The luciferase

reporter assay identified miR-539-5p as a miRNA that targets RGS10 in breast cancer cells

(Figure 3G–H).

To predict the potential effects of miR-539-5p on breast cancer cells, we transfected

SKBR3 and MDA-MB-231 cells with a miR-539-5p mimic to represent miR-539-5p

overexpression, a miR-539-5p inhibitor, or appropriate NCs (Figure 4A). RT-qPCR and

Western blotting validated transfection efficiency and showed that RGS10 mRNA and protein

levels were significantly decreased in SKBR3 cells overexpressing miR-539-5p compared to

SKBR3 cells transfected with miRNA-NC or the wild type (Figure 4B, C). Cell Counting Kit-8

(CCK-8) analysis, and colony formation and Transwell migration/invasion assays showed

SKBR3 cell proliferation, colony formation, migration, and invasion were significantly

increased in SKBR3 cells overexpressing miR-539-5p compared to SKBR3 cells transfected

with miRNA-NC or the wild type (Figure 4D–G). In contrast, RGS10 protein levels were

significantly increased (Figure 5A–C) and MDA-MB-231 cell proliferation, colony formation,

migration, and invasion were significantly decreased in MDA-MB-231 cells transfected with

miR-539-5p inhibitor compared to anti-miRNA-NC or the wild type (Figure 5D–G).

Based on these findings, for the first time, we propose a miR-539-5p/RGS10/LCN2

regulatory axis in breast cancer. Consistent with this, Western blotting and

immunofluorescence assays showed decreased E-cadherin protein expression and increased

LCN2, vimentin, and snail protein expression in SKBR3 cells overexpressing miR-539-5p

compared to SKBR3 cells transfected with miRNA-NC or the wild type (Figure 4H–I). These

effects were reversed in MDA-MB-231 cells transfected with a miR-539-5p inhibitor

compared to anti-miRNA-NC or the wild type (Figure 5H–I).

These findings identify miR-539-5p as a critical factor in breast cancer metastasis by

regulating RGS10/LCN2 expression.

RGS10 inhibits breast cancer growth by targeting LCN2 *in vivo*

To investigate tumorgenicity, we subcutaneously transplanted SKBR3 cells transfected

with shRNA-RGS10-506 or shRNA-NC into nude mice. The growth of tumors derived from

RGS10-depleted SKBR3 cells was significantly increased compared to NC, manifested by

larger tumor size compared to NC (Figure 6A–C). Immunohistochemistry showed LCN2,

snail, and vimentin protein expression were increased and E-cadherin protein expression was

decreased in tumor tissues derived from RGS10-depleted SKBR3 cells compared to NC

(Figure 6D).

These *in-vivo* data confirm the previous *in-vitro* observations that RGS10 deficiency

promotes invasion and metastasis by activating the LCN2 pathway to induce EMT in breast

cancer cells, and demonstrate that RGS10 has utility as a prognostic biomarker in breast

cancer.

Discussion

Distant metastasis is a main cause of death in patients with breast cancer. Identification of

prognostic biomarkers for early distant metastasis may inform clinical decision-making and

improve patient outcomes. To the best of our knowledge, this is the first study to characterize

the role of RGS10 as a tumor suppressor and biomarker of EMT in breast cancer. RGS10 was

expressed at lower levels in breast cancer tissues than in adjacent normal breast tissues. RGS10

expression was associated with molecular subtype of breast cancer, distant metastasis, and

survival status. Patients with high compared to low RGS10 mRNA expression in breast cancer

tissues had improved DFS and OS. RGS10 protein levels were lower in the highly aggressive

breast cancer cell line MDA-MB-231 compared to the poorly aggressive and less invasive

breast cancer cell lines MCF7 and SKBR3. RGS10 reduced breast cancer cell proliferation,

colony formation, invasion, and migration by inhibiting EMT via a novel mechanism

dependent on LCN2 and miR-539-5p.

EMT is involved in normal development and morphogenic processes, including

embryogenesis and tissue regeneration. Pathological EMT promotes invasion and metastasis in

tumors through intracellular signaling, transcription factors, miRNAs, and epigenetic and

posttranslational regulators (Huang et al., 2022; Lamouille et al., 2014) . Signaling pathways

such as TGF-β, Wnt, Notch, and phosphoinositide 3-kinase/AKT contribute to EMT, often

with cross talk at various levels and feedback activation/repression mechanisms (Deshmukh et

al., 2021; Huang et al., 2022). Transcription factors such as snail, slug, ZEBl/ZEB2, and

Twist1/Twist2 induce EMT by acting on the E-box sequence of the CDH1 promoter.

Noncoding miRNAs regulate EMT by selectively targeting mRNAs of cell receptors, signaling

pathways, the cell cycle, or cell adhesion (Górecki and Rak, 2021; Huang et al., 2022; Li et al.,

2019). Epigenetic modifications, including DNA methylation and histone modifications, alter

the expression of EMT transcription factors involved in the molecular pathways of

metabolism, transcription, differentiation, and apoptosis (Górecki and Rak, 2021).

The present study identifies RGS10 as an important mediator of EMT in breast cancer.

Previous studies have demonstrated links between several RGS proteins and various cancers,

with RGS proteins acting as tumor initiators or suppressors depending on the RGS protein and

type of cancer (Li et al., 2023). RGS10 is the smallest protein of the RGS D/12 subfamily,

which function as GAPs for the Gi family Gα subunits (Almutairi et al., 2020). RGS10 has

been linked to a poor prognosis in patients with laryngeal cancer (Yin et al., 2013), liver cancer

(Wen et al., 2015), and childhood acute myeloid leukemia (Chaudhury et al., 2018). RGS10

may represent a biomarker of clinical staging for ovarian cancer and is one of five signature

genes involved in the occurrence and development of ovarian cancer. This five-gene signature

(RGS11, RGS10, RGS13, RGS4, and RGS3) is overexpressed in ovarian cancer and involved

in extracellular matrix-receptor interaction, the TGF-β signaling pathway, the Wnt signaling

pathway, and the chemokine signaling pathway. These pathways mediate proliferation,

migration, and invasion of ovarian cancer cells; in particular, TGF-β signaling plays an

important role in EMT in ovarian cancer (Hu et al., 2021).

The novel action of RGS10 in EMT in breast cancer appears to be dependent on LNC2,

also known as neutrophil gelatinase-associated lipocalin, siderocalin, uterocalin, and oncogene

24p3. LNC2 is a secreted glycoprotein of the adipokine superfamily. LCN2 expression levels

are particularly high in breast, pancreatic, ovarian, colorectal, thyroid, and bile duct cancer

tissues and tumor cell lines. Previous studies show LCN2 can promote tumorigenesis by

increasing invasion, metastasis, and proliferation while decreasing apoptosis, possibly because

LCN2 can facilitate iron intake to cancer cells and form a heterodimer with matrix

metalloproteinase-9 (Santiago-Sánchez et al., 2020). In breast cancer, LCN2 can promote

progression by inducing EMT through the estrogen receptor alpha/Slug axis

(Morales-Valencia et al., 2022; Yang et al., 2009).

The regulation of EMT in breast cancer by RGS10 may rely on upstream regulation by

miR-539-5p. miRNAs play a critical role in the cellular processes of breast cancer, including

EMT (Zhang et al., 2013; Zhao et al., 2017). Previous reports indicate that miR-539 expression

is downregulated in breast cancer tissues and cell lines (Guo et al., 2018), and miR-539 acts as

a tumor suppressor by targeting epidermal growth factor receptor (Guo et al., 2018), specificity

protein 1 (Cai et al., 2020), or laminin subunit alpha 4 (Cai et al., 2020) expression. In patients

with breast cancer, decreased expression of miR-539 was significantly associated with lymph

node metastasis. In breast cancer cells, overexpression of miR-539 inhibited the proliferation

and promoted apoptosis of breast cancer cells, suppressed EMT, and sensitized cells to

cisplatin treatment (Cai et al., 2020). Further studies are required to fully elucidate miRNA

regulation of the gene expression networks that are essential to EMT in breast cancer.

Inflammation promotes EMT in tumors, and EMT induces the production of

proinflammatory factors by cancer cells (Suarez-Carmona et al., 2017). The present study

showed RGS10 expression in SKBR3 cells was associated with the inflammatory response.

Previous reports show RGS10 regulates cellular physiology and fundamental signaling

pathways in microglia, macrophages, and T-lymphocytes. In microglia and ovarian cancer

cells, RGS10 regulates inflammatory signaling by a G protein-independent mechanism,

linking RGS10 to the inflammatory signaling mediators tumor necrosis factor alpha (TNFα) and cyclooxygenase-2 (Alqinyah et al., 2018). In macrophages, RGS10 regulates activation and polarization by suppressing the production of the inflammatory cytokines TNFα and IL6 (Dean and Hooks, 2020). These findings suggest a potential role for RGS10 in the development of an inflamed and immunosuppressive tumor microenvironment, which may have important implications for progression in breast cancer (Suarez-Carmona et al., 2017).

Our study had some limitations. First, this was a retrospective study with a small sample size. Thus, the conclusions should be confirmed in meta-analyses or large randomized controlled trials. Second, the mechanism by which the miR-539-5p/RGS10/LCN2 axis may be related to outcomes in patients with breast cancer remains to be elucidated. Biochemical characterization of the molecular mechanisms of RGS10 in breast cancer should provide additional insight into the potential of RGS10 as a biomarker of EMT, metastasis, and prognosis in breast cancer and the role of RGS10 as a therapeutic target.

In conclusion, the results of this study show that RGS10 expression is related to survival outcomes in patients with breast cancer. RGS10 protein levels were lower in the highly aggressive breast cancer cell line MDA-MB-231 compared to the poorly aggressive and less invasive breast cancer cell lines MCF7 and SKBR3. Silencing RGS10 expression effectively increased the proliferation, colony formation, invasion, and migration ability of SKBR3 cells. The miR-539-5p/RGS10/LCN2 pathway was identified as an important regulatory axis of EMT in breast cancer. These data demonstrate RGS10 may play a tumor suppressor role and be considered a biomarker of EMT and prognosis in breast cancer.

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board of Shengjing Hospital of China

Medical University.

Consent for publication

Not applicable

Availability of data and material

These data and materials are available from the corresponding author for rational reasons.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Xi Gu and Yang Liu designed this research. Yi Jiang and Yang Liu analyzed the data derived

from public databases. Yang Liu, Yi Jiang, Peng Qiu, Tie Ma, Tong Zhu, Jiawen Bu, Yueting

Hu and Ming Jin performed the experiments. Xi Gu and Yi Jiang wrote the manuscript. All

authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science foundation of China (Grant No.82103468 to

YL), Basic Research Project for Universities of Liaoning Provincial Department of Education (Grant

No.LJKMZ20221155 to YL), 345 Talent Project of Shengjing Hospital of China Medical University (to

YL) and Clinical trial Tree planting project of Shengjing Hospital of China Medical University (Grant

No.M1953 to XG).

References

Adnan M, Morton G, Hadi S (2011) **Analysis of rpoS and bolA gene expression under**

various stress-induced environments in planktonic and biofilm phase using

2(-ΔΔCT) method *Mol Cell Biochem* **357**:275-282. https: //doi.

10.1007/s11010-011-0898-y

Almutairi F, Lee JK, Rada B (2020) **Regulator of G protein signaling 10: Structure,**

expression and functions in cellular physiology and diseases *Cell Signal* **75**:109765.

https: //doi. 10.1016/j.cellsig.2020.109765

Alqinyah M, Almutairi F, Wendimu MY, Hooks SB (2018) **RGS10 Regulates the Expression**

of Cyclooxygenase-2 and Tumor Necrosis Factor Alpha through a G

Protein-Independent Mechanism *Mol Pharmacol* **94**:1103-1113. https: //doi.

10.1124/mol.118.111674

Altman MK, Alshamrani AA, Jia W, Nguyen HT, Fambrough JM, Tran SK, et al. (2015)

Suppression of the GTPase-activating protein RGS10 increases Rheb-GTP and

mTOR signaling in ovarian cancer cells *Cancer Lett* **369**:175-183. https: //doi.

10.1016/j.canlet.2015.08.012

American Cancer Society (2023) **Survival Rates for Breast Cancer.**

https://www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-canc

er-diagnosis/breast-cancer-survival-rates.html. https: //doi.

Bianchini G, Balko JM, Mayer IA, Sanders ME, Gianni L (2016) **Triple-negative breast**

cancer: challenges and opportunities of a heterogeneous disease *Nat Rev Clin*

Oncol **13**:674-690. https: //doi. 10.1038/nrclinonc.2016.66

Cacan E, Ali MW, Boyd NH, Hooks SB, Greer SF (2014) **Inhibition of HDAC1 and DNMT1**

modulate RGS10 expression and decrease ovarian cancer chemoresistance *PLoS*

One **9**:e87455. https: //doi. 10.1371/journal.pone.0087455

Cai F, Chen L, Sun Y, He C, Fu D, Tang J (2020) **MiR-539 inhibits the malignant behavior**

of breast cancer cells by targeting SP1 *Biochem Cell Biol* **98**:426-433. https: //doi.

10.1139/bcb-2019-0111

Caldiran FY, Cacan E (2022) **RGS10 suppression by DNA methylation is associated with**

low survival rates in colorectal carcinoma *Pathol Res Pract* **236**:154007. https: //doi.

10.1016/j.prp.2022.154007

Chaudhury S, O'Connor C, Cañete A, Bittencourt-Silvestre J, Sarrou E, Prendergast Á, et al.

(2018) **Age-specific biological and molecular profiling distinguishes paediatric**

from adult acute myeloid leukaemias *Nat Commun* **9**:5280. https: //doi.

10.1038/s41467-018-07584-1

Coates AS, Winer EP, Goldhirsch A, Gelber RD, Gnant M, Piccart-Gebhart M, et al. (2015)

Tailoring therapies--improving the management of early breast cancer: St Gallen

International Expert Consensus on the Primary Therapy of Early Breast Cancer

2015 *Ann Oncol* **26**:1533-1546. https: //doi. 10.1093/annonc/mdv221

Dean P, Hooks SJTFJ (2020) **Role of RGS10 in the ovarian cancer microenvironment**

34:1-1. https: //doi.

Deshmukh AP, Vasaikar SV, Tomczak K, Tripathi S, den Hollander P, Arslan E, et al. (2021)

Identification of EMT signaling cross-talk and gene regulatory networks by

single-cell RNA sequencing *Proc Natl Acad Sci U S A* **118**. https: //doi.

10.1073/pnas.2102050118

Górecki I, Rak B (2021) **The role of microRNAs in epithelial to mesenchymal transition**

and cancers; focusing on mir-200 family *Cancer Treat Res Commun* **28**:100385.

https: //doi. 10.1016/j.ctarc.2021.100385

Gradishar WJ, Moran MS, Abraham J, Aft R, Agnese D, Allison KH, et al. (2021) **NCCN**

guidelines® insights: Breast cancer, version 4.2021 *J Natl Compr Canc Netw*

19:484-493. https: //doi. 10.6004/jnccn.2021.0023

Guo J, Gong G, Zhang B (2018) **miR-539 acts as a tumor suppressor by targeting**

epidermal growth factor receptor in breast cancer *Sci Rep* **8**:2073. https: //doi.

10.1038/s41598-018-20431-z

Győrffy B (2021) **Survival analysis across the entire transcriptome identifies biomarkers**

with the highest prognostic power in breast cancer *Comput Struct Biotechnol J*

19:4101-4109. https: //doi. 10.1016/j.csbj.2021.07.014

Hooks SB, Callihan P, Altman MK, Hurst JH, Ali MW, Murph MM (2010) **Regulators of**

G-Protein signaling RGS10 and RGS17 regulate chemoresistance in ovarian

cancer cells *Mol Cancer* **9**:289. https: //doi. 10.1186/1476-4598-9-289

Hooks SB, Murph MM (2015) **Cellular deficiency in the RGS10 protein facilitates**

chemoresistant ovarian cancer *Future Med Chem* **7**:1483-1489. https: //doi.

10.4155/fmc.15.81

Hu Y, Zheng M, Wang S, Gao L, Gou R, Liu O, et al. (2021) **Identification of a five-gene**

signature of the RGS gene family with prognostic value in ovarian cancer

Genomics **113**:2134-2144. https: //doi. 10.1016/j.ygeno.2021.04.012

Huang Y, Hong W, Wei X (2022) **The molecular mechanisms and therapeutic strategies of**

EMT in tumor progression and metastasis *J Hematol Oncol* **15**:129. https: //doi.

10.1186/s13045-022-01347-8

Hunt TW, Fields TA, Casey PJ, Peralta EG (1996) **RGS10 is a selective activator of G alpha**

i GTPase activity *Nature* **383**:175-177. https: //doi. 10.1038/383175a0

Lai X, Li Q, Wu F, Lin J, Chen J, Zheng H, et al. (2020) **Epithelial-mesenchymal transition**

and metabolic switching in cancer: Lessons from somatic cell reprogramming

Front Cell Dev Biol **8**:760. https: //doi. 10.3389/fcell.2020.00760

Lamouille S, Xu J, Derynck R (2014) **Molecular mechanisms of epithelial-mesenchymal**

transition *Nat Rev Mol Cell Biol* **15**:178-196. https: //doi. 10.1038/nrm3758

Lee AV, Oesterreich S, Davidson NE (2015) **MCF-7 cells--changing the course of breast**

cancer research and care for 45 years *J Natl Cancer Inst* **107**. https: //doi.

10.1093/jnci/djv073

Li CJ, Chu PY, Yiang GT, Wu MY (2019) **The molecular mechanism of**

epithelial-mesenchymal transition for breast carcinogenesis *Biomolecules* **9**. https:

//doi. 10.3390/biom9090476

Li L, Xu Q, Tang C (2023) **RGS proteins and their roles in cancer: friend or foe?** *Cancer*

Cell Int **23**:81. https: //doi. 10.1186/s12935-023-02932-8

Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP (2011)

Molecular signatures database (MSigDB) 3.0 *Bioinformatics* **27**:1739-1740. https:

//doi. 10.1093/bioinformatics/btr260

Liu F, Gu LN, Shan BE, Geng CZ, Sang MX (2016) **Biomarkers for EMT and MET in**

breast cancer: An update *Oncol Lett* **12**:4869-4876. https: //doi.

10.3892/ol.2016.5369

Merlin JL, Barberi-Heyob M, Bachmann N (2002) **In vitro comparative evaluation of**

trastuzumab (Herceptin) combined with paclitaxel (Taxol) or docetaxel

(Taxotere) in HER2-expressing human breast cancer cell lines *Ann Oncol*

13:1743-1748. https: //doi. 10.1093/annonc/mdf263

Miglietta F, Bottosso M, Griguolo G, Dieci MV, Guarneri V (2022) **Major advancements in**

metastatic breast cancer treatment: when expanding options means prolonging

survival *ESMO Open* **7**:100409. https: //doi. 10.1016/j.esmoop.2022.100409

Montel V, Huang TY, Mose E, Pestonjamasp K, Tarin D (2005) **Expression profiling of**

primary tumors and matched lymphatic and lung metastases in a xenogeneic

breast cancer model *Am J Pathol* **166**:1565-1579. https: //doi.

10.1016/s0002-9440(10)62372-3

Morales-Valencia J, Lau L, Martí-Nin T, Ozerdem U, David G (2022) **Therapy-induced**

senescence promotes breast cancer cells plasticity by inducing Lipocalin-2

expression *Oncogene* **41**:4361-4370. https: //doi. 10.1038/s41388-022-02433-4

Nguyen HT, Tian G, Murph MM (2014) **Molecular epigenetics in the management of**

ovarian cancer: are we investigating a rational clinical promise? *Front Oncol* **4**:71.

https: //doi. 10.3389/fonc.2014.00071

Rahman R, Dhruba SR, Matlock K, De-Niz C, Ghosh S, Pal R (2019) **Evaluating the**

consistency of large-scale pharmacogenomic studies *Brief Bioinform* **20**:1734-1753.

https: //doi. 10.1093/bib/bby046

Santiago-Sánchez GS, Pita-Grisanti V, Quiñones-Díaz B, Gumpper K, Cruz-Monserrate Z,

Vivas-Mejía PE (2020) **Biological functions and therapeutic potential of lipocalin 2**

in cancer *Int J Mol Sci* **21**. https: //doi. 10.3390/ijms21124365

Singh S, Chakrabarti R (2019) **Consequences of emt-driven changes in the immune**

microenvironment of breast cancer and therapeutic response of cancer cells *J Clin*

Med **8**. https: //doi. 10.3390/jcm8050642

Steeg PS (2005) **New insights into the tumor metastatic process revealed by gene**

expression profiling *Am J Pathol* **166**:1291-1294. https: //doi.

10.1016/s0002-9440(10)62348-6

Suarez-Carmona M, Lesage J, Cataldo D, Gilles C (2017) **EMT and inflammation:**

inseparable actors of cancer progression *Mol Oncol* **11**:805-823. https: //doi.

10.1002/1878-0261.12095

Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. (2021) **Global**

cancer statistics 2020: Globocan estimates of incidence and mortality worldwide

for 36 cancers in 185 countries *CA Cancer J Clin* **71**:209-249. https: //doi.

10.3322/caac.21660

Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. (2015)

STRING v10: protein-protein interaction networks, integrated over the tree of

life *Nucleic Acids Res* **43**:D447-452. https: //doi. 10.1093/nar/gku1003

Wang Z, Chen J, Wang S, Sun Z, Lei Z, Zhang HT, et al. (2022) **RGS6 suppresses**

TGF-β-induced epithelial-mesenchymal transition in non-small cell lung cancers

via a novel mechanism dependent on its interaction with SMAD4 *Cell Death Dis*

13:656. https: //doi. 10.1038/s41419-022-05093-0

Wen L, Li J, Guo H, Liu X, Zheng S, Zhang D, et al. (2015) **Genome-scale detection of**

hypermethylated CpG islands in circulating cell-free DNA of hepatocellular

carcinoma patients *Cell Res* **25**:1376. https: //doi. 10.1038/cr.2015.141

Wolff AC, Hammond ME, Hicks DG, Dowsett M, McShane LM, Allison KH, et al. (2013)

Recommendations for human epidermal growth factor receptor 2 testing in

breast cancer: American Society of Clinical Oncology/College of American

Pathologists clinical practice guideline update *J Clin Oncol* **31**:3997-4013. https:

//doi. 10.1200/jco.2013.50.9984

Yang J, Bielenberg DR, Rodig SJ, Doiron R, Clifton MC, Kung AL, et al. (2009) **Lipocalin 2**

promotes breast cancer progression *Proc Natl Acad Sci U S A* **106**:3913-3918. https:

//doi. 10.1073/pnas.0810617106

Yin W, Wang P, Wang X, Song W, Cui X, Yu H, et al. (2013) **Identification of microRNAs**

and mRNAs associated with multidrug resistance of human laryngeal cancer

Hep-2 cells *Braz J Med Biol Res* **46**:546-554. https: //doi.

10.1590/1414-431x20131662

Zhang K, Corsa CA, Ponik SM, Prior JL, Piwnica-Worms D, Eliceiri KW, et al. (2013) **The**

collagen receptor discoidin domain receptor 2 stabilizes SNAIL1 to facilitate

breast cancer metastasis *Nat Cell Biol* **15**:677-687. https: //doi. 10.1038/ncb2743

Zhao M, Ang L, Huang J, Wang J (2017) **MicroRNAs regulate the epithelial-mesenchymal**

transition and influence breast cancer invasion and metastasis *Tumour Biol*

39:1010428317691682. https: //doi. 10.1177/1010428317691682

Figure Legends

Figure 1. The expression and prognostic associations of RGS10 in breast cancer

(A) RGS10 mRNA levels in 31 normal human tissues. Data were derived from the Genotype-Tissue Expression database.

(B) RGS10 mRNA levels in cell lines representing 21 human cancers. Data were derived from the Cancer Cell Line Encyclopedia database.

(C) qRT-PCR showing RGS10 mRNA levels in freshly resected breast cancer tissues (n = 20) and matched adjacent normal breast tissues.

(D–E) Survival analyses showing disease-free survival (DFS) (D) and overall survival (OS) (E) in patients with breast cancer stratified by high versus low RGS10 mRNA levels. Data were derived from the Kaplan–Meier plotter database.

(F) Representative images showing immunohistochemical staining of RGS10 protein expression in breast cancer tissues or normal tissues (n = 133) (magnification: 200× and 400×).

(G–H) Kaplan–Meier analysis showing DFS (G) and OS (H) in patients with breast cancer stratified by presence versus absence of RGS10 protein in breast cancer tissues (n = 133).

Figure 2. RGS10 silencing increases the proliferation and migration of breast cancer cells *in vitro*

(A) Western blotting showing RGS10 protein levels in molecularly distinct breast cancer cell lines.

(B) Western blotting showing RGS10 protein levels in SKBR3 cells transfected with three independent shRNA constructs, shRNA-RGS10-161, shRNA-RGS10-321, and shRNA-RGS10-506, and shRNA-NC.

(C) CCK-8 assay showing the viability of SKBR3 cells transfected with shRNA-RGS10-161,

shRNA-RGS10-506, or shRNA-NC.

(D–F) Colony formation (D) and transwell migration/invasion (E, F) assays in SKBR3 cells

transfected with shRNA-RGS10-161, shRNA-RGS10-506, or shRNA-NC.

Figure 3. Protein–protein network interaction network and GO functional- and KEGG

pathway-enrichment analysis of genes co-expressed with RGS10

(A) Volcano plot showing differentially expressed genes between SKBR3 cells transfected

with shRNA-RGS10 or shRNA-NC.

(B–E) KEGG pathway analysis and GO enrichment analysis of differentially expressed genes

showing the ten most enriched terms. BP: biological processes; MF: molecular function; CC:

cellular compartment.

(F) LCN2, E-cadherin, and vimentin protein levels in SKBR3 cells transfected with

shRNA-RGS10 or shRNA-NC.

(G) Schematics of predicted miR-539-5p binding sites between wild-type and mutant RGS10

sequences in the 3′-untranslated regions.

(H) Relative luciferase activities detected after cotransfection of wild-type or mutant luciferase

reporter plasmids and an miR-539-mimic.

Figure 4. miR-539-5p regulates the migration, invasion, proliferation, and EMT of breast

cancer cells

(A) qPCR showing the transfection efficiency of the miR-539-5p mimic. *** $P < 0.001$,

one-way ANOVA.

(B–C) qRT-PCR and Western blotting showing RGS10 mRNA and protein levels in SKBR3

cells transfected with the miR-539-5p mimic, negative control (NC), or wild type (WT). *** P

< 0.001, one-way ANOVA.

(D) CCK-8 assay showing the viability of SKBR3 cells transfected with the miR-539-5p

mimic, NC, or WT. *** $P < 0.001$, one-way ANOVA.

(E–G) Colony formation (E) and transwell migration/invasion (F, G) assays in SKBR3 cells

transfected with the miR-539-5p mimic, NC, or WT. * $P < 0.05$, ** $P < 0.01$, Student's t test.

(H) Western blotting showing protein levels of LCN2 and biomarkers of EMT in SKBR3 cells

transfected with the miR-539-5p mimic or NC.

(I) Immunofluorescence staining showing E-cadherin, vimentin, and snail protein expression

in SKBR3 cells transfected with the miR-539-5p mimic or NC. Scale bar: 50 μm.

Figure 5. miR-539-5p inhibitor suppresses breast cancer cell proliferation and invasion

(A) qPCR showing transfection efficiency of the miR-539-5p inhibitor after 48 h. *** $P <$

0.001, one-way ANOVA.

(B–C) qRT-PCR and Western blotting showing RGS10 mRNA and protein levels in

MDA-MB-231 cells transfected with the miR-539-5p inhibitor, negative control (NC), or wild

type (WT). *** $P < 0.001$, one-way ANOVA.

(D) CCK-8 assay showing the viability of MDA-MB-231 cells transfected with the

miR-539-5p inhibitor, NC, or WT. *** $P < 0.001$, one-way ANOVA.

(E–G) Colony formation (E) and transwell migration/invasion (F, G) assays in MDA-MB-231

cells transfected with the miR-539-5p inhibitor, NC, or WT. * $P < 0.05$, ** $P < 0.01$, Student's

t test.

(H) Western blotting showing protein levels of LCN2 and biomarkers of EMT in

MDA-MB-231 cells transfected with the miR-539-5p inhibitor or NC.

(I) Immunofluorescence staining showing E-cadherin, vimentin, and snail protein expression

in MDA-MB-231 cells transfected with the miR-539-5p inhibitor or NC. Scale bar: 50 μm.

Figure 6. RGS10 inhibits breast cancer growth by targeting LCN2 *in vivo*

(A) Size of tumors derived from RGS10-depleted SKBR3 cells , negative control (NC) and

wild type(WT).

(B) Volume of tumors derived from RGS10-depleted SKBR3 cells , NC and WT.

(C) Hematoxylin and eosin staining of tumors derived from RGS10-depleted SKBR3 cells and

NC.

(D) Immunohistochemical staining showing LCN2, E-cadherin, snail, and vimentin protein

expression in tumors derived from RGS10-depleted SKBR3 cells and NC.

Table 1. Clinicopathological characteristics of 20 patients

Clinicopathological characteristic	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5
Sex	Female	Female	Female	Female	Female
Age (years)	58	40	70	38	58
T grade	2	1	2	1	1
N grade	0	1	3	0	1
Menopausal status	Postmenopausal	Premenopausal	Postmenopausal	Premenopausal	Postmenopausal
Subclass	Luminal A	Luminal A	Luminal A	Luminal A	Luminal A
Metastasis	No	No	No	No	No
	Patient 6	Patient 7	Patient 8	Patient 9	Patient 10
Sex	Female	Female	Female	Female	Female
Age (years)	59	56	80	55	37
T grade	2	2	3	2	1
N grade	3	0	2	2	3

Menopausal status	Postmenopausal	Postmenopausal	Postmenopausal	Postmenopausal	Premenopausal
Subclass	Luminal B	Luminal B	Luminal B	Luminal B	Luminal B
Metastasis	No	No	No	No	No
	Patient 11	Patient 12	Patient 13	Patient 14	Patient 15
Sex	Female	Female	Female	Female	Female
Age (years)	62	57	41	53	58
T grade	2	2	2	1	2
N grade	3	1	3	3	1
Menopausal status	Postmenopausal	Postmenopausal	Premenopausal	Postmenopausal	Postmenopausal
Subclass	HER2 positive	HER2 positive	HER2 positive	HER2 positive	HER2 positive
Metastasis	No	No	No	No	No
	Patient 16	Patient 17	Patient 18	Patient 19	Patient 20
Sex	Female	Female	Female	Female	Female
Age (years)	50	37	35	51	55
T grade	2	2	2	2	2
N grade	0	0	0	0	0

Menopausal status	Premenopausal	Premenopausal	Premenopausal	Premenopausal	Postmenopausal
Subclasses	Triple negative	Triple negative	Triple negative	Triple negative	Triple negative
Metastasis	No	No	No	No	No

732
733   HER2, human epidermal growth factor receptor 2.

Table 2. Correlations between RGS10 expression and clinicopathological characteristics

Variable	RGS10 expression (n, %)	RGS10 expression (n, %)	*P*-value
No. of patients	34 (25.6)	99 (74.4)	
Age (years)			0.590
≤45	8 (23.5)	28 (28.3)	
>45	26 (76.5)	71 (71.7)	
Other disease			0.746
Yes	28 (82.4)	79 (79.8)	
No	6 (17.6)	20 (20.2)	
Histological grade			0.744
I	12 (35.3)	30 (30.3)	
II	19 (55.9)	56 (56.6)	
III	3 (8.8)	13 (13.1)	
Tumor size (cm)			0.101
Median (range)	2.191 (1.0–8.0)	1.861 (0.6–4.0)	
No. of positive axillary lymph nodes			0.316
Median (range)	1.06 (0–19)	1.93 (0–27)	
No. of pregnancies			0.706
Median (range)	2.50 (1–6)	2.40 (0–7)	
No. of births			0.497
Median (range)	1.62 (1–3)	1.75 (0–7)	
Age at first pregnancy (years)			0.489

<21	1 (2.9)	6 (6.1)	
21–25	15 (44.1)	42 (42.4)	
26–30	18 (52.9)	46 (46.5)	
>30	0 (0)	5 (5.0)	
Menopausal status			0.216
Premenopausal	21 (61.8)	49 (49.5)	
Postmenopausal	13 (38.2)	50 (50.5)	
Estrogen receptor status			0.397
Positive	25 (73.5)	65 (65.7)	
Negative	9 (26.5)	34 (34.3)	
Progesterone receptor status			0.527
Positive	25 (73.5)	78 (78.8)	
Negative	9 (26.5)	21 (21.2)	
HER2 status			0.496
Positive	6 (17.6)	23 (23.2)	
Negative	28 (82.4)	76 (76.8)	
Ki67 status			0.773
>20	14 (41.2)	38 (38.4)	
≤20	20 (58.8)	61 (61.6)	
Molecular Type			0.043
Luminal A	6 (17.6)	30 (30.3)	
Luminal B	19 (55.9)	36 (36.3)	
HER2+	2 (5.9)	16 (16.2)	
Triple-negative breast cancer	7 (20.6)	17 (17.2)	

Distant metastasis			0.008
Yes	7 (20.6)	46 (46.5)	
No	27 (79.4)	53 (53.5)	
Disease-free survival (months)			<0.001
Median (range)	114.71 (53–140)	89.66 (9–132)	
Death			0.024
Yes	5 (14.7)	35 (35.4)	
No	29 (85.3)	64 (64.6)	
Overall survival (months)			0.030
Median (range)	120.41 (80–140)	107.64 (24–140)	

736
737 HER2, human epidermal growth factor receptor 2.
738

Table 3. Univariate and multivariate Cox regression analyses of clinicopathological risk factors for disease-free survival

Variable	DFS			
	Univariate analysis		Multivariate analysis	
	HR (95%CI)	*P*-value	HR (95%CI)	*P*-value
Age (years)	0.720 (0.400–1.296)	0.273	NA	
Other disease	0.885 (0.445–1.762)	0.728	NA	
Histological grade				
I		0.003		0.004
II	2.636 (1.263–5.503)	0.010	2.704 (1.294–5.649)	0.008
III	4.638 (1.877–11.462)	0.001	4.566 (1.846–11.294)	0.001
Tumor size (cm)	1.065 (0.840–1.350)	0.605	NA	
No. of positive axillary lymph nodes	1.012 (0.953–1.074)	0.698	NA	
No. of pregnancies	0.835 (0.662–1.054)	0.129	NA	
No. of births	0.694 (0.481–1.001)	0.051	NA	
Age at first pregnancy (years)			NA	
<21		0.501		
21–25	3.336 (0.449–24.810)	0.239		
26–30	3.886	0.182		

	(0.530–28.499)			
>30	1.838	0.667		
	(0.115–29.429)			
Menopausal status	0.844 (0.492–1.451)	0.540	NA	
Estrogen receptor status	1.534 (0.820–2.869)	0.180	NA	
Progesterone receptor status	1.098 (0.577–2.089)	0.776	NA	
HER2 status	0.841 (0.433–1.635)	0.610	NA	
Ki67 status	0.816 (0.465–1.431)	0.478	NA	
Molecular type			NA	
Luminal A		0.307		
Luminal B	1.940 (0.819–4.593)	0.132		
HER2+	1.147 (0.479–2.747)	0.758		
Triple-negative breast cancer	1.453 (0.527–4.007)	0.470		
RGS10 expression	0.323 (0.146–0.716)	0.005	0.321 (0.144–0.713)	0.005

742
743 HER2, human epidermal growth factor receptor 2; NA, not analyzed.

Table 4. Univariate and multivariate Cox regression analyses of clinicopathological risk factors for overall survival

Variable	Overall survival			
	Univariate analysis		Multivariate analysis	
	HR (95%CI)	*P*-value	HR (95%CI)	*P*-value
Age (years)	0.423 (0.226–0.792)	0.007	0.305 (0.154–0.603)	0.001
Other disease	0.984 (0.453–2.136)	0.967	NA	
Histological grade				
I		0.067		0.008
II	1.519 (0.703–3.284)	0.288	1.834 (0.840–4.007)	0.128
III	3.062 (1.180–7.943)	0.021	5.090 (1.812–14.300)	0.002
Tumor size (cm)	1.045 (0.788–1.385)	0.760	NA	
No. of positive axillary lymph nodes	1.011 (0.943–1.083)	0.759	NA	
No. of pregnancies	0.941 (0.731–1.213)	0.640	NA	
No. of births	0.896 (0.625–1.285)	0.550	NA	
Age at first pregnancy (years)			NA	
<21		0.580		
21–25	2.818 (0.378–21.001)	0.312		
26–30	2.021 (0.269–15.190)	0.494		
>30	3.255 (0.295–35.934)	0.335		
Menopausal status	0.769 (0.405–1.458)	0.420	NA	

Estrogen receptor status	0.769 (0.405–1.458)	0.420	NA	
Progesterone receptor status	1.091 (0.519–2.293)	0.818	NA	
HER2 status	0.587 (0.246–1.398)	0.229	NA	
Ki67 status	0.718 (0.370–1.392)	0.327	NA	
Molecular type			NA	
Luminal A		0.589		
Luminal B	0.639 (0.295–1.383)	0.255		
HER2+	0.756 (0.272–2.100)	0.592		
Triple-negative breast cancer	1.079 (0.467–2.493)	0.860		
RGS10 expression	0.353 (0.138–0.900)	0.029	0.364 (0.154–0.603)	0.001

747
748 HER2, human epidermal growth factor receptor 2; NA, not analyzed.

750   **Table 5. Gene sets enriched in phenotype 'High'**

MSigDB collection	Gene set name	NES	NOM *P*-value	FDR q-value
h.all.v6.2.symbols .gmt [Hallmarks]	HALLMARK_ALLOGRAFT_REJE CTION	-2.5 5	0	0
	HALLMARK_APOPTOSIS	-2.4 1	0	0
	HALLMARK_IL6_JAK_STAT3_SI GNALING	-2.4 0	0	0
	HALLMARK_IL2_STAT5_SIGNA LING	-2.3 9	0	0
	HALLMARK_INFLAMMATORY_ RESPONSE	-2.3 7	0	0
	HALLMARK_INTERFERON_GA MMA_RESPONSE	-2.3 3	0	0

	NES	NOM	FDR
HALLMARK_COMPLEMENT	-2.27	0	0
HALLMARK_KRAS_SIGNALING_UP	-2.25	0	0
HALLMARK_INTERFERON_ALPHA_RESPONSE	-2.22	0	0
HALLMARK_APICAL_JUNCTION	-2.13	0	0

MSigDB, Molecular Signatures Database; NES, normalized enrichment score; NOM, nominal; FDR, false discovery rate. Gene sets with NOM p-value <0.05 and FDR q-value <0.25 were considered as significant.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6